



INSTITUT
FOURIER

Probabilité

Excursion sets of correlated Random Fields

Application to continuum percolation theory

Emmanuel Roubin emmanuel.roubin@univ-grenoble-alpes.fr

05/03/2021

Laboratoire 3SR

Université Grenoble Alpes (France)




Motivations



Specimen

 O. Stamati, E. Roubin, E. Andó and Y. Malecot, *Phase segmentation of concrete x-ray tomographic images at meso-scale: Validation with neutron tomography*, 2018.

Motivations



Specimen


X-Ray Tomography

O. Stamati, E. Roubin, E. Andó and Y. Malecot, *Phase segmentation of concrete x-ray tomographic images at meso-scale: Validation with neutron tomography*, 2018.

Motivations



Specimen




X-Ray Tomography

 O. Stamati, E. Roubin, E. Andó and Y. Malecot, *Phase segmentation of concrete x-ray tomographic images at meso-scale: Validation with neutron tomography*, 2018.


Motivations



Specimen



X-Ray Tomography




Morphology Identification

 O. Stamati, E. Roubin, E. Andó and Y. Malecot, *Phase segmentation of concrete x-ray tomographic images at meso-scale: Validation with neutron tomography*, 2018.


Motivations



Specimen




X-Ray Tomography




Morphology Identification

Motivations




Specimen

X-Ray Tomography




Morphology Identification



Simulations (model)
Cracks / displacements

Motivations



 O. Stamati, E. Roubin, E. Andó and Y. Malecot, *Phase segmentation of concrete x-ray tomographic images at meso-scale: Validation with neutron tomography*, 2018.

Objectives being able to represent all kinds of morphologies

Objectives being able to represent all kinds of morphologies
... or at least dense granular morphologies

Outline

Morphological model based on correlated Random Fields

Morphological models

Excursions of correlated Random Fields

The Excursion Set Theory

$N + 1$ measures for N -dimensional spaces

Expectations of the measures

Application to percolation

Our positioning

The Euler Characteristic: percolation criterion

Morphological model based on correlated Random Fields

Morphological model based on correlated Random Fields

Morphological models

Excursions of correlated Random Fields

The Excursion Set Theory

$N + 1$ measures for N -dimensional spaces

Expectations of the measures

Application to percolation

Our positioning

The Euler Characteristic: percolation criterion

Goals:


- **Random** aspect of heterogeneities shape and positions
- **Discrete** aspect
- **Control** geometrical and topological quantities (volume, surface, size distribution of the heterogeneities. . .)

Morphological models

Goals:

- **Random** aspect of heterogeneities shape and positions
- **Discrete** aspect
- **Control** geometrical and topological quantities (volume, surface, size distribution of the heterogeneities. . .)

Hard spheres packing:




Morphological models

Goals:

- **Random** aspect of heterogeneities shape and positions
- **Discrete** aspect
- **Control** geometrical and topological quantities (volume, surface, size distribution of the heterogeneities. . .)

Hard spheres packing:




Simple, natural and efficient

Morphological models

Goals:



- **Random** aspect of heterogeneities shape and positions
- **Discrete** aspect
- **Control** geometrical and topological quantities (volume, surface, size distribution of the heterogeneities. . .)

Hard spheres packing:



Simple, natural and efficient, one “kind” of morphology, ideal shapes, heavy



Excursion set of correlated Random Fields:



 R. Adler, *Some new random field tools for spatial analysis*, 2008.

 E. Roubin, J.-B. Colliat N. Benkemoun, *Meso-scale modeling of concrete: a morphological description based on excursion sets of Random Fields*, 2015.

Excursion set of correlated Random Fields:





- Different “kinds” of morphologies, light, random shapes, evolutive

 R. Adler, *Some new random field tools for spatial analysis*, 2008.

 E. Roubin, J.-B. Colliat N. Benkemoun, *Meso-scale modeling of concrete: a morphological description based on excursion sets of Random Fields*, 2015.

Excursion set of correlated Random Fields:



- Different “kinds” of morphologies, light, random shapes, evolutive
- Hard to control, distribution less natural, smooth surfaces (for now)

▣ R. Adler, *Some new random field tools for spatial analysis*, 2008.

▣ E. Roubin, J.-B. Colliat N. Benkemoun, *Meso-scale modeling of concrete: a morphological description based on excursion sets of Random Fields*, 2015.

Morphological model based on correlated Random Fields

Morphological models

Excursions of correlated Random Fields

The Excursion Set Theory

$N + 1$ measures for N -dimensional spaces

Expectations of the measures

Application to percolation

Our positioning

The Euler Characteristic: percolation criterion

Technically to define a **strictly stationary correlated Random Field** we have to define:

- A constant probability distribution over the spatial parameter \mathbf{x} . $g(\mathbf{x})$ can be seen as a RV X . A classical distribution is the **Gaussian distribution** $\mathcal{N}(\mu, \sigma)$ where μ is the mean value and σ the standard deviation:

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-(x-\mu)^2/2\sigma^2}$$

- A covariance function which depends on the distance between two points in space $d = \|\mathbf{x} - \mathbf{y}\|$. A classical choice is the **Gaussian covariance function**:

$$C(d) = \sigma^2 e^{-d^2/L_c^2}$$



Correlated Random Fields

Correlation length L_c


The Gaussian correlation function

$$C(d) = \sigma^2 e^{-d^2/L_c^2}$$

has a single structural parameter L_c called the **correlation length**.



Large L_c




Small L_c

Correlated Random Fields


Out of topic... but other classes of covariance functions bring more flexibility.

With the **Matérn class** we can play with the roughness (additional parameter ν):


$$C(d) = \frac{\sigma^2}{\Gamma(\nu)2^{1-\nu}} \left(\frac{\sqrt{2\nu}d}{L_c} \right)^\nu K_\nu \left(\frac{\sqrt{2\nu}d}{L_c} \right)$$



$\nu \rightarrow \infty$
Gaussian



$\nu = 3/2$



$\nu = 1/2$
Exponential

An **excursion set** \mathcal{E}_s is the result of the “**threshold**” of a realisation of a RF:

$$\mathcal{E}_s = \{\boldsymbol{x} \in M \mid g(\boldsymbol{x}) \in \mathcal{H}_s\}$$

where M is the domain of definition of the RF and \mathcal{H}_s the so called **Hitting Set**.


Excursion sets

An **excursion set** \mathcal{E}_s is the result of the “**threshold**” of a realisation of a RF:

$$\mathcal{E}_s = \{\mathbf{x} \in M \mid g(\mathbf{x}) \in \mathcal{H}_s\}$$

where M is the domain of definition of the RF and \mathcal{H}_s the so called **Hitting Set**.

For example if we set $\mathcal{H}_s =]-\infty; \kappa]$ we have $\mathcal{E}_s(\kappa) = \{\mathbf{x} \in M \mid g(\mathbf{x}) \leq \kappa\}$



Excursion with “low” threshold


Excursion sets

An **excursion set** \mathcal{E}_s is the result of the “**threshold**” of a realisation of a RF:

$$\mathcal{E}_s = \{\mathbf{x} \in M \mid g(\mathbf{x}) \in \mathcal{H}_s\}$$

where M is the domain of definition of the RF and \mathcal{H}_s the so called **Hitting Set**.

For example if we set $\mathcal{H}_s =] - \infty; \kappa]$ we have $\mathcal{E}_s(\kappa) = \{\mathbf{x} \in M \mid g(\mathbf{x}) \leq \kappa\}$



Excursion with “low” threshold


Excursion sets

An **excursion set** \mathcal{E}_s is the result of the “**threshold**” of a realisation of a RF:


$$\mathcal{E}_s = \{\mathbf{x} \in M \mid g(\mathbf{x}) \in \mathcal{H}_s\}$$

where M is the domain of definition of the RF and \mathcal{H}_s the so called **Hitting Set**.

For example if we set $\mathcal{H}_s =] - \infty; \kappa]$ we have $\mathcal{E}_s(\kappa) = \{\mathbf{x} \in M \mid g(\mathbf{x}) \leq \kappa\}$



Excursion with “low” threshold



Excursion with “high” threshold


Excursion sets

An **excursion set** \mathcal{E}_s is the result of the “**threshold**” of a realisation of a RF:


$$\mathcal{E}_s = \{\mathbf{x} \in M \mid g(\mathbf{x}) \in \mathcal{H}_s\}$$

where M is the domain of definition of the RF and \mathcal{H}_s the so called **Hitting Set**.

For example if we set $\mathcal{H}_s =] - \infty; \kappa]$ we have $\mathcal{E}_s(\kappa) = \{\mathbf{x} \in M \mid g(\mathbf{x}) \leq \kappa\}$



Excursion with “low” threshold




Excursion with “high” threshold

Correlated Random Fields

$$g : \Omega \times \mathbb{R}^3 \mapsto \mathbb{R}$$

Continuous aspect
parametric variability

Medium L_c




Correlated Random Fields


$$g : \Omega \times \mathbb{R}^3 \mapsto \mathbb{R}$$

Continuous aspect
parametric variability


Large L_c



Medium L_c



Small L_c




Excursion sets

Correlated Random Fields


$$g : \Omega \times \mathbb{R}^3 \mapsto \mathbb{R}$$

Continuous aspect
parametric variability


Large L_c



Medium L_c



Small L_c




Heterogeneity sizes

Excursion sets

Correlated Random Fields

$$g : \Omega \times \mathbb{R}^3 \mapsto \mathbb{R}$$

Continuous aspect
parametric variability




Excursion sets

Correlated Random Fields

$$g : \Omega \times \mathbb{R}^3 \mapsto \mathbb{R}$$

Continuous aspect
parametric variability




Heterogeneity sizes

Observation scale

Excursion sets

$$\mathcal{E}_s = \{x \in M \mid g(x) \in \mathcal{H}_s\}$$

Discrete aspect
explicit morphology



Excursion sets

Correlated Random Fields


$$g : \Omega \times \mathbb{R}^3 \mapsto \mathbb{R}$$

Continuous aspect
parametric variability





Excursion sets

$$\mathcal{E}_s = \{x \in M \mid g(x) \in \mathcal{H}_s\}$$

Discrete aspect
explicit morphology



Excursion sets



The Excursion Set Theory

Morphological model based on correlated Random Fields

Morphological models

Excursions of correlated Random Fields

The Excursion Set Theory

$N + 1$ measures for N -dimensional spaces

Expectations of the measures

Application to percolation

Our positioning

The Euler Characteristic: percolation criterion

Family of measures

It exists several **family of measures** (Minkowski functionals, Lipschitz-Killing curvatures...). In an N -dimensional space, the size of the family is $N + 1$ where each element can be seen as a n -dimensional measure.

Each measure can be classified into two types:

- **geometrical measures** ($1 \leq n \leq N$)
- **topological measure** ($n = 0$)

Family of measures

It exists several **family of measures** (Minkowski functionals, Lipschitz-Killing curvatures...). In an N -dimensional space, the size of the family is $N + 1$ where each element can be seen as a n -dimensional measure.

Each measure can be classified into two types:

- **geometrical measures** ($1 \leq n \leq N$)
- **topological measure** ($n = 0$)

Here we simplify by using known **linear combinations** of those which gives:

In 2D

$n = 2$: Surface area

$n = 1$: Total curvature

$n = 0$: Euler Characteristic

Family of measures

It exists several **family of measures** (Minkowski functionals, Lipschitz-Killing curvatures...). In an N -dimensional space, the size of the family is $N + 1$ where each element can be seen as a n -dimensional measure.

Each measure can be classified into two types:

- **geometrical measures** ($1 \leq n \leq N$)
- **topological measure** ($n = 0$)

Here we simplify by using known **linear combinations** of those which gives:

In 2D

$n = 2$: Surface area

$n = 1$: Total curvature

$n = 0$: Euler Characteristic

In 3D

$n = 3$: Volume

$n = 2$: Surface area

$n = 1$: Total curvature

$n = 0$: Euler Characteristic

The Euler Characteristic: a topological measure

The Euler Characteristic is a mathematical measure that gives information on the topology of the morphology.

It enumerates n -dimensional features.


- In 2D

$$\chi = \#\{\text{connected components}\} - \#\{\text{holes}\}$$


- In 3D

$$\chi = \#\{\text{connected components}\} - \#\{\text{handles}\} + \#\{\text{holes}\}$$

Mean value of the measures over the threshold







Mean value of the measures over the threshold







Evolution of the 4 measures?





Mean value of the measures over the threshold







Mean value of the measures over the threshold







Mean value of the measures over the threshold







Mean value of the measures over the threshold







Mean value of the measures over the threshold







Mean value of the measures over the threshold







Mean value of the measures over the threshold



Mean value of the measures over the threshold



Mean value of the measures over the threshold



Morphological model based on correlated Random Fields

Morphological models

Excursions of correlated Random Fields

The Excursion Set Theory

$N + 1$ measures for N -dimensional spaces

Expectations of the measures

Application to percolation

Our positioning

The Euler Characteristic: percolation criterion

The expectation formula

In the context of **excursion sets of correlated Random Fields** each measure \mathcal{L}_j is a **Random Variable**.

They have a distribution that depends on:

- the parameters of the correlated Random Field ($C(\mathbf{x}, \mathbf{y}), f_X(x), M$)
- the hitting set (κ)

The expectation formula

In the context of **excursion sets of correlated Random Fields** each measure \mathcal{L}_j is a **Random Variable**.

They have a distribution that depends on:

- the parameters of the correlated Random Field ($C(\mathbf{x}, \mathbf{y}), f_X(x), M$)
- the hitting set (κ)

We don't know the distribution but we know its **expected value**:

$$\mathbb{E}(\mathcal{L}_j(\mathcal{E}_s)) = f(j, L_c, \mu, \sigma, M, \kappa)$$



The expectation formula

In the context of **excursion sets of correlated Random Fields** each measure \mathcal{L}_j is a **Random Variable**.

They have a distribution that depends on:

- the parameters of the correlated Random Field ($C(\mathbf{x}, \mathbf{y}), f_X(x), M$)
- the hitting set (κ)



We don't know the distribution but we know its **expected value**:

$$\mathbb{E}\{\mathcal{L}_j(\mathcal{E}_s)\} = \sum_{i=0}^{N-j} \binom{i+j}{i} \frac{\omega_{i+j}}{\omega_i \omega_j} \left(\frac{\lambda_2}{2\pi}\right)^{i/2} \mathcal{L}_{i+j}(M) \mathcal{M}_i^{\gamma}(\kappa)$$

The expectation formula

Gaussian Minkowski functionals: $\mathcal{M}_i^{\gamma_k}(\mathcal{H}_s)$

- They measure the probability of the Random Field to be in the hitting set $\mathcal{H}_s \subset \mathbb{R}^k$.
- They are Minkowski functionals associated with the measure of a Gaussian distribution γ_k .

The expectation formula

Gaussian Minkowski functionals: $\mathcal{M}_i^{\gamma_k}(\mathcal{H}_s)$

- They measure the probability of the Random Field to be in the hitting set $\mathcal{H}_s \subset \mathbb{R}^k$.
- They are Minkowski functionals associated with the measure of a Gaussian distribution γ_k .

Kinematic formula

If $\mathbf{X} = X_i$ is a standard Gaussian vector of size k in which $X_i \sim \mathcal{N}(0, \sigma^2)$ are independent and $\mathcal{H}_s \subset \mathbb{R}^k$:

$$\gamma_k(\mathcal{H}_s) = P(\mathbf{X} \in \mathcal{H}_s) = \frac{1}{\sigma^k (2\pi)^{k/2}} \int_{\mathcal{H}_s} e^{-\|\mathbf{x}\|^2 / 2\sigma^2} d\mathbf{x}$$

The expectation formula

Gaussian Minkowski functionals: $\mathcal{M}_i^{\gamma_k}(\mathcal{H}_s)$

- They measure the probability of the Random Field to be in the hitting set $\mathcal{H}_s \subset \mathbb{R}^k$.
- They are Minkowski functionals associated with the measure of a Gaussian distribution γ_k .

Kinematic formula

If $\mathbf{X} = X_i$ is a standard Gaussian vector of size k in which $X_i \sim \mathcal{N}(0, \sigma^2)$ are independent and $\mathcal{H}_s \subset \mathbb{R}^k$:

$$\gamma_k(\mathcal{H}_s) = P(\mathbf{X} \in \mathcal{H}_s) = \frac{1}{\sigma^k (2\pi)^{k/2}} \int_{\mathcal{H}_s} e^{-\|\mathbf{x}\|^2 / 2\sigma^2} d\mathbf{x}$$

If $\mathcal{K}(A, \rho)$ is the tube of A or ray ρ we have the following Taylor expansion:

$$\gamma_k(\mathcal{K}(\mathcal{H}_s, \rho)) = \sum_{j=0}^{\infty} \frac{\rho^j}{j!} \mathcal{M}_i^{\gamma_k}(\mathcal{H}_s)$$

The expectation formula

Application to scalar valued Gaussian Random Fields: $\mathcal{M}_i^\gamma(\mathcal{H}_s)$

Hitting set, Tube and expansions

$$\mathcal{H}_s \text{ and Tube } \mathcal{H}_s = [\kappa, \infty[\quad \text{and} \quad \mathcal{K}(\mathcal{H}_s) = [\kappa - \rho, \infty[$$

The expectation formula

Application to scalar valued Gaussian Random Fields: $\mathcal{M}_i^\gamma(\mathcal{H}_s)$

Hitting set, Tube and expansions

\mathcal{H}_s and Tube $\mathcal{H}_s = [\kappa, \infty[$ and $\mathcal{K}(\mathcal{H}_s) = [\kappa - \rho, \infty[$

Measures $\gamma(\mathcal{H}_s) = \frac{1}{\sigma\sqrt{2\pi}} \int_{\kappa}^{\infty} e^{-x^2/\sigma^2} dx = \bar{F}(\kappa)$ and $\gamma(\mathcal{K}(\mathcal{H}_s)) = \bar{F}(\kappa - \rho)$

The expectation formula

Application to scalar valued Gaussian Random Fields: $\mathcal{M}_i^\gamma(\mathcal{H}_s)$

Hitting set, Tube and expansions

\mathcal{H}_s and Tube $\mathcal{H}_s = [\kappa, \infty[$ and $\mathcal{K}(\mathcal{H}_s) = [\kappa - \rho, \infty[$

Measures $\gamma(\mathcal{H}_s) = \frac{1}{\sigma\sqrt{2\pi}} \int_{\kappa}^{\infty} e^{-x^2/\sigma^2} dx = \bar{F}(\kappa)$ and $\gamma(\mathcal{K}(\mathcal{H}_s)) = \bar{F}(\kappa - \rho)$

Expansions $\gamma(\mathcal{K}(\mathcal{H}_s)) = \bar{F}(\kappa - \rho) = \underbrace{\sum_{i=0}^{\infty} \frac{(-1\rho)^i}{j!} \bar{F}^{(i)}(\kappa)}_{\text{For small } \rho}$

The expectation formula

Application to scalar valued Gaussian Random Fields: $\mathcal{M}_i^\gamma(\mathcal{H}_s)$

Hitting set, Tube and expansions

\mathcal{H}_s and Tube $\mathcal{H}_s = [\kappa, \infty[$ and $\mathcal{K}(\mathcal{H}_s) = [\kappa - \rho, \infty[$

Measures $\gamma(\mathcal{H}_s) = \frac{1}{\sigma\sqrt{2\pi}} \int_{\kappa}^{\infty} e^{-x^2/\sigma^2} dx = \bar{F}(\kappa)$ and $\gamma(\mathcal{K}(\mathcal{H}_s)) = \bar{F}(\kappa - \rho)$

Expansions $\gamma(\mathcal{K}(\mathcal{H}_s)) = \bar{F}(\kappa - \rho) = \underbrace{\sum_{i=0}^{\infty} \frac{(-1)^i \rho^i}{i!} \bar{F}^{(i)}(\kappa)}_{\text{For small } \rho} = \underbrace{\sum_{i=0}^{\infty} \frac{\rho^i}{i!} \mathcal{M}_i^\gamma(\mathcal{H}_s)}_{\text{Kinematic formula}}$

The expectation formula

Application to scalar valued Gaussian Random Fields: $\mathcal{M}_i^\gamma(\mathcal{H}_s)$

Hitting set, Tube and expansions

\mathcal{H}_s and Tube $\mathcal{H}_s = [\kappa, \infty[$ and $\mathcal{K}(\mathcal{H}_s) = [\kappa - \rho, \infty[$

Measures $\gamma(\mathcal{H}_s) = \frac{1}{\sigma\sqrt{2\pi}} \int_{\kappa}^{\infty} e^{-x^2/\sigma^2} dx = \bar{F}(\kappa)$ and $\gamma(\mathcal{K}(\mathcal{H}_s)) = \bar{F}(\kappa - \rho)$

Expansions $\gamma(\mathcal{K}(\mathcal{H}_s)) = \bar{F}(\kappa - \rho) = \underbrace{\sum_{i=0}^{\infty} \frac{(-1\rho)^i}{j!} \bar{F}^{(i)}(\kappa)}_{\text{For small } \rho} = \underbrace{\sum_{i=0}^{\infty} \frac{\rho^i}{j!} \mathcal{M}_i^\gamma(\mathcal{H}_s)}_{\text{Kinematic formula}}$

Identification of the Gaussian Minkowski Functionals

$$\mathcal{M}_i^\gamma(\mathcal{H}_s) = (-1)^j \bar{F}^{(i)}(\kappa)$$

The expectation formula

Volume Fraction

$$\mathbb{E}\{\Phi\} = \frac{1}{\sqrt{\pi}} \int_{\kappa/\sigma}^{\infty} e^{-t^2} dt$$

Euler Characteristic

With the scale ratio $\beta = \text{size}(M)/L_c$

$$\mathbb{E}\{\chi\} = \left[\frac{\beta^3}{\sqrt{2}\pi^2} \left(\frac{\kappa^2}{\sigma^2} - 1 \right) + \frac{3\beta^2}{\sqrt{2}\pi^{3/2}} \frac{\kappa}{\sigma} + \frac{3\beta}{\sqrt{2}\pi} \right] e^{-\kappa^2/2\sigma^2} + \frac{1}{\sqrt{\pi}} \int_{\kappa/\sigma}^{\infty} e^{-t^2} dt$$

The expectation formula

Volume Fraction

$$\mathbb{E}\{\Phi\} = \frac{1}{\sqrt{\pi}} \int_{\kappa/\sigma}^{\infty} e^{-t^2} dt$$

Euler Characteristic

With the scale ratio $\beta = \text{size}(M)/L_c$

$$\mathbb{E}\{\chi\} = \left[\frac{\beta^3}{\sqrt{2\pi^2}} \left(\frac{\kappa^2}{\sigma^2} - 1 \right) + \frac{3\beta^2}{\sqrt{2\pi^{3/2}}} \frac{\kappa}{\sigma} + \frac{3\beta}{\sqrt{2\pi}} \right] e^{-\kappa^2/2\sigma^2} + \frac{1}{\sqrt{\pi}} \int_{\kappa/\sigma}^{\infty} e^{-t^2} dt$$

The expectation formula

Volume Fraction

$$\mathbb{E}\{\Phi\} = \frac{1}{\sqrt{\pi}} \int_{\kappa/\sigma}^{\infty} e^{-t^2} dt$$

Euler Characteristic





With the scale ratio $\beta = \text{size}(M)/L_c$

$$\mathbb{E}\{\chi\} = \left[\frac{\beta^3}{\sqrt{2\pi^2}} \left(\frac{\kappa^2}{\sigma^2} - 1 \right) + \frac{3\beta^2}{\sqrt{2\pi^{3/2}}} \frac{\kappa}{\sigma} + \frac{3\beta}{\sqrt{2\pi}} \right] e^{-\kappa^2/2\sigma^2} + \frac{1}{\sqrt{\pi}} \int_{\kappa/\sigma}^{\infty} e^{-t^2} dt$$

 R.J. Adler, *Random Fields and Geometry*, 1976.

 K.J. Worsley, *The geometry of random images*, 1996.

Mean value of the measures over the threshold



Application to percolation

Morphological model based on correlated Random Fields

Morphological models

Excursions of correlated Random Fields

The Excursion Set Theory

$N + 1$ measures for N -dimensional spaces

Expectations of the measures

Application to percolation

Our positioning

The Euler Characteristic: percolation criterion

Summary

A diagrammatic scheme has been presented for representing the structure of a three-dimensional polymer in a way which facilitates statistical analysis. It has been shown that the general condition for the formation of infinitely large molecules is expressed by $\alpha > 1/(f - 1)$, where f is the functionality of the branch units and α is the probability of chain branching as opposed to chain termination. Methods have been presented for

▣ Paul J. Flory, *Molecular size distribution in three dimensional polymers: Gelation*, 1941.

Summary

A diagrammatic scheme has been presented for representing the structure of a three-dimensional polymer in a way which facilitates statistical analysis. It has been shown that the general condition for the formation of infinitely large molecules is expressed by $\alpha > 1/(f - 1)$, where f is the functionality of the branch units and α is the probability of chain branching as opposed to chain termination. Methods have been presented for

■ Paul J. Flory, *Molecular size distribution in three dimensional polymers: Gelation*, 1941.

PERCOLATION PROCESSES

I. CRYSTALS AND MAZES

BY S. R. BROADBENT AND J. M. HAMMERSLEY

Received 15 August 1956

ABSTRACT. The paper studies, in a general way, how the random properties of a 'medium' influence the percolation of a 'fluid' through it. The treatment differs from conventional diffusion theory, in which it is the random properties of the fluid that matter. Fluid and medium bear general interpretations: for example, solute diffusing through solvent, electrons migrating over an atomic lattice, molecules penetrating a porous solid, disease infecting a community, etc.

■ S. R. Broadbent and J. M. Hammersley, *Percolation process I and II*, 1957.

The Critical Percolation Probabilities p_c




Figure 4.1 The randomly cut network as an example of percolation.

▣ R. Zallen, *The Physics of Amorphous Solids: Chapter 4 The Percolation Model*, 1983.

From (7.1), if K is singular at p_0 then it is also singular at $1 - p_0$, and if there is only one singularity these must be identical points, or

$$p_0 = \frac{1}{2}. \quad (7.2)$$

This establishes two important percolation probabilities as $\frac{1}{2}$ —that for the site problem on the triangular lattice and that for the bond problem on the simple quadratic lattice. The result (7.2) holds for any fully triangulated lattice.

▣ M. F. Sykes and J. W. Essam, *Exact Critical Percolation Probabilities for Site and Bond Problems in Two Dimensions*, 1964.

The Critical Percolation Probabilities p_c




Figure 4.1 The randomly cut network as an example of percolation.

▣ R. Zallen, *The Physics of Amorphous Solids: Chapter 4 The Percolation Model*, 1983.

From (7.1), if K is singular at p_0 then it is also singular at $1 - p_0$, and if there is only one singularity these must be identical points, or

$$p_0 = \frac{1}{2}. \quad (7.2)$$

This establishes two important percolation probabilities as $\frac{1}{2}$ —that for the site problem on the triangular lattice and that for the bond problem on the simple quadratic lattice. The result (7.2) holds for any fully triangulated lattice.

▣ M. F. Sykes and J. W. Essam, *Exact Critical Percolation Probabilities for Site and Bond Problems in Two Dimensions*, 1964.

- Lattices (graphs)
- Volumetric approach to regularise

▣ R. Zallen, *Critical density in percolation processes*, 1970.

- Boolean models

▣ K. Mecke, *Morphology of spatial patterns*, 1997.

▣ R. Schneider and W. Weil, *Stochastic and Integral Geometry*, 2008.

Links between percolation theory and topology

Percolation and topological quantification

They are **two different concepts**.

Percolation: find the existence of clusters of the size of the system

Topology: measure the connectivity

Links between percolation theory and topology

Percolation and topological quantification

They are **two different concepts**.

Percolation: find the existence of clusters of the size of the system

Topology: measure the connectivity

However it has been observed many times that **critical behaviour** takes place when the **Euler Characteristic changes sign**.

 B. L. Okun, *Euler Characteristic in Percolation Theory*, 1989.

 K. R. Mecke and H. Wagner, *Euler characteristic and related measures for random geometric sets*, 1991.

 H. Tomita and C. Murakami, *Percolation pattern in continuous media and its topology*, 1994.

Links between percolation theory and topology

Percolation and topological quantification

They are **two different concepts**.

Percolation: find the existence of clusters of the size of the system

Topology: measure the connectivity

However it has been observed many times that **critical behaviour** takes place when the **Euler Characteristic changes sign**.

- Often with analytical solutions
- Often limited to boolean problems in infinite spaces

 B. L. Okun, *Euler Characteristic in Percolation Theory*, 1989.

 K. R. Mecke and H. Wagner, *Euler characteristic and related measures for random geometric sets*, 1991.

 H. Tomita and C. Murakami, *Percolation pattern in continuous media and its topology*, 1994.

Outline

Morphological model based on correlated Random Fields

Morphological models

Excursions of correlated Random Fields

The Excursion Set Theory

$N + 1$ measures for N -dimensional spaces


Expectations of the measures

Application to percolation


Our positioning

The Euler Characteristic: percolation criterion


The Euler Characteristic




The Euler Characteristic




The Euler Characteristic




The Euler Characteristic




The Euler Characteristic




The Euler Characteristic




The Euler Characteristic




The Euler Characteristic




The Euler Characteristic




The Euler Characteristic




The Euler Characteristic




The Euler Characteristic: scale ratio




The Euler Characteristic: scale ratio




The Euler Characteristic: scale ratio




The Euler Characteristic: scale ratio




Phase diagram




Phase diagram




Phase diagram




Phase diagram




Phase diagram




Phase diagram




Phase diagram



Phase diagram



Percolation of the solid phase in N dimensions



Objectives being able to represent all kinds of morphologies

Objectives being able to represent all kinds of morphologies

Problems we're limited to certain morphologies

Objectives being able to represent all kinds of morphologies

Problems we're limited to certain morphologies

- $\exists (\mathcal{H}_s, C(X, Y), f_X)$ for a given set of \mathcal{L} ?

Objectives being able to represent all kinds of morphologies

Problems we're limited to certain morphologies

- $\exists (\mathcal{H}_s, C(X, Y), f_X)$ for a given set of \mathcal{L} ?
- $\exists!$ "type" of morphology for a given set of \mathcal{L} ?

Objectives being able to represent all kinds of morphologies

Problems we're limited to certain morphologies

- $\exists (\mathcal{H}_s, C(X, Y), f_X)$ for a given set of \mathcal{L} ?
- $\exists!$ "type" of morphology for a given set of \mathcal{L} ?
- ...

Objectives being able to represent all kinds of morphologies

Problems we're limited to certain morphologies

- $\exists (\mathcal{H}_s, C(X, Y), f_X)$ for a given set of \mathcal{L} ?
- $\exists!$ "type" of morphology for a given set of \mathcal{L} ?
- ...

Solutions ▪ Play with the distribution (or the hitting set)

Objectives being able to represent all kinds of morphologies

Problems we're limited to certain morphologies

- $\exists (\mathcal{H}_s, C(X, Y), f_X)$ for a given set of \mathcal{L} ?
- $\exists!$ "type" of morphology for a given set of \mathcal{L} ?
- ...

Solutions

- Play with the distribution (or the hitting set)
- Play with covariance function

Objectives being able to represent all kinds of morphologies

Problems we're limited to certain morphologies

- $\exists (\mathcal{H}_s, C(X, Y), f_X)$ for a given set of \mathcal{L} ?
- $\exists!$ "type" of morphology for a given set of \mathcal{L} ?
- ...

Solutions

- Play with the distribution (or the hitting set)
- Play with covariance function
- Seek help :)