


Contents lists available at [ScienceDirect](#)

## Environmental Pollution

journal homepage: [www.elsevier.com/locate/envpol](http://www.elsevier.com/locate/envpol)

## Corrigendum to ‘Varying modalities of perinatal exposure to a pesticide cocktail elicit neurological adaptations in mice and zebrafish’ [Environ. Pollut. 278 (2021) 116755]

Isabel Forner-Piquer<sup>a</sup>, Wendy Klement<sup>a</sup>, Giuseppe Gangarossa<sup>c</sup>, Emma Zub<sup>a</sup>, Frederic de Bock<sup>a</sup>, Marine Blaquiere<sup>a</sup>, Tangui Maurice<sup>d</sup>, Etienne Audinat<sup>a</sup>, Adèle Faucherre<sup>a</sup>, Frederic Lasserre<sup>b</sup>, Sandrine Ellero-Simatos<sup>b</sup>, Laurence Gamet-Payrastre<sup>b</sup>, Chris Jopling<sup>a</sup>, Nicola Marchi<sup>a,\*</sup>

<sup>a</sup> Institute of Functional Genomics, University of Montpellier, CNRS, INSERM, Montpellier, France

<sup>b</sup> Toxalim (Research Centre in Food Toxicology), Université de Toulouse, INRAE, ENVT, INP-Purpan, UPS, 31300, Toulouse, France

<sup>c</sup> Université de Paris, BFA, UMR 8251, CNRS, F-75014, Paris, France

<sup>d</sup> MMDN, University of Montpellier, EPHE, INSERM, UMR\_S1198, Montpellier, France

The authors regret a writing error in a paragraph of the Methods section (2.4. Freely moving telemetry video-electroencephalography), which we only recently noticed. We recognized that, during the manuscript assembly in 2020/21, we inadvertently included method text from an older document that referenced a discontinued practice.

Current one states:

Animals were anesthetized using an intraperitoneal injection with chloral hydrate and xylazine. Animals were placed on a heating pad during the procedure ensuring a physiological temperature.

Below is the amended text reflecting the correct and routine method

we used. The text should read:

For microelectrode implants, animals were in a sterile environment and deeply anesthetized using an intraperitoneal injection of ketamine/xylazine (100mg/kg; 10mg/kg) in sterile saline. Buprenorphine (subcutaneous, 0.05 mg/Kg) was used for analgesia and lidocaine for local anesthesia. During surgery, animals were placed on a heating pad, ensuring a physiological temperature and continuously controlling the adequate depth of anesthesia.

The authors would like to apologize for any inconvenience caused.

DOI of original article: <https://doi.org/10.1016/j.envpol.2021.116755>.

\* Corresponding author.

E-mail address: [nicola.marchi@igf.cnrs.fr](mailto:nicola.marchi@igf.cnrs.fr) (N. Marchi).

<https://doi.org/10.1016/j.envpol.2024.125432>

0269-7491/© 2024 Elsevier Ltd. All rights are reserved, including those for text and data mining, AI training, and similar technologies.