



HAL
open science

L 1 -THEORY FOR REACTION-DIFFUSION HELE-SHAW FLOW WITH LINEAR DRIFT

Noureddine Igbida

► **To cite this version:**

Noureddine Igbida. L 1 -THEORY FOR REACTION-DIFFUSION HELE-SHAW FLOW WITH LINEAR DRIFT. 2021. hal-03175798v1

HAL Id: hal-03175798

<https://hal.science/hal-03175798v1>

Preprint submitted on 21 Mar 2021 (v1), last revised 24 Mar 2023 (v4)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

L^1 -THEORY FOR REACTION-DIFFUSION HELE-SHAW FLOW WITH LINEAR DRIFT

NOUREDDINE IGBIDA[†]

ABSTRACT. The main goal of this paper is to prove L^1 -comparison and contraction principles for weak solutions (in the sense of distributions) of Hele-Shaw flow with a linear Drift. The flow is considered with a general reaction term including the Lipschitz continuous case, and subject to mixed homogeneous boundary conditions : Dirichlet and Neumann. Our approach combines a renormalization proceedings à la DiPerna-Lions with Kruzhkov device of doubling and de-doubling variables. The L^1 -contraction principle allows to handle the problem in a general framework of nonlinear semigroup theory in L^1 , taking thus advantage of this strong theory to study existence, uniqueness, comparison of weak solutions, L^1 -stability as well as many further questions.

1. INTRODUCTION AND PRELIMINARIES

1.1. **Introduction and main contributions.** Let $\Omega \subset \mathbb{R}^N$ be a bounded open set. We are interested in the existence and uniqueness as well as the L^1 -comparison principle for the weak solution concerning the PDE of the type

$$(1.1) \quad \begin{cases} \frac{\partial u}{\partial t} - \Delta p + \nabla \cdot (u V) = g(t, x, u) \\ u \in \text{Sign}(p) \end{cases} \quad \text{in } Q := (0, T) \times \Omega.$$

Here, Sign is the maximal monotone graph defined in \mathbb{R} by

$$\text{Sign}(r) = \begin{cases} 1 & \text{for any } r > 0 \\ [-1, 1] & \text{for } r = 0 \\ -1 & \text{for any } r < 0, \end{cases}$$

V and g are the given velocity field and the reaction term respectively, satisfying besides assumptions we precise next (including at least the common case of Lipschitz continuous reaction term). In the case of nonnegative solution (one phase problem), the problem may be written in the widespread form

$$(1.2) \quad \begin{cases} \frac{\partial u}{\partial t} - \Delta p + \nabla \cdot (u V) = g(t, x, u) \\ 0 \leq u \leq 1, p \geq 0, p(u - 1) = 0 \end{cases} \quad \text{in } Q.$$

The linear version of the problem which ties into the case where Sign 's graph is superseded by the identity ; that is to take $p = u$, the problem corresponds to Fokker-Planck equation. Existence, uniqueness and stability of weak solutions for this case is studied in [30] with possibly linear degenerate

Date: March 21, 2021.

[†]Institut de recherche XLIM-DMI, UMR-CNRS 7252, Faculté des Sciences et Techniques, Université de Limoges, France. E-mail : noureddine.igbida@unilim.fr .

diffusion, additive noise and BV -vector fields V in \mathbb{R}^N . One can see moreover the work [35] where the case of irregular coefficient and the associated stochastic differential equation are touch on.

In general, the problem is of elliptic-hyperbolic-parabolic type. Despite the broad results on this class of nonlinear PDE, the structure of (1.1) and (1.2) exclude them definitely out the scope of the current literature, at least concerning uniqueness (cf. [2, 7, 8, 19, 33, 41], see also the expository paper [9] for a complete list on refs). In spite of the "hard" non-linearity connected to the sign graph, the Fokker-Planck look of the equation with linear drift seems to be very fruitful for the analysis based on gradient flow in the space of probability measures equipped with Wasserstein distance (the distance arising in the Monge-Kantorovich optimal transport problem) in the one phase case (1.2) with no-flux Neumann boundary condition and no-reaction ; i.e. $g \equiv 0$. Numerous results on the existence, properties and estimates on the weak solution were elaborate in the last decade (one can see for instance [39] and the references therein). Other interesting progress on qualitative properties of a solution were obtained using the notion of viscosity solution (one can see for instance [1] and the references therein). The key utility of the viscosity concept is the skill to describe the pointwise behaviour of the free boundary evolution. Nonetheless, the uniqueness is still an open problem and seems to be harshness in general. In the case of monotone gradient like velocity field V , the uniqueness of absolute continuous solution in the set of probability equipped with Wasserstein distance (solutions in $AC([0; T], (P(\Omega), W_2))$) has been obtained craftily in [26] for (1.2), in the framework of gradient flow in euclidean Wasserstein space again in the case of Neumann boundary condition and $g \equiv 0$.

Our main results in this paper concern the L^1 -comparison and contraction principles for the reaction-diffusion-transport equation of the type (1.1) subject to mixed homogeneous boundary conditions : Dirichlet and Neumann. This allows to handle the problem in the context of nonlinear semigroup theory in $L^1(\Omega)$ to prove thus existence, uniqueness, comparison of weak solutions, L^1 -stability as well as many other questions. We'll not cover all the possibilities with this framework here, but let us mention at least its prospect to tackle the continuous dependence with respect to the Sign graph, including for instance the connection between this problem and the so called incompressible limit of the porous medium equation even in the singular case. This subject would be likely touch on in forthcoming works. We tackle the problem using a renormalisation approach of DiPerna-Lions type (cf. [27]) combined with Kruzhkov device of doubling variables. To avoid much more technicality of the paper, we restrict ourself to the case of outgoing vector field velocity on the boundary. Although, we do believe that our arguments may be generalized and used for general V with reasonable regularity. They can be extended also to the case where $\Omega = \mathbb{R}^N$ as well as to more general maximal monotone graph β instead of Sign.

Among the propose of this paper, let us mention the treatment of the equation with a reaction term g . Besides existence, uniqueness and comparison results for weak solutions, we supply sufficient condition on $g(t, x, 1)$ and $g(t, x, -1)$ to avoid a congested regime and cover the solution of (4.1) by the purely transport equation :

$$(1.3) \quad \frac{\partial u}{\partial t} + \nabla \cdot (u V) = g(t, x, u), \quad \text{in } Q.$$

The one phase problem (1.2) will be concerned as well with sufficient condition of the type $g(\cdot, 0) \geq 0$ a.e. in Q . For the application in crowd motion, the conditions on $g(\cdot, 1)$ may be heuristically connected to the reaction of the concerned population at the position $x \in \Omega$ and time $t > 0$ within "congested" circumstance.

1.2. Motivation and related works. By means of the diffusion term, the nonlinear equation of the type (1.1) is usually called Hele-Shaw Flow. Indeed, in the case where $V \equiv 0$, the equation is a

free-boundary problem modeling the evolution of a slow incompressible viscous fluid moving between slightly separated plates (cf. [29, 42, 43, 28, 25, 46] for physical and mathematical formulation). The equation appears also as the viscous incompressible limit of the porous medium equation (see for instance [32, 31, 13] and the references therein).

The study of the Hele-Shaw flow with a reaction term (without drift) goes back to [14] and [15] in the study of the limit of the m -porous medium equation as $m \rightarrow \infty$. Concrete models with a dissimilar typical reaction term appeared after in the study of the tumour growth (cf. [44]). The emergence of a drift in the Hele-Shaw flow appeared for the first time in the study of congested crowd motion (cf. [36, 37, 38]). In this case, the function u is assumptive to be nonnegative to model the density of population and the constraints on u prevents an evolution beyond a given threshold.

Completed with boundary condition (Dirichlet, Neumann or mixed Dirichlet-Neumann) and initial data, the questions of existence and uniqueness of a weak solution is well understood by now in the case where $V \equiv 0$, under general conditions on g (cf [14, 15], one can see also [44] for a different structure of a reaction term). For general V , though the existence of a solution seems to be more or less well understood, the uniqueness question seems to be very delicate and challenging. The main difficulties comes from the combination between the "hard" non-linearity Sign (so called hard-congestion) and the arbitrary pointing of velocity field for the drift. Although the first order term is linear, one notices the hyperbolic character of the equation (outside the congested region). This motivates the questions of existence and uniqueness of a weak solution, and challenges new as well as standard techniques for first order hyperbolic equations and second order hyperbolic-parabolic structures as well. In particular, one sees that setting

$$b = (I + \text{Sign}^{-1})^{-1} \quad \text{and} \quad \beta = (I + \text{Sign})^{-1},$$

$v := u - p$, the problem (1.1) falls into the scope of diffusion-transport equation of the type

$$(1.4) \quad \frac{\partial b(v)}{\partial t} - \Delta \beta(v) + \nabla \cdot (V(x)\Phi(v)) = f \quad \text{in } Q := (0, \infty) \times \Omega.$$

The study of this class of degenerate parabolic problems has under-gone a considerable progress in the last twenty years, thanks to the fundamental paper of J. Carrillo [19] in which the Kruzhkov device of doubling of variables was extended to this class of hyperbolic-parabolic-elliptic problems. In [19], the appropriate notion of solution was established : the so called weak-entropy solution. This definition (or, sometimes, parts of the uniqueness techniques of [19]) led to many developments, we refer to the expository paper [9] for a survey and complete references on this subject. As much as restrictive it is, the approach of [19] serves out primarily and ingeniously the case where V does not depend on space and homogeneous Dirichlet boundary condition. Strengthened with L^1 -nonlinear semi-group techniques (cf. [9]), the approach of [19] enables to achieve successful as well the uniqueness of weak solution in the "weakly degenerate" convection-diffusion problems of parabolic-elliptic type ; that is $[\beta = 0] \subseteq [\Phi = 0]$. As far as Dirichlet boundary condition are concerned (cf. [19, 33, 7, 7, 9]), the case of homogeneous Neumann was treated in [6]. Yet, the notion of entropic solution à la Carrillo is definitively the suitable notion for general problem of the type (1.4). Nevertheless, following the theory of hyperbolic equation, the case of linear drift is a particular case for which one expects the uniqueness of weak solution. As far as we are aware of, this case remains to be out of the scope of the current literature on this subject.

In spite of the serving of the second order term, we can not pass over the forcefulness of the first order term, which is here linear and carrying away along a vector field V with an arbitrary pointing in Ω . Heuristically, in the likely case $p \equiv 0$, the PDE turns into a linear initial-boundary value problem for continuity equation. In general, one guess the uniqueness of weak solution under reasonable assumption

on V , like bounded total variation coefficients. As far as the regularity of V is involved, the way how to handle the boundary condition is hooked to the proofs of uniqueness and adduce different difficulties to the problem. The treatment of boundary conditions is definitely a crucial step in the proof of uniqueness. This is closely connected in some sens to the regularity of the solution as well as to the pointing of V on the boundary. This affects the manner to treat the weak/strong trace of the flux on the boundary (see the papers [3], [5], [10], [20],[22]and by [5], [8], [6]). The approach based on the concept of renormalized solution introduced in 1989 by Lions and DiPerna [27] for tangential velocity field seems to be unmissable and powerful in general for the proof of uniqueness of weak solution for this kind of problem. This concept was extended to bounded domains in \mathbb{R}^N with inflow boundary conditions and velocity field with a kind of Sobolev regularity in [16], [17] and [40]. These results were generalized to velocity fields with BV regularity in [3, 4, 5, 23]. The concept was used also in [30] and [35] to tackle the existence and uniqueness of weak solutions for Fokker-Planck equation and its associated stochastic differential equation in some extreme cases like linear degenerate diffusion with irregular coefficient, additive noise and BV -vector fields V .

In this paper, we show how to use this approach in the presence of second order term of Hele-Shaw type to prove the uniqueness of weak solution in the contest of mixed Dirichlet-Neuman boundary condition with an outward pointing (possibly tangential) velocity fields V on the boundary. The case of strict inward pointing velocity fields V on the boundary need more careful, and possible delicate, treatment of the trace of the flux on the boundary. Taught, we do believe that it is possible adapt carefully our techniques to the case of strict inward pointing with the additional assumption $uV \cdot \nu = 0$ on the boundary, which can be heuristically interpreted as (a weak version of) $u = 0$ on the boundary.

1.3. Plan of the paper. In the following subsection, we give the main assumptions we'll use throughout the paper and the definition of weak solution we are dealing with. Section 2, is devoted to the L^1 -comparison principle for the weak solution. We introduce renormalized like formulations and use them with doubling and de-doubling variable techniques à la Kruzhkov inside Ω , to prove first that weak solutions satisfies some kind of local Kato's inequalities. Then, linking the outgoing assumption on the velocity field V on the boundary with the distance-to-boundary function, we prove that we can go with our Kato's inequalities up to the boundary proving thus L^1 -comparison principle, and then deduce the uniqueness. In section 3, we prove the existence of a weak solution by using nonlinear semigroup theory governed by L^1 -accretive operator. Here the main ingredient is to use the L^1 -contraction principle for weak solution of stationary problem associated with the ϵ -Euler implicit time discretization of the evolution problem. Then, we pass to the limit in the so called ϵ -approximate solution and prove that the limit is the weak solution of the evolution problem. In section 4, we treat the equation with reaction term $g(t, x, u)$. Using again nonlinear semigroup theory in $L^1(\Omega)$, we prove the existence and uniqueness of a weak solution under a general condition on g including the Lipschitz continuous case. Then, using the L^1 -comparison principle we show how some natural result concerning the existence and uniqueness of weak solution for some particular cases like one phase problem (1.2) as well the purely transport equation (1.3) (i.e. $p \equiv 0$) may be wrap up by this approach. Section 5 is devoted to some remarks, comments and possible extensions. At last, in Section 6, for completeness we give a complement for the proof of de-doubling variables process to handle space dependent vector field with arbitrary pointing out.

1.4. Preliminaries, remarks and and main assumptions. We assume that $\Omega \subset \mathbb{R}^N$ is a bounded open set, with regular boundary splitted into $\partial\Omega = \Gamma_D \cup \Gamma_N$, such that $\Gamma_D \cap \Gamma_N = \emptyset$ and

$$\mathcal{L}^{N-1}(\Gamma_D) > 0.$$

For any $h > 0$, we denote by

$$(1.1) \quad \xi_h(t, x) = \frac{1}{h} \min \left\{ h, d(x, \partial\Omega) \right\} \quad \text{and} \quad \nu_h(x) = -\nabla \xi_h, \quad \text{for any } x \in \Omega,$$

where $d(\cdot, \partial\Omega)$ names the euclidean distance-to-the-boundary function. We see that $\xi_h \in H_0^1(\Omega)$ is a regular (as well as the boundary is) concave function, $0 \leq \xi_h \leq 1$ and

$$\nu_h(x) = -\frac{1}{h} \nabla d(\cdot, \partial\Omega), \quad \text{for any } x \in \Omega \text{ s.t. } d(x, \partial\Omega) < h \leq h_0 \text{ (small enough)}.$$

In particular, for such x , we have $h\nu_h(x) = \nu(\pi(x))$, where $\pi(x)$ design the projection of x on the boundary $\partial\Omega$, and $\nu(y)$ represents the outward unitary normal to the boundary $\partial\Omega$ at y .

We denote by Sign^+ the maximal monotone graph given by

$$\text{Sign}^+(r) = \begin{cases} 1 & \text{for } r > 0 \\ [0, 1] & \text{for } r = 0 \\ 0 & \text{for } r < 0. \end{cases}$$

Moreover, we define Sign_0 and Sign_0^\pm , the discontinuous applications defined from \mathbb{R} to \mathbb{R} by

$$\text{Sign}_0(r) = \begin{cases} 1 & \text{for } r > 0 \\ 0 & \text{for } r = 0 \\ -1 & \text{for } r < 0, \end{cases} \quad \text{Sign}_0^+(r) = \begin{cases} 1 & \text{for } r > 0 \\ 0 & \text{for } r \leq 0 \end{cases} \quad \text{Sign}_0^-(r) = \begin{cases} 0 & \text{for } r \geq 0 \\ 1 & \text{for } r < 0 \end{cases}$$

Throughout the paper, we assume that u_0 and the velocity vector field V satisfy the following assumptions :

- $u_0 \in L^\infty(\Omega)$ satisfies $0 \leq |u_0| \leq 1$ a.e. in Ω .
- $V \in W_{loc}^{1,1}(\Omega)^N$, $\nabla \cdot V \in L^\infty(\Omega)$ and V satisfies (outward pointing velocity vector field condition on the boundary)

$$(1.2) \quad V \cdot \nu \geq 0 \quad \text{on } \partial\Omega.$$

Then, we denote by $H_D^1(\Omega)$ the Sobolev space

$$H_D^1(\Omega) = \left\{ u \in H^1(\Omega) : u = 0, \mathcal{L}^{N-1}\text{-a.e. in } \Gamma_D \right\}.$$

Remark 1. (1) *Since $V \in W_{loc}^{1,1}(\Omega)^N$ and $\nabla \cdot V \in L^\infty(\Omega)$, then $V \cdot \nu$ needs to be understood in a weak sense, like*

$$\int_{\Omega} V \cdot \nabla \xi \, dx + \int_{\Omega} \nabla \cdot V \, \xi \, dx \geq 0, \quad \text{for any } 0 \leq \xi \in \mathcal{D}(\bar{\Omega}).$$

One can see also the papers [20, 22] for more details and expressions in this direction. In particular, this condition implies that

$$(1.3) \quad \liminf_{h \rightarrow 0} \int_{D_h} \xi V(x) \cdot \nu_h(x) \, dx \geq 0, \quad \text{for any } 0 \leq \xi \in L^\infty(\Omega),$$

where, for any $0 < h \leq h_0$, D_h is a neighbour of $\partial\Omega$.

(2) *In some models of congested crowd motion, the vector field velocity may be given by*

$$(1.4) \quad V = -\nabla d(\cdot, \Gamma_D), \quad \text{in } \Omega.$$

This means that the pedestrian choose the euclidean geodesic trajectory to the escape door to get away from the environment Ω . See that in this case the velocity vector field is outward pointing on Γ_D as we need, but inward pointing on Γ_N . To fall into the scope of (1.2) one needs for instance to consider (1.4) outside a small neighbour of Γ_N and set V to be tangential in this neighbour. This example constitute a typical practical situation (among many others) for which all the results of this paper may be applied.

To begin with, we consider first the problem

$$(1.5) \quad \left\{ \begin{array}{l} \frac{\partial u}{\partial t} - \Delta p + \nabla \cdot (u V) = f \\ u \in \text{Sign}(p) \end{array} \right\} \quad \text{in } Q$$

$$\left\{ \begin{array}{l} p = 0 \\ (\nabla p + u V) \cdot \nu = 0 \\ u(0) = u_0 \end{array} \right\} \quad \begin{array}{l} \text{on } \Sigma_D := (0, T) \times \Gamma_D \\ \text{on } \Sigma_N := (0, T) \times \Gamma_N \\ \text{in } \Omega, \end{array}$$

where $f \in L^1(Q)$ is given.

Definition 1.1 (Notion of solution). *A couple (u, p) is said to be a weak solution of (1.5) if $(u, p) \in L^\infty(Q) \times L^2(0, T; H_D^1(\Omega))$, $u \in \text{Sign}(p)$ a.e. in Q and*

$$\frac{d}{dt} \int_{\Omega} u \xi + \int_{\Omega} (\nabla p - u V) \cdot \nabla \xi = \int_{\Omega} f \xi, \quad \text{in } \mathcal{D}'(0, T), \quad \forall \xi \in H_D^1(\Omega).$$

We'll say plainly that u is a solution of (1.5) if $u \in \mathcal{C}([0, T], L^1(\Omega))$, $u(0) = u_0$ and there exists $p \in L^2(0, T; H_D^1(\Omega))$ such that the couple (u, p) is a weak solution of (1.5).

Remark 2. *The assumptions we suppose on V look alike to be stronger regard to the literature on (linear) continuity equation. We think it is possible to extend our results for weak solution to more general V . But, in the presence of the second order term, the problem seems to be heavy and much more technical. Let us give here at least a short reminder on the uniqueness of weak solution for continuity equation, which could comprise some possible extensions (open questions !) for the Hele-Shaw flow with a linear Drift..*

- (1) *The analysis of the continuity equation (without a second order term) in the case when V has low regularity has drawn considerable attention : for an overview of some of the main contributions, we refer to the lecture notes by Ambrosio and Crippa [4] (see also [23] for more references on this subject). counter-example for the uniqueness when V is not enough regular (up to the boundary) may be found in [23]. Indeed, regardless the orientation of V at the boundary, uniqueness may be violated as soon as V enjoys BV regularity in every open set $\omega \subset\subset \Omega$, but not at the boundary of Ω (cf. [23] for more details and discussions on this subject).*
- (2) *When the vector filed is outward pointing as in (1.3), heuristically the solution would not be worst at/near the boundary. Indeed, at least in the smooth case the solution may simply "carried out" of the domain along the characteristics and, consequently, the behaviour of the solution*

inside the domain is not substantially affected by what happens close to the boundary. However, examples are given in [23] discussing that even if V is outward pointing at $\partial\Omega$, then uniqueness may be violated as soon as the BV regularity deteriorates at the boundary.

- (3) The case where the vector field V is inward pointing at the boundary, heuristically the solutions would be worst at/near the boundary. Indeed, the solution “enters” the domain Ω , although the characteristics are outward pointing at the boundary. This could explain at least partially the gain of technicality of this case in contrast of the outward pointing case.

2. L^1 -COMPARISON PRINCIPLE

In this section, we focus first on the uniqueness and L^1 -Comparison principle of weak solution. Our first main result is the following.

Theorem 2.1. *If (u_1, p_1) and (u_2, p_2) are two weak solutions of (1.5) associated with $f_1, f_2 \in L^1(Q)$ respectively, then there exists $\kappa \in L^\infty(Q)$, such that $\kappa \in \text{Sign}(u_1 - u_2)$ a.e. in Q and*

$$(2.1) \quad \frac{d}{dt} \int_{\Omega} (u_1 - u_2)^+ dx \leq \int_{\Omega} \kappa (f_1 - f_2) dx, \quad \text{in } \mathcal{D}'(0, T).$$

In particular, we have

$$\frac{d}{dt} \|u_1 - u_2\|_1 \leq \|f_1 - f_2\|_1, \quad \text{in } \mathcal{D}'(0, T).$$

Moreover, if $f_1 \leq f_2$, a.e. in Q , and u_1, u_2 are two corresponding solutions satisfying $u_1(0) \leq u_2(0)$ a.e. in Ω , then

$$u_1 \leq u_2, \quad \text{a.e. in } Q.$$

The main tool to prove this result is doubling and de-doubling variables. First we prove that a weak solution satisfies some kind of renormalized formulation. This enables us to use Kruzhkov device of doubling and de-doubling variables to prove (2.1).

Recall that, the merely transport equation case correspond to the situation where $p \equiv 0$. In this case, the renormalized formulation reads

$$\partial_t \beta(u) + V \cdot \nabla \beta(u) + u \nabla \cdot V \beta'(u) = f \beta'(u) \quad \text{in } \mathcal{D}'(Q),$$

for any $\beta \in C^1(\mathbb{R})$. From now on, for any $z \in L^1(\Omega)$, without abusing, the notation $V \cdot \nabla z$ needs to be understood in the sens of distribution as follows

$$(2.2) \quad V \cdot \nabla z = \nabla \cdot (z V) - z \nabla \cdot V.$$

Proposition 2.1. *If (u, p) is a weak solution of (1.5), then*

$$(2.3) \quad \begin{aligned} \partial_t \beta(u) - \Delta p^+ + V \cdot \nabla \beta(u) + u \nabla \cdot V(\beta'(u) \chi_{[p=0]} + \text{Sign}_0^+(p)) \\ \leq f(\beta'(u) \chi_{[p=0]} + \text{Sign}_0^+(p)) \quad \text{in } \mathcal{D}'(Q) \end{aligned}$$

and

$$(2.4) \quad \begin{aligned} \partial_t \beta(u) - \Delta p^- + V \cdot \nabla \beta(u) + u \nabla \cdot V(\beta'(u) \chi_{[p=0]} - \text{Sign}_0^-(p)) \\ \leq f(\beta'(u) \chi_{[p=0]} - \text{Sign}_0^-(p)) \quad \text{in } \mathcal{D}'(Q), \end{aligned}$$

for any $\beta \in C^1(\mathbb{R})$ such that $0 \leq \beta' \leq 1$.

Remark 3. *The formulations (2.3) and (2.4) describe some kind of renormalized formulation for the solution u . For the one phase Hele-Shaw problem this formulation reads simply as follows*

$$\begin{aligned} \partial_t \beta(u) - \Delta p + V \cdot \nabla \beta(u) + u \beta'(u) \nabla \cdot V \chi_{[p=0]} + \nabla \cdot V \chi_{[p \neq 0]} \\ \leq f(\beta'(u) \chi_{[p=0]} - \chi_{[p \neq 0]}) \quad \text{in } \mathcal{D}'(Q), \end{aligned}$$

for any $\beta \in C^1(\mathbb{R})$ such that $0 \leq \beta' \leq 1$.

We prove this results in two steps, we begin with the case where $\beta \equiv 0$. Then proceeding skillfully with the positive and negative part of p , and using a smoothing procedure with a commutator à la DiPerna-Lions (cf. [27], see also [3]), we prove that any weak solution u satisfies renormalized formulations (2.3) and (2.4).

Lemma 2.1. *If (u, p) is a weak solution of (1.5), then the renormalized formulations (2.3) and (2.4) are fulfilled with $\beta \equiv 0$ in $\mathcal{D}'((0, T) \times \bar{\Omega})$. That is*

$$(2.5) \quad -\Delta p^+ + (\nabla \cdot V - f) \text{Sign}_0^+(p) \leq 0 \quad \text{in } \mathcal{D}'((0, T) \times \bar{\Omega}),$$

$$(2.6) \quad -\Delta p^- + (\nabla \cdot V + f) \text{Sign}_0^-(p) \leq 0 \quad \text{in } \mathcal{D}'((0, T) \times \bar{\Omega}),$$

and

$$(2.7) \quad \partial_t u - \Delta p^+ + \nabla \cdot (u V) + \nabla \cdot V \text{Sign}_0^-(p) \leq f(1 - \text{Sign}_0^-(p)) \quad \text{in } \mathcal{D}'(Q),$$

Proof. We extend p onto $\mathbb{R} \times \Omega$ by 0 for any $t \notin (0, T)$. Then, for any $h > 0$ and nonnegative $\xi \in H^1(\Omega)$ and $\psi \in \mathcal{D}(\mathbb{R})$, let us consider

$$\Phi^h(t) = \xi \psi(t) \frac{1}{h} \int_t^{t+h} \mathcal{H}_\varepsilon^+(p(s)) ds,$$

where $\mathcal{H}_\varepsilon^+$ is given by

$$\mathcal{H}_\varepsilon^+(r) = \min\left(\frac{r^+}{\varepsilon}, 1\right), \quad \text{for any } r \in \mathbb{R},$$

for arbitrary $\varepsilon > 0$. It is clear that $\Phi_h \in W^{1,2}\left(0, T; H_D^1(\Omega)\right) \cap L^\infty(Q)$ is an admissible test function for the weak formulation, so that

$$(2.8) \quad - \iint_Q u \partial_t \Phi^h + \iint_Q (\nabla p - V u) \cdot \nabla \Phi^h = \iint_Q f \Phi^h.$$

See that

$$(2.9) \quad \iint_Q u \partial_t \Phi^h = \iint_Q u \partial_t \psi \frac{1}{h} \int_t^{t+h} \mathcal{H}_\varepsilon^+(p(s)) ds + \iint_Q u(t) \frac{\mathcal{H}_\varepsilon^+(p(t+h)) - \mathcal{H}_\varepsilon^+(p(t))}{h} \psi(t) \xi.$$

Moreover, using the fact that for a.e. $t \in (0, T)$, $-1 \leq u(t) \leq 1$, $\mathcal{H}_\varepsilon^+ \geq 0$ and $\mathcal{H}_\varepsilon^+(0) = 0$, we have $u(t, x) \mathcal{H}_\varepsilon^+(p(t, x)) = \mathcal{H}_\varepsilon^+(p(t, x))$ and $u(t, x) \mathcal{H}_\varepsilon^+(p(t+h, x)) \leq \mathcal{H}_\varepsilon^+(p(t+h, x))$ a.e. $(t, x) \in Q$. So,

$$\begin{aligned} \iint_Q u(t) \frac{\mathcal{H}_\varepsilon^+(p(t+h)) - \mathcal{H}_\varepsilon^+(p(t))}{h} \psi(t) \xi &\leq \iint_Q \frac{\mathcal{H}_\varepsilon^+(p(t+h)) - \mathcal{H}_\varepsilon^+(p(t))}{h} \psi(t) \xi \\ &\leq \iint_Q \frac{\psi(t-h) - \psi(t)}{h} \mathcal{H}_\varepsilon^+(p(t)) \xi \end{aligned}$$

This implies that

$$\limsup_{h \rightarrow 0} \iint_Q u(t) \frac{\mathcal{H}_\varepsilon^+(p(t+h)) - \mathcal{H}_\varepsilon^+(p(t))}{h} \psi(t) \xi \leq - \iint_Q \partial_t \psi \mathcal{H}_\varepsilon^+(p(t)) \xi,$$

and then, by letting $h \rightarrow 0$ in (2.8) and (2.9), we get

$$(2.13) \quad \iint_Q (\nabla p - V u) \cdot \nabla (\mathcal{H}_\varepsilon^+(p(t)) \xi) - \iint_Q f \mathcal{H}_\varepsilon^+(p(t)) \xi = \lim_{h \rightarrow 0} \iint_Q u \partial_t \Phi^h \leq 0.$$

On the other hand, using again the fact that $u \mathcal{H}_\varepsilon^+(p) = \mathcal{H}_\varepsilon^+(p)$, a.e. in Q , we have

$$\begin{aligned} \iint_Q (\nabla p - u V) \cdot \nabla (\mathcal{H}_\varepsilon^+(p) \xi) \psi &= \iint_Q |\nabla p|^2 \mathcal{H}_\varepsilon^+(p) \xi \psi + \iint_Q \mathcal{H}_\varepsilon^+(p) \nabla p \cdot \nabla \xi \psi \\ &\quad - \iint_Q V \cdot \nabla (\xi \mathcal{H}_\varepsilon^+(p)) \psi \\ &\leq 0. \end{aligned}$$

Thanks to (2.13), this implies that

$$\iint_Q |\nabla p|^2 \mathcal{H}_\varepsilon^+(p) \xi \psi - \iint_Q V \cdot \nabla (\xi \mathcal{H}_\varepsilon^+(p)) \leq \iint_Q f \mathcal{H}_\varepsilon^+(p(t)) \xi,$$

where we use also the fact that $|\nabla p|^2 \mathcal{H}_\varepsilon^+(p) \geq 0$. Now, see that using (1.2), we have

$$\iint_Q V \cdot \nabla (\xi \mathcal{H}_\varepsilon^+(p)) \psi \geq - \iint_Q \nabla \cdot V \xi \mathcal{H}_\varepsilon^+(p) \psi,$$

and then

$$\iint_Q |\nabla p|^2 \mathcal{H}_\varepsilon^+(p) \xi \psi + \iint_Q \nabla \cdot V \xi \mathcal{H}_\varepsilon^+(p) \psi \leq \iint_Q f \mathcal{H}_\varepsilon^+(p(t)) \xi.$$

Letting now $\varepsilon \rightarrow 0$, we get (2.5). As to (2.6), it follows by using the that $(-u, -p)$ is also a solution of (1.5) with f replaced by $-f$, and applying (2.5) to $(-u, -p)$. At last, recall that

$$(2.14) \quad \partial_t u - \Delta p + \nabla \cdot (u \nabla u) = f \quad \text{in } \mathcal{D}'(Q).$$

So, summing (2.6) restrained to $\mathcal{D}'(Q)$ and (2.14), we get (2.7). \square

Now, in order to prove the proposition by using (2.5) and (2.7), we prove the following technical lemma.

Lemma 2.2. *Let $u \in L^1_{loc}(Q)$, $F \in L^1_{loc}(Q)^N$ and $J_1 \in L^1_{loc}(Q)$ bu such that*

$$(2.15) \quad \partial_t u + V \cdot \nabla u - \nabla \cdot F \leq J_1 \quad \text{in } \mathcal{D}'(Q)$$

where $V \cdot \nabla u$ is taken in the sens $V \cdot \nabla u = \nabla \cdot (u V) - u \nabla \cdot V$, in $\mathcal{D}'(Q)$. If

$$(2.16) \quad -\nabla \cdot F \leq J_2 \quad \text{in } \mathcal{D}'(Q),$$

for some $J_2 \in L^1_{loc}(Q)$, then

$$(2.17) \quad \partial_t \beta(u) + V \cdot \nabla \beta(u) - \nabla \cdot F \leq J_1 \beta'(u) + J_2 (1 - \beta'(u)) \quad \text{in } \mathcal{D}'(\Omega),$$

for any $\beta \in C^1(\mathbb{R})$ such that $0 \leq \beta' \leq 1$.

Proof. We set $Q_\varepsilon := \{(t, x) \in Q : d((t, x), \partial Q) > \varepsilon\}$. Moreover, for any $z \in L^1(Q)$, we denote by z_ε the usual regularization of z by convolution given and denoted by

$$z_\varepsilon := z \star \rho_\varepsilon, \quad \text{in } Q_\varepsilon,$$

where ρ_ε is the usual mollifiers sequence defined here in $\mathbb{R} \times \mathbb{R}^N$. It is not difficult to see that (2.16) and (2.15) implies respectively

$$(2.18) \quad -\nabla \cdot F_\varepsilon \leq J_{2\varepsilon} \quad \text{in } Q_\varepsilon,$$

and

$$(2.19) \quad \partial_t u_\varepsilon + V \cdot \nabla u_\varepsilon - \nabla \cdot F_\varepsilon \leq J_{1\varepsilon} + C_\varepsilon \quad \text{in } Q_\varepsilon$$

and C_ε is the usual commutators given by

$$C_\varepsilon := V \cdot \nabla u_\varepsilon - (V \cdot \nabla u)_\varepsilon.$$

Using (2.2), here $(V \cdot \nabla u)_\varepsilon$ needs to be understood in the sense

$$(V \cdot \nabla u)_\varepsilon = (u V) \star \nabla \rho_\varepsilon - (u \nabla \cdot V) \star \rho_\varepsilon, \quad \text{in } Q_\varepsilon.$$

Multiplying (2.19) by $\beta'(u_\varepsilon)$ and (2.18) by $1 - \beta'(u_\varepsilon)$ and adding the resulting equations, we obtain

$$\beta'(u_\varepsilon) \partial_t u_\varepsilon + \beta'(u_\varepsilon) V \cdot \nabla u_\varepsilon - \nabla \cdot F_\varepsilon \leq C_\varepsilon \beta'(u_\varepsilon) + J_{1\varepsilon} \beta'(u_\varepsilon) + J_{2\varepsilon} (1 - \beta'(u_\varepsilon)) \quad \text{in } Q_\varepsilon$$

and then

$$(2.20) \quad \partial_t \beta(u_\varepsilon) + V \cdot \nabla \beta(u_\varepsilon) - \nabla \cdot F_\varepsilon \leq C_\varepsilon \beta'(u_\varepsilon) + J_{1\varepsilon} \beta'(u_\varepsilon) + J_{2\varepsilon} (1 - \beta'(u_\varepsilon)) \quad \text{in } Q_\varepsilon.$$

Since $V \in W_{loc}^{1,1}(\Omega)$ and $\nabla \cdot V \in L^\infty(\Omega)$, it is well known by now that, by taking a subsequence if necessary, the commutators converges to 0 in $L_{loc}^1(Q)$, as $\varepsilon \rightarrow 0$ (see for instance Ambrosio [3]). So, letting $\varepsilon \rightarrow 0$ in (2.20), we get (2.17). \square

Now, let us prove how to get the renormalized formulations (2.3) and (2.4).

Proof of Proposition 2.1. Thanks to Lemma 2.1, by using the fact that

$$\nabla \cdot V \text{Sign}_0^-(p) = -u \nabla \cdot V \text{Sign}_0^-(p) \quad \text{and} \quad \nabla \cdot V \text{Sign}_0^+(p) = u \nabla \cdot V \text{Sign}_0^+(p),$$

we see that (2.15) and (2.16) are fulfilled with

$$F := \nabla p^+, \quad J_1 := (f - u \nabla \cdot V)(1 - \text{Sign}_0^-(p))$$

and

$$J_2 := (f - u \nabla \cdot V) \text{Sign}_0^+(p).$$

Thanks to Lemma 2.1, for any $\beta \in \mathcal{C}^1(\mathbb{R})$ such that $0 \leq \beta' \leq 1$, we deduce that

$$\begin{aligned} & \partial_t \beta(u) - \Delta p^+ + V \cdot \nabla \beta(u) + (u \nabla \cdot V - f)(1 - \text{Sign}_0^-(p)) \beta'(u) \\ & + (f - u \nabla \cdot V) \text{Sign}_0^+(p) (\beta'(u) - 1) \leq 0 \quad \text{in } \mathcal{D}'(Q), \end{aligned}$$

Using again the fact that $\nabla \cdot V \text{Sign}_0^+(p) = u \nabla \cdot V \text{Sign}_0^+(p)$ this implies that

$$\begin{aligned} & \partial_t \beta(u) - \Delta p^+ + V \cdot \nabla \beta(u) + u \nabla \cdot V (1 - \text{Sign}_0^-(p)) \beta'(u) + u \nabla \cdot V \text{Sign}_0^+(p) (1 - \beta'(u)) \\ & \leq f(\text{Sign}_0^+(p)(1 - \beta'(u)) + (1 - \text{Sign}_0^-(p)) \beta'(u)) \quad \text{in } \mathcal{D}'(Q). \end{aligned}$$

That is

$$\begin{aligned} \partial_t \beta(u) - \Delta p^+ + V \cdot \nabla \beta(u) + u \nabla \cdot V(\beta'(u)(1 - \text{Sign}_0^-(p) - \text{Sign}_0^+(p)) + \text{Sign}_0^+(p)) \\ \leq f(\beta'(u)(1 - \text{Sign}_0^-(p) - \text{Sign}_0^+(p)) + \text{Sign}_0^+(p)) \quad \text{in } \mathcal{D}'(Q), \end{aligned}$$

and then

$$\begin{aligned} \partial_t \beta(u) - \Delta p^+ + V \cdot \nabla \beta(u) + u \nabla \cdot V(\beta'(u)\chi_{[p=0]} + \text{Sign}_0^+(p)) \\ \leq f(\beta'(u)\chi_{[p=0]} + \text{Sign}_0^+(p)) \quad \text{in } \mathcal{D}'(Q). \end{aligned}$$

Thus (2.3). At last, using the fact that the couple $(-u, -p)$ is a solution of (1.5) with f replaced by $-f$, and applying (2.3) to $(-u, -p)$, we deduce (2.4). \square

See here that the attendance of a nonlinear second order term creates an obstruction to use standard approaches for the uniqueness of weak solutions based effectively on the linearity of the first order term. In order to prove uniqueness for our case, we proceed by doubling and de-doubling variable techniques. To this aim, we'll use the fact that the renormalized formulation implies some kind of local entropic inequalities. This is the aim of the following Lemma.

Lemma 2.3. *If (u, p) is a weak solution of (1.5), then, for any $-1 \leq k \leq 1$, we have*

$$\begin{aligned} (2.21) \quad \partial_t(u - k)^+ - \Delta p^+ + \nabla \cdot ((u - k)^+ V) + k \nabla \cdot V \text{Sign}_0^+(u - k) \\ \leq f \text{Sign}_0^+(u - k) \quad \text{in } \mathcal{D}'(Q), \end{aligned}$$

and

$$\begin{aligned} (2.22) \quad \partial_t(u - k)^- - \Delta p^- + \nabla \cdot ((k - u)^+ V) - s \nabla \cdot V \text{Sign}_0^+(k - u) \\ \leq (u - f) \text{Sign}_0^+(k - u) \quad \text{in } \mathcal{D}'(Q) \end{aligned}$$

Remark 4. *See here that our entropic inequalities (2.21) and (2.22) are local and do not proceed up to the boundary like in [19]. Indeed, in contrast of Carillo's approach, we are intending here to handle the boundary condition separately. In particular, as we'll see this would reduce the technicality of de-doubling variables process (see Proposition 2.2).*

Proof of Lemma 2.3. See first that, since $|u| \leq 1$ a.e. in Q , (2.21) is trivially fulfilled when $k = 1$. Now, for any $-1 \leq k < 1$, we consider

$$\beta_\epsilon(r) = \tilde{\mathcal{H}}_\epsilon(r - k), \quad \text{for any } r \in \mathbb{R},$$

where

$$\tilde{\mathcal{H}}_\epsilon(r) = \begin{cases} \frac{1}{2\epsilon} r^{+2} & \text{for } r \leq \epsilon \\ r - \frac{\epsilon}{2} & \text{elsewhere.} \end{cases}$$

In this case, $\beta'_\epsilon(1) = 0$, for any $0 < \epsilon \leq \epsilon_0$ (small enough), and we have

$$\beta'_\epsilon(u) = \mathcal{H}_\epsilon^+(u - k) \rightarrow \text{Sign}_0^+(u - k), \quad \text{as } \epsilon \rightarrow 0.$$

Since

$$\text{Sign}_0^+(u - k) \chi_{[p=0]} + \text{Sign}_0^+(p) = \text{Sign}_0^+(u - k),$$

letting $\varepsilon \rightarrow 0$, in (2.3) where we replace β by $\tilde{\mathcal{H}}_\varepsilon$, we get

$$\begin{aligned} \partial_t(u-k)^+ - \Delta p^+ + V \cdot \nabla(u-k)^+ + u \nabla \cdot V \text{Sign}_0^+(u-k) \\ \leq f \text{Sign}_0^+(u-k) \quad \text{in } \mathcal{D}'(Q), \end{aligned}$$

which implies (2.21).

For the second part of the Lemma, we use again the fact that see that $(\tilde{u} := -u, \tilde{p} := -p)$ is a weak solution of (1.5) with f replaced by $\tilde{f} := -f$. So, using (2.21) for (\tilde{u}, \tilde{p}) with \tilde{f} , for any $k \leq 1$, we have

$$\begin{aligned} -\Delta \tilde{p}^+ + \nabla \cdot ((\tilde{u}-k)^+ V) + k \nabla \cdot V \text{Sign}_0^+(\tilde{u}-k) \\ \leq (\tilde{f} - \tilde{u}) \text{Sign}_0^+(\tilde{u}-k) \quad \text{in } \mathcal{D}'(Q) \end{aligned}$$

Which implies that, for any $-1 \leq s$, we have

$$\begin{aligned} \partial_t(s-u)^- - \Delta p^- + \nabla \cdot ((s-u)^+ V) - s \nabla \cdot V \text{Sign}_0^+(s-u) \\ \leq (u-f) \text{Sign}_0^+(s-u) \quad \text{in } \mathcal{D}'(Q). \end{aligned}$$

Thus the result of the lemma. \square

Now, we are able to prove some kind of "local" Kato's inequality which mold our L^1 -approach for the uniqueness.

Proposition 2.2 (Kato's inequality). *If (u_1, p_1) and (u_2, p_2) are two couples of $L^\infty(Q) \times L^2(0, T; H_D^1(\Omega))$ satisfying (2.21) and (2.22) corresponding to $f_1 \in L^1(Q)$ and $f_2 \in L^1(Q)$ respectively, then there exists $\kappa \in L^\infty(Q)$, such that $\kappa \in \text{Sign}^+(u_1 - u_2)$ a.e. in Q and*

$$(2.23) \quad \partial_t(u_1 - u_2)^+ - \Delta(p_1^+ + p_2^-) + \nabla \cdot ((u_1 - u_2)^+ V) \leq \kappa(f_1 - f_2) \text{ in } \mathcal{D}'(Q).$$

Proof. The proof of this lemma is based on a the usual doubling and de-doubling variable techniques. Let us give here briefly the arguments. To double the variables, we use the fact that (u_1, p_1) satisfies (2.21) with $k = u_1(s, y)$ we have

$$\begin{aligned} \frac{d}{dt} \int (u_1(t, x) - u_2(s, y))^+ \zeta + \int (\nabla_x p_1^+(t, x) - (u_1(t, x) - u_2(s, y))^+ V(x) \cdot \nabla_x \zeta \\ + \int_\Omega \nabla_x \cdot V u_2(s, y) \zeta \text{Sign}_0^+(u_1(t, x) - u_2(s, y)) \leq \int f_1(t, x) \text{Sign}_0^+(u_1(t, x) - u_2(s, y)) \zeta, \end{aligned}$$

for any $0 \leq \eta \in \mathcal{D}(\Omega \times \Omega)$. See that $\int \nabla_y p_2^-(s, y) \cdot \nabla_x \zeta dx = 0$, so that

$$\begin{aligned} \frac{d}{dt} \int (u_1(t, x) - u_2(s, y))^+ \zeta + \int (\nabla_x p_1^+(t, x) + \nabla_y p_2^-(t, x)) \cdot \nabla_x \zeta - \int (u_1(t, x) - u_2(s, y))^+ V(x) \cdot \nabla_x \zeta \\ + \int \nabla_x \cdot V u_2(s, y) \zeta \text{Sign}_0^+(u_1(t, x) - u_2(s, y)) \leq \int f_1(t, x) \text{Sign}_0^+(u_1(t, x) - u_2(s, y)) \zeta. \end{aligned}$$

Denoting by

$$u(t, s, x, y) = u_1(t, x) - u_2(s, y), \quad \text{and} \quad p(t, s, x, y) = p_1^+(t, x) + p_2^-(s, y),$$

and integrating with respect to y , we obtain

$$\begin{aligned} & \frac{d}{dt} \int u(t, s, x, y)^+ \zeta + \iint (\nabla_x + \nabla_y) p(t, s, x, y) \cdot \nabla_x \zeta - \iint u(t, s, x, y)^+ V(x) \cdot \nabla_x \zeta \\ & + \iint \nabla_x \cdot V u_2(s, y) \zeta \text{Sign}_0^+ u(t, s, x, y) \leq \iint f_1(t, x) \text{Sign}_0^+ u(t, s, x, y) \zeta. \end{aligned}$$

On the other hand, using the fact that (u_2, p_2) satisfies (2.22) with $k = u_1(t, x)$, we have

$$\begin{aligned} & \frac{d}{ds} \int u(t, s, x, y)^+ \zeta + \int (\nabla_y p_2^-(s, y) - u(t, s, x, y)^+ V(y) \cdot \nabla_y \zeta \\ & - \int_{\Omega} \nabla_y \cdot V u_1(t, x) \zeta \text{Sign}_0^+(u(t, s, x, y))) \leq - \int f_2(s, y) \text{Sign}_0^+(u(t, s, x, y)) \zeta. \end{aligned}$$

Working in the same way, we get

$$\begin{aligned} & \frac{d}{ds} \int u(t, s, x, y)^+ \zeta + \iint (\nabla_x + \nabla_y) p(t, s, x, y) \cdot \nabla_y \zeta - \iint u(t, s, x, y)^+ V(y) \cdot \nabla_y \zeta \\ & - \iint \nabla_y \cdot V(y) u_1(t, x) \zeta \text{Sign}_0^+(u(t, s, x, y)) \leq - \iint f_2(s, y) \text{Sign}_0^+(u(t, s, x, y)) \zeta. \end{aligned}$$

Adding both inequalities, we obtain

$$\begin{aligned} (2.24) \quad & \left(\frac{d}{dt} + \frac{d}{ds} \right) \iint u(t, s, x, y)^+ \zeta + \iint (\nabla_x + \nabla_y) p(t, s, x, y) \cdot (\nabla_x + \nabla_y) \zeta \\ & - \iint u(t, s, x, y)^+ (V(x) \cdot \nabla_x \zeta + V(y) \cdot \nabla_y \zeta) \\ & + \iint (\nabla_x \cdot V(x) u_2(s, y) - \nabla_y \cdot V(y) u_1(t, x)) \zeta \text{Sign}_0^+(u(t, s, x, y)) \\ & \leq \iint (f_1(t, x) - f_2(s, y)) \text{Sign}_0^+(u(t, s, x, y)) \zeta. \end{aligned}$$

Now, we can de-double the variables t and s , as well as x and y , by taking as usual the sequence of test functions

$$\psi_{\varepsilon}(t, s) = \psi \left(\frac{t+s}{2} \right) \rho_{\varepsilon} \left(\frac{t-s}{2} \right) \text{ and } \zeta_{\lambda}(x, y) = \xi \left(\frac{x+y}{2} \right) \delta_{\lambda} \left(\frac{x-y}{2} \right),$$

for any $t, s \in (0, T)$ and $x, y \in \Omega$. Here $\xi \in \mathcal{D}(\Omega)$, $\psi \in \mathcal{D}(0, T)$, ρ_{ε} and δ_{λ} are sequences of usual mollifiers in \mathbb{R} and \mathbb{R}^N respectively. See that

$$\left(\frac{d}{dt} + \frac{d}{ds} \right) \psi_{\varepsilon}(t, s) = \rho_{\varepsilon} \left(\frac{t-s}{2} \right) \dot{\psi} \left(\frac{t+s}{2} \right)$$

and

$$(\nabla_x + \nabla_y) \zeta_{\lambda}(x, y) = \delta_{\lambda} \left(\frac{x-y}{2} \right) \nabla \xi \left(\frac{x+y}{2} \right)$$

Moreover, for any $h \in L^1((0, T)^2 \times \Omega^2)$ and $\Phi \in L^1((0, T)^2 \times \Omega^2)^N$, it is not difficult to prove that

- $\lim_{\lambda \rightarrow 0} \lim_{\varepsilon \rightarrow 0} \int_0^T \int_0^T \iint_{\Omega} \iint_{\Omega} h(t, s, x, y) \zeta_{\lambda}(x, y) \rho_{\varepsilon}(t, s) = \int_0^T \int_{\Omega} h(t, t, x, x) \xi(x) \psi(t).$
- $\lim_{\lambda \rightarrow 0} \lim_{\varepsilon \rightarrow 0} \int_0^T \int_0^T \iint_{\Omega} \iint_{\Omega} h(t, s, x, y) \zeta_{\lambda}(x, y) \left(\frac{d}{dt} + \frac{d}{ds} \right) \rho_{\varepsilon}(t, s) = \int_0^T \int_{\Omega} h(t, t, x, x) \xi(x) \dot{\psi}(t).$
- $\lim_{\lambda \rightarrow 0} \lim_{\varepsilon \rightarrow 0} \int_0^T \int_0^T \iint_{\Omega} \iint_{\Omega} \Phi(t, s, x, y) \cdot (\nabla_x + \nabla_y) \zeta_{\lambda}(x, y) \rho_{\varepsilon}(t, s) = \int_0^T \int_{\Omega} \Phi(t, t, x, x) \cdot \nabla \xi(x) \psi(t) dt dx.$

So replacing ζ in (2.24) by ζ_λ , testing with ψ_ε and, letting $\varepsilon \rightarrow 0$ and $\lambda \rightarrow 0$, we get

$$\begin{aligned} & - \int_0^T \int_\Omega \left\{ (u_1 - u_2)^+ \xi \dot{\psi} + \nabla(p_1^+ + p_2^-) \cdot \nabla \xi \psi - (u_1 - u_2)^+ V \cdot \nabla \xi \psi + \nabla \cdot V (u_1 - u_2)^+ \xi \psi \right\} \\ & \leq \int_0^T \int_\Omega \kappa(x)(f_1 - f_2) \xi \psi + \lim_{\lambda \rightarrow 0} \int_0^T \int_\Omega \int_\Omega u(t, t, x, y)^+ (V(x) - V(y)) \cdot \nabla_y \zeta \psi, \end{aligned}$$

where $\kappa \in L^\infty(Q)$ is such that $\kappa \in \text{Sign}^+(u_1 - u_2)$ a.e. in Q . Notice here that the emergence of κ follows from the L^1 -compactness of the term u and the L^∞ -weak*-compactness of $\text{Sign}_0^+(u)$, as well as the monotonicity of the graph Sign^+ . Thus

$$(2.25) \quad \begin{aligned} & \frac{d}{dt} \int (u_1 - u_2)^+ \xi + \int \nabla(p_1^+ + p_2^-) \cdot \nabla \xi - \int (u_1 - u_2)^+ V \cdot \nabla \xi + \int \nabla \cdot V (u_1 - u_2)^+ \xi \\ & \leq \int \kappa(x)(f_1 - f_2) \xi + \lim_{\lambda \rightarrow 0} \int_\Omega \int_\Omega u(t, t, x, y)^+ (V(x) - V(y)) \cdot \nabla_y \zeta. \end{aligned}$$

To de-double the variables x and y in the terms corresponding to the vector field V , we use moreover the technical result of Lemma 6.8 (cf. Appendix) which is more or less well known. We put back its proof in the Appendix. This implies that

$$\begin{aligned} & \frac{d}{dt} \int (u_1(t, x) - u_2(t, x))^+ \xi dx + \int \nabla(p_1^+ + p_2^-) \cdot \nabla \xi dx \\ & - \int (u_1 - u_2)^+ V \cdot \nabla \xi dx \leq \int \kappa(f_1 - f_2) \xi dx. \end{aligned}$$

Thus the result of the proposition. \square

The aim now is to proceed with the sequence of test function ξ_h given by (1.1) in Kato's inequality and let $h \rightarrow 0$, to cover (2.1).

Lemma 2.4. *Under the assumption (1.3), if (u, p) is a weak solution of (1.5), then for any $\psi \in D(0, T)$ such that $\psi \geq 0$, we have*

$$\liminf_{h \rightarrow 0} \int_0^T \int_\Omega \nabla p^\pm \cdot \nabla \xi_h \psi(t) dt dx \geq 0.$$

Proof : Using (2.5), we see that, for any $0 \leq \psi \in D(0, T)$ we have

$$\begin{aligned} \int_0^T \int_\Omega \nabla p^+ \cdot \nabla \xi_h \psi dt dx & = - \int_0^T \int_\Omega \nabla p^+ \cdot \nabla (1 - \xi_h) \psi dt dx \\ & \geq \int_0^T \int_\Omega (f - \nabla \cdot V)(1 - \xi_h) \text{Sign}_0^+(p) \psi dt dx \end{aligned}$$

Then, letting $h \rightarrow 0$ and using the fact that $\xi_h \rightarrow 1$ in $L^\infty(\Omega)$ -weak*, we deduce that $\liminf_{h \rightarrow 0} \int_0^T \int_\Omega \nabla p^+(t) \cdot \nabla \xi_h \psi dt dx \geq 0$. The proof for p^- follows in a similar way by using (2.6). \square

Lemma 2.5. *If (u_1, p_1) and (u_2, p_2) are two couples of $L^\infty(Q) \times L^2(0, T; H_D^1(\Omega))$ satisfying (2.21) and (2.22) corresponding to $f_1 \in L^1(Q)$ and $f_2 \in L^1(Q)$ respectively, then there exists $\kappa \in L^\infty(Q)$, such that $\kappa \in \text{Sign}^+(u_1 - u_2)$ a.e. in Q and (2.1) is fulfilled.*

Proof. See that

$$\begin{aligned} \frac{d}{dt} \int_{\Omega} (u_1 - u_2)^+ dx - \int \kappa(f_1 - f_2) &= \lim_{h \rightarrow 0} \frac{d}{dt} \int_{\Omega} (u_1 - u_2)^+ \xi_h dx - \int \kappa(f_1 - f_2) \xi_h dx \\ &=: I(h). \end{aligned}$$

Taking ξ_h as a test function in (2.23), we have

$$I(h) \leq - \int (\nabla(p_1^+ + p_2^-) - (u_1 - u_2)^+ V) \cdot \nabla \xi_h dx.$$

Using Lemma 2.4, this implies that

$$\begin{aligned} \lim_{h \rightarrow 0} I(h) &\leq - \lim_{h \rightarrow 0} \int (u_1 - u_2)^+ V \cdot \nu(x) dx \\ &\leq 0 \end{aligned}$$

where we use the outgoing velocity vector field assumption (1.3). Thus (2.1). \square

Proof of Theorem 2.1. It clear that the first part follows by Lemma 2.5. The rest of the theorem is a straightforward consequence of (2.1). \square

3. EXISTENCE FOR THE EVOLUTION PROBLEM

Our main result here concerns existence of a solution.

Theorem 3.2. *For any $f \in L^2(Q)$ and $u_0 \in L^p \infty(\Omega)$ be such that $|u_0| \leq 1$ a.e. in Ω , the problem (1.5) has a unique weak solution (u, p) such that $u \in \mathcal{C}([0, T], L^1(\Omega))$ and $u(0) = u_0$. Moreover, the solution satisfies all the properties of Theorem 2.1.*

Remark 5. *Let $f \in L^2(Q)$, $u_0 \in L^\infty(\Omega)$ be such that $|u_0| \leq 1$ a.e. in Ω , and u be the unique weak solution of the problem (1.5). Using Theorem 3.2 and Theorem 2.1, we have*

- (1) *if $u_0 \geq 0$ and $f \geq 0$, then (u, p) is the unique weak solution of the one phase problem*

$$\left\{ \begin{array}{l} \frac{\partial u}{\partial t} - \Delta p + \nabla \cdot (u V) = f \\ 0 \leq u \leq 1, p \geq 0, p(u - 1) = 0 \end{array} \right\} \quad \begin{array}{l} \text{in } Q \\ \text{on } \Sigma_D \\ \text{on } \Sigma_N \\ \text{in } \Omega, \end{array}$$

in the sense of Definition 1.1.

- (2) *For each $n = 1, 2, \dots$, let $f_n \in L^2(Q)$, $u_{0n} \in L^\infty(\Omega)$ be such that $|u_{0n}| \leq 1$ a.e. in Ω , and u_n be the unique weak solution of the corresponding problem (1.5). If $u_{0n} \rightarrow u_0$ in $L^1(\Omega)$ and $f_n \rightarrow f$ in $L^1(Q)$, then $u_n \rightarrow u$, in $\mathcal{C}([0, T], L^1(\Omega))$ and $p_n \rightarrow p$, in $L^2(0, T; H_D^1(\Omega))$ – weak.*

To study the existence, we process in the framework of nonlinear semigroup theory in $L^1(\Omega)$. In connection with the Euler implicit discretization scheme of the evolution problem (1.5), we consider

the following stationary problem :

$$(3.1) \quad \left\{ \begin{array}{l} u - \Delta p + \nabla \cdot (u V) = f \\ u \in \text{Sign}(p) \end{array} \right\} \quad \text{in } \Omega$$

$$\left\{ \begin{array}{l} p = 0 \\ (\nabla p + u V) \cdot \eta = 0 \end{array} \right. \quad \begin{array}{l} \text{on } \Gamma_D \\ \text{on } \Gamma_N, \end{array}$$

where $f \in L^2(\Omega)$ and $\lambda > 0$ are given. A couple $(u, p) \in L^\infty(\Omega) \times H_D^1(\Omega)$ satisfying $u \in \text{Sign}(p)$ a.e. in Ω , is said to be a weak solution of (3.1) if

$$\int_{\Omega} u \xi + \int_{\Omega} \nabla p \cdot \nabla \xi - \int_{\Omega} u V \cdot \nabla \xi = \int_{\Omega} f \xi,$$

for any test function $\xi \in H_D^1(\Omega)$. As for the evolution problem, we'll say simply that u is a solution of (3.1) if there exists p such that the couple (u, p) is a weak solution of (3.1).

As a consequence of Theorem 2.1, we can deduce the following result.

Corollary 3.1. *If u_1 and u_2 are two solutions of (3.1) associated with $f_1, f_2 \in L^1(\Omega)$, respectively, then*

$$\int (u_1 - u_2)^+ \leq \int (f_1 - f_2)^+.$$

In particular, we have

$$\|u_1 - u_2\|_1 \leq \|f_1 - f_2\|_1$$

and, if $f_1 \leq f_2$, a.e. in Ω , then

$$u_1 \leq u_2, \quad \text{a.e. in } \Omega.$$

Proof. This is a simple consequence of the fact that if (u, p) is a solution of (3.1), then it is a solution of (1.5) with f replaced by $f - u$. \square

For the existence, we consider the regularized problem

$$(3.2) \quad \left\{ \begin{array}{l} u - \Delta p + \nabla \cdot (u V) = f \\ u = \mathcal{H}_\varepsilon(p) \end{array} \right\} \quad \text{in } \Omega$$

$$\left\{ \begin{array}{l} p = 0 \\ (\nabla p + u V) \cdot \eta = 0 \end{array} \right. \quad \begin{array}{l} \text{on } \Gamma_D \\ \text{on } \Gamma_N, \end{array}$$

for an arbitrary for any $\varepsilon > 0$. See that for any $\varepsilon > 0$, $|\mathcal{H}_\varepsilon| \leq 1$, \mathcal{H}_ε is Lipschitz continuous and satisfies

$$(I + \mathcal{H}_\varepsilon)^{-1}(r) \rightarrow (I + \text{Sign})^{-1}(r), \quad \text{as } \varepsilon \rightarrow 0, \quad \text{for any } r \in \mathbb{R}.$$

That is \mathcal{H}_ε converges to Sign in the sense of resolvent, which is equivalent to the convergence in the sens of graph (cf. [18]).

Proposition 3.3. *For any $f \in L^2(\Omega)$ and $\varepsilon > 0$, the problem (3.2) has a weak solution $(u_\varepsilon, p_\varepsilon)$ in the sense that $p_\varepsilon \in H_D^1(\Omega)$, $u_\varepsilon = \mathcal{H}_\varepsilon(p_\varepsilon)$ a.e. in Ω and*

$$(3.3) \quad \int u_\varepsilon \xi \, dx + \int \nabla p_\varepsilon \cdot \nabla \xi \, dx - \int u_\varepsilon V \cdot \nabla \xi \, dx = \int f \xi \, dx, \quad \text{for any } \xi \in H_D^1(\Omega).$$

Moreover, as $\varepsilon \rightarrow 0$, we have

$$(3.4) \quad \mathcal{H}_\varepsilon(p_\varepsilon) \rightarrow u \quad \text{in } L^\infty(\Omega) - \text{weak}^*,$$

$$p_\varepsilon \rightarrow p \quad \text{in } H_D^1(\Omega) - \text{weak}$$

and (u, p) is the weak solution of (3.1).

Proof. The existence of a solution for (3.2) is standard. For completeness we give the arguments. Let us denote by $H_D^1(\Omega)^*$ the topological dual space of $H_D^1(\Omega)$ and $\langle \cdot, \cdot \rangle$ the associate dual bracket. See that the operator $A_\varepsilon : H_D^1(\Omega) \rightarrow H_D^1(\Omega)^*$, given by

$$\langle A_\varepsilon p, \xi \rangle = \int \mathcal{H}_\varepsilon(p) \xi \, dx + \int \nabla p \cdot \nabla \xi \, dx - \int \mathcal{H}_\varepsilon(p) V \cdot \nabla \xi \, dx$$

is a bounded weakly continuous operator. Moreover, A_ε is coercive. Indeed, for any $u \in H_D^1(\Omega)$, we have

$$\begin{aligned} \langle A_\varepsilon p, p \rangle &= \int \mathcal{H}_\varepsilon(p) p \, dx + \int |\nabla p|^2 \, dx - \int \mathcal{H}_\varepsilon(p) V \cdot \nabla p \, dx \\ &\geq \int |\nabla p|^2 \, dx - \int |V \cdot \nabla p| \, dx \\ &\geq \frac{1}{2} \int |\nabla p|^2 \, dx - \frac{1}{2} \int |V|^2 \, dx, \end{aligned}$$

where we use Young inequality. So, for any $f \in H_D^1(\Omega)^*$ the problem $A_\varepsilon p = f$ has a solution $p_\varepsilon \in H_D^1(\Omega)$. To let $\varepsilon \rightarrow 0$, we see that

$$(3.9) \quad \int |\nabla p_\varepsilon|^2 \, dx \leq C(N, \Omega) \left(\int |V|^2 \, dx + \int |f|^2 \, dx \right).$$

Indeed, taking p as a test function we have

$$\int u p \, dx + \int |\nabla p|^2 \, dx = \int u V \cdot \nabla p \, dx + \int f p \, dx.$$

See that using Young inequality we have

$$\int u V \cdot \nabla p \, dx \leq \frac{3}{4} \int |V|^2 \, dx + \frac{1}{3} \int |\nabla p|^2 \, dx$$

and, by combining Pincaré inequality with Young inequality we have

$$\int f p \, dx \leq C(N, \Omega) \int |f|^2 \, dx + \frac{1}{3} \int |\nabla p|^2 \, dx.$$

Using the fact that $u p \geq 0$, we deduce (3.9). Now, it is clear that the sequences p_ε and $u_\varepsilon = \mathcal{H}_\varepsilon(p_\varepsilon)$ are bounded respectively in $H_D^1(\Omega)$ and in $L^\infty(\Omega)$. So, there exists a subsequence that we denote again by p_ε such that (3.4) and (3.12) are fulfilled. In particular, this implies that $u \in \text{Sign}(p)$, a.e. and letting $\varepsilon \rightarrow 0$ in (3.3), we obtain that (u, p) is a weak solution of (3.1). \square

To prove the existence, of a weak solution to (1.5), we fix $f \in L^2(Q)$, and for an arbitrary $0 < \varepsilon \leq \varepsilon_0$, and $n \in \mathbb{N}^*$ be such that $(n+1)\varepsilon = T$, we consider the sequence of (u_i, p_i) given by the the ε -Euler implicit schema associated with (1.5) :

$$(3.10) \quad \left\{ \begin{array}{ll} \left. \begin{array}{l} u_{i+1} - \varepsilon \Delta p_{i+1} + \varepsilon \nabla \cdot (u_{i+1} V) = u_i + \varepsilon f_i \\ u_{i+1} \in \text{Sign}(p_{i+1}) \end{array} \right\} & \text{in } \Omega \\ p_{i+1} = 0 & \text{on } \Gamma_D \\ (\nabla p_{i+1} + u_{i+1} V) \cdot \eta = 0 & \text{on } \Gamma_N, \end{array} \right. \quad i = 0, 1, \dots, n-1,$$

where, for each $i = 0, \dots, n-1$, f_i is given by

$$f_i = \frac{1}{\varepsilon} \int_{i\varepsilon}^{(i+1)\varepsilon} f(s) ds, \quad \text{a.e. in } \Omega.$$

Now, for a given ε -time discretization $0 = t_0 < t_1 < t_2 < \dots < t_n < t_{n+1} = T$, satisfying $t_{i+1} - t_i \leq \varepsilon$, we define the ε -approximate solution by

$$u_\varepsilon := \sum_{i=0}^n u_i \chi_{[t_i, t_{i+1})}, \quad \text{and} \quad p_\varepsilon := \sum_{i=1}^n p_i \chi_{[t_i, t_{i+1})}.$$

Thanks to Proposition 3.3 and the general theory of evolution problem governed by accretive operator (see for instance [12, 11]), we define the operator \mathcal{A} in $L^1(\Omega)$, by $\mu \in \mathcal{A}(z)$ if and only if $\mu, z \in L^1(\Omega)$ and z is a solution of the problem

$$\left\{ \begin{array}{ll} \left. \begin{array}{l} -\Delta p + \nabla \cdot (z V) = \mu \\ z \in \text{Sign}(p) \end{array} \right\} & \text{in } \Omega \\ p = 0 & \text{on } \Gamma_D \\ (\nabla p + u V) \cdot \eta = 0 & \text{on } \Gamma_N, \end{array} \right.$$

in the sens that $z \in L^\infty(\Omega)$, there exists $p \in H_D^1(\Omega)$ satisfying $z \in \text{Sign}(p)$ a.e. in Ω and

$$\int_{\Omega} \nabla p \cdot \nabla \xi - \int_{\Omega} z V \cdot \nabla \xi = \int_{\Omega} \mu \xi, \quad \forall \xi \in H_D^1(\Omega).$$

As a consequence of Corollary 3.1, we know that the operator \mathcal{A} is accretive in $L^1(\Omega)$. Moreover, since $R(I + \lambda \mathcal{A}) \supseteq L^2(\Omega)$, its closure $\overline{\mathcal{A}}$, in $L^1(\Omega)$, is m-accretive in $L^1(\Omega)$; i.e. $R(I + \lambda \overline{\mathcal{A}}) = L^1(\Omega)$, for any $\lambda > 0$. Furthermore, thanks to the general theory of nonlinear semigroup governed by accretive operator, we know that, as $\varepsilon \rightarrow 0$,

$$(3.11) \quad u_\varepsilon \rightarrow u, \quad \text{in } \mathcal{C}([0, T], L^1(\Omega)),$$

and u is the so called "Mild solution" of the evolution problem

$$\left\{ \begin{array}{l} u_t + \mathcal{A}u \ni f \quad \text{in } (0, T) \\ u(0) = u_0. \end{array} \right.$$

To accomplish the proof of existence for the problem (1.5), we prove that the mild solution u is in fact the solution of (1.5). To this aim, we use the limit of the sequence p_ε given by the ε -approximate solution.

Lemma 3.6. *As $\varepsilon \rightarrow 0$,*

$$(3.12) \quad p_\varepsilon \rightarrow p, \quad \text{in } L^2(0, T; H_D^1(\Omega))$$

and (u, p) yields a weak solution of (1.5).

Proof. Thanks to Proposition 3, the sequence $(u_i, p_i)_{i=1, \dots, n}$ given by (3.10) is well defined in $L^\infty(\Omega) \times H_D^1(\Omega)$ and satisfies $u_i \in \text{Sign}(p_i)$ and

$$(3.13) \quad \int_{\Omega} u_{i+1} \xi + \varepsilon \int_{\Omega} \nabla p_{i+1} \cdot \nabla \xi - \varepsilon \int_{\Omega} u_{i+1} V \cdot \nabla \xi = \varepsilon \int_{\Omega} f_i \xi.$$

Taking p_{i+1} as a test function in (3.13) and using the fact that $(u_{i+1} - u_i)p_{i+1} \geq 0$, it is not difficult to see that

$$\int |\nabla p_i|^2 dx \leq C(N, \Omega) \left(\int |V|^2 dx + \int |f_i|^2 dx \right),$$

where $C = C(N, \Omega)$. Thus

$$\int |\nabla p_\varepsilon|^2 dx \leq C(N, \Omega) \left(\int |V|^2 dx + \int |f_\varepsilon|^2 dx \right),$$

This implies that p_ε is bounded in $L^\infty(0, T; H_D^1(\Omega))$, and that there exists $p \in L^\infty(0, T; H_D^1(\Omega))$ such that

$$p_\varepsilon \rightarrow p, \quad \text{in } L^2(0, T; H_D^1(\Omega)) - \text{weak}.$$

Thanks to (3.11), $u \in \text{Sign}(p)$, a.e. in Q . Now, as usually used in nonlinear semigroup theory, we consider

$$\tilde{u}_\varepsilon = \sum_{i=0}^{n-1} \frac{(t - t_i)u_{i+1} - (t - t_{i+1})u_i}{\varepsilon} \chi_{[t_i, t_{i+1})}.$$

We see that for any test function $\xi \in H_D^1(\Omega)$, we have

$$\frac{d}{dt} \int_{\Omega} \tilde{u}_\varepsilon \xi + \int_{\Omega} (\nabla p_\varepsilon - u_\varepsilon V) \cdot \nabla \xi = \int_{\Omega} f_\varepsilon \xi, \quad \text{in } \mathcal{D}'([0, T]).$$

So, letting $\varepsilon \rightarrow 0$, and using the convergence of $(\tilde{u}_\varepsilon, u_\varepsilon, p_\varepsilon, f_\varepsilon)$ to (u, u, p, f) , we deduce that (u, p) is a weak solution of (1.5). □

To end up this section, let us mention some consequences of the Theorem 3.2 and Theorem 2.1 as well as the general theory of nonlinear semigroup theory.

4. REACTION-DIFFUSION CASE

In this section we consider the general case with a reaction term :

$$(4.1) \quad \left\{ \begin{array}{l} \frac{\partial u}{\partial t} - \Delta p + \nabla \cdot (u V) = g(\cdot, u) \\ u \in \text{Sign}(p) \end{array} \right\} \quad \text{in } Q$$

$$\left\{ \begin{array}{l} p = 0 \\ (\nabla p + u V) \cdot \nu = 0 \\ u(0) = u_0 \end{array} \right\} \quad \begin{array}{l} \text{on } \Sigma_D \\ \text{on } \Sigma_N \\ \text{in } \Omega, \end{array}$$

where $g : Q \times \mathbb{R} \rightarrow \mathbb{R}$ is a Caratheodory function ; i.e. continuous in $r \in \mathbb{R}$ and measurable in $(t, x) \in Q$. We assume that g satisfies moreover, the following assumptions :

$$(\mathcal{G}_1) \quad g^+(\cdot, -1) \in L^2(Q) \text{ and } g^-(\cdot, 1) \in L^2(Q)$$

$$(\mathcal{G}_2) \quad \text{There exists } 0 \leq \mathcal{R} \in L^2(0, T), \text{ s.t. for any } a, b \in [-1, +1], \text{ we have}$$

$$\text{Sign}_0^+(b - a) (g(t, x, b) - g(t, x, a)) \leq \mathcal{R}(t) (b - a)^+, \quad \text{for a.e. } (t, x) \in Q.$$

On sees in particular that (G2) implies that

$$(4.2) \quad g^-(\cdot, 1) - \mathcal{R}(1 - r) \leq g(\cdot, r) \leq g^+(\cdot, -1) + \mathcal{R}(1 + r), \quad \text{for any } r \in [-1, +1],$$

so that $g(\cdot, r) \in L^2(Q)$, for any $r \in [-1, +1]$.

Remark 6. See that the condition (G2) is fulfilled for instance whenever

$$\frac{\partial g}{\partial r}(t, x, \cdot) \leq \theta \quad \text{in } \mathcal{D}'([-1, 1]), \text{ for a.a. } (t, x) \in Q \text{ with } 0 \leq \theta \in \mathcal{C}([-1, 1]),$$

In this case $\mathcal{R} = \max_{-1 \leq r \leq 1} \theta(r)$.

Theorem 4.3. Under the assumption (\mathcal{G}_1) and (\mathcal{G}_2) , for any $u_0 \in L^\infty(\Omega)$ such that $|u_0| \leq 1$, a.e. in Ω , the problem (4.1) has a unique weak solution (u, p) in the sense of Definition (1.1) with $f = g(\cdot, u)$. Moreover,

$$(4.3) \quad (\mathcal{P}_1) \text{ for any } \omega_1, \omega_2 \in W^{1,1}(0, T) \text{ such that } u_0 \leq \omega_2(0) \text{ (resp. } \omega_1(0) \leq u_0) \text{ and, for any } t \in (0, T),$$

$$\dot{\omega}_2(t) + \omega_2(t) \nabla \cdot V \geq g(\cdot, \omega_2(t)) \quad \text{a.e. in } \Omega$$

$$(4.4) \quad (\text{resp. } \dot{\omega}_1(t) + \omega_1(t) \nabla \cdot V \leq g(\cdot, \omega_1(t)) \quad \text{a.e. in } \Omega),$$

we have

$$u \leq \omega_2 \quad (\text{resp. } \omega_1 \leq u) \quad \text{a.e. in } Q.$$

(\mathcal{P}_2) If g satisfies moreover

$$(\mathcal{G}_3) \quad \nabla \cdot V \geq g(\cdot, 1), \text{ a.e. in } Q \quad \text{and} \quad (\mathcal{G}_4) \quad \nabla \cdot V \leq g(\cdot, -1), \text{ a.e. in } Q,$$

then $p \equiv 0$, and u is given by the unique solution of

$$(4.5) \quad \left\{ \begin{array}{l} \frac{\partial u}{\partial t} + \nabla \cdot (u V) = g(t, x, u) \\ |u| \leq 1 \\ u V \cdot \nu = 0 \\ u(0) = u_0 \end{array} \right\} \quad \begin{array}{l} \text{in } Q \\ \text{on } \Sigma_N \\ \text{in } \Omega, \end{array}$$

in the sense that $u \in \mathcal{C}([0, T], L^1(\Omega))$, $|u| \leq 1$ a.e. in Q and

$$(4.6) \quad \frac{d}{dt} \int_{\Omega} u \xi - \int_{\Omega} u V \cdot \nabla \xi = \int_{\Omega} g(\cdot, u) \xi, \quad \text{in } \mathcal{D}'(0, T), \quad \forall \xi \in H_D^1(\Omega).$$

Corollary 4.2. *Under the assumptions of Theorem 4.3, assume moreover that $0 \leq u_0 \leq 1$ a.e. in Ω and*

$$(\mathcal{G}_5) \quad 0 \leq g(\cdot, 0) \text{ a.e. in } Q.$$

then

$$p \geq 0 \text{ and } 0 \leq u \leq 1, \quad \text{a.e. in } Q.$$

If, moreover, g satisfies (\mathcal{G}_3) , then the unique solution u is nonnegative and is given by (4.5).

Remark 7. *See that (\mathcal{G}_3) and (\mathcal{G}_4) constitute sufficient conditions to lessen the so called congestion phenomena in the transport process. This is an interesting property for the applications in crowded motion modeling. Indeed, heuristically the reaction term $g(t, x, 1)$ is connected to the reaction of the population at the position $x \in \Omega$ and time t , in a "congested" circumstance. Performing its value regarding to the divergence of the velocity vector field V , may avoid the congestion in the transportation phenomena for crowded motion.*

To prove Theorem 4.3, we use the following lemma which is a direct consequence of Lemma 1 of [14].

Lemma 4.7. *Let A be an m -accretive operator in $L^1(\Omega)$, $u_0 \in \overline{\mathcal{D}}(A)$ and $F : (0, T) \times \overline{\mathcal{D}}(A) \rightarrow L^1(\Omega)$ satisfying :*

- i) *F is Caratheodory, i.e. $t \rightarrow F(t, z)$ is measurable for any $x \in \overline{\mathcal{D}}(A)$, $z \rightarrow F(t, z)$ is continuous for a.a. $t \in (0, T)$*
- ii) *There exists $k \in L_{loc}^1(0, T)$ such that*

$$\int (F(t, z) - F(t, \hat{z})) \text{Sign}_0(z - \hat{z}) dx \leq k(t) \|z - \hat{z}\|_1, \quad \text{a.e. } t \in (0, T),$$

for every $z, \hat{z} \in \overline{\mathcal{D}}(A)$.

- iii) *There exists $c \in L_{loc}^1(0, T)$ such that*

$$\|F(t, z)\|_1 \leq c(t), \quad \text{a.e. } t \in (0, T)$$

for every $z \in \overline{\mathcal{D}}(A)$.

Then, there exists a unique $u \in \mathcal{C}([0, T]; X)$ such that u is the mild solution of

$$\begin{cases} \frac{du}{dt} + Au \ni f & \text{in } (0, T) \\ u(0) = u_0, \end{cases}$$

with $f(t) = F(t, u(t))$ a.e. $t \in (0, T)$.

Notice that, by the assumptions of Lemma 4.7, $F(\cdot, u) \in L_{loc}^1(0, T; X)$ for any $u \in \mathcal{C}([0, T]; X)$. The results of this lemma is well known in the case where $F(t, r) = f(t) + F_0(r)$, with $f(t) \in L^1(\Omega)$, for a.e. $t \in [0, T)$, and F_0 a Lipschitz continuous function in \mathbb{R} (since then $A - F_0 + kI$ is m -accretive in $L^1(\Omega)$). For the general case, one can see Lemma 1 of the paper [14] for the proof.

Proof of Theorem 4.3. Set $F : [0, T) \times L^1(\Omega) \rightarrow L^1(\Omega)$ defined by

$$F(t, z(\cdot)) = g(t, \cdot, z(\cdot)) \quad \text{a.e. in } \Omega, \text{ for any } (t, z) \in [0, T) \times L^1(\Omega).$$

See that F satisfies all the assumptions of Lemma 4.7. In particular, $k(t) = \mathcal{R}(t)$. Then, we consider $u \in \mathcal{C}([0, T), L^1(\Omega))$ the mild solution of the evolution problem

$$\begin{cases} u_t + \mathcal{A}u \ni f & \text{in } (0, T) \\ u(0) = u_0, \end{cases}$$

with $f = g(\cdot, u)$. Thanks to (4.2), it is clear that $f \in L^2(Q)$. Moreover, using Lemma 3.6, there exists p such that (u, p) is a weak solution of (4.1). The uniqueness follows from the equivalence between weak solution and mild solution as well as the uniqueness result of Theorem 2.1. Now let prove the properties (\mathcal{P}_1) and (\mathcal{P}_2) .

(\mathcal{P}_1) Thanks to the L^1 -comparison principal of Theorem 2.1, and using the fact that $(\omega_2, 0)$ is a weak solution of (1.5) with $f = \dot{\omega}_2 + \omega_2 \nabla \cdot V$, we see that

$$\begin{aligned} \frac{d}{dt} \int (u - \omega_2)^+ &\leq \int_{[u \geq \omega_2]} (g(\cdot, u) - \dot{\omega}_2 - \omega_2 \nabla \cdot V) \\ &\leq \int_{[u \geq \omega_2]} (g(\cdot, u) - g(\cdot, \omega_2))^+ \\ &\leq \mathcal{R} \int (u(t) - \omega_2)^+. \end{aligned}$$

Applying Gronwall and using the fact that $u(0) \leq \omega_2(0)$, we obtain $u(t) \leq \omega_2$ a.e. in Q . The proof of $u \geq \omega_1$ in Q follows in the same way by proving that

$$\frac{d}{dt} \int (\omega_1 - u)^+ \leq \mathcal{R} \int (\omega_1 - u)^+.$$

(\mathcal{P}_2) To prove this assertion, it is enough to prove that the problem (4.5) has a solution in the sense of (4.6). Then the conclusion follows by the uniqueness of the solution for (4.1). To this aim, we proceed by using the existence result as well as the comparison principle of the first part of the theorem. For an arbitrary $\alpha > 0$, we consider the reaction term g_α given by

$$g_\alpha(\cdot, r) = \alpha g(\cdot, r/\alpha), \quad \text{a.e. in } Q, \quad \text{for any } r \in \mathbb{R}.$$

It is clear that g_α satisfies all the assumptions of Theorem 4.3. So, we can consider (v_α, p_α) be a weak solution of

$$\begin{cases} \left. \begin{aligned} \frac{\partial v_\alpha}{\partial t} - \Delta p_\alpha + \nabla \cdot (v_\alpha V) &= g_\alpha(\cdot, v_\alpha) \\ \tilde{v}_\alpha &\in \text{Sign}(p_\alpha) \end{aligned} \right\} & \text{in } Q \\ p_\alpha &= 0 & \text{on } \Sigma_D \\ (\nabla p_\alpha + v_\alpha V) \cdot \nu &= 0 & \text{on } \Sigma_N \\ v_\alpha(0) &= \alpha u_0 & \text{in } \Omega. \end{cases}$$

On the other hand, thanks to (4.3), we see that

$$\partial_t \alpha + \alpha \nabla \cdot V \geq g_\alpha(\cdot, \alpha), \quad \text{in } Q.$$

Since $v_\alpha(0) \leq \alpha$, the assertion (\mathcal{P}_1) of the theorem implies that $v_\alpha \leq \alpha$ a.e. in Q . In a similar way, we can prove using (4.4), that $v_\alpha \geq -\alpha$ a.e. in Q . So, under the assumptions (4.3)- (4.4), we have

$$|v_\alpha| \leq \alpha, \quad \text{a.e. in } Q.$$

In particular, taking $0 < \alpha < 1$ arbitrary, this implies that $p_\alpha \equiv 0$ and v_α is in fact a solution of

$$\left\{ \begin{array}{l} \frac{\partial v_\alpha}{\partial t} + \nabla \cdot (v_\alpha V) = g_\alpha(t, x, v_\alpha) \\ |v_\alpha| \leq \alpha \end{array} \right\} \quad \text{in } Q$$

$$\left\{ \begin{array}{l} v_\alpha V \cdot \nu = 0 \\ v_\alpha(0) = \alpha u_0 \end{array} \right. \quad \begin{array}{l} \text{on } \Sigma_N \\ \text{in } \Omega. \end{array}$$

Thus, taking $u = v_\alpha/\alpha$, we generate the solution of (4.5) as claimed. □

5. EXTENSION, COMMENTS AND REMARKS

- (1) For the case where $\Gamma_N = \emptyset$, we do believe that one can combined our notion of renormalized solution with the entropic inequality à la Carillo [19] to handle the boundary condition $p = 0$ on $\partial\Omega$. Even more, we think also that in this case one can avoid the outgoing pointing condition of V on the boundary. Indeed, one sees that (2.3) and (2.4) holds to be true without any condition on $V \cdot \nu$ on the boundary. Using these formulation one can prove that weak solutions satisfy entropic inequality à la Carillo covering the boundary condition $p = 0$. Then, using doubling and de-doubling variables in the manner of [19] one can prove the L^1 -comparison principle for two weak solutions. Nevertheless, we assess that the problem would heighten in technicality. Notice here that in our case, the entropic inequalities (2.21) and (2.22) are local and do not proceed up to the boundary like in [19].
- (2) Thanks to the proof of Proposition 2.1, one sees that we can work with $\xi \in H_D^1(\Omega)$, so that any weak solution (u, p) satisfies

$$\begin{aligned} \frac{d}{dt} \int \beta(u) \xi \, dx + \int \nabla p^+ \cdot \nabla \xi \, dx + \langle V \cdot \nabla \beta(u), \xi \rangle + \int (u \nabla \cdot V(\beta'(u) \chi_{[p=0]} + \text{Sign}_0^+(p)) \xi \, dx \\ \leq \int f(\beta'(u) \chi_{[p=0]} + \text{Sign}_0^+(p)) \xi \, dx \quad \text{in } \mathcal{D}'(0, T), \end{aligned}$$

and

$$\begin{aligned} \frac{d}{dt} \int \beta(u) \xi \, dx + \int \nabla p^- \cdot \nabla \xi \, dx + \langle V \cdot \nabla \beta(u), \xi \rangle + \int (u \nabla \cdot V(\beta'(u) \chi_{[p=0]} - \text{Sign}_0^-(p)) \xi \, dx \\ \leq \int f(\beta'(u) \chi_{[p=0]} - \text{Sign}_0^-(p)) \xi \, dx \quad \text{in } \mathcal{D}'(0, T), \end{aligned}$$

for any $0 \leq \xi \in H_D^1(\Omega)$ and $\beta \in \mathcal{C}^1(\mathbb{R})$ such that $0 \leq \beta' \leq 1$. In particular, for the case of Neuman boundary condition ; i.e. $\Gamma_D = \emptyset$, these formulations implies that entropic inequalities

(2.21) and (2.22) hold to be true up to the boundary $\partial\Omega$. That is (2.21) and (2.22) are fulfilled in $\mathcal{D}'((0, T) \times \overline{\Omega})$. However, to de-double the variable by taking into account the boundary becomes unhandy since one needs to handle the trace on the boundary in a strong sense which is not guaranteed in general (one can see for instance [6] where this kind of approach is developed for elliptic-parabolic equation with homogeneous boundary condition). By the way, let us mention here that this case is fruitful for gradient flow approach in euclidean Wasserstein space. An elegant and promising proof of uniqueness is given in [26] for the one phase problem under a monotonicity assumption on V . The strength of this approach is the absolute continuity of weak solution in the set of probability equipped with Wasserstein distance.

- (3) For the case where $\Omega = \mathbb{R}^N$, one sees that working as in Section 2, it is possible to prove local Kato's inequality of the type (2.23) and achieve L^1 -comparison principle for weak solution like in Theorem (2.1). However, the existence of weak solutions in $L^1(\mathbb{R}^N)$ is not clear yet for us.
- (4) Beyond their role for the uniqueness, in our opinion formulations (2.3) and (2.4) remain to be interesting for qualitative descriptions of the weak solution. As usual for renormalized formulation, they enable to localize the description of the solution with respect to its the values. For similar description with respect to the values of p , one can see the following remarks which aims to widen the formulations (2.5) and (2.6).
- (5) One sees that the renormalization process we use in this paper concerns essentially u . The formulation (2.5) and (2.6) provide a particular renormalization for p in the region $[p \neq 0]$. Notice here, that it is possible to prove mended renormalized formulations for p . As a matter of fact, under the assumptions of Lemma 2.1, we can prove that for any $H \in \mathcal{C}^1(\mathbb{R})$ such that $H' \geq 0$ and $H(0) = 0$, we have

$$(5.10) \quad -\nabla \cdot (H(p)\nabla p) + |H(p)| \nabla \cdot V = -H'(p) |\nabla p|^2 + f H(p) \quad \text{in } \mathcal{D}'(Q).$$

If moreover, $H \geq 0$, then

$$(5.11) \quad -\nabla \cdot (H(p)\nabla p) + |H(p)| \nabla \cdot V \leq -H'(p) |\nabla p|^2 + f H(p) \quad \text{in } \mathcal{D}'((0, T) \times \overline{\Omega}).$$

These formulations improve the description of p with respect to the values of p , and yield renormalized formulations for p . As we see in Section 2, the case where H approximates Sign_0^+ is crucial for the proof of uniqueness. The general case may serve out other qualitative description of the solution. The proof of (5.10) and (5.11) follows more or less similar arguments as for the proof Lemma 2.1.

- (6) Coming back to the approximation (3.2), thanks to Proposition 3.3, we know that weak solutions of the stationary problem of (3.2) depends weak-continuously on the non-linearity \mathcal{H}_ε . For the evolution problem, working as in the proof of Proposition 3.3, one can prove that the problem

$$\left\{ \begin{array}{l} \frac{\partial u}{\partial t} - \Delta p + \nabla \cdot (u V) = f \\ u = \mathcal{H}_\varepsilon(p) \end{array} \right\} \quad \text{in } Q$$

$$\left\{ \begin{array}{l} p = 0 \\ (\nabla p + u V) \cdot \nu = 0 \end{array} \right. \quad \begin{array}{l} \text{on } \Sigma_D \\ \text{on } \Sigma_N \end{array}$$

$$\left\{ \begin{array}{l} u(0) = u_0 \end{array} \right. \quad \text{in } \Omega,$$

has a unique weak solution $(u_\varepsilon, p_\varepsilon) \in \mathcal{C}([0, T], L^1(\Omega)) \times L^2(0, T; H_D^1(\Omega))$, with $u_\varepsilon = \mathcal{H}_\varepsilon(p_\varepsilon)$ a.e. in Q , and as $\varepsilon \rightarrow 0$, we have $u_\varepsilon \rightarrow u$, in $L^\infty(Q) - \text{weak}^*$ and $p_\varepsilon \rightarrow p$, in $L^2(0, T; H_D^1(\Omega))$. We do believe that the convergence of u_ε remains to be true strongly in $L^1(\Omega)$, for both problems stationary and evolution one. We did not get into these further questions here. This need more fine estimates on u (like BV -estimates). We'll touch them in forthcoming works. The case where one replace \mathcal{H}_ε by an arbitrary non-linearity which converges to Sign, in the sense of resolvent, like for instance the porous medium non-linearity $\varphi(r) = |r|^{m-1}r$, for any $r \in \mathbb{R}$, with $m \rightarrow \infty$, is likely concerned.

- (7) It is clear that our approach is based on a renormalization process à la DiPerna-Lions. This corroborates that this concept is still fruitful for the uniqueness of weak solution (solution in the sense of distribution) for nonlinear versions of Fokker-Planck equation of the type (1.1). As we quote in the previous remarks, operating with the absolute continuity of weak solution in the set of probability equipped with Wasserstein distance seems to be promising (cf. [26]) even if it is actually hampered by conceptual technical assumptions like the sign of the solution, conservation of the mass and monotone transport field (connected to some kind of convexity conditions). We do not overlook the L^1 -kinetic approach, which was developed in [21] for Cauchy problems of general degenerate parabolic-hyperbolic equations with non-isotropic non-linearity. The approach could be applied to more general situations and, as far as we know, this approach has not yet been explored for problems of the type (1.1) and (1.2). At last let us quote here the very recent work of [24] (that we have just learned through Benoit Perthame) where the authors deal with the uniqueness of weak solutions for the one phase problem (1.2) in \mathbb{R}^N , by means of Hilbert's duality method, in the case where $V = \nabla\Phi$ is smooth enough (at least such that $\nabla\Delta\Phi$ is a $L_{loc}^{12/5}$ -Lebesgue function).

ACKNOWLEDGMENTS

We would like to thank Benoit Perthame for informing us about the papers [30], [35] and [24].

6. APPENDIX

The aim of this section is to complete the proof of Proposition 2.2 with the proof of de-doubling variable process concerning the vector field part of (2.25), which is

$$\lim_{\lambda \rightarrow 0} \int_{\Omega} \int_{\Omega} u(t, t, x, y)^+ (V(x) - V(y)) \cdot \nabla_y \zeta_\lambda \, dx dy.$$

Remember that

$$\zeta_\lambda(x, y) = \xi \left(\frac{x + y}{2} \right) \delta_\lambda \left(\frac{x - y}{2} \right),$$

for any $x, y \in \Omega$, where $\xi \in \mathcal{D}(\Omega)$, δ_λ is a sequence of usual mollifiers in \mathbb{R}^N . For instance, take

$$\delta_\lambda(x) = \frac{1}{\lambda^N} \delta \left(\frac{|x|^2}{\lambda^2} \right), \quad \text{for any } x \in \mathbb{R}^N$$

where $0 \leq \delta \in \mathcal{D}(0, 1)$ and $\frac{1}{2^N} \int \delta(|z|^2) \, dz = 1$. A typical example is given by $\delta(r) = C e^{\frac{-1}{1-r^2}} \chi_{[|r| \leq 1]}$, for any $r \in \mathbb{R}^N$.

Lemma 6.8. For any $h \in L^1((0, T) \times \Omega \times \Omega)$, $V \in W^{1,1}(\Omega)$ and $\psi \in \mathcal{D}(0, T)$, we have

$$\begin{aligned} & \lim_{\lambda \rightarrow 0} \int_0^T \int_{\Omega} \int_{\Omega} h(t, x, y) (V(x) - V(y)) \cdot \nabla_y \zeta_{\lambda}(x, y) \psi(t) dt dx dy \\ &= \int_0^T \int_{\Omega} h(x, x) \nabla \cdot V(x) \xi(x) \psi(t) dt dx. \end{aligned}$$

Proof. We see that

$$(\nabla_x + \nabla_y)\zeta = \delta_{\lambda} \left(\frac{x-y}{2} \right) \nabla \xi \left(\frac{x+y}{2} \right).$$

Since, for any $\eta \in \mathbb{R}^N$,

$$\nabla \delta_{\lambda}(\eta) = \frac{2}{\lambda^N} \frac{z}{\lambda^2} \delta' \left(\frac{|z|^2}{\lambda^2} \right),$$

we have

$$\nabla \delta_{\lambda} \left(\frac{x-y}{2} \right) = \frac{2}{\lambda^N} \frac{x-y}{2\lambda^2} \delta' \left(\frac{|x-y|^2}{(2\lambda)^2} \right), \quad \text{for any } x, y \in \Omega.$$

So,

$$\begin{aligned} \nabla_y \zeta(x, y) &= \frac{1}{2} \nabla \xi \left(\frac{x+y}{2} \right) \delta_{\lambda} \left(\frac{x-y}{2} \right) - \frac{1}{2} \xi \left(\frac{x+y}{2} \right) \nabla \delta_{\lambda} \left(\frac{x-y}{2} \right) \\ &= \nabla \xi \left(\frac{x+y}{2} \right) \delta_{\lambda} \left(\frac{x-y}{2} \right) - \frac{1}{\lambda^{N+1}} \xi \left(\frac{x+y}{2} \right) \delta' \left(\frac{|x-y|^2}{(2\lambda)^2} \right) \frac{x-y}{2\lambda} \end{aligned}$$

and

$$\begin{aligned} (V(x) - V(y)) \cdot \nabla_y \zeta &= \frac{1}{2} (V(x) - V(y)) \cdot \nabla \xi \left(\frac{x+y}{2} \right) \delta_{\lambda} \left(\frac{x-y}{2} \right) \\ &\quad - \frac{1}{\lambda^N} \xi \left(\frac{x+y}{2} \right) \delta' \left(\frac{|x-y|^2}{(2\lambda)^2} \right) \frac{V(x) - V(y)}{\lambda} \cdot \frac{x-y}{2\lambda}. \end{aligned}$$

This implies that

$$\begin{aligned} & \lim_{\lambda \rightarrow 0} \iint h(x, y) (V(x) - V(y)) \cdot \nabla_y \zeta dx dy \\ &= \lim_{\lambda \rightarrow 0} -\frac{1}{\lambda^N} \iint \xi \left(\frac{x+y}{2} \right) \delta' \left(\frac{|x-y|^2}{(2\lambda)^2} \right) \frac{V(x) - V(y)}{\lambda} \cdot \frac{x-y}{2\lambda} dx dy =: I(\lambda). \end{aligned}$$

Changing the variable by setting

$$z = \frac{x-y}{2\lambda}$$

we get $x = y + 2\lambda z$ and $dx = (2\lambda)^N dz$, so that

$$\begin{aligned} -I(\lambda) &= 2^N \iint h((y + 2\lambda z), y) \xi(y + \lambda z) \delta'(|z|^2) \frac{V(y + 2\lambda z) - V(y)}{\lambda} \cdot z \, dz dy \\ &= 2^N \sum_{i=1}^N \iint h((y + 2\lambda z), y) \xi(y + \lambda z) \delta'(|z|^2) \frac{V_i(y + 2\lambda z) - V_i(y)}{\lambda} z_i \, dz dy. \end{aligned}$$

Letting $\lambda \rightarrow 0$, we get

$$\begin{aligned} \lim_{\lambda \rightarrow 0} I(\lambda) &= -2^{N+1} \sum_{i=1}^N \iint \xi(y) h((y, y) \nabla V_i(y) \cdot z z_i \delta'(|z|^2) \, dz dy \\ &= -2^{N+1} \sum_{i,j=1}^N \iint \xi(y) h((y, y) \frac{\partial V_i(y)}{\partial y_j} z_j z_i \delta'(|z|^2) \, dz dy \\ &= -2^{N+1} \sum_{i=1}^N \iint \xi(y) h((y, y) \frac{\partial V_i(y)}{\partial y_j} z_i^2 \delta'(|z|^2) \, dz dy \\ &= -\frac{2^{N+1}}{N} \int \xi(y) h((y, y) \nabla \cdot V(y) \, dy \int |z|^2 \delta'(|z|^2) \, dz, \end{aligned}$$

where we use the fact that (since $z \in B(0, 1) \rightarrow \delta(|z|^2)$ is symmetric)

$$\int z_j z_i \delta'(|z|^2) \, dz = 0, \quad \text{for any } i \neq j$$

and

$$\sum_{i=1}^N \int z_i^2 \delta'(|z|^2) \, dz = N \int z_1^2 \delta'(|z|^2) \, dz = \int |z|^2 \delta'(|z|^2) \, dz.$$

At last, we use the fact that

$$\frac{2^{N+1}}{N} \int |z|^2 \delta'(|z|^2) \, dz = -1.$$

Indeed, recall that

$$\int |z|^2 \delta'(|z|^2) \, dz = \int_0^\infty \int_{S^{n-1}} |r\theta|^2 \delta'(|r\theta|^2) r^{n-1} dr \, ds_{n-1}(\theta),$$

where s_{n-1} is the probability measure defined on the sphere S_{n-1} by

$$s_{n-1}(A) = |\{rx : r \in [0, 1], x \in A\}|.$$

This implies that

$$\begin{aligned} \int |z|^2 \delta'(|z|^2) dz &= |B(0,1)| \int_0^1 r^2 \delta'(r^2) r^{n-1} dr \\ &= |B(0,1)| \int_0^1 \delta'(r^2) r^{n+1} dr \\ &= -\frac{N}{2} \int \delta(|z|^2) dz. \end{aligned}$$

Thus the result. □

REFERENCES

- [1] D. ALEXANDER, I. KIM and Y. YAO. Quasi-static evolution and congested crowd transport. non-linearity, 27 (2014), No.4, 823-858.
- [2] H. W. ALT and H. W. LUCKHAUS. Quasilinear elliptic-parabolic differential equations. *Math. Z.*, 183(1983), pp. 311-341.
- [3] L. AMBROSIO. Pairings between measures and bounded functions and compensated compactness. *Invent. Math.*, 158(2):227-260, 2004.
- [4] L. AMBROSIO and G. CRIPPA. Existence, uniqueness, stability and differentiability properties of the flow associated to weakly differentiable vector fields. In *Transport equations and multi-D hyperbolic conservation laws*, volume 5 of *Lect. Notes Unione Mat. Ital.*, pages 3-57. Springer, Berlin, 2008.
- [5] L. AMBROSIO, G. CRIPPA and S. MANIGLIA. Traces and fine properties of a BD class of vector fields and applications. *Ann. Fac. Sci. Toulouse Math.*, (6), 14(4):527-561, 2005.
- [6] B. ANDREIANOV and F. BOUHSSIS. Uniqueness for an elliptic-parabolic problem with Neumann boundary condition. *J. Evol. Equ.* 4 (2004), no. 2, 273-295.
- [7] B. ANDREIANOV and N. IGBIDA. Revising Uniqueness for a Nonlinear Diffusion-Convection Equation. *J. Diff. Eq.*, Vo. 227 (2006), no-1 69-79.
- [8] B. ANDREIANOV and N. IGBIDA. Uniqueness for the Inhomogeneous Dirichlet Problem for Elliptic-Parabolic Equations, . *Proc. Edinburgh Math. Society*, 137A, 1119-1133, 2007
- [9] B. ANDREIANOV and N. IGBIDA. On Uniqueness techniques for degenerate convection-diffusion problems. *Int. J. of Dynamical Systems and Differential Equations*, 2012 - Vol. 4, No.1/2 pp. 3 - 34
- [10] G. ANZELLOTTI. Transport equation and Cauchy problem for BV vector fields. *Ann. Mat. Pura Appl.*, (4), 135:293-318 (1984), 1983.
- [11] V. BARBU. Nonlinear differential equations of monotone types in Banach spaces. *Springer Mono-graphs in Mathematics*, Springer, New York, 2010.
- [12] Ph. BÉNILAN. Opérateurs accréatifs et semi-groupes dans les espaces L^p ($1 \leq p \leq \infty$). *Functional Analysis and Numerical Analysis*, Japan-France seminar, H.Fujita (ed.), Japan Society for the Advancement of Science 1978.
- [13] Ph. BÉNILAN and N. IGBIDA. Singular Limit of the Changing Sign Solutions of the Porous Medium Equation. *J. Evol. Equations*. 3 (2003), no. 2, 215-224
- [14] Ph. BÉNILAN and N. IGBIDA. Singular Limit of Perturbed Nonlinear Semigroups. *Comm. Appl. Nonlinear Anal.*, 3 (1996), no. 4, 23-42.
- [15] Ph. BÉNILAN and N. IGBIDA. The Mesa Problem for the Neumann Boundary Value Problem. *J. Differential Equations*, 196 (2004), no. 2, 301-315.
- [16] F. BOYER. Trace theorems and spatial continuity properties for the solutions of the transport equation. *Differential Integral Equations*, 18(8):891-934, 2005.
- [17] F. BOYER and P. FABRIE. Quasi-static evolution and congested crowd transport. *Mathematical tools for the study of the incompressible NavierStokes equations and related models*, volume 183 of *Applied mathematical sciences*. Springer, 2012.

- [18] H. BRÉZIS, *Opérateurs maximaux monotones et semigroups de contractions dans les espaces de Hilbert. (French)*, North-Holland Mathematics Studies, No. 5. Notas de Matematica (50). North-Holland Publishing Co., Amsterdam-London; American Elsevier Publishing Co., Inc., New York, 1973.
- [19] J. CARRILLO. Entropy solutions for nonlinear degenerate problems. *Arch. Rational Mech. Anal.*, 147(1999), pp. 269–361.
- [20] G.-Q. CHEN and H. FRID. Divergence-measure fields and hyperbolic conservation laws.. *Arch. Ration. Mech. Anal.*, 147(2):89–118, 1999.
- [21] G.-Q. CHEN and B. PERTHAME. Well-posedness for non-isotropic degenerate parabolic-hyperbolic equations. *Ann. I. H. Poincaré*, 20(4), 645-668, 2003.
- [22] G.-Q. CHEN, M. TORRES and W. P. ZIEMER. Traces and fine properties of a BD class of vector fields and applications. *Ann. Fac. Sci. Toulouse Math.*, (6), 14(4):527-561, 2005.
- [23] G. CRIPPA C. DONADELLO and L. SPINOLO. Initial-boundary value problems for continuity equations with BV coefficients. *Journal Math. Pure Appl.*, (102)1, 79-98, 2014.
- [24] N. DAVID and M. SCHMIDTCHEN. On the incompressible limit for a tumour growth model incorporating convective effects . Preprint. <https://hal.archives-ouvertes.fr/hal-03162169>.
- [25] E. DIBENEDETTO and A. FRIEDMAN. The Ill-Posed Hele-Shaw Model and the The Stefan Problem for Supercooled Water. *Trans. Amer. Math. Soc.*, 282(1984), 183–204.
- [26] S. DI MARINO and A.R. MÉSZÁROS. Uniqueness issues for evolution equations with density constraints. *Mathematical Mod. Meth. App. Sci.*, 26 (09), 1761-1783
- [27] R. J. DIPERNA and P.-L. LIONS Ordinary differential equations, transport theory and Sobolev spaces. *Ann. Fac. Sci. Toulouse Math.*, 98(3):511-547, 1989.
- [28] C. M. ELLIOT and V. JANOVSKÝ. Avariational Inequality Approach to the Hele-Shaw Flow With a Moving Boundary. *Proc. Roy. Soc. Edinburgh Sect. A*, 88(1981), 93–107.
- [29] C.M. ELLIOTT and J.R. OCKENDON. *Weak and Variational Methods for Moving Boundary Problems* Pitman, Boston, 1982.
- [30] A. FIGALI Existence and uniqueness of martingale solutions for SDEs with rough or degenerate coefficients *J. Funct. Anal.*, 254 (2008) 109-153.
- [31] O. GIL and F. QUIRÓS. Convergence of the porous media equation to Hele-Shaw. *Nonlinear Anal. Ser. A: Theory Methods*, 44 (2001), no. 8, 1111-1131.
- [32] N. IGBIDA. The Mesa-Limit of the Porous Medium-Equation and the Hele-Shaw Problem. *Differential Integral Equations*, 15 (2002), no. 2, 129–146
- [33] N. IGBIDA and J. M. URBANO. Uniqueness for nonlinear degenerate problems. *NoDEA Nonlin. Diff. Eq. Appl.* 10(2003), no.3, pp. 287–307.
- [34] S.N. KRUSHKOV. First order quasilinear equations with several space variables. *Mat. USSR-Sbornik*, 10(1970), pp. 217–242.
- [35] C. LEBRIS and P.L. LIONS. Existence and uniqueness of solutions to Fokker-Planck type equations with irregular coefficients. *Comm. Partial Differential Equations*, 33(2008), 1272-1317.
- [36] B. MAURY, A. ROUDNEFF-CHUPIN, F. SANTAMBROGIO, A macroscopic crowd motion model of gradient flow-type, *Math. Models and Meth. in Appl. Sci.*, 20 (2010), No. 10, 1787-1821.
- [37] B. MAURY, A. ROUDNEFF-CHUPIN, F. SANTAMBROGIO, Congestion-driven dendritic growth. *Discrete Contin. Dyn. Syst.*, 34 (2014), no. 4, 1575-1604.
- [38] B. MAURY, A. ROUDNEFF-CHUPIN, F. SANTAMBROGIO and J. VENEL. Handling congestion in crowd motion modeling, *Netw. Heterog. Media*, 6 (2011), No. 3, 485-519.
- [39] A. R. MÉSZÁROS and F. SANTAMBROGIO. Advection-diffusion equations with density constraints *Analysis and PDE*, 9(3):615-644.
- [40] S. MISCHLER. On the trace problem for solutions of the Vlasov equation. *Comm. Partial Differential Equations*, 25(7-8):1415–1443, 2000.
- [41] F. OTTO. L^1 contraction and uniqueness for quasilinear elliptic-parabolic equations. *J. Diff. Eq.*, 131(1996), pp. 20-38.
- [42] S. RICHARDSON. Hele Shaw Flows With a Free Boundary Produced by the Injection of Fluid in a Narrow Channel. *J. Fluid Mech.*, 56(1972), 609-618.
- [43] P.G. SAFFMAN and G.I. TAYLOR. The Penetration of a Fluid Into a Porous Medium or Hele Shaw Cell Containing a More Viscous Liquid, *Proc. Roy. Soc.* 245(1958), 312–329.

- [44] B. PERTHAME, F. QUIRÓS, J.L. VÁZQUEZ. The Hele-Shaw asymptotics for mechanical models of tumor growth. Arch. Rational Mech. Anal., 212 (2014), 93-127.
- [45] J. L. VÁZQUEZ. The porous medium equation : mathematical theory. Oxford University Press, 2007.
- [46] A. VISINTIN. Models of Phase Transitions. Progres in nonlinear Differential Equations and their Applications. Birkhauser, 1996.