



HAL
open science

A glycomic approach reveals a new mycobacterial polysaccharide

Gerald Larrouy-Maumus, Martine Gilleron, Henrieta Skovierová, Sophie Zuberogoitia, Patrick Brennan, Germain Puzo, Mary Jackson, Jérôme Nigou

► **To cite this version:**

Gerald Larrouy-Maumus, Martine Gilleron, Henrieta Skovierová, Sophie Zuberogoitia, Patrick Brennan, et al.. A glycomic approach reveals a new mycobacterial polysaccharide. *Glycobiology*, 2015, 25 (11), pp.1163-1171. 10.1093/glycob/cwv061 . hal-03178008

HAL Id: hal-03178008

<https://hal.science/hal-03178008>

Submitted on 23 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A glycomic approach reveals a new mycobacterial polysaccharide

Journal:	<i>Glycobiology</i>
Manuscript ID:	GLYCO-2015-00021.R1
Manuscript Type:	Regular Manuscripts
Date Submitted by the Author:	05-Aug-2015
Complete List of Authors:	<p>Larrouy-Maumus, Gérald; Institut de Pharmacologie et de Biologie Structurale, Tuberculosis & Infection Biology</p> <p>Gilleron, Martine; Institut de Pharmacologie et de Biologie Structurale, Tuberculosis & Infection Biology</p> <p>Skovierová, Henrieta; Colorado State University, Microbiology, Immunology and Path</p> <p>Zuberogoitia, Sophie; Institut de Pharmacologie et de Biologie Structurale, Tuberculosis & Infection Biology</p> <p>Brennan, Patrick; Colorado State University, Microbiology, Immunology and Pathology</p> <p>Puzo, Germain; Institut de Pharmacologie et de Biologie Structurale, Tuberculosis & Infection Biology</p> <p>Jackson, Mary; Colorado State University, Microbiology, Immunology and Pathology</p> <p>Nigou, Jerome; Institut de Pharmacologie et de Biologie Structurale, Tuberculosis & Infection Biology</p>
Key Words:	Glycomic, Mycobacterium, Polysaccharide

1
2
3 **A glycomic approach reveals a new mycobacterial polysaccharide**
4
5
6

7 **Gérald Larrouy-Maumus^{1,2,§}, Martine Gilleron^{1,2}, Henrieta Skovierová³, Sophie**
8 **Zuberogoitia^{1,2}, Patrick J. Brennan³, Germain Puzo^{1,2}, Mary Jackson³ and Jérôme**
9 **Nigou^{1,2,*}**
10
11
12
13

14
15
16
17
18 ¹ CNRS; Institut de Pharmacologie et de Biologie Structurale (IPBS); 205 route de Narbonne,
19 F-31077 Toulouse, France.
20

21
22 ² Université de Toulouse; Université Paul Sabatier; IPBS; F-31077 Toulouse, France.
23

24 ³ Mycobacteria Research Laboratories, Department of Microbiology, Immunology and
25 Pathology, Colorado State University, Fort Collins, CO 80523-1682, USA.
26
27
28

29
30
31 [§] Present address: MRC-Centre for Molecular Bacteriology and Infection, Imperial College,
32 Flowers Building, South Kensington, London, SW7 2AZ.
33
34
35
36
37

38 Address correspondence to:

39
40 Dr Jérôme Nigou, Institut de Pharmacologie et de Biologie Structurale, UMR 5089 CNRS /
41 Université Paul Sabatier, 205 route de Narbonne, BP 64182, 31077 Toulouse Cedex 4,
42 France. Tel: +33 5 61 17 55 54; Fax: +33 5 61 17 59 94; E-mail: jerome.nigou@ipbs.fr
43
44
45
46
47

48
49 Running title: Polysaccharide isolation and analysis by a glycomic approach
50
51

52
53
54 Keywords: Glycomic/Mycobacterium/Polysaccharide
55
56
57
58
59
60

Abstract

Mycobacterium tuberculosis lipoarabinomannan (LAM) and biosynthetically related lipoglycans and glycans play an important role in host-pathogen interactions. Therefore, the elucidation of the complete biosynthetic pathways of these important molecules is expected to afford novel therapeutic targets. The characterization of biosynthetic enzymes and transporters involved in the formation and localization of these complex macromolecules in the bacterial cell envelope largely relies on genetic manipulation of mycobacteria and subsequent analyses of lipoglycan structural alterations. However, lipoglycans are present in relatively low amounts. Their purification to homogeneity remains tedious and time-consuming. In order to overcome these issues and to reduce the biomass and time required for lipoglycan purification, we report here the development of a methodology to efficiently purify lipoglycans by sodium deoxycholate-PAGE. This faster purification method can be applied on a small amount of mycobacterial cells biomass (10-50 mg), resulting in tens of micrograms of purified lipoglycans. This amount of purified products was found to be sufficient to undertake structural analyses of lipoglycans and glycans carbohydrate domains by a combination of highly sensitive analytical procedures, involving cryoprobe NMR analysis of intact macromolecules and chemical degradations monitored by gas chromatography and capillary electrophoresis. This glycomic approach was successfully applied to the purification and structural characterization of a newly identified polysaccharide, structurally related to LAM, in the model fast-growing species *Mycobacterium smegmatis*.

Introduction

One-third of the world's population is infected by *Mycobacterium tuberculosis* (Mtb), the aetiological agent of human tuberculosis. Mtb cell envelope constituents, such as glycolipids, lipoglycans and glycans, play an important role in bacteria-host cell interactions (Angala et al, 2014; Ray et al, 2013). Biosynthetically related lipoglycans, comprising phosphatidyl-*myo*-inositol mannosides (PIMs), lipomannan (LM) and lipoarabinomannan (LAM) play essential roles in mycobacterial physiology and are important immunomodulatory molecules of the bacilli (Briken et al, 2004; Gilleron et al, 2008; Mishra et al, 2011; Torrelles & Schlesinger, 2010; Vergne et al, 2015). LAM are amphipathic macromolecules which possess a tripartite structure, including i) a mannosyl-phosphatidyl-*myo*-inositol (MPI) anchor, ii) a polysaccharide backbone composed of a D-mannan with a linear chain of $\alpha(1\rightarrow6)$ -linked D-Man_p units to which are attached single $\alpha(1\rightarrow2)$ -D-Man_p residues and a D-arabinan, with a linear chain of $\alpha(1\rightarrow5)$ -linked D-Araf units punctuated by branching produced by 3,5-*O*-linked α -D-Araf units, and finally iii) caps assigned either to small manno-oligosaccharides (ManLAM) or phosphoinositol (PILAM) (Berg et al, 2007; Mishra et al, 2011; Nigou et al, 2003). Lipid-free glycans structurally related to LAM and LM, i.e. arabinomannan (AM) and mannan, are also found in the outermost layers of the mycobacterial cell envelope (Nigou et al, 1999; Ortalo-Magne et al, 1995).

The elucidation of the complete biosynthetic pathways of these important molecules is expected to afford novel therapeutic targets (Angala et al, 2014; Berg et al, 2007; Gilleron et al, 2008). In support of this assumption, D-arabinan biosynthesis is the target of ethambutol (Deng et al, 1995), a first-line drug in the treatment of tuberculosis, as well as of benzothiazinones (Makarov et al, 2009), which are new antituberculous drug candidates in preclinical development. The characterization of biosynthetic enzymes as well as the transporters involved in the formation and localization of these complex macromolecules

1
2
3 largely relies on genetic manipulation of mycobacteria and subsequent analyses of lipoglycan
4
5 structural alteration (Angala et al, 2014; Gilleron et al, 2008). However, lipoglycans are
6
7 present in relatively low amounts (~1 mg/g of dry mycobacteria) and their purification to
8
9 homogeneity remains tedious and time-consuming (Laneelle et al, 2015; Nigou et al, 2004).
10
11 Conventional lipoglycan purification protocol involves several solvent extractions, followed
12
13 by enzymatic digestions and two chromatographic steps: hydrophobic interaction
14
15 chromatography to separate glycans and lipoglycans and then gel filtration to separate PIM,
16
17 LM and LAM individually. If homogenous fractions of PIM, LM or LAM are desired and
18
19 because gel filtration chromatography is not adapted to small quantities, a minimum of 1 g of
20
21 dry biomass is required to start with.
22
23

24
25 In order to reduce the biomass and the time required for lipoglycan purification, we aimed at
26
27 replacing the two chromatographic steps by a single step of purification using polyacrylamide
28
29 gel electrophoresis (PAGE). This approach has been successful for purification of LPS and
30
31 proteins (Claverol et al, 2003; Galvani et al, 2000; Hardy et al, 1997; Kurth & Stoffel, 1990;
32
33 Pupo et al, 2000; Pupo et al, 2004). Typically, after PAGE migration, macromolecules are
34
35 passively eluted from the gel by incubation in an elution buffer. The latter can be compatible
36
37 with mass spectrometry (MS), allowing subsequent structural analysis. This approach has
38
39 also proved to permit biological, including *in vivo*, studies as shown with LPS (Pupo et al,
40
41 1999).
42
43

44
45 Here, we report the development of a methodology to efficiently purify lipoglycans by
46
47 sodium deoxycholate-PAGE (DOC-PAGE). This faster purification method can be applied
48
49 on a small amount of mycobacterial cells biomass (10-50 mg), resulting in tens of
50
51 micrograms of purified lipoglycans. This amount is sufficient, as demonstrated, to undertake
52
53 structural analyses of lipoglycans and glycans carbohydrate domains by a combination of
54
55 highly sensitive analytical procedures, involving cryoprobe NMR analysis of intact
56
57
58
59
60

1
2
3 macromolecules and chemical degradations monitored by gas chromatography and capillary
4
5 electrophoresis. This glycomic approach was successfully applied to the structural
6
7 characterization of a newly identified polysaccharide in the model fast-growing species
8

9
10 *Mycobacterium smegmatis*.
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Results and Discussion

The development of a PAGE-based methodology to efficiently purify lipoglycans required to be able: i) to quickly localize lipoglycans in the gel in order to avoid their diffusion before elution, and ii) to efficiently elute the lipoglycans from the gel. *M. smegmatis* LAM purified by the conventional protocol was used as a standard to set up and optimize the different steps.

Direct detection of fluorescently-labelled lipoglycans in gels

LAM can be routinely detected on gels using either zinc sulfate negative staining or silver nitrate staining (Venisse et al, 1993). Zinc sulfate negative staining is reversible and non-degradative (Hardy et al, 1997) and reveals LAM as a translucent area. However, the presence of zinc sulfate hinders subsequent MS analyses. Silver nitrate staining is degradative and, once stained, the chemical structure of LAM is altered. So, none of the classical staining methods of LAM is compatible with a subsequent structural analysis. We thus resorted on an alternative strategy consisting in the detection of LAM on a standard lane and its use as a migration reference to excise gel bands containing unlabelled LAM (Figure 1). However, zinc sulfate and silver nitrate staining procedures take around 2 hrs (Hardy et al, 1997; Venisse et al, 1993), a period of time during which molecules can diffuse in the gel, precluding a precise excision. To overcome this limitation, we decided to fluorescently-label LAM prior to PAGE migration. The chromophore Lucifer Yellow (LY), which bears a hydrazide function enabling its coupling to lipoglycans after periodate oxidation (Pitarque et al, 2005), was selected. The labelling procedure was optimized by focusing on three parameters: i) oxidation reaction time, ii) LAM/LY ratio and iii) coupling reaction time. After optimization, the labelling procedure was set as follows: one-hour oxidation reaction, 10 min coupling reaction using a LAM/LY ratio of 1:100 mol/mol, and 1.5 hrs dialysis against water to remove most of the probe excess. Labelling 10 µg of LAM or less was sufficient to

1
2
3 visualize it on the gel under UV light (Figures 1 and 2). Importantly, LY-labelling did not
4
5 alter the migration of the lipoglycans. Indeed, LY-labelled LAM, when revealed by either
6
7 zinc sulphate negative staining (Figure 2A) or UV (Figure 2B), showed a migration identical
8
9 to that of native LAM. Altogether, this ‘real-time’ detection method allowed a precise
10
11 excision of the gel bands containing lipoglycans immediately after migration (Figure 1),
12
13 thereby reducing the risk of molecules diffusion in the gel and increasing their recovery
14
15 yields.
16
17

18 19 20 21 *Recovery of lipoglycans from gel by passive elution*

22
23 Several methods of proteins or LPS elution from gel fragments have been described (Claverol
24
25 et al, 2003; Kurth & Stoffel, 1990; Pupo et al, 1999; Pupo et al, 2000; Pupo et al, 2004). For
26
27 proteins, passive elution is usually employed using an extraction solution that can contain up
28
29 to 0.1 % of SDS in water (Claverol et al, 2003; Kurth & Stoffel, 1990). As for LPS, Hardy *et*
30
31 *al.* (Hardy et al, 1997; Pupo et al, 1999) developed a grinding process which involves passing
32
33 the gel pieces through 100 to 32 microns diameter pore metal filters to facilitate the elution of
34
35 LPS from polyacrylamide. The recovery yields reported were in the range of 70-80 % for
36
37 LPS without O-antigen chain but 10% only for O-antigen-containing LPS, which are
38
39 structurally closer to mycobacterial lipoglycans (De Castro et al, 2010; Raetz & Whitfield,
40
41 2002). Based on this set of previous studies, we focused on two types of extraction solutions,
42
43 one based on the use of detergents and the other on organic solvents. 20 µg of pure *M.*
44
45 *smegmatis* LAM were loaded on the gel and submitted to DOC-PAGE migration. DOC-
46
47 PAGE was selected over SDS-PAGE because lipoglycans migration was more focused (not
48
49 shown). The gel band containing LAM was excised according to the migration reference
50
51 provided by LY-labelled LAM (Figure 1) and cut into 0.5 mm sized pieces that were placed
52
53 in the elution solution. Initially, the passive elution condition was set to 3 hours at 37 °C. The
54
55
56
57
58
59
60

1
2
3 objective was to find an extraction solution more efficient than water used as a control.
4
5 Detergents are supposed to promote the disaggregation of the molecules, thereby facilitating
6
7 their elution through the mesh of the gel. Two detergents were tested: SDS and DOC, both at
8
9 concentrations of 0.1 % and 1 % in water. Higher concentrations make the elimination of
10
11 these detergents difficult and prevent any structural analysis (Claverol et al, 2003; Kurth &
12
13 Stoffel, 1990). The results presented in Table 1 show that the use of DOC did not provide any
14
15 improvement as compared to water alone (yield of 4%). 1% SDS increased the recovery yield
16
17 up to 9% but this remained insufficient. We thus concluded that passive elution using
18
19 detergents was not a suitable method for lipoglycans and then resorted to organic solvents.
20
21 Acetonitrile was first tested at concentrations of 15, 30 or 60 % in water. By dehydrating
22
23 gels, acetonitrile induces a leakage of water, draining out of the gel the other molecules
24
25 entrapped. However, elution yields were again comparable to those obtained with water alone
26
27 (Table 1). We finally tested the volatile base triethylamine (TEA), assuming that TEA may
28
29 increase mesh size, thereby facilitating the passive elution of molecules. TEA was previously
30
31 used at a concentration of 5 % in water for elution of LPS (Pupo et al, 2000). Here, we tested
32
33 it at 0.1, 1, 5 or 10 % in water. The recovery yield was improved by the use of 10% TEA as
34
35 compared to water alone, but remained in the same range than that obtained with 1 % SDS
36
37 (Table 1). Unlike SDS, however, TEA is volatile and can therefore be removed by
38
39 evaporation under vacuum. After optimization of the elution time, the final conditions were
40
41 set as incubation in 10 % TEA in water for 12 h at 37 °C, followed by two washes of 2 h with
42
43 the same solution. The recovery yield was ~60 % as compared to ~20 % using water only
44
45 (Table 1).
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Structural analysis of LAM eluted from gel

1
2
3 We next determined whether the new purification procedure developed herein did not alter
4 lipoglycans structure and whether it was compatible with the set of highly sensitive analytical
5 procedures we previously developed to investigate the latter (Nigou et al, 2000). 2 mg of
6 LAM were purified from *M. smegmatis* using the conventional protocol (LAMc). A portion
7 of this LAM batch (~200 µg) was submitted to the new purification procedure using DOC-
8 PAGE as described above and ~100 µg of LAM were recovered from the gel (LAMg). A
9 comparative structural analysis of both LAM samples was then performed. MALDI-TOF
10 mass spectrometry analysis exhibited for both deacylated LAM samples a broad unresolved
11 peak centered at m/z 14,900, indicating a molecular mass around 14.9 kDa for the major
12 molecular species (Figure 3). In addition, the intrinsic heterogeneity of LAM, as revealed by
13 the peak width, was similar for both samples. Thus, the overall structure of LAM did not
14 appear to be altered by the in-gel purification steps. To gain further insight into the fine
15 structure of the polysaccharidic moiety, both LAM samples were submitted to chemical
16 degradations, followed by capillary electrophoresis monitored by laser-induced fluorescence
17 (CE-LIF) or gas chromatography (GC) analyses. Total acid hydrolysis showed the presence
18 of Ara and Man, with an Ara/Man ratio of 0.8 ± 0.1 and 0.9 ± 0.05 for LAMc and LAMg
19 respectively (Table 2). Acetolysis experiment monitored CE-LIF (Nigou et al, 2000)
20 indicated that both LAM exhibited a mannan core with a similar degree of branching (Table
21 2). Finally, per-*O*-methylation analysis revealed the same glycosidic linkages for both LAMc
22 and LAMg and in similar relative abundance (Table 2).

23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47 Altogether, these data demonstrated that LAM structure was not altered by the in-gel
48 purification procedure and that it could be investigated after elution from the gel using an
49 adapted set of sensitive procedures.
50
51
52
53
54
55
56
57
58
59
60

1
2
3 *Identification and structural characterization of a new LAM-like polysaccharide in M.*
4
5 *smegmatis mc²155*

6
7 Mycobacterial lipoglycans and glycans are a complex family of biosynthetically related
8 molecules (Angala et al, 2014; Briken et al, 2004; Gilleron et al, 2008; Mishra et al, 2011;
9 Nigou et al, 2003). They consist in LM, LAM and their MPI anchor-free counterparts
10 mannan and AM. Interestingly, we identified in the crude fraction of lipoglycans and glycans
11 extracted from *M. smegmatis* a previously uncharacterized compound (named X) that could
12 be revealed by DOC-PAGE migration after alkali-treatment of the extract (Figure 4A)
13 (Skovierova et al, 2009). Monosaccharide analysis by GC after octanolysis and
14 trimethylsilylation showed that the crude fraction contained only D-Ara and D-Man,
15 suggesting that compound X harboured a structure reminiscent to that of lipoglycans or
16 glycans. Compound X was purified using the in-gel procedure described above and 100 µg
17 could be obtained by loading 2 mg of the extract. This amount was sufficient to investigate its
18 structure. Total acid hydrolysis followed by CE analysis indicated that compound X was
19 composed of Ara and Man, with an Ara/Man ratio of 0.25. This ratio value was 3 to 4 fold
20 lower than that of *M. smegmatis* LAM (Table 2). MALDI-TOF mass spectrometry analysis of
21 permethylated compound X showed a set of peaks centred around m/z 6000 and separated by
22 204 mass units corresponding to permethylated anhydro-Man residues (Figure 4B). As
23 previously observed for the other mycobacterial glycans and lipoglycans, compound X
24 exhibited a Gaussian distribution of glycoforms with the major ones showing a molecular
25 mass around 5 kDa, a value three times lower than that of LAM or AM. Compound X thus
26 contained an average of 9 Ara and 23 Man units, indicating the presence of an arabinan
27 domain much shorter than that of *M. smegmatis* LAM composed of a mean of ~55-70 Ara
28 and ~20-35 Man units (Angala et al, 2014; Kaur et al, 2014; Khoo et al, 1996; Mishra et al,
29 2011). Per-*O*-methylation analysis revealed the presence in compound X of 6-Man_p, 2,6-

1
2
3 Manp, t-Manp, 5-Araf, 3,5-Araf, 2-Araf and t-Araf residues, as observed in LAM (Table 2)
4
5 (Gilleron et al, 1997). However, in agreement with other chemical analyses, the relative
6
7 amount of arabinose residues was much lower in compound X than in LAM and consisted
8
9 mainly in 5-Araf units. In order to gain insight into the organisation of its arabinan domain,
10
11 10 µg of compound X were hydrolyzed with endoarabinanase from *Cellulomonas gelida* (Lee
12
13 et al, 2006; Shi et al, 2006), followed by the analysis of the released products by CE after
14
15 labelling by APTS or by MALDI-TOF MS after permethylation. As shown in Figure 5A,
16
17 digestion of compound X yielded the characteristic products previously described for LAM
18
19 (Shi et al, 2006), assigned to linear di-arabinofuranosides (Ara₂, peak II): Araf-α(1→5)-Araf-
20
21 α(1→, tetra-arabinofuranosides (Ara₄, peak III): Araf-β(1→2)-Araf-α(1→5)-Araf-α(1→5)-
22
23 Araf-α(1→ and biantennary hexa-arabinofuranosides (Ara₆, peak IV): [Araf-β(1→2)-Araf-
24
25 α(1-]₂→3 and →5)-Araf-α(1→5)-Araf-α(1→ motifs. This assignment was further confirmed
26
27 by MALDI-TOF MS analysis in the positive ion mode of the permethylated products (Figure
28
29 4B). The peaks at *m/z* 709.2 and *m/z* 1029.4 were assigned to (M+Na)⁺ ions corresponding to
30
31 the motifs Ara₄ and Ara₆ respectively (Figure 5B). Acetolysis experiment monitored by CE-
32
33 LIF (Nigou et al, 2000) showed a ratio of Manp-α(1→2)-Man to Man residues of 0.75:1
34
35 (Figure 5C), indicating a degree of branching of the mannan domain of 43% (Table 2). This
36
37 value is lower than that determined for *M. smegmatis* LAM (60% ± 5%) (Figure 5C; Table
38
39 2). The set of structural data obtained by chemical analyses were confirmed by NMR analysis
40
41 of the entire molecule. Indeed, the anomeric region of the 2D ¹H-¹³C HMQC spectrum of
42
43 compound X exhibited cross peaks assigned to 6-Manp (VI; δ_{H-1} 4.72; δ_{C-1} 97.8), 2,6-Manp
44
45 (VII; δ_{H-1} 4.92; δ_{C-1} 96.7), t-Manp (IV; δ_{H-1} 4.86; δ_{C-1} 97.8), 5-Araf (II; δ_{H-1} 4.89; δ_{C-1} 106.1),
46
47 3,5-Araf (I; δ_{H-1} 4.91; δ_{C-1} 106.0), and t-Araf (V; δ_{H-1} 4.95; δ_{C-1} 99.3) residues (Figure 6). In
48
49 agreement with chemical analyses, the signals corresponding to Ara units in compound X
50
51 spectrum (Figure 6B) were of much lower relative intensity that those recorded in LAM
52
53
54
55
56
57
58
59
60

1
2
3 spectrum (Figure 6A). In addition the cross peak corresponding to 2-Araf units (III; δ_{H-1} 4.98;
4
5 δ_{C-1} 106.9) present in lateral arabinan chains was very faint. Of note the signal corresponding
6
7 to 6-Man_p units in compound X showed a higher relative abundance, in agreement with a
8
9 weaker degree of branching of the mannan domain.
10
11 Altogether, these analyses allowed us to propose a structural model for compound X as
12
13 depicted in Figure 7.
14
15
16
17

18 Conclusion

19
20 We herein developed a methodology to efficiently purify lipoglycans by sodium
21
22 deoxycholate-PAGE that can be applied on a small amount of mycobacterial cells biomass
23
24 (10-50 mg). The resulting tens of micrograms of purified lipoglycans are sufficient to
25
26 undertake structural analyses of carbohydrate domains by a combination of highly sensitive
27
28 analytical procedures, involving cryoprobe NMR analysis of intact macromolecules and
29
30 chemical degradations monitored by gas chromatography and capillary electrophoresis. This
31
32 glycomic approach was successfully applied to the purification and structural characterization
33
34 of a newly identified polysaccharide in the model fast-growing species *M. smegmatis*. This
35
36 polysaccharide shows a structure reminiscent to that of LAM, bearing all the structural motifs
37
38 identified in this complex macromolecule, but with shorter arabinan and less branched
39
40 mannan domains. However, its biosynthetic origin remains unclear. Since it was isolated after
41
42 mild alkaline treatment, we investigated whether it might contain a MPI anchor in its native
43
44 state. Growing bacteria with *myo*-Inositol-(2-³H) resulted in a radiolabelling of LAM and LM
45
46 as previously described (Hunter & Brennan, 1990), but not of compound X, suggesting that
47
48 the structure of the latter does not contain *myo*-Inositol. A LM-like molecule based on a
49
50 glucuronosyl diacylglycerol, instead of a MPI, anchor was previously described in
51
52 corynebacteria (Tatituri et al, 2007), which along mycobacteria belongs to the suborder
53
54
55
56
57
58
59
60

1
2
3 *Corynebacterineae*. Disruption in *Corynebacterium glutamicum* of the *pimB/mgtA* gene,
4
5 coding for the α -mannosyltransferase involved in the addition of the first mannosyl unit on
6
7 the glucuronosyl residue, was found to result in the abrogation of the LM-like biosynthesis
8
9 (Tatituri et al, 2007). Interestingly, the glycolipid 3-(*O*- α -D-glucuronopyranosyl)-1,2-diacyl-
10
11 *sn*-glycerol, which corresponds to the free lipid anchor of this LM-like molecule, was also
12
13 characterized in *M. smegmatis* (Wolucka et al, 1993). However, analysis of a *M. smegmatis*
14
15 mutant disrupted for the orthologue of *pimB/mgtA* gene showed that compound X was still
16
17 produced, indicating that the structure of the latter is not likely to be based on a glucuronosyl
18
19 diacylglycerol lipid anchor. Understanding the biosynthetic origin of this newly identified
20
21 polysaccharide thus requires further investigations. Its identification asks again the question
22
23 of the biosynthetic link between lipoglycans on the one hand and their lipid-free glycan
24
25 counterparts on the other hand. Whether the latter are generated by action of an endogenous
26
27 endoglycosidase (α -mannosidase) still remains an open question (Pitarque et al, 2008).
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Material and Methods

Bacterial strains and culture conditions

M. smegmatis mc²155 was grown in Middlebrook 7H9 medium supplemented with ADC and 0.05% Tween 80.

Lipoglycans isolation and purification by the conventional protocol

LAM and LM were purified from *M. smegmatis* mc²155 as described previously (Gilleron et al, 1997; Nigou et al, 1997). Briefly, mycobacteria were delipidated using CHCl₃/CH₃OH 2:1 v/v, followed by CHCl₃/CH₃OH 1:1 v/v for 2 h at 50°C. The delipidated mycobacteria were then extracted 3 times in 50% ethanol at 50°C for 2h. The ethanol/water extracts were pooled, dried and submitted to digestion by DNase (D4263, Sigma-Aldrich), RNase (R4875, Sigma-Aldrich), α -amylase (A6814, Sigma-Aldrich) and a cocktail of proteases [α -chymotrypsin (C4129, Sigma-Aldrich), *Streptomyces griseus* proteases (P8811, Sigma-Aldrich) and trypsin, (T3914, Sigma-Aldrich)] to remove nucleic acids, α -glucan and proteins. The digested solution was dialysed against dH₂O and dried, resulting in a fraction containing glycans and lipoglycans. This fraction was then either used for DOC-PAGE purification of lipoglycans or used for conventional purification. For conventional purification, the fraction containing glycans and lipoglycans was loaded onto an octyl-sepharose column, eluted first with 10% isopropanol and then with 80% isopropanol at a flow rate of 0.3 mL/min to separate glycans and lipoglycans. After dialysis against dH₂O of the 80% isopropanol fraction, the lipoglycans were then loaded into a gel filtration column Biogel P100 (1.5 x 50 cm) in a DOC buffer (Tris-HCl 10 mM, NaCl 200 mM, 1mM EDTA, 0,25% sodium deoxycholate, pH 8.0) at a flow rate of 1 mL/min to separate LAM from LM. The fractions were analysed by SDS-

1
2
3 PAGE and the fraction containing LAM or LM were pooled, dialysed against dH₂O, dried
4
5 and precipitated with ethanol.
6
7

8
9
10 *Lipoglycans labelling with Lucifer Yellow for PAGE-UV*

11
12 Glycans and lipoglycans, typically 1 to 10 µg, were submitted to oxidation by sodium
13
14 metaperiodate 1 M for 1 h at 4°C in the dark (Pitarque et al, 2005). The reaction was
15
16 quenched by addition of sodium bisulphite 80 mM and settled for 5 min at room temperature.
17
18 Oxidized glycans and lipoglycans were then labelled for 10 min at 37°C using a 5 mg/mL
19
20 solution of Lucifer Yellow (861502, Aldrich) solubilized in dH₂O. Excess dye was removed
21
22 by micro-dialysis against dH₂O for 2 h at room temperature. The labelled glycans and
23
24 lipoglycans were then dried under speed-vacuum and submitted to DOC-PAGE migration.
25
26
27

28
29
30 *DOC-PAGE passive elution of lipoglycans*

31
32 Lucifer Yellow-labelled and unlabelled glycans and lipoglycans samples were loaded onto a
33
34 DOC-PAGE gel 15% and submitted to electrophoresis. After migration, the gel was placed
35
36 onto a glass surface and labelled lipoglycans were detected under UV (λ 254 nm). By
37
38 assuming that the migration of the labelled lipoglycans was similar to that of the non-labelled
39
40 one, the bands corresponding to LAM, LM or glycans were excised from the gel. The band
41
42 was then cut into 0.5 mm sized pieces and placed into 1.5 mL microtubes. Lipoglycans were
43
44 then passively eluted using a solution of triethylamine (TEA) 10% in dH₂O at 37°C overnight
45
46 followed by 2 washes of 2 h at 37°C with fresh elution solution. The solution containing
47
48 passively eluted lipoglycans were pooled, passed through a 0.22 µm non-pyrogenic filter and
49
50 dried under speed-vacuum. The eluted lipoglycans were then re-suspended with dH₂O and
51
52 submitted to dialysis against dH₂O for 2 h at room temperature prior to use for structural
53
54
55
56
57
58
59
60 determination.

1
2
3 *Determination of the absolute configuration of the monosaccharides*
4

5 1 mg of the lipoglycans and glycans crude fraction was octanolysed by 2M trifluoroacetic in
6
7 (*R*)-(-)-2-Butanol for 16 hr at 110°C (Gerwig et al, 1978). Octyl-glycosides were dried and
8
9 trimethylsilylated by treating with hexamethyldisilazane/chlorotrimethylsilylamine/pyridine,
10
11 4:2:1, v/v/v, for 15 min at room temperature. The resulting derivatives were solubilized in
12
13 cyclohexane before analysis by GC and comparison with standards of D-Ara, L-Ara, D-Man,
14
15 L-Man submitted to the same procedure.
16
17

18
19
20
21 *Monosaccharide composition and mannan core branching degree determination by capillary*
22
23 *electrophoresis*
24

25 These analyses were performed as previously described (Nigou et al, 2000). Briefly, for
26
27 monosaccharide composition, lipoglycans or glycans were submitted to strong acid
28
29 hydrolysis with 2 M trifluoroacetic acid at 110°C for 2 h and then dried under speed-vacuum.
30
31 For mannan core degree determination, lipoglycans or glycans were treated with an acetic
32
33 anhydride/acetic acid/sulfuric acid (10:10:1, v/v/v) mixture for 3 h at 40°C. After extraction
34
35 with water, acetolysis products were deacetylated with a methanol/20% aqueous ammonia
36
37 solution (1:1, v/v) at 37°C for 18 h and dried under a stream of nitrogen. The resulting mono-
38
39 or oligo-saccharides were derivatized for 90 min at 55°C using a solution of 1-aminopyrene-
40
41 3,6,8-trisulfonate (APTS) at 0.2 M in 15% acetic acid and 1 M sodium cyanoborohydride
42
43 solution dissolved in tetrahydrofuran. The APTS-labelled monosaccharides were suspended
44
45 in dH₂O and subjected to capillary electrophoresis. Analysis was performed on a P/ACE 5000
46
47 capillary electrophoresis in reverse polarity (Beckman Coulter) equipped with a laser-induced
48
49 fluorescence (LIF) detector with a 4 mW argon-ion laser (λ_{ext} 488 nm, λ_{em} 520 nm).
50
51 Separations were performed using an uncoated fused-silica capillary column (50 μ m internal
52
53
54
55
56
57
58
59
60

1
2
3 diameter, 40 cm effective length) and 30 mM triethylamine 1% (w/v) acetic acid running
4
5 buffer at pH 3.5, at 25°C with an applied voltage of 20 kV.
6
7

8
9
10 *Digestion of lipoglycans by the endoarabinanase from Cellulomonas gelida*

11
12 10 µg of dried purified lipoglycans or glycans were treated with 4 µL of endoarabinanase
13
14 from *Cellulomonas gelidas* (suspended at 5 mg/mL in Tris-HCl 10 mM pH 7.0 for 12h at
15
16 37°C) prepared as described in (Lee et al, 2006; Shi et al, 2006). The released products were
17
18 purified through a 3 kDa microcon® centrifugal filter unit and analysed by CE-LIF after
19
20 tagging by APTS (Nigou et al, 2000) or MALDI-TOF MS in positive ion mode after
21
22 permethylation (Morelle et al, 2004).
23
24

25
26
27 *Glycosidic linkage analysis*

28
29 Glycosyl linkage composition was performed according to the modified procedure of
30
31 Ciucanu and Kerek (Ciucanu & Kerek, 1984). The per-O-methylated lipoglycans or glycans
32
33 were hydrolyzed using 500 µl of 2 M trifluoroacetic acid at 110°C for 2 h, reduced using 350
34
35 µl of a 10mg/ml solution of NaBD₄ (NH₄OH 1M / C₂H₅OH, 1:1, v/v) and per-O-acetylated
36
37 using 300 µl of acetic anhydride for 1 h at 110°C. The resulting alditol acetates were
38
39 solubilized in cyclohexane before analysis by GC and GC/MS.
40
41
42
43
44

45
46 *MALDI-TOF analysis*

47
48 Deacylated lipoglycans were analysed on a Voyager DE-STR MALDI-TOF instrument
49
50 (PerSeptive Biosystems, Framingham, MA) using linear mode detection. Mass spectra were
51
52 recorded in negative ion mode using 300 ns time delay, with a grid of 80% of full
53
54 acceleration voltage (25kV) and a guide wire voltage of 0.15%. 2,5 dihydroxybenzoic acid
55
56 was used as the matrix at a concentration of 10 mg/mL in 50% ethanol in dH₂O.
57
58
59
60

1
2
3 Methylated oligosaccharides were analysed on a 4700 Proteomics Analyser (with TOF-TOF
4
5 Optics, Applied Biosystems) using the reflectron mode. Ionization was effected by irradiation
6
7 with pulsed UV light (355 nm) from an Nd:YAG laser. Samples were analysed by the
8
9 instrument operating at 20 kV in the positive ion modes using an extraction delay time set at
10
11 20 ns. Typically, spectra from 1,000 to 2,500 laser shots were summed to obtain the final
12
13 spectrum.
14
15

16 17 18 *NMR analysis* 19

20
21 NMR spectra were recorded at 298K on a Bruker Avance 600 MHz spectrometer equipped
22
23 with a cryogenic probe TCI (Bruker Biospin, Germany). Purified lipoglycans or glycans were
24
25 exchanged in D₂O (D, 99.97% from Euriso-top, Saint-Aubin, France), with intermediate
26
27 lyophilisation, and then dissolved in 0.5 ml D₂O. Samples were analysed in a 200 x 5 mm
28
29 UL-5 NMR tubes (Euriso-Top, Saint-Aubin, France). Proton and carbon chemical shifts are
30
31 expressed in part per million and referenced relative to internal acetone signals at $\delta_{\text{H/TMS}}$
32
33 2.225 and $\delta_{\text{C/TMS}}$ 31.45 ppm.
34
35

36
37 All 2D NMR data sets were recorded without sample spinning. ¹H-¹³C correlation spectra
38
39 were acquired in the echo/anti-echo-TPPI gradient selection mode (256 data points) recorded
40
41 in the proton-detected mode using the “hsqcetgpsisp2” sequence from the Topspin v2.1
42
43 software (Bruker Biospin) (Schleucher et al, 1994). The GARP sequence (Shaka et al, 1985)
44
45 at the carbon frequency was used as a composite pulse decoupling during acquisition.
46

47
48 The HSQC spectra were recorded with a spectral width of 25,000 Hz in ¹³C and 4807 Hz in
49
50 ¹H dimensions in order to collect a 1024 x 256 (TPPI) point data matrix with 2 scans per t₁
51
52 value expanded to 2048 x 1024 by zero filling. The relaxation delay was 2.0 s. A sine bell
53
54 window shifted by $\pi/2$ was applied in both dimensions.
55
56
57
58
59
60

Acknowledgments

Dr Pascal Ludwiczak (IPBS) is gratefully acknowledged for his initial work on lipoglycan extraction from gels. We thank Dr. Thomas Lerner (MRC-NIMR) and Ms Deborah Hunt (MRC-NIMR) for careful reading of the manuscript.

Conflict of interest statement

None declared

Funding

This work was supported by grants from The National Institute of Allergy and Infectious Diseases/National Institutes of Health (AI064798 and AI018357). The IPBS NMR equipment was financed by the French Research Ministry, CNRS, Université Paul Sabatier, the Région Midi-Pyrénées and the European structural funds.

Abbreviations

CE-LIF, capillary electrophoresis monitored by laser-induced fluorescence; DOC: sodium deoxycholate; HMQC, heteronuclear multiple quantum coherence; LAM, lipoarabinomannan; LM, lipomannan; LY, Lucifer Yellow; MALDI-TOF, matrix-assisted laser desorption ionization time of flight; MS, mass spectrometry; PAGE, polyacrylamide gel electrophoresis.

REFERENCES

- 1
2
3
4
5
6 Angala SK, Belardinelli JM, Huc-Claustre E, Wheat WH, Jackson M (2014) The cell
7 envelope glycoconjugates of *Mycobacterium tuberculosis*. *Crit Rev Biochem Mol Biol* 49:
8 361-399
9
- 10 Berg S, Kaur D, Jackson M, Brennan PJ (2007) The glycosyltransferases of *Mycobacterium*
11 *tuberculosis* - roles in the synthesis of arabinogalactan, lipoarabinomannan, and other
12 glycoconjugates. *Glycobiology* 17: 35-56R
13
- 14 Briken V, Porcelli SA, Besra GS, Kremer L (2004) Mycobacterial lipoarabinomannan and
15 related lipoglycans: from biogenesis to modulation of the immune response. *Mol Microbiol*
16 53: 391-403
17
- 18 Ciucanu I, Kerek F (1984) A simple and rapid method for the permethylation of
19 carbohydrates. *Carbohydr Res* 131: 209-217
20
- 21 Claverol S, Burllet-Schiltz O, Gairin JE, Monsarrat B (2003) Characterization of protein
22 variants and post-translational modifications: ESI-MSn analyses of intact proteins eluted
23 from polyacrylamide gels. *Mol Cell Proteomics* 2: 483-493
24
- 25 De Castro C, Parrilli M, Holst O, Molinaro A (2010) Microbe-associated molecular patterns
26 in innate immunity: Extraction and chemical analysis of gram-negative bacterial
27 lipopolysaccharides. *Methods Enzymol* 480: 89-115
28
- 29 Deng L, Mikusova K, Robuck KG, Scherman M, Brennan PJ, McNeil MR (1995)
30 Recognition of multiple effects of ethambutol on metabolism of mycobacterial cell envelope.
31 *Antimicrob Agents Chemother* 39: 694-701
32
- 33 Galvani M, Hamdan M, Righetti PG (2000) Investigating the reaction of a number of gel
34 electrophoresis cross-linkers with beta-lactoglobulin by matrix assisted laser
35 desorption/ionization-mass spectrometry. *Electrophoresis* 21: 3684-3692
36
- 37 Gerwig GJ, Kamerling JP, Vliegthart JFG (1978) Determination of the D and L
38 configuration of neutral monosaccharides by high-resolution capillary G.L.C. *Carbohydr Res*
39 62: 349-357
40
- 41 Gilleron M, Himoudi N, Adam O, Constant P, Venisse A, Riviere M, Puzo G (1997)
42 *Mycobacterium smegmatis* phosphoinositols-glyceroarabinomannans. Structure and
43 localization of alkali-labile and alkali-stable phosphoinositides. *J Biol Chem* 272: 117-124
44
- 45 Gilleron M, Jackson M, Nigou J, Puzo G (2008) Structure, Biosynthesis, and Activities of the
46 phosphatidyl-myo-Inositol-Based Lipoglycans. In *The mycobacterial Cell Envelope*, Daffe
47 M, Reytrat J (eds), pp 75-105. Washington DC: ASM Press
48
- 49 Hardy E, Pupo E, Castellanos-Serra L, Reyes J, Fernandez-Patron C (1997) Sensitive reverse
50 staining of bacterial lipopolysaccharides on polyacrylamide gels by using zinc and imidazole
51 salts. *Anal Biochem* 244: 28-32
52
- 53
54
55
56 Hunter SW, Brennan PJ (1990) Evidence for the presence of a phosphatidylinositol anchor on
57 the lipoarabinomannan and lipomannan of *Mycobacterium tuberculosis*. *J Biol Chem* 265:
58 9272-9279
59
60

- 1
2
3 Kaur D, Angala SK, Wu SW, Khoo KH, Chatterjee D, Brennan PJ, Jackson M, McNeil MR
4 (2014) A Single Arabinan Chain Is Attached to the Phosphatidylinositol Mannosyl Core of
5 the Major Immunomodulatory Mycobacterial Cell Envelope Glycoconjugate,
6 Lipoarabinomannan. *J Biol Chem* **289**: 30249-56
7
- 8 Khoo KH, Douglas E, Azadi P, Inamine JM, Besra GS, Mikusova K, Brennan PJ, Chatterjee
9 D (1996) Truncated structural variants of lipoarabinomannan in ethambutol drug-resistant
10 strains of *Mycobacterium smegmatis*. Inhibition of arabinan biosynthesis by ethambutol. *The*
11 *J Biol Chem* **271**: 28682-28690
12
- 13 Kurth J, Stoffel W (1990) A facile method for the isolation and preparation of proteins and
14 peptides for sequence analysis in the picomolar range. *Biol Chem Hoppe Seyler* **371**: 675-685
15
- 16 Laneelle MA, Nigou J, Daffe M (2015) Lipid and lipoarabinomannan isolation and
17 characterization. *Methods Mol Biol* **1285**: 77-103
18
- 19 Lee A, Wu SW, Scherman MS, Torrelles JB, Chatterjee D, McNeil MR, Khoo KH (2006)
20 Sequencing of oligoarabinosyl units released from mycobacterial arabinogalactan by
21 endogenous arabinanase: identification of distinctive and novel structural motifs.
22 *Biochemistry* **45**: 15817-15828
23
- 24 Makarov V, Manina G, Mikusova K, Mollmann U, Ryabova O, Saint-Joanis B, Dhar N,
25 Pasca MR, Buroni S, Lucarelli AP, Milano A, De Rossi E, Belanova M, Bobovska A,
26 Dianiskova P, Kordulakova J, Sala C, Fullam E, Schneider P, McKinney JD, Brodin P,
27 Christophe T, Waddell S, Butcher P, Albrethsen J, Rosenkrands I, Brosch R, Nandi V,
28 Bharath S, Gaonkar S, Shandil RK, Balasubramanian V, Balganesht T, Tyagi S, Grosset J,
29 Riccardi G, Cole ST (2009) Benzothiazinones kill *Mycobacterium tuberculosis* by blocking
30 arabinan synthesis. *Science* **324**: 801-804
31
32
- 33 Mishra AK, Driessen NN, Appelmeik BJ, Besra GS (2011) Lipoarabinomannan and related
34 glycoconjugates: structure, biogenesis and role in *Mycobacterium tuberculosis* physiology
35 and host-pathogen interaction. *FEMS Microbiol Rev* **35**: 1126-1157
36
- 37 Morelle W, Slomianny MC, Diemer H, Schaeffer C, van Dorselaer A, Michalski JC (2004)
38 Fragmentation characteristics of permethylated oligosaccharides using a matrix-assisted laser
39 desorption/ionization two-stage time-of-flight (TOF/TOF) tandem mass spectrometer. *Rapid*
40 *Commun Mass Spectrom* **18**: 2637-2649
41
- 42 Nigou J, Gilleron M, Brando T, Puzo G (2004) Structural analysis of mycobacterial
43 lipoglycans. *Appl Biochem Biotechnol* **118**: 253-267
44
- 45 Nigou J, Gilleron M, Brando T, Vercellone A, Puzo G (1999) Structural definition of
46 arabinomannans from *Mycobacterium bovis* BCG. *Glycoconj J* **16**: 257-264
47
- 48 Nigou J, Gilleron M, Cahuzac B, Bounery JD, Herold M, Thurnher M, Puzo G (1997) The
49 phosphatidyl-myo-inositol anchor of the lipoarabinomannans from *Mycobacterium bovis*
50 bacillus Calmette Guerin. Heterogeneity, structure, and role in the regulation of cytokine
51 secretion. *J Biol Chem* **272**: 23094-23103
52
- 53 Nigou J, Gilleron M, Puzo G (2003) Lipoarabinomannans: from structure to biosynthesis.
54 *Biochimie* **85**: 153-166
55
56
57
58
59
60

1
2
3 Nigou J, Vercellone A, Puzo G (2000) New structural insights into the molecular deciphering
4 of mycobacterial lipoglycan binding to C-type lectins: lipoarabinomannan glycoform
5 characterization and quantification by capillary electrophoresis at the subnanomole level. *J*
6 *Mol Biol* 299: 1353-1362
7

8 Ortalo-Magne A, Dupont MA, Lemassu A, Andersen AB, Gounon P, Daffe M (1995)
9 Molecular composition of the outermost capsular material of the tubercle bacillus.
10 *Microbiology* 141 (Pt 7): 1609-1620
11

12 Pitarque S, Herrmann JL, Duteyrat JL, Jackson M, Stewart GR, Lecointe F, Payre B,
13 Schwartz O, Young DB, Marchal G, Lagrange PH, Puzo G, Gicquel B, Nigou J, Neyrolles O
14 (2005) Deciphering the molecular bases of *Mycobacterium tuberculosis* binding to the lectin
15 DC-SIGN reveals an underestimated complexity. *Biochem J* 392: 615-624
16

17 Pitarque S, Larrouy-Maumus G, Payre B, Jackson M, Puzo G, Nigou J (2008) The
18 immunomodulatory lipoglycans, lipoarabinomannan and lipomannan, are exposed at the
19 mycobacterial cell surface. *Tuberculosis (Edinb)* 88: 560-565
20

21 Pupo E, Aguila A, Santana H, Nunez JF, Castellanos-Serra L, Hardy E (1999) Mice
22 immunization with gel electrophoresis-micropurified bacterial lipopolysaccharides.
23 *Electrophoresis* 20: 458-461
24

25 Pupo E, Lopez CM, Alonso M, Hardy E (2000) High-efficiency passive elution of bacterial
26 lipopolysaccharides from polyacrylamide gels. *Electrophoresis* 21: 526-530
27

28 Pupo E, Phillips NJ, Gibson BW, Apicella MA, Hardy E (2004) Matrix-assisted laser
29 desorption/ionization-time of flight-mass spectrometry of lipopolysaccharide species
30 separated by slab-polyacrylamide gel electrophoresis: high-resolution separation and
31 molecular weight determination of lipooligosaccharides from *Vibrio fischeri* strain HMK.
32 *Electrophoresis* 25: 2156-2164
33

34 Raetz CR, Whitfield C (2002) Lipopolysaccharide endotoxins. *Annu Rev Biochem* 71: 635-
35 700
36

37 Ray A, Cot M, Puzo G, Gilleron M, Nigou J (2013) Bacterial cell wall macroamphiphiles:
38 pathogen-/microbe-associated molecular patterns detected by mammalian innate immune
39 system. *Biochimie* 95: 33-42
40

41 Schleucher J, Schwendinger M, Sattler M, Schmidt P, Schedletzky O, Glaser SJ, Sorensen
42 OW, Griesinger C (1994) A general enhancement scheme in heteronuclear multidimensional
43 NMR employing pulsed field gradients. *J Biomol NMR* 4: 301-306
44

45 Shaka AJ, Barker PB, Freeman R (1985) Computer-optimized decoupling scheme for
46 wideband applications and low level operation. *J Magn Reson* 64: 547-552
47

48 Shi L, Berg S, Lee A, Spencer JS, Zhang J, Vissa V, McNeil MR, Khoo KH, Chatterjee D
49 (2006) The carboxy terminus of EmbC from *Mycobacterium smegmatis* mediates chain
50 length extension of the arabinan in lipoarabinomannan. *J Biol Chem* 281: 19512-19526
51

52 Skovierova H, Larrouy-Maumus G, Zhang J, Kaur D, Barilone N, Kordulakova J, Gilleron
53 M, Guadagnini S, Belanova M, Prevost MC, Gicquel B, Puzo G, Chatterjee D, Brennan PJ,
54 Nigou J, Jackson M (2009) AftD, a novel essential arabinofuranosyltransferase from
55 mycobacteria. *Glycobiology* 19: 1235-1247
56
57
58
59
60

1
2
3 Tatituri RV, Illarionov PA, Dover LG, Nigou J, Gilleron M, Hitchen P, Krumbach K, Morris
4 HR, Spencer N, Dell A, Eggeling L, Besra GS (2007) Inactivation of *Corynebacterium*
5 *glutamicum* NCgl0452 and the role of MgtA in the biosynthesis of a novel mannosylated
6 glycolipid involved in lipomannan biosynthesis. *J Biol Chem* 282: 4561-4572
7

8
9 Torrelles JB, Schlesinger LS (2010) Diversity in *Mycobacterium tuberculosis* mannosylated
10 cell wall determinants impacts adaptation to the host. *Tuberculosis (Edinb)* 90: 84-93

11
12 Venisse A, Berjeaud JM, Chaurand P, Gilleron M, Puzo G (1993) Structural features of
13 lipoarabinomannan from *Mycobacterium bovis* BCG. Determination of molecular mass by
14 laser desorption mass spectrometry. *J Biol Chem* 268: 12401-12411

15
16 Vergne I, Gilleron M, Nigou J (2015) Manipulation of the endocytic pathway and phagocyte
17 functions by *Mycobacterium tuberculosis* lipoarabinomannan. *Front Cell Infect Microbiol* 4:
18 187

19
20 Wolucka BA, McNeil MR, Kalbe L, Cocito C, Brennan PJ (1993) Isolation and
21 characterization of a novel glucuronosyl diacylglycerol from *Mycobacterium smegmatis*.
22 *Biochim Biophys Acta* 1170: 131-136
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **Legends to figures**
4

5
6 **Figure 1: DOC-PAGE-UV detection of LY-labelled lipoglycans and work-flow showing**
7
8 **the strategy developed to purify lipoglycans from gel and to investigate their structure.**
9

10
11
12 **Figure 2: DOC-PAGE migration of native and LY-labelled LAM.**

13
14 5 µg of native (lane 1) and LY-labelled (lane 2) LAM were submitted to DOC-PAGE
15
16 migration. The same gel was revealed first by zinc sulphate negative staining (A) and then by
17
18 UV (B). std, protein ladder.
19
20

21
22
23 **Figure 3: MALDI-TOF MS spectra of deacylated LAM from *M. smegmatis* purified by**
24 **the conventional protocol (A) or by DOC-PAGE (B).**
25
26

27
28
29
30 **Figure 4: DOC-PAGE (A) and MALDI-TOF MS (B) analyses of compound X.**

31
32 A) 10 µg of the *M. smegmatis* glycans and lipoglycans fraction were loaded on the gel before
33
34 (lane 1) or after alkali-treatment (lane 2). The gel was revealed by zinc sulphate negative
35
36 staining. B) Linear MALDI-TOF MS spectrum in the positive mode of permethylated
37
38 compound X.
39
40

41
42
43 **Figure 5. Structural analysis of compound X arabinan (A, B) and mannan (C) domains.**

44
45 **A, B)** Digestion of arabinan domain (10 µg of compound X) by endoarabinanase from
46
47 *Cellulomonas gelida* monitored by CE-LIF (A) and MALDI-TOF MS analysis (B). A) I,
48
49 Internal Standard-APTS; II, Ara₂-APTS; III, Ara₄-APTS; IV, Ara₆-APTS; **a**, compound X;
50
51 **b**, LAM from *M. smegmatis*; **c**, co-injection. B) MALDI-TOF MS analysis in positive ion
52
53 mode of the permethylated products of hydrolysis of LAM (upper spectrum) and Compound
54
55
56
57
58
59
60

1
2
3 X (lower spectrum) (50 μg of each). Ions at m/z 709.3 correspond to Ara₄ motif and m/z
4
5 1029.4 correspond to Ara₆ motifs.
6


7 C) Mannan core degree of branching determined by acetolysis treatment (10 μg of compound
8
9 X). I, Ara-APTS; II, Man-APTS; III, Internal Standard-APTS; IV, Man- $\alpha(1\rightarrow2)$ -Man-APTS;
10
11 **a**, compound X; **b**, LAM from *M. smegmatis*; **c**, co-injection.
12
13
14
15

16 **Figure 6: 2D ^1H - ^{13}C HMQC NMR spectra expansion ($\delta^1\text{H}$: 4.60-5.15 and ^{13}C : 97.0-**
17
18 **113.0) of *M. smegmatis* LAM (A) and compound X (B) in D_2O at 298K.**
19

20 I : 3,5- α -Araf; II : 5- α -Araf; III : 2- α -Araf; IV : t- α -Manp ; V : t- β -Araf; VI : 6- α -Manp ;
21
22 VII : 2,6- α -Manp.
23
24
25
26

27 **Figure 7: Structural model of compound X from *M. smegmatis*.**

28
29 Compound X is comprised of a linear $\alpha(1\rightarrow6)$ -mannan domain that is substituted at almost
30
31 half of the *O*-2 positions by one Manp unit. It contains a short arabinan chain, possibly linked
32
33 at *O*-2 position of a Manp unit as recently described in LAM by Kaur *et al* (2014), that
34
35 harbours the Ara₄ and Ara₆ motifs typically found in LAM and AM.
36
37

38 ●, D-Manp; , D-Araf.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **Tables**
4
5

6 **Table 1: LAM recovery yield from DOC-PAGE using different elution conditions.**
7

8

Elution solution	Time of elution (hours)	% of detergent or organic solvent	Recovery (%) ^(a)
water	3	-	4.0±0.2
	12	-	20.0±2.5
SDS in water	3	0.1 %	6.2±0.5
	3	1%	9.1±0.4
DOC in water	3	0.1%	3.5±0.3
	3	1%	3.8±0.6
ACN in water	3	15%	3.1±0.5
	3	30%	3.4±0.8
	3	60%	2.5±1.1
TEA in water	3	0.1%	4.1±0.4
	3	1%	5.3±0.2
	3	5%	5.7±0.4
	3	10%	7.8±0.3
	12	10%	60.0±4.0

9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

^(a) Values were determined after loading 20 µg of LAM and represent the average of 2 independent experiments

Table 2: Structural features of LAMc, LAMg and Compound X.

	LAMc	LAMg	Compound X
Ara/Man ratio^(a)	0.8±0.1	0.9±0.05	0.25±0.08
Mannan degree of branching^(a) (%)	55±8	60±5	43±5
Glycosyl linkage^(b) (%)			
t-Araf	4.4	4.3	0.3
2-Araf	11.2	9.5	3.2
5-Araf	39.3	41.5	18.8
3,5-Araf	13.7	12.0	6.5
t-Manp	19.4	21.6	28.5
6-Manp	6.7	6.0	19.5
2,6-Manp	5.3	5.1	23.2
^(a) Monitored by CE-LIF			
^(b) Monitored by GC			




Figure 1: DOC-PAGE-UV detection of LY-labelled lipoglycans and work-flow showing the strategy developed to purify lipoglycans from gel and to investigate their structure.
234x82mm (300 x 300 DPI)

Peer Review




Figure 2: DOC-PAGE migration of native and LY-labelled LAM. 5 μg of native (lane 1) and LY-labelled (lane 2) LAM were submitted to DOC-PAGE migration. The same gel was revealed first by zinc sulphate negative staining (A) and then by UV (B). std, protein ladder.

69x43mm (600 x 600 DPI)

Review




Figure 3: MALDI-TOF MS spectra of deacylated LAM from *M. smegmatis* purified by the conventional protocol (A) or by DOC-PAGE (B).
124x151mm (300 x 300 DPI)




Figure 4: DOC-PAGE (A) and MALDI-TOF MS (B) analyses of compound X. A) 10 μg of the *M. smegmatis* glycans and lipoglycans fraction were loaded on the gel before (lane 1) or after alkali-treatment (lane 2). The gel was revealed by zinc sulphate negative staining. B) Linear MALDI-TOF MS spectrum in the positive mode of permethylated compound X.

169x89mm (300 x 300 DPI)

Review




Figure 5. Structural analysis of compound X arabinan (A, B) and mannan (C) domains. A, B) Digestion of arabinan domain (10 μg of compound X) by endoarabinanase from *Cellulomonas gelida* monitored by CE-LIF (A) and MALDI-TOF MS analysis (B). A) I, Internal Standard-APTS; II, Ara2-APTS; III, Ara4-APTS; IV, Ara6-APTS; a, compound X; b, LAM from *M. smegmatis*; c, co-injection. B) MALDI-TOF MS analysis in positive ion mode of the permethylated products of hydrolysis of LAM (upper spectrum) and Compound X (lower spectrum) (50 μg of each). Ions at m/z 709.3 correspond to Ara4 motif and m/z 1029.4 correspond to Ara6 motifs. C) Mannan core degree of branching determined by acetolysis treatment (10 μg of compound X). I, Ara-APTS; II, Man-APTS; III, Internal Standard-APTS; IV, Man- $\alpha(1\rightarrow2)$ Man-APTS; a, compound X; b, LAM from *M. smegmatis*; c, co-injection.

176x156mm (96 x 96 DPI)





Figure 6: 2D ^1H - ^{13}C HMQC NMR spectra expansion ($\delta^1\text{H}$: 4.60-5.15 and ^{13}C : 97.0- 113.0) of *M. smegmatis* LAM (A) and compound X (B) in D_2O at 298K.
 I : 3,5- α -Araf ; II : 5- α -Araf ; III : 2- α -Araf ; IV : t- α -Manp ; V : t- β -Araf ; VI : 6- α -Manp ; VII : 2,6- α -Manp.

164x388mm (600 x 600 DPI)



24 Figure 7: Structural model of compound X from *M. smegmatis*.

25 Compound X is comprised of a linear $\alpha(1\rightarrow6)$ -mannan domain that is substituted at almost half of the O-2
 26 positions by one Manp unit. It contains a short arabinan chain, possibly linked at O-2 position of a Manp unit
 27 as recently described in LAM by Kaur et al (2014), that harbours the Ara4 and Ara6 motifs typically found in
 28 LAM and AM. \circ , D-Manp; \square , D-Araf.

29 99x49mm (300 x 300 DPI)

30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Review