



HAL
open science

A Note on the Join of Varieties of Monoids with LI

Nathan Grosshans

► **To cite this version:**

| Nathan Grosshans. A Note on the Join of Varieties of Monoids with LI. 2021. hal-03183838v2

HAL Id: hal-03183838

<https://hal.science/hal-03183838v2>

Preprint submitted on 10 May 2021 (v2), last revised 27 Aug 2021 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A Note on the Join of Varieties of Monoids With \mathbf{LI}

Nathan Grosshans   

Fachbereich Elektrotechnik/Informatik, Universität Kassel, Kassel, Germany

Abstract

In this note, we give a characterisation in terms of identities of the join of \mathbf{V} with the variety of finite locally trivial semigroups \mathbf{LI} for several well-known varieties of finite monoids \mathbf{V} by using classical algebraic-automata-theoretic techniques. To achieve this, we use the new notion of essentially- \mathbf{V} stamps defined by Grosshans, McKenzie and Segoufin and show that it actually coincides with the join of \mathbf{V} and \mathbf{LI} precisely when some natural condition on the variety of languages corresponding to \mathbf{V} is verified.

This work is a kind of rediscovery of the work of J. C. Costa around 20 years ago from a rather different angle, since Costa's work relies on the use of advanced developments in profinite topology, whereas what is presented here essentially uses an algebraic, language-based approach.

2012 ACM Subject Classification Theory of computation \rightarrow Formal languages and automata theory; Theory of computation \rightarrow Algebraic language theory

Keywords and phrases Varieties of monoids, join, \mathbf{LI}

1 Introduction

One of the most fundamental problems appearing in finite automata theory is the one of *characterisation*: given some subclass of the class of regular languages, find out whether there is a way to characterise those languages using some class of finite objects. This problem is often linked to and motivated by the problem of *decidability*: given some subclass of the class of regular languages, find out whether there exists an algorithm testing the membership of any regular language in that subclass. The obvious approach to try to find a characterisation of a class of regular languages would be to look for properties shared by all the minimal finite automata of those languages. If we find such characterising properties, we can then ask whether they can be checked by an algorithm to answer the problem of decidability for this class of languages. However, one of the most fruitful approaches in face of those two problems has been the *algebraic approach*, in which we basically replace automata with morphisms into monoids: a language L over some alphabet Σ is then said to be recognised by a morphism φ into a monoid M if and only if L is the inverse image by φ of a subset of M . Under this notion of recognition, each language has a minimal morphism recognising it, the *syntactic morphism* into the *syntactic monoid* of that language, that are minimal under some notion of division. The fundamental result on which this algebraic approach relies is that a language is regular if and only if its syntactic monoid is finite. One can thus try to find a characterisation of some class of regular languages by looking at the algebraic properties of the syntactic monoids of these languages.

And many such characterisations that are decidable were indeed successfully obtained since Schützenberger's seminal work in 1965 [16]. His famous result, that really started the field of *algebraic automata theory*, states that the star-free regular languages are exactly those whose syntactic monoids are finite and aperiodic. Another important early result in that vein is the one of Simon [17] characterising the piecewise testable languages as exactly those having a finite \mathfrak{J} -trivial syntactic monoid. Eilenberg [11] was the first to prove that such algebraic characterisations actually come as specific instances of a general bijective correspondence between varieties of finite monoids and varieties of languages — classes

46 of, respectively, finite monoids and regular languages closed under some respective natural
 47 operations. Thus, a class of regular languages can indeed be characterised by the syntactic
 48 monoids of these languages, as soon as it verifies some nice properties. Eilenberg's result was
 49 later completed by Reiterman's theorem [15], that uses a notion of identities defined using
 50 profinite topology and states that a class of finite monoids is a variety of finite monoids if and
 51 only if it is defined by a set of profinite identities. Therefore, one can always characterise the
 52 variety of finite monoids associated to a variety of languages by a set of profinite identities
 53 and, additionally, this characterisation often leads to decidability, especially when this set is
 54 finite. Several research works have been conducted to characterise varieties of finite monoids
 55 or semigroups by profinite identities.

56 A kind of varieties of finite monoids or semigroups that has attracted many research
 57 efforts aiming for characterisations through identities are the varieties defined as the join
 58 of two other varieties. Given two varieties of finite monoids \mathbf{V} and \mathbf{W} , the *join of \mathbf{V}*
 59 *and \mathbf{W}* , denoted by $\mathbf{V} \vee \mathbf{W}$, is the smallest variety of finite monoids containing both \mathbf{V}
 60 and \mathbf{W} . One of the main motivations to try to understand $\mathbf{V} \vee \mathbf{W}$ is that the variety
 61 of languages corresponding to it by the Eilenberg correspondence, $\mathcal{L}(\mathbf{V} \vee \mathbf{W})$, is the one
 62 obtained by considering parallel computations of automata recognising languages from both
 63 $\mathcal{L}(\mathbf{V})$ and $\mathcal{L}(\mathbf{W})$, the varieties of languages corresponding to, respectively, \mathbf{V} and \mathbf{W} . This
 64 is a fundamental operation on automata, and while it is straightforward that $\mathcal{L}(\mathbf{V} \vee \mathbf{W})$
 65 is simply the smallest variety of languages containing both $\mathcal{L}(\mathbf{V})$ and $\mathcal{L}(\mathbf{W})$, this does
 66 not at all furnish a decidable characterisation of $\mathcal{L}(\mathbf{V} \vee \mathbf{W})$, let alone a set of identities
 67 defining $\mathbf{V} \vee \mathbf{W}$. Generally speaking, the problem of finding a set of identities defining
 68 $\mathbf{V} \vee \mathbf{W}$ is difficult (see [2, 21]), but it has been solved for many specific cases: have a look
 69 at [1, 3, 5, 20, 19, 6, 8, 9] for some examples.

70 In this paper, we give a general method to find a set of identities defining the join of an
 71 arbitrary variety of finite monoids \mathbf{V} and the *variety of finite locally trivial semigroups* **LI**, as
 72 soon as one has a set of identities defining \mathbf{V} and \mathbf{V} verifies some criterion. Joins of that sort
 73 have been studied quite a lot in the literature we mentioned in the previous paragraph (e.g.
 74 in [5, 19, 8, 9]), but while these works usually rely heavily on profinite topology with some
 75 in-depth understanding of the structure of the elements of the so-called free pro- \mathbf{V} monoids
 76 and free pro-**LI** semigroups, we present a method that reduces the use of profinite topology
 77 to the minimum and that relies mainly on algebraic and language-theoretic techniques. The
 78 variety **LI** is well-known to correspond to the class of languages for which membership only
 79 depends on bounded-length prefixes and suffixes of words. In [12], McKenzie, Segoufin and
 80 the author introduced the notion of *essentially- \mathbf{V}* stamps (surjective morphisms $\varphi: \Sigma^* \rightarrow M$
 81 for Σ an alphabet and M a finite monoid) to characterise the built-in ability that programs
 82 over monoids in \mathbf{V} have to treat separately some constant-length beginning and ending of a
 83 word. Informally said, a stamp is *essentially- \mathbf{V}* when it behaves like a stamp into a monoid
 84 of \mathbf{V} as soon as a sufficiently long beginning and ending of the input word has been fixed.
 85 Our method builds on two results, that we prove in this article.

- 86 1. The first result is a characterisation in terms of identities of the class **EV** of essentially- \mathbf{V}
 87 stamps given a set of identities E defining \mathbf{V} : a stamp is in **EV** if and only if it satisfies
 88 all identities $x^\omega y u z t^\omega = x^\omega y v z t^\omega$ for $u = v$ an identity in E and where x, y, z, t do appear
 89 neither in u nor in v .
- 90 2. The second result says that **EV** and $\mathbf{V} \vee \mathbf{LI}$ do correspond if and only if \mathbf{V} verifies some
 91 criterion, that can be formulated in terms of quotient-expressibility in $\mathcal{L}(\mathbf{V})$: any language
 92 $L \in \mathcal{L}(\mathbf{V})$ must, for an arbitrary choice of x, y , be such that the quotient $u^{-1} L v^{-1}$ for u
 93 and v long enough can be expressed as the quotient $(xu)^{-1} K (vy)^{-1}$ for some $K \in \mathcal{L}(\mathbf{V})$.

94 Using these results, we can find a set of identities defining the so-called *ne*-variety of stamps
 95 corresponding to $\mathbf{V} \vee \mathbf{LI}$ as soon as a set of identities defining \mathbf{V} is known by proving that \mathbf{V}
 96 verifies the criterion in point 2. Note that, for technical reasons, we work with the *ne*-variety
 97 of stamps corresponding to $\mathbf{V} \vee \mathbf{LI}$ rather than directly with the variety of finite semigroups
 98 $\mathbf{V} \vee \mathbf{LI}$, but this is not a problem since they correspond bijectively. We apply this method
 99 to reprove characterisations of the join of \mathbf{LI} with each of the well-known varieties of finite
 100 monoids \mathbf{R} , \mathbf{L} , \mathbf{J} and any variety of finite groups.

101 The author noticed after proving those results that his work actually forms a kind of
 102 rediscovery of the work of J. C. Costa in [8]. He defines an operator U associating to each set
 103 of identities E the exact same new set $U(E)$ of identities as in point 1. Costa then defines a
 104 property of cancellation for varieties of finite semigroups such that for any \mathbf{V} verifying it,
 105 $U(E)$ defines $\mathbf{V} \vee \mathbf{LI}$ for E defining \mathbf{V} . He finally uses this result to derive characterisations
 106 of $\mathbf{V} \vee \mathbf{LI}$ for all the cases we are treating in our paper and many more.

107 What is, then, the contribution of our article? In a nutshell, it does mainly use algebraic
 108 and language-theoretic techniques while Costa's work relies heavily on profinite topology. In
 109 our setting, once the stage is set, all proofs are quite straightforward without real difficulties
 110 and rely on classical language-theoretic characterisations of the varieties under consideration.
 111 This is to contrast with Costa's work, that for instance draws upon the difficult analysis of
 112 the elements of free pro- \mathbf{R} monoids by Almeida and Weil [4] to characterise $\mathbf{R} \vee \mathbf{LI}$.

113 **Organisation of the article.** Section 2 is dedicated to the necessary preliminaries. In
 114 Section 3, we recall the definition of essentially- \mathbf{V} stamps and prove the characterisation
 115 by identities of point 1 above. Section 4 is then dedicated to the necessary and sufficient
 116 criterion for \mathbf{EV} and $\mathbf{V} \vee \mathbf{LI}$ to correspond presented in point 2 and finally those results are
 117 applied to specific cases in Section 5. We finish with a short conclusion.

118 **2 Preliminaries**

119 We briefly introduce the mathematical material used in this paper. For the basics and the
 120 classical results of automata theory, we refer the reader to the two classical references of the
 121 domain by Eilenberg [10, 11] and Pin [13]. For definitions and results specific to varieties of
 122 stamps and associated profinite identities, see the articles by Straubing [18] and by Pin and
 123 Straubing [14]. We also assume some basic knowledge of topology.

124 **General notations.** Let $i \in \mathbb{N}$ be a natural number. We shall denote by $[i]$ the set of all
 125 natural numbers $n \in \mathbb{N}$ verifying $1 \leq n \leq i$.

126 **Words and languages.** Let Σ be a finite alphabet. We denote by Σ^* the set of all finite
 127 words over Σ . We also denote by Σ^+ the set of all finite non empty words over Σ , the empty
 128 word being denoted by ε . All our alphabets and words will always be finite, without further
 129 mention of this fact. Given some word $w \in \Sigma^*$, we denote its length by $|w|$ and the set of
 130 letters it contains by $\text{alph}(w)$. Given some $n \in \mathbb{N}$, we denote by $\Sigma^{\geq n}$, Σ^n and $\Sigma^{< n}$ the set of
 131 words over Σ of length, respectively, at least n , exactly n and less than n .

132 A *language over Σ* is a subset of Σ^* . A language is *regular* if it is recognised by a
 133 deterministic finite automaton. The *quotient of a language L over Σ relative to the words u*
 134 *and v over Σ* is the language, denoted by $u^{-1}Lv^{-1}$, of the words w such that $uwv \in L$.

135 **Monoids, semigroups and varieties.** A *semigroup* is a non-empty set equipped with an
 136 associative law that we will write multiplicatively. A *monoid* is a semigroup with an identity.
 137 An example of a semigroup is Σ^+ , the free semigroup over Σ . Similarly Σ^* is the free monoid
 138 over Σ . A *morphism* φ from a semigroup S to a semigroup T is a function from S to T such
 139 that $\varphi(xy) = \varphi(x)\varphi(y)$ for all $x, y \in S$. A morphism of monoids additionally requires that
 140 the identity is preserved. A semigroup T is a *subsemigroup* of a semigroup S if T is a subset
 141 of S and is equipped with the restricted law of S . Additionally the notion of submonoids
 142 requires the presence of the identity. A semigroup T *divides* a semigroup S if T is the image
 143 by a semigroup morphism of a subsemigroup of S . Division of monoids is defined in the same
 144 way. The *Cartesian (or direct) product* of two semigroups is simply the semigroup given by
 145 the Cartesian product of the two underlying sets equipped with the Cartesian product of
 146 their laws. An element s of a semigroup is *idempotent* if $ss = s$.

147 A *variety of finite monoids* is a non-empty class of finite monoids closed under Cartesian
 148 product and monoid division. A *variety of finite semigroups* is defined similarly. When
 149 dealing with varieties, we consider only finite monoids and semigroups, so we will drop the
 150 adjective finite when talking about varieties in the rest of this article.

151 **Varieties of stamps** Let $f: \Sigma^* \rightarrow \Gamma^*$ be a morphism from the free monoid over an alphabet
 152 Σ to the free monoid over an alphabet Γ , that we might call an *all-morphism*. We say that
 153 f is an *ne-morphism* (non-erasing morphism) whenever $f(\Sigma) \subseteq \Gamma^+$.

154 We call *stamp* a surjective morphism $\varphi: \Sigma^* \rightarrow M$ for Σ an alphabet and M a finite
 155 monoid. We say that a stamp $\varphi: \Sigma^* \rightarrow M$ *all-divides* (respectively *ne-divides*) a stamp
 156 $\psi: \Gamma^* \rightarrow N$ whenever there exists an *all-morphism* (respectively *ne-morphism*) $f: \Sigma^* \rightarrow \Gamma^*$
 157 and a surjective morphism $\alpha: \mathfrak{Im}(\psi \circ f) \rightarrow M$ such that $\varphi = \alpha \circ \psi \circ f$. The *direct product*
 158 of two stamps $\varphi: \Sigma^* \rightarrow M$ and $\psi: \Gamma^* \rightarrow N$ is the stamp $\varphi \times \psi: \Sigma^* \rightarrow K$ such that K is the
 159 submonoid of $M \times N$ generated by $\{(\varphi(a), \psi(a)) \mid a \in \Sigma\}$ and $\varphi \times \psi(a) = (\varphi(a), \psi(a))$ for
 160 all $a \in \Sigma$.

161 An *all-variety of stamps* (respectively *ne-variety of stamps*) is a non-empty class of stamps
 162 closed under direct product and *all-division* (respectively *ne-division*).

163 We will often use the following characteristic index of stamps, defined in [7]. Consider
 164 some stamp $\varphi: \Sigma^* \rightarrow M$. As M is finite there is a $k \in \mathbb{N}_{>0}$ such that $\varphi(\Sigma^{2k}) = \varphi(\Sigma^k)$: this
 165 implies that $\varphi(\Sigma^k)$ is a semigroup. The smallest such k is called the *stability index* of φ .

166 **Varieties of languages.** A language L over some alphabet Σ is *recognised by a monoid* M
 167 if there is a morphism $\varphi: \Sigma^* \rightarrow M$ and $F \subseteq M$ such that $L = \varphi^{-1}(F)$. We also say that φ
 168 *recognises* L . It is well known that a language is regular if and only if it is recognised by a
 169 finite monoid. The *syntactic congruence* of L , denoted by \sim_L , is the equivalence relation
 170 on Σ^* defined by $u \sim_L v$ for $u, v \in \Sigma^*$ whenever for all $x, y \in \Sigma^*$, $xuy \in L$ if and only if
 171 $xvy \in L$. The quotient Σ^*/\sim_L is a monoid, called *the syntactic monoid of L* , that recognises
 172 L via the *syntactic morphism* η_L of L sending any word u to its equivalence class $[u]_{\sim_L}$ for
 173 \sim_L . A stamp $\varphi: \Sigma^* \rightarrow M$ recognises L if and only if there exists a surjective morphism
 174 $\varphi: M \rightarrow \Sigma^*/\sim_L$ verifying $\eta_L = \alpha \circ \varphi$.

175 A *class of languages* \mathcal{C} is a correspondence that associates a set $\mathcal{C}(\Sigma)$ to each alphabet Σ .
 176 A (all-)variety of languages (respectively an *ne-variety of languages*) \mathcal{V} is a non-empty class
 177 of regular languages closed under Boolean operations, quotients and inverses of *all-morphisms*
 178 (respectively *ne-morphisms*). A classical result of Eilenberg [11, Chapter VII, Section 3]
 179 says that there is a bijective correspondence between varieties of monoids and varieties of
 180 languages: to each variety of monoids \mathbf{V} we can bijectively associate $\mathcal{L}(\mathbf{V})$ the variety of

languages whose syntactic monoids belong to \mathbf{V} . This was generalised by Straubing [18] to the case of varieties of stamps: to each *all*-variety (respectively *ne*-variety) of stamps \mathbf{V} we can bijectively associate $\mathcal{L}(\mathbf{V})$ the *all*-variety (respectively *ne*-variety) of languages whose syntactic morphisms belong to \mathbf{V} . Given two *all*-varieties (respectively *ne*-varieties) of stamps \mathbf{V}_1 and \mathbf{V}_2 , we have $\mathbf{V}_1 \subseteq \mathbf{V}_2 \Leftrightarrow \mathcal{L}(\mathbf{V}_1) \subseteq \mathcal{L}(\mathbf{V}_2)$.

For \mathbf{V} some variety of monoids, we define $\langle \mathbf{V} \rangle_{all}$ the *all*-variety of all stamps $\varphi: \Sigma^* \rightarrow M$ such that $M \in \mathbf{V}$. Of course, in that case $\mathcal{L}(\mathbf{V}) = \mathcal{L}(\langle \mathbf{V} \rangle_{all})$. Similarly, for \mathbf{V} some variety of semigroups, we define $\langle \mathbf{V} \rangle_{ne}$ the *ne*-variety of all stamps $\varphi: \Sigma^* \rightarrow M$ such that $\varphi(\Sigma^+) \in \mathbf{V}$. In that case, we consider $\mathcal{L}(\mathbf{V})$ to be the *ne*-variety of languages corresponding to $\langle \mathbf{V} \rangle_{ne}$. The operations $\langle \cdot \rangle_{all}$ and $\langle \cdot \rangle_{ne}$ form bijective correspondences (see [18]).

Identities. Every finite monoid M is considered to be a complete metric space equipped with the discrete metric d defined by $d(m, n) = \begin{cases} 0 & \text{if } m = n \\ 1 & \text{otherwise} \end{cases}$ for all $m, n \in M$.

Let Σ be an alphabet. Given $u, v \in \Sigma^*$, we set

$$r(u, v) = \min\{|M| \mid \exists \varphi: \Sigma^* \rightarrow M \text{ stamp s.t. } \varphi(u) \neq \varphi(v)\}$$

and $d(u, v) = 2^{-r(u, v)}$, using the conventions that $\min \emptyset = +\infty$ and $2^{-\infty} = 0$. Then d is a metric on Σ^* . The completion of the metric space (Σ^*, d) , denoted by $(\widehat{\Sigma^*}, \widehat{d})$, is a metric monoid called the *free profinite monoid on Σ^** . Its elements are all the formal limits $\lim_{n \rightarrow \infty} x_n$ of Cauchy sequences $(x_n)_{n \geq 0}$ in (Σ^*, d) and the metric d on Σ^* extends to a metric \widehat{d} on $\widehat{\Sigma^*}$ defined by $\widehat{d}(\lim_{n \rightarrow \infty} x_n, \lim_{n \rightarrow \infty} y_n) = \lim_{n \rightarrow \infty} d(x_n, y_n)$ for Cauchy sequences $(x_n)_{n \geq 0}$ and $(y_n)_{n \geq 0}$ in (Σ^*, d) . Note that, when it is clear from the context, we usually do not make the metric explicit when talking about a metric space. One important example of elements of $\widehat{\Sigma^*}$ is given by the elements $x^\omega = \lim_{n \rightarrow \infty} x^{n!}$ for all $x \in \Sigma^*$.

Every stamp $\varphi: \Sigma^* \rightarrow M$ extends uniquely to a uniformly continuous morphism $\widehat{\varphi}: \widehat{\Sigma^*} \rightarrow M$ with $\widehat{\varphi}(\lim_{n \rightarrow \infty} x_n) = \lim_{n \rightarrow \infty} \varphi(x_n)$ for every Cauchy sequence $(x_n)_{n \geq 0}$ in Σ^* . Similarly, every *all*-morphism $f: \Sigma^* \rightarrow \Gamma^*$ extends uniquely to a uniformly continuous morphism $\widehat{f}: \widehat{\Sigma^*} \rightarrow \widehat{\Gamma^*}$ with $\widehat{f}(\lim_{n \rightarrow \infty} x_n) = \lim_{n \rightarrow \infty} f(x_n)$ for every Cauchy sequence $(x_n)_{n \geq 0}$ in Σ^* .

For $u, v \in \widehat{A^*}$ with A an alphabet, we say that a stamp $\Sigma^* \rightarrow M$ *all-satisfies* (respectively *ne-satisfies*) the identity $u = v$ if for every *all*-morphism (respectively *ne*-morphism) $f: A^* \rightarrow \Sigma^*$, it holds that $\widehat{\varphi} \circ \widehat{f}(u) = \widehat{\varphi} \circ \widehat{f}(v)$. Given some set of identities E , we denote by $\llbracket E \rrbracket_{all}$ (respectively $\llbracket E \rrbracket_{ne}$) the class of stamps *all*-satisfying (respectively *ne*-satisfying) all the identities of E . When $\llbracket E \rrbracket_{all}$ (respectively $\llbracket E \rrbracket_{ne}$) is equal to some *all*-variety (respectively *ne*-variety) of stamps \mathbf{V} , we say that E *all-defines* (respectively *ne-defines*) \mathbf{V} .

► **Theorem 1.** *A class of stamps is an all-variety (respectively ne-variety) of stamps if and only if it can be all-defined (respectively ne-defined) by a set of identities.*

To give some examples, the classical varieties of monoids \mathbf{J} , \mathbf{R} and \mathbf{L} can be characterised by identities in the following way:

$$\langle \mathbf{R} \rangle_{all} = \llbracket (ab)^\omega a = (ab)^\omega \rrbracket_{all} = \llbracket (ab)^\omega a = (ab)^\omega \rrbracket_{ne}$$

$$\langle \mathbf{L} \rangle_{all} = \llbracket b(ab)^\omega = (ab)^\omega \rrbracket_{all} = \llbracket b(ab)^\omega = (ab)^\omega \rrbracket_{ne}$$

$$\langle \mathbf{J} \rangle_{all} = \llbracket (ab)^\omega a = (ab)^\omega, b(ab)^\omega = (ab)^\omega \rrbracket_{all} = \llbracket (ab)^\omega a = (ab)^\omega, b(ab)^\omega = (ab)^\omega \rrbracket_{ne}.$$

Finite locally trivial semigroups and the join operation. The variety \mathbf{LI} of finite locally trivial semigroups is well-known to verify $\langle \mathbf{LI} \rangle_{ne} = \llbracket x^\omega y x^\omega = x^\omega \rrbracket_{ne}$ and to be such that for

223 any alphabet Σ , the set $\mathcal{L}(\mathbf{LI})(\Sigma)$ contains exactly all Boolean combinations of languages of
 224 the form $u\Sigma^*$ or Σ^*u for $u \in \Sigma^*$, or equivalently exactly all languages of the form $U\Sigma^*V \cup W$
 225 with $U, V, W \subseteq \Sigma^*$ finite (see [13, p. 38]).

226 Given some variety of monoids \mathbf{V} , the join of \mathbf{V} and \mathbf{LI} , denoted by $\mathbf{V} \vee \mathbf{LI}$, is the
 227 inclusion-wise smallest variety of semigroups containing both \mathbf{V} and \mathbf{LI} . In fact, a finite
 228 semigroup S belongs to $\mathbf{V} \vee \mathbf{LI}$ if and only if there exist $M \in \mathbf{V}$ and $T \in \mathbf{LI}$ such that
 229 S divides the semigroup $M \times T$. (See [11, Chapter V, Exercise 1.1].) We can prove the
 230 following (see the appendix for the proof).

231 ► **Proposition 2.** *Let \mathbf{V} be a variety of monoids. Then $\langle \mathbf{V} \vee \mathbf{LI} \rangle_{\text{ne}}$ is the inclusion-wise
 232 smallest ne-variety of stamps containing both $\langle \mathbf{V} \rangle_{\text{all}}$ and $\langle \mathbf{LI} \rangle_{\text{ne}}$. Moreover, $\mathcal{L}(\mathbf{V} \vee \mathbf{LI})$ is the
 233 inclusion-wise smallest ne-variety of languages containing both $\mathcal{L}(\mathbf{V})$ and $\mathcal{L}(\mathbf{LI})$ and verifies
 234 that $\mathcal{L}(\mathbf{V} \vee \mathbf{LI})(\Sigma)$ is the Boolean closure of $\mathcal{L}(\mathbf{V})(\Sigma) \cup \mathcal{L}(\mathbf{LI})(\Sigma)$ for each alphabet Σ .*

235 3 Essentially- \mathbf{V} stamps

236 In this section, we give a characterisation of essentially- \mathbf{V} stamps (defined in [12]), for \mathbf{V} a
 237 variety of monoids, in terms of identities. We first recall the definition.

238 ► **Definition 3.** *Let \mathbf{V} be a variety of monoids. Let $\varphi: \Sigma^* \rightarrow M$ be a stamp from an alphabet
 239 Σ to a finite monoid M . Let s be the stability index of φ .*

240 *We say that φ is essentially- \mathbf{V} whenever there exists a stamp $\mu: \Sigma^* \rightarrow N$ with $N \in \mathbf{V}$
 241 such that for all $u, v \in \Sigma^*$, we have*

$$242 \quad \mu(u) = \mu(v) \Rightarrow (\varphi(xuy) = \varphi(xvy) \quad \forall x, y \in \Sigma^s) .$$

243 *We will denote by \mathbf{EV} the class of all essentially- \mathbf{V} stamps.*

244 Now, we give an equivalent condition for a stamp to be essentially- \mathbf{V} , based on a specific
 245 congruence depending on that stamp.

246 ► **Definition 4.** *Let $\varphi: \Sigma^* \rightarrow M$ for Σ an alphabet and M a finite monoid be a stamp and let
 247 s be its stability index. We define the equivalence relation \equiv_φ on Σ^* by $u \equiv_\varphi v$ for $u, v \in \Sigma^*$
 248 whenever $\varphi(xuy) = \varphi(xvy)$ for all $x, y \in \Sigma^{\geq s}$.*

249 ► **Proposition 5.** *Let $\varphi: \Sigma^* \rightarrow M$ for Σ an alphabet and M a finite monoid be a stamp.
 250 Then \equiv_φ is a congruence of finite index and for any variety of monoids \mathbf{V} , we have $\varphi \in \mathbf{EV}$
 251 if and only if $\Sigma^*/\equiv_\varphi \in \mathbf{V}$.*

252 **Proof.** Let us denote by s the stability index of φ .

253 The equivalence relation \equiv_φ is a congruence because given $u, v \in \Sigma^*$ verifying $u \equiv_\varphi v$,
 254 for all $\alpha, \beta \in \Sigma^*$, we have $\alpha u \beta \equiv_\varphi \alpha v \beta$ since for any $x, y \in \Sigma^{\geq s}$, it holds that $\varphi(x\alpha u \beta y) =$
 255 $\varphi(x\alpha v \beta y)$ because $x\alpha, \beta y \in \Sigma^{\geq s}$. Furthermore, this congruence is of finite index because for
 256 all $u, v \in \Sigma^*$, we have that $\varphi(u) = \varphi(v)$ implies $u \equiv_\varphi v$.

257 Let now \mathbf{V} be a variety of monoids. Assume first that $\Sigma^*/\equiv_\varphi \in \mathbf{V}$. It is quite direct to
 258 see that $\varphi \in \mathbf{EV}$, as the stamp $\mu: \Sigma^* \rightarrow \Sigma^*/\equiv_\varphi$ defined by $\mu(w) = [w]_{\equiv_\varphi}$ for all $w \in \Sigma^*$
 259 witnesses this fact. Assume then that $\varphi \in \mathbf{EV}$. This means that there exists a stamp
 260 $\mu: \Sigma^* \rightarrow N$ with $N \in \mathbf{V}$ such that for all $u, v \in \Sigma^*$, we have

$$261 \quad \mu(u) = \mu(v) \Rightarrow (\varphi(xuy) = \varphi(xvy) \quad \forall x, y \in \Sigma^s) .$$

262 Now consider $u, v \in \Sigma^*$ such that $\mu(u) = \mu(v)$. For any $x, y \in \Sigma^{\geq s}$, we have that $x = x_1x_2$
 263 with $x_1 \in \Sigma^*$ and $x_2 \in \Sigma^s$ as well as $y = y_1y_2$ with $y_1 \in \Sigma^s$ and $y_2 \in \Sigma^*$, so that

264 $\varphi(xuy) = \varphi(x_1)\varphi(x_2uy_1)\varphi(y_2) = \varphi(x_1)\varphi(x_2vy_1)\varphi(y_2) = \varphi(xvy)$. Hence, $u \equiv_\varphi v$. Therefore,
 265 for all $u, v \in \Sigma^*$, we have that $\mu(u) = \mu(v)$ implies $u \equiv_\varphi v$, so we can define the mapping
 266 $\alpha: N \rightarrow \Sigma^*/\equiv_\varphi$ such that $\alpha(\mu(w)) = [w]_{\equiv_\varphi}$ for all $w \in \Sigma^*$. It is easy to check that α
 267 is actually a surjective morphism. Thus, we can conclude that Σ^*/\equiv_φ , which divides N ,
 268 belongs to \mathbf{V} . \blacktriangleleft

269 Using this equivalent condition, we prove that given a set of identities ne -defining $\langle \mathbf{V} \rangle_{all}$
 270 for some variety of monoids \mathbf{V} , we get a set of identities ne -defining \mathbf{EV} , which actually
 271 shows that \mathbf{EV} is an ne -variety of stamps along the way.

272 **► Proposition 6.** *Let \mathbf{V} be a variety of monoids and let E be a set of identities such that*
 273 $\langle \mathbf{V} \rangle_{all} = \llbracket E \rrbracket_{ne}$. *Then*

$$274 \quad \mathbf{EV} = \llbracket x^\omega yuzt^\omega = x^\omega yvzt^\omega \mid u = v \in E, x, y, z, t \notin \text{alph}(u) \cup \text{alph}(v) \rrbracket_{ne} .$$

275 **Proof.** Let

$$276 \quad F = \{x^\omega yuzt^\omega = x^\omega yvzt^\omega \mid u = v \in E, x, y, z, t \notin \text{alph}(u) \cup \text{alph}(v)\} .$$

277 Central to the proof is the following claim.

278 \triangleright **Claim 7.** Let $\varphi: \Sigma^* \rightarrow M$ for Σ an alphabet and M a finite monoid be a stamp. Consider
 279 the stamp $\mu: \Sigma^* \rightarrow \Sigma^*/\equiv_\varphi$ defined by $\mu(w) = [w]_{\equiv_\varphi}$ for all $w \in \Sigma^*$. It holds that for all
 280 $u, v \in \widehat{\Sigma}^*$,

$$281 \quad \widehat{\mu}(u) = \widehat{\mu}(v) \Leftrightarrow (\widehat{\varphi}(\alpha^\omega \beta u \gamma \delta^\omega) = \widehat{\varphi}(\alpha^\omega \beta v \gamma \delta^\omega) \quad \forall \alpha, \beta, \gamma, \delta \in \Sigma^+) .$$

282 Using this claim, we can prove that $\mathbf{EV} = \llbracket F \rrbracket_{ne}$.

283 **Inclusion from left to right.** Let $\varphi: \Sigma^* \rightarrow M$ for Σ an alphabet and M a finite monoid
 284 be a stamp in \mathbf{EV} . Consider the stamp $\mu: \Sigma^* \rightarrow \Sigma^*/\equiv_\varphi$ defined by $\mu(w) = [w]_{\equiv_\varphi}$ for all
 285 $w \in \Sigma^*$. Since $\varphi \in \mathbf{EV}$, Proposition 5 tells us that $\Sigma^*/\equiv_\varphi \in \mathbf{V}$, hence $\mu \in \langle \mathbf{V} \rangle_{all}$.

286 Let us consider any identity $x^\omega yuzt^\omega = x^\omega yvzt^\omega \in F$. It is thus defined on an alphabet
 287 B with $u = v \in E$ defined on an alphabet $A \subseteq B$ and $x, y, z, t \in B \setminus A$. Let $f: B^* \rightarrow \Sigma^*$ be
 288 an ne -morphism. Since $\mu \in \langle \mathbf{V} \rangle_{all}$, we have that μ ne -satisfies the identity $u = v$, so that
 289 $\widehat{\mu}(\widehat{f}(u)) = \widehat{\mu}(\widehat{f}(v))$. Notice that we have that $\widehat{f}(x^\omega) = f(x)^\omega$ as well as $\widehat{f}(t^\omega) = f(t)^\omega$ and
 290 that $f(x), f(y), f(z), f(t) \in \Sigma^+$ because f is non-erasing. Therefore, we have

$$291 \quad \begin{aligned} \widehat{\varphi}(\widehat{f}(x^\omega yuzt^\omega)) &= \widehat{\varphi}(f(x)^\omega f(y)\widehat{f}(u)f(z)f(t)^\omega) \\ 292 \quad &= \widehat{\varphi}(f(x)^\omega f(y)\widehat{f}(v)f(z)f(t)^\omega) \\ 293 \quad &= \widehat{\varphi}(\widehat{f}(x^\omega yvzt^\omega)) \end{aligned}$$

295 by Claim 7. As this holds for any ne -morphism $f: B^* \rightarrow \Sigma^*$, we can conclude that φ
 296 ne -satisfies the identity $x^\omega yuzt^\omega = x^\omega yvzt^\omega$.

297 This is true for any identity in F , so $\varphi \in \llbracket F \rrbracket_{ne}$. In conclusion, $\mathbf{EV} \subseteq \llbracket F \rrbracket_{ne}$.

298 **Inclusion from right to left.** Let $\varphi: \Sigma^* \rightarrow M$ for Σ an alphabet and M a finite monoid
 299 be a stamp in $\llbracket F \rrbracket_{ne}$. Consider the stamp $\mu: \Sigma^* \rightarrow \Sigma^*/\equiv_\varphi$ defined by $\mu(w) = [w]_{\equiv_\varphi}$ for all
 300 $w \in \Sigma^*$. We are now going to show that $\mu \in \langle \mathbf{V} \rangle_{all}$.

301 Take any identity $u = v \in E$ defined on an alphabet A . There is thus an identity
 302 $x^\omega yuzt^\omega = x^\omega yvzt^\omega \in F$ defined on an alphabet B such that $A \subseteq B$ and $x, y, z, t \in B \setminus A$.
 303 Let $f: A^* \rightarrow \Sigma^*$ be an ne -morphism.

304 Take any $\alpha, \beta, \gamma, \delta \in \Sigma^+$. Let us define the *ne*-morphism $g: B^* \rightarrow \Sigma^*$ as the unique one
305 such that for all $b \in B$,

$$306 \quad g(b) = \begin{cases} f(b) & \text{if } b \in A \\ \alpha & \text{if } b = x \\ \beta & \text{if } b = y \\ \gamma & \text{if } b = z \\ \delta & \text{if } b = t \end{cases} .$$

307 Observe in particular that $\widehat{g}(w) = \widehat{f}(w)$ for any $w \in \widehat{A}^*$ and that $\widehat{g}(x^\omega) = g(x)^\omega = \alpha^\omega$ as
308 well as $\widehat{g}(t^\omega) = \delta^\omega$. Now, as φ *ne*-satisfies $x^\omega y u z t^\omega = x^\omega y v z t^\omega$, we have that

$$309 \quad \widehat{\varphi}(\alpha^\omega \beta \widehat{f}(u) \gamma \delta^\omega) = \widehat{\varphi}(\widehat{g}(x^\omega y u z t^\omega)) = \widehat{\varphi}(\widehat{g}(x^\omega y v z t^\omega)) = \widehat{\varphi}(\alpha^\omega \beta \widehat{f}(v) \gamma \delta^\omega) .$$

310 Since this holds for any $\alpha, \beta, \gamma, \delta \in \Sigma^+$, by Claim 7, we have that $\widehat{\mu}(\widehat{f}(u)) = \widehat{\mu}(\widehat{f}(v))$.

311 Therefore, $\widehat{\mu}(\widehat{f}(u)) = \widehat{\mu}(\widehat{f}(v))$ for any *ne*-morphism $f: A^* \rightarrow \Sigma^*$, which means that μ
312 *ne*-satisfies $u = v$.

313 So, to conclude, since this is true for any $u = v \in E$, we have that $\mu \in \langle \mathbf{V} \rangle_{all}$, which
314 implies that $\Sigma^* / \equiv_\varphi \in \mathbf{V}$ and thus $\varphi \in \mathbf{EV}$ by Proposition 5. In conclusion, $\llbracket F \rrbracket_{ne} \subseteq \mathbf{EV}$.

315 The claim still needs to be proved.

316 Proof of Claim 7. Let $\varphi: \Sigma^* \rightarrow M$ for Σ an alphabet and M a finite monoid be a stamp
317 of stability index s . Consider the stamp $\mu: \Sigma^* \rightarrow \Sigma^* / \equiv_\varphi$ defined by $\mu(w) = [w]_{\equiv_\varphi}$ for all
318 $w \in \Sigma^*$. We now want to show that for all $u, v \in \widehat{\Sigma}^*$,

$$319 \quad \widehat{\mu}(u) = \widehat{\mu}(v) \Leftrightarrow (\widehat{\varphi}(\alpha^\omega \beta u \gamma \delta^\omega) = \widehat{\varphi}(\alpha^\omega \beta v \gamma \delta^\omega) \quad \forall \alpha, \beta, \gamma, \delta \in \Sigma^+) .$$

320 Let $u, v \in \widehat{\Sigma}^*$. There exist two Cauchy sequences $(u_n)_{n \geq 0}$ and $(v_n)_{n \geq 0}$ in Σ^* such that
321 $u = \lim_{n \rightarrow \infty} u_n$ and $v = \lim_{n \rightarrow \infty} v_n$. As $\Sigma^* / \equiv_\varphi$ and M are discrete, we have that all
322 four Cauchy sequences $(\mu(u_n))_{n \geq 0}$, $(\varphi(u_n))_{n \geq 0}$, $(\mu(v_n))_{n \geq 0}$ and $(\varphi(v_n))_{n \geq 0}$ are ultimately
323 constant. So there exists $k \in \mathbb{N}$ such that $\widehat{\mu}(u) = \mu(u_k)$, $\widehat{\varphi}(u) = \varphi(u_k)$, $\widehat{\mu}(v) = \mu(v_k)$ and
324 $\widehat{\varphi}(v) = \varphi(v_k)$.

325 Assume first that $\widehat{\mu}(u) = \widehat{\mu}(v)$. Take any $\alpha, \beta, \gamma, \delta \in \Sigma^+$. Since M is discrete, both
326 Cauchy sequences $(\varphi(\alpha^{n!}))_{n \geq 0}$ and $(\varphi(\delta^{n!}))_{n \geq 0}$ are ultimately constant. So there exists
327 $l \in \mathbb{N}$ such that for all $m \in \mathbb{N}, m \geq l$, we have $\widehat{\varphi}(\alpha^\omega) = \varphi(\alpha^{m!})$ and $\widehat{\varphi}(\delta^\omega) = \varphi(\delta^{m!})$. Hence,
328 taking some $m \in \mathbb{N}, m \geq l$ such that $|\alpha^{m!} \beta| \geq s$ and $|\gamma \delta^{m!}| \geq s$, it follows that

$$329 \quad \widehat{\varphi}(\alpha^\omega \beta u \gamma \delta^\omega) = \varphi(\alpha^{m!} \beta u_k \gamma \delta^{m!}) = \varphi(\alpha^{m!} \beta v_k \gamma \delta^{m!}) = \widehat{\varphi}(\alpha^\omega \beta v \gamma \delta^\omega)$$

330 because $[u_k]_{\equiv_\varphi} = \widehat{\mu}(u) = \widehat{\mu}(v) = [v_k]_{\equiv_\varphi}$. Thus, we have that

$$331 \quad \widehat{\varphi}(\alpha^\omega \beta u \gamma \delta^\omega) = \widehat{\varphi}(\alpha^\omega \beta v \gamma \delta^\omega)$$

332 for all $\alpha, \beta, \gamma, \delta \in \Sigma^+$.

333 Assume then that $\widehat{\varphi}(\alpha^\omega \beta u \gamma \delta^\omega) = \widehat{\varphi}(\alpha^\omega \beta v \gamma \delta^\omega)$ for all $\alpha, \beta, \gamma, \delta \in \Sigma^+$. Take any $\alpha, \beta \in$
334 $\Sigma^{\geq s}$. Since $\varphi(\Sigma^s)$ is a finite semigroup and verifies that $\varphi(\Sigma^s) = \varphi(\Sigma^s)^2$, by a classical result
335 in finite semigroup theory (see e.g. [13, Chapter 1, Proposition 1.12]), we have that there
336 exist $\alpha_1, e, f, \beta_2 \in \Sigma^s$ and $\alpha_2, \beta_1 \in \Sigma^{\geq s}$ such that $\varphi(\alpha_1 e \alpha_2) = \varphi(\alpha)$ and $\varphi(\beta_1 f \beta_2) = \varphi(\beta)$
337 with $\varphi(e)$ and $\varphi(f)$ idempotents. Now, since $\varphi(e)$ is idempotent, we have that

$$338 \quad \widehat{\varphi}(e^\omega) = \widehat{\varphi}(\lim_{n \rightarrow \infty} e^{n!}) = \lim_{n \rightarrow \infty} \varphi(e^{n!}) = \lim_{n \rightarrow \infty} \varphi(e)^{n!} = \varphi(e)$$

339 and similarly, $\widehat{\varphi}(f^\omega) = \varphi(f)$. So it follows that

$$\begin{aligned}
 340 \quad \varphi(\alpha u_k \beta) &= \varphi(\alpha_1 e \alpha_2 u_k \beta_1 f \beta_2) \\
 341 \quad &= \widehat{\varphi}(\alpha_1 e^\omega \alpha_2 u \beta_1 f^\omega \beta_2) \\
 342 \quad &= \widehat{\varphi}(\alpha_1 e^\omega \alpha_2 v \beta_1 f^\omega \beta_2) \\
 343 \quad &= \varphi(\alpha_1 e \alpha_2 v_k \beta_1 f \beta_2) \\
 344 \quad &= \varphi(\alpha v_k \beta) . \\
 345
 \end{aligned}$$

346 As this is true for any $\alpha, \beta \in \Sigma^{\geq s}$, by definition it holds that $u_k \equiv_\varphi v_k$, hence $\widehat{\mu}(u) = \mu(u_k) =$
 347 $\mu(v_k) = \widehat{\mu}(v)$. \triangleleft

348 This concludes the proof of the proposition. \blacktriangleleft

349 **4 Essentially-V stamps and the join of V and LI**

350 In this section, we establish the link between essentially-V stamps and $\mathbf{V} \vee \mathbf{LI}$ and give a
 351 criterion that characterises exactly when they do correspond.

352 More precisely, consider the following criterion for a variety of monoids \mathbf{V} .

353 **► Criterion (A).** For any $L \in \mathcal{L}(\mathbf{V})(\Sigma)$ with Σ some alphabet, we have $xLy \in \mathcal{L}(\mathbf{V} \vee \mathbf{LI})(\Sigma)$
 354 for all $x, y \in \Sigma^*$.

355 It is a kind of mild closure condition that appears to be a sufficient and necessary condition
 356 for \mathbf{EV} and $\mathbf{V} \vee \mathbf{LI}$ to correspond.

357 **► Proposition 8.** *Let \mathbf{V} be a variety of monoids. Then $\langle \mathbf{V} \vee \mathbf{LI} \rangle_{\text{ne}} \subseteq \mathbf{EV}$ and there is
 358 equality if and only if \mathbf{V} verifies criterion (A).*

359 Why this proposition is actually useful to give characterisations of $\mathbf{V} \vee \mathbf{LI}$ in terms of
 360 identities will become clear in the next section. For now, we focus on its proof, that entirely
 361 relies on the following characterisation of the languages recognised by essentially-V stamps.

362 **► Proposition 9.** *Let \mathbf{V} be a variety of monoids. For any alphabet Σ , the set $\mathcal{L}(\mathbf{EV})(\Sigma)$
 363 contains exactly all Boolean combinations of languages of the form xLy for $L \in \mathcal{L}(\mathbf{V})(\Sigma)$
 364 and $x, y \in \Sigma^*$.*

365 **Proof.** Let \mathcal{C} be the class of languages such that for any alphabet Σ , the set $\mathcal{C}(\Sigma)$ contains
 366 exactly all Boolean combinations of languages of the form xLy for $L \in \mathcal{L}(\mathbf{V})(\Sigma)$ and $x, y \in \Sigma^*$.

367 Let Σ be an alphabet. We are now going to show that $\mathcal{L}(\mathbf{EV})(\Sigma) = \mathcal{C}(\Sigma)$, which proves
 368 the proposition as it holds for any alphabet Σ .

369 **Inclusion from right to left.** Let $L \in \mathcal{L}(\mathbf{V})(\Sigma)$ and $x, y \in \Sigma^*$. Let $\mu: \Sigma^* \rightarrow N$ be the
 370 syntactic morphism of L : this means that $N \in \mathbf{V}$ and that there exists $F \subseteq N$ such
 371 that $L = \mu^{-1}(F)$. Let also $\varphi: \Sigma^* \rightarrow M$ be the syntactic morphism of the language
 372 $xLy = x\Sigma^*y \cap \Sigma^{|x|}\mu^{-1}(F)\Sigma^{|y|}$ and let s be its stability index. We then consider some
 373 $u, v \in \Sigma^*$ such that $\mu(u) = \mu(v)$. Take any $x', y' \in \Sigma^*$ such that $|x'| \geq |x|$ and $|y'| \geq |y|$. We
 374 clearly have that $x'uy' \in x\Sigma^*y$ if and only if $x'vy' \in x\Sigma^*y$. Moreover, $x' = x'_1x'_2$ for some
 375 $x'_1 \in \Sigma^{|x|}$ and $x'_2 \in \Sigma^*$ and $y' = y'_1y'_2$ for some $y'_1 \in \Sigma^*$ and $y'_2 \in \Sigma^{|y|}$, so that

$$\begin{aligned}
 376 \quad x'uy' \in \Sigma^{|x|}\mu^{-1}(F)\Sigma^{|y|} &\Leftrightarrow \mu(x'_2uy'_1) \in F \\
 377 \quad &\Leftrightarrow \mu(x'_2vy'_1) \in F \\
 378 \quad &\Leftrightarrow x'vy' \in \Sigma^{|x|}\mu^{-1}(F)\Sigma^{|y|} .
 \end{aligned}$$

380 Hence, $x'uy' \in xLy$ if and only if $x'vy' \in xLy$ for all $x', y' \in \Sigma^*$ such that $|x'| \geq |x|$ and
 381 $|y'| \geq |y|$, so that, by definition of the stability index s of φ and as φ is the syntactic morphism
 382 of xLy , we have $\varphi(x'uy') = \varphi(x'vy')$ for all $x', y' \in \Sigma^s$. Thus, it follows that $\varphi \in \mathbf{EV}$.

383 This implies that $xLy \in \mathcal{L}(\mathbf{EV})(\Sigma)$. Therefore, since this is true for any $L \in \mathcal{L}(\mathbf{V})(\Sigma)$
 384 and $x, y \in \Sigma^*$ and since $\mathcal{L}(\mathbf{EV})(\Sigma)$ is closed under Boolean operations, we can conclude that
 385 $\mathcal{C}(\Sigma) \subseteq \mathcal{L}(\mathbf{EV})(\Sigma)$.

386 **Inclusion from left to right.** Let $L \in \mathcal{L}(\mathbf{EV})(\Sigma)$ and let $\varphi: \Sigma^* \rightarrow M$ be its syntactic
 387 morphism: it is an essentially- \mathbf{V} stamp. Given s its stability index, this means there exists a
 388 stamp $\mu: \Sigma^* \rightarrow N$ with $N \in \mathbf{V}$ such that for all $u, v \in \Sigma^*$, we have

$$389 \quad \mu(u) = \mu(v) \Rightarrow (\varphi(xuy) = \varphi(xvy) \quad \forall x, y \in \Sigma^s) .$$

390 For each $m \in N$ and $x, y \in \Sigma^s$ consider the language $x\mu^{-1}(m)y$: it verifies that for any
 391 two words $w, w' \in x\mu^{-1}(m)y$, we have $w = xuy$ and $w' = xvy$ with $\mu(u) = \mu(v) = m$, so
 392 that $\varphi(w) = \varphi(w')$. By definition of the syntactic morphism, this means that for all $m \in N$
 393 and $x, y \in \Sigma^s$, either $x\mu^{-1}(m)y \subseteq L$ or $x\mu^{-1}(m)y \cap L = \emptyset$. Therefore, there exists a set
 394 $E \subseteq N \times \Sigma^s \times \Sigma^s$ such that $L \cap \Sigma^{\geq 2s} = \bigcup_{(m,x,y) \in E} x\mu^{-1}y$, hence

$$395 \quad L = \bigcup_{(m,x,y) \in E} x\mu^{-1}(m)y \cup F$$

396 for some $F \subseteq \Sigma^{< 2s}$.

397 Take some $w \in F$. We have that $\{w\} = w\Sigma^* \cap \bigcap_{a \in \Sigma} (wa\Sigma^*)^c$ with necessarily $\Sigma^* \in$
 398 $\mathcal{L}(\mathbf{V})(\Sigma)$. Thus, the singleton language $\{w\}$ belongs to $\mathcal{C}(\Sigma)$ and since this is true for any
 399 $w \in F$ and F is finite, we can deduce from this that F is in $\mathcal{C}(\Sigma)$, as the latter is trivially
 400 closed under Boolean operations.

401 Now, for all $m \in N$, the language $\mu^{-1}(m)$ belongs to $\mathcal{L}(\mathbf{V})(\Sigma)$, so we finally have $L \in \mathcal{C}(\Sigma)$.
 402 This is true for any $L \in \mathcal{L}(\mathbf{EV})(\Sigma)$, so in conclusion, $\mathcal{L}(\mathbf{EV})(\Sigma) \subseteq \mathcal{C}(\Sigma)$. \blacktriangleleft

403 Proposition 8 then follows from the two next lemmata, that are both easy consequences
 404 of Proposition 9. For completeness, we give the proofs in the appendix.

405 **► Lemma 10.** *Let \mathbf{V} be a variety of monoids. Then $\langle \mathbf{V} \vee \mathbf{LI} \rangle_{\text{ne}} \subseteq \mathbf{EV}$.*

406 **► Lemma 11.** *Let \mathbf{V} be a variety of monoids. Then $\mathbf{EV} \subseteq \langle \mathbf{V} \vee \mathbf{LI} \rangle_{\text{ne}}$ if and only if \mathbf{V}
 407 verifies criterion (A).*

408 **5 Applications**

409 In this last section, we use the link between essentially- \mathbf{V} stamps and $\mathbf{V} \vee \mathbf{LI}$ to reprove some
 410 characterisations of joins between \mathbf{LI} and some well-known varieties of monoids in terms of
 411 identities.

412 One thing that seems at a first glance a bit problematic about proving that some variety
 413 of monoids \mathbf{V} satisfies criterion (A) when one's goal is precisely to characterise $\mathbf{V} \vee \mathbf{LI}$ is
 414 that one needs to prove that some languages belong to $\mathcal{L}(\mathbf{V} \vee \mathbf{LI})$, a class about which one
 415 a priori only knows what is given by Proposition 2. But understanding $\mathcal{L}(\mathbf{V})$ enough is
 416 actually sufficient: if it holds that given any language $L \in \mathcal{L}(\mathbf{V})(\Sigma)$ for some alphabet Σ and
 417 any $x, y \in \Sigma^*$, there exists some language $K \in \mathcal{L}(\mathbf{V})(\Sigma)$ such that L is equal to the quotient
 418 $x^{-1}Ky^{-1}$, then \mathbf{V} verifies criterion (A). We don't know whether this quotient-expressibility
 419 condition that solely depends on the variety \mathbf{V} (without explicit reference to \mathbf{LI}) is actually

420 equivalent to it satisfying criterion (A), but we can prove such an equivalence for a weaker
421 quotient-expressibility condition for \mathbf{V} . The proof is to be found in the appendix.

422 ► **Proposition 12.** *Let \mathbf{V} be a variety of monoids. Then \mathbf{V} satisfies criterion (A) if and*
423 *only if for any $L \in \mathcal{L}(\mathbf{V})(\Sigma)$ for some alphabet Σ and $x, y \in \Sigma^*$, there exist $k, l \in \mathbb{N}$*
424 *such that for all $u \in \Sigma^k, v \in \Sigma^l$, there exists some language $K \in \mathcal{L}(\mathbf{V})(\Sigma)$ verifying*
425 *$u^{-1}Lv^{-1} = (xu)^{-1}K(vy)^{-1}$.*

426 This quotient-expressibility condition appears to be particularly useful to prove that a
427 variety of monoids \mathbf{V} does not satisfy criterion (A) without needing to understand what
428 $\mathcal{L}(\mathbf{V} \vee \mathbf{LI})$ is. We demonstrate this for the variety of finite commutative and idempotent
429 monoids \mathbf{J}_1 .

430 ► **Proposition 13.** *\mathbf{J}_1 does not satisfy criterion (A).*

431 **Proof.** Given an alphabet Σ , the set $\mathcal{L}(\mathbf{J}_1)(\Sigma)$ contains exactly all Boolean combinations of
432 languages of the form $\Sigma^*a\Sigma^*$ for $a \in \Sigma$ (see [13, Chapter 2, Proposition 3.10]).

433 Let $L = \{a, b\}^*b\{a, b\}^* \in \mathcal{L}(\mathbf{J}_1)(\{a, b\})$ and $x = b, y = \varepsilon$. Take any $k, l \in \mathbb{N}$ and set
434 $u = a^k$ and $v = a^l$. Consider some $K \in \mathcal{L}(\mathbf{J}_1)(\{a, b\})$. We have that $xuavy \in K \Leftrightarrow$
435 $xuabvy \in K$ so that $a \in (xu)^{-1}K(vy)^{-1} \Leftrightarrow ab \in (xu)^{-1}K(vy)^{-1}$. But $a \notin u^{-1}Lv^{-1}$
436 and $ab \in u^{-1}Lv^{-1}$, hence $u^{-1}Lv^{-1} \neq (xu)^{-1}K(vy)^{-1}$ and this holds for any choice of K .
437 So for any $k, l \in \mathbb{N}$, there exists $u \in \Sigma^k, v \in \Sigma^l$ such that no $K \in \mathcal{L}(\mathbf{J}_1)(\{a, b\})$ verifies
438 $u^{-1}Lv^{-1} = (xu)^{-1}K(vy)^{-1}$.

439 In conclusion, by Proposition 12, \mathbf{J}_1 does not satisfy criterion (A). ◀

440 We now prove the announced characterisations of joins between \mathbf{LI} and some well-known
441 varieties of monoids in terms of identities.

442 ► **Theorem 14.** *We have the following.*

- 443 1. $\langle \mathbf{R} \vee \mathbf{LI} \rangle_{\text{ne}} = \mathbf{ER} = \llbracket x^\omega y(ab)^\omega azt^\omega = x^\omega y(ab)^\omega zt^\omega \rrbracket_{\text{ne}}$.
- 444 2. $\langle \mathbf{L} \vee \mathbf{LI} \rangle_{\text{ne}} = \mathbf{EL} = \llbracket x^\omega yb(ab)^\omega zt^\omega = x^\omega y(ab)^\omega zt^\omega \rrbracket_{\text{ne}}$.
- 445 3. $\langle \mathbf{J} \vee \mathbf{LI} \rangle_{\text{ne}} = \mathbf{EJ} = \llbracket x^\omega y(ab)^\omega azt^\omega = x^\omega y(ab)^\omega zt^\omega, x^\omega yb(ab)^\omega zt^\omega = x^\omega y(ab)^\omega zt^\omega \rrbracket_{\text{ne}}$.
- 446 4. $\langle \mathbf{H} \vee \mathbf{LI} \rangle_{\text{ne}} = \mathbf{EH}$ for any variety of groups \mathbf{H} .

447 **Proof.** In each case, we prove that the variety of monoids under consideration satisfies
448 criterion (A) using Proposition 12. We then use Proposition 8.

449 **Proof of 1.** It is well-known that given an alphabet Σ , the set $\mathcal{L}(\mathbf{R})(\Sigma)$ contains all
450 languages that are disjoint unions of languages that are of the form $A_0^*a_1A_1^* \cdots a_kA_k^*$ where
451 $k \in \mathbb{N}$, $a_1, \dots, a_k \in \Sigma$, $A_0, A_1, \dots, A_k \subseteq \Sigma$ and $a_i \notin A_{i-1}$ for all $i \in [k]$ (see [13, Chapter 4,
452 Theorem 3.3]).

453 Let Σ be an alphabet and take some language $A_0^*a_1A_1^* \cdots a_kA_k^*$ where $k \in \mathbb{N}$, $a_1, \dots, a_k \in$
454 Σ , $A_0, A_1, \dots, A_k \subseteq \Sigma$ and $a_i \notin A_{i-1}$ for all $i \in [k]$. Take $x, y \in \Sigma^*$. Observe that y can be
455 uniquely written as $y = zt$ where $z \in A_k^*$ and $t \in \{\varepsilon\} \cup (\Sigma \setminus A_k)\Sigma^*$. We have

$$456 A_0^*a_1A_1^* \cdots a_kA_k^* = x^{-1} \left(xA_0^*a_1A_1^* \cdots a_kA_k^*t \cap \bigcap_{v \in A_k^{<|z|}} (xA_0^*a_1A_1^* \cdots a_kvt) \right) y^{-1}$$

457 using the convention that $xA_0^*a_1A_1^* \cdots a_kvt = xvt$ for all $v \in A_k^{<|z|}$ when $k = 0$. The language
458 $xA_0^*a_1A_1^* \cdots a_kA_k^*t \cap \bigcap_{v \in A_k^{<|z|}} (xA_0^*a_1A_1^* \cdots a_kvt)$ does belong to the set $\mathcal{L}(\mathbf{R})(\Sigma)$ because
459 the latter is closed under Boolean operations and by definition of z and t . Thus, we can
460 conclude that for each $L \in \mathcal{L}(\mathbf{R})(\Sigma)$ and $x, y \in \Sigma^*$, there exists $K \in \mathcal{L}(\mathbf{R})(\Sigma)$ such that
461 $L = x^{-1}Ky^{-1}$ by using the characterisation of $\mathcal{L}(\mathbf{R})(\Sigma)$, the fact that quotients commute
462 with unions [13, p. 20] and closure of $\mathcal{L}(\mathbf{R})(\Sigma)$ under unions.

463 **Proof of 2.** It is also well-known that given an alphabet Σ , the set $\mathcal{L}(\mathbf{L})(\Sigma)$ contains all
 464 languages that are disjoint unions of languages that are of the form $A_0^*a_1A_1^*\cdots a_kA_k^*$ where
 465 $k \in \mathbb{N}$, $a_1, \dots, a_k \in \Sigma$, $A_0, A_1, \dots, A_k \subseteq \Sigma$ and $a_i \notin A_i$ for all $i \in [k]$ (see [13, Chapter 4,
 466 Theorem 3.4]). The proof is then dual to the previous case.

467 **Proof of 3.** Given an alphabet Σ , for each $k \in \mathbb{N}$, we define the equivalence relation \sim_k
 468 on Σ^* by $u \sim_k v$ for $u, v \in \Sigma^*$ whenever u and v have the same set of subwords of length
 469 at most k . This relation is a congruence of finite index on Σ^* . Simon proved [17] that a
 470 language belongs to $\mathcal{L}(\mathbf{J})(\Sigma)$ if and only if it is equal to a union of \sim_k -classes for some $k \in \mathbb{N}$.

471 Let Σ be an alphabet and take some $L \in \mathcal{L}(\mathbf{J})(\Sigma)$ as well as $x, y \in \Sigma^*$. Thus, there exists
 472 $k \in \mathbb{N}$ such that L is a union of \sim_k -classes. Define the language $K = \bigcup_{w \in L} [xwy]_{\sim_{|xy|+k}}$: it
 473 belongs to $\mathcal{L}(\mathbf{J})(\Sigma)$ by construction. We now show that $L = x^{-1}Ky^{-1}$, which concludes
 474 the proof. Let $w \in L$: we have that $xwy \in [xwy]_{\sim_{|xy|+k}} \subseteq K$, so that $w \in x^{-1}Ky^{-1}$. Let
 475 conversely $w \in x^{-1}Ky^{-1}$. This means that $xwy \in K$, which implies that there exists $w' \in L$
 476 such that $xwy \sim_{|xy|+k} xw'y$. Actually, it holds that any $u \in \Sigma^*$ of length at most k is a
 477 subword of w if and only if it is a subword of w' , because xuy is a subword of xwy if and
 478 only if it is a subword of $xw'y$. Hence, $w \sim_k w'$, which implies that $w \in L$.

479 **Proof of 4.** Consider some variety of groups \mathbf{H} . Take some language $L \in \mathcal{L}(\mathbf{H})(\Sigma)$ for
 480 an alphabet Σ and let $x, y \in \Sigma^*$. Consider the syntactic morphism $\eta: \Sigma^* \rightarrow M$ of L : we
 481 have that M is a group in \mathbf{H} . Define the language $K = \eta^{-1}(\eta(x)\eta(L)\eta(y))$: it belongs to
 482 $\mathcal{L}(\mathbf{H})(\Sigma)$. We now show that $L = x^{-1}Ky^{-1}$, which concludes the proof. Let $w \in L$: we
 483 have that $\eta(xwy) \in \eta(x)\eta(L)\eta(y)$, so that $w \in x^{-1}Ky^{-1}$. Conversely, let $w \in x^{-1}Ky^{-1}$. We
 484 have that $xwy \in K$, which means that $\eta(xwy) = \eta(x)\eta(w')\eta(y)$ for some $w' \in L$, so that
 485 $\eta(w) = \eta(w') \in \eta(L)$, as any element in \mathbf{H} is invertible. Thus, $w \in L$. ◀

486 **6 Conclusion**

487 The general method presented in this paper actually allows to reprove in a straightforward
 488 language-theoretic way even more characterisations of the join of **LI** with some variety of
 489 finite monoids. This can for instance be done for the variety of finite commutative monoids
 490 **Com** or the variety of finite commutative aperiodic monoids **ACom**.

491 In fact, as already observed in some sense by Costa [8], many varieties of finite monoids
 492 seem to verify criterion (A). The main question left open by this present work is to understand
 493 better what exactly those varieties are. Another question left open is whether Proposition 12
 494 can be refined by using the stronger quotient-expressibility condition alluded to before the
 495 statement of the proposition. The answers to both questions are unclear to the author, but
 496 making progress on them may also lead to a better understanding of joins of varieties of
 497 finite monoids with **LI**.

498 **Acknowledgements.** The author wants to thank Thomas Place, who suggested the link
 499 between essentially-**V** stamps and $\mathbf{V} \vee \mathbf{LI}$, but also Luc Segoufin who started the discussion
 500 with Thomas Place and encouraged the author to write the present article.

501 ——— **References** ———

- 502 1 Jorge Almeida. Some pseudovariety joins involving the pseudovariety of finite groups. In
503 *Semigroup Forum*, volume 37, pages 53–57. Springer, 1988.
- 504 2 Jorge Almeida. *Finite semigroups and universal algebra*, volume 3. World Scientific, 1994.
- 505 3 Jorge Almeida and Assis Azevedo. The join of the pseudovarieties of R-trivial and L-trivial
506 monoids. *Journal of Pure and Applied Algebra*, 60(2):129–137, 1989.
- 507 4 Jorge Almeida and Pascal Weil. Free profinite $\mathit{mathscrR}$ -trivial monoids. *Int. J. Algebra*
508 *Comput.*, 7(5):625–672, 1997. doi:10.1142/S0218196797000289.
- 509 5 Assis Azevedo. *The Join of the Pseudovariety J with Permutative Pseudovarieties*, pages 1–11.
510 Springer US, Boston, MA, 1990. doi:10.1007/978-1-4899-2608-1_1.
- 511 6 Assis Azevedo and Marc Zeitoun. Three examples of join computations. In *Semigroup Forum*,
512 volume 57, pages 249–277. Springer, 1998.
- 513 7 Laura Chaubard, Jean-Eric Pin, and Howard Straubing. First order formulas with modular
514 predicates. In *21th IEEE Symposium on Logic in Computer Science (LICS 2006), 12-15*
515 *August 2006, Seattle, WA, USA, Proceedings*, pages 211–220. IEEE Computer Society, 2006.
516 doi:10.1109/LICS.2006.24.
- 517 8 José Carlos Costa. Some pseudovariety joins involving locally trivial semigroups. In *Semigroup*
518 *Forum*, volume 64, pages 12–28. Springer, 2001.
- 519 9 José Carlos Costa. Some pseudovariety joins involving groups and locally trivial semigroups.
520 In *Semigroups, Algorithms, Automata And Languages*, pages 341–348. World Scientific, 2002.
- 521 10 Samuel Eilenberg. *Automata, Languages, and Machines*, volume A. Academic Press, New
522 York, 1974.
- 523 11 Samuel Eilenberg. *Automata, Languages, and Machines*, volume B. Academic Press, New
524 York, 1976.
- 525 12 Nathan Grosshans, Pierre McKenzie, and Luc Segoufin. Tameness and the power of programs
526 over monoids in DA. *CoRR*, abs/2101.07495, 2021. URL: [https://arxiv.org/abs/2101.](https://arxiv.org/abs/2101.07495)
527 [07495](https://arxiv.org/abs/2101.07495), arXiv:2101.07495.
- 528 13 Jean-Éric Pin. *Varieties of formal languages*. North Oxford, London and Plenum, New-York,
529 1986. (Traduction de Variétés de langages formels).
- 530 14 Jean-Éric Pin and Howard Straubing. Some results on C-varieties. *RAIRO-Theor. Inf. Appl.*,
531 39(1):239–262, 2005.
- 532 15 Jan Reiterman. The Birkhoff theorem for finite algebras. *algebra universalis*, 14(1):1–10, 1982.
- 533 16 Marcel Paul Schützenberger. On finite monoids having only trivial subgroups. *Inf. Control.*,
534 8(2):190–194, 1965. doi:10.1016/S0019-9958(65)90108-7.
- 535 17 Imre Simon. Piecewise testable events. In *Automata Theory and Formal Languages, 2nd GI*
536 *Conference*, volume 33 of *Lecture Notes in Computer Science*, pages 214–222. Springer, 1975.
- 537 18 Howard Straubing. On logical descriptions of regular languages. In Sergio Rajsbaum, editor,
538 *LATIN 2002: Theoretical Informatics, 5th Latin American Symposium, Cancun, Mexico, April*
539 *3-6, 2002, Proceedings*, volume 2286 of *Lecture Notes in Computer Science*, pages 528–538.
540 Springer, 2002. doi:10.1007/3-540-45995-2_46.
- 541 19 Marc Zeitoun. The join of the pseudovarieties of idempotent semigroups and locally trivial
542 semigroups. In *Semigroup Forum*, volume 50, pages 367–381. Springer, 1995.
- 543 20 Marc Zeitoun. On the decidability of the membership problem of the pseudovariety JVB. *Int.*
544 *J. Algebra Comput.*, 5(1):47–64, 1995. doi:10.1142/S0218196795000057.
- 545 21 Marc Zeitoun. On the join of two pseudovarieties. *Semigroups, Automata and Languages, eds.*
546 *J. Almeida, GMS Gomes, and PV Silva. World Scientific*, pages 281–288, 1996.

547 **A** **Missing proofs**

548 **Proof of Proposition 2.** Let \mathbf{W} be some ne -variety of stamps such that $\langle \mathbf{V} \rangle_{all} \cup \langle \mathbf{LI} \rangle_{ne} \subseteq \mathbf{W}$.
549 There exists some variety of semigroups \mathbf{W}' such that $\langle \mathbf{W}' \rangle_{ne} = \mathbf{W}$.

550 Let $S \in \mathbf{V} \cup \mathbf{LI}$. We denote by S^1 the monoid S if S is already a monoid and the monoid
 551 $S \cup \{1\}$ otherwise. Then the evaluation morphism $\eta_S: S^* \rightarrow S^1$ such that $\eta_S(s) = s$ for
 552 all $s \in S$ verifies $\eta_S(S^+) = S$ and additionally $S^1 = S$ when $S \in \mathbf{V}$. This implies that
 553 $\eta_S \in \langle \mathbf{V} \rangle_{all} \cup \langle \mathbf{LI} \rangle_{ne} \subseteq \mathbf{W}$. But by definition of \mathbf{W}' , it must be that $S = \eta_S(S^+) \in \mathbf{W}'$.

554 Therefore, \mathbf{W}' contains both \mathbf{V} and \mathbf{LI} , which implies that $\mathbf{V} \vee \mathbf{LI} \subseteq \mathbf{W}'$ by inclusion-wise
 555 minimality of $\mathbf{V} \vee \mathbf{LI}$. By definition, we can then conclude that $\langle \mathbf{V} \vee \mathbf{LI} \rangle_{ne} \subseteq \langle \mathbf{W}' \rangle_{ne} = \mathbf{W}$.
 556 So $\langle \mathbf{V} \vee \mathbf{LI} \rangle_{ne}$ is the inclusion-wise smallest ne -variety of stamps containing both $\langle \mathbf{V} \rangle_{all}$ and
 557 $\langle \mathbf{LI} \rangle_{ne}$.

558 Let now \mathcal{W} be some ne -variety of languages such that $\mathcal{L}(\mathbf{V}) \cup \mathcal{L}(\mathbf{LI}) \subseteq \mathcal{W}$. It holds that
 559 $\mathcal{W} = \mathcal{L}(\mathbf{W})$ for some ne -variety of stamps \mathbf{W} . We have that $\langle \mathbf{V} \rangle_{all}$, which is in particular an
 560 ne -variety of stamps, is included in \mathbf{W} because $\mathcal{L}(\langle \mathbf{V} \rangle_{all}) = \mathcal{L}(\mathbf{V}) \subseteq \mathcal{W} = \mathcal{L}(\mathbf{W})$, but also
 561 that $\langle \mathbf{LI} \rangle_{ne}$ is included in \mathbf{W} because $\mathcal{L}(\langle \mathbf{LI} \rangle_{ne}) = \mathcal{L}(\mathbf{LI}) \subseteq \mathcal{W} = \mathcal{L}(\mathbf{W})$. By inclusion-wise
 562 minimality of $\langle \mathbf{V} \vee \mathbf{LI} \rangle_{ne}$, it follows that $\langle \mathbf{V} \vee \mathbf{LI} \rangle_{ne} \subseteq \mathbf{W}$. Hence, using again the above
 563 fact on the Eilenberg correspondence, we can conclude that $\mathcal{L}(\mathbf{V} \vee \mathbf{LI}) = \mathcal{L}(\langle \mathbf{V} \vee \mathbf{LI} \rangle_{ne}) \subseteq$
 564 $\mathcal{L}(\mathbf{W}) = \mathcal{W}$. So $\mathcal{L}(\mathbf{V} \vee \mathbf{LI})$ is the inclusion-wise smallest ne -variety of languages containing
 565 both $\mathcal{L}(\mathbf{V})$ and $\mathcal{L}(\mathbf{LI})$.

566 Consider now the class of languages \mathcal{C} such that $\mathcal{C}(\Sigma)$ is the Boolean closure of $\mathcal{L}(\mathbf{V})(\Sigma) \cup$
 567 $\mathcal{L}(\mathbf{LI})(\Sigma)$ for each alphabet Σ . By closure under Boolean operations of $\mathcal{L}(\mathbf{V} \vee \mathbf{LI})$, we
 568 have that $\mathcal{C} \subseteq \mathcal{L}(\mathbf{V} \vee \mathbf{LI})$. Now, as Boolean operations commute with both quotients [13,
 569 p. 20] and inverses of ne -morphisms [13, Proposition 0.4], by closure of $\mathcal{L}(\mathbf{V})$ and $\mathcal{L}(\mathbf{LI})$
 570 under quotients and inverses of ne -morphisms, we actually have that \mathcal{C} is an ne -variety
 571 of languages. Therefore, by inclusion-wise minimality of $\mathcal{L}(\mathbf{V} \vee \mathbf{LI})$, we can conclude that
 572 $\mathcal{L}(\mathbf{V} \vee \mathbf{LI}) = \mathcal{C}$. ◀

573 **Proof of Lemma 10.** We actually have that $\mathcal{L}(\mathbf{V}) \cup \mathcal{L}(\mathbf{LI}) \subseteq \mathcal{L}(\mathbf{EV})$, which allows us to
 574 conclude by inclusion-wise minimality of $\mathcal{L}(\mathbf{V} \vee \mathbf{LI})$ (Proposition 2) and by the fact that
 575 $\mathcal{L}(\mathbf{EV})$ is an ne -variety of languages (Proposition 6).

576 Let Σ be an alphabet. The fact that $\mathcal{L}(\mathbf{V})(\Sigma) \subseteq \mathcal{L}(\mathbf{EV})(\Sigma)$ follows trivially from
 577 Proposition 9. Moreover, for all $u \in \Sigma^*$, since necessarily $\Sigma^* \in \mathcal{L}(\mathbf{V})(\Sigma)$, we have that both
 578 $u\Sigma^*$ and Σ^*u belong to $\mathcal{L}(\mathbf{LI})(\Sigma)$. Thus, as $\mathcal{L}(\mathbf{EV})(\Sigma)$ is closed under Boolean operations,
 579 it follows that $\mathcal{L}(\mathbf{LI})(\Sigma) \subseteq \mathcal{L}(\mathbf{EV})(\Sigma)$.

580 This concludes the proof, since it holds for any alphabet Σ . ◀

581 **Proof of Lemma 11.** Assume that $\mathbf{EV} \subseteq \langle \mathbf{V} \vee \mathbf{LI} \rangle_{ne}$. For any $L \in \mathcal{L}(\mathbf{V})(\Sigma)$ for Σ some al-
 582 phabet and any $x, y \in \Sigma^*$, by Proposition 9, we have that $xLy \in \mathcal{L}(\mathbf{EV})(\Sigma) \subseteq \mathcal{L}(\mathbf{V} \vee \mathbf{LI})(\Sigma)$.
 583 Hence, \mathbf{V} verifies criterion (A).

584 Conversely, assume that \mathbf{V} verifies criterion (A). For any alphabet Σ , the set $\mathcal{L}(\mathbf{V} \vee \mathbf{LI})(\Sigma)$
 585 contains all languages of the form xLy for $L \in \mathcal{L}(\mathbf{V})(\Sigma)$ and $x, y \in \Sigma^*$, so it contains all
 586 Boolean combinations of languages of that form, since it is closed under Boolean operations.
 587 Therefore, by Proposition 9, we have $\mathcal{L}(\mathbf{EV}) \subseteq \mathcal{L}(\mathbf{V} \vee \mathbf{LI})$, so that $\mathbf{EV} \subseteq \langle \mathbf{V} \vee \mathbf{LI} \rangle_{ne}$. ◀

588 **Proof of Proposition 12.** Let us first observe that given any alphabet Σ , given any language
 589 K on that alphabet and given any two words $x, y \in \Sigma^*$, we have that $x(x^{-1}Ky^{-1})y =$
 590 $x\Sigma^*y \cap K$ and $x^{-1}(xKy)y^{-1} = K$.

591 **Implication from right to left.** Assume that for any $L \in \mathcal{L}(\mathbf{V})(\Sigma)$ for some alphabet Σ and
 592 $x, y \in \Sigma^*$, there exist $k, l \in \mathbb{N}$ such that for all $u \in \Sigma^k, v \in \Sigma^l$, there exists some language
 593 $K \in \mathcal{L}(\mathbf{V})(\Sigma)$ verifying $u^{-1}Lv^{-1} = (xu)^{-1}K(vy)^{-1}$. Take some $L \in \mathcal{L}(\mathbf{V})(\Sigma)$ for some

594 alphabet Σ and take $x, y \in \Sigma^*$. Consider also some $k, l \in \mathbb{N}$ that are guaranteed to exist by
 595 the assumption we just made.

596 For all $u \in \Sigma^k, v \in \Sigma^l$, there exists some language $K \in \mathcal{L}(\mathbf{V})(\Sigma)$ verifying $u^{-1}Lv^{-1} =$
 597 $(xu)^{-1}K(vy)^{-1}$, so that by our observation at the beginning of the proof, we have

$$598 \quad x(u\Sigma^*v \cap L)y = xu(u^{-1}Lv^{-1})vy = xu((xu)^{-1}K(vy)^{-1})vy = xu\Sigma^*vy \cap K .$$

599 Using Proposition 2, we thus have that $x(u\Sigma^*v \cap L)y \in \mathcal{L}(\mathbf{V} \vee \mathbf{LI})(\Sigma)$ for all $u \in \Sigma^k, v \in \Sigma^l$.
 600 Moreover, since we have that the set of words of L of length at least $k + l$ is

$$601 \quad \Sigma^{\geq k+l} \cap L = \bigcup_{u \in \Sigma^k, v \in \Sigma^l} (u\Sigma^*v \cap L)$$

602 and since

$$603 \quad L = (\Sigma^{\geq k+l} \cap L) \cup F$$

604 where F is a finite set of words on Σ of length less than $k + l$, we have that

$$605 \quad xLy = x((\Sigma^{\geq k+l} \cap L) \cup F)y = \bigcup_{u \in \Sigma^k, v \in \Sigma^l} x(u\Sigma^*v \cap L)y \cup xFy .$$

606 We can thus conclude that $xLy \in \mathcal{L}(\mathbf{V} \vee \mathbf{LI})(\Sigma)$ since $xFy \in \mathcal{L}(\mathbf{LI})(\Sigma)$ and because
 607 $\mathcal{L}(\mathbf{V} \vee \mathbf{LI})(\Sigma)$ is closed under unions.

608 **Implication from left to right.** Assume that \mathbf{V} satisfies criterion (A). Take some $L \in$
 609 $\mathcal{L}(\mathbf{V})(\Sigma)$ for some alphabet Σ and take $x, y \in \Sigma^*$. By hypothesis, we know that $xLy \in$
 610 $\mathcal{L}(\mathbf{V} \vee \mathbf{LI})(\Sigma)$.

611 By Proposition 2, this means that xLy is a Boolean combination of languages in $\mathcal{L}(\mathbf{V})(\Sigma) \cup$
 612 $\mathcal{L}(\mathbf{LI})(\Sigma)$. Further, this implies that xLy can be written as the union of intersections of
 613 languages of $\mathcal{L}(\mathbf{V})(\Sigma)$ and $\mathcal{L}(\mathbf{LI})(\Sigma)$ or their complements, which in turn implies, by closure
 614 of $\mathcal{L}(\mathbf{V})(\Sigma)$ and $\mathcal{L}(\mathbf{LI})(\Sigma)$ under Boolean operations, that xLy can be written as a finite
 615 union of languages of the form $K \cap (U\Sigma^*V \cup W)$ with $K \in \mathcal{L}(\mathbf{V})(\Sigma)$ and $U, V, W \subseteq \Sigma^*$ finite.
 616 Since any word in xLy must be of length at least $|xy|$ and have x as a prefix and y as a
 617 suffix, we can assume that any language $K \cap (U\Sigma^*V \cup W)$ appearing in a finite union as
 618 described above verifies that $U \subseteq x\Sigma^*$, that $V \subseteq \Sigma^*y$ and that $W \subseteq x\Sigma^*y$. Now, if we take
 619 $k, l \in \mathbb{N}$ big enough, we thus have that

$$620 \quad xLy = \bigcup_{u \in \Sigma^k, v \in \Sigma^l} (K_{u,v} \cap xu\Sigma^*vy) \cup F$$

621 where $K_{u,v} \in \mathcal{L}(\mathbf{V})(\Sigma)$ for all $u \in \Sigma^k, v \in \Sigma^l$ and $F \subseteq \Sigma^{<|xy|+k+l}$. Hence, for all $u \in \Sigma^k, v \in$
 622 Σ^l , we have

$$\begin{aligned} 623 \quad u^{-1}Lv^{-1} &= u^{-1}(x^{-1}(xLy)y^{-1})v^{-1} \\ 624 \quad &= (xu)^{-1} \left(\bigcup_{u' \in \Sigma^k, v' \in \Sigma^l} (K_{u',v'} \cap xu'\Sigma^*v'y) \cup F \right) (vy)^{-1} \\ 625 \quad &= \bigcup_{u' \in \Sigma^k, v' \in \Sigma^l} (xu)^{-1} \left(xu'((xu')^{-1}K_{u',v'}(v'y)^{-1})v'y \right) (vy)^{-1} \cup \\ &\quad (xu)^{-1}F(vy)^{-1} \\ 626 \quad &= (xu)^{-1}K_{u,v}(vy)^{-1} , \end{aligned}$$

628 using classical formulae for quotients [13, p. 20] and observing that $(xu)^{-1}K(vy)^{-1} = \emptyset$ for
 629 any $K \subseteq \Sigma^*$ such that $K \cap xu\Sigma^*vy = \emptyset$. ◀