

# A Note on the Join of Varieties of Monoids with LI Nathan Grosshans

## ▶ To cite this version:

Nathan Grosshans. A Note on the Join of Varieties of Monoids with LI. 2021. hal-03183838v2

## HAL Id: hal-03183838 https://hal.science/hal-03183838v2

Preprint submitted on 10 May 2021 (v2), last revised 27 Aug 2021 (v3)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

### 3 Nathan Grosshans 🖂 🏠 💿

<sup>4</sup> Fachbereich Elektrotechnik/Informatik, Universität Kassel, Kassel, Germany

#### 5 — Abstract

In this note, we give a characterisation in terms of identities of the join of V with the variety of finite
locally trivial semigroups LI for several well-known varieties of finite monoids V by using classical
algebraic-automata-theoretic techniques. To achieve this, we use the new notion of essentially-V
stamps defined by Grosshans, McKenzie and Segoufin and show that it actually coincides with the
join of V and LI precisely when some natural condition on the variety of languages corresponding
to V is verified.

This work is a kind of rediscovery of the work of J. C. Costa around 20 years ago from a rather different angle, since Costa's work relies on the use of advanced developments in profinite topology, whereas what is presented here essentially uses an algebraic, language-based approach.

<sup>15</sup> 2012 ACM Subject Classification Theory of computation  $\rightarrow$  Formal languages and automata theory; <sup>16</sup> Theory of computation  $\rightarrow$  Algebraic language theory

17 Keywords and phrases Varieties of monoids, join, LI

### 18 **1** Introduction

One of the most fundamental problems appearing in finite automata theory is the one of 19 characterisation: given some subclass of the class of regular languages, find out whether there 20 is a way to characterise those languages using some class of finite objects. This problem is 21 often linked to and motivated by the problem of *decidability*: given some subclass of the class 22 of regular languages, find out whether there exists an algorithm testing the membership of 23 any regular language in that subclass. The obvious approach to try to find a characterisation 24 of a class of regular languages would be to look for properties shared by all the minimal 25 finite automata of those languages. If we find such characterising properties, we can then 26 ask whether they can be checked by an algorithm to answer the problem of decidability 27 for this class of languages. However, one of the most fruitful approaches in face of those 28 two problems has been the *algebraic approach*, in which we basically replace automata with 29 morphisms into monoids: a language L over some alphabet  $\Sigma$  is then said to be recognised by 30 a morphism  $\varphi$  into a monoid M if and only if L is the inverse image by  $\varphi$  of a subset of M. 31 Under this notion of recognition, each language has a minimal morphism recognising it, the 32 syntatic morphism into the syntactic monoid of that language, that are minimal under some 33 notion of division. The fundamental result on which this algebraic approach relies is that a 34 language is regular if and only if its syntactic monoid is finite. One can thus try to find a 35 characterisation of some class of regular languages by looking at the algebraic properties of 36 the syntactic monoids of these languages. 37

And many such characterisations that are decidable were indeed successfully obtained 38 since Schützenberger's seminal work in 1965 [16]. His famous result, that really started the 39 field of algebraic automata theory, states that the star-free regular languages are exactly 40 those whose syntactic monoids are finite and aperiodic. Another important early result in 41 that vain is the one of Simon [17] characterising the piecewise testable languages as exactly 42 those having a finite  $\mathfrak{J}$ -trivial syntactic monoid. Eilenberg [11] was the first to prove that 43 such algebraic characterisations actually come as specific instances of a general bijective 44 correspondence between varieties of finite monoids and varieties of languages — classes 45

of, respectively, finite monoids and regular languages closed under some respective natural 46 operations. Thus, a class of regular languages can indeed be characterised by the syntactic 47 monoids of these languages, as soon as it verifies some nice properties. Eilenberg's result was 48 later completed by Reiterman's theorem [15], that uses a notion of identities defined using 49 profinite topology and states that a class of finite monoids is a variety of finite monoids if and 50 only if it is defined by a set of profinite identities. Therefore, one can always characterise the 51 variety of finite monoids associated to a variety of languages by a set of profinite identities 52 and, additionally, this characterisation often leads to decidability, especially when this set is 53 finite. Several research works have been conducted to characterise varieties of finite monoids 54 or semigroups by profinite identities. 55

A kind of varieties of finite monoids or semigroups that has attracted many research 56 efforts aiming for characterisations through identities are the varieties defined as the join 57 of two other varieties. Given two varieties of finite monoids  $\mathbf{V}$  and  $\mathbf{W}$ , the join of  $\mathbf{V}$ 58 and  $\mathbf{W}$ , denoted by  $\mathbf{V} \lor \mathbf{W}$ , is the smallest variety of finite monoids containing both  $\mathbf{V}$ 59 and W. One of the main motivations to try to understand  $\mathbf{V} \vee \mathbf{W}$  is that the variety 60 of languages corresponding to it by the Eilenberg correspondence,  $\mathcal{L}(\mathbf{V} \vee \mathbf{W})$ , is the one 61 obtained by considering parallel computations of automata recognising languages from both 62  $\mathcal{L}(\mathbf{V})$  and  $\mathcal{L}(\mathbf{W})$ , the varieties of languages corresponding to, respectively, V and W. This 63 is a fundamental operation on automata, and while it is straightforward that  $\mathcal{L}(\mathbf{V} \lor \mathbf{W})$ 64 is simply the smallest variety of languages containing both  $\mathcal{L}(\mathbf{V})$  and  $\mathcal{L}(\mathbf{W})$ , this does 65 not at all furnish a decidable characterisation of  $\mathcal{L}(\mathbf{V} \lor \mathbf{W})$ , let alone a set of identities 66 defining  $\mathbf{V} \vee \mathbf{W}$ . Generally speaking, the problem of finding a set of identities defining 67  $\mathbf{V} \vee \mathbf{W}$  is difficult (see [2, 21]), but it has been solved for many specific cases: have a look 68 at [1, 3, 5, 20, 19, 6, 8, 9] for some examples. 69

In this paper, we give a general method to find a set of identities defining the join of an 70 arbitrary variety of finite monoids  $\mathbf{V}$  and the variety of finite locally trivial semigroups  $\mathbf{LI}$ , as 71 soon as one has a set of identities defining V and V verifies some criterion. Joins of that sort 72 have been studied quite a lot in the literature we mentioned in the previous paragraph (e.g. 73 in [5, 19, 8, 9]), but while these works usually rely heavily on profinite topology with some 74 in-depth understanding of the structure of the elements of the so-called free pro-V monoids 75 and free pro-LI semigroups, we present a method that reduces the use of profinite topology 76 to the minimum and that relies mainly on algebraic and language-theoretic techniques. The 77 variety LI is well-known to correspond to the class of languages for which membership only 78 depends on bounded-length prefixes and suffixes of words. In [12], McKenzie, Segoufin and 79 80 the author introduced the notion of essentially-V stamps (surjective morphisms  $\varphi \colon \Sigma^* \to M$ for  $\Sigma$  an alphabet and M a finite monoid) to characterise the built-in ability that programs 81 over monoids in  $\mathbf{V}$  have to treat separately some constant-length beginning and ending of a 82 word. Informally said, a stamp is essentially- $\mathbf{V}$  when it behaves like a stamp into a monoid 83 of  $\mathbf{V}$  as soon as a sufficiently long beginning and ending of the input word has been fixed. 84 Our method builds on two results, that we prove in this article. 85

1. The first result is a characterisation in terms of identities of the class **EV** of essentially-**V** stamps given a set of identites E defining **V**: a stamp is in **EV** if and only if it satisfies all identities  $x^{\omega}yuzt^{\omega} = x^{\omega}yvzt^{\omega}$  for u = v an identity in E and where x, y, z, t do appear neither in u nor in v.

<sup>90</sup> 2. The second result says that **EV** and **V**  $\vee$  **LI** do correspond if and only if **V** verifies some <sup>91</sup> criterion, that can be formulated in terms of quotient-expressibility in  $\mathcal{L}(\mathbf{V})$ : any language <sup>92</sup>  $L \in \mathcal{L}(\mathbf{V})$  must, for an arbitrary choice of x, y, be such that the quotient  $u^{-1}Lv^{-1}$  for u<sup>93</sup> and v long enough can be expressed as the quotient  $(xu)^{-1}K(vy)^{-1}$  for some  $K \in \mathcal{L}(\mathbf{V})$ .

<sup>94</sup> Using these results, we can find a set of identities defining the so-called *ne*-variety of stamps <sup>95</sup> corresponding to  $\mathbf{V} \lor \mathbf{LI}$  as soon as a set of identities defining  $\mathbf{V}$  is known by proving that  $\mathbf{V}$ <sup>96</sup> verifies the criterion in point 2. Note that, for technical reasons, we work with the *ne*-variety <sup>97</sup> of stamps corresponding  $\mathbf{V} \lor \mathbf{LI}$  rather than directly with the variety of finite semigroups <sup>98</sup>  $\mathbf{V} \lor \mathbf{LI}$ , but this is not a problem since they correspond bijectively. We apply this method <sup>99</sup> to reprove characterisations of the join of  $\mathbf{LI}$  with each of the well-known varieties of finite <sup>100</sup> monoids  $\mathbf{R}$ ,  $\mathbf{L}$ ,  $\mathbf{J}$  and any variety of finite groups.

The author noticed after proving those results that his work actually forms a kind of rediscovery of the work of J. C. Costa in [8]. He defines an operator U associating to each set of identities E the exact same new set U(E) of identities as in point 1. Costa then defines a property of cancellation for varieties of finite semigroups such that for any  $\mathbf{V}$  verifying it, U(E) defines  $\mathbf{V} \vee \mathbf{LI}$  for E defining  $\mathbf{V}$ . He finally uses this result to derive characterisations of  $\mathbf{V} \vee \mathbf{LI}$  for all the cases we are treating in our paper and many more.

<sup>107</sup> What is, then, the contribution of our article? In a nutshell, it does mainly use algebraic <sup>108</sup> and language-theoretic techniques while Costa's work relies heavily on profinite topology. In <sup>109</sup> our setting, once the stage is set, all proofs are quite straightforward without real difficulties <sup>110</sup> and rely on classical language-theoretic characterisations of the varieties under consideration. <sup>111</sup> This is to contrast with Costa's work, that for instance draws upon the difficult analysis of <sup>112</sup> the elements of free pro-**R** monoids by Almeida and Weil [4] to characterise **R**  $\lor$  **LI**.

<sup>113</sup> **Organisation of the article.** Section 2 is dedicated to the necessary preliminaries. In <sup>114</sup> Section 3, we recall the definition of essentially-V stamps and prove the characterisation <sup>115</sup> by identities of point 1 above. Section 4 is then dedicated to the necessary and sufficient <sup>116</sup> criterion for **EV** and **V**  $\vee$  **LI** to correspond presented in point 2 and finally those results are <sup>117</sup> applied to specific cases in Section 5. We finish with a short conclusion.

## <sup>118</sup> 2 Preliminaries

We briefly introduce the mathematical material used in this paper. For the basics and the classical results of automata theory, we refer the reader to the two classical references of the domain by Eilenberg [10, 11] and Pin [13]. For definitions and results specific to varieties of stamps and associated profinite identities, see the articles by Straubing [18] and by Pin and Straubing [14]. We also assume some basic knowledge of topology.

General notations. Let  $i \in \mathbb{N}$  be a natural number. We shall denote by [i] the set of all natural numbers  $n \in \mathbb{N}$  verifying  $1 \le n \le i$ .

**Words and languages.** Let  $\Sigma$  be a finite alphabet. We denote by  $\Sigma^*$  the set of all finite words over  $\Sigma$ . We also denote by  $\Sigma^+$  the set of all finite non empty words over  $\Sigma$ , the empty word being denoted by  $\varepsilon$ . All our alphabets and words will always be finite, without further mention of this fact. Given some word  $w \in \Sigma^*$ , we denote its length by |w| and the set of letters it contains by alph(w). Given some  $n \in \mathbb{N}$ , we denote by  $\Sigma^{\geq n}$ ,  $\Sigma^n$  and  $\Sigma^{< n}$  the set of words over  $\Sigma$  of length, respectively, at least n, exactly n and less than n.

<sup>132</sup> A language over  $\Sigma$  is a subset of  $\Sigma^*$ . A language is regular if it is recognised by a <sup>133</sup> deterministic finite automaton. The quotient of a language L over  $\Sigma$  relative to the words u<sup>134</sup> and v over  $\Sigma$  is the language, denoted by  $u^{-1}Lv^{-1}$ , of the words w such that  $uwv \in L$ .

**Monoids, semigroups and varieties.** A *semigroup* is a non-empty set equipped with an 135 associative law that we will write multiplicatively. A monoid is a semigroup with an identity. 136 An example of a semigroup is  $\Sigma^+$ , the free semigroup over  $\Sigma$ . Similarly  $\Sigma^*$  is the free monoid 137 over  $\Sigma$ . A morphism  $\varphi$  from a semigroup S to a semigroup T is a function from S to T such 138 that  $\varphi(xy) = \varphi(x)\varphi(y)$  for all  $x, y \in S$ . A morphism of monoids additionally requires that 139 the identity is preserved. A semigroup T is a subsemigroup of a semigroup S if T is a subset 140 of S and is equipped with the restricted law of S. Additionally the notion of submonoids 141 requires the presence of the identity. A semigroup T divides a semigroup S if T is the image 142 by a semigroup morphism of a subsemigroup of S. Division of monoids is defined in the same 143 way. The *Cartesian (or direct) product* of two semigroups is simply the semigroup given by 144 the Cartesian product of the two underlying sets equipped with the Cartesian product of 145 their laws. An element s of a semigroup is *idempotent* if ss = s. 146

A variety of finite monoids is a non-empty class of finite monoids closed under Cartesian product and monoid division. A variety of finite semigroups is defined similarly. When dealing with varieties, we consider only finite monoids and semigroups, so we will drop the adjective finite when talking about varieties in the rest of this article.

<sup>151</sup> Varieties of stamps Let  $f: \Sigma^* \to \Gamma^*$  be a morphism from the free monoid over an alphabet <sup>152</sup>  $\Sigma$  to the free monoid over an alphabet  $\Gamma$ , that we might call an *all*-morphism. We say that <sup>153</sup> f is an ne-*morphism* (non-erasing morphism) whenever  $f(\Sigma) \subseteq \Gamma^+$ .

We call stamp a surjective morphism  $\varphi \colon \Sigma^* \to M$  for  $\Sigma$  an alphabet and M a finite monoid. We say that a stamp  $\varphi \colon \Sigma^* \to M$  all-divides (respectively ne-divides) a stamp  $\psi \colon \Gamma^* \to N$  whenever there exists an all-morphism (respectively ne-morphism)  $f \colon \Sigma^* \to \Gamma^*$ and a surjective morphism  $\alpha \colon \mathfrak{Im}(\psi \circ f) \to M$  such that  $\varphi = \alpha \circ \psi \circ f$ . The direct product of two stamps  $\varphi \colon \Sigma^* \to M$  and  $\psi \colon \Gamma^* \to N$  is the stamp  $\varphi \times \psi \colon \Sigma^* \to K$  such that K is the submonoid of  $M \times N$  generated by  $\{(\varphi(a), \psi(a)) \mid a \in \Sigma\}$  and  $\varphi \times \psi(a) = (\varphi(a), \psi(a))$  for all  $a \in \Sigma$ .

An all-variety of stamps (respectively ne-variety of stamps) is a non-empty class of stamps closed under direct product and all-division (respectively *ne*-division).

We will often use the following characteristic index of stamps, defined in [7]. Consider some stamp  $\varphi \colon \Sigma^* \to M$ . As M is finite there is a  $k \in \mathbb{N}_{>0}$  such that  $\varphi(\Sigma^{2k}) = \varphi(\Sigma^k)$ : this implies that  $\varphi(\Sigma^k)$  is a semigroup. The smallest such k is called the *stability index* of  $\varphi$ .

**Varieties of languages.** A language L over some alphabet  $\Sigma$  is recognised by a monoid M 166 if there is a morphism  $\varphi \colon \Sigma^* \to M$  and  $F \subseteq M$  such that  $L = \varphi^{-1}(F)$ . We also say that  $\varphi$ 167 recognises L. It is well known that a language is regular if and only if it is recognised by a 168 finite monoid. The syntactic congruence of L, denoted by  $\sim_L$ , is the equivalence relation 169 on  $\Sigma^*$  defined by  $u \sim_L v$  for  $u, v \in \Sigma^*$  whenever for all  $x, y \in \Sigma^*$ ,  $xuy \in L$  if and only if 170  $xvy \in L$ . The quotient  $\Sigma^*/\sim_L$  is a monoid, called the syntactic monoid of L, that recognises 171 L via the syntactic morphism  $\eta_L$  of L sending any word u to its equivalence class  $[u]_{\sim_L}$  for 172  $\sim_L$ . A stamp  $\varphi \colon \Sigma^* \to M$  recognises L if and only if there exists a surjective morphism 173  $\varphi \colon M \to \Sigma^* / \sim_L$  verifying  $\eta_L = \alpha \circ \varphi$ . 174

A class of languages C is a correspondence that associates a set  $C(\Sigma)$  to each alphabet  $\Sigma$ . A (all-)variety of languages (respectively an ne-variety of languages)  $\mathcal{V}$  is a non-empty class of regular languages closed under Boolean operations, quotients and inverses of all-morphisms (respectively *ne*-morphisms). A classical result of Eilenberg [11, Chapter VII, Section 3] rays that there is a bijective correspondence between varieties of monoids and varieties of languages: to each variety of monoids  $\mathbf{V}$  we can bijectively associate  $\mathcal{L}(\mathbf{V})$  the variety of

<sup>181</sup> languages whose syntactic monoids belong to **V**. This was generalised by Straubing [18] <sup>182</sup> to the case of varieties of stamps: to each *all*-variety (respectively *ne*-variety) of stamps <sup>183</sup> **V** we can bijectively associate  $\mathcal{L}(\mathbf{V})$  the *all*-variety (respectively *ne*-variety) of languages <sup>184</sup> whose syntactic morphisms belong to **V**. Given two *all*-varieties (respectively *ne*-varieties) <sup>185</sup> of stamps  $\mathbf{V_1}$  and  $\mathbf{V_2}$ , we have  $\mathbf{V_1} \subseteq \mathbf{V_2} \Leftrightarrow \mathcal{L}(\mathbf{V_1}) \subseteq \mathcal{L}(\mathbf{V_2})$ .

For **V** some variety of monoids, we define  $\langle \mathbf{V} \rangle_{all}$  the *all*-variety of all stamps  $\varphi \colon \Sigma^* \to M$ such that  $M \in \mathbf{V}$ . Of course, in that case  $\mathcal{L}(\mathbf{V}) = \mathcal{L}(\langle \mathbf{V} \rangle_{all})$ . Similarly, for **V** some variety of semigroups, we define  $\langle \mathbf{V} \rangle_{ne}$  the *ne*-variety of all stamps  $\varphi \colon \Sigma^* \to M$  such that  $\varphi(\Sigma^+) \in \mathbf{V}$ . In that case, we consider  $\mathcal{L}(\mathbf{V})$  to be the *ne*-variety of languages corresponding to  $\langle \mathbf{V} \rangle_{ne}$ . The operations  $\langle \cdot \rangle_{all}$  and  $\langle \cdot \rangle_{ne}$  form bijective correspondences (see [18]).

<sup>191</sup> **Identities.** Every finite monoid M is considered to be a complete metric space equipped <sup>192</sup> with the discrete metric d defined by  $d(m,n) = \begin{cases} 0 & \text{if } m = n \\ 1 & \text{otherwise} \end{cases}$  for all  $m, n \in M$ . <sup>193</sup> Let  $\Sigma$  be an alphabet. Given  $u, v \in \Sigma^*$ , we set

$$_{^{194}} \qquad r(u,v) = \min\{|M| \mid \exists \varphi \colon \Sigma^* \to M \text{ stamp s.t. } \varphi(u) \neq \varphi(v)\}$$

and  $d(u,v) = 2^{-r(u,v)}$ , using the conventions that  $\min \emptyset = +\infty$  and  $2^{-\infty} = 0$ . Then d 195 is a metric on  $\Sigma^*$ . The completion of the metric space  $(\Sigma^*, d)$ , denoted by  $(\widehat{\Sigma^*}, \widehat{d})$ , is a 196 metric monoid called the *free profinite monoid on*  $\Sigma^*$ . Its elements are all the formal 197 limits  $\lim_{n\to\infty} x_n$  of Cauchy sequences  $(x_n)_{n\geq 0}$  in  $(\Sigma^*, d)$  and the metric d on  $\Sigma^*$  extends 198 to a metric  $\widehat{d}$  on  $\widehat{\Sigma^*}$  defined by  $\widehat{d}(\lim_{n\to\infty} x_n, \lim_{n\to\infty} y_n) = \lim_{n\to\infty} d(x_n, y_n)$  for Cauchy 199 sequences  $(x_n)_{n\geq 0}$  and  $(y_n)_{n\geq 0}$  in  $(\Sigma^*, d)$ . Note that, when it is clear from the context, we 200 usually do not make the metric explicit when talking about a metric space. One important 201 example of elements of  $\widehat{\Sigma^*}$  is given by the elements  $x^{\omega} = \lim_{n \to \infty} x^{n!}$  for all  $x \in \Sigma^*$ . 202

Every stamp  $\varphi \colon \Sigma^* \to M$  extends uniquely to a uniformly continuous morphism  $\widehat{\varphi} \colon \widehat{\Sigma^*} \to M$ 203 M with  $\widehat{\varphi}(\lim_{n\to\infty} x_n) = \lim_{n\to\infty} \varphi(x_n)$  for every Cauchy sequence  $(x_n)_{n>0}$  in  $\Sigma^*$ . Similarly, 204 every all-morphism  $f: \Sigma^* \to \Gamma^*$  extends uniquely to a uniformly continuous morphism 205  $\widehat{f}: \widehat{\Sigma^*} \to \widehat{\Gamma^*}$  with  $\widehat{f}(\lim_{n \to \infty} x_n) = \lim_{n \to \infty} f(x_n)$  for every Cauchy sequence  $(x_n)_{n \ge 0}$  in  $\Sigma^*$ . 206 For  $u, v \in \widehat{A^*}$  with A an alphabet, we say that a stamp  $\Sigma^* \to M$  all-satisfies (respectively 207 ne-satisfies) the identity u = v if for every all-morphism (respectively ne-morphism)  $f: A^* \to A^*$ 208  $\Sigma^*$ , it holds that  $\widehat{\varphi} \circ \widehat{f}(u) = \widehat{\varphi} \circ \widehat{f}(v)$ . Given some set of identities E, we denote by  $\llbracket E \rrbracket_{all}$ 209 (respectively  $\llbracket E \rrbracket_{ne}$ ) the class of stamps *all*-satisfying (respectively *ne*-satisfying) all the 210 identities of E. When  $\llbracket E \rrbracket_{all}$  (respectively  $\llbracket E \rrbracket_{ne}$ ) is equal to some all-variety (respectively 211 *ne*-variety) of stamps  $\mathbf{V}$ , we say that E all-*defines* (respectively ne-*defines*)  $\mathbf{V}$ . 212

▶ **Theorem 1.** A class of stamps is an all-variety (respectively ne-variety) of stamps if and only if it can be all-defined (respectively ne-defined) by a set of identities.

To give some examples, the classical varieties of monoids J, R and L can be characterised by identities in the following way:

<sup>217</sup> 
$$\langle \mathbf{R} \rangle_{all} = [\![(ab)^{\omega}a = (ab)^{\omega}]\!]_{all} = [\![(ab)^{\omega}a = (ab)^{\omega}]\!]_{ne}$$

 $\langle \mathbf{L} \rangle_{all} = \llbracket b(ab)^\omega = (ab)^\omega \rrbracket_{all} = \llbracket b(ab)^\omega = (ab)^\omega \rrbracket_{ne}$ 

218

$$\langle \mathbf{J} \rangle_{all} = \llbracket (ab)^{\omega} a = (ab)^{\omega}, \\ b(ab)^{\omega} = (ab)^{\omega} \rrbracket_{all} = \llbracket (ab)^{\omega} a = (ab)^{\omega}, \\ b(ab)^{\omega} = (ab)^{\omega} \rrbracket_{ne} .$$

Finite locally trivial semigroups and the join operation. The variety LI of finite locally trivial semigroups is well-known to verify  $\langle \mathbf{LI} \rangle_{ne} = [\![x^{\omega}yx^{\omega} = x^{\omega}]\!]_{ne}$  and to be such that for

any alphabet  $\Sigma$ , the set  $\mathcal{L}(\mathbf{LI})(\Sigma)$  contains exactly all Boolean combinations of languages of the form  $u\Sigma^*$  or  $\Sigma^*u$  for  $u \in \Sigma^*$ , or equivalently exactly all languages of the form  $U\Sigma^*V \cup W$ with  $U, V, W \subseteq \Sigma^*$  finite (see [13, p. 38]).

Given some variety of monoids  $\mathbf{V}$ , the join of  $\mathbf{V}$  and  $\mathbf{LI}$ , denoted by  $\mathbf{V} \vee \mathbf{LI}$ , is the inclusion-wise smallest variety of semigroups containing both  $\mathbf{V}$  and  $\mathbf{LI}$ . In fact, a finite semigroup S belongs to  $\mathbf{V} \vee \mathbf{LI}$  if and ony if there exist  $M \in \mathbf{V}$  and  $T \in \mathbf{LI}$  such that S divides the semigroup  $M \times T$ . (See [11, Chapter V, Exercise 1.1].) We can prove the following (see the appendix for the proof).

Proposition 2. Let V be a variety of monoids. Then  $\langle \mathbf{V} \vee \mathbf{LI} \rangle_{ne}$  is the inclusion-wise smallest ne-variety of stamps containing both  $\langle \mathbf{V} \rangle_{all}$  and  $\langle \mathbf{LI} \rangle_{ne}$ . Moreover,  $\mathcal{L}(\mathbf{V} \vee \mathbf{LI})$  is the inclusion-wise smallest ne-variety of languages containing both  $\mathcal{L}(\mathbf{V})$  and  $\mathcal{L}(\mathbf{LI})$  and verifies that  $\mathcal{L}(\mathbf{V} \vee \mathbf{LI})(\Sigma)$  is the Boolean closure of  $\mathcal{L}(\mathbf{V})(\Sigma) \cup \mathcal{L}(\mathbf{LI})(\Sigma)$  for each alphabet  $\Sigma$ .

## 235 **3** Essentially-V stamps

In this section, we give a characterisation of essentially-V stamps (defined in [12]), for V a
variety of monoids, in terms of identities. We first recall the definition.

▶ Definition 3. Let V be a variety of monoids. Let  $\varphi \colon \Sigma^* \to M$  be a stamp from an alphabet  $\Sigma$  to a finite monoid M. Let s be the stability index of  $\varphi$ .

We say that  $\varphi$  is essentially-**V** whenever there exists a stamp  $\mu: \Sigma^* \to N$  with  $N \in \mathbf{V}$ such that for all  $u, v \in \Sigma^*$ , we have

$$\mu(u) = \mu(v) \Rightarrow (\varphi(xuy) = \varphi(xvy) \quad \forall x, y \in \Sigma^s) .$$

<sup>243</sup> We will denote by **EV** the class of all essentially-V stamps.

Now, we give an equivalent condition for a stamp to be essentially-V, based on a specific congruence depending on that stamp.

<sup>246</sup> ► **Definition 4.** Let  $\varphi: \Sigma^* \to M$  for  $\Sigma$  an alphabet and M a finite monoid be a stamp and let <sup>247</sup> s be its stability index. We define the equivalence relation  $\equiv_{\varphi}$  on  $\Sigma^*$  by  $u \equiv_{\varphi} v$  for  $u, v \in \Sigma^*$ <sup>248</sup> whenever  $\varphi(xuy) = \varphi(xvy)$  for all  $x, y \in \Sigma^{\geq s}$ .

Proposition 5. Let  $\varphi \colon \Sigma^* \to M$  for  $\Sigma$  an alphabet and M a finite monoid be a stamp. Then  $\equiv_{\varphi}$  is a congruence of finite index and for any variety of monoids  $\mathbf{V}$ , we have  $\varphi \in \mathbf{EV}$ if and only if  $\Sigma^* / \equiv_{\varphi} \in \mathbf{V}$ .

<sup>252</sup> **Proof.** Let us denote by *s* the stability index of  $\varphi$ .

The equivalence relation  $\equiv_{\varphi}$  is a congruence because given  $u, v \in \Sigma^*$  verifying  $u \equiv_{\varphi} v$ , for all  $\alpha, \beta \in \Sigma^*$ , we have  $\alpha u\beta \equiv_{\varphi} \alpha v\beta$  since for any  $x, y \in \Sigma^{\geq s}$ , it holds that  $\varphi(x\alpha u\beta y) = \varphi(x\alpha v\beta y)$  because  $x\alpha, \beta y \in \Sigma^{\geq s}$ . Furthermore, this congruence is of finite index because for all  $u, v \in \Sigma^*$ , we have that  $\varphi(u) = \varphi(v)$  implies  $u \equiv_{\varphi} v$ .

Let now **V** be a variety of monoids. Assume first that  $\Sigma^* / \equiv_{\varphi} \in \mathbf{V}$ . It is quite direct to see that  $\varphi \in \mathbf{EV}$ , as the stamp  $\mu \colon \Sigma^* \to \Sigma^* / \equiv_{\varphi}$  defined by  $\mu(w) = [w]_{\equiv_{\varphi}}$  for all  $w \in \Sigma^*$ witnesses this fact. Assume then that  $\varphi \in \mathbf{EV}$ . This means that there exists a stamp  $\mu \colon \Sigma^* \to N$  with  $N \in \mathbf{V}$  such that for all  $u, v \in \Sigma^*$ , we have

$$\mu(u) = \mu(v) \Rightarrow (\varphi(xuy) = \varphi(xvy) \quad \forall x, y \in \Sigma^s)$$
.

Now consider  $u, v \in \Sigma^*$  such that  $\mu(u) = \mu(v)$ . For any  $x, y \in \Sigma^{\geq s}$ , we have that  $x = x_1 x_2$ with  $x_1 \in \Sigma^*$  and  $x_2 \in \Sigma^s$  as well as  $y = y_1 y_2$  with  $y_1 \in \Sigma^s$  and  $y_2 \in \Sigma^*$ , so that

 $\begin{aligned} \varphi(xuy) &= \varphi(x_1)\varphi(x_2uy_1)\varphi(y_2) = \varphi(x_1)\varphi(x_2vy_1)\varphi(y_2) = \varphi(xvy). \text{ Hence, } u \equiv_{\varphi} v. \text{ Therefore,} \\ \text{for all } u, v \in \Sigma^*, \text{ we have that } \mu(u) = \mu(v) \text{ implies } u \equiv_{\varphi} v, \text{ so we can define the mapping} \\ \alpha \colon N \to \Sigma^* / \equiv_{\varphi} \text{ such that } \alpha(\mu(w)) = [w]_{\equiv_{\varphi}} \text{ for all } w \in \Sigma^*. \text{ It is easy to check that } \alpha \\ \text{is actually a surjective morphism. Thus, we can conclude that } \Sigma^* / \equiv_{\varphi}, \text{ which divides } N, \\ \text{belongs to } \mathbf{V}. \end{aligned}$ 

Using this equivalent condition, we prove that given a set of identities *ne*-defining  $\langle \mathbf{V} \rangle_{all}$ for some variety of monoids  $\mathbf{V}$ , we get a set of identities *ne*-defining  $\mathbf{EV}$ , which actually shows that  $\mathbf{EV}$  is an *ne*-variety of stamps along the way.

Proposition 6. Let V be a variety of monoids and let E be a set of identities such that  $\langle V \rangle_{all} = \llbracket E \rrbracket_{ne}$ . Then

$$\mathbf{EV} = \llbracket x^{\omega}yuzt^{\omega} = x^{\omega}yvzt^{\omega} \mid u = v \in E, x, y, z, t \notin \mathrm{alph}(u) \cup \mathrm{alph}(v) \rrbracket_{\mathrm{ne}}$$

275 Proof. Let

$$F = \{x^{\omega}yuzt^{\omega} = x^{\omega}yvzt^{\omega} \mid u = v \in E, x, y, z, t \notin alph(u) \cup alph(v)\}.$$

<sup>277</sup> Central to the proof is the following claim.

<sup>278</sup>  $\triangleright$  Claim 7. Let  $\varphi \colon \Sigma^* \to M$  for  $\Sigma$  an alphabet and M a finite monoid be a stamp. Consider <sup>279</sup> the stamp  $\mu \colon \Sigma^* \to \Sigma^* / \equiv_{\varphi}$  defined by  $\mu(w) = [w]_{\equiv_{\varphi}}$  for all  $w \in \Sigma^*$ . It holds that for all <sup>280</sup>  $u, v \in \widehat{\Sigma^*}$ ,

$$\widehat{\mu}(u) = \widehat{\mu}(v) \Leftrightarrow \left(\widehat{\varphi}(\alpha^{\omega}\beta u\gamma\delta^{\omega}) = \widehat{\varphi}(\alpha^{\omega}\beta v\gamma\delta^{\omega}) \quad \forall \alpha, \beta, \gamma, \delta \in \Sigma^{+}\right) \,.$$

Using this claim, we can prove that  $\mathbf{EV} = \llbracket F \rrbracket_{ne}$ .

Inclusion from left to right. Let  $\varphi \colon \Sigma^* \to M$  for  $\Sigma$  an alphabet and M a finite monoid be a stamp in **EV**. Consider the stamp  $\mu \colon \Sigma^* \to \Sigma^* / \equiv_{\varphi}$  defined by  $\mu(w) = [w]_{\equiv_{\varphi}}$  for all  $w \in \Sigma^*$ . Since  $\varphi \in \mathbf{EV}$ , Proposition 5 tells us that  $\Sigma^* / \equiv_{\varphi} \in \mathbf{V}$ , hence  $\mu \in \langle \mathbf{V} \rangle_{all}$ .

Let us consider any identity  $x^{\omega}yuzt^{\omega} = x^{\omega}yvzt^{\omega} \in F$ . It is thus defined on an alphabet *B* with  $u = v \in E$  defined on an alphabet  $A \subseteq B$  and  $x, y, z, t \in B \setminus A$ . Let  $f: B^* \to \Sigma^*$  be an *ne*-morphism. Since  $\mu \in \langle \mathbf{V} \rangle_{all}$ , we have that  $\mu$  *ne*-satisfies the identity u = v, so that  $\hat{\mu}(\hat{f}(u)) = \hat{\mu}(\hat{f}(v))$ . Notice that we have that  $\hat{f}(x^{\omega}) = f(x)^{\omega}$  as well as  $\hat{f}(t^{\omega}) = f(t)^{\omega}$  and that  $f(x), f(y), f(z), f(t) \in \Sigma^+$  because f is non-erasing. Therefore, we have

$$\widehat{\varphi}\big(\widehat{f}(x^{\omega}yuzt^{\omega})\big) = \widehat{\varphi}\big(f(x)^{\omega}f(y)\widehat{f}(u)f(z)f(t)^{\omega}\big)$$

$$=\widehat{\varphi}\big(\widehat{f}(x^{\omega}yvzt^{\omega})\big)$$

<sup>295</sup> by Claim 7. As this holds for any *ne*-morphism  $f: B^* \to \Sigma^*$ , we can conclude that  $\varphi$ <sup>296</sup> *ne*-satisfies the identity  $x^{\omega}yuzt^{\omega} = x^{\omega}yvzt^{\omega}$ .

<sup>297</sup> This is true for any identity in F, so  $\varphi \in \llbracket F \rrbracket_{ne}$ . In conclusion,  $\mathbf{EV} \subseteq \llbracket F \rrbracket_{ne}$ .

 $=\widehat{\varphi}\big(f(x)^{\omega}f(y)\widehat{f}(v)f(z)f(t)^{\omega}\big)$ 

Inclusion from right to left. Let  $\varphi \colon \Sigma^* \to M$  for  $\Sigma$  an alphabet and M a finite monoid be a stamp in  $\llbracket F \rrbracket_{ne}$ . Consider the stamp  $\mu \colon \Sigma^* \to \Sigma^* / \equiv_{\varphi}$  defined by  $\mu(w) = [w]_{\equiv_{\varphi}}$  for all  $w \in \Sigma^*$ . We are now going to show that  $\mu \in \langle \mathbf{V} \rangle_{all}$ .

Take any identity  $u = v \in E$  defined on an alphabet A. There is thus an identity  $x^{\omega}yuzt^{\omega} = x^{\omega}yvzt^{\omega} \in F$  defined on an alphabet B such that  $A \subseteq B$  and  $x, y, z, t \in B \setminus A$ . Let  $f: A^* \to \Sigma^*$  be an *ne*-morphism.

Take any  $\alpha, \beta, \gamma, \delta \in \Sigma^+$ . Let us define the *ne*-morphism  $g: B^* \to \Sigma^*$  as the unique one such that for all  $b \in B$ ,

$$g(b) = \begin{cases} f(b) & \text{if } b \in A \\ \alpha & \text{if } b = x \\ \beta & \text{if } b = y \\ \gamma & \text{if } b = z \\ \delta & \text{if } b = t \end{cases}$$

Observe in particular that  $\widehat{g}(w) = \widehat{f}(w)$  for any  $w \in \widehat{A^*}$  and that  $\widehat{g}(x^{\omega}) = g(x)^{\omega} = \alpha^{\omega}$  as well as  $\widehat{g}(t^{\omega}) = \delta^{\omega}$ . Now, as  $\varphi$  ne-satisfies  $x^{\omega}yuzt^{\omega} = x^{\omega}yvzt^{\omega}$ , we have that

$$\widehat{\varphi}\left(\alpha^{\omega}\beta\widehat{f}(u)\gamma\delta^{\omega}\right) = \widehat{\varphi}\left(\widehat{g}(x^{\omega}yuzt^{\omega})\right) = \widehat{\varphi}\left(\widehat{g}(x^{\omega}yvzt^{\omega})\right) = \widehat{\varphi}\left(\alpha^{\omega}\beta\widehat{f}(v)\gamma\delta^{\omega}\right)$$

Since this holds for any  $\alpha, \beta, \gamma, \delta \in \Sigma^+$ , by Claim 7, we have that  $\widehat{\mu}(\widehat{f}(u)) = \widehat{\mu}(\widehat{f}(v))$ .

Therefore,  $\hat{\mu}(\hat{f}(u)) = \hat{\mu}(\hat{f}(v))$  for any *ne*-morphism  $f: A^* \to \Sigma^*$ , which means that  $\mu$ *ne*-satisfies u = v.

So, to conclude, since this is true for any  $u = v \in E$ , we have that  $\mu \in \langle \mathbf{V} \rangle_{all}$ , which implies that  $\Sigma^* = \varphi \in \mathbf{V}$  and thus  $\varphi \in \mathbf{EV}$  by Proposition 5. In conclusion,  $[\![F]\!]_{ne} \subseteq \mathbf{EV}$ .

<sup>315</sup> The claim still needs to be proved.

Proof of Claim 7. Let  $\varphi \colon \Sigma^* \to M$  for  $\Sigma$  an alphabet and M a finite monoid be a stamp of stability index s. Consider the stamp  $\mu \colon \Sigma^* \to \Sigma^* / \equiv_{\varphi}$  defined by  $\mu(w) = [w]_{\equiv_{\varphi}}$  for all  $w \in \Sigma^*$ . We now want to show that for all  $u, v \in \widehat{\Sigma^*}$ ,

$$\widehat{\mu}(u) = \widehat{\mu}(v) \Leftrightarrow \left(\widehat{\varphi}(\alpha^{\omega}\beta u\gamma\delta^{\omega}) = \widehat{\varphi}(\alpha^{\omega}\beta v\gamma\delta^{\omega}) \quad \forall \alpha, \beta, \gamma, \delta \in \Sigma^+\right) \,.$$

Let  $u, v \in \widehat{\Sigma^*}$ . There exist two Cauchy sequences  $(u_n)_{n\geq 0}$  and  $(v_n)_{n\geq 0}$  in  $\Sigma^*$  such that  $u = \lim_{n\to\infty} u_n$  and  $v = \lim_{n\to\infty} v_n$ . As  $\Sigma^*/\equiv_{\varphi}$  and M are discrete, we have that all four Cauchy sequences  $(\mu(u_n))_{n\geq 0}$ ,  $(\varphi(u_n))_{n\geq 0}$ ,  $(\mu(v_n))_{n\geq 0}$  and  $(\varphi(v_n))_{n\geq 0}$  are ultimately constant. So there exists  $k \in \mathbb{N}$  such that  $\widehat{\mu}(u) = \mu(u_k)$ ,  $\widehat{\varphi}(u) = \varphi(u_k)$ ,  $\widehat{\mu}(v) = \mu(v_k)$  and  $\widehat{\varphi}(v) = \varphi(v_k)$ .

Assume first that  $\hat{\mu}(u) = \hat{\mu}(v)$ . Take any  $\alpha, \beta, \gamma, \delta \in \Sigma^+$ . Since M is discrete, both Cauchy sequences  $(\varphi(\alpha^{n!}))_{n\geq 0}$  and  $(\varphi(\delta^{n!}))_{n\geq 0}$  are ultimately constant. So there exists  $l \in \mathbb{N}$  such that for all  $m \in \mathbb{N}, m \geq l$ , we have  $\hat{\varphi}(\alpha^{\omega}) = \varphi(\alpha^{m!})$  and  $\hat{\varphi}(\delta^{\omega}) = \varphi(\delta^{m!})$ . Hence, taking some  $m \in \mathbb{N}, m \geq l$  such that  $|\alpha^{m!}\beta| \geq s$  and  $|\gamma\delta^{m!}| \geq s$ , it follows that

$$\widehat{\varphi}(\alpha^{\omega}\beta u\gamma\delta^{\omega}) = \varphi(\alpha^{m!}\beta u_k\gamma\delta^{m!}) = \varphi(\alpha^{m!}\beta v_k\gamma\delta^{m!}) = \widehat{\varphi}(\alpha^{\omega}\beta v\gamma\delta^{\omega})$$

because  $[u_k]_{\equiv_{\varphi}} = \widehat{\mu}(u) = \widehat{\mu}(v) = [v_k]_{\equiv_{\varphi}}$ . Thus, we have that

$$\widehat{\varphi}(\alpha^{\omega}\beta u\gamma\delta^{\omega}) = \widehat{\varphi}(\alpha^{\omega}\beta v\gamma\delta^{\omega})$$

for all  $\alpha, \beta, \gamma, \delta \in \Sigma^+$ .

Assume then that  $\widehat{\varphi}(\alpha^{\omega}\beta u\gamma\delta^{\omega}) = \widehat{\varphi}(\alpha^{\omega}\beta v\gamma\delta^{\omega})$  for all  $\alpha, \beta, \gamma, \delta \in \Sigma^+$ . Take any  $\alpha, \beta \in \Sigma^{\geq s}$ . Since  $\varphi(\Sigma^s)$  is a finite semigroup and verifies that  $\varphi(\Sigma^s) = \varphi(\Sigma^s)^2$ , by a classical result in finite semigroup theory (see e.g. [13, Chapter 1, Proposition 1.12]), we have that there exist  $\alpha_1, e, f, \beta_2 \in \Sigma^s$  and  $\alpha_2, \beta_1 \in \Sigma^{\geq s}$  such that  $\varphi(\alpha_1 e \alpha_2) = \varphi(\alpha)$  and  $\varphi(\beta_1 f \beta_2) = \varphi(\beta)$ with  $\varphi(e)$  and  $\varphi(f)$  idempotents. Now, since  $\varphi(e)$  is idempotent, we have that

$$\widehat{\varphi}(e^{\omega}) = \widehat{\varphi}(\lim_{n \to \infty} e^{n!}) = \lim_{n \to \infty} \varphi(e^{n!}) = \lim_{n \to \infty} \varphi(e)^{n!} = \varphi(e)$$

and similarly,  $\widehat{\varphi}(f^{\omega}) = \varphi(f)$ . So it follows that

- 340  $\varphi(\alpha u_k\beta) = \varphi(\alpha_1 e \alpha_2 u_k\beta_1 f \beta_2)$
- $=\widehat{\varphi}(\alpha_1 e^{\omega} \alpha_2 u \beta_1 f^{\omega} \beta_2)$
- $=\widehat{\varphi}(\alpha_1 e^{\omega} \alpha_2 v \beta_1 f^{\omega} \beta_2)$

 $=\varphi(\alpha v_k\beta)$ .

- $_{\mathbf{343}} \qquad \qquad = \varphi(\alpha_1 e \alpha_2 v_k \beta_1 f \beta_2)$
- 344 345

As this is true for any  $\alpha, \beta \in \Sigma^{\geq s}$ , by definition it holds that  $u_k \equiv_{\varphi} v_k$ , hence  $\widehat{\mu}(u) = \mu(u_k) = \mu(v_k) = \mu(v_k) = \widehat{\mu}(v)$ .

<sup>348</sup> This concludes the proof of the proposition.

## $_{349}$ **4** Essentially-V stamps and the join of V and LI

In this section, we establish the link between essentially-V stamps and  $V \vee LI$  and give a criterion that characterises exactly when they do correspond.

<sup>352</sup> More precisely, consider the following criterion for a variety of monoids **V**.

Solution (A). For any  $L \in \mathcal{L}(\mathbf{V})(\Sigma)$  with  $\Sigma$  some alphabet, we have  $xLy \in \mathcal{L}(\mathbf{V} \vee \mathbf{LI})(\Sigma)$ for all  $x, y \in \Sigma^*$ .

It is a kind of mild closure condition that appears to be a sufficient and necessary condition for  $\mathbf{EV}$  and  $\mathbf{V} \lor \mathbf{LI}$  to correspond.

<sup>357</sup> ► Proposition 8. Let V be a variety of monoids. Then  $\langle V \lor LI \rangle_{ne} \subseteq EV$  and there is <sup>358</sup> equality if and only if V verifies criterion (A).

Why this proposition is actually useful to give characterisations of  $\mathbf{V} \lor \mathbf{LI}$  in terms of identities will become clear in the next section. For now, we focus on its proof, that entirely relies on the following characterisation of the languages recognised by essentially- $\mathbf{V}$  stamps.

<sup>362</sup> ▶ Proposition 9. Let V be a variety of monoids. For any alphabet Σ, the set  $\mathcal{L}(\mathbf{EV})(\Sigma)$ <sup>363</sup> contains exactly all Boolean combinations of languages of the form xLy for  $L \in \mathcal{L}(\mathbf{V})(\Sigma)$ <sup>364</sup> and  $x, y \in \Sigma^*$ .

Proof. Let C be the class of languages such that for any alphabet  $\Sigma$ , the set  $C(\Sigma)$  contains exactly all Boolean combinations of languages of the form xLy for  $L \in \mathcal{L}(\mathbf{V})(\Sigma)$  and  $x, y \in \Sigma^*$ . Let  $\Sigma$  be an alphabet. We are now going to show that  $\mathcal{L}(\mathbf{EV})(\Sigma) = C(\Sigma)$ , which proves

the proposition as it holds for any alphabet  $\Sigma$ .

Inclusion from right to left. Let  $L \in \mathcal{L}(\mathbf{V})(\Sigma)$  and  $x, y \in \Sigma^*$ . Let  $\mu \colon \Sigma^* \to N$  be the syntactic morphism of L: this means that  $N \in \mathbf{V}$  and that there exists  $F \subseteq N$  such that  $L = \mu^{-1}(F)$ . Let also  $\varphi \colon \Sigma^* \to M$  be the syntactic morphism of the language  $xLy = x\Sigma^*y \cap \Sigma^{|x|}\mu^{-1}(F)\Sigma^{|y|}$  and let s be its stability index. We then consider some  $u, v \in \Sigma^*$  such that  $\mu(u) = \mu(v)$ . Take any  $x', y' \in \Sigma^*$  such that  $|x'| \ge |x|$  and  $|y'| \ge |y|$ . We clearly have that  $x'uy' \in x\Sigma^*y$  if and only if  $x'vy' \in x\Sigma^*y$ . Moreover,  $x' = x'_1x'_2$  for some  $x'_1 \in \Sigma^{|x|}$  and  $x'_2 \in \Sigma^*$  and  $y' = y'_1y'_2$  for some  $y'_1 \in \Sigma^*$  and  $y'_2 \in \Sigma^{|y|}$ , so that

F

376 
$$x'uy' \in \Sigma^{|x|} \mu^{-1}(F)\Sigma^{|y|} \Leftrightarrow \mu(x'_2 uy'_1) \in F$$

377 
$$\Leftrightarrow \mu(x'_2 v y'_1) \in$$

$$\Leftrightarrow x'vy' \in \Sigma^{|x|}\mu^{-1}(F)\Sigma^{|y|} .$$

Hence,  $x'uy' \in xLy$  if and only if  $x'vy' \in xLy$  for all  $x', y' \in \Sigma^*$  such that  $|x'| \ge |x|$  and  $|y'| \ge |y|$ , so that, by definition of the stability index s of  $\varphi$  and as  $\varphi$  is the syntactic morphism of xLy, we have  $\varphi(x'uy') = \varphi(x'vy')$  for all  $x', y' \in \Sigma^s$ . Thus, it follows that  $\varphi \in \mathbf{EV}$ .

This implies that  $xLy \in \mathcal{L}(\mathbf{EV})(\Sigma)$ . Therefore, since this is true for any  $L \in \mathcal{L}(\mathbf{V})(\Sigma)$ and  $x, y \in \Sigma^*$  and since  $\mathcal{L}(\mathbf{EV})(\Sigma)$  is closed under Boolean operations, we can conclude that  $\mathcal{L}(\Sigma) \subseteq \mathcal{L}(\mathbf{EV})(\Sigma)$ .

Inclusion from left to right. Let  $L \in \mathcal{L}(\mathbf{EV})(\Sigma)$  and let  $\varphi \colon \Sigma^* \to M$  be its syntactic morphism: it is an essentially-V stamp. Given s its stability index, this means there exists a stamp  $\mu \colon \Sigma^* \to N$  with  $N \in \mathbf{V}$  such that for all  $u, v \in \Sigma^*$ , we have

389 
$$\mu(u) = \mu(v) \Rightarrow \left( \varphi(xuy) = \varphi(xvy) \quad \forall x, y \in \Sigma^s \right)$$
.

For each  $m \in N$  and  $x, y \in \Sigma^s$  consider the language  $x\mu^{-1}(m)y$ : it verifies that for any two words  $w, w' \in x\mu^{-1}(m)y$ , we have w = xuy and w' = xvy with  $\mu(u) = \mu(v) = m$ , so that  $\varphi(w) = \varphi(w')$ . By definition of the syntactic morphism, this means that for all  $m \in N$ and  $x, y \in \Sigma^s$ , either  $x\mu^{-1}(m)y \subseteq L$  or  $x\mu^{-1}(m)y \cap L = \emptyset$ . Therefore, there exists a set  $E \subseteq N \times \Sigma^s \times \Sigma^s$  such that  $L \cap \Sigma^{\geq 2s} = \bigcup_{(m,x,y) \in E} x\mu^{-1}y$ , hence

$$L = \bigcup_{(m,x,y)\in E} x\mu^{-1}(m)y \cup F$$

<sup>396</sup> for some  $F \subseteq \Sigma^{<2s}$ .

Take some  $w \in F$ . We have that  $\{w\} = w\Sigma^* \cap \bigcap_{a \in \Sigma} (wa\Sigma^*)^{\complement}$  with necessarily  $\Sigma^* \in \mathcal{L}(\mathbf{V})(\Sigma)$ . Thus, the singleton language  $\{w\}$  belongs to  $\mathcal{C}(\Sigma)$  and since this is true for any  $w \in F$  and F is finite, we can deduce from this that F is in  $\mathcal{C}(\Sigma)$ , as the latter is trivially closed under Boolean operations.

Now, for all  $m \in N$ , the language  $\mu^{-1}(m)$  belongs to  $\mathcal{L}(\mathbf{V})(\Sigma)$ , so we finally have  $L \in \mathcal{C}(\Sigma)$ . This is true for any  $L \in \mathcal{L}(\mathbf{EV})(\Sigma)$ , so in conclusion,  $\mathcal{L}(\mathbf{EV})(\Sigma) \subseteq \mathcal{C}(\Sigma)$ .

Proposition 8 then follows from the two next lemmata, that are both easy consequences
 of Proposition 9. For completeness, we give the proofs in the appendix.

<sup>405</sup> ► Lemma 10. Let V be a variety of monoids. Then  $\langle V \lor LI \rangle_{ne} \subseteq EV$ .

▶ Lemma 11. Let V be a variety of monoids. Then  $\mathbf{EV} \subseteq \langle \mathbf{V} \lor \mathbf{LI} \rangle_{ne}$  if and only if V verifies criterion (A).

### 408 **5** Applications

<sup>409</sup> In this last section, we use the link between essentially-V stamps and  $V \vee LI$  to reprove some <sup>410</sup> characterisations of joins between LI and some well-known varieties of monoids in terms of <sup>411</sup> identities.

One thing that seems at a first glance a bit problematic about proving that some variety 412 of monoids V satisfies criterion (A) when one's goal is precisely to characterise  $\mathbf{V} \vee \mathbf{LI}$  is 413 that one needs to prove that some languages belong to  $\mathcal{L}(\mathbf{V} \vee \mathbf{LI})$ , a class about which one 414 a priori only knows what is given by Proposition 2. But understanding  $\mathcal{L}(\mathbf{V})$  enough is 415 actually sufficient: if it holds that given any language  $L \in \mathcal{L}(\mathbf{V})(\Sigma)$  for some alphabet  $\Sigma$  and 416 any  $x, y \in \Sigma^*$ , there exists some language  $K \in \mathcal{L}(\mathbf{V})(\Sigma)$  such that L is equal to the quotient 417  $x^{-1}Ky^{-1}$ , then V verifies criterion (A). We don't know whether this quotient-expressibility 418 condition that solely depends on the variety  $\mathbf{V}$  (without explicit reference to  $\mathbf{LI}$ ) is actually 419

equivalent to it satisfying criterion (A), but we can prove such an equivalence for a weaker quotient-expressibility condition for **V**. The proof is to be found in the appendix.

▶ Proposition 12. Let V be a variety of monoids. Then V satifies criterion (A) if and and only if for any  $L \in \mathcal{L}(\mathbf{V})(\Sigma)$  for some alphabet  $\Sigma$  and  $x, y \in \Sigma^*$ , there exist  $k, l \in \mathbb{N}$ such that for all  $u \in \Sigma^k, v \in \Sigma^l$ , there exists some language  $K \in \mathcal{L}(\mathbf{V})(\Sigma)$  verifying  $u^{-1}Lv^{-1} = (xu)^{-1}K(vy)^{-1}$ .

This quotient-expressibility condition appears to be particularly useful to prove that a variety of monoids V does not satisfy criterion (A) without needing to understand what  $\mathcal{L}(\mathbf{V} \vee \mathbf{LI})$  is. We demonstrate this for the variety of finite commutative and idempotent monoids  $\mathbf{J}_1$ .

**430 • Proposition 13.**  $J_1$  does not satisfy criterion (A).

<sup>431</sup> **Proof.** Given an alphabet  $\Sigma$ , the set  $\mathcal{L}(\mathbf{J}_1)(\Sigma)$  contains exactly all Boolean combinations of <sup>432</sup> languages of the form  $\Sigma^* a \Sigma^*$  for  $a \in \Sigma$  (see [13, Chapter 2, Proposition 3.10]).

Let  $L = \{a, b\}^* b\{a, b\}^* \in \mathcal{L}(\mathbf{J_1})(\{a, b\})$  and  $x = b, y = \varepsilon$ . Take any  $k, l \in \mathbb{N}$  and set  $u = a^k$  and  $v = a^l$ . Consider some  $K \in \mathcal{L}(\mathbf{J_1})(\{a, b\})$ . We have that  $xuavy \in K \Leftrightarrow$   $xuabvy \in K$  so that  $a \in (xu)^{-1}K(vy)^{-1} \Leftrightarrow ab \in (xu)^{-1}K(vy)^{-1}$ . But  $a \notin u^{-1}Lv^{-1}$ and  $ab \in u^{-1}Lv^{-1}$ , hence  $u^{-1}Lv^{-1} \neq (xu)^{-1}K(vy)^{-1}$  and this holds for any choice of K. So for any  $k, l \in \mathbb{N}$ , there exists  $u \in \Sigma^k, v \in \Sigma^l$  such that no  $K \in \mathcal{L}(\mathbf{J_1})(\{a, b\})$  verifies  $u^{-1}Lv^{-1} = (xu)^{-1}K(vy)^{-1}$ .

439 In conclusion, by Proposition 12,  $J_1$  does not satisfy criterion (A).

◀

We now prove the announced characterisations of joins between **LI** and some well-known varieties of monoids in terms of identities.

<sup>442</sup> ► **Theorem 14.** We have the following.

443 1.  $\langle \mathbf{R} \vee \mathbf{L} \mathbf{I} \rangle_{\mathrm{ne}} = \mathbf{E} \mathbf{R} = \llbracket x^{\omega} y(ab)^{\omega} azt^{\omega} = x^{\omega} y(ab)^{\omega} zt^{\omega} \rrbracket_{\mathrm{ne}}.$ 

444 2.  $\langle \mathbf{L} \vee \mathbf{L} \mathbf{I} \rangle_{\mathrm{ne}} = \mathbf{E} \mathbf{L} = \llbracket x^{\omega} y b(ab)^{\omega} z t^{\omega} = x^{\omega} y(ab)^{\omega} z t^{\omega} \rrbracket_{\mathrm{ne}}.$ 

445 **3.**  $\langle \mathbf{J} \vee \mathbf{L} \mathbf{I} \rangle_{ne} = \mathbf{E} \mathbf{J} = \llbracket x^{\omega} y(ab)^{\omega} azt^{\omega} = x^{\omega} y(ab)^{\omega} zt^{\omega}, x^{\omega} yb(ab)^{\omega} zt^{\omega} = x^{\omega} y(ab)^{\omega} zt^{\omega} \rrbracket_{ne}.$ 

446 4.  $\langle \mathbf{H} \lor \mathbf{L} \mathbf{I} \rangle_{ne} = \mathbf{E} \mathbf{H}$  for any variety of groups  $\mathbf{H}$ .

447 Proof. In each case, we prove that the variety of monoids under consideration satisfies
 448 criterion (A) using Proposition 12. We then use Proposition 8.

**Proof of 1.** It is well-known that given an alphabet  $\Sigma$ , the set  $\mathcal{L}(\mathbf{R})(\Sigma)$  contains all languages that are disjoint unions of languages that are of the form  $A_0^*a_1A_1^*\cdots a_kA_k^*$  where  $k \in \mathbb{N}, a_1, \ldots, a_k \in \Sigma, A_0, A_1, \ldots, A_k \subseteq \Sigma$  and  $a_i \notin A_{i-1}$  for all  $i \in [k]$  (see [13, Chapter 4, Theorem 3.3]).

Let  $\Sigma$  be an alphabet and take some language  $A_0^* a_1 A_1^* \cdots a_k A_k^*$  where  $k \in \mathbb{N}, a_1, \dots, a_k \in \Sigma$ ,  $A_0, A_1, \dots, A_k \subseteq \Sigma$  and  $a_i \notin A_{i-1}$  for all  $i \in [k]$ . Take  $x, y \in \Sigma^*$ . Observe that y can be uniquely written as y = zt where  $z \in A_k^*$  and  $t \in \{\varepsilon\} \cup (\Sigma \setminus A_k)\Sigma^*$ . We have

$$_{456} \qquad A_0^* a_1 A_1^* \cdots a_k A_k^* = x^{-1} \Big( x A_0^* a_1 A_1^* \cdots a_k A_k^* t \cap \bigcap_{v \in A_k^{<|z|}} (x A_0^* a_1 A_1^* \cdots a_k v t)^{\complement} \Big) y^{-1}$$

using the convention that  $xA_0^*a_1A_1^*\cdots a_kvt = xvt$  for all  $v \in A_k^{<|z|}$  when k = 0. The language  $xA_0^*a_1A_1^*\cdots a_kA_k^*t \cap \bigcap_{v \in A_k^{<|z|}} (xA_0^*a_1A_1^*\cdots a_kvt)^{\complement}$  does belong to the set  $\mathcal{L}(\mathbf{R})(\Sigma)$  because the latter is closed under Boolean operations and by definition of z and t. Thus, we can conclude that for each  $L \in \mathcal{L}(\mathbf{R})(\Sigma)$  and  $x, y \in \Sigma^*$ , there exists  $K \in \mathcal{L}(\mathbf{R})(\Sigma)$  such that  $L = x^{-1}Ky^{-1}$  by using the characterisation of  $\mathcal{L}(\mathbf{R})(\Sigma)$ , the fact that quotients commute with unions [13, p. 20] and closure of  $\mathcal{L}(\mathbf{R})(\Sigma)$  under unions.

**Proof of 2.** It is also well-known that given an alphabet  $\Sigma$ , the set  $\mathcal{L}(\mathbf{L})(\Sigma)$  contains all languages that are disjoint unions of languages that are of the form  $A_0^*a_1A_1^*\cdots a_kA_k^*$  where  $k \in \mathbb{N}, a_1, \ldots, a_k \in \Sigma, A_0, A_1, \ldots, A_k \subseteq \Sigma$  and  $a_i \notin A_i$  for all  $i \in [k]$  (see [13, Chapter 4, Theorem 3.4]). The proof is then dual to the previous case.

**Proof of 3.** Given an alphabet  $\Sigma$ , for each  $k \in \mathbb{N}$ , we define the equivalence relation  $\sim_k$ 467 on  $\Sigma^*$  by  $u \sim_k v$  for  $u, v \in \Sigma^*$  whenever u and v have the same set of subwords of length 468 at most k. This relation is a congruence of finite index on  $\Sigma^*$ . Simon proved [17] that a 469 language belongs to  $\mathcal{L}(\mathbf{J})(\Sigma)$  if and only it is equal to a union of  $\sim_k$ -classes for some  $k \in \mathbb{N}$ . 470 Let  $\Sigma$  be an alphabet and take some  $L \in \mathcal{L}(\mathbf{J})(\Sigma)$  as well as  $x, y \in \Sigma^*$ . Thus, there exists 471  $k \in \mathbb{N}$  such that L is a union of  $\sim_k$ -classes. Define the language  $K = \bigcup_{w \in L} [xwy]_{\sim_{|xy|+k}}$ : it 472 belongs to  $\mathcal{L}(\mathbf{J})(\Sigma)$  by construction. We now show that  $L = x^{-1}Ky^{-1}$ , which concludes 473 the proof. Let  $w \in L$ : we have that  $xwy \in [xwy]_{\sim |xy|+k} \subseteq K$ , so that  $w \in x^{-1}Ky^{-1}$ . Let conversely  $w \in x^{-1}Ky^{-1}$ . This means that  $xwy \in K$ , which implies that there exists  $w' \in L$ 474 475 such that  $xwy \sim_{|xy|+k} xw'y$ . Actually, it holds that any  $u \in \Sigma^*$  of length at most k is a 476 subword of w if and only if it is a subword of w', because xuy is a subword of xwy if and 477 only if it is a subword of xw'y. Hence,  $w \sim_k w'$ , which implies that  $w \in L$ . 478

**Proof of 4.** Consider some variety of groups **H**. Take some language  $L \in \mathcal{L}(\mathbf{H})(\Sigma)$  for an alphabet  $\Sigma$  and let  $x, y \in \Sigma^*$ . Consider the syntactic morphism  $\eta \colon \Sigma^* \to M$  of L: we have that M is a group in **H**. Define the language  $K = \eta^{-1}(\eta(x)\eta(L)\eta(y))$ : it belongs to  $\mathcal{L}(\mathbf{H})(\Sigma)$ . We now show that  $L = x^{-1}Ky^{-1}$ , which concludes the proof. Let  $w \in L$ : we have that  $\eta(xwy) \in \eta(x)\eta(L)\eta(y)$ , so that  $w \in x^{-1}Ky^{-1}$ . Conversely, let  $w \in x^{-1}Ky^{-1}$ . We have that  $xwy \in K$ , which means that  $\eta(xwy) = \eta(x)\eta(w')\eta(y)$  for some  $w' \in L$ , so that  $\eta(w) = \eta(w') \in \eta(L)$ , as any element in **H** is invertible. Thus,  $w \in L$ .

## 486 6 Conclusion

The general method presented in this paper actually allows to reprove in a straightforward language-theoretic way even more characterisations of the join of **LI** with some variety of finite monoids. This can for instance be done for the variety of finite commutative monoids **Com** or the variety of finite commutative aperiodic monoids **ACom**.

In fact, as already observed in some sense by Costa [8], many varieties of finite monoids seem to verify criterion (A). The main question left open by this present work is to understand better what exactly those varieties are. Another question left open is whether Proposition 12 can be refined by using the stronger quotient-expressibility condition alluded to before the statement of the proposition. The answers to both questions are unclear to the author, but making progress on them may also lead to a better understanding of joins of varieties of finite monoids with **LI**.

<sup>498</sup> **Acknowledgements.** The author wants to thank Thomas Place, who suggested the link <sup>499</sup> between essentially-V stamps and  $V \vee LI$ , but also Luc Segoufin who started the discussion <sup>500</sup> with Thomas Place and encouraged the author to write the present article.

501		References
502	1	Jorge Almeida. Some pseudovariety joins involving the pseudovariety of finite groups. In
503		Semigroup Forum, volume 37, pages 53–57. Springer, 1988.
504	2	Jorge Almeida. Finite semigroups and universal algebra, volume 3. World Scientific, 1994.
505	3	Jorge Almeida and Assis Azevedo. The join of the pseudovarieties of R-trivial and L-trivial
506		monoids. Journal of Pure and Applied Algebra, 60(2):129–137, 1989.
507	4	Jorge Almeida and Pascal Weil. Free profinite mathscrR-trivial monoids. Int. J. Algebra
508		Comput., 7(5):625-672, 1997. doi:10.1142/S0218196797000289.
509	5	Assis Azevedo. The Join of the Pseudovariety J with Permutative Pseudovarieties, pages 1–11.
510		Springer US, Boston, MA, 1990. doi:10.1007/978-1-4899-2608-1_1.
511	6	Assis Azevedo and Marc Zeitoun. Three examples of join computations. In Semigroup Forum,
512		volume 57, pages 249–277. Springer, 1998.
513	7	Laura Chaubard, Jean-Eric Pin, and Howard Straubing. First order formulas with modular
514		predicates. In 21th IEEE Symposium on Logic in Computer Science (LICS 2006), 12-15
515		August 2006, Seattle, WA, USA, Proceedings, pages 211–220. IEEE Computer Society, 2006.
516	0	doi:10.1109/LICS.2006.24.
517	8	José Carlos Costa. Some pseudovariety joins involving locally trivial semigroups. In <i>Semigroup</i>
518	0	Forum, volume 64, pages 12–28. Springer, 2001.
519	9	Jose Carlos Costa. Some pseudovariety joins involving groups and locally trivial semigroups.
520	10	In Semigroups, Algorithms, Automata Ana Languages, pages 341–348. World Scientific, 2002.
521	10	Samuel Enemberg. Automata, Languages, and Machines, volume A. Academic Press, New Vorle 1074
522	11	Samuel Filenborg Automata Languages and Machines volume B. Academic Press New
523	11	Vork 1976
524	12	Nathan Grosshans Pierre McKenzie and Luc Segoufin Tameness and the power of programs
526		over monoids in DA. CoBR. abs/2101.07495, 2021. UBL: https://arxiv.org/abs/2101.
527		07495. arXiv:2101.07495.
528	13	Jean-Éric Pin. Varieties of formal languages. North Oxford, London and Plenum, New-York,
529		1986. (Traduction de Variétés de langages formels).
530	14	Jean-Éric Pin and Howard Straubing. Some results on C-varieties. RAIRO-Theor. Inf. Appl.,
531		39(1):239–262, 2005.
532	15	Jan Reiterman. The Birkhoff theorem for finite algebras. <i>algebra universalis</i> , 14(1):1–10, 1982.
533	16	Marcel Paul Schützenberger. On finite monoids having only trivial subgroups. Inf. Control.,
534		8(2):190-194, 1965. doi:10.1016/S0019-9958(65)90108-7.
535	17	Imre Simon. Piecewise testable events. In Automata Theory and Formal Languages, 2nd GI
536		Conference, volume 33 of Lecture Notes in Computer Science, pages 214–222. Springer, 1975.
537	18	Howard Straubing. On logical descriptions of regular languages. In Sergio Rajsbaum, editor,
538		LATIN 2002: Theoretical Informatics, 5th Latin American Symposium, Cancun, Mexico, April
539		3-6, 2002, Proceedings, volume 2286 of Lecture Notes in Computer Science, pages 528–538.
540		Springer, 2002. doi:10.1007/3-540-45995-2\_46.
541	19	Marc Zeitoun. The join of the pseudovarieties of idempotent semigroups and locally trivial
542		semigroups. In Semigroup Forum, volume 50, pages 367–381. Springer, 1995.
543	20	Marc Zeitoun. On the decidability of the membership problem of the pseudovariety $J \vee B$ . Int.
544	01	J. Algebra Comput., 5(1):47-64, 1995. doi:10.1142/S0218196795000057.
545	21	Marc Zeitoun. On the join of two pseudovarieties. Semigroups, Automata and Languages, eds.
546		J. Atmeiaa, GMS Gomes, and PV Silva. World Scientific, pages 281–288, 1996.

## 547 **A** Missing proofs

Front of Proposition 2. Let W be some *ne*-variety of stamps such that  $\langle \mathbf{V} \rangle_{all} \cup \langle \mathbf{LI} \rangle_{ne} \subseteq \mathbf{W}$ . There exists some variety of semigroups  $\mathbf{W}'$  such that  $\langle \mathbf{W}' \rangle_{ne} = \mathbf{W}$ .

Let  $S \in \mathbf{V} \cup \mathbf{LI}$ . We denote by  $S^1$  the monoid S if S is already a monoid and the monoid  $S \cup \{1\}$  otherwise. Then the evaluation morphism  $\eta_S \colon S^* \to S^1$  such that  $\eta_S(s) = s$  for all  $s \in S$  verifies  $\eta_S(S^+) = S$  and additionally  $S^1 = S$  when  $S \in \mathbf{V}$ . This implies that  $\eta_S \in \langle \mathbf{V} \rangle_{all} \cup \langle \mathbf{LI} \rangle_{ne} \subseteq \mathbf{W}$ . But by definition of  $\mathbf{W}'$ , it must be that  $S = \eta_S(S^+) \in \mathbf{W}'$ .

Therefore,  $\mathbf{W}'$  contains both  $\mathbf{V}$  and  $\mathbf{LI}$ , which implies that  $\mathbf{V} \lor \mathbf{LI} \subseteq \mathbf{W}'$  by inclusion-wise minimality of  $\mathbf{V} \lor \mathbf{LI}$ . By definition, we can then conclude that  $\langle \mathbf{V} \lor \mathbf{LI} \rangle_{ne} \subseteq \langle \mathbf{W}' \rangle_{ne} = \mathbf{W}$ . So  $\langle \mathbf{V} \lor \mathbf{LI} \rangle_{ne}$  is the inclusion-wise smallest *ne*-variety of stamps containing both  $\langle \mathbf{V} \rangle_{all}$  and  $\langle \mathbf{LI} \rangle_{ne}$ .

Let now  $\mathcal{W}$  be some *ne*-variety of languages such that  $\mathcal{L}(\mathbf{V}) \cup \mathcal{L}(\mathbf{LI}) \subseteq \mathcal{W}$ . It holds that 558  $\mathcal{W} = \mathcal{L}(\mathbf{W})$  for some *ne*-variety of stamps **W**. We have that  $\langle \mathbf{V} \rangle_{all}$ , which is in particular an 559 *ne*-variety of stamps, is included in **W** because  $\mathcal{L}(\langle \mathbf{V} \rangle_{all}) = \mathcal{L}(\mathbf{V}) \subseteq \mathcal{W} = \mathcal{L}(\mathbf{W})$ , but also 560 that  $\langle \mathbf{L}\mathbf{I} \rangle_{ne}$  is included in W because  $\mathcal{L}(\langle \mathbf{L}\mathbf{I} \rangle_{ne}) = \mathcal{L}(\mathbf{L}\mathbf{I}) \subseteq \mathcal{W} = \mathcal{L}(\mathbf{W})$ . By inclusion-wise 561 minimality of  $\langle \mathbf{V} \vee \mathbf{L} \mathbf{I} \rangle_{ne}$ , it follows that  $\langle \mathbf{V} \vee \mathbf{L} \mathbf{I} \rangle_{ne} \subseteq \mathbf{W}$ . Hence, using again the above 562 fact on the Eilenberg correspondence, we can conclude that  $\mathcal{L}(\mathbf{V} \vee \mathbf{LI}) = \mathcal{L}(\langle \mathbf{V} \vee \mathbf{LI} \rangle_{ne}) \subseteq$ 563  $\mathcal{L}(\mathbf{W}) = \mathcal{W}$ . So  $\mathcal{L}(\mathbf{V} \vee \mathbf{LI})$  is the inclusion-wise smallest *ne*-variety of languages containing 564 both  $\mathcal{L}(\mathbf{V})$  and  $\mathcal{L}(\mathbf{LI})$ . 565

<sup>566</sup> Consider now the class of languages C such that  $C(\Sigma)$  is the Boolean closure of  $\mathcal{L}(\mathbf{V})(\Sigma) \cup$ <sup>567</sup>  $\mathcal{L}(\mathbf{LI})(\Sigma)$  for each alphabet  $\Sigma$ . By closure under Boolean operations of  $\mathcal{L}(\mathbf{V} \vee \mathbf{LI})$ , we <sup>568</sup> have that  $C \subseteq \mathcal{L}(\mathbf{V} \vee \mathbf{LI})$ . Now, as Boolean operations commute with both quotients [13, <sup>569</sup> p. 20] and inverses of *ne*-morphisms [13, Proposition 0.4], by closure of  $\mathcal{L}(\mathbf{V})$  and  $\mathcal{L}(\mathbf{LI})$ <sup>570</sup> under quotients and inverses of *ne*-morphisms, we actually have that C is an *ne*-variety <sup>571</sup> of languages. Therefore, by inclusion-wise minimality of  $\mathcal{L}(\mathbf{V} \vee \mathbf{LI})$ , we can conclude that <sup>572</sup>  $\mathcal{L}(\mathbf{V} \vee \mathbf{LI}) = C$ .

<sup>573</sup> **Proof of Lemma 10.** We actually have that  $\mathcal{L}(\mathbf{V}) \cup \mathcal{L}(\mathbf{LI}) \subseteq \mathcal{L}(\mathbf{EV})$ , which allows us to <sup>574</sup> conclude by inclusion-wise minimality of  $\mathcal{L}(\mathbf{V} \vee \mathbf{LI})$  (Proposition 2) and by the fact that <sup>575</sup>  $\mathcal{L}(\mathbf{EV})$  is an *ne*-variety of languages (Proposition 6).

Let  $\Sigma$  be an alphabet. The fact that  $\mathcal{L}(\mathbf{V})(\Sigma) \subseteq \mathcal{L}(\mathbf{EV})(\Sigma)$  follows trivially from Proposition 9. Moreover, for all  $u \in \Sigma^*$ , since necessarily  $\Sigma^* \in \mathcal{L}(\mathbf{V})(\Sigma)$ , we have that both  $u\Sigma^*$  and  $\Sigma^*u$  belong to  $\mathcal{L}(\mathbf{LI})(\Sigma)$ . Thus, as  $\mathcal{L}(\mathbf{EV})(\Sigma)$  is closed under Boolean operations, it follows that  $\mathcal{L}(\mathbf{LI})(\Sigma) \subseteq \mathcal{L}(\mathbf{EV})(\Sigma)$ .

This concludes the proof, since it holds for any alphabet  $\Sigma$ .

◄

<sup>581</sup> **Proof of Lemma 11.** Assume that  $\mathbf{EV} \subseteq \langle \mathbf{V} \lor \mathbf{LI} \rangle_{ne}$ . For any  $L \in \mathcal{L}(\mathbf{V})(\Sigma)$  for  $\Sigma$  some al-<sup>582</sup> phabet and any  $x, y \in \Sigma^*$ , by Proposition 9, we have that  $xLy \in \mathcal{L}(\mathbf{EV})(\Sigma) \subseteq \mathcal{L}(\mathbf{V} \lor \mathbf{LI})(\Sigma)$ . <sup>583</sup> Hence,  $\mathbf{V}$  verifies criterion (A).

<sup>584</sup> Conversely, assume that **V** verifies criterion (A). For any alphabet  $\Sigma$ , the set  $\mathcal{L}(\mathbf{V} \vee \mathbf{LI})(\Sigma)$ <sup>585</sup> contains all languages of the form xLy for  $L \in \mathcal{L}(\mathbf{V})(\Sigma)$  and  $x, y \in \Sigma^*$ , so it contains all <sup>586</sup> Boolean combinations of languages of that form, since it is closed under Boolean operations. <sup>587</sup> Therefore, by Proposition 9, we have  $\mathcal{L}(\mathbf{EV}) \subseteq \mathcal{L}(\mathbf{V} \vee \mathbf{LI})$ , so that  $\mathbf{EV} \subseteq \langle \mathbf{V} \vee \mathbf{LI} \rangle_{ne}$ .

<sup>588</sup> **Proof of Proposition 12.** Let us first observe that given any alphabet  $\Sigma$ , given any language <sup>589</sup> K on that alphabet and given any two words  $x, y \in \Sigma^*$ , we have that  $x(x^{-1}Ky^{-1})y =$ <sup>590</sup>  $x\Sigma^*y \cap K$  and  $x^{-1}(xKy)y^{-1} = K$ .

<sup>591</sup> Implication from right to left. Assume that for any  $L \in \mathcal{L}(\mathbf{V})(\Sigma)$  for some alphabet  $\Sigma$  and <sup>592</sup>  $x, y \in \Sigma^*$ , there exist  $k, l \in \mathbb{N}$  such that for all  $u \in \Sigma^k, v \in \Sigma^l$ , there exists some language <sup>593</sup>  $K \in \mathcal{L}(\mathbf{V})(\Sigma)$  verifying  $u^{-1}Lv^{-1} = (xu)^{-1}K(vy)^{-1}$ . Take some  $L \in \mathcal{L}(\mathbf{V})(\Sigma)$  for some

alphabet  $\Sigma$  and take  $x, y \in \Sigma^*$ . Consider also some  $k, l \in \mathbb{N}$  that are guaranteed to exist by the assumption we just made.

For all  $u \in \Sigma^k, v \in \Sigma^l$ , there exists some language  $K \in \mathcal{L}(\mathbf{V})(\Sigma)$  verifying  $u^{-1}Lv^{-1} = (xu)^{-1}K(vy)^{-1}$ , so that by our observation at the beginning of the proof, we have

$$x(u\Sigma^*v \cap L)y = xu(u^{-1}Lv^{-1})vy = xu((xu)^{-1}K(vy)^{-1})vy = xu\Sigma^*vy \cap K$$

Using Proposition 2, we thus have that  $x(u\Sigma^*v \cap L)y \in \mathcal{L}(\mathbf{V} \vee \mathbf{LI})(\Sigma)$  for all  $u \in \Sigma^k, v \in \Sigma^l$ . Moreover, since we have that the set of words of L of length at least k + l is

$$\Sigma^{\geq k+l} \cap L = \bigcup_{u \in \Sigma^k, v \in \Sigma^l} (u\Sigma^* v \cap L)$$

602 and since

5

$$L = (\Sigma^{\geq k+l} \cap L) \cup F$$

where F is a finite set of words on  $\Sigma$  of length less than k + l, we have that

$$algoremute{1.5} xLy = x \big( (\Sigma^{\geq k+l} \cap L) \cup F \big) y = \bigcup_{u \in \Sigma^k, v \in \Sigma^l} x (u\Sigma^* v \cap L) y \cup xFy \; .$$

We can thus conclude that  $xLy \in \mathcal{L}(\mathbf{V} \vee \mathbf{LI})(\Sigma)$  since  $xFy \in \mathcal{L}(\mathbf{LI})(\Sigma)$  and because  $\mathcal{L}(\mathbf{V} \vee \mathbf{LI})(\Sigma)$  is closed under unions.

Implication from left to right. Assume that **V** satisfies criterion (A). Take some  $L \in \mathcal{L}(\mathbf{V})(\Sigma)$  for some alphabet  $\Sigma$  and take  $x, y \in \Sigma^*$ . By hypothesis, we know that  $xLy \in \mathcal{L}(\mathbf{V} \vee \mathbf{LI})(\Sigma)$ .

By Proposition 2, this means that xLy is a Boolean combination of languages in  $\mathcal{L}(\mathbf{V})(\Sigma) \cup$ 611  $\mathcal{L}(\mathbf{LI})(\Sigma)$ . Further, this implies that xLy can be written as the union of intersections of 612 languages of  $\mathcal{L}(\mathbf{V})(\Sigma)$  and  $\mathcal{L}(\mathbf{LI})(\Sigma)$  or their complements, which in turn implies, by closure 613 of  $\mathcal{L}(\mathbf{V})(\Sigma)$  and  $\mathcal{L}(\mathbf{LI})(\Sigma)$  under Boolean operations, that xLy can be written as a finite 614 union of languages of the form  $K \cap (U\Sigma^*V \cup W)$  with  $K \in \mathcal{L}(\mathbf{V})(\Sigma)$  and  $U, V, W \subseteq \Sigma^*$  finite. 615 Since any word in xLy must be of length at least |xy| and have x as a prefix and y as a 616 suffix, we can assume that any language  $K \cap (U\Sigma^* V \cup W)$  appearing in a finite union as 617 described above verifies that  $U \subseteq x\Sigma^*$ , that  $V \subseteq \Sigma^* y$  and that  $W \subseteq x\Sigma^* y$ . Now, if we take 618  $k, l \in \mathbb{N}$  big enough, we thus have that 619

$$_{620} \qquad xLy = \bigcup_{u \in \Sigma^k, v \in \Sigma^l} (K_{u,v} \cap xu\Sigma^*vy) \cup F$$

where  $K_{u,v} \in \mathcal{L}(\mathbf{V})(\Sigma)$  for all  $u \in \Sigma^k, v \in \Sigma^l$  and  $F \subseteq \Sigma^{<|xy|+k+l}$ . Hence, for all  $u \in \Sigma^k, v \in \Sigma^{l}$  $\Sigma^l$ , we have

$$u^{-1}Lv^{-1} = u^{-1} \left( x^{-1} (xLy)y^{-1} \right) v^{-1}$$

$$= (xu)^{-1} \Big(\bigcup_{u' \in \Sigma^k, v' \in \Sigma^l} (K_{u',v'} \cap xu'\Sigma^*v'y) \cup F\Big)(vy)^{-1}$$

625

624

$$= \bigcup_{\substack{u' \in \Sigma^k, v' \in \Sigma^l \\ (xu)^{-1} F(vy)^{-1}}} (xu'((xu')^{-1}K_{u',v'}(v'y)^{-1})v'y)(vy)^{-1} \cup$$
  
$$= (xu)^{-1}F(vy)^{-1}$$
  
$$= (xu)^{-1}K_{u,v}(vy)^{-1} ,$$

626 627

using classical formulae for quotients [13, p. 20] and observing that  $(xu)^{-1}K(vy)^{-1} = \emptyset$  for any  $K \subseteq \Sigma^*$  such that  $K \cap xu\Sigma^*vy = \emptyset$ .