



HAL
open science

Forward analysis for WSTS, part I: completions

Alain Finkel, Jean Goubault-Larrecq

► **To cite this version:**

Alain Finkel, Jean Goubault-Larrecq. Forward analysis for WSTS, part I: completions. *Mathematical Structures in Computer Science*, 2020, 30 (7), pp.752-832. 10.1017/S0960129520000195 . hal-03186488

HAL Id: hal-03186488

<https://hal.science/hal-03186488v1>

Submitted on 10 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Forward Analysis for WSTS, Part I: Completions

Alain Finkel and Jean Goubault-Larrecq
Université Paris-Saclay, ENS Paris-Saclay, CNRS, LSV,
91190, Gif-sur-Yvette, France

Email: `\protect\T1\textbraceleftfinkel, goubault\protect\T1\textbrace`

Abstract

We define representations for downwards-closed subsets of a rich family of well-quasi-orders, and more generally for closed subsets of an even richer family of Noetherian topological spaces. This includes the cases of finite words, of multisets, of finite trees, notably. Those representations are given as finite unions of ideals, or more generally of irreducible closed subsets. All the representations we explore are computable, in the sense that we exhibit algorithms that decide inclusion, and compute finite unions and finite intersections. The origin of this work lies in the need for computing finite representations of sets of successors of the downward closure of one state, or more generally of a downwards-closed set of states, in a well-structured transition system, and this is where we start: we define adequate notions of completions of well quasi-orders, and more generally, of Noetherian spaces. For verification purposes, we argue that the required completions must be ideal completions, or more generally sobrifications, that is, spaces of irreducible closed subsets.

Contents

1	Introduction	2
2	Preliminaries	5
2.1	Order	6
2.2	Domain Theory	7
2.3	Topology	8
2.4	Sobriety	11
3	Completions of Wqos	12
4	S-Representations	17
5	Completing Ring Ideals	23

*An extended abstract already appeared in Proc. 26th International Symposium on Theoretical Aspects of Computer Science (STACS'09).

6	Completing Words	27
7	Completing Multisets	33
8	Completing Words, Prefix Topology	37
9	Completing Finite Trees in a Simple Case	41
10	Completing Finite Trees: the General Case	50
	10.1 Tree Steps	51
	10.2 Tree Iterators	52
	10.3 Checking Inclusion	55
	10.4 Intersections of Tree Steps and Tree Iterators	58
	10.5 STREs, Tree-Products	60
11	Conclusion	67
A	Auxiliary Proofs on Irreducible Closed Sets	72
B	Proofs of Results on Words (Section 6)	74
C	Proofs of Results on Multisets (Section 7)	80
D	Proofs of Results on Words with the Prefix Topology (Section 8)	83
E	Proofs of Results on Trees (Section 10)	86
	E.1 Tree Steps	90
	E.2 Tree Iterators	91
	E.3 Checking Inclusion between Tree Steps	96
	E.4 Checking Inclusion between Tree Steps and Tree Iterators	97
	E.5 Checking Inclusion between Tree Iterators	98
	E.6 Intersections of Tree-Products	101

1 Introduction

Well-structured transition systems (WSTS) are a paradigmatic class of infinite-state transition systems on which many properties of interest in verification are decidable [Finkel, 1987, Abdulla et al., 1996, Finkel and Schnoebelen, 2001]. They include Petri nets, affine counter systems, lossy channel systems, data nets, and many more.

Briefly put, a *WSTS* is a triple (X, \rightarrow, \leq) where \leq is a well quasi-order on the (possibly infinite) state space X , and \rightarrow is a monotonic transition relation on X . (We define well quasi-orders in Section 2.1.) To simplify things slightly, by *monotonic* we mean strongly monotonic, namely that if $x \rightarrow x'$ and $x \leq y$, then there is a state y' such that $x' \leq y'$ and $y \rightarrow y'$. The set of one-step predecessors $\text{Pre}(E) = \{x \in X \mid \exists x' \in E, x \rightarrow x'\}$ of any upwards-closed subset E is then upwards-closed again, where E is *upwards-closed* if and only if $x \in E$ and $x \leq y$ imply

$y \in E$. Similarly, the sets $\text{Pre}^k(E)$ of k -step predecessors, $\text{Pre}^{\leq k}(E)$ of at-most- k -step predecessors, and $\text{Pre}^*(E) = \bigcup_{k \in \mathbb{N}} \text{Pre}^{\leq k}(E)$ of iterated predecessors of the upwards-closed set E are upwards-closed. The fact that \leq is a well quasi-order implies (see Section 2.1 again) that every upwards-closed subset is the upward closure $\uparrow A$ of a *finite* set of points A (a *basis* of the set). This implies that $\text{Pre}^*(E) = \text{Pre}^{\leq k}(E)$ for some $k \in \mathbb{N}$: write $\text{Pre}^*(E)$ as $\uparrow A$ with $A = \{x_1, \dots, x_n\}$, realize that for each i , x_i must be in $\text{Pre}^{\leq k}(E)$ for some k , and take the largest of these k s. In particular, there is a simple algorithm that decides *coverability* in WSTS, namely, which decides whether, given $x \in X$ and a basis for an upwards-closed subset E , whether one can reach an element of E in finitely many \rightarrow steps starting from x [Abdulla et al., 2000]: compute $\text{Pre}^{\leq 0}(E) = E$, iterate using $\text{Pre}^{\leq k+1}(E) = E \cup \text{Pre}(\text{Pre}^{\leq k}(E))$ until $\text{Pre}^{\leq k+1}(E) \subseteq \text{Pre}^{\leq k}(E)$, at which point $\text{Pre}^{\leq k}(E) = \text{Pre}^*(E)$; then test whether $x \in \text{Pre}^*(E)$. For this to work, we need the WSTS to be effective, which means that we can compute $\text{Pre}(E)$ for E upwards-closed, and we can test inclusion.

This was generalized to topological WSTS by the second author [Goubault-Larrecq, 2010]; the algorithm is the same, but X must now be a Noetherian space (see Section 2.3), the sets E are required to be open, and \rightarrow is a lower semi-continuous relation. Topological WSTS include WSTS, but also some other infinite-state systems, among which the class of lossy concurrent polynomial games is probably the most interesting new instance—see Section 6 of [Goubault-Larrecq, 2010].

The algorithm described above works *backwards*, but sometimes we would prefer a forward algorithm that would compute $\downarrow \text{Post}^*({x})$, where \downarrow denotes downward closure, $\text{Post}(E) = \{x' \in X \mid \exists x \in E, x \rightarrow x'\}$ is the set of one-step successors of E and $\text{Post}^*(E) = \bigcup_{k \in \mathbb{N}} \text{Post}^k(E)$. The set $\downarrow \text{Post}^*({x})$ is called the *cover* of x , and can be used to decide coverability as well: one can reach $\uparrow E$ from x if and only if $\downarrow \text{Post}^*({x})$ intersects E .

However, although the backward procedure always terminates, it is often slow. Forward procedures, when they exist, may fail to terminate: on lossy channel systems, any terminating forward procedure would enable us to decide boundedness, which is undecidable [Mayr, 2003]. But they often give results faster in practice. For this reason, only the non-terminating forward procedure is implemented in the tool TREX [Abdulla et al., 1998].

The cover also provides more useful information than the set computed by the backward algorithm. For example, the cover is a good first approximation of the reachability set $\text{Post}^*({x})$, and the original reachability algorithms for Petri nets rely on the computation of covers [Kosaraju, 1982, Mayr, 1981, Lambert, 1992]. This can also serve as a first step toward model checking liveness properties, as in [Emerson and Namjoshi, 1998] and more recently in [Blondin et al., 2017a, Blondin et al., 2017b].

For Petri nets, the cover can be computed by the so-called coverability tree algorithm of Karp and Miller [Karp and Miller, 1969]. Part II of this paper generalizes this to a large class of WSTS [Finkel and Goubault-Larrecq, 2012]. Part III of this paper defines and studies very-WSTS, a subclass of WSTS, for which the cover is computable and for which linear temporal logic (introduced in [Pnueli, 1977]) model checking is decidable. The present part I deals with an important preparatory step: characterizing downwards-closed subsets of well quasi-ordered sets X . We shall see that such downwards-closed subsets can always be written as the downward closure of *finitely*

many points in a completion \widehat{X} of X . In fact, we start by defining the possible relevant completions from a verification perspective, and realize that the smallest possible one is the ideal completion, or equivalently (in the more general, Noetherian case) the sobrification of X . We shall then explore concrete computer representations for elements of \widehat{X} , for a large class of Noetherian spaces X (in particular, well quasi-orders) that includes most of the spaces needed in the verification of WSTS today.

This paper is an extended version of [Finkel and Goubault-Larrecq, 2009]. Before this paper, and except for some partial results [Finkel, 1990, Emerson and Namjoshi, 1998, Geeraerts et al., 2006], a general theory of downwards-closed sets was missing. This may explain the scarcity of forward algorithms for WSTS. Quoting Abdulla *et al.* [Abdulla et al., 2004b]: “Finally, we aim at developing generic methods for building downwards-closed languages, in a similar manner to the methods we have developed for building upwards-closed languages in [Abdulla et al., 2000]. This would give a general theory for forward analysis of infinite state systems, in the same way the work in [Abdulla et al., 2000] is for backward analysis.” Our contribution is to provide such a theory of downwards-closed sets.

Related Work

The coverability for general WSTS was shown decidable by using a backward algorithm presented in 1996 [Abdulla et al., 1996]; this algorithm was an abstraction of the coverability algorithm for lossy channel systems [Abdulla and Jonsson, 1993]. Coverability for vector addition systems with resets had been shown decidable by Arnold and Latteux [Arnold and Latteux, 1978, Theorem 5, p. 391]. Interestingly, the latter algorithm is an early instance of the backward algorithm presented in [Abdulla et al., 1996], and applied to \mathbb{N}^n .

While this paper is not about algorithms, it is worth recalling that the inspiration for our line of work, which culminates in part III [Blondin et al., 2017a, Blondin et al., 2017b], comes from Karp and Miller’s celebrated finite coverability tree algorithm [Karp and Miller, 1969] for Petri nets. This arguably computes a finite representation of the cover $\downarrow \text{Post}^*(\{x\})$, and we expand on that in [Finkel and Goubault-Larrecq, 2012, Section 4.1] and in [Blondin et al., 2017a, Blondin et al., 2017b]. Further related work on this issue can be found in that paper. What matters to us here is that, while the state space of a Petri net is \mathbb{N}^k , Karp and Miller’s finite representation is given by finitely many points in the *completion* \mathbb{N}_ω^k , where \mathbb{N}_ω is \mathbb{N} plus a fresh, infinite element ω .

The focus of this paper is on *finite representations* of downwards-closed subsets of well quasi-ordered sets, and more generally of closed subsets of Noetherian spaces. In computer speak, we focus on data structures rather than algorithms. Mathematically, we shall need to define the right notion of completion \widehat{X} for well quasi-ordered sets, resp., Noetherian spaces X —these will be the familiar constructions of ideal completion, resp. sobrification—and to study finite representations of their (downwards-)closed subsets.

Data structures are a prerequisite to define algorithms. In our context, one may argue that what we need is a so-called *adequate domain of limits (ADL)*, as defined by Ganty, Geeraerts, Raskin and van Begin [Geeraerts et al., 2006, Ganty et al., 2006].

An ADL is an axiomatization of a data structure on which the authors’ expand, enlarge and check procedure, which computes the cover, works. Alternatively, an ADL is an axiomatization for a relevant completion \widehat{X} . We shall see that these completions have strong ties with the ideal completion, resp., sobrification, mentioned above.

In the special case of finite words, such finite representations were developed by Abdulla, Colomb-Annichini, Bouajjani and Jonsson [Abdulla et al., 1998] as specific regular expressions called SREs (simple regular expressions) and word products. In their case, the alphabet is finite, with equality as well quasi-order. Similar representations also apply to certain more complex well quasi-ordered sets of letters, as demonstrated in [Abdulla et al., 2004b] for example. More generally, it had been shown by Kabil and Pouzet that this representation is in fact valid for any arbitrary well quasi-ordered set of letters [Kabil and Pouzet, 1992]. We improve on this slightly by showing that this even works for all Noetherian sets of letters (Section 6). Interestingly, one of the key notions we use in the proof is that of an irreducible (closed) subset, which comes from topology. This is also a central concept in Kabil and Pouzet’s proof, narrowed down to well quasi-orders.

We define finite representations for (downwards-)closed subsets of a large class of Noetherian data types, including tuples of natural numbers or finite words, as mentioned above, but also many more. Some of them are representations of (downwards-)closed subsets in well-known well quasi-orders, as in the case of finite multisets (Section 7), or of finite labeled trees (Section 10; by far the most technical part of this paper. The fact that the right completion for languages of trees can be described as certain regular tree languages was also observed by Wies, Zufferey and Henzinger, for finite sets of labels [Wies et al., 2010]. They did not characterize what kind of regular tree language is required precisely, which we do.) Some others are representations of closed subsets of Noetherian spaces that do not arise from well quasi-orders. For instance, we deal with polynomial ideals in Section 5, with finite words again but with a different topology, the prefix topology, in Section 8.

Acknowledgments

We thank Philippe Schnoebelen, Simon Halfon, Sylvain Schmitz, Ranko Lazic, and all those who expressed interest in this work since 2009, and encouraged us into submitting this paper again—not just out of friendship, but also because they need the results. Simon Halfon found the polynomial time decision procedure mentioned in Lemma 7.6, and is gratefully acknowledged for this. The referees have done a remarkably thorough job in rereading this paper, and we thank them for that. Any remaining mistake is, of course, ours.

2 Preliminaries

We shall borrow from theories of well quasi-orderings, as used classically in well-structured transition systems [Abdulla et al., 2000, Finkel and Schnoebelen, 2001], from domain theory [Abramsky and Jung, 1994, Gierz et al., 2003], as well as from topol-

ogy [Goubault-Larrecq, 2013]. We recap most of what we need. The purpose is not to give a crash course on these three fields, rather to fix notations and notions.

2.1 Order

A *quasi-ordering* \leq is a reflexive and transitive relation on a set X . It is a (partial) *ordering* iff it is antisymmetric. A set X equipped with a partial ordering is a *poset*.

We write \geq for the opposite quasi-ordering, \approx for the equivalence relation $\leq \cap \geq$, $<$ for the associated strict ordering ($\leq \setminus \approx$), and $>$ for the converse ($\geq \setminus \approx$) of $<$. The *upward closure* $\uparrow E$ of a set E is $\{y \in X \mid \exists x \in E, x \leq y\}$. The *downward closure* $\downarrow E$ is $\{y \in X \mid \exists x \in E, y \leq x\}$. A subset E of X is *upwards-closed* if and only if $E = \uparrow E$, i.e., any element greater than or equal to some element in E is again in E , which was the definition we gave in the introduction. The notion of *downwards-closed* sets is defined similarly. When the ambient space X is not clear from context, we shall write $\downarrow_X E$, $\uparrow_X E$ instead of $\downarrow E$, $\uparrow E$. We also write $\uparrow x$ instead of $\uparrow\{x\}$, and $\downarrow x$ instead of $\downarrow\{x\}$.

A quasi-ordering is *well-founded* iff it has no infinite strictly descending chain, i.e., $x_0 > x_1 > \dots > x_i > \dots$. An *antichain* is a set of pairwise incomparable elements. A quasi-ordering is *well* if and only if it is well-founded and has no infinite antichain.

There are a number of equivalent definitions for *well quasi-orderings* (*wqo*). One is that, from any infinite sequence $x_0, x_1, \dots, x_i, \dots$, one can extract an infinite ascending chain $x_{i_0} \leq x_{i_1} \leq \dots \leq x_{i_k} \leq \dots$, with $i_0 < i_1 < \dots < i_k < \dots$. Another one is that any upwards-closed subset can be written $\uparrow E$, with E *finite*. Such a finite E is called a *finite basis* for the upwards-closed set. In a wqo, every upwards-closed set has a minimal finite basis, composed of the subset of its pairwise incomparable, minimal elements. We shall see another, topological, characterization of wqos below.

There is a rich supply of wqos. First, for any $k \in \mathbb{N}$, \mathbb{N}^k is a wqo in the product ordering ($(x_1, \dots, x_k) \leq (y_1, \dots, y_k)$ iff $x_i \leq y_i$ for every i , $1 \leq i \leq k$): this is *Dickson's Lemma* [Dickson, 1913]. \mathbb{N}^k is the set of configurations of Petri nets, or more generally, of counter machines.

For every well-quasi ordered alphabet Σ , Σ^* with the embedding (a.k.a. scattered subword, a.k.a. divisibility) quasi-ordering is wqo: this is *Higman's Lemma* [Higman, 1952]. This is instrumental in the backward analysis of lossy channel systems [Abdulla and Jonsson, 1993]. Under the same assumptions, the collection of finite trees labeled with elements from Σ , with the tree embedding quasi-ordering, is wqo: this is *Kruskal's Tree Theorem* [Kruskal, 1960].

A map f from a quasi-ordered set X to a quasi-ordered set Y is *monotonic* if and only if $x \leq x'$ implies $f(x) \leq f(x')$, for all $x, x' \in X$. (We write \leq for the underlying ordering of any poset, unless mentioned otherwise.) We call it a *quasi-order embedding* if and only if $x \leq x'$ is *equivalent* to $f(x) \leq f(x')$. The *order embeddings* are the injective quasi-order embeddings; there is no difference between the two notions when X is a poset. An *order isomorphism* is a surjective (hence bijective) order embedding. Hence $f: X \rightarrow Y$ is an order embedding if and only if f is an order isomorphism onto its image.

Given any quasi-ordered set X , the *order quotient* of X is defined as the set of equivalence classes $[x]$ of elements $x \in X$ under \approx , quasi-ordered by letting $[x]$ be

below $[y]$ iff $x \leq y$. (We then write $[x] \leq [y]$.) This is well-defined, and a partial order.

We shall say that a set is *well-ordered* by \leq iff it is well-quasi-ordered by \leq and \leq is an ordering. The well-ordered posets are exactly the order quotients of wqos.

In a quasi-ordered set X , an *upper bound* of a family $(x_i)_{i \in I}$ of points of X is an element $x \in X$ such that $x_i \leq x$ for every $i \in I$. A *least upper bound* is one that is less than or equal to all other upper bounds of the same family. If X is a poset, then the least upper bound of a family is unique if it exists at all.

2.2 Domain Theory

Domain theory is, *prima facie*, concerned with certain posets, called dcpos, where certain least upper bounds exist, and so-called Scott-continuous maps, which are not just monotonic but also preserve these least upper bounds. Over the years, domain theory has revealed itself as having firm grounds in general topology as well. Let us start with the order-theoretic view.

A *directed family* in a poset X is any non-empty family $(x_i)_{i \in I}$ such that, for all $i, j \in I$, there is a $k \in I$ with $x_i, x_j \leq x_k$. A *directed-complete partial order*, in short a *dcpo*, is a poset X in which every directed family $(x_i)_{i \in I}$ of points of X has a least upper bound $\sup_{i \in I} x_i$.

A map f from a poset X to a poset Y is *Scott-continuous* if and only if it is monotonic and preserves least upper bounds of directed families, i.e., if $(x_i)_{i \in I}$ is a directed family in X with least upper bound x , then $f(x)$ is the least upper bound of the (necessarily directed) family $(f(x_i))_{i \in I}$.

An element $x \in X$ is *finite* iff, for every directed family $(z_i)_{i \in I}$ that has a least upper bound $z \geq x$, then $z_i \geq x$ for some $i \in I$ already. The poset X is *algebraic* iff the family of finite elements below any given element x is directed, and admits x as least upper bound. The finite elements are often much simpler to describe than arbitrary elements, and act as approximants to the latter.

Let us give a few examples. The power $\mathbb{P}(X)$ of a set X is a dcpo, in fact a complete lattice, under inclusion \subseteq . Its finite elements, in the sense above, are the finite subsets of X , in the usual sense of the word, and $\mathbb{P}(X)$ is algebraic. We write $\mathbb{P}_{\text{fin}}(X)$ for the set of finite subsets of X , ordered by inclusion.

Neither $\mathbb{P}(X)$ nor $\mathbb{P}_{\text{fin}}(X)$ is wqo under inclusion, unless X is finite. \mathbb{N} , with its natural ordering, is an algebraic poset, which is also a wqo. \mathbb{N} is not a dcpo, since \mathbb{N} itself is a directed family without a least upper bound. However, \mathbb{N}_ω , obtained by adjoining a new top element ω to \mathbb{N} , is a dcpo. Its finite elements are the elements of \mathbb{N} , and \mathbb{N}_ω is algebraic.

Some dcpos fail to be algebraic. E.g., the only finite element of $[0, 1]$, with its natural ordering, is 0. However, $[0, 1]$ is continuous, in the following sense.

Define the *way below* relation \ll on a poset X by $x \ll y$ iff, for every directed family $(z_i)_{i \in I}$ that has a least upper bound $z \geq y$, then $z_i \geq x$ for some $i \in I$ already. So, in particular, the finite elements are those that are way below themselves.

Note that $x \ll y$ implies $x \leq y$, and that $x' \leq x \ll y \leq y'$ implies $x' \ll y'$. However, \ll is not reflexive or irreflexive in general. Write $\uparrow E = \{y \in X \mid \exists x \in E, x \ll y\}$, $\downarrow E = \{y \in X \mid \exists x \in E, y \ll x\}$.

The poset X is *continuous* iff, for every $x \in X$, $\downarrow x$ is a directed family, and has x as least upper bound. More finely, call a *basis* (not to be confused with the finite bases of upwards-closed subsets of wqos) any subset B of X such that any element $x \in X$ is the least upper bound of a directed family of elements way below x in B . Then X is continuous if and only if it has a basis, and in this case X itself is the largest basis. On the other hand, every algebraic poset is continuous, and has a least basis, namely its set of finite elements.

An essential property of continuous posets is *interpolation* [Mislove, 1998, Lemma 4.16]: if $x \ll y$, then $x \ll z \ll y$ for some $z \in X$. We may even choose z to be in any prescribed basis B . For example, in $[0, 1]$, $x \ll y$ iff $x < y$ or $x = 0$, and we may choose B to be, say, the set of rational points in $[0, 1]$. Interpolation fails in general, non-continuous posets, even non-continuous dcpos.

Any finite product of dcpos is a dcpo, where product is taken in the order-theoretic sense, i.e., with the product ordering. Then any finite product of algebraic (resp., continuous) posets is again algebraic (resp., continuous).

Given a poset X , which might fail to be a dcpo, there is a canonical way to obtain a completion, called the *ideal completion* $\mathbf{I}(X)$ of X . An *ideal* I of X is any downwards-closed set that is also directed. $\mathbf{I}(X)$ is defined as the poset of all ideals of X , ordered by inclusion. $\mathbf{I}(X)$ is then a dcpo, where directed suprema are computed as unions, and X order-embeds into $\mathbf{I}(X)$ through the function $\eta^{\mathbf{I}}: X \rightarrow \mathbf{I}(X)$ that maps x to $\downarrow x$. For example, $\mathbf{I}(\mathbb{N})$ consists of all the ideals $\downarrow n$, $n \in \mathbb{N}$, plus a fresh element above all others, which we write ω and, as an ideal, is just the whole of \mathbb{N} . In this sense, $\mathbf{I}(\mathbb{N})$ is the completion \mathbb{N}_ω we have already mentioned in the context of the Karp-Miller algorithm.

$\mathbf{I}(X)$ is the *free dcpo* over X , meaning that for every monotonic map f from X to a dcpo Y extends to a unique Scott-continuous map g from $\mathbf{I}(X)$ to Y —namely, $f = g \circ \eta^{\mathbf{I}}$, see [Goubault-Larrecq, 2013, Exercise 5.5.3, or comment pages 175–176]. $\mathbf{I}(X)$ is also an algebraic dcpo [Goubault-Larrecq, 2013, Proposition 5.1.46], with the elements of X forming a basis.

2.3 Topology

A *topology* \mathcal{O} on a set X is a collection of subsets (the *opens*) of X that is closed under arbitrary unions and finite intersections. In particular, considering empty unions and empty intersections, both \emptyset and X itself are open. We say that X itself is a topological space, leaving \mathcal{O} implicit. The complements of opens are the *closed sets*. The largest open contained in A is its *interior*, the smallest closed subset $cl(A)$ containing it is its *closure*.

A famous topology in domain theory is the *Scott topology* on a poset X . Its opens, the *Scott opens*, are all upwards-closed subsets U such that every directed family $(x_i)_{i \in I}$ that has a least upper bound x in U intersects U , i.e., $x_i \in U$ for some $i \in I$. In other words, the closed subsets of the topology, namely the *Scott closed* subsets, are the downwards-closed subsets F that are stable under taking least upper bounds of directed families of elements of F . The non-empty Scott closed subsets of $[0, 1]$ are the intervals $[0, t]$, $0 \leq t \leq 1$, and its Scott open subsets are the half-open intervals $(t, 1]$, $0 \leq t \leq 1$, plus $[0, 1]$ itself.

Order	Topology
upwards-closed	open
downwards-closed	closed
monotonic	continuous
wqo	Noetherian
ideal	irreducible closed
ideal completion $I(X)$	sobrification $\mathcal{S}(X)$

Figure 1: Informal, order vs. topology glossary

A topology is *coarser* than another iff it contains less opens. Conversely, a topology is *finer* than another iff it contains more opens.

For example, consider the *Alexandroff topology* of a quasi-order X , whose opens are all upwards-closed subsets. This is finer than the Scott topology, and in general strictly finer: on $[0, 1]$, $[1/2, 1]$ is Alexandroff open but not Scott open. On \mathbb{N} , the Scott and Alexandroff topologies agree, and the non-empty opens are of the form $\uparrow n$, $n \in \mathbb{N}$. The *discrete topology* is the finest possible topology, where all subsets are open. Note that this is also the Alexandroff topology of the equality ordering.

The Alexandroff topology converts a quasi-order into a topological space, and suggests a glossary of generalizations of order-theoretic notions as topological notions, see Figure 1. We have just explained the first row: the upwards-closed subsets of a quasi-order are the opens in the Alexandroff topology. We shall explain the other rows below.

We shall write X_σ for X with its Scott topology, and X_a for X with its Alexandroff topology. It is easy to see that the downwards-closed subsets of X are exactly the closed subsets of X_a , and we shall use this fact several times. This is the second row of Figure 1.

A map f from a topological space X to a topological space Y is *continuous* if and only if $f^{-1}(V)$ is open in X for every open subset V of Y . When both X and Y are posets equipped with the Alexandroff topology, a map $f: X \rightarrow Y$ is continuous if and only if it is monotonic. This is the third row of Figure 1. When X and Y are posets equipped with their Scott topology, then f is continuous if and only if it is Scott-continuous.

A *homeomorphism* is a topological isomorphism, i.e., a continuous, bijective map whose inverse is also continuous.

Given any collection \mathcal{C} of subsets of a set X , there is a smallest (*coarsest*) topology containing all elements of \mathcal{C} . This is the topology *generated by* \mathcal{C} , \mathcal{C} is then called a *subbase* for the topology, and the elements of \mathcal{C} are *subbasic opens*. Their complements are the *subbasic closed subsets*. Any open in the topology is then a (possibly infinite) union of finite intersections of subbasic opens. If any open can be written as a union of elements of \mathcal{C} , then one says that \mathcal{C} is a *base* of the topology, and the elements of \mathcal{C} are *basic opens*. This occurs typically when \mathcal{C} contains X and is closed under binary intersections.

In a continuous poset, $\uparrow x$ is Scott-open for all x , and every Scott-open set U is a union of such sets, viz. $U = \bigcup_{x \in U} \uparrow x$ [Abramsky and Jung, 1994]. I.e., the subsets

$\uparrow x$ form a base of the Scott topology. Note that the subsets $\uparrow x$ form a base of the Alexandroff topology instead.

Every topology comes with a *specialization quasi-ordering* \leq , defined as $x \leq y$ iff every open that contains x also contains y ; equivalently, iff $x \in cl\{y\}$. It is easy to see that every open is upwards-closed with respect to \leq . The converse need not hold. A subset A of X is *saturated* iff A equals the intersection of all opens U containing A , equivalently iff it is upwards-closed with respect to \leq . The specialization quasi-ordering of both the Scott and Alexandroff topologies of a poset X ordered by \leq is \leq again.

In fact, the Alexandroff topology is the finest having this property. The coarsest is called the *upper topology*; its opens are arbitrary unions of complements of sets of the form $\downarrow E$, E finite. And the Scott topology is somewhere inbetween. The sets $\downarrow E$, with E finite, will play an important role, and we call them the *finitary closed* subsets. These are closed in the upper, Scott, and Alexandroff topologies.

Paralleling the notations X_σ , X_a , we write X_u for X with its upper topology.

A topological space X is T_0 iff for any two distinct points $x, y \in X$, there is an open subset containing x but not y , or conversely. X is T_0 if and only if its specialization quasi-ordering \leq is a partial ordering, i.e., $x \leq y$ and $y \leq x$ imply $x = y$.

A *subspace* of a topological space X is a subset A of X with the so-called *subspace topology*, whose opens are $A \cap U$, U open in X .

The *product* $\prod_{i \in I} X_i$ of a family $(X_i)_{i \in I}$ of topological spaces is the space of tuples $\vec{x} = (x_i)_{i \in I}$ where each x_i is in X_i , and with the *product topology*. The latter is the coarsest that makes the projection maps $\pi_i: \vec{x} \mapsto x_i$ continuous. In other words, the sets $\pi_i^{-1}(U)$, $i \in I$, U open in X_i , form a subbase of the product topology. The binary product of X and Y is written $X \times Y$, and the open subsets of the product topology on the latter are the unions $\bigcup_{i \in I} U_i \times V_i$, where I is an arbitrary index set, U_i open in X and V_i open in Y .

A *topological embedding* f of X into Y is a map from X to Y that is a homeomorphism of X onto the image $f[X] = \{f(x) \mid x \in X\}$ of f , seen as a subspace of Y . Equivalently, f is a topological embedding if and only if it is injective, continuous, and *almost open* in the sense that every open subset U of X is the inverse image $f^{-1}(V)$ of some open subset V of Y . A trivial example is the canonical injection $i: A \rightarrow X$ of a subspace A of X into X . Up to homeomorphism, these are the only topological embeddings: any topological embedding $f: X \rightarrow Y$ is by definition the composition of the canonical injection of $f[X]$ into Y with the homeomorphism $f: X \rightarrow f[X]$.

Given an equivalence relation \equiv on a topological space X , we can form the *quotient space* X/\equiv . Its elements are the equivalence classes of elements of X modulo \equiv . The map $q: X \rightarrow X/\equiv$ sending every element to its equivalence class is called the *quotient map*, and X/\equiv is then given the *quotient topology*, defined as the finest topology on X/\equiv that makes q continuous. Explicitly, the opens of the quotient topology are exactly the subsets V of X/\equiv such that $q^{-1}(V)$ is open in X .

A crucial notion in topology is compactness. A subset K of X is *compact* iff every *open cover* $(U_i)_{i \in I}$ (a family of opens U_i whose union contains K) contains a finite subcover. Alternatively, K is compact iff, for every directed family $(U_i)_{i \in I}$ of opens (directed with respect to inclusion) such that $K \subseteq \bigcup_{i \in I} U_i$, then $K \subseteq U_i$ for some $i \in I$ already.

A topological space X is *Noetherian* iff every open subset of X is compact [Grothendieck, 1960, chapitre 0, § 2]. A less intimidating definition is that X is Noetherian if and only if its lattice of open subsets has the *ascending chain condition*: every properly ascending chain $U_0 \subsetneq U_1 \subsetneq \cdots \subsetneq U_n \subsetneq$ of opens must be finite [Goubault-Larrecq, 2013, Proposition 9.7.6].

There is a strong link between Noetherian spaces and wqos: a poset X is wqo if and only if X is Noetherian in its Alexandroff topology [Goubault-Larrecq, 2013, Proposition 9.7.17]. So wqos are a special case of Noetherian spaces, yielding the fourth row of Figure 1. But there are more Noetherian spaces. We shall see a few of them in this paper, and we only mention two examples for now.

One of the simplest examples, although somehow artificial, is \mathbb{N} with the *cofinite topology*, whose closed subsets are \mathbb{N} plus all finite subsets of \mathbb{N} . This is Noetherian, because every properly descending chain of closed sets must be finite; by taking complements, every properly ascending chain of open sets is finite. If that were a wqo with the Alexandroff topology of some quasi-ordering, that quasi-ordering would have to be the specialization quasi-ordering of the space, which is equality. However, equality on \mathbb{N} is not wqo, since \mathbb{N} itself is an infinite antichain. In fact, the Alexandroff topology of $=$ is the discrete topology, which is much finer than the cofinite topology.

The primary example of a Noetherian space, \mathbb{C}^k with its Zariski topology [Goubault-Larrecq, 2013, Exercise 9.7.53], is far from arising from a wqo as well: its specialization quasi-ordering is equality = again, and the whole space is an infinite antichain. This is one of the ingredients used in the study of the lossy concurrent polynomial games mentioned in the introduction.

2.4 Sobriety

For this section, we refer to [Abramsky and Jung, 1994, Section 7.2.1] or to Chapter 8 of [Goubault-Larrecq, 2013].

A closed subset C of a topological space is *irreducible* if and only if C is non-empty, and whenever $C \subseteq F_1 \cup F_2$ with F_1, F_2 closed, then $C \subseteq F_1$ or $C \subseteq F_2$. Equivalently, if C is included in a finite union of closed subsets F_1, \dots, F_n (whatever $n \in \mathbb{N}$), then $C \subseteq F_i$ for some $i, 1 \leq i \leq n$.

The finitary closed subset $\downarrow x = cl(\{x\})$ ($x \in X$) is always irreducible. (When we write $\downarrow x$ in a topological space, this is relative to its specialization quasi-ordering.) A space X is *sober* iff every irreducible closed subset C is the closure of a unique point, i.e., $C = \downarrow x$ for some unique x . Every sober space is T_0 , and every continuous dcpo is sober in its Scott topology, see [Abramsky and Jung, 1994, Proposition 7.2.27] or [Goubault-Larrecq, 2013, Proposition 8.2.12 (b)].

Much as we could complete a poset X to a dcpo $\mathbf{I}(X)$, we can complete a topological space to its *sobrification* $\mathcal{S}(X)$. The elements of $\mathcal{S}(X)$ are the irreducible closed subsets of X . Its opens are the subsets of the form $\diamond U = \{C \in \mathcal{S}(X) \mid C \cap U \neq \emptyset\}$, U open in X . (This is a topology, not just a subbase.)

As an example, in a poset X with its Alexandroff topology, not only all sets of the form $\downarrow x$ are irreducible closed, but every ideal is irreducible closed, too. We let the reader check this, and also that the converse holds: the ideals of a poset X are exactly the irreducible closed subsets of X_a , leading to the fifth row of Figure 1. This goes

much further: by *Hoffmann's Theorem* [Hoffmann, 1979b], for a poset X , the sobrification $\mathcal{S}(X_a)$ coincides with the ideal completion $\mathbf{I}(X)$ exactly [Goubault-Larrecq, 2013, Fact 8.2.49]. This means that the points are the same, but also the topologies, that is, the topology of $\mathcal{S}(X_a)$ is the Scott topology of $\mathbf{I}(X)$. This justifies the sixth row of Figure 1.

$\mathcal{S}(X)$ is always sober, and the map $\eta_X^{\mathcal{S}} : x \mapsto \downarrow x$ is a topological embedding of X inside $\mathcal{S}(X)$ as soon as X is T_0 . I.e., up to isomorphism, any T_0 space can be seen as a subspace of its sobrification $\mathcal{S}(X)$, equating $x \in X$ with $\downarrow x$ in $\mathcal{S}(X)$.

The sobrification $\mathcal{S}(X)$ of X can be thought of as X together with all missing limits from X . Note in particular that a sober space is always a dcpo in its specialization ordering, see [Abramsky and Jung, 1994, Proposition 7.2.13] or [Goubault-Larrecq, 2013, Corollary 8.2.23].

A topological space X is Noetherian if and only if $\mathcal{S}(X)$ is Noetherian [Goubault-Larrecq, 2013, Lemma 9.7.9]. This is clear from the fact that, up to natural order isomorphism, X and $\mathcal{S}(X)$ have the same opens, see [Gierz et al., 2003, Proposition V-4.7(i)] or [Goubault-Larrecq, 2013, Lemma 8.2.26]. Actually, $\mathcal{S}(X)$ is the *free sober space* above the topological space X , meaning that every continuous map f from X to a sober space Y extends to a unique continuous map g from $\mathcal{S}(X)$ to Y , in the sense that $f = g \circ \eta_X^{\mathcal{S}}$, see [Gierz et al., 2003, Exercise V-4.9] or [Goubault-Larrecq, 2013, Theorem 8.2.44]. This is a form of extension by continuity theorem, and is the proper categorical way of saying that $\mathcal{S}(X)$ is X plus all missing limits.

$\mathcal{S}(X)$, as a space of specific closed subsets of X , embeds into the *Hoare powerdomain* $\mathcal{H}_V(X)$, namely the space of all non-empty closed subsets of X . Let also $\mathcal{H}_V(X)_{\perp}$ be the *lifted Hoare powerdomain* of X , which one can see either as $\mathcal{H}_V(X)$ plus a fresh bottom element \perp added, or as the set of all closed subsets of X , including the empty set. The topology of both $\mathcal{H}_V(X)$ and $\mathcal{H}_V(X)_{\perp}$ is the so-called *lower Vietoris topology*, whose subbasic opens are $\diamond U = \{F \in \mathcal{H}_V(X) \mid F \cap U \neq \emptyset\}$, U open in X . With this topology, $\mathcal{S}(X)$ can be considered a subspace of $\mathcal{H}_V(X)$, and the latter as a subspace of $\mathcal{H}_V(X)_{\perp}$. We use a slightly different symbol $\diamond U$ here, compared to the open subsets $\diamond U$ of $\mathcal{S}(X)$: although they denote very similar sets (and $\diamond U = \diamond U \cap \mathcal{S}(X)$), the sets $\diamond U$ only form a subbase of the lower Vietoris topology on $\mathcal{H}_V(X)$ and $\mathcal{H}_V(X)_{\perp}$, while the sets $\diamond U$ are *exactly* all the open subsets of $\mathcal{S}(X)$.

Remarkably, $\mathcal{H}_V(X)$ and $\mathcal{H}_V(X)_{\perp}$ are Noetherian for every Noetherian space X [Goubault-Larrecq, 2013, Exercise 9.7.14], even though their specialization quasi-ordering, which is inclusion, is in general not wqo.

3 Completions of Wqos

We have announced that the proper completion \widehat{X} of a wqo, or more generally of a Noetherian space X , would be its ideal completion, or more generally its sobrification. Before we compute finite representations, it is in order to vindicate this choice.

A rational way to define a completion is to state the properties we need for it first, and then derive what it should be. In our case, there are various properties we might want for a completion, depending on the point of view we take. In the conference

version of this paper [Finkel and Goubault-Larrecq, 2009], we had explored several of these points of view.

Let us concentrate on just one: Geeraerts *et al.*'s axiomatization of so-called *adequate domain of limits* for well-quasi-ordered sets X , used in their expand, enlarge, and check forward procedure [Geeraerts et al., 2006]. We stress that this notion is independent of their algorithm, and of any particular algorithm: adequate domains of limits are merely an axiomatization of some basic requirements on the representability of downwards-closed subsets. These requirements are also needed in our own approach [Finkel and Goubault-Larrecq, 2012].

An *adequate domain of limits* [Geeraerts et al., 2006] (*ADL*) for a well-ordered set X is a triple (L, \preceq, γ) where L is a set disjoint from X (the set of *limits*); (L_1) the map $\gamma: L \cup X \rightarrow \mathbb{P}(X)$ is such that $\gamma(z)$ is downwards-closed for all $z \in L \cup X$, and $\gamma(x) = \downarrow_X x$ for all non-limit points $x \in X$; (L_2) there is a limit point $\top \in L$ such that $\gamma(\top) = X$; (L_3) for all $z, z' \in L \cup X$, $z \preceq z'$ if and only if $\gamma(z) \subseteq \gamma(z')$; and (L_4) for any downwards-closed subset D of X , there is a finite subset $E \subseteq L \cup X$ such that $\hat{\gamma}(E) = D$. Here $\hat{\gamma}(E) = \bigcup_{z \in E} \gamma(z)$.

No explicit construction for such adequate domains of limits is given by Geeraerts *et al.*, and they have to be found by trial and error. Our first result, below, is that there is a unique least (weak) adequate domain of limits of X , and this is $\mathbf{I}(X) = \mathcal{S}(X_a)$ minus X . This not only gives a concrete construction of such an adequate domain of limits, but also shows that we do not have much freedom in defining one: any other one must contain $\mathcal{S}(X_a)$.

The definition of ADLs above is slightly awkward. Let us simplify it.

Requirement (L_2) in [Geeraerts et al., 2006] only serves to ensure that all closed subsets of $L \cup X$ can be represented as $\downarrow_{L \cup X} E$ for some finite subset E : the closed subset $L \cup X$ itself is then exactly $\downarrow_{L \cup X} \{\top\}$. However, (L_2) is unnecessary for this, since $L \cup X$ already equals $\downarrow_{L \cup X} E$ by (L_3), where E is the finite subset of $L \cup X$ such that $\hat{\gamma}(E) = L \cup X$ as ensured by (L_4). We will not need (L_2) either in our own subsequent work [Finkel and Goubault-Larrecq, 2012], and shall call *weak adequate domain of limits* (*WADL*) any triple (L, \preceq, γ) satisfying (L_1), (L_3), and (L_4).

Even so, this definition remains awkward. First, the real space of interest is not L , but $Z = L \cup X$; L can always be recovered as $Z \setminus X$. Then $\gamma(z)$ should be downwards-closed for every $z \in Z$, i.e., it should be closed in X_a . The space of downwards-closed subsets is the Hoare powerdomain $\mathcal{H}_V(X_a)_\perp$. As a consequence, (L_1) can be expressed more succinctly by requiring that γ be a map from Z to $\mathcal{H}_V(X_a)_\perp$, and that every subset of the form $\downarrow_X x$, $x \in X$, is obtained as $\gamma(x)$. Requirement (L_3) means that γ is a quasi-order embedding. In other words, the elements z of Z can be thought as *syntax* for particular elements $\gamma(z)$ of $\mathcal{H}_V(X_a)_\perp$, and we define \preceq on syntax by $z \preceq z'$ iff $\gamma(z) \subseteq \gamma(z')$. So we may safely omit \preceq from the definition, and remove requirement (L_3).

The only important requirement is (L_4), which states that every downwards-closed subset of X should be describable as a finite union of representable subsets, i.e., of elements of the form $\gamma(z)$, $z \in Z$. (L_1) also requires all elements of the form $\downarrow_X x$, $x \in X$, to be representable. However, this is a consequence of (L_4): $\downarrow_X x$ is a finite union of representable subsets $\gamma(z_1), \dots, \gamma(z_n)$; then $x \in \gamma(z_i)$ for some i , $1 \leq i \leq n$, from which we deduce that $\gamma(z_i) = \downarrow_X x$.

We therefore arrive at the following definition.

Definition 3.1 (ADL, WADL) *Let X be a quasi-ordered set. A weak adequate domain of limits, or WADL, on X is a pair (Z, γ) of a set Z and a map $\gamma: Z \rightarrow \mathcal{H}_V(X_a)_\perp$ (the representation map) such that every downwards-closed subset of X is a finite union of representables. A representable subset of X is by definition one of the form $\gamma(z)$ for some $z \in Z$.*

(Z, γ) is an adequate domain of limits (ADL) iff, additionally, the whole set X is representable.

In any case, the limit points of Z are those $z \in Z$ such that $\gamma(z)$ is not of the form $\downarrow_X x$, $x \in X$.

We check the formal relationship with Geeraerts *et al.*'s conditions. The easy proof is left to the reader.

Lemma 3.2 *Let X be a quasi-ordered set.*

If (L, \sqsubseteq, γ) satisfies (L_1) , (L_3) and (L_4) , then $(L \cup X, \gamma)$ is a WADL.

If (L, \sqsubseteq, γ) satisfies (L_1) , (L_2) , (L_3) and (L_4) , then $(L \cup X, \gamma)$ is an ADL.

Conversely, if (Z, γ) is a WADL (resp., ADL) on X , then $(L, \sqsubseteq, \gamma')$ satisfies (L_1) , (resp., and (L_2)), (L_3) and (L_4) where L is the set of limit points of Z , \sqsubseteq is defined by $z \sqsubseteq z'$ iff $\gamma'(z) \subseteq \gamma'(z')$, and γ' is defined by $\gamma'(z) = \gamma(z)$ if $z \in L$, $\gamma'(x) = \downarrow_X x$ if $x \in X$.

Definition 3.1 displays a tension between mathematical practice and computer science needs. That every downwards-closed subset of X be a finite union of representables $\gamma(z_1), \dots, \gamma(z_n)$ means that we can represent any downwards-closed set by finitely many pieces of information z_1, \dots, z_n . However, from a computer science perspective, we have not (yet) put any computability conditions on WADL. We repair this now.

Definition 3.3 (Effective WADL) *A WADL (Z, γ) on X is an effective WADL iff the relation \sqsubseteq on Z , defined by $z \sqsubseteq z'$ iff $\gamma(z) \subseteq \gamma(z')$, is decidable.*

This naturally assumes that Z is a domain of objects representable on a computer, e.g., a word, or a natural number.

From a mathematical standpoint, on the other hand, one usually reasons up to order quotients and order isomorphisms. Then γ and Z are useless in Definition 3.1, and the only relevant part of a WADL is the collection of representable subsets. I.e., a WADL is, up to these details, a collection of downwards-closed subsets, the representable subsets, such that every downwards-closed subset is a finite union of representables.

We can then bound precisely all WADLs between two well-known spaces of downwards-closed subsets (i.e., closed in X_a).

Lemma 3.4 *Let X be a poset, and (Z, γ) be a WADL on X . The set $\gamma[Z]$ of representable subsets is such that:*

$$\mathcal{S}(X_a) \subseteq \gamma[Z] \subseteq \mathcal{H}_V(X_a)_\perp$$

Proof. We must show that $\mathcal{S}(X_a) \subseteq \gamma[Z]$, the other inclusion being by definition. Let $C \in \mathcal{S}(X_a)$, i.e., assume C is irreducible closed. C must be a finite union of representables $\bigcup_{i=1}^n \gamma(z_i)$ by the definition of WADLs. So $C \subseteq \gamma(z_i)$ for some i , $1 \leq i \leq n$, by irreducibility (and since each $\gamma(z_i)$ is closed in X_a). It follows that $C = \gamma(z_i)$, hence $C \in \gamma[Z]$. \square

So, up to order quotients and order isomorphisms, there cannot be any WADL smaller than the sobrification $\mathcal{S}(X_a)$. We shall see later that the latter is effective in a large number of practical cases.

Naturally, the statement of Lemma 3.4 does not require any topology. Purely order-theoretically, Lemma 3.4 states that the collection of representable subsets must lie between the collection of ideals ($\mathbf{I}(X) = \mathcal{S}(X_a)$) and the collection of all downwards-closed subsets ($\mathcal{H}_V(X_a)_\perp$).

The interest in using topology is in the proof of Lemma 3.4: the point is that the key notion is irreducibility, a topological notion. In turn, these notions and proofs will generalize to the topological, Noetherian case with no effort later.

When X is a wqo, the ideal completion $\mathbf{I}(X) = \mathcal{S}(X_a)$ is not just a lower bound below any WADL, it is itself a WADL. This follows from the following more general topological statement.

Proposition 3.5 *Let X be a Noetherian space. Then $\mathcal{S}(X)$ is the least collection \mathcal{C} of closed subsets of X such that every closed subset of X can be expressed as a finite union of elements of \mathcal{C} .*

Proof. First, if \mathcal{C} is as above, then $\mathcal{S}(X) \subseteq \mathcal{C}$. The proof is as in Lemma 3.4, which is in fact a topological proof: every element C of $\mathcal{S}(X_a)$ must be written as a finite union of elements of \mathcal{C} , and by irreducibility it must equal one of them.

Conversely, we need to show that every closed subset of X is a finite union of irreducible closed subsets, provided that X is Noetherian. This is a well-known fundamental result, and occurs as part of [Goubault-Larrecq, 2013, Theorem 9.7.12]. We give an elementary proof of it in Lemma 3.6 below, for the sake of completeness. \square

Lemma 3.6 *In a Noetherian space, every closed subset is a finite union of irreducibles.*

Proof. By taking complements, in a Noetherian space X every properly descending chain $F_0 \supsetneq F_1 \supsetneq \dots \supsetneq F_n \supsetneq \dots$ of closed subsets must be finite, in other words $\mathcal{H}_V(X_a)$, ordered by inclusion, is well-founded. Imagine there were a closed subset C that cannot be written as a finite union of irreducibles. By well-foundedness, we can choose C minimal. C is not empty, since the empty set can be written as a finite union of irreducibles, namely none. C cannot be irreducible either, so there are two closed subsets F_1 and F_2 such that $C \subseteq F_1 \cup F_2$, but $C \not\subseteq F_1$ and $C \not\subseteq F_2$. Because of the latter, $C \cap F_1$ and $C \cap F_2$ are strictly smaller than C . By the minimality of C , $C \cap F_1$ and $C \cap F_2$ can be written as finite unions of irreducibles, so $C = (C \cap F_1) \cup (C \cap F_2)$ is also a finite union of irreducibles: contradiction. \square

As a special case, we obtain that every downwards-closed subset of a wqo X is a finite union of ideals. This can also be deduced from the observation by Erdős and Tarski [Erdős and Tarski, 1943] that a poset has no infinite antichain if and only if every downwards-closed subset is a finite union of ideals.

Remark 3.7 *Generalizing the above cited result of Erdős and Tarski to the topological setting, we have the following: a topological space has no infinite discrete subspace if and only if every closed subset is a finite union of irreducibles [Goubault-Larrecq, 2019].*

Proposition 3.5, once toned down to wqos, translates to the following.

Corollary 3.8 (Least WADL) *Let X be a wqo. The ideal completion $\mathbf{I}(X) = \mathcal{S}(X_a)$ is the least WADL, in the sense that:*

- 1) *for any WADL (Z, γ) on X , every element of $\mathcal{S}(X_a)$ is representable;*
- 2) *$(\mathcal{S}(X_a), i)$ is itself a WADL, where i is the canonical injection of $\mathcal{S}(X_a)$ into $\mathcal{H}_V(X_a)_\perp$.*

In other words, up to the coding function γ , there is a unique *minimal* WADL on any given wqo X . We contend that $\mathcal{S}(X_a)$ is, in all practical cases, the sole WADL worth considering, and will in particular be effective.

Our treatment so far uses topology for no particular good reason apart from mathematical elegance. Our presentation, however, lends itself to the following natural topological extension of WADLs. We have claimed that the additional generality obtained by shifting focus from wqos to the larger class of Noetherian spaces was useful in [Goubault-Larrecq, 2010]. Notably, the class of polynomial concurrent programs introduced there is naturally seen as a *topological* WSTS, that is, as a pair (X, \rightarrow) where X is Noetherian space (instead of a wqo) and \rightarrow is a lower semi-continuous binary relation—this is the natural generalization of WSTS to a topological setting. Using Noetherianness, and algorithms and proof arguments that are variants of WSTS arguments, it was shown in that same paper that the reachability of sets of states defined by so-called forbidden patterns is decidable for polynomial concurrent programs. Note that the latter are *not* WSTS. In that context, studying topological WADLs instead of WADLs is the logical next step.

Definition 3.9 *Let X be a topological space. A topological WADL on X is a pair (Z, γ) of a set Z and a map $\gamma: Z \rightarrow \mathcal{H}_V(X)_\perp$ (the representation map) such that every closed subset of X is a finite union of representables. A representable subset of X is by definition one of the form $\gamma(z)$ for some $z \in Z$.*

So WADLs are topological WADLs, in the special case where X comes with the Alexandroff topology of some quasi-ordering. We have just seen (Proposition 3.5) that, when X is Noetherian, $\mathcal{S}(X)$ is the least topological WADL. We state it as follows.

Proposition 3.10 (Least Topological WADL) *Let X be a Noetherian space. The sobrification $\mathcal{S}(X)$ is the least topological WADL, in the sense that:*

- 1) *for any topological WADL (Z, γ) on X , every element of $\mathcal{S}(X)$ is representable;*
- 2) *$(\mathcal{S}(X), i)$ is itself a topological WADL, where i is the canonical injection of $\mathcal{S}(X)$ into $\mathcal{H}_V(X)_\perp$.*

$D ::=$	A	(finite poset; Theorem 4.3)	
	\mathbb{N}	(natural numbers; Theorem 4.4)	
	$D_1 \times D_2 \times \cdots \times D_n$	(product; Theorem 4.5)	
	$D_1 + D_2 + \cdots + D_n$	(coproduct; Theorem 4.6)	
	$\mathcal{S}(D)$	(sobrification; Theorem 4.7)	*
	$\mathbb{P}(D)$	(powerset; Theorem 4.11)	*
	$\mathbb{P}^*(D)$	(non-empty powerset; Theorem 4.11)	*
	$\mathcal{H}_V(D)$	(Hoare powerdomain; Theorem 4.8)	*
	$\mathcal{H}_V(D)_\perp$	(lifted Hoare powerdomain; Theorem 4.8)	*
	$\text{Spec}(R)$	(spectrum of ring R ; Proposition 5.1)	*
	D^*	(finite words; Theorem 6.15)	
	D^\otimes	(finite multisets; Theorem 7.7)	
	$\triangleright_{n=1}^{+\infty} D_n$	(words, prefix; Theorem 8.10)	*
	$\mathcal{T}(D)$	(finite trees; Theorem 10.36)	

Figure 2: An algebra of Noetherian datatypes

4 S-Representations

We shall devote the rest of this paper to describe completions $\widehat{D} = \mathcal{S}(D)$ for those datatypes of Figure 2. As we shall see, these datatypes include most of the datatypes encountered in the literature on WSTS (e.g, Petri nets and more generally counter machines, lossy channel systems, data nets), and contain several new ones.

All the datatypes in this Figure are Noetherian spaces, as can be gathered from Section 9.7 of [Goubault-Larrecq, 2013]. We also state the relevant theorem in each case. Stars indicate constructs that, while preserving Noetherianness, do not preserve well-quasi-orderedness. What the values of these types are, what their topologies are (and associated specialization quasi-orderings) will also be made precise in each corresponding section.

The completion process is modular: the completion \widehat{D} of a type D , built from D_1, \dots, D_n , will be defined as a function of $\widehat{D}_1, \dots, \widehat{D}_n$. In each case, we shall show that if $\widehat{D}_1, \dots, \widehat{D}_n$ are effective, then so is \widehat{D} .

As a result, all the datatypes defined in Figure 2 will be effective. This is important: Definition 3.3, applied to the WADL $\widehat{X} = \mathcal{S}(X_a)$ (when X is wqo), requires us to decide the ordering (i.e., inclusion) on $\mathcal{S}(X)$. We shall require—and obtain—more: we shall be able to compute finite intersections of closed subsets (i.e., downwards-closed subsets in wqos) as well.

We consider topological WADLs—e.g., the starred rows in Figure 2—for added generality, but also because the topological approach, relying on the notion of irreducibility, provides a unifying perspective on the matter. This leads to the following notion of an effective, finite representation of irreducible closed subsets. The closed, not necessarily irreducible, subsets are all finite unions of irreducibles (Lemma 3.6) and can therefore be represented as finite sets of codes. Below, this is how we represent the

closed sets X (item D) and $\llbracket a \rrbracket \cap \llbracket b \rrbracket$ (item E).

Definition 4.1 (S-Representation) *Let X be a topological space. An S-representation of X is a tuple $(S, \llbracket _ \rrbracket, \sqsubseteq, \tau, \wedge)$ where:*

- (A) S is a recursively enumerable set of so-called codes (of irreducible closed subsets);
- (B) $\llbracket _ \rrbracket$ is a surjective map from S to $\mathcal{S}(X)$;
- (C) \sqsubseteq is a decidable relation such that, for all codes $a, b \in S$, $a \sqsubseteq b$ iff $\llbracket a \rrbracket \subseteq \llbracket b \rrbracket$;
- (D) τ is a finite subset of S , such that $X = \bigcup_{a \in \tau} \llbracket a \rrbracket$;
- (E) \wedge is a computable map from $S \times S$ to the collection $\mathbb{P}_{fm}(S)$ of finite subsets of S (and we write $a \wedge b$ for $\wedge(a, b)$) such that $\llbracket a \rrbracket \cap \llbracket b \rrbracket = \bigcup_{c \in a \wedge b} \llbracket c \rrbracket$.

We call \wedge the intersection map.

The idea is that codes represent irreducible closed subsets, through the semantic function $\llbracket _ \rrbracket$, that \sqsubseteq implements inclusion, τ denotes the whole set X , and \wedge implements intersection.

We justify this now, in a more precise way. We represent closed subsets F through finite sets $\{a_1, \dots, a_m\}$ of codes. The denotation of such a finite set is the union $\bigcup_{i=1}^m \llbracket a_i \rrbracket$. Since $\llbracket _ \rrbracket$ is surjective, and using Lemma 3.6, every closed subset of a Noetherian space X can be represented this way. This is in particular true for the whole set X (item D) and the intersection $\llbracket a \rrbracket \cap \llbracket b \rrbracket$ in item E.

While \sqsubseteq allows us to test two (codes of) irreducible closed subsets for inclusion, one can extend the inclusion test to arbitrary closed subsets: this is what we show now.

Lemma 4.2 *Given irreducible closed subsets $C_1, \dots, C_m, C'_1, \dots, C'_n$ of a topological space X , the following are equivalent:*

- $C_1 \cup \dots \cup C_m \subseteq C'_1 \cup \dots \cup C'_n$;
- $\{C_1, \dots, C_m\} \sqsubseteq^b \{C'_1, \dots, C'_n\}$, that is, for every i ($1 \leq i \leq m$), there is a j ($1 \leq j \leq n$) with $C_i \subseteq C'_j$.

Proof. $C_1 \cup \dots \cup C_m \subseteq C'_1 \cup \dots \cup C'_n$ if and only if for every i , C_i is included in $C'_1 \cup \dots \cup C'_n$. Since C_i is irreducible, the latter is equivalent to the existence of j such that $C_i \subseteq C'_j$. \square

In general, \leq^b is the *Hoare quasi-ordering* on subsets, also called the *domination quasi-ordering*: $A \leq^b B$ iff for every $a \in A$, there is a $b \in B$ such that $a \subseteq b$. We will use \leq^b for various quasi-orderings \leq , and will accordingly use the notations \sqsubseteq^b as above, or \leq^b later.

Given an S-representation, we can then test two closed sets for inclusion: given two finite sets $\{a_1, \dots, a_m\}$ and $\{b_1, \dots, b_n\}$ of codes, $\bigcup_{i=1}^m \llbracket a_i \rrbracket$ is included in $\bigcup_{j=1}^n \llbracket b_j \rrbracket$ iff for every i , there is a j such that $a_i \sqsubseteq b_j$.

Finite intersections are computable, too, using \wedge : the intersection of two closed sets represented by finite sets $\{a_1, \dots, a_m\}$ and $\{b_1, \dots, b_n\}$ of codes is $\bigcup_{i=1}^m \llbracket a_i \rrbracket \cap \bigcup_{j=1}^n \llbracket b_j \rrbracket = \bigcup_{i,j} \llbracket a_i \rrbracket \cap \llbracket b_j \rrbracket$, and is therefore represented by the finite set $\bigcup_{i,j} a_i \wedge b_j$. Finite unions are, of course, easily computable as well.

Our purpose is to show that every space X that occurs as the space of values of some type D in Figure 2 has an S-representation. In each case, we will actually define an S-representation of D as a function of given S-representation of its constituent datatypes, and we shall use a uniform presentation: each result will be given in the form of a proposition of the following shape.

“Proposition XXX (S-Representation, datatype D) Let X_i be Noetherian spaces, and $(S_i, \llbracket _ \rrbracket_i, \leq_i, \tau_i, \wedge_i)$ be an S-representation of D_i for each i . Then $(S', \llbracket _ \rrbracket', \leq', \tau', \wedge')$ is an S-representation of D , where:

- (A) S' is ...
- (B) $\llbracket \cdot \cdot \rrbracket'$ is defined as ...
- (C) \leq' is defined as ...
- (D) τ' is defined as ...
- (E) \wedge' is defined by $a' \wedge' b' = \dots$ ”

We start with the easiest cases. The more difficult cases will be dealt with in separate sections. Our first instance is trivial: in a finite quasi-ordered set A , irreducible closed subsets, that is, ideals, are all of the form $\downarrow x, x \in A$, so $\widehat{A} = \mathcal{S}(X_a) = \mathbf{I}(X)$ is isomorphic to A .

Theorem 4.3 (S-Representation, Finite Quasi-Orders) *Let A be any finite quasi-ordered set. An S-representation of A is $(S, \llbracket _ \rrbracket, \leq, \tau, \wedge)$ where:*

- (A) S is A itself,
- (B) $\llbracket _ \rrbracket$ is the identity map,
- (C) \leq is the given ordering on A ,
- (D) τ is the set of maximal elements of A ,
- (E) $a \wedge b$ is the set of maximal lower bounds of a and b .

All this is computable, by just maintaining all needed information in tables. As a particular case, one finds the finite sets: these are the posets whose ordering is equality. In particular, the above provides an S-representation for finite sets Q of control states of various kinds of machines. In this case, $\tau = Q, a \wedge b = \{a\}$ if $a = b$, and $a \wedge b = \emptyset$ otherwise.

The next case is an easy exercise.

Theorem 4.4 (S-Representation, \mathbb{N}) *An S-representation of \mathbb{N} is $(S, \llbracket _ \rrbracket, \leq, \tau, \wedge)$ where:*

- (A) $S = \mathbb{N}_\omega$,

(B) $\llbracket _ \rrbracket$ maps n to $\downarrow n$ and ω to \mathbb{N} ,

(C) \leq is the usual ordering on \mathbb{N}_ω ,

(D) $\tau = \{\omega\}$,

(E) $m \wedge n = \{\min(m, n)\}$.

Theorem 4.5 (S-Representation, Products) *Let X_1, \dots, X_n be n Noetherian spaces, $X = X_1 \times \dots \times X_n$, and $(S_i, \llbracket _ \rrbracket_i, \leq_i, \tau_i, \wedge_i)$ an S-representation of X_i for each i , $1 \leq i \leq n$. Then $(S, \llbracket _ \rrbracket, \leq, \tau, \wedge)$ is an S-representation of X , where:*

(A) $S = S_1 \times \dots \times S_n$;

(B) $\llbracket (a_1, \dots, a_n) \rrbracket = \llbracket a_1 \rrbracket \times \dots \times \llbracket a_n \rrbracket$;

(C) $(a_1, \dots, a_n) \leq (b_1, \dots, b_n)$ iff $a_1 \leq_1 b_1$ and ... and $a_n \leq_n b_n$;

(D) $\tau = \tau_1 \times \dots \times \tau_n$;

(E) $(a_1, \dots, a_n) \wedge (b_1, \dots, b_n) = (a_1 \wedge_1 b_1) \times \dots \times (a_n \wedge_n b_n)$.

Proof. The elements of $S(X)$ are the products $C_1 \times \dots \times C_n$ of irreducible closed subsets C_1 of X_1, \dots, C_n of X_n (see Lemma A.2 in the Appendix), which justifies items A and B: we represent $C_1 \times \dots \times C_n$ as the n -tuple of codes for C_1, C_2, \dots, C_n . The if direction of item C follows from the fact that product is monotonic with respect to inclusion. Conversely, if $\prod_{i=1}^n \llbracket a_i \rrbracket \subseteq \prod_{i=1}^n \llbracket b_i \rrbracket$, then $\llbracket a_i \rrbracket \subseteq \llbracket b_i \rrbracket$ for every i : since $\llbracket a_j \rrbracket$ is non-empty, we can pick an element x_j from $\llbracket a_j \rrbracket$ for every $j \neq i$; then, for every $x \in \llbracket a_i \rrbracket$, the tuple $(x_1, \dots, x_{i-1}, x, x_{i+1}, \dots, x_n)$ is in $\prod_{i=1}^n \llbracket a_i \rrbracket$, hence in $\prod_{i=1}^n \llbracket b_i \rrbracket$, showing that x is in $\llbracket b_i \rrbracket$. Items D and E are clear. \square

So, for example, an S-representation for \mathbb{N}^k , the datatype of configurations of Petri nets, and more generally, of counter machines, is as expected: $S = \mathbb{N}_\omega^k, (m_1, \dots, m_k) \leq (n_1, \dots, n_k)$ iff $m_i \leq n_i$ for every $i, 1 \leq i \leq k, \tau = \{(\omega, \dots, \omega)\}$, and $(m_1, \dots, m_k) \wedge (n_1, \dots, n_k) = \{\min(m_1, n_1), \dots, \min(m_k, n_k)\}$.

Theorem 4.6 (S-Representation, Coproducts) *Let X_1, \dots, X_n be n Noetherian spaces, and $X = X_1 + \dots + X_n$. Then $S(X)$ is homeomorphic to $S(X_1) + \dots + S(X_n)$.*

Let $(S_i, \llbracket _ \rrbracket_i, \leq_i, \tau_i, \wedge_i)$ be an S-representation of X_i for each $i, 1 \leq i \leq n$. Then $(S, \llbracket _ \rrbracket, \leq, \tau, \wedge)$ is an S-representation of $X = X_1 + \dots + X_n$, where:

(A) $S = \{(i, a) \mid 1 \leq i \leq n, a \in S_i\}$;

(B) $\llbracket (i, a) \rrbracket = \llbracket a \rrbracket_i$ (up to the homeomorphism between $S(X)$ and $S(X_1) + \dots + S(X_n)$);

(C) $(i, a) \leq (j, b)$ iff $i = j$ and $a \leq_i b$;

(D) $\tau = \bigcup_{i=1}^n \{i\} \times \tau_i$;

(E) $(i, a) \wedge (j, b) = \emptyset$ if $i \neq j$, $(i, a) \wedge (i, b) = \{(i, c) \mid c \in a \wedge_i b\}$.

Proof. For the first part, see [Goubault-Larrecq, 2013, Fact 8.4.3], which states that \mathcal{S} commutes with coproducts (in fact with all colimits, since \mathcal{S} is a left adjoint). The rest is clear. \square

A trivial case of \mathcal{S} -representation is provided by sobrifications themselves, because $\mathcal{S}(\mathcal{S}(X))$ is canonically isomorphic to $\mathcal{S}(X)$. Indeed, $Y = \mathcal{S}(X)$ is sober, and for every sober space Y , $\eta^{\mathcal{S}}$ is an isomorphism between Y and $\mathcal{S}(Y)$ [Goubault-Larrecq, 2013, Fact 8.2.5]. It follows that any \mathcal{S} -representation $(S, \llbracket _ \rrbracket, \preceq, \tau, \wedge)$ for X yields an \mathcal{S} -representation for $\mathcal{S}(X)$ with the same set S of codes and the same operations \preceq, τ , and \wedge , namely $(S, \eta_X^{\mathcal{S}} \circ \llbracket _ \rrbracket, \preceq, \tau, \wedge)$:

Theorem 4.7 (S-Representation, Sobrifications) *Let X be a Noetherian space, and $X' = \mathcal{S}(X)$. Let $(S, \llbracket _ \rrbracket, \preceq, \tau, \wedge)$ be an \mathcal{S} -representation of X . Then $(S', \llbracket _ \rrbracket', \preceq', \tau', \wedge')$ is an \mathcal{S} -representation of X' where:*

- (A) $S' = S$;
- (B) for every $a \in S$, $\llbracket a \rrbracket' = \downarrow_{X'} \llbracket a \rrbracket$;
- (C) $a \preceq' b$ iff $a \preceq b$;
- (D) $\tau' = \tau$;
- (E) $a \wedge' b = a \wedge b$.

Let us deal with the Hoare powerdomain $\mathcal{H}_V(X)$ of X and its lifted version.

Theorem 4.8 (S-Representation, Hoare Powerdomains) *Let X be a Noetherian space, and $X' = \mathcal{H}_V(X)_\perp$ (resp., $X' = \mathcal{H}_V(X)$). Let $(S, \llbracket _ \rrbracket, \preceq, \tau, \wedge)$ be an \mathcal{S} -representation of X . Then $(S', \llbracket _ \rrbracket', \preceq', \tau', \wedge')$ is an \mathcal{S} -representation of X' where:*

- (A) $S' = \mathbb{P}_{fm}(S)$ (resp., $S' = \mathbb{P}_{fm}^*(S)$);
- (B) for every $a' \in S'$, $\llbracket a' \rrbracket' = \downarrow_{X'} \{\bigcup_{a \in a'} \llbracket a \rrbracket\}$;
- (C) $a' \preceq' b'$ iff $a' \preceq^b b'$, where $a' \preceq^b b'$ iff for every $a \in a'$, there is a $b \in b'$ such that $a \preceq b$ (compare Lemma 4.2);
- (D) $\tau' = \{\tau\}$;
- (E) $a' \wedge' b' = \{\bigcup_{a \in a', b \in b'} (a \wedge b)\}$.

Proof. By Lemma 3.6, every element F of $\mathcal{H}_V(X)_\perp$ is a finite union of irreducible closed subsets, which are each of the form $\llbracket a \rrbracket$ with $a \in S$ by assumption, since $\llbracket _ \rrbracket$ is surjective. So $\llbracket _ \rrbracket'$ is surjective.

Next, $a' \preceq' b'$ iff $\llbracket a' \rrbracket' \subseteq \llbracket b' \rrbracket'$, iff $\bigcup_{a \in a'} \llbracket a \rrbracket \subseteq \bigcup_{b \in b'} \llbracket b \rrbracket$, iff $a' \preceq^b b'$, by Lemma 4.2.

We must check that $X' = \bigcup_{a' \in \tau'} \llbracket a' \rrbracket'$. The right-hand side is $\llbracket \tau \rrbracket' = \downarrow_{X'} \{\bigcup_{a \in \tau} \llbracket a \rrbracket\} = \downarrow_{X'} \{X\} = X'$.

Finally, let us compute $\llbracket a' \rrbracket' \cap \llbracket b' \rrbracket'$. This is $\downarrow_{X'} \{\bigcup_{a \in a'} \llbracket a \rrbracket\} \cap \downarrow_{X'} \{\bigcup_{b \in b'} \llbracket b \rrbracket\} = \downarrow_{X'} \{\bigcup_{a \in a'} \llbracket a \rrbracket \cap \bigcup_{b \in b'} \llbracket b \rrbracket\} = \downarrow_{X'} \{\bigcup_{a \in a', b \in b'} (\llbracket a \rrbracket \cap \llbracket b \rrbracket)\} = \downarrow_{X'} \{\bigcup_{a \in a', b \in b'} \bigcup_{c \in a \wedge b} \llbracket c \rrbracket\} = \downarrow_{X'} \{\bigcup_{c \in \bigcup_{a \in a', b \in b'} (a \wedge b)} \llbracket c \rrbracket\} = \llbracket \bigcup_{a \in a', b \in b'} (a \wedge b) \rrbracket'$. \square

Let $\mathbb{P}^*(X)$ be the set of non-empty subsets of X . We topologize $X' = \mathbb{P}(X)$ (resp., $X' = \mathbb{P}^*(X)$) by the lower Vietoris topology, generated by the subbasic opens $\diamond U = \{A \in X' \mid A \cap U \neq \emptyset\}$, where U ranges over the open subsets of X . (Although there is a risk of confusion with the lower Vietoris topology on $\mathcal{H}_V(X)_\perp$, we shall see that the two are strongly tied.)

It is worth to point out that A is below B in the specialization quasi-ordering of those spaces if and only if $cl(A) \subseteq cl(B)$. This is well-known, and appears for example as part of Proposition 7.3 of [Goubault-Larrecq, 2007]. We include the proof for completeness.

Lemma 4.9 *The specialization quasi-ordering of $\mathbb{P}(X)$, resp. $\mathbb{P}^*(X)$, is inclusion of closures.*

Proof. Let us temporarily write \preceq for that specialization quasi-ordering.

If $A \preceq B$ in $\mathbb{P}(X)$ (resp., $\mathbb{P}^*(X)$), then consider the (open) complement U of $cl(B)$. Since $A \preceq B$, if $A \in \diamond U$ then $B \in \diamond U$. However, B does not intersect U , since U is the complement of $cl(B)$, so B is not in $\diamond U$. It follows that A is not in $\diamond U$ either. This means that A does not intersect U , and therefore that it is included in its complement, $cl(B)$. Since $cl(B)$ is closed, contains A , and $cl(A)$ is by definition the smallest closed subset of X containing A , $cl(A)$ is included in $cl(B)$.

In the converse direction, we use the standard fact that, for an open subset U of X , $cl(A)$ intersects U if and only if A intersects U . Let us assume that $cl(A) \subseteq cl(B)$. Let $\mathcal{U} = \bigcup_{i \in I} \bigcap_{j \in J_i} \diamond U_{ij}$ (where each J_i is finite) be any open subset of $\mathbb{P}(X)$ (resp., $\mathbb{P}^*(X)$) containing A . Then, for some $i \in I$, A intersects U_{ij} for every $j \in J_i$. The larger set $cl(B)$ must then intersect each U_{ij} as well. Hence (see standard fact) B also intersects each U_{ij} . It follows that B is in \mathcal{U} . Since \mathcal{U} is arbitrary, $A \preceq B$. \square

When X is equipped with the Alexandroff topology of a quasi-ordering \leq , $cl(A) = \downarrow A$ and $cl(B) = \downarrow B$, and therefore the specialization quasi-ordering of $\mathbb{P}(X)$ and $\mathbb{P}^*(X)$ is the familiar Hoare quasi-ordering \leq^b in that case.

Even when \leq is wqo, \leq^b fails to be wqo in general. However, $\mathbb{P}(X)$ and $\mathbb{P}^*(X)$ are Noetherian for X Noetherian, as first remarked in [Goubault-Larrecq, 2007]. This stems from the following result, and the fact that sobrifications of Noetherian spaces are Noetherian.

Lemma 4.10 *For a topological space X , $\mathcal{H}_V(X)_\perp$ is the sobrification of $\mathbb{P}(X)$, and $\mathcal{H}_V(X)$ is the sobrification of $\mathbb{P}^*(X)$, up to homeomorphism.*

Proof. We deal with the first claim, by exhibiting a homeomorphism \square between $\mathcal{H}_V(X)_\perp$ and $\mathcal{S}(\mathbb{P}(X))$. To distinguish closure in X and closure in $\mathbb{P}(X)$, let us write cl_X for the former and $cl_{\mathbb{P}(X)}$ for the latter.

For every closed subset C of X , $cl_{\mathbb{P}(X)}(\{C\})$ is the closure of the point C in $\mathbb{P}(X)$, and that is equal to the downward closure of $\{C\}$ with respect to the specialization quasi-ordering of $\mathbb{P}(X)$. By Lemma 4.9, this is the set of subsets A of X such that $cl_X(A) \subseteq cl_X(C)$. Since $cl_X(C) = C$, and since $cl_X(A) \subseteq C$ is equivalent to $A \subseteq C$ (since C is closed in X), $cl_{\mathbb{P}(X)}(\{C\})$ is therefore just the set $\square C$ of subsets of C .

The notation $\square C$ is justified by the fact that it is the complement of $\diamond U$ where U is the complement of C . (Admittedly, we could also have written it as $\downarrow C$.) Since

$\square C$ is equal to $cl_{\mathbb{P}(X)}(\{C\})$, it is in particular irreducible closed. This defines a map $\square: \mathcal{H}_V(X)_\perp \rightarrow \mathcal{S}(\mathbb{P}(X))$.

Conversely, let \mathcal{I} be an irreducible closed subset of $\mathbb{P}(X)$. As a closed set, we can write it as $\bigcap_{i \in I} \bigcup_{j \in J_i} \square C_{ij}$, where each J_i is finite and each C_{ij} is closed. For each $i \in I$, $\mathcal{I} \subseteq \bigcup_{j \in J_i} \square C_{ij}$, and since \mathcal{I} is irreducible, there is a $j_i \in J_i$ such that $\mathcal{I} \subseteq \square C_{ij_i}$. Therefore $\mathcal{I} \subseteq \bigcap_{i \in I} \square C_{ij_i}$, and as the right-hand side is clearly included in \mathcal{I} , this inequality is in fact an equality. It is easy to see that \square commutes with arbitrary intersections. As a consequence, \mathcal{I} is of the form $\square C$, where $C = \bigcap_{i \in I} C_{ij_i}$. It follows that \square is surjective.

Note furthermore that C is unique: if $\mathcal{I} = \square C$, then C is necessarily the largest closed set that is an element of \mathcal{I} . Hence the map $\square: \mathcal{H}_V(X)_\perp \rightarrow \mathcal{S}(\mathbb{P}(X))$ is bijective.

To show that a map is continuous, it is enough to show that the inverse image of a subbasic open is open. A subbase of opens of $\mathcal{S}(\mathbb{P}(X))$ consists of the sets of the form $\diamond \diamond U$, U open in X , because the outer \diamond commutes with all unions and finite intersections. (Both \diamond and \square commute with unions; we let the reader check that the outer \diamond also commutes with finite intersections, as a consequence of irreducibility.)

To show that \square is continuous, it therefore suffices to show that $\square^{-1}(\diamond \diamond U)$ is open for every open subset U of X . For every C in $\mathcal{H}_V(X)_\perp$, $C \in \square^{-1}(\diamond \diamond U)$ if and only if $\square C$ intersects $\diamond \diamond U$, if and only if some closed subset C' of C intersects U , if and only if C itself intersects U . So $\square^{-1}(\diamond \diamond U) = \diamond U$, showing that \square is continuous.

Conversely, the inverse image of $\diamond U$ by the inverse map \square^{-1} is $\diamond \diamond U$, showing that \square^{-1} , too, is continuous. Therefore \square is a homeomorphism. \square

This not only shows that $\mathbb{P}(X)$ and $\mathbb{P}^*(X)$ are Noetherian for X Noetherian, but also that they have the same sobrifications as $\mathcal{H}_V(X)_\perp$ and $\mathcal{H}_V(X)$ respectively, hence can be given the same S-representations:

Theorem 4.11 (S-Representation, Powersets) *Let X be a Noetherian space, and $X' = \mathbb{P}(X)$ (resp., $X' = \mathbb{P}^*(X)$). Let $(S, \llbracket _ \rrbracket, \leq, \tau, \wedge)$ be an S-representation of X . Then $(S', \llbracket _ \rrbracket', \leq', \tau', \wedge')$ is an S-representation of X' where:*

- (A) $S' = \mathbb{P}_{fin}(S)$ (resp., $S' = \mathbb{P}_{fin}^*(S)$);
- (B) for every $a' \in S'$, $\llbracket a' \rrbracket' = \downarrow_{X'} \{ \bigcup_{a \in a'} \llbracket a \rrbracket \}$;
- (C) $a' \leq' b'$ iff $a' \leq^b b'$;
- (D) $\tau' = \{ \tau \}$;
- (E) $a' \wedge' b' = \{ \bigcup_{a \in a', b \in b'} (a \wedge b) \}$.

5 Completing Ring Ideals

The primary example of Noetherian spaces, historically, are the spectra of Noetherian rings. Mentioning them is therefore mandatory. The non-algebraically inclined reader is invited to proceed to finite words (Section 6).

Let R be a commutative ring (with unit). Recall that an *ideal* I is any additive subgroup of R such that for any $r \in I, r' \in R$, the product rr' is in I . A *prime ideal* p is an ideal that does not contain the multiplicative unit 1 of R (equivalently, which is different from the whole of R), and such that whenever $rr' \in p$, then r or r' is in p . The *spectrum* $\text{Spec}(R)$ of R is the set of all prime ideals of R . It is equipped with the *Zariski topology*, whose closed subsets are $F_I = \{p \in \text{Spec}(R) \mid I \subseteq p\}$, where I ranges over the ideals of R .

Union and intersection is computed on such sets by $F_I \cap F_{I'} = F_{I+I'}$, where $I + I' = \{r + r' \mid r \in I, r' \in I'\}$, and $F_I \cup F_{I'} = F_{I \cap I'}$.

A ring R is *Noetherian* iff every \subseteq -increasing sequence of ideals $I_0 \subseteq I_1 \subseteq \dots \subseteq I_n \subseteq \dots$ in R is stationary: for some $n \in \mathbb{N}$, all the ideals I_n, I_{n+1}, \dots , are equal. For example, the ring $K[X_1, \dots, X_k]$ of all polynomials over the variables X_1, \dots, X_k with coefficients in K is Noetherian for any field K , in fact even for any Noetherian ring K . For any Noetherian ring R , $\text{Spec}(R)$ is a Noetherian topological space [Grothendieck, 1960, corollaire 1.1.6, p.81]. The specialization ordering of $\text{Spec}(R)$ is reverse inclusion \supseteq [Grothendieck, 1960, corollaire 1.1.7, p.81]. By [Grothendieck, 1960, proposition 1.1.10, (i), p.82], the sets $\text{Spec}(R) \setminus \downarrow(r)$ form a base of the Zariski topology, where (r) is the (prime) ideal generated by $r \in R$, so that $\downarrow(r) = \{p \mid p \supseteq (r)\} = \{p \mid r \in p\}$. In particular, the Zariski topology coincides with the upper topology of \supseteq (even when R is not Noetherian).

There are in general several ideals I that yield the same closed set F_I . In fact, two ideals yield the same closed set if and only if they have the same radical; the *radical* \sqrt{I} is defined as $\{r \in R \mid \exists k \geq 1, r^k \in I\}$.

Whatever the ring R , $\text{Spec}(R)$ is always sober [Grothendieck, 1960, corollaire 1.1.14, (ii), p.82]. It follows that its irreducible closed subsets are exactly its subsets of the form F_p , p a prime ideal, which are exactly the downward closure (with respect to \supseteq) of p . When $\text{Spec}(R)$ is also Noetherian, it follows from Lemma 3.6 that every closed subset F_I of $\text{Spec}(R)$ is a finite union of irreducible closed subsets $F_{p_1} \cup \dots \cup F_{p_n}$. Since the latter is equal to $F_{p_1 \cap \dots \cap p_n}$, $\sqrt{I} = \sqrt{p_1 \cap \dots \cap p_n}$, and the latter equals $p_1 \cap \dots \cap p_n$ since radical commutes with intersections and since $\sqrt{p} = p$ for every prime ideal p . So, every radical ideal I in a Noetherian ring is the intersection of finitely many prime ideals: this is *Kaplansky's Theorem* [Faith, 1999, Theorem 14.34]. Applying this to the closed subset $F_p \cap F_{p'}$, where p and p' are prime ideals, we obtain that $F_p \cap F_{p'} = F_{p+p'}$ is a finite union of irreducible closed subsets $F_{p_1} \cup \dots \cup F_{p_n}$ by Lemma 3.6. So $\sqrt{p+p'} = p_1 \cap \dots \cap p_n$. Applying this to the whole space $\text{Spec}(R) = F_{\{0\}}$, we obtain that $\sqrt{\{0\}} = \{0\} = p_1 \cap \dots \cap p_n$ for finitely many prime ideals p_1, \dots, p_n .

We therefore obtain an S-representation for $\text{Spec}(R)$, with enough computability assumptions on the ring R . The following proposition is almost vacuous, and only reflects our needs for S-representations at the level of rings.

Proposition 5.1 ($\text{Spec}(R)$) *Let R be a Noetherian ring, and assume that the set $\text{Spec}(R)$ of prime ideals of R is recursively enumerable, that the relation \trianglelefteq defined by $p \trianglelefteq p'$ iff $F_p \subseteq F_{p'}$ iff $\sqrt{p} \supseteq \sqrt{p'}$ is decidable, and that given $p, p' \in \text{Spec}(R)$ one can compute a finite set $p \wedge p'$ of elements $p_1, \dots, p_n \in \text{Spec}(R)$ such that $\sqrt{p+p'} = p_1 \cap \dots \cap p_n$. Let also τ be a finite set of prime ideals whose intersection is $\{0\}$.*

Then $(\text{Spec}(R), \text{id}_{\text{Spec}(R)}, \preceq, \tau, \wedge)$ is an S -representation of $\text{Spec}(R)$.

An important special case is given by taking the polynomial ring $K[X_1, \dots, X_k]$ for R , where K is a Noetherian ring. For the purpose of computability, we shall even concentrate on $\mathbb{Q}[X_1, \dots, X_k]$. The latter is an interesting space as far as verification of so-called polynomial programs is concerned [Müller-Olm and Seidl, 2002]: such programs have k rational-valued variables, and the only allowed operations are $+$, $-$, \times , assigning an arbitrary value to a variable non-deterministically, and testing for non-equality. The natural state space for such programs is \mathbb{Q}^k . However, \mathbb{Q}^k embeds into $\text{Spec}(\mathbb{Q}[X_1, \dots, X_k])$, by mapping every tuple (v_1, \dots, v_k) of values to the prime ideal generated by the polynomials $X_1 - v_1, \dots, X_k - v_k$. While Müller-Olm and Seidl computed with polynomial ideals directly [Müller-Olm and Seidl, 2002], one can alternatively notice that polynomial programs form a topological WSTS, where the state space \mathbb{Q}^k has the subspace topology from $\text{Spec}(\mathbb{Q}[X_1, \dots, X_k])$ [Goubault-Larrecq, 2010]¹.

To satisfy the requirements of Proposition 5.1 for $\text{Spec}(\mathbb{Q}[X_1, \dots, X_k])$, we represent polynomial ideals using *Gröbner bases* [Buchberger and Loos, 1983, Section 11], which are certain finite sets of polynomials $u = \{P_1, \dots, P_n\}$ representing the ideal $(u) = \{Q_1 P_1 + \dots + Q_n P_n \mid Q_1, \dots, Q_n \in \mathbb{Q}[X_1, \dots, X_k]\}$. Given a Gröbner basis u , one can decide whether (u) is a prime ideal: see [Adams and Loustaunau, 1994, Algorithm 4.4.1, p.244] or [Grieco and Zucchetti, 1989, Section 5, end]. So the set S of all Gröbner bases u such that (u) is prime is recursively enumerable.

We can now define $\llbracket u \rrbracket$ as $F_{(u)}$.

Given two Gröbner bases u and v , it is easy to check whether $(u) \supseteq (v)$. It suffices to check whether $P \in (u)$ for every $P \in v$, and this proceeds by using the polynomials of u as rewriting rules and checking whether P rewrites to 0 [Buchberger and Loos, 1983, Section 11]. However, one needs to decide whether $\sqrt{(u)} \supseteq \sqrt{(v)}$, equivalently, $\sqrt{(u)} \supseteq (v)$, i.e., to decide whether $P \in \sqrt{(u)}$ for every $P \in v$. The easiest way to decide this is to use the *Rabinowitch trick* [Rabinowitch, 1929]: $P \in \sqrt{(u)}$ iff $1 \in (u \cup \{1 - YP\})$, where Y is a fresh variable.

It is clear that one can take $\tau = \{\{0\}\}$, the ideal generated by 0, or equivalently by the empty family of polynomials, since $\{0\}$ is a prime ideal in $\mathbb{Q}[X_1, \dots, X_k]$, and in fact the minimal prime ideal, so $F_{\{0\}}$ is the unique largest element of $\text{Spec}(\mathbb{Q}[X_1, \dots, X_k])$.

The really tricky part is in defining the intersection map \wedge , i.e., to give an effective version of Kaplansky's Theorem. The algorithms that allow us to do this are too complicated to even give a glimpse of here. One may consult [Laplagne, 2006].

Theorem 5.2 (S-Representation, Spectrum of a Polynomial Ring) *An S -representation $(S, \llbracket _ \rrbracket, \preceq, \tau, \wedge)$ of $\text{Spec}(\mathbb{Q}[X_1, \dots, X_k])$ in its Zariski topology is given by:*

- (A) S is the collection of Gröbner bases u on $\mathbb{Q}[X_1, \dots, X_k]$ such that (u) is a prime ideal.
- (B) $\llbracket u \rrbracket = F_{(u)}$.
- (C) $u \preceq v$ iff $1 \in (u \cup \{1 - YP\})$, where Y is a fresh variable, for every $P \in v$.

¹The state space was erroneously claimed to be \mathbb{C}^k in that paper.

(D) $\tau = \{\{0\}\}$.

(E) $u \wedge v$ is a finite collection of Gröbner bases u_1, \dots, u_n such that $(u_1), \dots, (u_n)$ are prime ideals and $\sqrt{u+v} = (u_1) \cap \dots \cap (u_n)$, computed by Laplagne's algorithm [Laplagne, 2006].

An alternative S-representation of $\text{Spec}(R)$ is given by using for S the set of all those finite sets u of polynomials such that (u) is a *primary ideal*, instead of a prime ideal. A primary ideal p is such that whenever rr' is in p , then $r \in p$ or some power of r' is in p . Every prime ideal is primary, but the converse fails. The radical \sqrt{p} of a primary ideal is always prime. Given a set u of polynomials, one can decide whether (u) is primary [Grieco and Zucchetti, 1989, Theorem 3.2], and in fact one can compute a Gröbner basis for $\sqrt{(u)}$ in this case. So S is again, in particular, recursively enumerable. We define again $\llbracket u \rrbracket$ as $F_{(u)}$. Since $F_{(u)} = F_{\sqrt{(u)}}$ and $\sqrt{(u)}$ is prime, $F_{(u)}$ is certainly an irreducible closed subset. Next, \preceq and τ are defined as above, while \wedge is now based on a computable variant of the *Lasker-Noether Theorem*, instead of Kaplansky's Theorem. This states that every ideal I in a Noetherian ring R can be written as the intersection of finitely many primary ideals. When $R = \mathbb{Q}[X_1, \dots, X_k]$, then one can even compute a finite collection of Gröbner bases w_1, \dots, w_m such that $(u+v) = (w_1) \cap \dots \cap (w_m)$ and $(w_1), \dots, (w_m)$ are primary ideals, see [Sturmfels, 2002, Chapter 5]. Now given $u, v \in S$, $\llbracket u \rrbracket \cap \llbracket v \rrbracket = F_{(u)} \cap F_{(v)} = F_{(u+v)} = F_{\sqrt{(u+v)}}$. One can compute a finite collection of Gröbner bases w_1, \dots, w_k such that $\sqrt{(u+v)} = \sqrt{(w_1)} \cap \dots \cap \sqrt{(w_k)}$ and $(w_1), \dots, (w_k)$ are primary ideals. Then $\llbracket u \rrbracket \cap \llbracket v \rrbracket = F_{\sqrt{(w_1)}} \cup \dots \cup F_{\sqrt{(w_k)}} = F_{(w_1)} \cup \dots \cup F_{(w_k)} = \bigcup_{i=1}^k \llbracket w_i \rrbracket$: define $u \wedge v$ as $\{w_1, \dots, w_k\}$.

To sum up:

Theorem 5.3 (S-Representation, Spectrum of a Polynomial Ring, Alternate) *An S-representation $(S, \llbracket _ \rrbracket, \preceq, \tau, \wedge)$ of $\text{Spec}(\mathbb{Q}[X_1, \dots, X_k])$ in its Zariski topology is given by:*

(A) S is the collection of Gröbner bases u on $\mathbb{Q}[X_1, \dots, X_k]$ such that (u) is a primary ideal.

(B) $\llbracket u \rrbracket = F_{(u)}$.

(C) $u \preceq v$ iff $1 \in (u \cup \{1 - YP\})$, where Y is a fresh variable, for every $P \in v$.

(D) $\tau = \{1\}$.

(E) $u \wedge v$ is a finite collection of Gröbner bases u_1, \dots, u_n such that $(u_1), \dots, (u_n)$ are primary ideals and $\sqrt{u+v} = (u_1) \cap \dots \cap (u_n)$, computed as in [Sturmfels, 2002, Chapter 5].

We finish this section by mentioning an issue with our polynomial program example. We really think of the state space as \mathbb{Q}^k , not the larger space $\text{Spec}(\mathbb{Q}[X_1, \dots, X_k])$. To make things formal, this means equipping \mathbb{Q}^k with the subspace topology, whose closed subsets are exactly those sets of the form $\mathcal{Z}(u) = \{\vec{x} \in \mathbb{Q}^k \mid \forall P \in u, P(\vec{x}) = 0\}$.

$0\}$, for u an ideal in $\mathbb{Q}[X_1, \dots, X_k]$. That topology is usually called the *Zariski topology* on \mathbb{Q}^k , and makes polynomial programs topological WSTS. Whether we use \mathbb{Q}^k or $\text{Spec}(\mathbb{Q}[X_1, \dots, X_k])$ is of little consequence if we use the backward algorithm mentioned in the introduction, because the only thing it cares about is open subsets, which can be encoded as complements of sets $\mathcal{Z}(u)$, namely as ideals u .

The situation is different with S-representations, since S-representations do not encode closed sets, but *irreducible* closed subsets, and $\text{Spec}(\mathbb{Q}[X_1, \dots, X_k])$ contains many more irreducible closed subsets than \mathbb{Q}^k . This boils down to the fact that we do not know of an S-representation for \mathbb{Q}^k with its Zariski topology: the situation for the apparently more complex space $\text{Spec}(\mathbb{Q}[X_1, \dots, X_k])$ is simpler.

The situation is the following: we have two spaces $X = \mathbb{Q}^k$ and $Y = \text{Spec}(\mathbb{Q}[X_1, \dots, X_k])$, and X is a subspace of Y ; we know of an S-representation for Y , can we infer one for X ?

Proposition 5.4 *Let X be a subspace of a topological space Y . Then $\mathcal{S}(X)$ embeds into $\mathcal{S}(Y)$, i.e., every irreducible closed subset of X can be equated, in a canonical way, with some irreducible closed subset of Y .*

Proof. Let $m: X \rightarrow Y$ be the inclusion map. Then $\mathcal{S}(m)$ is a topological embedding [Goubault-Larrecq, 2013, Lemma 8.4.11]. In other words, every irreducible closed subset C of X can be equated with $\mathcal{S}(m)(C)$, namely the closure of C in Y , and that is irreducible closed in Y . \square

In our case, this means that an S-representation for \mathbb{Q}^k consists in a subset of either set of codes considered in Proposition 5.2 or in Proposition 5.3. Characterizing those codes remains to be elucidated.

6 Completing Words

If X is a wqo, then X^* is a wqo again under the embedding quasi-ordering by Higman's Lemma. This is often used when X is a finite alphabet Σ , with equality as quasi-ordering, but more general wqos are sometimes needed. For instance, Abdulla *et al.* [Abdulla *et al.*, 2004b] need to use X^* where $X = \Sigma^{\otimes}$, the set of finite multisets on a finite alphabet Σ . (We will deal with multisets in Section 7.) In that case, X itself is infinite. That paper is also one where a suitable theory of downwards-closed subsets was first developed, on $(\Sigma^{\otimes})^*$, and our constructions will generalize theirs. Data nets [Lazič *et al.*, 2008] are transition systems on a state space of the form X^* with $X = \mathbb{N}^k$, for some $k \in \mathbb{N}$, and again X is infinite in this case. More recently, Leroux and Schmitz have analyzed the question of reachability in Petri nets [Leroux and Schmitz, 2015], and required to work on ideals in the space of runs of Petri nets, which is a subspace of $(\mathbb{N}^k)^*$.

We work at the more general level of Noetherian spaces. In that context, the analogue of Higman's Lemma reads: for every Noetherian space X , the set X^* of finite words over X taken as alphabet, is Noetherian again, with the so-called *word topology* [Goubault-Larrecq, 2013, Theorem 9.7.33]. (The converse also holds.) The latter topology is generated by basic open subsets $X^*U_1X^*U_2X^*\dots X^*U_nX^*$, where

$n \in \mathbb{N}$ and U_1, \dots, U_n are open subsets of X . We write AB for the sets of concatenations ww' of words $w \in A$ and $w' \in B$, and equate subsets of X such as U_i with the set of one-letter words whose letter is in U_i . So $X^*U_1X^*U_2X^*\dots X^*U_nX^*$ is the (open) subset of words containing a not necessarily contiguous word $a_1a_2\dots a_n$ with $a_1 \in U_1, a_2 \in U_2, \dots, a_n \in U_n$. We stress that such subsets form a *base*, not just a subbase:

Lemma 6.1 *Let X be a topological space. Call elementary open of X^* any subset of the form $X^*U_1X^*U_2X^*\dots X^*U_nX^*$, with all U_i open in X . Every finite intersection of elementary opens can be expressed as a finite union of elementary opens. In particular, the elementary opens form a base of the word topology.*

Proof. This is Exercise 9.7.28 of [Goubault-Larrecq, 2013]. An empty intersection is just X^* , and the intersection of $X^*U_1X^*U_2X^*\dots X^*U_mX^*$ and $X^*V_1X^*V_2X^*\dots X^*V_nX^*$ is computed by induction on $m+n$ using the auxiliary formulae $X^*\cap\mathcal{V} = \mathcal{V}$, $\mathcal{U}\cap X^* = \mathcal{U}$, and $X^*U_1\mathcal{U}\cap X^*V_1\mathcal{V} = X^*U_1(\mathcal{U}\cap X^*V_1\mathcal{V}) \cup X^*V_1(X^*U_1\mathcal{U}\cap\mathcal{V}) \cup X^*(U_1\cap V_1)(\mathcal{U}\cap\mathcal{V})$. \square

If \leq is the specialization quasi-ordering of X , then the specialization quasi-ordering of X^* is the standard *embedding quasi-ordering* \leq^* , a.k.a. Higman's divisibility quasi-ordering [Higman, 1952]: $w \leq^* w'$ iff, writing w as the sequence of m letters $a_1a_2\dots a_m$, one can write w' as $w_0a'_1w_1a'_2w_2\dots w_{m-1}a'_mw'_m$ with $a_1 \leq a'_1, a_2 \leq a'_2, \dots, a_m \leq a'_m$. Higman's Lemma states that if X is well-quasi-ordered by \leq , then X^* is well-quasi-ordered by \leq^* [Higman, 1952]. The fact that X^* is Noetherian if and only if X is Noetherian is a natural generalization of Higman's Lemma: the latter can be obtained as a special case by considering Alexandroff topologies [Goubault-Larrecq, 2013, Exercise 9.7.34].

The completion $\mathcal{S}(X^*)$ is well-known in case X is wqo. As mentioned in the introduction, this is due to Kabil and Pouzet [Kabil and Pouzet, 1992]. Kabil and Pouzet also look at the (ideal) completion of spaces of finite words over more general ordered sets X . We explore another direction, that where X is Noetherian. This will include the result by Kabil and Pouzet in the wqo case as a by-product. Additionally, we give a simple, dynamic programming algorithm for deciding inclusion between irreducible closed subsets, and computing intersections, retrieving formulae that were known in the case where X is finite [Abdulla et al., 1998].

To study the completion $\mathcal{S}(X^*)$, we start by examining the shape of closed subsets of X^* . For any subset A of X , let A^* denote the set of all words $a_1a_2\dots a_n$ with $a_1, a_2, \dots, a_n \in A$, $n \in \mathbb{N}$ (n is possibly equal to 0). Let $A^?$ be $A \cup \{\epsilon\}$. We delegate the proof of the following Lemma to Appendix B, and similarly for a certain number of other results of this Section. Our aim is to avoid disrupting the flow of arguments, and to proceed as fast as we can to the final result.

Lemma 6.2 *Let X be a topological space. The complement of $X^*U_1X^*U_2X^*\dots X^*U_nX^*$ ($n \in \mathbb{N}, U_1, U_2, \dots, U_n$ open in X) in X^* is \emptyset when $n = 0$, and $F_1^*X^?F_2^*X^?\dots X^?F_{n-1}^*X^?F_n^*$ otherwise, where $F_1 = X \setminus U_1, \dots, F_n = X \setminus U_n$.*

*If X is Noetherian, then this complement can be expressed as a finite union of sets of the form $F_1^*C_1^?F_2^*C_2^?\dots C_{n-1}^?F_n^*$, where C_1, C_2, \dots, C_{n-1} range over irreducible closed subsets of X .*

Definition 6.3 (Word-Product, Word-SRE) *Let X be a topological space. Call a word-product P on X any expression of the form $e_1 e_2 \cdots e_n$, where $n \geq 0$, and each e_i is an atomic expression, i.e., either F_i^* with F_i closed in X , or $F_i^?$ with F_i irreducible closed in X . The components of P are the closed sets F_1, \dots, F_n . Word-products are interpreted as the obvious subsets of X^* . When $n = 0$, this notation is abbreviated as ϵ , and denotes the one-element set $\{\epsilon\}$.*

Call word-SRE any finite sum of word-products, where sum is interpreted as union.

There is no harm in requiring F_i non-empty in addition, in atomic expressions F_i^* : indeed $\emptyset^* = \{\epsilon\}$, so such atomic expressions can simply be erased.

This definition is inspired from the products and *simple regular expressions* (SRE) of [Abdulla et al., 2004a]. Indeed, we get back the latter from Definition 6.3 in the case where X is a finite alphabet Σ , with the discrete topology (hence its specialization quasi-ordering is $=$). Then each closed subset F_i is just a finite subset, and each irreducible closed subset C_i is just a singleton.

Lemma 6.4 *Let X be a topological space. For every closed subset F of X , for every closed subset \mathcal{F} of X^* , $F^? \mathcal{F}$ is closed in X^* .*

Lemma 6.5 *Let X be a topological space. For every closed subset F of X , for every closed subset \mathcal{F} of X^* , $F^* \mathcal{F}$ is closed in X^* .*

Corollary 6.6 *Let X be a topological space. Every word-product, every word-SRE is closed in X^* .*

We can in fact say more:

Lemma 6.7 *Let X be a topological space. Every word-product is irreducible closed in X^* .*

It is instructive to see how X^* embeds in its completion $\mathcal{S}(X)$. Recall that the topological closure $\eta_X^{\mathcal{S}}(x)$ of a point $x \in X$ is also its downward closure $\downarrow x$, for the specialization quasi-ordering of X .

Lemma 6.8 (Embedding) *Let X be a topological space. The closure $\eta_{X^*}^{\mathcal{S}}(x_1 x_2 \cdots x_n)$ of the word $x_1 x_2 \cdots x_n$ in X^* is the word-product $\eta_X^{\mathcal{S}}(x_1)^? \eta_X^{\mathcal{S}}(x_2)^? \cdots \eta_X^{\mathcal{S}}(x_n)^?$.*

Proof. The latter is easily seen to be the downward closure of $x_1 x_2 \cdots x_n$ with respect to \leq^* , which is the specialization quasi-ordering of X^* . \square

We shall see that the converse of Lemma 6.7 holds: the irreducible closed subsets of X^* , i.e., the elements of $\mathcal{S}(X^*)$, are exactly the word-products when X is Noetherian. The following lemmas will serve to show this, as well as to give some ways of computing on word-products. We do not make an explicit distinction between syntax and semantics, on purpose, so as to avoid excessively formal notation.

Lemma 6.9 *Let X be a topological space. Inclusion between word-products can be checked in polynomial time (precisely in time proportional to the product of the lengths of the two word-products), modulo an oracle testing inclusion of closed subsets of X .*

Explicitly, we have: $\epsilon \subseteq P$ for any word-product P , $P \not\subseteq \epsilon$ unless all the atomic expressions in P are syntactically equal to \emptyset^ , and for all $C, C' \in \mathcal{S}(X)$, for all $F, F' \in \mathcal{H}_{\vee}(X)$, for all word-products P, P' :*

- $C^?P \subseteq C'^?P'$ if and only if $C \subseteq C'$ and $P \subseteq P'$, or $C \not\subseteq C'$ and $C^?P \subseteq P'$.
- $C^?P \subseteq F'^*P'$ if and only if $C \subseteq F'$ and $P \subseteq F'^*P'$, or $C \not\subseteq F'$ and $C^?P \subseteq P'$.
- $F^*P \subseteq C'^?P'$ if and only if F is empty and $P \subseteq C'^?P'$, or F is non-empty and $F^*P \subseteq P'$.
- $F^*P \subseteq F'^*P'$ if and only if $F \subseteq F'$ and $P \subseteq F'^*P'$, or $F \not\subseteq F'$ and $F^*P \subseteq P'$.

The above formulae lend themselves immediately to a dynamic programming algorithm, modulo an oracle O testing inclusion of closed subsets of X . Assume that we wish to test whether $P \subseteq P'$, where $P = e_1e_2 \cdots e_m$ and $P' = e_1e_2 \cdots e_n$. We create an $(m+1) \times (n+1)$ array $A = (a_{ij})_{0 \leq i \leq m, 0 \leq j \leq n}$. At the end of the algorithm, a_{ij} will be true if and only if $e_{i+1} \cdots e_m \subseteq e_{j+1} \cdots e_n$. We initialize A by letting a_{mj} be true for every j , $0 \leq j \leq n$. For every i , $0 \leq i < m$, we set a_{in} to false, unless e_{i+1}, \dots, e_m are all equal to \emptyset^* , in which case a_{in} is set to true; explicitly, we initialize a flag b to true, and enumerating i from $m-1$ to 0 , we do the following: if e_{i+1} is not of the form F^* , or is of the form F^* with $F \not\subseteq \emptyset$ (which we can decide using the oracle O), then set b to false, otherwise leave b unchanged; then set a_{in} to b . This completes the initialization phase. Then, using two nested loops on i and j , one enumerating i from $m-1$ to 0 , the other one enumerating j from $n-1$ to 0 (for each value of i), we set a_{ij} to true if and only if:

- e_{i+1} is of the form $C^?$, e_{j+1} is of the form $C'^?$, and either $C \subseteq C'$ (which we decide using the oracle O) and $a_{(i+1)(j+1)}$ is true, or $C \not\subseteq C'$ and $a_{i(j+1)}$ is true;
- or e_{i+1} is of the form $C^?$, e_{j+1} is of the form F'^* , and either $C \subseteq F'$ and $a_{(i+1)j}$ is true or $C \not\subseteq F'$ and $a_{i(j+1)}$ is true;
- or e_{i+1} is of the form F^* , e_{j+1} is of the form $C'^?$, and either F is empty (which we decide using O on F and \emptyset , as in the second part of the initialization phase) and $a_{(i+1)j}$ is true, or F is non-empty and $a_{i(j+1)}$ is true;
- or e_{i+1} is of the form F^* , e_{j+1} is of the form F'^* , and either $F \subseteq F'$ and $a_{(i+1)j}$ is true, or $F \not\subseteq F'$ and $a_{i(j+1)}$ is true.

Otherwise, we set a_{ij} to false. At the end of the nested loops, we return a_{00} , which is true if and only if $P \subseteq P'$. Alternatively to dynamic programming, we may use a directed recursive implementation, with memoization [Michie, 1968].

We can rephrase the equations of Lemma 6.9 in the slightly more synthetic, following form. This happens to be the inclusion of products as specified in [Abdulla et al., 2004a], in the case where X is a finite set. The fourth case is not needed if we first remove all atomic expressions \emptyset^* . We will refer to this specific formulation in the proof of Lemma 10.22, and in Definition 10.31.

Lemma 6.10 *Let X be a topological space. Given two atomic expressions e_1 and e'_1 , and two word-products P_1 and P'_1 , letting $P = e_1P_1$ and $P' = e'_1P'_1$, then $P \subseteq P'$ if and only if:*

- 1) $e_1 \not\sqsubseteq e'_1$ and $P \subseteq P'_1$,
- 2) or $e_1 = C^?$, $e'_1 = C'^?$, $C \subseteq C'$ and $P_1 \subseteq P'_1$,
- 3) or $e_1 = F'^*$, $e_1 \sqsubseteq e'_1$ and $P_1 \subseteq P'$,
- 4) or $e_1 = \emptyset^*$ and $P_1 \subseteq P'$.

The relation \sqsubseteq on atomic expressions is defined by: $C^? \sqsubseteq C'^?$ if and only if $C \subseteq C'$; $F^* \sqsubseteq F'^*$ if and only if $F \subseteq F'$; $C^? \sqsubseteq F'^*$ if and only if $C \subseteq F'$; and $F^* \sqsubseteq C'^?$ if and only if F is empty.

Corollary 6.11 *Let X be a topological space. Inclusion between word-SREs can be checked in polynomial time, modulo an oracle testing inclusion of closed subsets of X .*

Proof. By Lemma 4.2, and since word-products are irreducible closed (Lemma 6.7), inclusion of word-SREs $P_1 \cup \dots \cup P_m$ and $P'_1 \cup \dots \cup P'_n$ reduces to mn inclusion tests $P_i \subseteq P'_j$ between word-products, which we decide using the dynamic programming algorithm mentioned after Lemma 6.9. \square

We can also compute intersections of word-products.

Lemma 6.12 *Let X be a topological space. Any finite intersection of word-products is expressible as a finite union of word-products. Specifically, the intersection of two word-products is given by: $\epsilon \cap P = \epsilon$ for every word-product P , and by the recursive formulae:*

- $C^? P \cap C'^? P' = (C^? P \cap P') \cup (P \cap C'^? P') \cup (C \cap C')^? (P \cap P')$;
- $C^? P \cap F'^* P' = (C \cap F')^? (P \cap F'^* P') \cup (C^? P \cap P')$;
- $F^* P \cap F'^* P' = (F \cap F')^* (P \cap F'^* P') \cup (F^* \cap F')^* (F^* P \cap P')$.

Recall that the components of a word-product $P = e_1 e_2 \dots e_n$ are the components of each e_i , where the component of $C^?$ is C , and the component of F^* is F . Lemma 6.12 yields the following, more computation-oriented description of the intersection algorithm for word-products. Our particular way of presenting it will be helpful in Theorem 6.15, in Lemma 10.26, and in the proof of Lemma 10.34.

Lemma 6.13 *Let X be a Noetherian space. Define the finite set $\text{Meet}^{\mathcal{E}}(P, P')$ of word-products as follows, where P and P' are word-products, and the oracle \mathcal{E} maps pairs (F, F') of a component F of P and a component F' of P' to a finite set of irreducible closed subsets of X .*

First, let $\text{Meet}^{\mathcal{E}}(\epsilon, P') = \{\epsilon\}$, $\text{Meet}^{\mathcal{E}}(P, \epsilon) = \{\epsilon\}$. Then, let:

$$\begin{aligned} \text{Meet}^{\mathcal{E}}(C^?P, C'^?P') &= \{C''^?P'' \mid C'' \in \mathcal{E}(C, C'), P'' \in \text{Meet}^{\mathcal{E}}(P, P')\} \\ &\quad \cup \text{Meet}^{\mathcal{E}}(C^?P, P') \cup \text{Meet}^{\mathcal{E}}(P, C'^?P') \\ \text{Meet}^{\mathcal{E}}(C^?P, F'^*P') &= \begin{cases} \{C''^?P'' \mid C'' \in \mathcal{E}(C, F'), \\ P'' \in \text{Meet}^{\mathcal{E}}(P, F'^*P')\} \cup \text{Meet}^{\mathcal{E}}(C^?P, P') & \text{if } \mathcal{E}(C, F') \neq \emptyset, \\ \text{Meet}^{\mathcal{E}}(P, F'^*P') \cup \text{Meet}^{\mathcal{E}}(C^?P, P') & \text{otherwise} \end{cases} \\ \text{Meet}^{\mathcal{E}}(F^*P, C'^?P') &= \begin{cases} \{C''^?P'' \mid C'' \in \mathcal{E}(F, C'), \\ P'' \in \text{Meet}^{\mathcal{E}}(F^*P, P')\} \cup \text{Meet}^{\mathcal{E}}(P, C'^?P') & \text{if } \mathcal{E}(F, C') \neq \emptyset, \\ \text{Meet}^{\mathcal{E}}(F^*P, P') \cup \text{Meet}^{\mathcal{E}}(P, C'^?P') & \text{otherwise} \end{cases} \\ \text{Meet}^{\mathcal{E}}(F^*P, F'^*P') &= \left\{ \left(\bigcup_{C'' \in \mathcal{E}(F, F')} C'' \right)^* P'' \mid P'' \in \text{Meet}^{\mathcal{E}}(F^*P, P') \cup \text{Meet}^{\mathcal{E}}(P, F'^*P') \right\}. \end{aligned}$$

If \mathcal{E} computes intersections of closed subsets of X , i.e., is such that for any component F of P and any component F' of P' , $\mathcal{E}(F, F')$ is a finite family of irreducible closed subsets of X whose union is $F \cap F'$, then $\text{Meet}^{\mathcal{E}}(P, P')$ is a finite family of word-products whose union is $P \cap P'$.

Note that the map $(F, F') \mapsto \mathcal{E}(F, F')$ is well-defined, by Lemma 3.6. We will later require to be able to compute it.

Proof. The lemma is a simple consequence of Lemma 6.12.

In the first case, if $C \cap C'$ is non-empty, we conclude since $(C \cap C')^?(P \cap P') = \bigcup_{C'' \in \mathcal{E}(C, C')} C''^?(P \cap P')$, $P \cap P'$ is the union of the word-products in $\text{Meet}^{\mathcal{E}}(P, P')$, and unions distribute over concatenation. There is a subtle issue when $C \cap C'$ is empty. In that subcase, $(C \cap C')^?(P \cap P')$ is equal to $P \cap P'$, and that is different from $\bigcup_{C'' \in \mathcal{E}(C, C')} C''^?(P \cap P')$, which is empty; however, $(C \cap C')^?(P \cap P')$ is equal to $(P \cap P') \cup (C^?P \cap P') \cup (P \cap C'^?P')$, hence also to $(C^?P \cap P') \cup (P \cap C'^?P')$, because $P \cap P'$ is included in $C^?P \cap P'$ (or in $P \cap C'^?P'$), and that justifies the indicated formula again. In the second case (and symmetrically, the third case), we rely on $(C \cap F')^?(P \cap F'^*P') = \bigcup_{C'' \in \mathcal{E}(C, F')} C''^?(P \cap F'^*P')$, which is valid if $\mathcal{E}(C, F')$ is non-empty. If $\mathcal{E}(C, F')$ is empty, then $C \cap F'$ is empty, and then $(C \cap F')^?$ is not equal to $\bigcup_{C'' \in \mathcal{E}(C, F')} C''^?$, rather to $\{\epsilon\}$; so $(C \cap F')^?(P \cap F'^*P') = P \cap F'^*P'$ in that (sub)case. In the final case, we use the fact that $F \cap F' = \bigcup_{C'' \in \mathcal{E}(F, F')} C''$.

Finally, the definition of $\text{Meet}^{\mathcal{E}}(P, P')$ is well-founded, by induction on the number of atomic expressions in P and P' . \square

Proposition 6.14 *Let X be a Noetherian space. The closed subsets of X^* are the (languages of) word-SREs, and the irreducible closed subsets of X^* are the (languages of) word-products.*

Proof. Lemma 6.7 states that every word-product is irreducible closed.

Conversely, we observe that, in a Noetherian space Y with a base \mathcal{B} of opens, every open is a finite union of elements of \mathcal{B} . This is an easy consequence of the fact that every open, which is a union of elements of \mathcal{B} , is also compact.

Consider $Y = X^*$, \mathcal{B} consisting of the subsets of the form $X^*U_1X^*U_2X^*\cdots X^*U_nX^*$, where each U_i is open in X (Lemma 6.1). Taking the complements of finite unions of such basic opens, and using Lemma 6.2, one obtains that every closed subset of X^* is a finite intersection of finite unions of word-products. Distributing unions over intersections, and using Lemma 6.12, we conclude that every closed subset F is expressible as a word-SRE, i.e., as a finite union of word-products.

If F is also irreducible, it follows immediately that F is one of these word-products. \square

We now state the final S-representation we obtain, in a way that we hope will be readable. The pedantic, formal statement is given in the Appendix (Proposition B.2).

Theorem 6.15 (S-Representation, Words) *Let X be a Noetherian space, $X' = X^*$, and $(S, \llbracket _ \rrbracket, \sqsubseteq, \tau, \wedge)$ be an S-representation of X . Then $(S', \llbracket _ \rrbracket', \sqsubseteq', \tau', \wedge')$ is an S-representation of X' , where:*

- (A) S' is the collection of all word-products over the alphabet S , and $\llbracket _ \rrbracket'$ is defined in the obvious way.
- (B) \sqsubseteq' is defined using the procedure of Lemma 6.9, where inclusion of finite sets of elements of S is tested by: u is included in u' iff for every $a \in u$, there is an $a' \in u'$ such that $a \sqsubseteq a'$ (Lemma 4.2).
- (C) τ' is $\{\tau^*\}$.
- (D) \wedge' is implemented by the procedure $\text{Meet}^{\mathcal{E}}$ of Lemma 6.13, where the oracle \mathcal{E} is defined by $\mathcal{E}(u, u') = \bigcup_{a \in u, a' \in u'} (a \wedge a')$.

Note that $\llbracket _ \rrbracket'$ is surjective, as required: the irreducible closed subsets of X' are the word-products by Proposition 6.14.

7 Completing Multisets

If X is a wqo, then the space of finite multisets X^{\circledast} of elements of X , with a quasi-ordering \leq^{\circledast} to be defined below, is a wqo again. This is again typically used when X is a finite alphabet Σ : the multiset language generators of [Abdulla et al., 2004b] are the ideals of such a wqo Σ^{\circledast} .

Beyond finite alphabets, branching vector addition systems with states (BVASS) are a generalization of Petri nets with a form of branching, with applications in security [Verma and Goubault-Larrecq, 2005], in linear logic [de Groote et al., 2004], in structured databases [Bojańczyk et al., 2009, Jacquemard et al., 2016], and are a rediscovery of Rambow's multiset-valued linear indexed grammars [Rambow, 1994] in computational linguistics, see [Schmitz, 2010]. BVASS, and some of their extensions, can be conveniently represented as transition systems on the space $(\mathbb{N}^k)^{\circledast}$ of finite multisets of k -tuples of natural numbers [Jacobé de Naurois, 2014]. Note that the alphabet $(\mathbb{N}^k)^{\circledast}$ is infinite in this case.

As a final example, the synchronous polyadic π -calculus processes investigated in [Acciai and Boreale, 2012] are encoded as trees, which can be seen as nested multisets,

with bounded nesting depth. Encoding processes by trees of this form was pioneered by Meyer [Meyer, 2008]. Precisely, for a finite set Σ , which consists of channel names and so-called unit processes in that case, let $\mathcal{T}_0^\circledast(\Sigma)$ be defined recursively as the set of finite trees $f(m)$ where $f \in \Sigma$ and m is a finite multiset of elements of $\mathcal{T}_0^\circledast(\Sigma)$, quasi-ordered by the universal relation ($s \leq_0 t$ is always true). Define $\mathcal{T}_{k+1}^\circledast(\Sigma)$, for every $k \in \mathbb{N}$, as the set of finite trees $f(m)$ where $f \in \Sigma$ and m is a finite multiset of elements of $\mathcal{T}_k^\circledast(\Sigma)$, quasi-ordered by \leq_{k+1} , defined by $f(m) \leq_{k+1} f'(m')$ if and only if $f = f'$ and $m(\leq_k)^\circledast m'$. All processes are encoded as elements of $\mathcal{T}_k^\circledast(\Sigma)$ for some $k \in \mathbb{N}$. Equivalently, $\mathcal{T}_k^\circledast(\Sigma)$ is $\Sigma \times (\Sigma \times (\Sigma \times \dots \times (\Sigma \times Y^\circledast)^\circledast \dots)^\circledast)^\circledast$, where there are k nested uses of $_\circledast$, and $Y = \mathcal{T}_0^\circledast(\Sigma)$.

We again turn to a more general topological setting. Given any topological space, let X^\circledast be the set of all finite multisets on X . We shall write $\{x_1, \dots, x_n\}$ for the multiset containing exactly the elements x_1, \dots, x_n . We write \emptyset for the empty multiset, and $m \uplus m'$ for the multiset union of m and m' .

On the order-theoretic side, we quasi-order X^\circledast , not with the multiset extension \leq^{mul} of the specialization quasi-ordering \leq of X , rather with the following quasi-ordering.

Definition 7.1 (Sub-multiset) *The sub-multiset quasi-ordering \leq^\circledast is defined by: $\{x_1, x_2, \dots, x_m\} \leq^\circledast \{y_1, y_2, \dots, y_n\}$ if and only if there is an injective map $r: \{1, 2, \dots, m\} \rightarrow \{1, 2, \dots, n\}$ such that $x_i \leq y_{r(i)}$ for every i , $1 \leq i \leq m$.*

When \leq is just equality, this quasi-ordering makes $m \leq^\circledast m'$ if and only if every element of m occurs at least as many times in m' as it occurs in m : this is the \leq^m quasi-ordering considered, on finite sets X , by Abdulla *et al.* [Abdulla et al., 2004b, Section 2].

The multiset extension \leq^{mul} of \leq is more mainstream than \leq^\circledast . One usually defines $m \leq^{mul} m'$ if and only if one can obtain m from m' in finitely many steps, repeatedly replacing one element by finitely many strictly smaller ones.

Clearly, $m \leq^\circledast m'$ implies $m \leq^{mul} m'$. It turns out that \leq^\circledast is wqo for every wqo \leq . This implies that \leq^{mul} is wqo, too: any infinite sequence of multisets m_n , $n \in \mathbb{N}$, is such that there are indices $i < j$ such that $m_i \leq^\circledast m_j$, and therefore $m_i \leq^{mul} m_j$.

On the topological side, we simply observe that multisets are equivalence classes of finite words up to permutation. Accordingly, we topologize X^\circledast with the quotient topology [Goubault-Larrecq, 2013, Exercise 9.7.35]. The quotient map $\Psi: X^* \rightarrow X^\circledast$ sends every word $x_1 x_2 \dots x_n$ to the multiset $\{x_1, x_2, \dots, x_n\}$ and is sometimes called the *Parikh mapping* [Parikh, 1966]. We have the following results.

Proposition 7.2 *For every Noetherian space X , X^\circledast is Noetherian.*

A base of the topology on X^\circledast is given by the sets $\langle U_1, U_2, \dots, U_n \rangle$ with U_1, U_2, \dots, U_n open in X . The set $\langle U_1, U_2, \dots, U_n \rangle$ is defined as containing all multisets that contain one element from U_1 , another one from U_2 , ..., another one from U_n , or more precisely all multisets of the form $\{x_1, x_2, \dots, x_n\} \uplus m$ with $x_1 \in U_1, x_2 \in U_2, \dots, x_n \in U_n$.

The specialization quasi-ordering of X^\circledast is \leq^\circledast , where \leq is the specialization quasi-ordering of X .

If X has the Alexandroff topology of \leq , then X^\otimes has the Alexandroff topology of \leq^\otimes .

If \leq is wqo, then \leq^\otimes is wqo.

Proof. If X is Noetherian, then X^\otimes is, too, since every quotient of a Noetherian space is Noetherian [Goubault-Larrecq, 2013, Proposition 9.7.18 (v)].

The inverse image $\Psi^{-1}(\langle U_1, U_2, \dots, U_n \rangle)$ is equal to the union over all permutations π of $\{1, 2, \dots, n\}$ of the opens $X^*U_{\pi(1)}X^* \cdots X^*U_{\pi(n)}X^*$, hence is open in X^* (see Section 6). For every subset V of X^\otimes , V is open (in the quotient topology) if and only if $\Psi^{-1}(V)$ is open, so $\langle U_1, U_2, \dots, U_n \rangle$ is open in X^\otimes . To show that these sets form a base, take a multiset m and an open neighborhood V of m in X^\otimes . Write m as $\Psi(w)$ for some word $w \in X^*$. Since $w \in \Psi^{-1}(V)$, and we know of a base of the topology of X^* (Lemma 6.1), we can find open subsets U_1, U_2, \dots, U_n of X such that $w \in X^*U_1X^*U_2X^* \cdots X^*U_nX^* \subseteq \Psi^{-1}(V)$. Then $m = \Psi(w)$ is in $\Psi[X^*U_1X^*U_2X^* \cdots X^*U_nX^*] = \langle U_1, U_2, \dots, U_n \rangle$, which is included in V .

If $m \leq^\otimes m'$, then every basic open subset $\langle U_1, U_2, \dots, U_n \rangle$ that contains m also contains m' , so m is below m' in the specialization quasi-ordering. Conversely, we shall show that the downward closure $\downarrow^\otimes m'$ of m' with respect to \leq^\otimes is closed: if m is below m' in the specialization quasi-ordering, then m will be in the closure of m' , hence in $\downarrow^\otimes m'$, and this will imply that $m \leq^\otimes m'$. To show that $\downarrow^\otimes m'$ is closed, it is enough to show that $\Psi^{-1}(\downarrow^\otimes m')$ is closed in X^* , since Ψ is quotient. Write m' as $\Psi(w')$, where w' is the word $x_1x_2 \cdots x_n$. Then $\Psi^{-1}(\downarrow^\otimes m')$ is the union over all permutations π of $\{1, 2, \dots, n\}$ of the sets $(\downarrow x_{\pi(1)})^? (\downarrow x_{\pi(2)})^? \cdots (\downarrow x_{\pi(n)})^?$, which are closed by Lemma 6.7.

Assume now that X has the Alexandroff topology of \leq . The upward closure of a multiset $m = \{x_1, x_2, \dots, x_n\}$ in X^\otimes is equal to the open subset $\langle \uparrow x_1, \uparrow x_2, \dots, \uparrow x_n \rangle$. Every upwards-closed subset is the union of the upward closures of its points, hence is open, too, so X^\otimes has the Alexandroff topology of \leq^\otimes .

In particular, if \leq is wqo, then X with the Alexandroff topology of \leq is Noetherian, so X^\otimes is Noetherian, too. Since X^\otimes has the Alexandroff topology of \leq^\otimes , the latter is wqo. \square

Our candidates for (irreducible) closed subsets of X^\otimes are the Parikh images of word-products and word-SREs. Write $F \mid C_1, C_2, \dots, C_n$ for the family of all multisets that one can obtain by picking at most one element from C_1 (possibly zero), at most one from C_2, \dots , at most one from C_n , and as many as we wish from F . We think of the enumeration C_1, C_2, \dots, C_n as a multiset itself, hence invariant under permutation. Formally, $m \in F \mid C_1, C_2, \dots, C_n$ if and only if one can write m as $m_0 \uplus \{x_i \mid i \in I\}$, where all the elements of m_0 are in F , I is a subset of $\{1, 2, \dots, n\}$ and for each $i \in I$, x_i is in C_i (implicitly, up to permutation of the x_i , or equivalently of the C_i).

Definition 7.3 (M-Product, M-SRE) *Let X be a topological space. Call an m-product on X any subset of the form $F \mid C_1, C_2, \dots, C_n$, where $n \in \mathbb{N}$, F is a closed subset of X , and C_1, C_2, \dots, C_n range over irreducible closed subsets of X .*

When F is empty, we shall also write this as simply $\mid C_1, C_2, \dots, C_n$. When $n = 0$, we just write $F \mid$, and when $n = 0$ and $F = \emptyset$, we write this \mid .

An m-SRE is any finite union of m-products.

The proof of the following Proposition, as well as for most other results of this Section, are to be found in Appendix C.

Proposition 7.4 *Let X be a topological space. Then the m-SREs are closed in X^\otimes , and the m-products are irreducible closed.*

If X is Noetherian, then every irreducible closed subset of X^\otimes is an m-product, and every closed subset of X^\otimes is an m-SRE.

Again, it is instructive to see how X^\otimes embeds in its completion $\mathcal{S}(X^\otimes)$.

Lemma 7.5 (Embedding) *Let X be a topological space. The closure $\eta_{X^\otimes}^{\mathcal{S}} \{x_1, x_2, \dots, x_n\}$ of the multiset $\{x_1, x_2, \dots, x_n\}$ in X^\otimes is the m-product $|\eta_X^{\mathcal{S}}(x_1), \eta_X^{\mathcal{S}}(x_2), \dots, \eta_X^{\mathcal{S}}(x_n)$.*

Proof. By Proposition 7.4, $|\eta_X^{\mathcal{S}}(x_1), \eta_X^{\mathcal{S}}(x_2), \dots, \eta_X^{\mathcal{S}}(x_n)$ is (irreducible) closed, and is clearly the downward closure of $\{x_1, x_2, \dots, x_n\}$ with respect to \leq^\otimes . \square

One can decide inclusion between m-products by using Ψ again. This leads to the following algorithm.

Lemma 7.6 *Let X be a topological space. Inclusion between m-products can be checked in polynomial time, modulo an oracle testing inclusion of closed subsets of X .*

Explicitly, let $P = F | C_1, C_2, \dots, C_m$ and $P' = F' | C'_1, C'_2, \dots, C'_n$ be two m-products. Let $I = \{i_1, i_2, \dots, i_k\}$ be the subset of those indices i , $1 \leq i \leq m$, such that $C_i \not\subseteq F'$.

Then $P \subseteq P'$ if and only if $F \subseteq F'$ and there is an injective map $r: I \rightarrow \{1, 2, \dots, n\}$ such that $C_i \subseteq C'_{r(i)}$ for every $i \in I$ —in other words, $\{C_{i_1}, C_{i_2}, \dots, C_{i_k}\} \subseteq^\otimes \{C'_{1}, C'_{2}, \dots, C'_{n}\}$.

It may not be immediately obvious why this leads to a polynomial time algorithm. The reason is the following observation, due to Simon Halfon [Halfon, 2018, Corollary 7.14]. Let G be the bipartite graph whose vertex set is the disjoint union of I and of $\{1, 2, \dots, n\}$, and such that there is an edge from $i \in I$ to $j \in \{1, 2, \dots, n\}$ if and only if $C_i \subseteq C'_j$. Finding r means finding a matching of G that covers all the vertices in I . Let $N(G)$ be the number of edges in any maximum matching of G . $N(G)$ can be computed in polynomial time, say by the Ford-Fulkerson algorithm [Cormen et al., 2001, Section 26.2], and r exists if and only if $N(G) \geq k$, where k is the cardinality of I .

We now turn to S-representations.

Theorem 7.7 (S-Representation, Multisets) *Let X be a Noetherian space, $X' = X^\otimes$, and $(S, \llbracket _ \rrbracket, \preceq, \tau, \wedge)$ be an S-representation of X . Then $(S', \llbracket _ \rrbracket', \preceq', \tau', \wedge')$ is an S-representation of X' , where:*

- (A) *S' is the collection of all m-product notations, i.e., of all expressions of the form $A | u$, where A is a finite subset of S , and u is a multiset of elements of S . When $u = \{b_1, \dots, b_n\}$, we also write $A | b_1, \dots, b_n$ for $A | u$.*

- (B) $\llbracket A \mid b_1, \dots, b_n \rrbracket' = (\bigcup_{a \in A} \llbracket a \rrbracket) \mid \llbracket b_1 \rrbracket, \dots, \llbracket b_n \rrbracket$.
- (C) $A \mid u \leq' A' \mid u'$ if and only if $A \leq^b A'$ and $u_1 \leq^\otimes u'$ where u_1 is the subset of those elements $a \in u$ such that $a \leq a'$ for no $a' \in A'$.
- (D) τ' is $\{\tau \mid \emptyset\}$.
- (E) \wedge' is defined as follows. A matching $f: \{1, \dots, m\} \rightarrow \{1, \dots, n\}$ is any bijection from some subset of $\{1, \dots, m\}$ (the domain $\text{dom } f$) to some subset of $\{1, \dots, n\}$ (the codomain $\text{cod } f$). Then $(A \mid a_1, \dots, a_m) \wedge' (A' \mid a'_1, \dots, a'_n)$ is the collection of all m -product notations of the form $A'' \mid m_{1f} \uplus m_{2f} \uplus m_{3f}$, where:

- $A'' = \bigcup_{\substack{a \in A \\ a' \in A'}} (a \wedge a')$;
- f ranges over all matchings from $\{1, \dots, m\}$ to $\{1, \dots, n\}$;
- m_{1f} ranges over all multisets of the form $\{c_i \mid i \in \text{dom } f\}$ where $c_i \in a_i \wedge a'_{f(i)}$ for every $i \in \text{dom } f$;
- m_{2f} ranges over all multisets of the form $\{c_i \mid 1 \leq i \leq m, i \notin \text{dom } f\}$, where $c_i \in \bigcup_{a' \in A'} (a_i \wedge a')$ for each $i, 1 \leq i \leq m, i \notin \text{dom } f$;
- m_{3f} ranges over all multisets of the form $\{c'_j \mid 1 \leq j \leq n, j \notin \text{cod } f\}$, where $c'_j \in \bigcup_{a \in A} (a \wedge a'_j)$ for each $j, 1 \leq j \leq n, j \notin \text{cod } f$.

As a final note to this section, Abdulla *et al.* [Abdulla et al., 2004b] required a completion of $(A^\otimes)^*$, for some finite set A . We note that the elements of $\mathcal{S}((A^\otimes)^*)$ are exactly their *word language generators*, which we retrieve here in a principled way.

8 Completing Words, Prefix Topology

The word topology is not the only interesting topology on X^* that makes it Noetherian, assuming X Noetherian. The prefix topology is another [Goubault-Larrecq, 2013, Exercise 9.7.36], and its specialization quasi-ordering is a form of the prefix ordering. We mention that topology because its specialization quasi-ordering is *never* a wqo, unless X is trivial. Also, this is the topology needed to decide reachability of sets defined by forbidden patterns in the so-called oblivious k -stack system model of [Goubault-Larrecq, 2010, Section 5].

The prefix topology is defined not just on X^* , but more generally on sets of *heterogeneous words*, that is, words whose letters are taken from possibly distinct spaces, depending on their position. This of course includes the case of words in X^* , but heterogeneous words are a natural generalization to consider, and incur no additional difficulty.

Let $X_1, X_2, \dots, X_n, \dots$ be countably many topological spaces. A heterogeneous word over these spaces is any tuple (x_1, x_2, \dots, x_m) in $X_1 \times X_2 \times \dots \times X_m, m \in \mathbb{N}$. We write it as $x_1 x_2 \dots x_m$, and call $m = |w|$ the *length* of the form $w = x_1 x_2 \dots x_m$.

A *telescope* on $(X_n)_{n \geq 1}$ is a sequence $\mathcal{U} = U_0, U_1, \dots, U_n, \dots$ of opens, where U_n is open in $\prod_{i=1}^n X_i$ for each $n \in \mathbb{N}$, and such that $U_n X_{n+1} \subseteq U_{n+1}$ for every

$n \in \mathbb{N}$. (We write $U_n X_{n+1}$ instead of $U_n \times X_{n+1}$. When $n = 0$, $\prod_{i=1}^n X_i$ just contains the empty word ϵ , so that $\prod_{i=1}^n X_i$ is just $\{\epsilon\}$ in that case, with the only possible topology. U_0 must be open in that, and that means that U_0 must be empty or equal to $\{\epsilon\}$ itself.)

A telescope is a *wide telescope* iff $U_n = \prod_{i=1}^n X_i$ for some $n \in \mathbb{N}$ —equivalently, for all sufficiently large $n \in \mathbb{N}$. Given any telescope $\mathcal{U} = U_0, U_1, \dots, U_n, \dots$ on $(X_n)_{n \geq 1}$, let $[\mathcal{U}]$ be the set of heterogeneous words w over $X_1, X_2, \dots, X_k, \dots$, such that $w \in U_{|w|}$.

We write $\triangleright_{n=1}^{+\infty} X_n$ for the space of all *heterogeneous words* over $(X_n)_{n \geq 1}$, i.e., the disjoint union of all spaces $\prod_{i=1}^n X_i$, $n \in \mathbb{N}$, with the *prefix topology*, which is given by the trivial open \emptyset , plus all subsets of the form $[\mathcal{U}]$, \mathcal{U} a wide telescope on $(X_n)_{n \geq 1}$. One checks easily that those form a topology, and we shall say so explicitly in Proposition 8.1 below.

The point of the definition of the prefix topology is that its specialization quasi-ordering is the prefix quasi-ordering, defined by $a_1 a_2 \cdots a_m \leq^p b_1 b_2 \cdots b_n$, where $a_i, b_i \in X_i$ for all i , iff $m \leq n$, $a_1 \leq b_1$, $a_2 \leq b_2$, \dots , and $a_m \leq b_m$. (Here $a_i \leq b_i$ means that a_i is less than or equal to b_i in the specialization quasi-ordering of X_i .) This is part of the following result, which appears as [Goubault-Larrecq, 2013, Exercise 9.7.36].

Proposition 8.1 *Let $X_1, X_2, \dots, X_n, \dots$ be countably many topological spaces. The prefix topology on $\triangleright_{n=1}^{+\infty} X_n$ is indeed a topology. Its specialization quasi-ordering is the prefix quasi-ordering \leq^p .*

If $X_1, X_2, \dots, X_n, \dots$ are all Noetherian, then $\triangleright_{n=1}^{+\infty} X_n$ is Noetherian.

Proof. Let us abbreviate $X_1 X_2 \cdots X_n$ as All_n , for each $n \in \mathbb{N}$. When $n = 0$, $All_0 = \{\epsilon\}$.

It is easy to see that given wide telescopes $\mathcal{U}_i = U_{i0}, U_{i1}, \dots, U_{in}, \dots$, $i \in I$, the sequence $\bigcup_{i \in I} \mathcal{U}_i$ defined as $\bigcup_{i \in I} U_{i0}, \bigcup_{i \in I} U_{i1}, \dots, \bigcup_{i \in I} U_{in}, \dots$ is an infinite sequence of empty sets (if I is empty), or a wide telescope (if $I \neq \emptyset$) and $[\bigcup_{i \in I} \mathcal{U}_i] = \bigcup_{i \in I} [\mathcal{U}_i]$; moreover, if I is finite, then the sequence $\bigcap_{i \in I} \mathcal{U}_i$ defined as $\bigcap_{i \in I} U_{i0}, \bigcap_{i \in I} U_{i1}, \dots, \bigcap_{i \in I} U_{in}, \dots$ is also a wide telescope (when $I = \emptyset$, this is the wide telescope $All_0, All_1, \dots, All_n, \dots$), and $[\bigcap_{i \in I} \mathcal{U}_i] = \bigcap_{i \in I} [\mathcal{U}_i]$. Therefore, the prefix topology, as we have defined it, is indeed a topology.

Write \sqsubseteq temporarily for the specialization quasi-ordering of $\triangleright_{n=1}^{+\infty} X_n$. For every telescope $\mathcal{U} = U_0, U_1, \dots, U_n, \dots$, if $a_1 a_2 \cdots a_m \in [\mathcal{U}]$, and $a_1 a_2 \cdots a_m \leq^p b_1 b_2 \cdots b_n$, then $b_1 b_2 \cdots b_m$ is in U_m , since U_m is open hence upwards-closed in \leq^p . Since $U_m X \subseteq U_{m+1}$ by the definition of telescopes, $b_1 b_2 \cdots b_m b_{m+1}$ is in U_{m+1} , and by an easy induction, $b_1 b_2 \cdots b_m b_{m+1} \cdots b_n$ is in U_n . So \mathcal{U} is upwards-closed in \leq^p : $w \leq^p w'$ implies $w \sqsubseteq w'$.

Conversely, assume $w \sqsubseteq w'$, where $w = a_1 a_2 \cdots a_m$, and $w' = b_1 b_2 \cdots b_n$. We shall examine various wide telescopes \mathcal{U} such that $w \in [\mathcal{U}]$, and draw consequences from the fact that $w' \in [\mathcal{U}]$. Considering the telescope $\emptyset, \dots, \emptyset, All_m, All_{m+1}, \dots$, one sees that $m \leq n$. Considering the telescopes $\emptyset, \dots, \emptyset, U_m, All_{m+1}, All_{m+2}, \dots$, where U_m is an arbitrary open set of X^m of which w is a member, one sees that $b_1 b_2 \cdots b_m$ is in U_m , so $a_1 a_2 \cdots a_m$ is less than or equal to $b_1 b_2 \cdots b_m$ in the special-

ization quasi-ordering $\leq_1 \times \leq_2 \times \cdots \times \leq_m$ of $X_1 X_2 \cdots X_m$. As a consequence, $w \leq^\triangleright w'$.

Let us show that $X = \bigtriangleright_{n=1}^{+\infty} X_n$ is Noetherian, assuming that $X_1, X_2, \dots, X_n, \dots$, all are. For every non-empty wide telescope $\mathcal{U} = U_0, U_1, \dots, U_n, \dots$, there is a least number m such that U_m is non-empty, and a least number n such that $U_n = \text{All}_n$. Moreover, $m \leq n$. Call m the *small end* $m(\mathcal{U})$ of \mathcal{U} , n its *big end* $n(\mathcal{U})$. If $\lfloor \mathcal{U} \rfloor \subseteq \lfloor \mathcal{V} \rfloor$, then $m(\mathcal{U}) \geq m(\mathcal{V})$ (consider any word of length $m(\mathcal{U})$ in $\lfloor \mathcal{U} \rfloor$), and $n(\mathcal{U}) \geq n(\mathcal{V})$ (otherwise, consider any word of length $n(\mathcal{U})$ that is not in $\lfloor \mathcal{V} \rfloor$). It follows that, in any infinite ascending chain $\lfloor \mathcal{U}_1 \rfloor \subseteq \lfloor \mathcal{U}_2 \rfloor \subseteq \cdots \subseteq \lfloor \mathcal{U}_k \rfloor \subseteq \cdots$, all small ends coincide, and all big ends coincide, for all k large enough, say $k \geq p$. Let m be this common small end, n be the common big end. Then, for each $k \geq p$, \mathcal{U}_k is a telescope of the form $\emptyset, \dots, \emptyset, U_{km}, U_{k(m+1)}, \dots, U_{k(n-1)}, \text{All}_n, \text{All}_{n+1}, \dots$. In addition, for each j with $m \leq j < n$, $U_{pj} \subseteq U_{(p+1)j} \subseteq \cdots \subseteq \cdots U_{kj} \subseteq \cdots$ is an infinite ascending chain of opens in $X_1 X_2 \cdots X_j$. The latter is Noetherian, so the chain stabilizes, say at $k_j \geq p$. Therefore $\lfloor \mathcal{U}_1 \rfloor \subseteq \lfloor \mathcal{U}_2 \rfloor \subseteq \cdots \subseteq \lfloor \mathcal{U}_k \rfloor \subseteq \cdots$ stabilizes at $\max(p, k_m, k_{m+1}, \dots, k_{n-1})$. This holds for ascending chains of opens that exclude the empty open subset; the general case is easy. It follows that $\bigtriangleright_{n=1}^{+\infty} X_n$ is Noetherian. \square

When $X_1 = X_2 = \cdots = X_n = \cdots$ are the same space X , we write X^\triangleright for the space $\bigtriangleright_{n=1}^{+\infty} X_n$. Although it has the same elements as X^* , it has a definitely distinct topology. E.g., while \leq^* is wqo when the specialization ordering \leq of X is, \leq^\triangleright is well-founded but not well, as soon as X contains two incomparable elements a and b . Indeed, in this case $a, ba, bba, bbba, \dots$ is an infinite antichain.

In order to characterize the completion $\mathcal{S}(\bigtriangleright_{n=1}^{+\infty} X_n)$, we define the subset $\lceil F_1 F_2 \cdots F_k \rceil$ of $\bigtriangleright_{n=1}^{+\infty} X_n$, where $k \in \mathbb{N}$ and each F_i is closed in X_i , as the set of all heterogeneous words $a_1 a_2 \cdots a_m$ of $\prod_{i=1}^m X_i$ such that $m \leq k$, $a_1 \in F_1, a_2 \in F_2, \dots, a_m \in F_m$. (When $k = 0$, $\lceil F_1 F_2 \cdots F_k \rceil$ is just $\{\epsilon\}$.) The following, as well as other results of this Section, are proved in Appendix D.

Proposition 8.2 *Let $X_1, X_2, \dots, X_n, \dots$ be countably many topological spaces. The sets of the form $\lceil F_1 F_2 \cdots F_n \rceil$, where each F_i is closed in X_i , form a subbase of closed sets for $\bigtriangleright_{n=1}^{+\infty} X_n$: these sets are closed, and every closed subset is an intersection of finite unions of such sets.*

Lemma 8.3 *Let $X_1, X_2, \dots, X_n, \dots$ be countably many topological spaces. The subsets of the form $\lceil C_1 C_2 \cdots C_n \rceil$, where C_i is irreducible closed in X_i for each i , $1 \leq i \leq n$, are irreducible closed in $\bigtriangleright_{n=1}^{+\infty} X_n$.*

This is enough to state how $\bigtriangleright_{n=1}^{+\infty} X_n$ embeds in its completion.

Lemma 8.4 (Embedding) *Let $X_1, X_2, \dots, X_n, \dots$ be countably many topological spaces, and $Y = \bigtriangleright_{n=1}^{+\infty} X_n$. The closure $\eta_Y^{\mathcal{S}}(x_1 x_2 \cdots x_n)$ of the word $x_1 x_2 \cdots x_n$ in $X = \bigtriangleright_{n=1}^{+\infty} X_n$ is $\lceil \eta_X^{\mathcal{S}}(x_1) \eta_X^{\mathcal{S}}(x_2) \cdots \eta_X^{\mathcal{S}}(x_n) \rceil$.*

Proof. By Lemma 8.3, $\lceil \eta_X^{\mathcal{S}}(x_1) \eta_X^{\mathcal{S}}(x_2) \cdots \eta_X^{\mathcal{S}}(x_n) \rceil$ is irreducible closed, and contains $x_1 x_2 \cdots x_n$, so must contain $\eta_Y^{\mathcal{S}}(x_1 x_2 \cdots x_n)$. Conversely, every element of

$\langle \eta_X^S(x_1)\eta_X^S(x_2)\cdots\eta_X^S(x_n) \rangle$ is a (\leq^\triangleright -)prefix of $x_1x_2\cdots x_n$, and must therefore be in $\eta_Y^S(x_1x_2\cdots x_n)$. \square

There is just one extra irreducible closed subset in $\triangleright_{n=1}^{+\infty} X_n$, unless some X_n is empty:

Lemma 8.5 *Let $X_1, X_2, \dots, X_n, \dots$, be countably many non-empty topological spaces. The whole space $\triangleright_{n=1}^{+\infty} X_n$ is irreducible closed in itself.*

So we obtain the following description of all irreducible closed subsets.

Lemma 8.6 *Let $X_1, X_2, \dots, X_n, \dots$, be countably many non-empty topological spaces. The only irreducible closed subsets of $\triangleright_{n=1}^{+\infty} X_n$ are $\triangleright_{n=1}^{+\infty} X_n$ itself, and the subsets of the form $\langle C_1C_2\cdots C_n \rangle$, where C_i is irreducible closed in X_i for each i , $1 \leq i \leq n$.*

This suggests that $\mathcal{S}(\triangleright_{n=1}^{+\infty} X_n)$ coincides with $\triangleright_{n=1}^{+\infty} \mathcal{S}(X_n)$, with a new top element \top added, at least up to isomorphism. For any space Y , let Y^\top be the space obtained by adding a fresh element \top to Y , and whose closed subsets are those of Y , plus Y^\top itself. The specialization quasi-ordering of Y^\top is given by: $y \preceq^\top y'$ iff $y, y' \in Y$ and $y \preceq y'$, or $y' = \top$, where \preceq is the specialization quasi-ordering of Y .

Proposition 8.7 *Let $X_1, X_2, \dots, X_n, \dots$, be countably many non-empty topological spaces. The map $i: (\triangleright_{n=1}^{+\infty} \mathcal{S}(X_n))^\top \rightarrow \mathcal{S}(\triangleright_{n=1}^{+\infty} X_n)$ that sends \top to $\triangleright_{n=1}^{+\infty} X_n$ and the word $C_1C_2\cdots C_n$ (where $C_i \in \mathcal{S}(X_i)$ for each i) to $\langle C_1C_2\cdots C_n \rangle$ is an order-isomorphism and a homeomorphism.*

To complete the picture, we examine the case where some of the spaces X_n are empty. Taking n to be the largest index such that X_1, \dots, X_n are non-empty (and 0 if every X_i is empty), we then write $X_1\triangleright X_2\triangleright\cdots\triangleright X_n$, or $\triangleright_{k=1}^n X_k$, instead of $\triangleright_{k=1}^{+\infty} X_k$. Since there cannot be any $(n+1)$ st letter, this is a space of words of length at most n . Clearly as well $\triangleright_{k=1}^n X_k$ then does not depend on the actual spaces X_{n+1}, X_{n+2}, \dots , provided X_{n+1} is empty, which justifies the notation.

Lemma 8.8 *Let X_1, X_2, \dots, X_n be non-empty topological spaces. The only irreducible closed subsets of $\triangleright_{k=1}^n X_k$ are the subsets of the form $\langle C_1C_2\cdots C_m \rangle$, where C_i is irreducible closed in X_i for each i , $1 \leq i \leq m$, and $m \leq n$.*

We then obtain an isomorphism as in Proposition 8.7, without the need to add a top element \top .

Proposition 8.9 *Let X_1, X_2, \dots, X_n be non-empty topological spaces. The map $i: \triangleright_{k=1}^n \mathcal{S}(X_k) \rightarrow \mathcal{S}(\triangleright_{k=1}^n X_k)$ that sends the word $C_1C_2\cdots C_k$ (where $k \leq n$ and $C_i \in \mathcal{S}(X_i)$ for each i) to $\langle C_1C_2\cdots C_k \rangle$ is an order-isomorphism and a homeomorphism.*

We therefore obtain:

Theorem 8.10 (S-Representation, Prefix) *Let $X_1, X_2, \dots, X_n, \dots$ be countably many Noetherian spaces, $X' = \bigtriangleright_{n=1}^{+\infty} X_n$, and $(S_i, \llbracket _ \rrbracket_i, \leq_i, \tau_i, \wedge_i)$ be an S -representation of X_i for each $i \geq 1$. Assume that the disjoint sum $\coprod_{i=1}^{+\infty} S_i$ is recursively enumerable, that $a \leq_i b$ is decidable in a, b, i , that τ_i is computable in i , that $a \wedge_i b$ is computable in a, b, i .*

Then $(S', \llbracket _ \rrbracket', \leq', \tau', \wedge')$ is an S -representation of X' , where:

- (A) S' is the set of all heterogeneous words over $S_1, S_2, \dots, S_n, \dots$, plus a fresh element ω in case no S_n is empty (i.e., no X_n is empty).
- (B) $\llbracket a_1 a_2 \cdots a_n \rrbracket' = \llbracket a_1 \rrbracket_1 \llbracket a_2 \rrbracket_2 \cdots \llbracket a_n \rrbracket_n$ where $a_1 \in S_1, a_2 \in S_2, \dots, a_n \in S_n$, and $\llbracket \omega \rrbracket' = \bigtriangleright_{n=1}^{+\infty} X_n$ (if no X_n is empty).
- (C) \leq' is defined by: $u \leq' \omega$ for all $u \in S'$ and $\omega \leq' u$ for no word $u \neq \omega$ in S' (in case no X_n is empty), and $a_1 a_2 \cdots a_m \leq' a'_1 a'_2 \cdots a'_n$ iff $m \leq n, a_1 \leq_1 a'_1, a_2 \leq_2 a'_2, \dots$, and $a_m \leq_m a'_m$.
- (D) τ' is $\{\omega\}$ if no X_n is empty, $\{a_1 a_2 \cdots a_n \mid a_1 \in \tau_1, a_2 \in \tau_2, \dots, a_n \in \tau_n\}$ otherwise, where n is the largest index such that X_n is non-empty.
- (E) \wedge' is defined by: $\omega \wedge' u' = \{u'\}$, $u \wedge' \omega = \{u\}$ (if no X_n is empty), and $a_1 a_2 \cdots a_m \wedge' a'_1 a'_2 \cdots a'_n = \{c_1 c_2 \cdots c_{\min(m,n)} \mid c_1 \in a_1 \wedge_1 a'_1, c_2 \in a_2 \wedge_2 a'_2, \dots, c_{\min(m,n)} \in a_{\min(m,n)} \wedge_{\min(m,n)} a'_{\min(m,n)}\}$.

Proof. Follows easily from Proposition 8.7 in case no X_n is empty, or from Proposition 8.9 otherwise. \square

When $X_1 = X_2 = \dots = X_n = \dots$ are all the same space, one can drop the subscripts to S, \leq, τ, \wedge . Then $u \leq' u'$ is decidable in polynomial time modulo an oracle for \leq .

Remark 8.11 *We have already seen an example of spaces that are Noetherian, but not wqo in their specialization preordering, for example $\mathbb{P}(X)$ where X is Noetherian. The construction X^\triangleright is another example: while X^\triangleright is Noetherian for X Noetherian, the prefix ordering \leq^\triangleright is not wqo, even if \leq is wqo.*

9 Completing Finite Trees in a Simple Case

The case of finite trees is by far the most complex one. We start with a simple case: that of *ranked* trees, whose vertices are decorated by function symbols from a *finite* set, with equality as ordering. The ordering on trees is the so-called homeomorphic embedding \preceq , a wqo by Kruskal's Tree Theorem [Kruskal, 1960]. In that case, the topology will be the Alexandroff topology, and irreducible closed subsets will be ideals. We shall also take advantage of the assumptions to give short, automata-theoretic proofs of some of our results.

The completions of sets of trees over a finite set of function symbols were already considered in [Wies et al., 2010], where it was used to decide coverability for depth-bounded processes without requiring one to know the depth bound in advance, and in

[Goubault-Larrecq and Schmitz, 2016], where it was used to decide piecewise testable separability of regular tree languages.

The more general case of unranked trees with function symbols taken from a Noetherian space, will be dealt with in Section 10. This will definitely be more complicated, but will share many similarities with the present case—apart from the fact that we will not be able to use automata-theoretic methods.

Let Σ be a finite *signature*, namely a finite set of so-called *function symbols* f, g, \dots , each coming with a natural number called its *arity*. Let Σ_r be the subset of those elements of Σ that have arity exactly r .

The set $\mathcal{T}(\Sigma)$ of *terms* over Σ [Baader and Nipkow, 1998] is the smallest such that, for every $f \in \Sigma_r$, for all $t_1, \dots, t_r \in \mathcal{T}(\Sigma)$, $f(t_1, \dots, t_r)$ is in $\mathcal{T}(\Sigma)$. By $f(t_1, \dots, t_n)$, we mean the tree whose root is labeled f , and has a list of r subtrees t_1, \dots, t_r , from left to right.

Our terms are ground: there is no variable involved here, although one may code variables as specific constants, i.e., as specific function symbols of arity 0.

The embedding ordering \preceq is defined by $s \preceq t$ if and only if:

- either $t = g(t_1, \dots, t_n)$ and $s \preceq t_j$ for some $j, 1 \leq j \leq n$;
- or $s = f(s_1, \dots, s_m), t = g(t_1, \dots, t_n), f = g$ (hence $m = n$) and $s_1 \preceq t_1, \dots, s_n \preceq t_n$.

Since the canonical S-representation for finite words under the word topology (or the \leq^* quasi-ordering) consists in certain regular expressions, we shall similarly define an S-representation for $\mathcal{T}(\Sigma)$ as certain tree regular expressions [Comon et al., 2004, Section 2.2].

We define *simple tree regular expressions* (over Σ), a.k.a. *STREs*, by the following abstract syntax:

$$\begin{aligned} S &::= 0 \mid P \mid S + S & P &::= f^{\bar{?}}(S, \dots, S) \mid \mathfrak{C}^{\bar{*}}.S \\ \mathfrak{C} &::= 0 \mid A \mid \mathfrak{C} + \mathfrak{C} & A &::= f(S_{\square}, \dots, S_{\square}) & S_{\square} &::= S \mid \square \end{aligned}$$

where $f \in \Sigma_r$ in $f^{\bar{?}}(P_1, \dots, P_r)$ and in $f(S_{\square_1}, \dots, S_{\square_r})$, the sum operation $+$ is associative and commutative (we shall sometimes write $\sum_{i=1}^m P_i$ for $P_1 + \dots + P_m$) with 0 denoting the empty sum, and $\square \notin \Sigma$ is a placeholder called the *hole*. Note that \square is not meant to denote a family of placeholders, rather a single one. The extended trees over the signature $\Sigma \cup \{\square\}$, where \square has arity 0, are called *contexts*.

The standard notations for $\bar{?}$ and $\bar{*}$ are $?$ and $*$. We want to distinguish $\bar{?}$ visually from $?$, since we will need both in Section 10; and similarly for $\bar{*}$ and $*$.

The STREs of the form P are called *tree pre-products*. Among them, the *tree products* will be our notations for ideals. They will satisfy additional constraints, which we shall define later (Definition 9.5).

To define the *semantics* of STREs, we write $t \in S$ as an abbreviation for ‘ t is in the language of S ,’ and define this language by structural induction on S .

Accordingly, $t \in f^{\bar{?}}(S_1, \dots, S_n)$ if and only if either t is of the form $f(t_1, \dots, t_n)$ with $t_i \in S_i$ for every $i, 1 \leq i \leq n$, or if $t \in \bigcup_{i=1}^n S_i$. The latter is necessary for S to denote a downwards-closed language.

As the notation suggests, for $S = P_1 + \dots + P_m$, $t \in S$ iff $t \in P_j$ for some j , $1 \leq j \leq m$, and the language of 0 is empty.

The productions of \mathfrak{C} , A , and S_\square serve to form *iterators* $\mathfrak{C}^{\bar{\cdot}}$. S . The language of $A = f(S_{\square_1}, \dots, S_{\square_n})$ consists of those contexts in $\mathcal{T}(\Sigma \cup \{\square\})$ of the form $f(c_1, \dots, c_n)$ where $c_i \in S_{\square_i}$ for every i . In turn, $c \in S_\square$ if and only if either $S_\square = \square$ and c is the trivial context \square , or S_\square is an STRE S , c is a tree t in $\mathcal{T}(\Sigma)$, and $t \in S$. The language of $\mathfrak{C} = A_1 + \dots + A_m$ is the union of the languages of A_j , $1 \leq j \leq m$.

Intuitively, the language of $\mathfrak{C}^{\bar{\cdot}}$. S should consist of all trees obtained by applying contexts in \mathfrak{C} , repeatedly, until one reaches a tree in S . For example, $(f(\square))^{\bar{\cdot}}$. $a^{\bar{\cdot}}$ will recognize all trees of the form $f^n(a)$, $n \in \mathbb{N}$. There are however two catches.

- 1) The first one has to do with patterns A where \square occurs more than once: as usual with tree regular expressions, in replacing \square by a tree from S , several occurrences of \square can be replaced by *different* trees from S . Hence $(f(\square, \square))^{\bar{\cdot}}$. $(a^{\bar{\cdot}} + b^{\bar{\cdot}})$ consists of all binary-branching trees with inner nodes labelled f and leaves labelled a or b , including $f(f(a, a), a)$ and $f(f(b, b), b)$ but also $f(f(a, b), a)$ or $f(f(b, b), a)$ among others. (We assume f binary, and a and b of arity 0.) For a context c , and a set of trees S , accordingly, we shall write $c[S]$ for the set of trees obtained from c by replacing each occurrence of \square by a (possibly different) tree in S .
- 2) The second catch has to do with downward closure. It is tempting to define the trees of $\mathfrak{C}^{\bar{\cdot}}$. S as those in $c_1[\dots[c_k[S]]\dots]$, for some $k \in \mathbb{N}$ and some $c_1, \dots, c_k \in \mathfrak{C}$. However, there are cases where that language would fail to be downwards-closed, e.g., $(f(a^{\bar{\cdot}}, \square))^{\bar{\cdot}}$. $b^{\bar{\cdot}}$ would contain $f(a, b)$ but not a , according to that semantics.

We repair that as follows. For $A = f(S_{\square_1}, \dots, S_{\square_n})$, define $\text{args } A$, the *argument support* of A , as the set of trees $t \in \mathcal{T}(\Sigma)$ such that some context $f(\dots, t, \dots)$ (i.e., with one of its arguments equal to t) is in the language of A . Alternatively: if those S_{\square_i} , $1 \leq i \leq n$, that are different from \square define non-empty languages, then $\text{args } A$ is the union of those languages; if some $S_{\square_i} \neq \square$ has an empty language, then $\text{args } A = \emptyset$. Hence, for example, $\text{args } f(\square, \square) = \emptyset$, $\text{args } f(a^{\bar{\cdot}}, \square) = a^{\bar{\cdot}} = \{a\}$, and $\text{args } f'(a^{\bar{\cdot}}, \square, 0) = \emptyset$. For $\mathfrak{C} = A_1 + \dots + A_m$, let $\text{args } \mathfrak{C} = \bigcup_{j=1}^m \text{args } A_j$.

For every $c \in \mathfrak{C}$, let us write $c^{\bar{\cdot}}[S]$ for $c[S] \cup S$. We are now ready to define the language of $\mathfrak{C}^{\bar{\cdot}}$. S , as the language of trees in $c_1^{\bar{\cdot}}[\dots[c_k^{\bar{\cdot}}[S \cup \text{args } \mathfrak{C}]]\dots]$, for some $k \in \mathbb{N}$ and some $c_1, \dots, c_k \in \mathfrak{C}$ —in writing $S \cup \text{args } \mathfrak{C}$, we equate the STRE S with the language it defines.

Proposition 9.1 *Every STRE defines a downwards-closed language of $\mathcal{T}(\Sigma)$ with respect to \preceq . Conversely, every downwards-closed language of $\mathcal{T}(\Sigma)$ with respect to \preceq is the language of some STRE.*

Proof. For the first part, we use induction on the size of the STRE, and the main point consists in checking that if $t \in \mathfrak{C}^{\bar{\cdot}}$. S then any smaller tree s (w.r.t. \preceq) is also in $\mathfrak{C}^{\bar{\cdot}}$. S . By induction hypothesis, S is downwards-closed. We use a secondary induction on the k used in the definition of the language of $\mathfrak{C}^{\bar{\cdot}}$. S as $c_1^{\bar{\cdot}}[\dots[c_k^{\bar{\cdot}}[S \cup \text{args } \mathfrak{C}]]\dots]$. If $k = 0$, then $t \in S \cup \text{args } \mathfrak{C}$, hence $s \in S \cup \text{args } \mathfrak{C}$ as well since both S and $\text{args } \mathfrak{C}$




Figure 3: Converting tree automata to STREs.

are downwards-closed—args \mathcal{C} is downwards-closed by the outer induction hypothesis. Otherwise $k \geq 1$. If t is in $c_2^{\bar{c}}[\dots[c_k^{\bar{c}}[S \cup \text{args } \mathcal{C}]] \dots]$, then we conclude by the inner induction hypothesis directly. Otherwise, one of the summands in c_1 is of the form $f(S_{\square_1}, \dots, S_{\square_n})$, and $t = f(t_1, \dots, t_n)$. If s is smaller than some t_i , then either $S_{\square_i} = \square$ and $t_i \in c_2^{\bar{c}}[\dots[c_k^{\bar{c}}[S \cup \text{args } \mathcal{C}]] \dots]$, which allows us to conclude by the inner induction hypothesis; or S_{\square_i} is of the form P , and is downwards-closed by the outer induction hypothesis. If instead $s = f(s_1, \dots, s_n)$ where s_i is smaller than t_i for each i , we conclude similarly that $s_i \in \mathcal{C}^*.S$ for each position i such that $S_{\square_i} = \square$. For all other positions i , S_{\square_i} is an STRE, which is downwards-closed by the outer induction hypothesis, so $s_i \in S_{\square_i}$. Hence s is in $c_1[\mathcal{C}^*.S]$, and therefore in $\mathcal{C}^*.S$.

For the converse direction, let L be a downwards-closed language. The complement $\mathcal{T}(\Sigma) \setminus L$ of L is upwards-closed, and since \preceq is a wqo, $\mathcal{T}(\Sigma) \setminus L$ can be written as the upward closure $\uparrow\{t_1, t_2, \dots, t_n\}$ of finitely many trees. For each i , $\uparrow t_i$ is easily seen to be recognizable by a finite (bottom-up) tree automaton. Since finite unions and complements of recognizable languages are recognizable, L is recognized by some finite tree automaton \mathcal{A} .

We now convert \mathcal{A} to an STRE. In general, we describe a procedure that converts any (ε -free) finite tree automaton \mathcal{A} to an STRE whose language is the downward closure $\downarrow L(\mathcal{A})$ of the language recognized by \mathcal{A} . This is best explained on an example: see Figure 3a, where there is one transition a of arity 0 (from no state) to state q_1 , one binary transition f from the pair of states q_1, q_2 to q_3 , and so on. In textual form, we write these transitions as rewrite rules [Comon et al., 2004]: $a \rightarrow q_1$, $f(q_1, q_2) \rightarrow q_3$, $h(q_3, q_4) \rightarrow q_2$, $d \rightarrow q_3$, $g(q_4) \rightarrow q_4$, $b \rightarrow q_4$. A tree t is *recognized at a state* s if and

only if $t \rightarrow^* s$, using the rewrite rules of \mathcal{A} . There is a set of final states, marked with an outgoing arrow with dangling end: in our example, just q_3 . The language $L(\mathcal{A})$ of \mathcal{A} is the set of trees recognized at some final state.

We first extend our automaton with ε -transitions. An ε -transition from s to s' will be drawn as a dashed arrow, see Figure 3b, and is just a rewrite rule of the new form $s \rightarrow s'$. This implies that every tree recognized at s is also recognized at s' . For each transition, say $f(s_1, s_2, \dots, s_n) \rightarrow s$, of \mathcal{A} , we add n ε -transitions $s_1 \rightarrow s$, $s_2 \rightarrow s$, \dots , $s_n \rightarrow s$. (To make things clear, we are assuming that \mathcal{A} does not originally contain any ε -transition.) Call the resulting automaton $\downarrow \mathcal{A}$. It is an easy exercise to show that $L(\downarrow \mathcal{A}) = \downarrow L(\mathcal{A})$.

There is a graph underlying $\downarrow \mathcal{A}$, whose vertices are the states of $\downarrow \mathcal{A}$, and whose edges are the ε -transitions. Build its strongly connected components: on Figure 3b, they are shown against a grey background. By construction, any two states in the same strongly connected component C recognize exactly the same trees, so it makes sense to talk of the language $L_C(\downarrow \mathcal{A})$ of those trees recognized at any state of C . Let $C \rightarrow C'$ if and only if $s \rightarrow s'$ for some $s \in C$, $s' \in C'$, $C \neq C'$. The strict ordering $<$ defined as the transitive closure \rightarrow^+ is well-founded, and we shall build an STRE S_C whose language is $L_C(\downarrow \mathcal{A})$, by induction along $<$.

If C is a trivial strongly connected component (one state s , no self-edge), then enumerate its incoming non- ε transitions $f_i(s_{i1}, s_{i2}, \dots, s_{in_i}) \rightarrow s$, $1 \leq i \leq m$. Let S_{ij} be an STRE whose language is the set of trees recognized at s_{ij} , which is given by induction hypothesis. Then $S_C = \sum_{i=1}^m f_i^{\bar{?}}(S_{i1}, S_{i2}, \dots, S_{in_i})$ is the desired STRE. For instance, the set of trees recognized at the leftmost state q_1 is the language of $a^{\bar{?}}$.

If C is a non-trivial strongly connected component, then enumerate the non- ε transitions $f_i(s_{i1}, s_{i2}, \dots, s_{in_i}) \rightarrow s_i$, $1 \leq i \leq m$, whose end state s_i is in C . For each pair i, j , if s_{ij} is in C , then let $S_{\square ij} = \square$; otherwise, let $S_{\square ij}$ be an STRE whose language is the set of trees recognized at s_{ij} , which we obtain by induction hypothesis. It is not too hard to see that $S_C = (\sum_{i=1}^m f_i(S_{\square i1}, S_{\square i2}, \dots, S_{\square in_i}))^{\bar{*}}.0$ is an STRE that suits our needs. For example, the rightmost strongly connected component $\{q_4\}$ yields the STRE $S_4 = (b + g(\square))^{\bar{*}}.0$. One might have expected an STRE of the more intuitive form $(g(\square))^{\bar{*}}.b^{\bar{?}}$, however note that they define exactly the same language.

Finally, $\downarrow L(\mathcal{A})$ is the union of the languages of the strongly connected components containing a final state; in our example the strongly connected component in the middle yields the final STRE $(d + f(a^{\bar{?}}, \square) + h(\square, S_4))^{\bar{*}}.0$. \square

We characterize the STREs that define ideals of $\mathcal{T}(\Sigma)$ with respect to \preceq . Let us define a rewrite relation \rightarrow_1 on STREs that moves all $+$ signs to the outside: for a \rightarrow_1 -normal STRE $S = P_1 + \dots + P_m$, each P_i will be irreducible, hence S will be an ideal, i.e., an irreducible closed subset, if and only if $m = 1$. (Recall that $\mathcal{S}(P_a) = \mathbf{I}(P)$ for every poset P .)

The rewrite relation \rightarrow_1 is defined in Figure 4. Recall that $+$ is understood modulo associativity and commutativity. Letters matter, too: S, S', S_1, \dots, S_n are STREs, while P, P' are those special STREs of the form $f^{\bar{?}}(S_1, \dots, S_n)$ or $\mathfrak{C}^{\bar{*}}.S$, etc. In particular, rule (R6) applies provided the pattern $f(S_1, \dots, S_n)$ does not contain \square at all. Similarly, in rule (R10), S and S' cannot contain \square .

For rule (R12), we need some auxiliary definitions.

(R1)	$P + P'$	\rightarrow_1	P'	if $P \subseteq P'$
(R2)	$A + A'$	\rightarrow_1	A'	if $A \subseteq A'$
(R3)	$0 + P$	\rightarrow_1	P	
(R4)	$0 + A$	\rightarrow_1	A	
(R5)	$0^*.S$	\rightarrow_1	S	
(R6)	$(\mathfrak{C} + f(S_1, \dots, S_n))^*.S$	\rightarrow_1	$\mathfrak{C}^*. (S + f^{\vec{?}}(S_1, \dots, S_n))$	
(R7)	$f^{\vec{?}}(\vec{S}_1, 0, \vec{S}_2)$	\rightarrow_1	0	
(R8)	$f^{\vec{?}}(\vec{S}_1, S + S', \vec{S}_2)$	\rightarrow_1	$f^{\vec{?}}(\vec{S}_1, S, \vec{S}_2) + f^{\vec{?}}(\vec{S}_1, S', \vec{S}_2)$	
(R9)	$f(\vec{S}_{\square 1}, 0, \vec{S}_{\square 2})$	\rightarrow_1	0	
(R10)	$f(\vec{S}_{\square 1}, S + S', \vec{S}_{\square 2})$	\rightarrow_1	$f(\vec{S}_{\square 1}, S, \vec{S}_{\square 2}) + f(\vec{S}_{\square 1}, S', \vec{S}_{\square 2})$	
(R11)	$\mathfrak{C}^*.0$	\rightarrow_1	0	if $\mathfrak{C} = \sum_{i=1}^m f_i(\square, \dots, \square)$ and no f_i has arity 0
(R12)	$\mathfrak{C}^*. (S + S')$	\rightarrow_1	$\mathfrak{C}^*.S + \mathfrak{C}^*.S'$	if \mathfrak{C} is \square -linear

Figure 4: The rewrite relation \rightarrow_1 .

Definition 9.2 A pattern $A = f(S_{\square 1}, \dots, S_{\square n})$ is \square -linear if and only if at most one $S_{\square i}$ is the hole \square .

Writing \mathfrak{C} as $A_1 + \dots + A_m$, we say that \mathfrak{C} is \square -linear if and only if every non-empty A_i is \square -linear.

The \square -linearity restriction imposed on the last rule is needed for the following to hold.

Lemma 9.3 If $S \rightarrow_1^* S'$ then S and S' define the same language. □

Lemma 9.4 Every STRE S has a normal form with respect to \rightarrow_1 .

Proof. Using Bachmair and Plaisted's associative path ordering $>_{apo}$ [Bachmair and Plaisted, 1985] on a precedence where $+$ is minimal, $f > f^{\vec{?}}$ for each symbol f , and the $(_)^* _$ operator has lexicographic status, we see that \rightarrow_1 is even a terminating relation: every sequence of rewrite steps terminates. (Bachmair and Plaisted's ordering has been improved upon many times, but is sufficient in the case of just one associative commutative symbol $+$.) □

Definition 9.5 A tree-product is any \rightarrow_1 -normal tree pre-product P .

Lemma 9.6 Every ideal, i.e., every irreducible closed subset of $\mathcal{T}(\Sigma)$ is the language of some tree-product.

Proof. By Proposition 9.1, an ideal I is the language of some STRE S . S has a \rightarrow_1 -normal form by Lemma 9.4, write it $P_1 + \dots + P_m$. Since I is non-empty, $m \geq 1$, and since I is irreducible closed, it is included in, hence equal to, the language of some P_i . □

Conversely, we check that the language of every tree-product is irreducible closed. In the special case that we are in, it is easier to show that they are directed sets.

Let us introduce some additional notation.

Definition 9.7 A pattern $A = f(S_{\square_1}, \dots, S_{\square_n})$ is \square -generated if and only if at least one S_{\square_i} is the hole \square ; it is empty if and only if some S_{\square_i} is different from \square and has an empty language.

Writing C as $A_1 + \dots + A_m$, we say that C is \square -generated if and only if every non-empty A_i is \square -generated, and is empty if and only if every A_i is empty.

By inspection of the rules defining \rightarrow_1 , we see:

Lemma 9.8 The tree-products are exactly the STREs of the form:

- $f^{\bar{?}}(P_1, \dots, P_n)$ where P_1, \dots, P_n are tree-products;
- or $\mathfrak{C}^{\bar{*}}.(P_1 + \dots + P_n)$ where $n \in \mathbb{N}$, P_1, \dots, P_n are pairwise incomparable tree-products, $\mathfrak{C} = \sum_{i=1}^m f_i(P_{\square_{i1}}, \dots, P_{\square_{in_i}})$, $m \geq 1$, each summand $f_i(P_{\square_{i1}}, \dots, P_{\square_{in_i}})$ is incomparable with any other, each pattern $P_{\square_{ij}}$ is either a tree-product or the hole \square , \mathfrak{C} is \square -generated, and one of the following conditions is satisfied: (a) \mathfrak{C} is not \square -linear and $n \geq 1$, or (b) \mathfrak{C} is not \square -linear, $n = 0$, and $P_{\square_{ij}} \neq \square$ for some i, j , or (c) \mathfrak{C} is \square -linear and $n \leq 1$.

Proof. A tree pre-product of the form $f^{\bar{?}}(S_1, \dots, S_n)$ is \rightarrow_1 -normal if and only if S_1, \dots, S_n are \rightarrow_1 -normal and neither (R7) nor (R8) applies. The latter means that each S_i is a tree-product P_i .

Next, consider a tree pre-product of the form $\mathfrak{C}^{\bar{*}}.S$. Write \mathfrak{C} as $\sum_{i=1}^m A_i$ where $A_i = f_i(S_{\square_{i1}}, \dots, S_{\square_{in_i}})$, and S as a sum of tree pre-products $P_1 + \dots + P_n$.

If $\mathfrak{C}^{\bar{*}}.S$ is \rightarrow_1 -normal, then (R5) does not apply, so $m \geq 1$. Since (R1) and (R3) do not apply (and $+$ is understood modulo associativity and commutativity), P_1, \dots, P_n are pairwise incomparable (and different from 0, but that is implied). Similarly, since (R2) and (R4) do not apply, each summand $f_i(P_{\square_{i1}}, \dots, P_{\square_{in_i}})$ is incomparable with any other. Additionally, (R9) and (R10) do not apply, so each $S_{\square_{ij}}$ is either equal to \square or equal to some tree-product. Henceforth write $S_{\square_{ij}}$ as $P_{\square_{ij}}$. If \mathfrak{C} were not \square -generated, then for some non-empty A_i , $P_{\square_{ij}}$ would be different from \square for every j : then rule (R6) would apply. We now prove that (a), (b), or (c) holds depending on the shape of \mathfrak{C} . If \mathfrak{C} is not \square -linear but (a) does not hold, then $n = 0$. Since (R11) does not apply, some $P_{\square_{ij}}$ is different from \square , or some f_i has arity 0. However, if some f_i has arity 0, then A_i would just be $f_i()$, and (R6) would apply: so (b) holds. It remains to consider the case where \mathfrak{C} is \square -linear. Since (R12) does not apply, it must be that $n \leq 1$, so (c) holds.

Conversely, it is easy to check that if the conditions listed in the statement of the lemma for $\mathfrak{C}^{\bar{*}}.(P_1 + \dots + P_n)$ are satisfied, then $\mathfrak{C}^{\bar{*}}.(P_1 + \dots + P_n)$ is \rightarrow_1 -normal. \square

Lemma 9.9 If S_1, \dots, S_n are directed STREs, then so is $f^{\bar{?}}(S_1, \dots, S_n)$.

Proof. Non-emptiness is clear, since S_1, \dots, S_n are non-empty. Let t, t' be any two trees in $f^{\bar{?}}(S_1, \dots, S_n)$. If $t = f(t_1, \dots, t_n)$ and $t' = f(t'_1, \dots, t'_n)$ with $t_i, t'_i \in S_i$ for every i , then we can find $t''_i \geq t_i, t'_i$ in S_i , and then $f(t''_1, \dots, t''_n) \geq t, t'$ is in $f^{\bar{?}}(S_1, \dots, S_n)$.

If t is in some S_j and $t' = f(t'_1, \dots, t'_n)$ with $t'_i \in S_i$ for every i , then build the tree $s = f(s_1, \dots, s_{j-1}, t, s_{j+1}, \dots, s_n)$, where s_i is an arbitrary tree from the non-empty

set S_i , $i \neq j$. Clearly, s is in $f^{\bar{?}}(S_1, \dots, S_n)$, so, by reduction to the previous case, there is a tree $t'' \in f^{\bar{?}}(S_1, \dots, S_n)$ such that $s, t' \preceq t''$. Since $t \preceq s$, we obtain that $t, t' \preceq t''$.

Similarly if t' is in some S_k , we build a new tree $s' = f(s'_1, \dots, s'_{k-1}, t', s'_{k+1}, \dots, s'_n)$ and conclude by a similar argument that there is a tree $t'' \in f^{\bar{?}}(S_1, \dots, S_n)$ such that $t' \preceq s' \preceq t''$ and $t \preceq t''$. \square

The case of STREs of the form $\mathfrak{C}^{\bar{?}}.S$ is more complex. The three cases (a), (b) and (c) in the lemma below match those of Lemma 9.8, second item.

Lemma 9.10 *Let $\mathfrak{C} = A_1 + \dots + A_m$ be a sum of patterns $A_i = f_i(S_{\square_{i1}}, \dots, S_{\square_{in_i}})$, where each $S_{\square_{ij}}$ that is different from \square has a non-empty language. If any of the following conditions is satisfied, then $\mathfrak{C}^{\bar{?}}.S$ is directed:*

- (a) \mathfrak{C} is not \square -linear, and S has a non-empty language, or
- (b) \mathfrak{C} is not \square -linear, and some $S_{\square_{ij}}$ is different from \square , or
- (c) \mathfrak{C} is \square -linear and \square -generated, and S is irreducible.

Proof. The fact that $\mathfrak{C}^{\bar{?}}.S$ is non-empty is an easy exercise: in case (a), $\mathfrak{C}^{\bar{?}}.S$ contains S ; in case (b), it contains $S_{\square_{ij}}$; in case (c), it contains S , which, as an irreducible subset, is necessarily non-empty. Let t, t' be any two trees in $\mathfrak{C}^{\bar{?}}.S$.

In case (a), some non-empty A_i is of the form $f(S_{\square_1}, \dots, S_{\square_n})$, where \square occurs at least twice, say at positions j and j' , $j \neq j'$. For every k , $1 \leq k \leq n$, define a tree t_k as follows: if $S_{\square_k} = \square$ and $k \neq j'$ (including the case $k = j$), let $t_k = t$, if $k = j'$ then let $t_k = t'$, and if $S_{\square_k} \neq \square$ then pick any tree for t_k from the language of S_{\square_k} , which is non-empty by assumption. We check that $f(t_1, \dots, t_n)$ is in $\mathfrak{C}^{\bar{?}}.S$, and $t = t_j, t' = t_{j'}$ both embed into $f(t_1, \dots, t_n)$.

Case (b) reduces to (a), since if $S_{\square_{ij}} \neq \square$, then $\mathfrak{C}^{\bar{?}}.S$ defines the same language as $\mathfrak{C}^{\bar{?}}.(S + S_{\square_{ij}})$.

In case (c), every non-empty A_i is of the form $f(\dots, \square, \dots)$ with a unique occurrence of \square . In that case, the language of $\mathfrak{C}^{\bar{?}}.S$ can be described more simply: it consists of those trees of the form $c_1[c_2[\dots c_k[s]\dots]]$, where $k \in \mathbb{N}$, each c_k is a context in the language of \mathfrak{C} , with just one occurrence of \square each, and s is a tree in $S \cup \text{args } \mathfrak{C}$. For short, say that a context c is in $\mathfrak{C}^{\bar{?}}$ if and only if it is of the form $c_1[c_2[\dots c_k[\square]\dots]]$, where $k \in \mathbb{N}$ and each c_k is in \mathfrak{C} . Such contexts have exactly one occurrence of \square . Hence the language of $\mathfrak{C}^{\bar{?}}.S$ consists of those trees $c[s]$ where c is in $\mathfrak{C}^{\bar{?}}$ and $s \in S \cup \text{args } \mathfrak{C}$. Given any two such trees $t = c[s]$ and $t' = c'[s']$, we find a tree $t'' \in \mathfrak{C}^{\bar{?}}.S$ in which both t and t' embed, as follows.

If both s and s' are in S , then by directedness there is an $s'' \in S$ such that $s, s' \preceq s''$: we define t'' as $c[c'[s'']]$.

If $s \in S$ and $s' \in \text{args } \mathfrak{C}$, then there is an $A_i = f_i(S_{\square_{i1}}, \dots, S_{\square_{in_i}})$ and a position j' , $1 \leq j' \leq n_i$, such that $S_{\square_{ij'}} \neq \square$ and the language of $S_{\square_{ij'}}$ contains s' . The unique position j at which $S_{\square_{ij}} = \square$ is different from j' . Let $u = f_i(u_1, \dots, u_n)$ where: $u_j = s, u_{j'} = s'$, for every position $k \neq j, j'$ u_k is an arbitrary tree from $S_{\square_{ik}}$. By construction, $u \in \mathfrak{C}^{\bar{?}}.S$, hence so is $t'' = c[c'[u]]$. Additionally, since s and s' both embed in u , t and t' both embed in t'' .

The same argument applies when $s \in \text{args } \mathcal{C}$ and $s' \in S$. Finally, we consider the case where s, s' are both in $\text{args } \mathcal{C}$. Then $s \in \text{args } A_i$ for some i , say $A_i = f_i(S_{\square_{i1}}, \dots, S_{\square_{in_i}})$, $S_{\square_{ij'}} \neq \square$ and s is in the language of $S_{\square_{ij'}}$. Since S is irreducible, it is non-empty, hence we can pick a tree s_0 from it. Let $u_{j'} = s$, $u_j = s_0$ where j is the unique position where $S_{\square_{ij}} = \square$, and pick u_k from the non-empty language $S_{\square_{ik}}$ for every $k \neq j, j'$; define $u = f_i(u_1, \dots, u_{n_i})$, a tree from $\mathcal{C}^*.S$ in which s embeds. Similarly, since $s' \in \text{args } \mathcal{C}$, s' is in the support of some $A_{i'}$, say $A_{i'} = f_{i'}(S_{\square_{i'1}}, \dots, S_{\square_{i'n_{i'}}$), $S_{\square_{i'j''}} \neq \square$ and s' is in the language of $S_{\square_{i'j''}}$. Let $v_{j''} = s'$ (instead of s in our previous step), $v_{j''} = u$ (instead of s_0) where j'' is the unique position where $S_{\square_{i'j''}} = \square$, and pick v_k from the non-empty language $S_{\square_{i'k}}$ for every $k \neq j'', j'''$. The tree $v = f_{i'}(v_1, \dots, v_{n_{i'}})$ is again in $\mathcal{C}^*.S$, and now both s and s' embed into it. Finally, we define t'' as $c[c'[v]]$. \square

Theorem 9.11 *The ideals, i.e., the irreducible closed subsets of $\mathcal{T}(\Sigma)$ are exactly the languages of tree-products.*

Proof. One direction is Lemma 9.6. In the other direction, we show that the language of every tree-product P is directed by structural induction on P , using Lemma 9.9 or Lemma 9.10, depending on its shape, as given by Lemma 9.8. In doing the proof, one needs to observe that any \rightarrow_1 -normal STRE $S = P_1 + \dots + P_m$ (where the language of each P_i is an ideal by induction hypothesis) has an empty language if and only if $m = 0$ —because ideals are never empty. \square

Exceptionally, let us dispense with the traditional shape of our S-representation theorem, and let us just state the following.

Theorem 9.12 (S-Representation, Finite Ranked Trees) *Let Σ be a finite signature. There is an S-representation $(S, \llbracket _ \rrbracket, \sqsubseteq, \tau, \wedge)$ of $\mathcal{T}(\Sigma)$, where S is the collection of all tree-products over Σ .*

Proof. Tree-products are regular tree languages. Hence inclusion (\sqsubseteq) is decidable [Thatcher and Wright, 1968, Section 2], see also [Comon et al., 2004]. Since $\mathcal{T}(\Sigma)$ is equal to the language of $(\sum_{f \in \Sigma} f(\square, \dots, \square))^{\bar{x}}.0$ (where each $f \in \Sigma_r$ is applied to a list of r boxes), we obtain τ by normalizing with respect to \rightarrow_1 . To compute finite intersections (\wedge) of two tree-products, we first convert those tree-products to tree automata, compute their intersection, convert the result back to an STRE by the construction of Proposition 9.1, and normalize it by \rightarrow_1 to obtain its expression as a sum of tree-products. \square

The above procedures are not optimal, and notably the inclusion procedure takes exponential time. As in the word case (see Lemma 6.9), there is a polynomial time inclusion test, but it is complex, and its correctness proof is difficult. We shall study it—by necessity—in the general case of unranked trees over a Noetherian signature. (See Corollary 10.33.) We let the interested reader do the required modifications to adapt it to the finite, ranked case.

Anticipating on that general case, the ranked trees in $\mathcal{T}(\Sigma)$ embed into the set of all (unranked) trees over the finite set Σ (seen as a Noetherian space, with the discrete topology). Proposition 5.4 tells us that the sobrification of the former can be seen,

up to isomorphism, as a subspace of the sobrification of the latter. We leave it as an exercise to check that this embedding $S \mapsto S^\circ$ is defined by:

- if $S = P_1 + \cdots + P_m$, then $S^\circ = P_1^\circ + \cdots + P_m^\circ$;
- $(f^{\vec{t}}(P_1, \dots, P_n))^\circ = f^{\vec{t}}(P_1^{\circ?} \cdots P_n^{\circ?})$;
- $((\sum_{i=1}^m f_i(P_{\square i_1}, \dots, P_{\square i_n}))^{\bar{*}} \cdot S)^\circ = (\sum_{i=1}^m f_i(P_{\square i_1}^{\circ?} \cdots P_{\square i_n}^{\circ?}))^{\bar{*}} \cdot S^\circ$
- $\square^\circ = \square$.

Combining this with the polynomial time inclusion test we shall see in Definition 10.31, this provides us with a polynomial time inclusion test for the tree-products and STREs over Σ .

10 Completing Finite Trees: the General Case

For every set X , let $\mathcal{T}(X)$ denote the set of all (ground, first-order) terms built using function symbols from X . Function symbols are now unranked, and may be applied to arbitrarily long lists of arguments. Since lists can be seen as finite words, $\mathcal{T}(X) = X \times \mathcal{T}(X)^*$. The leaves of such terms are the constants $f()$, formed from a function symbol $f \in X$ and an empty list of arguments; we shall also simply write f instead of $f()$. In general, a term t will be written as $f(t_1, t_2, \dots, t_n)$. When we wish to stress that the list t_1, t_2, \dots, t_n is a word, we shall also write $f(t_1 t_2 \cdots t_n)$ or $f(\vec{t})$ for a word $\vec{t} \in \mathcal{T}(X)^*$.

Again, our terms are ground. We shall later use the notation $\mathcal{T}(X, \{\square\})$ to refer to the set of terms with exactly one variable \square . These are the terms of $\mathcal{T}(X \cup \{\square\})$ where \square is always applied to the empty list of arguments.

The *subterms* of a term t are defined inductively as usual: writing t as $f(t_1, \dots, t_n)$, they are t itself plus all subterms of t_i , $1 \leq i \leq n$.

Given any quasi-ordering \leq on X , the *embedding quasi-ordering* \preceq_{\leq} on $\mathcal{T}(X, \mathcal{V})$ is defined by induction on the sum of the sizes of the terms s, t , by $s \preceq_{\leq} t$ iff:

- either $t = g(t_1, \dots, t_n)$ and $s \preceq_{\leq} t_j$ for some j , $1 \leq j \leq n$;
- or $s = f(\vec{s}), t = g(\vec{t}), f \leq g$ and $\vec{s} \preceq_{\leq}^* \vec{t}$;

where we recall that \preceq_{\leq}^* is the embedding quasi-ordering on words, understanding that the letters (which are terms themselves here) are quasi-ordered by \preceq_{\leq} (this is a recursive definition). As with several other notions here, we reuse freely some notations and some notions that we had introduced in the special case considered in Section 9.

Kruskal's Tree Theorem—in a more complete form than stated earlier—states that \preceq_{\leq} is wqo on $\mathcal{T}(X)$ iff \leq is wqo on X .

The latter extends to a topological setting: for a topological space X , $\mathcal{T}(X)$ is Noetherian iff X is Noetherian [Goubault-Larrecq, 2013, Theorem 9.7.46]. For this to make sense, we need to put a topology on $\mathcal{T}(X)$, and this is the *tree topology* [Goubault-Larrecq, 2013, Definition 9.7.39], defined as follows. For short, let $Y = \mathcal{T}(X)$. The *simple tree expressions* on X are given by the grammar $\pi ::= \diamond U(\pi_1 \mid$

$\dots | \pi_n$) where U is open in X , and $n \in \mathbb{N}$. (The base case is obtained when $n = 0$.) Such a simple tree expression denotes the set of terms t that have a subterm of the form $f(\vec{t})$ with $f \in U$ and $\vec{t} \in Y^* \pi_1 Y^* \dots Y^* \pi_n Y^*$. The simple tree expressions generate a topology, called the tree topology.

Here are a few basic facts about the tree topology. These can be found in [Goubault-Larrecq, 2013, Exercise 9.7.40, Exercise 9.7.43]. The proof of the following proposition, and of most subsequent results, can be found in Appendix E.

Proposition 10.1 *Let X be a topological space. Every finite intersection of simple tree expressions can be rewritten as a finite union of simple tree expressions. In particular, the simple tree expressions form a base of the tree topology.*

Letting \leq be the specialization quasi-ordering of X , the specialization quasi-ordering of $\mathcal{T}(X)$ is the embedding quasi-ordering \preceq_{\leq} .

The reader might be under the impression that the tree topology is far removed from the embedding quasi-ordering \preceq_{\leq} . Not so: the situation is exactly as for words (Proposition 7.2), and when X is a poset, then the tree topology is the Alexandroff topology of \preceq_{\leq} :

Proposition 10.2 *Let X be a set quasi-ordered by \leq . The tree topology on $\mathcal{T}(X_a)$ is exactly the Alexandroff topology of \preceq_{\leq} on $\mathcal{T}(X)$.*

Proof. This is the first part of Exercise 9.7.48 of [Goubault-Larrecq, 2013]. We give a proof just before Section E.1 in Appendix E. \square

This means, as for most other cases studied in this paper (except the case of rings, of words under the prefix ordering, and of powersets), that in the familiar case where X is a poset, the completion $\widehat{\mathcal{T}(X)}$ is both the sobrification of $\mathcal{T}(X)$ with the tree topology, and the ideal completion of $\mathcal{T}(X)$ with the \preceq_{\leq} quasi-ordering. (Recall Hoffmann's Theorem that for a poset Y , $\mathcal{S}(Y_a) = \mathbf{I}(Y)$ [Hoffmann, 1979b].) The present section is therefore merely an extension of Section 9 to a more general format of trees, and more general spaces of function symbols.

10.1 Tree Steps

To characterize $\mathcal{S}(X)$, we rely again on specific forms of regular tree expressions, this time for terms. We start with regular expressions based on the $\vec{?}$ operator, which we call *tree steps*.

Definition 10.3 (Tree Step) *Let X be a topological space.*

For every word-product \vec{P} on $\mathcal{T}(X)$, the support $\text{supp } \vec{P}$ of \vec{P} is defined as the set of terms t such that the one-element sequence t is in the language of \vec{P} . Equivalently, when $\vec{P} = e_1 e_2 \dots e_n$, $\text{supp } \vec{P} = \bigcup_{i=1}^n \text{supp } e_i$, where $\text{supp } S^ = S$ and $\text{supp } P^? = P$.*

For every closed subset F of X , let $F^{\vec{?}}(\vec{P})$ denote the union of $\text{supp } \vec{P}$ with the set of all terms of the form $f(\vec{t})$, $f \in F$, $\vec{t} \in \vec{P}$.

The tree steps are the expressions of the form $C^{\vec{?}}(\vec{P})$ where C is irreducible closed in X and \vec{P} is a word-product on $\mathcal{T}(X)$.

E.g., when $C = \{f\}$, $\vec{P} = \{a\}^? \{b\}^?$, then $C^{\vec{P}}$ is the set of all terms $f(a, b)$, $f(a)$, $f(b)$, $f()$, but also the terms a and b from $\text{supp } \vec{P}$.

Lemma 10.4 *Let X be a topological space. For every closed subset F of X , and every word-product \vec{P} on $\mathcal{T}(X)$, $\text{supp } \vec{P}$ and $F^{\vec{P}}$ are closed in $\mathcal{T}(X)$. If moreover $F = C$ is irreducible, then so is the tree step $C^{\vec{P}}$.*

Proof. See Appendix E, as for most other results of this section. \square

10.2 Tree Iterators

We turn to the needed generalization of tree regular expressions of the form $\mathfrak{C}^*.S$. Recall that $\mathcal{T}(X, \{\square\})$ is the set of terms of $\mathcal{T}(X \cup \{\square\})$ where \square is always applied to the empty list of arguments— \square acts as a hole, meant to be replaced by terms.

Definition 10.5 (Context) *Let X be a topological space, and \square be an element called the hole, and assumed not to be in X . A context is a term of $\mathcal{T}(X, \{\square\})$.*

Given any context \mathcal{C} , and any subset S of $\mathcal{T}(X)$, $\mathcal{C}[S]$ denotes the set of all terms in $\mathcal{T}(X)$ obtained by replacing each occurrence of \square in \mathcal{C} by (possibly different) terms from S .

The hole \square can be replaced by different terms. E.g., when $S = \{a, b\}$, $f(\square, \square)[S]$ denotes $\{f(a, a), f(a, b), f(b, a), f(b, b)\}$. Notice that terms without the hole are also considered as contexts: take $\mathcal{C} = f(c)$ for example, then $\mathcal{C}[S]$ contains just $f(c)$, for any subset S .

By extension, where t is a single term, $\mathcal{C}[t]$ denotes the unique term obtained by replacing every hole \square in \mathcal{C} by t . One should beware that $\mathcal{C}[S]$ is *not* in general equal to the set of all terms $\mathcal{C}[t]$, $t \in S$, as the example $S = \{a, b\}$, $\mathcal{C} = f(\square, \square)$ demonstrates.

Given two contexts \mathcal{C} and \mathcal{C}' , the notation $\mathcal{C}[\mathcal{C}'[S]]$ makes sense: this is $\mathcal{C}[S']$, where $S' = \mathcal{C}'[S]$. One can also read this as $\mathcal{C}[\mathcal{C}'][S]$, where $\mathcal{C}[\mathcal{C}']$ is the context obtained by replacing each occurrence of \square in \mathcal{C} by the (same) context \mathcal{C}' .

Given a single hole \square , we equip $\{\square\}$ with the only possible topology, and write $\mathcal{T}(X) + \{\square\}$ for the topological coproduct. Its open subsets are the sets of the form U or $U \cup \{\square\}$, where U is open in $\mathcal{T}(X)$, and similarly for closed subsets. Its *irreducible* closed subsets are the irreducible closed subsets of $\mathcal{T}(X)$, and $\{\square\}$.

Definition 10.6 (Tree Iterators) *Let X be a topological space, and \square be a distinguished element outside X , called the hole.*

The tree iterators are formal expressions of the form $\mathfrak{C}^.S$, where \mathfrak{C} is a closed subset of $X \times (\mathcal{T}(X) + \{\square\})^*$, and S is a closed subset of $\mathcal{T}(X)$. We equate pairs (f, \vec{u}) in \mathfrak{C} with the contexts $c = f(\vec{u})$ —such contexts are elementary contexts, in that \square can only occur directly under f .*

Call argument support $\text{args } \mathfrak{C}$ of \mathfrak{C} the set of all terms $t \in \mathcal{T}(X)$ such that $f(t) \in \mathfrak{C}$ for some $f \in X$. (Equivalently, such that some term of the form $f(\dots, t, \dots)$ belongs to \mathfrak{C} .) Then $\mathfrak{C}^.S$ denotes the smallest set of terms in $\mathcal{T}(X)$ such that:*

- 1) every term in S is in $\mathfrak{C}^*.S$;

2) every term in $\text{args } \mathfrak{C}$ is in $\mathfrak{C}^{\bar{\cdot}}.S$;

3) for every elementary context $c \in \mathfrak{C}$, $c[\mathfrak{C}^{\bar{\cdot}}.S]$ is included in $\mathfrak{C}^{\bar{\cdot}}.S$.

For example, let $\mathfrak{C} = \{f\} \times \{\square\}^*$. Its argument support is the set of all terms $t \in \mathcal{T}(X)$ such that $t \in \{\square\}^*$, i.e., $t = \square$: this is impossible since \square is not a term of $\mathcal{T}(X)$. So $\text{args } \mathfrak{C}$ is empty. Let $S = \{a, b\}$. $\mathfrak{C}^{\bar{\cdot}}.S$ is the following set of terms. First, there are the terms from S , namely a and b . Then, there are the terms obtained from the latter by applying the elementary contexts $c \in \mathfrak{C}$ to the above terms. These contexts are $f()$, $f(\square)$, $f(\square\square)$, etc. So we obtain the terms f , $f(a)$, $f(b)$, $f(a, a)$, $f(a, b)$, $f(b, a)$, $f(b, b)$, etc. In a third step, we find the terms obtained by applying the contexts $f()$, $f(\square)$, $f(\square\square)$, etc., to the latter terms: we obtain $f(f)$, $f(f(a))$, $f(f(b))$, $f(f(a, a))$, \dots , $f(a, f(a))$, etc. Continuing this way, we realize that $\mathfrak{C}^{\bar{\cdot}}.S$ is the set of terms whose non-leaf nodes are labeled with f , and whose leaves are labeled by a or $b \dots$ or f .

For a more complex example, take $\mathfrak{C} = \{f\} \times \{c\}^? \{\square\}^*$. Now $\text{args } \mathfrak{C} = \{c\} - c$ is the only part of $\{c\}^? \{\square\}^*$ that does not contain the hole \square . Take $S = \{a, b\}$ again. Then $\mathfrak{C}^{\bar{\cdot}}.S$ is the following set of terms. First, we find the terms from S , namely a and b . Second, we find the terms from $\text{args } \mathfrak{C}$, that is, c . Third, we find the results of applying contexts of the form $f()$, $f(c)$, $f(\square)$, $f(c\square)$, $f(\square\square)$, $f(c\square\square)$, $f(\square\square\square)$, etc., to the above terms: these are the terms of the form $f(t_1, t_2, \dots, t_n)$ where each t_i is in $\{a, b, c\}$. Continuing this way, we obtain that $\mathfrak{C}^{\bar{\cdot}}.S$ is the set of all terms built using f , a , b , c , where a , b , c are applied to no argument. In particular, $\mathfrak{C}^{\bar{\cdot}}.S$ is the same set as $\mathfrak{C}'^{\bar{\cdot}}.S'$, where $\mathfrak{C}' = \{f\} \times \{\square\}^*$, and $S' = \{a, b, c\}$.

Remark 10.7 *One should beware that $\mathfrak{C}^{\bar{\cdot}}.S$ can be non-empty even when S is empty, and even when $\text{args } \mathfrak{C}$ is empty. For example, for $\mathfrak{C} = \{f\} \times \{\square\}^*$, $\mathfrak{C}^{\bar{\cdot}}.\emptyset$ is the set of terms whose only function symbol is f , at all positions: $f()$, $f(f())$, $f(f(), f())$, \dots . This unexpected kind of non-empty STRE will be used in case 3 of Lemma 10.11, and in the definition of τ' in Theorem 10.36, notably.*

The argument support $\text{args } \mathfrak{C}$ mimics the eponymous notion of Section 9. While we do not commit yet to a specific syntax for closed subsets \mathfrak{C} of $X \times (\mathcal{T}(X) + \{\square\})^*$, the case that will be of most interest to us—and which will be the general case when X is Noetherian—is when \mathfrak{C} is of the form $\bigcup_{i=1}^m C_i \times Q_i$, where C_i is irreducible closed in X and each Q_i is a word-product over $\mathcal{T}(X) + \{\square\}$ for each i , $1 \leq i \leq m$. In that case, $\text{args } \mathfrak{C}$ simplifies:

Lemma 10.8 *Let $\mathfrak{C} = \bigcup_{i=1}^m C_i \times Q_i$, where C_i is a closed non-empty subset of X and each Q_i is a word-product over $\mathcal{T}(X) + \{\square\}$ for each i , $1 \leq i \leq m$. Then $\text{args } \mathfrak{C} = \bigcup_{i=1}^m \text{supp } Q_i \cap \mathcal{T}(X)$.*

Each Q_i is closed in $\mathcal{T}(X) + \{\square\}$. In particular, $\text{args } \mathfrak{C}$ is closed in $\mathcal{T}(X)$.

Proof. Let $t \in \text{args } \mathfrak{C}$. For some i , $1 \leq i \leq m$, there is a term $f(t)$ with $f \in C_i$ and such that the one-element word t is in Q_i . The latter means that $t \in \text{supp } Q_i$. Also, $t \in \mathcal{T}(X)$ by definition. Conversely, any element of $\text{supp } Q_i \cap \mathcal{T}(X)$ is a term (not the hole) t in $\text{supp } Q_i$, i.e., such that the one-element word t is in Q_i . Since C_i is non-empty, pick f from C_i . Then $f(t)$ is in \mathfrak{C} , so t is in $\text{args } \mathfrak{C}$.

For any space Y , the function $i: Y \rightarrow Y^*$ that maps any $y \in Y$ to the one-letter word y is continuous (Lemma B.1 in the Appendix). Since $\text{supp } Q_i = i^{-1}(Q_i)$, it is closed in $Y = \mathcal{T}(X) + \{\square\}$. It follows that $\text{supp } Q_i \cap \mathcal{T}(X)$ is closed in $\mathcal{T}(X)$, hence also $\text{args } \mathcal{C}$. \square

Lemma 10.9 *Let X be a topological space, and \square be a hole outside $\mathcal{T}(X)$. Every tree iterator $\mathcal{C}^{\bar{\cdot}}.S$ such that $\text{args } \mathcal{C}$ is closed in $\mathcal{T}(X)$ denotes a closed subset of $\mathcal{T}(X)$.*

The cases where tree iterators are irreducible require an analysis of the number of holes that can occur in each context, which parallels the analysis we ran in Lemma 9.8, leading to three cases (a), (b) and (c).

Definition 10.10 *Let X be a topological space, and \square be a hole outside $\mathcal{T}(X)$. Let \mathcal{C} be a subset of $X \times (\mathcal{T}(X) + \{\square\})^*$. \mathcal{C} is \square -linear iff its elements have at most one occurrence of \square each, i.e., for every elementary context $f(\vec{t}) \in \mathcal{C}$, at most one element of \vec{t} equals \square . \mathcal{C} is \square -generated iff for every $f(\vec{u}) \in \mathcal{C}$ such that \square does not occur in \vec{u} , one can split \vec{u} as a concatenation $\vec{u}_1\vec{u}_2$ so that $f(\vec{u}_1\square\vec{u}_2)$ is in \mathcal{C} .*

The \square -generated closed sets \mathcal{C} are those such that every element in \mathcal{C} contains the hole \square , or has a larger element in \mathcal{C} that contains the hole. So, for example, $\mathcal{C} = \{f\} \times \{\square\}^*$ is \square -generated. Indeed, its elements are the pairs (f, \square^n) , $n \in \mathbb{N}$; when $n \geq 1$, this contains \square , and when $n = 0$, (f, ϵ) is below (f, \square) , which is inside \mathcal{C} and contains \square : i.e., we apply the above definition, picking $\vec{u}_1 = \vec{u}_2 = \epsilon$. \mathcal{C} is not \square -linear, since, say, $(f, \square\square)$ is in it, and contains two holes.

On the other hand, $\{f\} \times \{\square\}^?$ is both \square -linear and \square -generated.

The tree iterators $\mathcal{C}^{\bar{\cdot}}.S$ are easier to understand when \mathcal{C} is \square -linear, just as in Section 9. Let $\mathcal{C}^{\square*}$ be the set of all contexts of the form $c_1[c_2[\dots[c_k]\dots]]$, where $k \in \mathbb{N}$, and $c_1, c_2, \dots, c_k \in \mathcal{C}$. (When $k = 0$, this denotes \square .) Whenever \mathcal{C} is \square -linear, all these contexts have at most occurrence of \square . Then $\mathcal{C}^{\bar{\cdot}}.S$ is the set of all terms obtained from a context c in $\mathcal{C}^{\square*}$ by replacing the unique occurrence of the hole \square (if any) by a term from $S \cup \text{args } \mathcal{C}$.

Lemma 10.11 *Let X be a topological space, and \square be a hole outside $\mathcal{T}(X)$, \mathcal{C} be a closed subset of $X \times (\mathcal{T}(X) + \{\square\})^*$, S be a closed subset of $\mathcal{T}(X)$, and assume that $\text{args } \mathcal{C}$ is closed. Then the tree iterator $\mathcal{C}^{\bar{\cdot}}.S$ is irreducible in the following cases:*

- 1) if \mathcal{C} is non- \square -linear, and S is non-empty;
- 2) or if \mathcal{C} is \square -generated and \square -linear and S is irreducible;
- 3) or if \mathcal{C} is non-empty, \square -generated, and S is empty.

In case 3, $\mathcal{C}^{\bar{\cdot}}.\emptyset$ is in particular non-empty. That should not be a surprise, considering Remark 10.7.

Using these two forms of closed subsets—tree steps and tree iterators—one can express the complements of all simple tree expressions.

Lemma 10.12 *Let X be a topological space. The complement \mathcal{C}_π of the open subset denoted by the simple tree expression $\pi = \diamond U(\pi_1 \mid \pi_2 \mid \dots \mid \pi_n)$ is given by structural induction on π by:*

- $\mathcal{C}_\pi = ((F \times \{\square\}^*) \cup (X \times (\mathcal{C}_{\pi_1^* \{\square\}^?} \mathcal{C}_{\pi_2^* \{\square\}^?} \dots \{\square\}^? \mathcal{C}_{\pi_n^*})))^* . \emptyset$ if $n \geq 1$, where F is the complement of U in X ;
- if $n = 0$, then $\mathcal{C}_\pi = (F \times \{\square\}^*)^* . \emptyset$.

Lemma 10.12 can be made explicit, especially when $U = X$ and $n = 1$:

Lemma 10.13 *Let X be a topological space. The complement \mathcal{C}_π of the open subset denoted by the simple tree expression $\pi = \diamond U(\pi_1 \mid \pi_2 \mid \dots \mid \pi_n)$, is given by structural induction on π by:*

- if $U = X$ and $n = 0$, then \mathcal{C}_π is empty;
- if $U = X$ and $n = 1$, then \mathcal{C}_π is $X^?(\mathcal{C}_{\pi_1^*})$;
- if $U = X$ and $n \geq 2$, then \mathcal{C}_π is $(X \times (\mathcal{C}_{\pi_1^* \{\square\}^?} \mathcal{C}_{\pi_2^* \{\square\}^?} \dots \{\square\}^? \mathcal{C}_{\pi_n^*}))^* . \emptyset$;
- if $U \neq X$ and $n = 0$, then \mathcal{C}_π is $(F \times \{\square\}^*)^* . \emptyset$, where F is the complement of U ;
- if $U \neq X$ and $n = 1$, then \mathcal{C}_π is $((F \times \{\square\}^*)^* . X^?(\mathcal{C}_{\pi_1^*}))$, where F is the complement of U ;
- if $U \neq X$ and $n \geq 2$, then \mathcal{C}_π is $((F \times \{\square\}^*) \cup (X \times (\mathcal{C}_{\pi_1^* \{\square\}^?} \mathcal{C}_{\pi_2^* \{\square\}^?} \dots \{\square\}^? \mathcal{C}_{\pi_n^*})))^* . \emptyset$, where F is the complement of U .

Proof. The case $U = X$, $n = 0$ is clear. When $U = X$ and $n = 1$, Lemma 10.12 tells us that $\mathcal{C}_\pi = (X \times \mathcal{C}_{\pi_1^*})^* . \emptyset$. This is by definition the smallest set A of terms containing $\mathcal{C}_{\pi_1} = \text{supp}(X \times \mathcal{C}_{\pi_1^*})$ and such that for every elementary context $f(\vec{u}) \in X \times \mathcal{C}_{\pi_1^*}$, $f(\vec{u})$ (which is a term, i.e., \square does not occur in it) is in A . Thus $A = X^?(\mathcal{C}_{\pi_1^*})$. When $U \neq X$ and $n = 1$, by Lemma 10.12 $\mathcal{C}_\pi = ((F \times \{\square\}^*) \cup (X \times (\mathcal{C}_{\pi_1^*})))^* . \emptyset$. The argument support of $\mathcal{C} = (F \times \{\square\}^*) \cup (X \times (\mathcal{C}_{\pi_1^*}))$ is \mathcal{C}_{π_1} , so the elements of \mathcal{C}_π are those of \mathcal{C}_{π_1} , those of the form $f(\vec{t})$ where f is arbitrary and $\vec{t} \in \mathcal{C}_{\pi_1^*}$ (hence all the terms of $X^?(\mathcal{C}_{\pi_1^*})$), and those obtained from the latter by applying any number of function symbols from F . The other cases follow directly from Lemma 10.12. \square

10.3 Checking Inclusion

We start with tree step inclusion.

Lemma 10.14 *Let X be a topological space, C and C' be two irreducible closed subsets of X , \vec{P} and \vec{P}' be two word-products over $\mathcal{T}(X)$. Then $C^{\vec{P}}(\vec{P}) \subseteq C'^{\vec{P}'}(\vec{P}')$ iff $C \subseteq C'$ and $\vec{P} \subseteq \vec{P}'$, or $C^{\vec{P}}(\vec{P}) \subseteq \text{supp } \vec{P}'$.*

We turn to the cases where one of the closed set to compare is of the form $\mathcal{C}^* . S$. For every closed subset \mathcal{C} of $X \times (\mathcal{T}(X) + \{\square\})^*$ and every closed subset S of $\mathcal{T}(X)$, we write $\mathcal{C}[S]$ for the set of pairs (f, \vec{t}) , where (f, \vec{u}) ranges over \mathcal{C} and where \vec{t} is obtained from \vec{u} by replacing each occurrence of \square by possible different terms from S .

Lemma 10.15 *Let X be a topological space, C be an irreducible closed subset of X , \vec{P} be a word-product over $\mathcal{T}(X)$, \mathfrak{C} be a closed subset of $X \times (\mathcal{T}(X) + \{\square\})^*$ such that $\text{args } \mathfrak{C}$ is closed in $\mathcal{T}(X)$, where \square is a hole outside $\mathcal{T}(X)$, and S be a closed subset of $\mathcal{T}(X)$.*

Then $C^{\vec{P}}(\vec{P}) \subseteq \mathfrak{C}^.S$ iff $C \times \vec{P} \subseteq \mathfrak{C}[\mathfrak{C}^*.S]$ and $\text{supp } \vec{P} \subseteq \mathfrak{C}^*.S$, or $C^{\vec{P}}(\vec{P}) \subseteq \text{args } \mathfrak{C} \cup S$.*

The expression $\mathfrak{C}[\mathfrak{C}^*.S]$ in Lemma 10.15 is arguably not syntactically smaller than $\mathfrak{C}^*.S$, and this would cause some problems in designing an algorithm for inclusion testing.

We shall show below that we can replace $\mathfrak{C}[\mathfrak{C}^*.S]$ by $\mathfrak{C}[C]$ for any irreducible closed set C containing $\mathfrak{C}^*.S$. The one that will suit us best is a set $\mathcal{T}^\square(X)$, which we now define. While $\mathcal{T}^\square(X)$ is semantically very large—larger than the set of all terms!—, one can think of it denoted by some specific piece of syntax that one would naturally call the *wildcard*.

Definition 10.16 (Wildcard $\mathcal{T}^\square(X)$) *Let $\mathcal{T}^\square(X)$ be the disjoint union of $\mathcal{T}(X)$ with a fresh element \square . Its topology is described by letting the closed subsets of $\mathcal{T}^\square(X)$ be those of $\mathcal{T}(X)$, plus $\mathcal{T}^\square(X)$ itself.*

That is, $\mathcal{T}^\square(X)$ is obtained from $\mathcal{T}(X)$ by adding a new top element to it. Notice that the whole of $\mathcal{T}^\square(X)$ is then irreducible closed in $\mathcal{T}^\square(X)$.

$\mathcal{T}^\square(X)$ is different from $\mathcal{T}(X) + \{\square\}$. While both spaces have the same elements, the topology of $\mathcal{T}^\square(X)$ is strictly coarser than that of $\mathcal{T}(X) + \{\square\}$: the only closed subset of $\mathcal{T}^\square(X)$ that contains \square is $\mathcal{T}^\square(X)$ itself, while any set $F \cup \{\square\}$, F closed in $\mathcal{T}(X)$, is closed in $\mathcal{T}(X) + \{\square\}$. In terms of opens, every non-empty open subset of $\mathcal{T}^\square(X)$ contains \square , while any set $U \cup \{\square\}$, with U open in $\mathcal{T}(X)$, is also open in $\mathcal{T}(X) + \{\square\}$.

In the following, $\mathfrak{C}[\mathcal{T}^\square(X)]$ is just what it looks like it should be: the set of contexts obtained by instantiating a context from \mathfrak{C} by replacing some (but not necessarily all) of its holes by terms.

Lemma 10.17 *Let X be a topological space, C be an irreducible closed subset of X , \vec{P} be a word-product over $\mathcal{T}(X)$, \mathfrak{C} be a closed subset of $X \times (\mathcal{T}(X) + \{\square\})^*$ such that $\text{args } \mathfrak{C}$ is closed in $\mathcal{T}(X)$, where \square is a hole outside $\mathcal{T}(X)$, and S be a closed subset of $\mathcal{T}(X)$.*

Then $C^{\vec{P}}(\vec{P}) \subseteq \mathfrak{C}^.S$ iff $C \times \vec{P} \subseteq \mathfrak{C}[\mathcal{T}^\square(X)]$ and $\text{supp } \vec{P} \subseteq \mathfrak{C}^*.S$, or $C^{\vec{P}}(\vec{P}) \subseteq \text{args } \mathfrak{C} \cup S$.*

Proof. Using Lemma 10.15, it is enough to show that if $C \times \vec{P} \subseteq \mathfrak{C}[\mathcal{T}^\square(X)]$ and $\text{supp } \vec{P} \subseteq \mathfrak{C}^*.S$, then $C \times \vec{P} \subseteq \mathfrak{C}[\mathfrak{C}^*.S]$. Indeed, let (f, \vec{t}) be any element of $C \times \vec{P}$. By assumption, $f(\vec{t})$ is in $\mathfrak{C}[\mathcal{T}^\square(X)]$. So there is an elementary context $f(\vec{u})$ in \mathfrak{C} such that \vec{t} is obtained from \vec{u} by replacing those u_j that are equal to \square by some terms t_j , $1 \leq j \leq m$. Since $t_j \in \text{supp } \vec{P}$, $t_j \in \mathfrak{C}^*.S$. So (f, \vec{t}) is in $\mathfrak{C}[\mathfrak{C}^*.S]$. \square

One may wonder above why we used $\mathcal{T}^\square(X)$ instead of, e.g., $\mathcal{T}(X)$. This would work, too. But using $\mathcal{T}^\square(X)$ will be what we shall need, in particular in the case where we need to compare two tree iterators.

Let us refine our understanding of the construction $\mathfrak{C}[S]$.

Lemma 10.18 *Let X be a Noetherian space, and S be a closed subset of $\mathcal{T}^\square(X)$. Let \mathfrak{C} be a closed subset of $X \times (\mathcal{T}(X) + \{\square\})^*$ of the form $\bigcup_{i=1}^m C_i \times Q_i$, where each C_i is irreducible closed in X and each Q_i is a word-product over $\mathcal{T}(X) + \{\square\}$.*

Then $\mathfrak{C}[S]$ equals $\bigcup_{i=1}^m C_i \times Q_i[S]$, where for each word-product $Q = e_1 e_2 \cdots e_n$, $Q[S] = e_1[S] e_2[S] \cdots e_n[S]$, and for each atomic expression e , $e[S]$ equals $S^?$ if $e = \{\square\}^?$, $I^?$ if $e = I^?$ for some irreducible closed subset I of $\mathcal{T}(X)$, $((\mathcal{F} \setminus \{\square\}) \cup S)^$ if $e = \mathcal{F}^*$ and $\square \in \mathcal{F}$, \mathcal{F}^* if $e = \mathcal{F}^*$ and $\square \notin \mathcal{F}$. \square*

That is, $\mathfrak{C}[S]$ is obtained from \mathfrak{C} by literally replacing \square by S throughout. In particular, when $S = \mathcal{T}^\square(X)$: $\{\square\}^?[\mathcal{T}^\square(X)] = \mathcal{T}^\square(X)^?$, $I^?[\mathcal{T}^\square(X)] = I^?$ when I is irreducible closed in $\mathcal{T}(X)$, $\mathcal{F}^*[\mathcal{T}^\square(X)] = \mathcal{F}^*$ when \mathcal{F} is closed in $\mathcal{T}(X)$, and $\mathcal{F}^*[\mathcal{T}^\square(X)] = \mathcal{T}^\square(X)^*$ when $\square \in \mathcal{F}$.

This allows us to give our final characterization of inclusion between tree steps and tree iterators, in a way that will lend itself more directly to a recursive algorithm.

Lemma 10.19 *Let X be a topological space, C be an irreducible closed subset of X , \vec{P} be a word-product over $\mathcal{T}(X)$, \mathfrak{C} be a closed subset of $X \times (\mathcal{T}(X) + \{\square\})^*$ of the form $\bigcup_{i=1}^m C_i \times Q_i$, where each C_i is irreducible closed in X and each Q_i is a word-product over $\mathcal{T}(X) + \{\square\}$, where \square is a hole outside $\mathcal{T}(X)$, and S be a closed subset of $\mathcal{T}(X)$.*

Then $C^{\vec{P}}(\vec{P}) \subseteq \mathfrak{C}^{\bar{S}}$ iff either: $\text{supp } \vec{P} \subseteq \mathfrak{C}^{\bar{S}}$, $C \subseteq C_i$ and $\vec{P} \subseteq Q_i[\mathcal{T}^\square(X)]$ for some i , $1 \leq i \leq m$; or $C^{\vec{P}}(\vec{P}) \subseteq \text{args } \mathfrak{C} \cup S$.

Proof. Lemma 10.17 applies since $\text{args } \mathfrak{C} = \bigcup_{i=1}^m \text{supp } Q_i \cap \mathcal{T}(X)$ is closed. Then \vec{P} is irreducible closed (in $\mathcal{T}^\square(X)^*$, not just in $\mathcal{T}(X)^*$) by Lemma 6.7. So $C \times \vec{P}$ is irreducible closed in $X \times \mathcal{T}^\square(X)^*$, whence $C \times \vec{P} \subseteq \mathfrak{C}[\mathcal{T}^\square(X)] = \bigcup_{i=1}^m C_i \times Q_i[\mathcal{T}^\square(X)]$ iff $C \times \vec{P} \subseteq C_i \times Q_i[\mathcal{T}^\square(X)]$ for some i , $1 \leq i \leq m$, iff $C \subseteq C_i$ and $\vec{P} \subseteq Q_i[\mathcal{T}^\square(X)]$ for some i , $1 \leq i \leq m$. \square

Let us now turn to the third case, where we compare tree iterators and tree steps in the other order.

Lemma 10.20 *Let X be a topological space, C be an irreducible closed subset of X , \vec{P} be a word-product over $\mathcal{T}(X)$, \mathfrak{C} be a closed subset of $X \times (\mathcal{T}(X) + \{\square\})^*$ such that $\text{args } \mathfrak{C}$ is closed in $\mathcal{T}(X)$, where \square is a hole outside $\mathcal{T}(X)$, and S be a closed subset of $\mathcal{T}(X)$. Assume also that there is an elementary context $f(\vec{u})$ in \mathfrak{C} such that \square occurs as some element of \vec{u} .*

Then $\mathfrak{C}^{\bar{S}} \subseteq C^{\vec{P}}(\vec{P})$ iff $\mathfrak{C}^{\bar{S}} \subseteq \text{supp } \vec{P}$.

Proof. Assume $\mathfrak{C}^{\bar{S}} \subseteq C^{\vec{P}}(\vec{P})$. By assumption, there is an elementary context $f(\vec{u})$ in \mathfrak{C} such that \square occurs as some element of \vec{u} . To make things simpler, observe that this implies that $f(\square)$ is in \mathfrak{C} . For every $t \in \mathfrak{C}^{\bar{S}}$, $f(t)$ is then again in $\mathfrak{C}^{\bar{S}}$. So $f(t)$ is in $C^{\vec{P}}(\vec{P})$. It follows that either $f \in C$ and $t \in \text{supp } \vec{P}$, or $f(t) \in \text{supp } \vec{P}$. Since $\text{supp } \vec{P}$ is closed, it is downwards-closed in \preceq , so if $f(t) \in \text{supp } \vec{P}$, then $t \in \text{supp } \vec{P}$. In any case, $t \in \text{supp } \vec{P}$. Since t is arbitrary, $\mathfrak{C}^{\bar{S}} \subseteq \text{supp } \vec{P}$. The converse inclusion is obvious. \square

Note that the condition that there is an elementary context $f(\vec{u})$ in \mathfrak{C} such that \square occurs as some element of \vec{u} is satisfied in all of the cases 1–3 where we proved $\mathfrak{C}^{\bar{\cdot}}.S$ to be irreducible (Lemma 10.11), since \mathfrak{C} is not \square -linear in the first case, and \square -generated in the remaining cases.

Finally, we deal with the case where we try to compare two tree iterators. We do this in several steps, and start, as above, with a lemma with no obvious algorithmic content, but which gives the basic characterization we need. The proof of this heavily depends on irreducibility, as usual. Let $@$ be the application map from $X \times \mathcal{T}(X)^*$ to $\mathcal{T}(X)$. This sends (f, \vec{t}) to $f(\vec{t})$, and is continuous (see Exercise 9.7.47 of [Goubault-Larrecq, 2013], or Lemma E.6 in Appendix E).

Lemma 10.21 *Let X be a topological space, \mathfrak{C} and \mathfrak{C}' be two closed subsets of $X \times (\mathcal{T}(X) + \{\square\})^*$ such that $\text{args } \mathfrak{C}$ and $\text{args } \mathfrak{C}'$ are closed in $\mathcal{T}(X)$, where \square is a hole outside $\mathcal{T}(X)$, and let S, S' be two closed subsets of $\mathcal{T}(X)$.*

Then $\mathfrak{C}^{\bar{\cdot}}.S \subseteq \mathfrak{C}'^{\bar{\cdot}}.S'$ iff $\mathfrak{C}[\mathfrak{C}^{\bar{\cdot}}.S] \subseteq @^{-1}(\text{args } \mathfrak{C}' \cup S') \cup \mathfrak{C}'[\mathcal{T}^{\square}(X)]$ and $\text{args } \mathfrak{C} \cup S \subseteq \mathfrak{C}'^{\bar{\cdot}}.S'$.

Again, we need to make the above lemma clearer, in the case that is of primary interest to us.

Lemma 10.22 *Let X be a topological space, \mathfrak{C} and \mathfrak{C}' be two closed subsets of $X \times (\mathcal{T}(X) + \{\square\})^*$, where \square is a hole outside $\mathcal{T}(X)$, and let S, S' be two closed subsets of $\mathcal{T}(X)$. Assume also that \mathfrak{C} is of the form $\bigcup_{i=1}^m C_i \times Q_i$, and that \mathfrak{C}' is of the form $\bigcup_{j=1}^n C'_j \times Q'_j$, where each C_i and each C'_j is irreducible closed in X , and Q_i and Q'_j are word-products over $\mathcal{T}(X) + \{\square\}$ for each $i, 1 \leq i \leq m$, and each $j, 1 \leq j \leq n$. Assume finally that $\mathfrak{C}^{\bar{\cdot}}.S$ is irreducible, and that $\square \in Q_i$ for every $i, 1 \leq i \leq m$.*

Then $\mathfrak{C}^{\bar{\cdot}}.S \subseteq \mathfrak{C}'^{\bar{\cdot}}.S'$ iff:

- *either $\mathfrak{C}^{\bar{\cdot}}.S \subseteq \text{args } \mathfrak{C}' \cup S'$,*
- *or $\text{args } \mathfrak{C} \cup S \subseteq \mathfrak{C}'^{\bar{\cdot}}.S'$, and for every $i, 1 \leq i \leq m$, there is a $j, 1 \leq j \leq n$, such that $C_i \subseteq C'_j$ and $Q_i[\mathcal{T}^{\square}(X)] \subseteq Q'_j[\mathcal{T}^{\square}(X)]$.*

Note that the assumptions that $\mathfrak{C}^{\bar{\cdot}}.S$ is irreducible, and that $\square \in Q_i$ for every $i, 1 \leq i \leq m$, are satisfied as soon as any of the cases 1–3 of Lemma 10.11 hold.

10.4 Intersections of Tree Steps and Tree Iterators

We now compute intersections, and we start with tree steps. All missing proofs are in Appendix E.6.

In the following lemma, recall that, by Lemma 6.13, the intersection of any two word-products \vec{P} and \vec{P}' (here, on $\mathcal{T}(X)$) can be expressed as a finite union of word-products \vec{P}''_j .

Lemma 10.23 *Let X be a topological space. The intersection of two tree steps $P = C^{\bar{\cdot}}(\vec{P})$ and $P' = C'^{\bar{\cdot}}(\vec{P}')$ is equal to $\bigcup_{j=1}^n (C \cap C')^{\bar{\cdot}}(\vec{P}''_j) \cup (\text{supp } \vec{P} \cap P') \cup (P \cap \text{supp } \vec{P}')$, where $\vec{P} \cap \vec{P}'$ is expressed as a finite union $\bigcup_{j=1}^n \vec{P}''_j$ of word-products on*

$\mathcal{T}(X)$. If $C \cap C'$ can be written as the union of finitely many irreducible closed subsets C_i , $1 \leq i \leq m$, then $P \cap P'$ is also equal to the union of the tree steps $C_i^{\vec{P}}(\vec{P}_j'')$ ($1 \leq i \leq m, 1 \leq j \leq m$), of $\text{supp } \vec{P} \cap P'$, and of $P \cap \text{supp } \vec{P}'$.

Lemma 10.24 *Let X be a Noetherian space, and S be a closed subset of $\mathcal{T}(X)$. Let $C^{\vec{P}}$ be a tree step, \mathfrak{C} be a closed subset of $X \times (\mathcal{T}(X) + \{\square\})^*$ of the form $\bigcup_{j=1}^n C_j \times Q_j$, where each C_j is irreducible closed in X and each Q_j is a word-product over $\mathcal{T}(X) + \{\square\}$.*

The intersection of the tree step $P = C^{\vec{P}}$ and of the tree iterator $P' = \mathfrak{C}^.S$ is the union of $\text{supp } \vec{P} \cap P'$, of $P \cap (S \cup \text{args } \mathfrak{C})$, and of $(C \cap C_j)^{\vec{P}}(\vec{P} \cap Q_j[P'])$, $1 \leq j \leq n$.*

If, for each j , one can write $C \cap C_j$ as the union of finitely many irreducible subsets C_{ij} , $1 \leq i \leq m_j$, and if $\vec{P} \cap Q_j[P']$ can be expressed as the union of finitely many word-products $\vec{P}_{\ell j}$, $1 \leq \ell \leq q_j$, then $P \cap P'$ is also equal to the union of $\text{supp } \vec{P} \cap P'$, of $P \cap (S \cup \text{args } \mathfrak{C})$, and of $C_{ij}^{\vec{P}}(\vec{P}_{\ell j})$, $1 \leq j \leq n, 1 \leq i \leq m_j, 1 \leq \ell \leq q_j$.

Lemma 10.26 below, which deals with intersections of tree iterators, is only valid if the word-products Q_i and Q'_j on $\mathcal{T}(X) + \{\square\}$ are normalized. A *normalized word-product* on $\mathcal{T}(X) + \{\square\}$ is of the form $e_1 e_2 \cdots e_n$, where each atomic expression e_i is of the form $P^?$ (with P irreducible closed in $\mathcal{T}(X)$), $\{\square\}^?$, F^* (with F closed in $\mathcal{T}(X)$), or $\{\square\}^*$. In other words, we forbid atomic expressions of the form $(F \cup \{\square\})^*$ where F is a non-empty closed subset of $\mathcal{T}(X)$. Note that the components of a normalized product are either closed subsets of $\mathcal{T}(X)$ (not containing \square) or just $\{\square\}$.

Lemma 10.25 *Let X be a Noetherian space, Q and Q' be two normalized word-products on $\mathcal{T}(X) + \{\square\}$, and P and P' be two closed subsets of $\mathcal{T}(X)$. The intersection $Q[P] \cap Q'[P']$ can be written as a finite union $\bigcup_{i=1}^n Q_i[P \cap P']$, where each Q_i is a normalized word-product over $\mathcal{T}(X) + \{\square\}$. Explicitly, the set $\{Q_i \mid 1 \leq i \leq n\}$ is obtained as $\text{Meet}^{\mathcal{E}}(Q, Q')$, where, for all components F of Q and F' of Q' , $\mathcal{E}(F, F')$ is a finite set of irreducible closed subsets whose union is:*

- $F \cap F'$ if $F, F' \neq \{\square\}$;
- $\{\square\}$ if $F = F' = \{\square\}$;
- $P \cap F'$ if $F = \{\square\}$ and $F' \neq \{\square\}$;
- $F \cap P'$ if $F \neq \{\square\}$ and $F' = \{\square\}$.

Moreover, for every i , $\text{supp } Q_i$ is included in $(\text{supp } Q \cap \text{supp } Q') \cup (\text{supp } Q \cap P') \cup (P \cap \text{supp } Q') \cup \{\square\}$.

Proof. Since X is Noetherian, $\mathcal{T}(X)$ is, too. By Lemma 6.13, $Q[P] \cap Q'[P']$ is equal to the union of the finitely many elements of $\text{Meet}^{\cap}(Q[P], Q'[P'])$.

For every set F that is either closed in $\mathcal{T}(X)$ or equal to $\{\square\}$, and every closed subset S of $\mathcal{T}(X)$, we write $F[S]$ for F if $F \neq \{\square\}$, for S otherwise. We observe that for all components F of Q and F' of Q' , $F[P] \cap F'[P']$ is the union of the elements

$$\begin{array}{ll}
S ::= 0 \mid P \mid S + S & P ::= C^{\bar{?}}(\vec{P}) \mid \mathfrak{C}^{\bar{*}}.S \\
& \vec{P} ::= e_1 e_2 \cdots e_n & e ::= P^? \mid S^* \\
\mathfrak{C} ::= 0 \mid A \mid \mathfrak{C} + \mathfrak{C} & A ::= C(Q) \\
& Q ::= \square^* \mid \vec{P}_1 \square^? \vec{P}_2 \square^? \cdots \square^? \vec{P}_m
\end{array}$$

Figure 5: STREs, tree products ($C \in \mathcal{S}(X)$, $n \geq 0$, $m \geq 1$)

$C[P \cap P']$, where C ranges over the elements of $\mathcal{E}(F, F')$: indeed both are equal to $F \cap F'$ in the first case defining \mathcal{E} , to $P \cap P'$ in the second case, to $P \cap F'$ in the third case and to $F \cap P'$ in the fourth case. It follows, by induction on the definition of $Meet$, that $Meet^\cap(Q[P], Q'[P'])$ is equal to (the union of the finitely many elements of) $Meet^\mathcal{E}(Q, Q')[P \cap P']$. It is also easy to check that $Meet^\mathcal{E}(Q, Q')$ consists of normalized word-products only, because \mathcal{E} only returns (sets of irreducible) closed subsets of $\mathcal{T}(X)$, or $\{\square\}$.

For the final part of the Lemma, $\text{supp } Q_i$ consists of unions of closed sets as returned by $\mathcal{E}(F, F')$ on components F of Q and F' of Q' , as inspection of the $Meet^\mathcal{E}$ procedure reveals. If $F, F' \neq \{\square\}$, then $\mathcal{E}(F, F') = F \cap F'$ is included in $\text{supp } Q \cap \text{supp } Q'$. If $F = F' = \{\square\}$, then $\mathcal{E}(F, F') = \{\square\}$. If $F = \{\square\}$ and $F' \neq \{\square\}$, then $\mathcal{E}(F, F') = P \cap F'$ is included in $P \cap \text{supp } Q'$. Finally, if $F \neq \{\square\}$ and $F' = \{\square\}$, then $\mathcal{E}(F, F') = F \cap P'$ is included in $\text{supp } Q \cap P'$. \square

Lemma 10.26 *Let X be a Noetherian space, S and S' be closed subsets of $\mathcal{T}(X)$. Let also \mathfrak{C} (resp., \mathfrak{C}') be a closed subset of $X \times (\mathcal{T}(X) + \{\square\})^*$ of the form $\bigcup_{i=1}^m C_i \times Q_i$ (resp., $\bigcup_{j=1}^n C'_j \times Q'_j$), where each C_i and each C'_j is irreducible closed in X and each Q_i and each Q'_j is a normalized word-product over $\mathcal{T}(X) + \{\square\}$. For all i, j , write $C_i \cap C_j$ as $\bigcup_{k=1}^{p_{ij}} C''_{ijk}$ where each C''_{ijk} is irreducible closed in X , and let $Q''_{ij\ell}$, $1 \leq \ell \leq q_{ij}$ enumerate the elements of $Meet^\mathcal{E}(Q_i, Q'_j)$, where the oracle \mathcal{E} is defined in Lemma 10.25.*

Then the intersection of the tree iterators $P = \mathfrak{C}^{\bar{}}.S$ and $P' = \mathfrak{C}'^{\bar{*}}.S'$ is the tree iterator $\mathfrak{C}''^{\bar{*}}.S''$, where $\mathfrak{C}'' = \bigcup_{i,j,k,\ell} C''_{ijk} \times Q''_{ij\ell}$ and where S'' is the union of $P \cap (\text{args } \mathfrak{C}' \cup S')$ and of $(\text{args } \mathfrak{C} \cup S) \cap P'$.*

10.5 STREs, Tree-Products

Assume X Noetherian. We now claim that the closed subsets of $\mathcal{T}(X)$ are exactly those denoted by simple tree regular expressions (STREs), and those that are irreducible are exactly those denoted by tree-products, which we now define.

The STREs S and the tree-products P are defined in Figure 5, with additional constraints that we describe in Requirement 10.28 below. We understand those up to associativity and commutativity for $+$, and the fact that 0 is neutral for $+$. Hence, for example, every STRE S can be written as a sum of finitely many tree products P_1, \dots, P_n , and those are unique up to permutation; in particular $S = 0$ if and only if $n = 0$.

Those expressions have the obvious semantics, once we understand 0 as the empty set, $+$ as union, and in productions such as Q , \square as the set $\{\square\}$. The semantics of $A = C(Q)$ is the product of the semantics of C and of Q . We have:

Lemma 10.27 *The languages of S , P are closed subsets of $\mathcal{T}(X)$, the languages of \vec{P} are closed subsets of $\mathcal{T}(X)^*$, the languages of \mathfrak{C} , A are closed subsets of $X \times (\mathcal{T}(X) + \{\square\})^*$, the languages of Q are closed subsets of $(\mathcal{T}(X) + \{\square\})^*$.*

Proof. By induction on syntax. If $P = C^{\vec{P}}$, then P is closed by Lemma 10.4. If $P = \mathfrak{C}^{\vec{S}}$, then P is closed by Lemma 10.9, which applies since args \mathfrak{C} is closed, due to Lemma 10.8. The case of S follows from the fact that finite unions of closed sets are closed. The case of \vec{P} follows from Corollary 6.6, and similarly for Q , noticing that $\{\square\}$ is closed in $\mathcal{T}(X) + \{\square\}$. The case of A is because products of closed sets are closed, and the case of \mathfrak{C} follows, again, from the fact that finite unions of closed sets are closed. \square

Requirement 10.28 *We say that $A = C(Q)$ is syntactically \square -linear if and only if Q is of the form $\vec{P}_1 \square^? \vec{P}_2 \square^? \dots \square^? \vec{P}_m$ with $m \leq 2$, and is syntactically \square -generated if and only if it is of this form with $m \geq 2$, or $Q = \square^*$.*

$\mathfrak{C} = \sum_{i=1}^m C_i(Q_i)$ is syntactically \square -linear (resp., generated) if and only if every $C_i(Q_i)$ is syntactically \square -linear (resp., generated).

We require every subexpression $\mathfrak{C}^{\vec{S}}$ to be such that \mathfrak{C} is syntactically \square -generated, $\mathfrak{C} \neq 0$, and one of the following conditions hold:

- 1) \mathfrak{C} is not syntactically \square -linear, and $S \neq 0$, or
- 2) \mathfrak{C} is syntactically \square -linear and S is a tree-product P ;
- 3) or $S = 0$.

Lemma 10.29 *The language of every tree-product is an irreducible closed subset of $\mathcal{T}(X)$.*

Proof. We first show that the language of every tree-product P is non-empty. If P is of the form $C^{\vec{P}}$, then its language contains $f()$, for any f in C . If P is of the form $\mathfrak{C}^{\vec{S}}$, where $\mathfrak{C} = C_1(Q_1) + \dots + C_n(Q_n)$, then since $\mathfrak{C} \neq 0$, n is non-zero. (Remember that we reason up to the fact that $+$ is associative and commutative, and 0 is neutral for $+$.) In that case, pick f from C_1 , and we note that $f()$ is in the language of $\mathfrak{C}^{\vec{S}}$.

It follows that: (*) for every STRE S such that $S \neq 0$, the language of S is non-empty.

We now prove the claim by induction on expressions. We need to show both that the language of every tree-product is irreducible closed, and that every expression of the form Q is an irreducible closed subset of $(\mathcal{T}(X) + \{\square\})^*$. The latter follows from Lemma 6.7 (every word-product is irreducible closed). The fact that every expression of the form $\mathfrak{C}^{\vec{S}}$ is irreducible closed follows from Lemma 10.11, once we observe that every non-syntactically \square -linear context \mathfrak{C} has a non- \square -linear language, and using (*) in case 1; and for the remaining cases, that every syntactically \square -linear context \mathfrak{C}

has a \square -linear language, and similarly for \square -generatedness. Every expression of the form $C^{\vec{P}}$ is irreducible closed by Lemma 10.4. \square

We shall see below that the converse holds.

An example of tree-products is given in the following result, which also serves to state how $\mathcal{T}(X)$ embeds into its completion $\widehat{\mathcal{T}(X)} = \mathcal{S}(\mathcal{T}(X))$.

Lemma 10.30 (Embedding) *Let X be a topological space. The closure $\eta_{\mathcal{T}(X)}^{\mathcal{S}}(t)$ of the term t in $\mathcal{T}(X)$ is defined by structural induction on t by: if $t = f(t_1, t_2, \dots, t_m)$, then $\eta_{\mathcal{T}(X)}^{\mathcal{S}}(t) = (\eta_X^{\mathcal{S}} f)^{\vec{P}} (\eta_{\mathcal{T}(X)}^{\mathcal{S}}(t_1))^{\vec{P}} (\eta_{\mathcal{T}(X)}^{\mathcal{S}}(t_2))^{\vec{P}} \cdots (\eta_{\mathcal{T}(X)}^{\mathcal{S}}(t_m))^{\vec{P}}$.*

Proof. By Lemma 10.4, $(\eta_X^{\mathcal{S}} f)^{\vec{P}} (\eta_{\mathcal{T}(X)}^{\mathcal{S}}(t_1))^{\vec{P}} (\eta_{\mathcal{T}(X)}^{\mathcal{S}}(t_2))^{\vec{P}} \cdots (\eta_{\mathcal{T}(X)}^{\mathcal{S}}(t_m))^{\vec{P}}$ is closed, since $\eta_{\mathcal{T}(X)}^{\mathcal{S}}(t_1), \eta_{\mathcal{T}(X)}^{\mathcal{S}}(t_2), \dots, \eta_{\mathcal{T}(X)}^{\mathcal{S}}(t_m)$ are closed by induction hypothesis. It also contains t , so it contains the closure of t . Conversely, it is easy to see that whenever $s \in (\eta_X^{\mathcal{S}} f)^{\vec{P}} (\eta_{\mathcal{T}(X)}^{\mathcal{S}}(t_1))^{\vec{P}} (\eta_{\mathcal{T}(X)}^{\mathcal{S}}(t_2))^{\vec{P}} \cdots (\eta_{\mathcal{T}(X)}^{\mathcal{S}}(t_m))^{\vec{P}}$, then $s \preceq t$, so s is in $\eta_{\mathcal{T}(X)}^{\mathcal{S}}(t)$. \square

We can define a syntactic inclusion test \leq as follows, following Lemma 10.14, Lemma 10.19, Lemma 10.20, and Lemma 10.22. We write semantic inclusion as \subseteq . The following notion of syntactic support is meant to mimic the semantical notion of support of Definition 10.3 in the syntax, while the notion of syntactic argument support mimics the argument support of Definition 10.6; accordingly, we use the same notations $\text{supp } \vec{P}$ and $\text{args } \mathfrak{C}$.

Definition 10.31 *Let the syntactic support of $\vec{P} = e_1 e_2 \cdots e_n$ be $\text{supp } \vec{P} = \sum_{i=1}^n \text{supp } e_i$, where $\text{supp } P^? = P$, $\text{supp } S^* = S$.*

Let $\text{supp}' Q$ be 0 if $Q = \square^$, $\sum_{i=1}^m \text{supp } \vec{P}_i$ if $Q = \vec{P}_1 \square^? \vec{P}_2 \square^? \cdots \square^? \vec{P}_m$. (This denotes the intersection of the support of Q with $\mathcal{T}(X)$.) The syntactic argument support \mathfrak{C} of the context $\mathfrak{C} = \sum_{i=1}^n C_i(Q_i)$ is $\sum_{i=1}^n \text{supp}' Q_i$.*

We define $S \leq S'$, $P \leq P'$, etc., by induction on the sum of the sizes of the two expressions involved, by:

- 1) for $S = \sum_{i=1}^m P_i$ and $S' = \sum_{j=1}^n P'_j$, $S \leq S'$ if and only if for every i , there is a j such that $P_i \leq P'_j$;
- 2) for $P = C^{\vec{P}}$ and $P' = C'^{\vec{P}'}$, $P \leq P'$ if and only if $C \subseteq C'$ and $\vec{P} \leq \vec{P}'$, or $P \leq \text{supp } \vec{P}'$;
- 3) for $P = C^{\vec{P}}$ and $P' = \mathfrak{C}' . S$ with $\mathfrak{C} = \sum_{i=1}^m C_i(Q_i)$, $P \leq P'$ if and only if $\text{supp } \vec{P} \leq P'$ and $C \subseteq C_i$, $\vec{P} \leq Q_i$ for some i , $1 \leq i \leq m$, or $P \leq \text{args } \mathfrak{C} + S$;
- 4) for $P = \mathfrak{C}' . S$ and $P' = C^{\vec{P}}$, $P \leq P'$ if and only if $P \leq \text{supp } \vec{P}$;
- 5) for $P = \mathfrak{C}' . S$ and $P' = \mathfrak{C}'' . S'$ with $\mathfrak{C} = \sum_{i=1}^m C_i(Q_i)$ and $\mathfrak{C}' = \sum_{j=1}^n C'_j(Q'_j)$, $P \leq P'$ if and only if $P \leq \text{args } \mathfrak{C}' + S'$, or $\text{args } \mathfrak{C} + S \leq P'$ and for every i there is a j such that $C_i \subseteq C'_j$ and $Q_i \leq Q'_j$;

- 6) $Q \leq Q'$ if and only if Q is less than Q' as a word-product (see Lemma 6.10), comparing letters by \leq , recursively, and considering \square as a letter above all others. (This case subsumes tests of the form $\vec{P} \leq Q$ as well, for instance, since any \vec{P} is a Q .)

One easily sees that $S \leq S'$ if and only if (the language of) S is included in S' , $P \leq P'$ if and only if P is included in P' , etc., by induction. One should pay attention to the fact that $Q \leq Q'$ is equivalent to the inclusion of $Q[\mathcal{T}^\square(X)]$ into $Q'[\mathcal{T}^\square(X)]$, as needed in cases 3 and 5 above. This is why the comparison of expressions of the form Q assumes that \square is a letter outside $\mathcal{T}(X)$, and above all elements of $\mathcal{T}(X)$ (see Item 6): indeed, \square denotes the whole set $\mathcal{T}^\square(X)$ there.

In case Item 6 of the definition is not clear enough, here is a complete explanation. This will also help understand how the conditions $\vec{P} \leq Q_i$ of Item 3 and $Q_i \leq Q_j$ of Item 5 are checked. We let $\epsilon \leq Q'$ for every Q' , $Q \leq \epsilon$ iff $Q = \epsilon$, and, if $Q = e_1 Q_1$ and $Q' = e'_1 Q'_1$, then $Q \leq Q'$ if and only if:

- 1) $e_1 \not\sqsubseteq e'_1$ and $Q \leq Q'_1$,
- 2) or e_1 is of the form $E^?$, e'_1 is of the form $E'^?$, $e_1 \sqsubseteq e'_1$ and $Q_1 \leq Q'_1$,
- 3) or e'_1 is of the form E'^* , $e_1 \sqsubseteq e'_1$ and $Q_1 \leq Q'$,
- 4) or $e_1 = \emptyset^*$ and $Q_1 \leq Q'$.

Additionally, \sqsubseteq is defined by: $P^? \sqsubseteq P'^?$ iff $P \leq P'$, $P^? \sqsubseteq \square^?$ and $\square^? \sqsubseteq \square^?$ are always true, $\square^? \sqsubseteq P'^?$ is always false; $S^* \sqsubseteq S'^*$ iff $S \leq S'$, $S^* \sqsubseteq \square^*$ and $\square^* \sqsubseteq \square^*$ are always true, $\square^* \sqsubseteq S'^*$ is always false; $P^? \sqsubseteq S^*$ iff $P \leq S$, $P^? \sqsubseteq \square^*$ and $\square^? \sqsubseteq \square^*$ are always true, $\square^? \sqsubseteq S^*$ is always false; $S^* \sqsubseteq P^?$ if and only if $S = \emptyset$; similarly $S^* \sqsubseteq \square^?$ if and only if $S = \emptyset$; and $\square^* \sqsubseteq P^?$, $S^* \sqsubseteq \square^?$ (with S non-empty), $\square^* \sqsubseteq \square^?$ are always false. One should be conscious that, for instance, $P^? \sqsubseteq \square^?$ is always true (and the converse inequality is false unless $P = \square$), even when P contains further subexpressions with occurrences of \square , e.g., $((\{f\} \times \{\square\}^*)^* . S)^? \sqsubseteq \square^?$.

Lemma 10.32 *Let X be a Noetherian space. Then the language of P is included in that of P' if and only if $P \leq P'$. Similarly for $S \leq S'$. \square*

Corollary 10.33 *Let X be a Noetherian space. Inclusion of word-products, resp. of STREs, on X , can be checked in polynomial time modulo an oracle testing inclusion of closed subsets of X .*

Proof. Here memoization [Michie, 1968] is easier to implement and analyze than dynamic programming. We write a recursive program `incl` that takes two word-products P and P' and returns true if $P \leq P'$, false otherwise. This program maintains a table A , initially empty, that maps pairs of word-products to Booleans. The program `incl`, applied to P and P' , first checks whether (P, P') is in A , and if so returns immediately the Boolean associated with it in A ; otherwise, it tests whether $P \leq P'$, depending on the shape of P and P' , using the corresponding case (1 through 6), as given in Definition 10.31; once this is done, it adds (P, P') to A , associated with the Boolean

value true if P was found to be below P' , false otherwise. Each case, 1 through 6, involves further inclusion tests between word-products, which are evaluated by calling `incl` recursively.

Instead of describing all cases, we will focus on case 3 below. It will be practical to consider another program `Incl`, defined in mutual recursion with `incl` by: `Incl(S, S')` is true, where $S = P_1 + \dots + P_m$ and $S' = P'_1 + \dots + P'_n$, if and only if for every $i \in \{1, \dots, m\}$, there is a $j \in \{1, \dots, n\}$ such that `incl(Pi, P'j)` returns true; `Incl` returns true on every pair whose second component is \square , and false on every pair (\square, S') where S' is an STRE.

In case 3, then P is of the form $C^{\vec{P}}(\vec{P})$ and P' is of the form $\mathfrak{C}^*.S$ with $\mathfrak{C} = \sum_{i=1}^m C_i(Q_i)$. Following the definition of case 3, `incl` does the following.

- 1) First, it tests whether $\text{supp } \vec{P} \leq P'$. We refer to Definition 10.3 for the definition of $\text{supp } \vec{P}$. Accordingly, writing \vec{P} as $e_1 e_2 \dots e_n$, testing whether $\text{supp } \vec{P} \leq P'$ means testing whether, for every $i \in \{1, \dots, n\}$:
 - $P'' \leq P'$, in case e_i is of the form $P''^?$;
 - or $P_j'' \leq P'$ for every $j \in \{1, \dots, m\}$, in case e_i is of the form S^* , where $S = P_1'' + \dots + P_m''$.

All the tests $P'' \leq P'$ (in the first case), and $P_j'' \leq P'$ (in the second case) are done by calling `incl` recursively.

- 2) If item 1 succeeded, `incl` then tests whether $C \subseteq C_i$ and $\vec{P} \leq Q_i$ for some i , $1 \leq i \leq m$. This is done by enumerating the indices $i \in \{1, \dots, m\}$, testing whether $C \subseteq C_i$ for each one (using the given oracle that tests inclusion of closed subsets of X) and whether $\vec{P} \leq Q_i$. For the latter, we use Item 6 of Definition 10.3: we need to compare \vec{P} and Q_i as word-products; this allows us to use the dynamic programming procedure given after Lemma 6.9, fed with `Incl` as an oracle. (You may wish to return to the explanation before Lemma 10.32 in order to better understand where recursive calls to `Incl` are involved; namely in the \leq tests involved in the definition of \sqsubseteq .)
- 3) If item 1 failed, `incl` instead tests whether $P \leq \text{args } \mathfrak{C} + S$. In other words, using the fact that P is irreducible, it tests whether $P \leq S$ (using `Incl`), or whether $P \leq \text{args } \mathfrak{C}$. For the latter, we use Lemma 10.8, and therefore we test whether $P \leq \text{supp } Q_i \cap \mathcal{T}(X)$ for some i , $1 \leq i \leq m$. Writing Q_i as $e_1 e_2 \dots e_n$, this means testing whether $P \leq \text{supp } e_j$ for some j , $1 \leq j \leq n$, such that $e_j \neq \square^?, \square^*$. Finally, if e_j is of the form $P''^?$, we test whether $P \leq \text{supp } e_j = P'$ by calling `incl` recursively on (P, P') , and if it is of the form S'^* , we test whether $P \leq \text{supp } e_j = S'$ by calling `Incl` on (P, S') .

All other cases are dealt with similarly. Up to a multiplicative constant, the time complexity a memoized procedure such as `incl` on input (P, P') is bounded by the product of two values: the first one is the number of pairs on which `incl` can be called recursively, and this is bounded by the product of the sizes of P and P' ; the second one is the time complexity of each case (such as case 3, explained above), including table

lookups and table updates on A , but counting the complexity of each recursive call to `incl` (including those obtained through an intermediate call to `Incl`) as one. Analyzing each case shows that the second value is also polynomial in the sizes of P and P' , whence the conclusion. \square

Lemma 10.34 *Let X be a Noetherian space. Given two tree-products P, P' , one can express their intersection as a finite sum of tree-products. If $(S, \llbracket _ \rrbracket, \preceq, \tau, \wedge)$ is an S -representation of X , then this intersection can be implemented as a map \wedge' modulo an oracle that implements \wedge .*

Proof. Let $|E|$ denote the number of subexpressions of E that are tree-products, for every kind of expression in Figure 5. Explicitly: $|0| = 0$, $|S_1 + S_2| = |S_1| + |S_2|$, $|C^{\bar{?}}(\vec{P})| = 1 + |\vec{P}|$, $|\mathfrak{C}^{\bar{*}}.S| = 1 + |\mathfrak{C}| + |S|$, $|e_1 e_2 \cdots e_n| = \sum_{i=1}^n |e_i|$, $|P^{\bar{?}}| = |P|$, $|S^*| = |S|$, $|\mathfrak{C}_1 + \mathfrak{C}_2| = |\mathfrak{C}_1| + |\mathfrak{C}_2|$, $|C(Q)| = |Q|$, $|\square^*| = 0$, $|\vec{P}_1 \square^{\bar{?}} \vec{P}_2 \square^{\bar{?}} \cdots \square^{\bar{?}} \vec{P}_m| = \sum_{j=1}^m |\vec{P}_j|$.

The fact that $P \cap P'$ can be expressed as a finite sum of tree-products is by induction on $|P| + |P'|$. When both are tree steps, say $P = C^{\bar{?}}(\vec{P})$ and $P' = C'^{\bar{?}}(\vec{P}')$, we use Lemma 10.23, using the subprocedure $Meet^{\mathcal{E}}$ of Lemma 6.13 to compute the intersection of the word-products \vec{P} and \vec{P}' as a finite union $\bigcup_{j=1}^n \vec{P}''_j$, where \mathcal{E} merely computes the intersection of a component of \vec{P} and of a component of \vec{P}' , by induction.

When P is a tree step $C^{\bar{?}}(\vec{P})$ and P' is a tree iterator $P' = \mathfrak{C}^{\bar{*}}.S$ (or conversely), we use Lemma 10.24 instead. Again, we use $Meet^{\mathcal{E}}$ in order to compute $\vec{P} \cap Q_j[P']$, where \mathcal{E} is intersection again. Note that the components of $Q_j[P']$ are either components of Q_j , which are finite sums of tree-products that are strictly smaller than P' , or equal to P' (at the positions where Q_j holds a $\square^{\bar{?}}$ or a \square^*). In any case, \mathcal{E} is only ever applied to pairs of word-products (P_0, P'_0) such that $|P_0| < |P|$ and $|P'_0| \leq |P'|$, so the induction hypothesis applies and $\mathcal{E}(P_0, P'_0)$ therefore computes $P_0 \cap P'_0$, as desired. The fact that the induction hypothesis really applies is probably clearer with an example. Imagine that $\vec{P} = P_1^{\bar{?}} S_1^*$, and $Q_j = P_2^{\bar{?}} \square^{\bar{?}} S_2^* \square^* P_3^{\bar{?}}$. Following Lemma 6.13, we compute $Meet^{\mathcal{E}}(\vec{P}, Q_j[P'])$ as follows. We use the first clause that defines it, and we are led (among other things) to compute $\mathcal{E}(P_1, P_2)$ —since $|P_1| < |P|$ and $|P_2| < |P'|$, the induction hypothesis applies—and $Meet^{\mathcal{E}}(S_1^*, (\square^{\bar{?}} S_2^* \square^* P_3^{\bar{?}})[P'])$. For the latter, we use the third clause, and we are led to compute $\mathcal{E}(S_1, P')$ —here $|S_1| < |P|$ and the size of the second argument is equal to $|P'|$, so the induction hypothesis applies again—and then $Meet^{\mathcal{E}}(S_1^*, (S_2^* \square^* P_3^{\bar{?}})[P'])$ and $Meet^{\mathcal{E}}(\epsilon, (\square^{\bar{?}} S_2^* \square^* P_3^{\bar{?}})[P'])$. The latter is trivial. For the former, we are led to compute $\mathcal{E}(S_1, S_2)$ (by induction hypothesis, since $|S_1| < |P|$ and $|S_2| < |P'|$) and then $Meet^{\mathcal{E}}(S_1^*, (\square^* P_3^{\bar{?}})[P'])$ and $Meet^{\mathcal{E}}(\epsilon, (S_2^* \square^* P_3^{\bar{?}})[P'])$. The latter is again trivial, so we look at the former. This requires us to compute $\mathcal{E}(S_1, P')$ —and the induction hypothesis applies again since $|S_1| < |P|$ and the size of the second argument is equal to $|P'|$ —and then $Meet^{\mathcal{E}}(S_1^*, (P_3^{\bar{?}})[P'])$ and $Meet^{\mathcal{E}}(\epsilon, (\square^* P_3^{\bar{?}})[P'])$. The last call to \mathcal{E} that we need has arguments S_1 and P_3 , with $|S_1| < |P|$ and $|P_3| < |P'|$.

Finally, when P is a tree iterator $\mathfrak{C}^{\bar{*}}.S$ and P' is a tree iterator $\mathfrak{C}'^{\bar{*}}.S'$, we use Lemma 10.26. (Note that the syntax of Figure 5 forces all word-products Q to be normalized, so that Lemma 10.26 indeed applies.) This involves calling $Meet^{\mathcal{E}}$ again,

where \mathcal{E} is now the oracle of Lemma 10.25. The latter is only ever applied to pairs of word-products (P_0, P'_0) such that $|P_0| < |P|$ and $|P'_0| \leq |P'|$, or such that $|P_0| \leq |P|$ and $|P'_0| < |P'|$, so that the induction hypothesis applies again.

However, the result $\mathfrak{C}''^{\bar{*}}.S''$ given by Lemma 10.26 may fail to satisfy Requirement 10.28. We repair this by using a rewriting process akin to the relation \rightarrow_1 of Figure 4. (This is very similar, except that (R11) does not have an equivalent here, since there is no such thing as a function of arity 0.) First, we split \mathfrak{C}'' into a sum $\mathfrak{C}''_0 + \mathfrak{C}''_1$, where \square does not occur at all in \mathfrak{C}''_0 (hence \mathfrak{C}''_0 is of the form $\sum_a C''_a(\vec{P}_a)$) and \square occurs in each summand of \mathfrak{C}''_1 (so \mathfrak{C}''_1 is syntactically \square -generated). We can then rewrite $\mathfrak{C}''^{\bar{*}}.S''$ as $\mathfrak{C}''_1^{\bar{*}}.(S'' + \sum_a C''_a^{\bar{*}}(\vec{P}_a))$ (mimicking rule (R6)). Modulo this rewrite, we can assume that we have expressed $P \cap P'$ as $\mathfrak{C}''^{\bar{*}}.S''$, where \mathfrak{C}'' is syntactically \square -generated. If $\mathfrak{C}'' = 0$, then we simplify that to S'' . Otherwise, if \mathfrak{C}'' is syntactically \square -linear, then $\mathfrak{C}''^{\bar{*}}.$ distributes over $+$ (as with rule (R12)): writing S'' as $\sum_b P''_b$, we return $\sum_b \mathfrak{C}''^{\bar{*}}.P''_b$. In all other cases, we return $\mathfrak{C}''^{\bar{*}}.S''$. Whatever the situation, the expression we return satisfies requirement 10.28. \square

Proposition 10.35 *Let X be a Noetherian space. The closed subsets of $\mathcal{T}(X)$ are exactly the languages of STREs, and the irreducible closed subsets of $\mathcal{T}(X)$ are exactly the languages of tree-products.*

Proof. One direction follows from Lemma 10.29. Conversely, we show that every closed subset of terms is expressible as some STRE.

For every simple tree expression π , the complement of π can be expressed as an STRE, by induction on (the syntax of) π and following Lemma 10.13. We deal with the second and sixth items of that lemma; the other cases are similar. In the second case, the complement of π is $X^{\bar{*}}(\mathfrak{C}\pi_1^{\bar{*}})$. Since X is Noetherian, we can express X itself as a finite union of irreducible closed subsets C_i , $1 \leq i \leq m$. By induction hypothesis, $\mathfrak{C}\pi_1$ is expressible as some STRE S . Then π is expressible as $\sum_{i=1}^m C_i^{\bar{*}}(S^{\bar{*}})$. In the sixth case, the complement of π is equal to $((F \times \{\square\}^*) \cup (X \times (\mathfrak{C}\pi_1^{\bar{*}}\{\square\}^{\bar{*}}\mathfrak{C}\pi_2^{\bar{*}}\{\square\}^{\bar{*}} \dots \{\square\}^{\bar{*}}\mathfrak{C}\pi_n^{\bar{*}})))^{\bar{*}}.\emptyset$, where F is closed in X and $n \geq 2$. Write F as a finite union of irreducible closed subsets C'_k of X , $1 \leq k \leq p$, and each $\mathfrak{C}\pi_j$, $1 \leq j \leq n$, as some STRE S_j . Recall that $X = \bigcup_{i=1}^m C_i$. Then π is expressible as the expression $\mathfrak{C}''^{\bar{*}}.S''$, where $\mathfrak{C}'' = \sum_{k=1}^p C'_k(\square^*) + \sum_{i=1}^m C_i(S_1^* \square^? S_2^* \square^? \dots \square^? S_n^*)$ and $S'' = 0$. Observe that this satisfies Requirement 10.28.

Since X is Noetherian, and simple tree expressions form a base of the tree topology (Proposition 10.1), every open subset of $\mathcal{T}(X)$ is a finite union of tree expressions. Therefore every closed subset is a finite intersection of STREs, hence itself an STRE by Lemma 10.34.

Finally, given an irreducible closed subset of $\mathcal{T}(X)$, expressed as an STRE $S = \sum_{i=1}^n P_i$, that irreducible closed subset must be the language of some P_i —by irreducibility. \square

We can now conclude.

Theorem 10.36 (S-Representation, Trees) *Let X be a Noetherian space, $X' = \mathcal{T}(X)$, and $(S, \llbracket _ \rrbracket, \trianglelefteq, \tau, \wedge)$ be an S -representation of X . Then $(S', \llbracket _ \rrbracket', \trianglelefteq', \tau', \wedge')$ is an S -*

representation of X' , where $\llbracket _ \rrbracket'$ is defined in terms of auxiliary maps that we write $\llbracket _ \rrbracket^\circ$:

- (A) S' is the collection of all tree-products over the signature S ;
- (B) \leq' is implemented by using the procedure outlined in Definition 10.31, where the inclusions between closed subsets of X are decided using \leq (and Lemma 4.2);
- (C) τ' is $(\sum_{i=1}^n C_i(\square^*))^*. \emptyset$, where $\tau = \{C_1, \dots, C_n\}$;
- (D) \wedge' is defined by the procedure of Lemma 10.34.

One should not be surprised of the shape of τ' , which is non-empty despite the mention of the empty set: see Remark 10.7.

11 Conclusion

We have developed the first comprehensive theory of (downwards-)closed subsets, as required for a general understanding of forward analysis techniques of WSTS. This generalizes previous domain proposals on tuples of natural numbers, on words, on multisets, allowing for nested datatypes, and infinite alphabets.

We have also done this on new domains such as trees, words under prefix, infinite powersets, and Noetherian rings or \mathbb{Q}^k .

Each of these domains is effective, in the sense that each has finite presentations with a decidable ordering.

We have also shown how the notion of sobrification $\mathcal{S}(X)$ was in a sense inevitable (Section 3). In the special case of wqos, the latter coincides with the ideal completion of X , and it is important to stress that we have, in particular, characterized the shapes of ideals in several wqos of interest in verification (\mathbb{N}^k , X^* , X^\otimes , $\mathcal{T}(X)$). Ideals in wqos are growingly appearing to be a central concept in verification.

The natural generalization from wqos to Noetherian spaces was also naturally in order, not only because it allows us to deal with extra constructions (words under prefix, rings, infinite powersets), but also because the proofs, as well as the computations (witness Lemma 4.2) intimately rely on irreducibility, a concept that occurs naturally from the study of sobrification.

References

- [Abdulla et al., 1998] Abdulla, P. A., Bouajjani, A., and Jonsson, B. (1998). On-the-fly analysis of systems with unbounded, lossy fifo channels. In *CAV'98*, pages 305–318, Vancouver, Canada. Springer Verlag LNCS 1427.
- [Abdulla et al., 1996] Abdulla, P. A., Čerans, K., Jonsson, B., and Tsay, Y.-K. (1996). General decidability theorems for infinite-state systems. In *LICS*, pages 313–321.
- [Abdulla et al., 2000] Abdulla, P. A., Čerans, K., Jonsson, B., and Tsay, Y.-K. (2000). Algorithmic analysis of programs with well quasi-ordered domains. *Inf. Comput.*, 160(1-2):109–127.

- [Abdulla et al., 2004a] Abdulla, P. A., Collomb-Annichini, A., Bouajjani, A., and Jonsson, B. (2004a). Using forward reachability analysis for verification of lossy channel systems. *Formal Methods in System Design*, 25(1):39–65.
- [Abdulla et al., 2004b] Abdulla, P. A., Deneux, J., Mahata, P., and Nylén, A. (2004b). Forward reachability analysis of timed Petri nets. In Lakhnech, Y. and Yovine, S., editors, *FORMATS/FTRTFT*, pages 343–362. Springer Verlag LNCS 3253.
- [Abdulla and Jonsson, 1993] Abdulla, P. A. and Jonsson, B. (1993). Verifying programs with unreliable channels. In *LICS'93*, pages 160–170.
- [Abramsky and Jung, 1994] Abramsky, S. and Jung, A. (1994). Domain theory. In Abramsky, S., Gabbay, D. M., and Maibaum, T. S. E., editors, *Handbook of Logic in Comp. Sci.*, volume 3, pages 1–168. OUP.
- [Acciai and Boreale, 2012] Acciai, L. and Boreale, M. (2012). Deciding safety properties in infinite-state pi-calculus via behavioural types. *Information and Computation*, 212:92–117.
- [Adams and Loustaunau, 1994] Adams, W. W. and Loustaunau, P. (1994). *An introduction to Gröbner bases*, volume 3 of *Graduate Studies in Mathematics*. American Mathematical Society. 289 pages.
- [Arnold and Latteux, 1978] Arnold, A. and Latteux, M. (1978). Récursivité et cônes rationnels fermés par intersection. *CALCOLO*, 15(4):381–394.
- [Baader and Nipkow, 1998] Baader, F. and Nipkow, T., editors (1998). *Term Rewriting and All That*. Cambridge University Press. 301 pages.
- [Bachmair and Plaisted, 1985] Bachmair, L. and Plaisted, D. A. (1985). Termination orderings for associative-commutative rewriting systems. *Journal of Logic and Computation*, 1(4):329–349.
- [Blondin et al., 2017a] Blondin, M., Finkel, A., and Goubault-Larrecq, J. (2017a). Forward analysis for WSTS, Part III: Karp-Miller trees. In Lokam, S. and Ramanujam, R., editors, *Proceedings of the 37th Conference on Foundations of Software Technology and Theoretical Computer Science (FSTTCS'17)*, volume 93 of *Leibniz International Proceedings in Informatics*, pages 16:1–16:15, Kanpur, India. Leibniz-Zentrum für Informatik.
- [Blondin et al., 2017b] Blondin, M., Finkel, A., and Goubault-Larrecq, J. (2017b). Forward analysis for wsts, part III: Karp-Miller trees. *CoRR*, abs/1710.07258. Long and improved version of [Blondin et al., 2017a].
- [Bojańczyk et al., 2009] Bojańczyk, M., Muscholl, A., Schwentick, T., and Ségoufin, L. (2009). Two-variable logic on data trees and XML reasoning. *Journal of the ACM*, 56(3):13:1–13:48.
- [Buchberger and Loos, 1983] Buchberger, B. and Loos, R. (1982-1983). Algebraic simplification. In Buchberger, B., Collins, G. E., Loos, R., and Albrecht, R., editors, *Computer Algebra, Symbolic and Algebraic Computation*. Springer Verlag.

- [Comon et al., 2004] Comon, H., Dauchet, M., Gilleron, R., Jacquemard, F., Lugiez, D., Tison, S., and Tommasi, M. (2004). Tree automata techniques and applications. www.grappa.univ-lille3.fr/tata.
- [Cormen et al., 2001] Cormen, T. H., Leiserson, C. E., Rivest, R. L., and Stein, C. (2001). *Introduction to Algorithms*. MIT Press and McGraw-Hill, 2nd edition.
- [de Groote et al., 2004] de Groote, P., Guillaume, B., and Salvati, S. (2004). Vector addition tree automata. In *LICS'04*, pages 64–73. IEEE Computer Society.
- [Dickson, 1913] Dickson, L. E. (1913). Finiteness of the odd perfect and primitive abundant numbers with n distinct prime factors. *American Journal of Mathematics*, 35(4):413–422.
- [Emerson and Namjoshi, 1998] Emerson, E. A. and Namjoshi, K. S. (1998). On model checking for non-deterministic infinite-state systems. In *LICS'98*, pages 70–80.
- [Erdős and Tarski, 1943] Erdős, P. and Tarski, A. (1943). On families of mutually exclusive sets. *Annals of Mathematics*, 44:315–329.
- [Faith, 1999] Faith, C. C. (1999). *Rings and Things and a Fine Array of Twentieth Century Associative Algebra*. American Mathematical Society.
- [Finkel, 1987] Finkel, A. (1987). A generalization of the procedure of Karp and Miller to well structured transition system. In Ottmann, T., editor, *Proceedings of the 14th International Colloquium on Automata, Languages and Programming (ICALP'87)*, volume 267 of *Lecture Notes in Computer Science*, pages 499–508, Karlsruhe, Germany. Springer-Verlag.
- [Finkel, 1990] Finkel, A. (1990). Reduction and covering of infinite reachability trees. *Inf. Comput.*, 89(2):144–179.
- [Finkel and Goubault-Larrecq, 2009] Finkel, A. and Goubault-Larrecq, J. (2009). Forward analysis for WSTS, part I: Completions. In Albers, S. and Marion, J.-Y., editors, *Proceedings of the 26th Annual Symposium on Theoretical Aspects of Computer Science (STACS'09)*, volume 3 of *Leibniz International Proceedings in Informatics*, pages 433–444, Freiburg, Germany. Leibniz-Zentrum für Informatik.
- [Finkel and Goubault-Larrecq, 2012] Finkel, A. and Goubault-Larrecq, J. (2012). Forward analysis for WSTS, part II: Complete WSTS. *Logical Methods in Computer Science*, 8(3:28).
- [Finkel and Schnoebelen, 2001] Finkel, A. and Schnoebelen, P. (2001). Well-structured transition systems everywhere! *Theor. Comp. Sci.*, 256(1–2):63–92.
- [Ganty et al., 2006] Ganty, P., Raskin, J.-F., and van Begin, L. (2006). A complete abstract interpretation framework for coverability properties of WSTS. In *VMCAI'06*, pages 49–64. Springer Verlag LNCS 3855.

- [Geeraerts et al., 2006] Geeraerts, G., Raskin, J.-F., and van Begin, L. (2006). Expand, enlarge and check: New algorithms for the coverability problem of WSTS. *J. Comp. Sys. Sci.*, 72(1):180–203.
- [Gierz et al., 2003] Gierz, G., Hofmann, K. H., Keimel, K., Lawson, J. D., Mislove, M., and Scott, D. S. (2003). Continuous lattices and domains. In *Encyclopedia of Mathematics and its Applications*, volume 93. CUP.
- [Goubault-Larrecq, 2007] Goubault-Larrecq, J. (2007). On Noetherian spaces. In *LICS'07*, pages 453–462.
- [Goubault-Larrecq, 2010] Goubault-Larrecq, J. (2010). Noetherian spaces in verification. In Abramsky, S., Meyer auf der Heide, F., and Spirakis, P., editors, *Proceedings of the 37th International Colloquium on Automata, Languages and Programming (ICALP'10) – Part II*, volume 6199 of *Lecture Notes in Computer Science*, pages 2–21, Bordeaux, France. Springer.
- [Goubault-Larrecq, 2013] Goubault-Larrecq, J. (2013). *Non-Hausdorff Topology and Domain Theory, Selected Topics in Point-Set Topology*, volume 22 of *New Mathematical Monographs*. Cambridge University Press.
- [Goubault-Larrecq, 2019] Goubault-Larrecq, J. (2019). Spaces with no Infinite Discrete Subspace. *Topology Proceedings*, 53:1–10. To appear.
- [Goubault-Larrecq and Schmitz, 2016] Goubault-Larrecq, J. and Schmitz, S. (2016). Deciding piecewise testable separability for regular tree languages. In Calamoneri, T., Gorla, D., Rabani, Y., Sangiorgi, D., and Mitzenmacher, M., editors, *43rd Intl. Coll. Automata, Languages, and Programming*, volume 95, pages 97:1–97:15. Leibniz-Zentrum für Informatik, Leibniz International Proceedings in Informatics.
- [Grieco and Zucchetti, 1989] Grieco, M. and Zucchetti, B. (1989). How to decide whether a polynomial ideal is primary or not. In *Proc. 5th International Conference on Applied Algebra, Algebraic Algorithms and Error-Correcting Codes (AAECC-5), 1987*, pages 258–268, Menorca, Spain. Springer-Verlag LNCS 356.
- [Grothendieck, 1960] Grothendieck, A. (1960). *Éléments de géométrie algébrique (rédigés avec la collaboration de Jean Dieudonné): I. Le langage des schémas*, volume 4. Publications mathématiques de l’I.H.É.S. pages 5–228.
- [Halfon, 2018] Halfon, S. (2018). *On Effective Representations of Well Quasi-Orderings*. PhD thesis, ENS Paris-Saclay, Université Paris-Saclay.
- [Higman, 1952] Higman, G. (1952). Ordering by divisibility in abstract algebras. *Proceedings of the London Mathematical Society*, 2(7):326–336.
- [Hoffmann, 1979a] Hoffmann, R.-E. (1979a). On the sobrification remainder ${}^s X - X$. *Pacific Journal of Mathematics*, 83(1):145–156.
- [Hoffmann, 1979b] Hoffmann, R.-E. (1979b). Sobrification of partially ordered sets. *Semigroup Forum*, 17:123–138.

- [Jacobé de Naurois, 2014] Jacobé de Naurois, P. (2014). Coverability in a nonfunctional extension of BVASS. hal-00947136. <https://hal.archives-ouvertes.fr/hal-00947136>.
- [Jacquemard et al., 2016] Jacquemard, F., Ségoufin, L., and Dimino, J. (2016). $fo^2(<, +1, \sim)$ on data trees, data tree automata and branching vector addition systems. *LMCS*, 12(2):1–28.
- [Kabil and Pouzet, 1992] Kabil, M. and Pouzet, M. (1992). Une extension d’un théorème de P. Julien sur les âges de mots. *Informatique théorique et applications*, 26(5):449–482.
- [Karp and Miller, 1969] Karp, R. M. and Miller, R. E. (1969). Parallel program schemata. *J. Comp. Sys. Sci.*, 3(2):147–195.
- [Kosaraju, 1982] Kosaraju, S. R. (1982). Decidability of reachability in vector addition systems (preliminary version). In *Proc. 14th Annual ACM Symposium on the Theory of Computing (STOC’82)*, pages 267–281, San Francisco, California, USA. ACM Press.
- [Kruskal, 1960] Kruskal, J. B. (1960). Well-quasi-ordering, the tree theorem, and vazonnyi’s conjecture. *Transactions of the American Mathematical Society*, 95(2):210–225.
- [Lambert, 1992] Lambert, J.-L. (1992). A structure to decide reachability in Petri nets. *Theoretical Computer Science*, 99(1):79–104.
- [Laplagne, 2006] Laplagne, S. (2006). Computation of the minimal associated primes. In Decker, W., Dewar, M., Kaltofen, E., and Watt, S., editors, *Challenges in Symbolic Computation Software*, number 06271 in Dagstuhl Seminar Proceedings. Internationales Begegnungs- und Forschungszentrum fuer Informatik (IBFI), Schloss Dagstuhl, Germany. <<http://drops.dagstuhl.de/opus/volltexte/2006/774>> [date of citation: 2006-01-01].
- [Lazič et al., 2008] Lazič, R., Newcomb, T., Ouaknine, J., Roscoe, A. W., and Worrell, J. (2008). Nets with tokens which carry data. *Fundamenta Informaticae*, 88(3):251–274.
- [Leroux and Schmitz, 2015] Leroux, J. and Schmitz, S. (2015). Demystifying reachability in vector addition systems. In *LICS’15*, pages 56–67.
- [Mayr, 1981] Mayr, E. W. (1981). An algorithm for the general Petri net reachability problem. In *Proc. 13th Annual ACM Symposium on the Theory of Computing (STOC’81)*, pages 238–246, Milwaukee, Wisconsin, USA. ACM Press.
- [Mayr, 2003] Mayr, R. (2003). Undecidable problems in unreliable computations. *Theor. Comput. Sci.*, 297(1-3):337–354.
- [Meyer, 2008] Meyer, R. (2008). On boundedness in depth in the pi-calculus. *IFIP TCS*, 273:477–489.

- [Michie, 1968] Michie, D. (1968). Memo functions and machine learning. *Nature*, 218:19–22.
- [Mislove, 1998] Mislove, M. (1998). Topology, domain theory and theoretical computer science. *Topology and Its Applications*, 89:3–59.
- [Müller-Olm and Seidl, 2002] Müller-Olm, M. and Seidl, H. (2002). Polynomial constants are decidable. In Hermenegildo, M. V. and Puebla, G., editors, *Proc. 9th International Symposium on Static Analysis (SAS'02)*, pages 4–19. Springer-Verlag LNCS 2477.
- [Parikh, 1966] Parikh, R. (1966). On context-free languages. *Journal of the ACM*, 13(4):570–581.
- [Pnueli, 1977] Pnueli, A. (1977). The temporal logic of programs. In *18th Annual Symposium on Foundations of Computer Science*, pages 46–57, Providence, RI, USA. IEEE Computer Society.
- [Rabinowitch, 1929] Rabinowitch, S. (1929). Zum Hilbertschen Nullstellensatz. *Math. Ann.*, 102:520.
- [Rambow, 1994] Rambow, O. (1994). Multiset-valued linear index grammars: Imposing dominance constraints on derivations. In *ACL'94*, pages 263–270. ACL Press.
- [Schmitz, 2010] Schmitz, S. (2010). On the computational complexity of dominance links in grammatical formalisms. In *ACL'10*, pages 514–524. Association for Computer Linguistics.
- [Sturmfels, 2002] Sturmfels, B. (2002). *Solving Systems of Polynomial Equations*, volume 97 of *CBMS Regional Conferences Series*. American Mathematical Society.
- [Thatcher and Wright, 1968] Thatcher, J. W. and Wright, J. B. (1968). Generalized finite automata theory with an application to a decision problem of second-order logic. *Mathematical Systems Theory*, 2(1):57–81.
- [Verma and Goubault-Larrecq, 2005] Verma, K. N. and Goubault-Larrecq, J. (2005). Karp-Miller trees for a branching extension of VASS. *Discrete Mathematics & Theoretical Computer Science*, 7(1):217–230.
- [Wies et al., 2010] Wies, T., Zufferey, D., and Henzinger, T. A. (2010). Forward analysis of depth-bounded processes. In Ong, L., editor, *Proc. 13th Intl. Conf. Foundations of Software Science and Computational Structures (FoSSaCS'10)*, pages 94–108. Springer Verlag LNCS 6014.

A Auxiliary Proofs on Irreducible Closed Sets

The results of this section are well-known. We include their proofs for convenience.

Lemma A.1 *For every continuous map $f: Y \rightarrow Z$ between topological spaces Y and Z , for every irreducible closed subset C of Y , $cl(f[C])$ is irreducible closed in Z .*

Proof. This is a direct consequence of the fact that \mathcal{S} is a functor, in particular that $\mathcal{S}(f)$ maps every $C \in \mathcal{S}(Y)$ to an element of $\mathcal{S}(Z)$, and of the fact that $\mathcal{S}(f)(C) = cl(f[C])$. Here is a direct argument. Assume that $cl(f[C])$ is included in the union of two closed subsets F_1 and F_2 of Z . Then $f[C] \subseteq F_1 \cup F_2$, so $C \subseteq f^{-1}(F_1 \cup F_2) = f^{-1}(F_1) \cup f^{-1}(F_2)$. Since C is irreducible, C is included in $f^{-1}(F_1)$ or in $f^{-1}(F_2)$. Assume without loss of generality that $C \subseteq f^{-1}(F_1)$. Then $f[C]$ is included in F_1 , and since $cl(f[C])$ is the smallest closed set containing $f[C]$, and F_1 is a closed set containing $f[C]$, $cl(f[C])$ is included in F_1 . \square

Lemma A.2 *Let $(X_i)_{i \in I}$ be any family of topological spaces. The irreducible closed subsets of $\prod_{i \in I} X_i$ are exactly the products $\prod_{i \in I} C_i$ where each C_i is irreducible closed in X_i .*

Proof. This is Proposition 8.4.7 of [Goubault-Larrecq, 2013], and leads to Hoffmann's theorem [Hoffmann, 1979a, Theorem 1.4] that the sobrification functor preserves products. We give a direct proof. Let $X = \prod_{i \in I} X_i$. This has a base of open sets of the form $\bigcap_{i \in J} \pi_i^{-1}(U_i)$, where J is a finite subset of I and each U_i is open in X_i .

We first note that a closed set C is irreducible if and only if it is non-empty, and whenever it intersects two opens U_1 and U_2 , then it also intersects their intersection $U_1 \cap U_2$. (Just take the contrapositive of the definition of irreducibility, with the complement of U_1 for F_1 and the complement of U_2 for F_2 .)

If each C_i is closed in X_i , then $\prod_{i \in I} C_i$ is closed, being the intersection $\bigcap_{i \in I} \pi_i^{-1}(C_i)$. If additionally each C_i is irreducible, we claim that $C = \prod_{i \in I} C_i$ is also irreducible. It is certainly non-empty.

Assume that C intersects two open subsets W and W' of X . Write W as a union of basic open subsets $\bigcap_{j \in J_k} \pi_j^{-1}(U_{jk})$, $k \in K$, where each J_k is a finite subset of I , and each U_{jk} is open in X_j . Then C intersects $\bigcap_{j \in J_k} \pi_j^{-1}(U_{jk})$ for some $k \in K$, say at $(x_i)_{i \in I}$. Similarly, write W' as a union of basic open subsets $\bigcap_{j \in J_{k'}} \pi_j^{-1}(U_{jk'})$, $k' \in K'$, where each $J_{k'}$ is a finite subset of I , and each $U_{jk'}$ is open in X_j . Then C intersects $\bigcap_{j \in J_{k'}} \pi_j^{-1}(U_{jk'})$ for some $k' \in K'$, say at $(x'_i)_{i \in I}$. Without loss of generality, we may assume that $J_k = J_{k'}$: otherwise replace J_k , resp. $J_{k'}$, by $J_k \cup J_{k'}$, and define the missing sets U_{jk} , resp. $U_{jk'}$, as X_j . For every $j \in J_k$, C_j intersects U_{jk} (at x_j) and also $U_{jk'}$ (at x'_j). Since C_j is irreducible, it therefore also intersects their intersection $U_{jk} \cap U_{jk'}$, say at y_j . For every $j \in I \setminus J_k$, define y_j as some arbitrary point from C_j . Then $(y_j)_{j \in I}$ is in C , and in $\bigcap_{j \in J_k} \pi_j^{-1}(U_{jk} \cap U_{jk'})$, hence in $W \cap W'$.

Conversely, we claim that every irreducible closed subset C of X must be a product $\prod_{i \in I} C_i$ of irreducible closed subsets C_i of X_i . We define C_i as $cl(\pi_i[C])$: by Lemma A.1, C_i is irreducible closed in X_i . Clearly, C is included in $\prod_{i \in I} C_i$. Assume for the sake of contradiction that the inclusion were strict: there is a point $\vec{x} = (x_i)_{i \in I}$ in $\prod_{i \in I} C_i$ and not in C . Then \vec{x} is in the open complement W of C . By definition of the product topology, W contains a basic open subset $\bigcap_{j \in J} \pi_j^{-1}(U_j)$ containing \vec{x} , where J is a finite subset of I , and each U_j is open in X_j . Since it contains \vec{x} , x_j is in U_j for every $j \in J$. Since it is included in W , $\bigcap_{j \in J} \pi_j^{-1}(U_j)$ is disjoint from C . Since C is irreducible, if C intersected $\pi_j^{-1}(U_j)$ for every $j \in J$, it would also intersect $\bigcap_{j \in J} \pi_j^{-1}(U_j)$, which is impossible, as we have just seen. Hence C is disjoint

from $\pi_j^{-1}(U_j)$ for some $j \in J$. This implies that $\pi_j[C]$ is disjoint from U_j , hence that $C_j = cl(\pi_j[C])$ is also disjoint from U_j . (A set intersects an open set if and only if its closure does.) However, since $\vec{x} \in \prod_{i \in I} C_i$, x_j is in C_j , and x_j is also in U_j , contradiction. \square

B Proofs of Results on Words (Section 6)

Lemma 6.2 (recap). Let X be a topological space. The complement of $X^*U_1X^*U_2X^*\cdots X^*U_nX^*$ ($n \in \mathbb{N}$, U_1, U_2, \dots, U_n open in X) in X^* is \emptyset when $n = 0$, and $F_1^*X^?F_2^*X^?\cdots X^?F_{n-1}^*X^?F_n^*$ otherwise, where $F_1 = X \setminus U_1, \dots, F_n = X \setminus U_n$.

If X is Noetherian, then this complement can be expressed as a finite union of sets of the form $F_1^*C_1^?F_2^*C_2^?\cdots C_{n-1}^?F_n^*$, where C_1, C_2, \dots, C_{n-1} range over irreducible closed subsets of X .

Proof. When $n = 0$, this is clear: the complement of $X^*U_1X^*U_2X^*\cdots X^*U_nX^*$ is the empty set. So let $n \geq 1$.

We first claim that the complement of $X^*U_1X^*U_2X^*\cdots X^*U_nX^*$ is $F_1^*X^?F_2^*X^?\cdots X^?F_{n-1}^*X^?F_n^*$. We show this by induction on n . If $n = 1$, then the complement of $X^*U_1X^*$ is the set of words that contain no letter from U_1 , i.e., F_1^* . If $n \geq 1$, let w be an arbitrary element of the complement of $X^*U_1X^*U_2X^*\cdots X^*U_nX^*$. Let w_1 be the longest prefix of w comprised of letters not in U_1 . Note that w_1 is in F_1^* . If $w_1 = w$, then certainly w is in $F_1^* \subseteq F_1^*X^?F_2^*X^?\cdots X^?F_{n-1}^*X^?F_n^*$. Otherwise, w is of the form w_1xw' , where $x \in U_1$ and w' is not in $X^*U_2X^*\cdots X^*U_nX^*$. By induction hypothesis w' is in $F_2^*X^?\cdots X^?F_{n-1}^*X^?F_n^*$, hence again w is in $F_1^*X^?F_2^*X^?\cdots X^?F_{n-1}^*X^?F_n^*$.

Conversely, let w be any word in $F_1^*X^?F_2^*X^?\cdots X^?F_{n-1}^*X^?F_n^*$. Let w_1 be the longest prefix of w that lies in F_1^* . Then either $w = w_1$, then $w \in F_1^*$ cannot be in $X^*U_1X^*U_2X^*\cdots X^*U_nX^*$, since all the words in the latter set must contain at least one letter in U_1 ; or $w = w_1xw'$ for some $x \notin U_1$, i.e. $x \in U_1$, and $w' \in F_2^*X^?\cdots X^?F_{n-1}^*X^?F_n^*$. By induction hypothesis, w' cannot be in $X^*U_2X^*\cdots X^*U_nX^*$. By construction, x would be the first occurrence of an element of U_1 in w . If $w = w_1xw'$ were in $X^*U_1X^*U_2X^*\cdots X^*U_nX^*$, then, some suffix w'' of w' would be in $X^*U_2X^*\cdots X^*U_nX^*$. Then $w'' \leq^* w'$, hence w' would be in $X^*U_2X^*\cdots X^*U_nX^*$, which is open hence upwards-closed: contradiction.

By Lemma 3.6, X , as a closed subset of itself, is a finite union of irreducible closed subsets. I.e., there is a finite subset E of $\mathcal{S}(X)$ such that $X = \bigcup_{C \in E} C$. Distributing across the $_?$ operator and concatenation in the expression $F_1^*X^?F_2^*X^?\cdots X^?F_{n-1}^*X^?F_n^*$ yields that the complement of $X^*U_1X^*U_2X^*\cdots X^*U_nX^*$ equals:

$$\bigcup_{C_1, \dots, C_{n-1} \in E} F_1^*C_1^?F_2^*C_2^?\cdots C_{n-2}^?F_{n-1}^*C_{n-1}^?$$

from which the desired conclusion follows. \square

Lemma 6.4 (recap). Let X be a topological space. For every closed subset F of X , for every closed subset \mathcal{F} of X^* , $F^?\mathcal{F}$ is closed in X^* .

This is part of Exercise 9.7.29 of [Goubault-Larrecq, 2013], but the argument is non-trivial. Here is a complete proof. This requires the following construction, which

we shall also require in the next Lemma. For any open \mathcal{U} of X^* , and any open U of X , define \mathcal{U}/U as follows. If $\mathcal{U} = X^*$, then $\mathcal{U}/U = \emptyset$; otherwise, \mathcal{U} is a union of basic opens of the form $X^*U_{i_1}X^*U_{i_2}X^*\cdots X^*U_{i_{n_i}}X^*$, $i \in I$, where $n_i \geq 1$ for every $i \in I$, then we let \mathcal{U}/U be the union of all basic opens $X^*(U_{i_1} \cap U)X^*U_{i_2}X^*\cdots X^*U_{i_{n_i}}X^*$. In all formality, \mathcal{U}/U depends on a presentation of \mathcal{U} as a union of basic opens, not on \mathcal{U} itself, but this will cause no problem in the sequel. Note that this definition is made possible by the fact that $X^*U_1X^*U_2X^*\cdots X^*U_nX^*$ form a base, not just a subbase, of the topology (Lemma 6.1).

Proof. If \mathcal{F} is empty, then $F^? \mathcal{F}$ is empty hence closed. Henceforth we assume that \mathcal{F} is non-empty, so that the complement \mathcal{U} is different from X^* . Then \mathcal{U}/U is open. Also, $X^*U\mathcal{U}$ is open. We claim that the complement of $F^? \mathcal{F}$ in X^* is $X^*X\mathcal{U} \cup \mathcal{U}/U$, which will show the claim. We first make the following remark. Let L_1 and L_2 be two subsets of X^* that are downwards-closed with respect to \leq^* . For any word w not in L_1L_2 , we can write w as $w_1w'w_2$, where w_1 is the longest prefix of w in L_1 , w_2 is the longest suffix of w in L_2 , and w' is not empty. Indeed, any prefix of a word in L_1 is again in L_1 , and any suffix of a word in L_2 is in L_2 , since both are downwards-closed with respect to \leq^* . That remark applies, notably, to $L_1 = F^?$ and $L_2 = \mathcal{F}$ fit, in the second case because \leq^* is the specialization quasi-ordering of X^* and every closed subset is downwards-closed.

If $\mathcal{F} = X^*$, then the complement of $F^? \mathcal{F} = X^*$ is empty, so the claim is proved. Otherwise, write \mathcal{U} as the union of basic opens $X^*U_{i_1}X^*U_{i_2}X^*\cdots X^*U_{i_{n_i}}X^*$, $i \in I$, $n_i \geq 1$.

Assume w is in the complement of $F^? \mathcal{F}$, and write w as $w_1w'w_2$, as above. Since w' is not empty, it starts with some letter $x \in X$. Then by the maximality property of w_1 , w_1 is in $F^?$, but w_1x is not. Again, by the maximality property of w_2 , $w'w_2$ is not in \mathcal{F} , hence in \mathcal{U} . If $w_1 \neq \varepsilon$, then w_1 is a single letter, so w is in $X\mathcal{U} \subseteq X^*X\mathcal{U}$. If $w_1 = \varepsilon$, then x is not in F (otherwise w_1x would be in $F^?$), hence is in U . So $w'w_2$ starts with a letter in U ; since $w'w_2$ is in \mathcal{U} , $w'w_2$ is in $X^*U_{i_1}X^*U_{i_2}X^*\cdots X^*U_{i_{n_i}}X^*$ for some $i \in I$. If $w'w_2$ is in $U_{i_1}X^*U_{i_2}X^*\cdots X^*U_{i_{n_i}}X^*$, then $w'w_2$ is in $(U \cap U_{i_1})X^*U_{i_2}X^*\cdots X^*U_{i_{n_i}}X^*$, so $w = w_1w'w_2$ is in $X^*(U \cap U_{i_1})X^*U_{i_2}X^*\cdots X^*U_{i_{n_i}}X^* \subseteq \mathcal{U}/U$; otherwise, $w'w_2$ is in $UX^*U_{i_1}X^*U_{i_2}X^*\cdots X^*U_{i_{n_i}}X^*$, so $w = w_1w'w_2$ is in $X^*UX^*U_{i_1}X^*U_{i_2}X^*\cdots X^*U_{i_{n_i}}X^* \subseteq X^*U\mathcal{U} \subseteq X^*X\mathcal{U}$.

Conversely, assume w is in $X^*X\mathcal{U} \cup \mathcal{U}/U$. If $w \in X^*X\mathcal{U}$, then w contains a subsequence of the form $a_0a_1a_2 \cdots a_{n_i}$, for some $i \in I$, where a_0 is arbitrary, $a_1 \in U_{i_1}$, $a_2 \in U_{i_2}, \dots, a_{n_i} \in U_{i_{n_i}}$. Note that $a_1a_2 \cdots a_{n_i}$ is in $U_{i_1}U_{i_2} \cdots U_{i_{n_i}} \subseteq X^*U_{i_1}X^*U_{i_2}X^*\cdots X^*U_{i_{n_i}}X^* \subseteq \mathcal{U}$. If w were in $F^? \mathcal{F}$, then since $F^? \mathcal{F}$ is downwards-closed with respect to \leq^* , $a_0a_1a_2 \cdots a_{n_i}$ would be in $F^? \mathcal{F}$, hence $a_1a_2 \cdots a_{n_i}$ would be in \mathcal{F} : contradiction. (More slowly: from $a_0a_1a_2 \cdots a_{n_i} \in F^? \mathcal{F}$ we deduce that $a_0 \in F$ and $a_1a_2 \cdots a_{n_i} \in \mathcal{F}$, or $a_0a_1a_2 \cdots a_{n_i} \in \mathcal{F}$; but the latter also implies $a_1a_2 \cdots a_{n_i} \in \mathcal{F}$, since \mathcal{F} is downwards-closed.) So w is in the complement of $F^? \mathcal{F}$. If, on the other hand, $w \in \mathcal{U}/U$, then w contains a subsequence of the form $a_1a_2 \cdots a_{n_i}$, for some $i \in I$, where $a_1 \in U \cap U_{i_1}$, $a_2 \in U_{i_2}, \dots, a_{n_i} \in U_{i_{n_i}}$. In particular, $a_1a_2 \cdots a_{n_i}$ is in $U_{i_1}U_{i_2} \cdots U_{i_{n_i}} \subseteq X^*U_{i_1}X^*U_{i_2}X^*\cdots X^*U_{i_{n_i}}X^* \subseteq \mathcal{U}$. If w were in $F^? \mathcal{F}$, then since $F^? \mathcal{F}$ is downwards-closed with respect to \leq^* , $a_1a_2 \cdots a_{n_i}$ would be in $F^? \mathcal{F}$. However, a_1 is in U , so is not in F , and this implies that $a_1a_2 \cdots a_{n_i}$ would be in \mathcal{F} :

contradiction. So, again, w is in the complement of $F^? \mathcal{F}$. □

Lemma 6.5 (recap). Let X be a topological space. For every closed subset F of X , for every closed subset \mathcal{F} of X^* , $F^* \mathcal{F}$ is closed in X^* .

More specifically, in the case where \mathcal{F} is non-empty, let U be the open complement of F in X , \mathcal{U} be the open complement of \mathcal{F} in X^* . Then the complement of $F^* \mathcal{F}$ is $X^* \mathcal{U} \mathcal{U} \cup \mathcal{U} / \mathcal{U}$, and is therefore open.

Proof. Write \mathcal{U} is a union of basic opens of the form $X^* U_{i_1} X^* U_{i_2} X^* \cdots X^* U_{i_{n_i}} X^*$, $i \in I$, as above.

Assume w is in the complement of $F^* \mathcal{F}$, and write w as $w_1 w' w_2$, where w_1 is the longest prefix of w in F^* , w_2 is the longest suffix of w in \mathcal{F} , and w' is not empty. Let x be the first letter of w' , and note that $w_1 \in F^*$ but $w_1 x$ is not in F^* : so x is in U ; and that $w' w_2$ is in \mathcal{U} , so $w' w_2$ is in some basic open set $X^* U_{i_1} X^* U_{i_2} X^* \cdots X^* U_{i_{n_i}} X^*$, $i \in I$. Depending on whether the first letter x of $w' w_2$ is in U_{i_1} or not, $w' w_2$ is in $(U \cap U_{i_1}) X^* U_{i_2} X^* \cdots X^* U_{i_{n_i}} X^*$ or in $U X^* U_{i_1} X^* U_{i_2} X^* \cdots X^* U_{i_{n_i}} X^*$, so that w is in $X^* \mathcal{U} \mathcal{U}$ or in $\mathcal{U} / \mathcal{U}$.

Conversely, if $w \in X^* \mathcal{U} \mathcal{U}$, then w contains a subword $a_0 a_1 a_2 \cdots a_{n_i}$ for some $i \in I$, $a_0 \in U$, $a_1 \in U_{i_1}$, $a_2 \in U_{i_2}$, \dots , $a_{n_i} \in U_{i_{n_i}}$. If w were in $F^* \mathcal{F}$, then $a_0 a_1 a_2 \cdots a_{n_i}$ would be, too: it is indeed clear that $F^* \mathcal{F}$ is downwards-closed, since F^* is and \mathcal{F} is closed. Since $a_0 \in U$, a_0 is not in F , so $a_0 a_1 a_2 \cdots a_{n_i}$ is in \mathcal{F} . Again by downward closure, $a_1 a_2 \cdots a_{n_i}$ is in \mathcal{F} : contradiction. So w is in the complement of $F^* \mathcal{F}$. And if $w \in \mathcal{U} / \mathcal{U}$, then w contains a subword of the form $a_1 a_2 \cdots a_{n_i}$, for some $i \in I$, where $a_1 \in U \cap U_{i_1}$, $a_2 \in U_{i_2}$, \dots , $a_{n_i} \in U_{i_{n_i}}$. In particular, $a_1 a_2 \cdots a_{n_i}$ is in $U_{i_1} U_{i_2} \cdots U_{i_{n_i}} \subseteq X^* U_{i_1} X^* U_{i_2} X^* \cdots X^* U_{i_{n_i}} X^* \subseteq \mathcal{U}$. If w were in $F^* \mathcal{F}$, then so would be this subword, and as $a_1 \in U$ is not in F , $a_1 a_2 \cdots a_{n_i}$ would be in \mathcal{F} : contradiction. So w is in the complement of $F^* \mathcal{F}$. □

Lemma B.1 *The concatenation function $cat: X^* \times X^* \rightarrow X^*$ is continuous. The function $i: X \rightarrow X^*$ that maps the letter x to x as a word, is also continuous.*

Proof. This is Exercise 9.7.27 of [Goubault-Larrecq, 2013]. To show that i is continuous, observe that $i^{-1}(X^* U_1 X^* U_2 X^* \cdots X^* U_n X^*) = U_1$ if $n = 1$, X if $n = 0$, and \emptyset otherwise. As far as cat is concerned, $cat^{-1}(X^* U_1 X^* U_2 X^* \cdots X^* U_n X^*) = \bigcup_{i=0}^n (X^* U_1 X^* U_2 X^* \cdots X^* U_i X^*) \times (X^* U_{i+1} X^* U_2 X^* \cdots X^* U_n X^*)$. □

Lemma 6.7 (recap). Let X be a topological space. Every word-product is irreducible closed in X^* .

Proof. By induction on syntax, starting with the fact that the base case ϵ denotes a one-element downwards-closed set, hence is trivially irreducible closed. It suffices to show that sets of the form F^* and $C^?$ are irreducible closed, where F is closed and C is irreducible closed, and that if \mathcal{C}_1 and \mathcal{C}_2 are irreducible closed and $\mathcal{C}_1 \mathcal{C}_2$ is closed, then $\mathcal{C}_1 \mathcal{C}_2$ is irreducible closed.

1. Let us show that F^* is irreducible closed in X^* , for any closed subset F of X . Assume $F^* \subseteq \mathcal{F}_1 \cup \mathcal{F}_2$, where \mathcal{F}_1 and \mathcal{F}_2 are closed in X^* . If F^* was not contained in \mathcal{F}_1 or in \mathcal{F}_2 , then there would be a word $w_1 \in F^* \setminus \mathcal{F}_1$ and a word $w_2 \in F^* \setminus \mathcal{F}_2$. Then $w_1 w_2$ would again be in F^* , hence either in \mathcal{F}_1 or in \mathcal{F}_2 . Assume by symmetry that

w_1w_2 is in \mathcal{F}_1 . Since $w_1 \leq^* w_1w_2$, and closed sets such as \mathcal{F}_1 are downwards-closed, we would have $w_1 \in \mathcal{F}_1$: contradiction. So F^* is irreducible.

2. We claim that $C^?$ is irreducible closed in X^* whenever C is irreducible closed in X . It is enough to observe that $C^? = cl(i[C])$ —recall that i is continuous by Lemma B.1—and to use Lemma A.1. The inclusion $cl(i[C]) \subseteq C^?$ stems for the fact that $C^?$ is closed, and $i[C] \subseteq C^?$, which is clear. The converse inclusion is obvious.

3. Finally, we show that whenever \mathcal{C}_1 and \mathcal{C}_2 are irreducible closed in X^* , and $\mathcal{C}_1\mathcal{C}_2$ is closed, it is irreducible. Indeed, $\mathcal{C}_1 \times \mathcal{C}_2$ is irreducible closed by Lemma A.2, and since cat is continuous by Lemma B.1, $cl(cat[\mathcal{C}_1 \times \mathcal{C}_2])$ is irreducible by Lemma A.1. Then $cat[\mathcal{C}_1 \times \mathcal{C}_2] = \mathcal{C}_1\mathcal{C}_2$, and since the latter is closed by assumption, it is equal to $cl(cat[\mathcal{C}_1 \times \mathcal{C}_2])$, hence irreducible closed. \square

Lemma 6.9 (recap). Let X be a topological space. Inclusion between word-products can be checked in polynomial time (precisely in time proportional to the product of the lengths of the two word-products), modulo an oracle testing inclusion of closed subsets of X .

Explicitly, we have: $\epsilon \subseteq P$ for any word-product P , $P \not\subseteq \epsilon$ unless all the atomic expressions in P are syntactically equal to \emptyset^* , and for all $C, C' \in \mathcal{S}(X)$, for all $F, F' \in \mathcal{H}_V(X)$, for all word-products P, P' :

- $C^?P \subseteq C'^?P'$ if and only if $C \subseteq C'$ and $P \subseteq P'$, or $C \not\subseteq C'$ and $C^?P \subseteq P'$.
- $C^?P \subseteq F'^*P'$ if and only if $C \subseteq F'$ and $P \subseteq F'^*P'$, or $C \not\subseteq F'$ and $C^?P \subseteq P'$.
- $F^*P \subseteq C'^?P'$ if and only if F is empty and $P \subseteq C'^?P'$, or F is non-empty and $F^*P \subseteq P'$.
- $F^*P \subseteq F'^*P'$ if and only if $F \subseteq F'$ and $P \subseteq F'^*P'$, or $F \not\subseteq F'$ and $F^*P \subseteq P'$.

Proof. The cases $\epsilon \subseteq P$ and $P \not\subseteq \epsilon$ are obvious.

We first examine when $C^?P \subseteq C'^?P'$ holds. The if direction is easy. Conversely, assume $C^?P \subseteq C'^?P'$. For every $x \in C$, either $x \in C'$ or xw is in P' for every $w \in P$: indeed, when $w \in P$, then $xw \in C^?P \subseteq C'^?P'$, and if $x \notin C'$ this can only happen if $xw \in P'$. This means that C is contained in $C' \cup F$, where $F = \{x \in X \mid \forall w \in P \cdot xw \in P'\} = \bigcap_{w \in P} f_w^{-1}(P')$, and f_w is the map $x \mapsto xw$. Note that $f_w(x) = cat(i(x), w)$, hence f_w is continuous. Using the fact that P' is closed (Corollary 6.6), F is closed. Since $C \subseteq C' \cup F$ and C is irreducible, we have proved: (*) either $C \subseteq C'$ or $C \subseteq F$. We also note that: (**) $P \subseteq P'$, in any case: fixing some element $x \in C$ (recall that C is non-empty), for every $w \in P$, xw is in $C^?P$, hence in $C'^?P'$; so w or xw is in P' ; since P' is closed and $w \leq^* xw$, w must be in P' in any case. Using (*) and (**), we now have two cases: either $C \subseteq C'$ and $P \subseteq P'$, or $C \not\subseteq C'$, $C \subseteq F$, and $P \subseteq P'$. In the latter case, $C \subseteq F$ entails $CP \subseteq P'$ by definition of F , so $C^?P = P \cup CP \subseteq P'$.

We now examine when $C^?P \subseteq F'^*P'$ holds. This is similar. The if direction is obvious. Conversely, if $C^?P \subseteq F'^*P'$, then for every $x \in C$, either $x \in F'$ or $xw \in P'$ for every $w \in P$. So $C \subseteq F' \cup F$, where F is the closed set $\{x \in X \mid$

$\forall w \in P \cdot xw \in P'\}$. Since C is irreducible, (*) either $C \subseteq F'$ or $C \subseteq F$. Also, (**) $P \subseteq F'^*P'$, because $P \subseteq C^?P \subseteq F'^*P'$. If $C \not\subseteq F'$, then $C \subseteq F$, hence $CP \subseteq P'$ by the definition of F ; since P' is downwards-closed, P is also included in P' , so $C^?P = P \cup CP \subseteq P'$.

Let us proceed to the case $F^*P \subseteq C'^?P'$. When F is empty, since $F^*P = P$, the equivalence between $F^*P \subseteq C'^?P'$ and $P \subseteq C'^?P'$ is obvious. Otherwise, since F is non-empty, let x be some element in F . For any $w \in F^*P$, xw is also in F^*P , so is in $C'^?P'$. This implies that xw or w is in P' . But, as P' is downwards-closed, $w \in P'$ in any case. So $F^*P \subseteq P'$. The converse is again easy.

Finally, assume $F^*P \subseteq F'^*P'$. If $F \subseteq F'$, then $P \subseteq F'^*P'$, since $P \subseteq F^*P$. Otherwise, let x be in F but not in F' . For any word $w \in F^*P$, xw is again in F^*P , hence in F'^*P' . Since $x \notin F'$, xw must be in P' , hence also $w \in P'$. So $F^*P \subseteq P'$.

We obtain the desired algorithm (up to an oracle) by dynamic programming. \square

Lemma 6.12 (recap). Let X be a topological space. Any finite intersection of word-products is expressible as a finite union of word-products. Specifically, the intersection of two word-products is given by: $\epsilon \cap P = \epsilon$ for every word-product P , and by the recursive formulae:

- $C^?P \cap C'^?P' = (C^?P \cap P') \cup (P \cap C'^?P') \cup (C \cap C')^?(P \cap P')$;
- $C^?P \cap F'^*P' = (C \cap F')^?(P \cap F'^*P') \cup (C^?P \cap P')$;
- $F^*P \cap F'^*P' = (F \cap F')^*(P \cap F'^*P') \cup (F \cap F')^*(F^*P \cap P')$.

Proof. Let us deal with the first case. Any word w in $C^?P \cap C'^?P'$ is either in $P \cap P'$, or is in CP and in P' , or in P and in $C'P'$, or is of the form xw' , with $x \in C \cap C'$ and $w' \in P \cap P'$. So $C^?P \cap C'^?P' \subseteq (P \cap P') \cup (C^?P \cap P') \cup (P \cap C'^?P') \cup (C \cap C')^?(P \cap P') = (C^?P \cap P') \cup (P \cap C'^?P') \cup (C \cap C')^?(P \cap P')$. It is easy to see that conversely, $(C^?P \cap P') \cup (P \cap C'^?P') \cup (C \cap C')^?(P \cap P')$ is included in $C^?P \cap C'^?P'$.

Next, any word w in $C^?P \cap F'^*P'$ is either in $P \cap F'^*P'$, or is of the form xw' with $x \in C$, $w' \in P$, and $xw' \in F'^*P'$. In the latter case, either $x \in C \cap F'$ and $w' \in P \cap F'^*P'$, so $w \in (C \cap F')^?(P \cap F'^*P')$; or $x \in C$, x is not in F' so $w = xw'$ is in P' , hence w is in $C^?P \cap P'$. In any case, $C^?P \cap F'^*P' \subseteq (P \cap F'^*P') \cup (C \cap F')(P \cap F'^*P') \cup (C^?P \cap P') = (C \cap F')^?(P \cap F'^*P') \cup (C^?P \cap P')$. The converse inclusion is clear.

Finally, for every word w in $F^*P \cap F'^*P'$, write w as w_1w_2 where w_1 is the longest prefix of w in F^* , and $w_2 \in P$; also, as $w'_1w'_2$ where w'_1 is the longest prefix of w in F'^* , and $w'_2 \in P'$. If w_1 is shorter than w'_1 , then w_2 is also in F'^*P' , so $w \in (F \cap F')^*(P \cap F'^*P')$, otherwise $w \in (F \cap F')^*(F^*P \cap P')$. So $F^*P \cap F'^*P' \subseteq (F \cap F')^*(P \cap F'^*P') \cup (F \cap F')^*(F^*P \cap P')$. The converse inclusion is obvious.

These formulae define the intersection of two word-products, by induction on the sum of the number of atomic formulae in each of them. So the intersection of two word-products is a finite union of word-products. The empty intersection, the space X^* itself, is clearly a word-product. By induction on the number of word-products, any finite intersection of word-products can then be rewritten as a finite union of word-products. \square

Proposition B.2 Let X be a Noetherian space, $X' = X^*$, and $(S, \llbracket _ \rrbracket, \leq, \tau, \wedge)$ be an S -representation of X . Then $(S', \llbracket _ \rrbracket', \leq', \tau', \wedge')$ is an S -representation of X' , where:

- S' is the collection of all word-product notations, i.e., of all expressions of the form $e_1 e_2 \cdots e_n$ where each e_i is either of the form u^* where u is a finite subset of S , or of the form $a^?$ where $a \in S$. We write ϵ for the empty word-product notation.
- $\llbracket e_1 e_2 \cdots e_n \rrbracket' = \llbracket e_1 \rrbracket' \llbracket e_2 \rrbracket' \cdots \llbracket e_n \rrbracket' = \{x_1 x_2 \cdots x_n \mid x_1 \in \llbracket e_1 \rrbracket', x_2 \in \llbracket e_2 \rrbracket', \dots, x_n \in \llbracket e_n \rrbracket'\}$, where $\llbracket u^* \rrbracket'$ is the set of all finite words whose letters are in $\bigcup_{a \in u} \llbracket a \rrbracket'$, and $\llbracket a^? \rrbracket'$ is the set of words containing at most one letter, and this letter is in $\llbracket a \rrbracket'$;
- \leq' is the relation $\text{star}[\leq]$, defined recursively from \leq by: $\epsilon \leq' w$ for every word-product notation w , $w \leq' \epsilon$ iff $w = \epsilon$, and for all $a, a' \in S$, for all non-empty finite subsets u, u' of S :
 - $a^? w \leq' a'^? w'$ iff $a \leq a'$ and $w \leq' w'$, or $a \not\leq a'$ and $a^? w \leq' w'$;
 - $a^? w \leq' u'^* w'$ iff $a \leq a'$ for some $a' \in u'$ and $w \leq' u'^* w'$, or $a \leq a'$ for no $a' \in u'$ and $a^? w \leq' w'$;
 - $u^* w \leq' a'^? w'$ iff u is empty and $w \leq' a'^? w'$, or u is non-empty and $u^* w \leq' w'$;
 - $u^* w \leq' u'^* w'$ iff either for every $a \in u$, there is an $a' \in u'$ such that $a \leq a'$ and $w \leq' u'^* w'$, or there is an $a \in u$ such that $a \leq a'$ for no $a' \in u'$, and $u^* w \leq' w'$.
- τ' is $\{\tau^*\}$.
- \wedge' is the map $\text{meet}[\wedge]$, parametrized by \leq , and defined recursively by: $\epsilon \wedge' w' = \{\epsilon\}$, $w \wedge' \epsilon = \{\epsilon\}$, and for all $a, a' \in S$, for all non-empty finite subsets u, u' of S :
 - $a^? w \wedge' a'^? w' = \{a''^? w'' \mid a'' \in a \wedge a', w'' \in w \wedge' w'\} \cup (a^? w \wedge' w') \cup (w \wedge' a'^? w')$;
 - $a^? w \wedge' u'^* w' = \{a''^? w'' \mid a'' \in u', a'' \in a \wedge a', w'' \in w \wedge' u'^* w'\} \cup (a^? w \wedge' w')$ if there is an $a' \in u'$ such that $a \wedge a' \neq \emptyset$, $a^? w \wedge' u'^* w' = (w \wedge' u'^* w') \cup (a^? w \wedge' w')$ otherwise; and similarly for $u^* w \wedge' a'^? w'$;
 - $u^* w \wedge' u'^* w' = \left\{ \left(\bigcup_{\substack{a \in u \\ a' \in u'}} a \wedge a' \right)^* w'' \mid w'' \in (u^* w \wedge' w') \cup (w \wedge' u'^* w') \right\}$.

Proof. $\llbracket _ \rrbracket'$ is surjective: the irreducible closed subsets of X' are the word-products by Proposition 6.14, i.e., of the form $\llbracket w \rrbracket'$ for some word-product notation $w \in X'$.

The formulae defining $\text{star}[\leq]$ are obtained from Lemma 6.9. The formula defining τ' is justified by the fact that $\bigcup_{w \in \tau'} \llbracket w \rrbracket' = \llbracket \tau^* \rrbracket'$ is the set of all words whose letters are all in $\bigcup_{a \in \tau} \llbracket a \rrbracket' = X$, i.e., this is the whole set X^* . The formulae defining \wedge' are obtained from Lemma 6.13. \square

C Proofs of Results on Multisets (Section 7)

Proposition 7.4 (recap). Let X be a topological space. Then the m-SREs are closed in X^{\otimes} , and the m-products are irreducible closed.

If X is Noetherian, then every irreducible closed subset of X^{\otimes} is an m-product, and every closed subset of X^{\otimes} is an m-SRE.

Proof. Consider any m-product $P = F \mid C_1, C_2, \dots, C_n$. We observe that $\Psi^{-1}(P)$ is the union over all permutations π of $\{1, 2, \dots, n\}$ of the word-products $F^* C_{\pi(1)}^? F^* C_{\pi(2)}^? F^* \dots F^* C_{\pi(n)}^? F^*$. This means that the words whose multiset of letters can be split as at most one letter from each of C_1, C_2, \dots, C_n , plus remaining letters from F , are just the words that are comprised of letters from F , except for zero or one letter from $C_i, i \in \{1, 2, \dots, n\}$, sprinkled here and there in some order. So $\Psi^{-1}(P)$ is closed in X^* . Because Ψ is quotient, P is closed in X^{\otimes} .

It also follows that any m-SRE is closed in X^{\otimes} .

Next we show that the m-products $F \mid C_1, C_2, \dots, C_n$ are indeed irreducible. By Lemma 6.7, $F^* C_1^? C_2^? \dots C_n^?$ is irreducible closed in X^* . So the closure of its image by Ψ is irreducible closed by Lemma A.1. However, the image of $F^* C_1^? C_2^? \dots C_n^?$ by Ψ is $F \mid C_1, C_2, \dots, C_n$. We have seen that it is closed, hence equal to its closure. Therefore it is irreducible closed.

Now assume X Noetherian. Let \mathcal{F} be any closed subset of X^{\otimes} . Since Ψ is continuous, $\Psi^{-1}(\mathcal{F})$ is closed in X^* , hence a finite union of word-products, by Proposition 6.14. Since Ψ is surjective, \mathcal{F} is equal to the image $\Psi[\Psi^{-1}(\mathcal{F})]$ of $\Psi^{-1}(\mathcal{F})$ by Ψ . So \mathcal{F} is a finite union of subsets $\Psi[P_i], i \in I$, where each P_i is a word-product. We calculate $\Psi[P_i]$ as follows. Write P_i as $e_1 e_2 \dots e_n$, and since it will not change its image by Ψ , reorder the atomic expressions e_i in $e_1 e_2 \dots e_n$ so that the starred ones come first. Doing so allows us to write our word-product as $F_1^* F_2^* \dots F_m^* C_1^? C_2^? \dots C_p^?$, up to permutation of factors. Its image by Ψ is the set of multisets obtained by picking at most one element from C_1 , at most one from C_2, \dots , at most one C_p , then arbitrarily many from F_1 , arbitrarily many from F_2, \dots , arbitrarily many from F_m . It follows that $\Psi[e_1 e_2 \dots e_n] = (F_1 \cup F_2 \cup \dots \cup F_m) \mid C_1, C_2, \dots, C_p$. In particular, $\Psi[P_i]$ is an m-product. Therefore \mathcal{F} is a finite union of m-products, hence an m-SRE.

If \mathcal{F} is also irreducible, then this finite union must be the union of a single m-product, hence is an m-product. \square

Lemma 7.6 (recap). Let X be a topological space. Inclusion between m-products can be checked in polynomial time, modulo an oracle testing inclusion of closed subsets of X .

Explicitly, let $P = F \mid C_1, C_2, \dots, C_m$ and $P' = F' \mid C'_1, C'_2, \dots, C'_n$ be two m-products. Let $I = \{i_1, i_2, \dots, i_k\}$ be the subset of those indices $i, 1 \leq i \leq m$, such that $C_i \not\subseteq F'$.

Then $P \subseteq P'$ if and only if $F \subseteq F'$ and there is an injective map $r: I \rightarrow \{1, 2, \dots, n\}$ such that $C_i \subseteq C'_{r(i)}$ for every $i \in I$ —in other words, $\{C_{i_1}, C_{i_2}, \dots, C_{i_k}\} \subseteq^{\otimes} \{C'_{r(i_1)}, C'_{r(i_2)}, \dots, C'_{r(i_k)}\}$.

Proof. Assume $P \subseteq P'$. If $F \not\subseteq F'$, then pick $x \in F \setminus F'$: the multiset consisting of $n + 1$ copies of x is in P but not in P' : contradiction. So $F \subseteq F'$.

Let now $I = \{i_1, i_2, \dots, i_k\}$ be as above. Let $D_1 = C_{i_1}, D_2 = C_{i_2}, \dots, D_k =$

C_{i_k} . Let also E_1, E_2, \dots, E_{m-k} be an enumeration of those C_i , $1 \leq i \leq n$, with $i \notin I$. Consider the word-product P_1 defined as $E_1^? E_2^? \dots E_{m-k}^? F^* D_1^? D_2^? \dots D_k^?$. Note that $P_1 \subseteq \Psi^{-1}(P)$, so $P_1 \subseteq \Psi^{-1}(P')$. On the other hand, $\Psi^{-1}(P')$ is the union over all permutations π of $\{1, 2, \dots, n\}$ of $F'^* C'_{\pi(1)} F'^* C'_{\pi(2)} F'^* \dots F'^* C'_{\pi(n)} F'^*$. Since P_1 is irreducible (Lemma 6.7), there a permutation π of $\{1, 2, \dots, n\}$ such that $P_1 \subseteq F'^* C'_{\pi(1)} F'^* C'_{\pi(2)} F'^* \dots F'^* C'_{\pi(n)} F'^*$. Using Lemma 6.9, and the fact that E_1, E_2, \dots, E_{m-k} are contained in F' , and $F \subseteq F'$, and recalling the definition of P_1 , we obtain that $D_1^? D_2^? \dots D_k^?$ is included in $F'^* C'_{\pi(1)} F'^* C'_{\pi(2)} F'^* \dots F'^* C'_{\pi(n)} F'^*$.

We show that there is an injective map $r: I \rightarrow \{\pi(1), \pi(2), \dots, \pi(n)\}$ such that $C_i \subseteq C'_{r(i)}$ for every $i \in I$, by induction on $k + n$. If $k = 0$, the empty map fits. Otherwise, since $D_1 \not\subseteq F'$, using Lemma 6.9, we must have $D_1^? D_2^? \dots D_k^? \subseteq C'_{\pi(1)} F'^* C'_{\pi(2)} F'^* \dots F'^* C'_{\pi(n)} F'^*$. Now we have two cases, again following Lemma 6.9. In the first case $D_1 = C_{i_1} \subseteq C'_{\pi(1)}$ and $D_2^? \dots D_k^? \subseteq F'^* C'_{\pi(2)} F'^* \dots F'^* C'_{\pi(n)} F'^*$, so there is an injective map $r': \{i_2, \dots, i_k\} \rightarrow \{\pi(2), \dots, \pi(n)\}$ such that $C_i \subseteq C'_{r'(i)}$ for every $i \in \{i_2, \dots, i_k\}$, by induction hypothesis. Then taking $r(i_1) = \pi(1)$ and $r(i) = r'(i)$ for every $i \in \{i_2, \dots, i_k\}$ fits. In the second case, $D_1^? D_2^? \dots D_k^? \subseteq F'^* C'_{\pi(2)} F'^* \dots F'^* C'_{\pi(n)} F'^*$, and we conclude directly by the induction hypothesis.

Conversely, if there is an injective map $r: I \rightarrow \{1, 2, \dots, n\}$ such that $C_i \subseteq C'_{r(i)}$ for every $i \in I$, it is clear that $P \subseteq P'$. \square

Theorem 7.7 (recap). Let X be a Noetherian space, $X' = X^\otimes$, and $(S, \llbracket _ \rrbracket, \trianglelefteq, \tau, \wedge)$ be an S-representation of X . Then $(S', \llbracket _ \rrbracket', \trianglelefteq', \tau', \wedge')$ is an S-representation of X' , where:

- (A) S' is the collection of all *m-product notations*, i.e., of all expressions of the form $A \mid u$, where A is a finite subset of S , and u is a multiset of elements of S . When $u = \{b_1, \dots, b_n\}$, we also write $A \mid b_1, \dots, b_n$ for $A \mid u$.
- (B) $\llbracket A \mid b_1, \dots, b_n \rrbracket' = (\bigcup_{a \in A} \llbracket a \rrbracket \mid \llbracket b_1 \rrbracket, \dots, \llbracket b_n \rrbracket)$.
- (C) $A \mid u \trianglelefteq' A' \mid u'$ if and only if $A \trianglelefteq^b A'$ and $u_1 \trianglelefteq^\otimes u'$ where u_1 is the subset of those elements $a \in u$ such that $a \trianglelefteq a'$ for no $a' \in A'$.
- (D) τ' is $\{\tau \mid \emptyset\}$.
- (E) \wedge' is defined as follows. A *matching* $f: \{1, \dots, m\} \rightarrow \{1, \dots, n\}$ is any bijection from some subset of $\{1, \dots, m\}$ (the *domain* $\text{dom } f$) to some subset of $\{1, \dots, n\}$ (the *codomain* $\text{cod } f$). Then $(A \mid a_1, \dots, a_m) \wedge' (A' \mid a'_1, \dots, a'_n)$ is the collection of all m-product notations of the form $A'' \mid m_{1f} \uplus m_{2f} \uplus m_{3f}$, where:

- $A'' = \bigcup_{\substack{a \in A \\ a' \in A'}} (a \wedge a')$;
- f ranges over all matchings from $\{1, \dots, m\}$ to $\{1, \dots, n\}$;
- m_{1f} ranges over all multisets of the form $\{c_i \mid i \in \text{dom } f\}$ where $c_i \in a_i \wedge a'_{f(i)}$ for every $i \in \text{dom } f$;

- m_{2f} ranges over all multisets of the form $\{c_i \mid 1 \leq i \leq m, i \notin \text{dom } f\}$, where $c_i \in \bigcup_{a' \in A'} (a_i \wedge a')$ for each $i, 1 \leq i \leq m, i \notin \text{dom } f$;
- m_{3f} ranges over all multisets of the form $\{c'_j \mid 1 \leq j \leq n, j \notin \text{cod } f\}$, where $c'_j \in \bigcup_{a \in A} (a \wedge a'_j)$ for each $j, 1 \leq j \leq n, j \notin \text{cod } f$.

Proof. First, $\llbracket _ \rrbracket'$ is surjective by Proposition 7.4. The formula for \leq' is justified by Lemma 7.6. The fact that $X^\otimes = \bigcup_{A|u \in \tau'} \llbracket A \mid u \rrbracket'$ is clear: the union on the right-hand side is $\llbracket \tau \mid \emptyset \rrbracket'$, which is by definition the set of multisets whose elements are all in $\bigcup_{a \in \tau} \llbracket a \rrbracket = X$.

To justify the formula for \wedge' , finally, compute the intersection of $\llbracket A \mid a_1, \dots, a_m \rrbracket' = F \mid \llbracket a_1 \rrbracket, \dots, \llbracket a_m \rrbracket$ (where $F = \bigcup_{a \in A} \llbracket a \rrbracket$) and of $\llbracket A' \mid a'_1, \dots, a'_n \rrbracket' = F' \mid \llbracket a'_1 \rrbracket, \dots, \llbracket a'_n \rrbracket$ (where $F' = \bigcup_{a' \in A'} \llbracket a' \rrbracket$). Any multiset m in the intersection can be split in four parts: first, the multiset m_0 of those elements that are in $F \cap F'$; then, among the remaining elements, the multiset m_1 of those elements that are both in some $\llbracket a_i \rrbracket$ and in some $\llbracket a'_j \rrbracket$: reasoning on indices, there must be a matching $f: \{1, \dots, m\} \rightarrow \{1, \dots, n\}$ such that m_1 is a multiset of elements $x_i, i \in \text{dom } f$, where x_i is in $\llbracket a_i \rrbracket \cap \llbracket a'_{f(i)} \rrbracket$; then, the remaining elements are obtained by taking at most one element from each $\llbracket a_i \rrbracket, i \notin \text{dom } f$, provided they are in F' , and at most one element from each $\llbracket a'_j \rrbracket, j \notin \text{cod } f$, provided they are in F . Let E_f be the set $(F \cap F') \mid \underbrace{\llbracket a_i \rrbracket \cap \llbracket a'_{f(i)} \rrbracket}_{i \in \text{dom } f}, \underbrace{\llbracket a_i \rrbracket \cap F'}_{\substack{1 \leq i \leq m \\ i \notin \text{dom } f}}, \underbrace{F \cap \llbracket a'_j \rrbracket}_{\substack{1 \leq j \leq n \\ j \notin \text{cod } f}}$. We have just shown that $\llbracket A \mid a_1, \dots, a_m \rrbracket' \cap$

$\llbracket A' \mid a'_1, \dots, a'_n \rrbracket'$ was contained in the union of all E_f , when f ranges over the matchings from $\{1, \dots, m\}$ to $\{1, \dots, n\}$. The converse inclusion is obvious.

We then observe that:

- $F \cap F' = \bigcup_{\substack{a \in A \\ a' \in A'}} (\llbracket a \rrbracket \cap \llbracket a' \rrbracket) = \bigcup_{\substack{a \in A \\ a' \in A' \\ c \in a \wedge a'}} \llbracket c \rrbracket = \bigcup_{c \in A''} \llbracket c \rrbracket$;
- for each $i \in \text{dom } f, \llbracket a_i \rrbracket \cap \llbracket a'_{f(i)} \rrbracket = \bigcup_{c_i \in a_i \wedge a'_{f(i)}} \llbracket c_i \rrbracket$;
- for each $i, 1 \leq i \leq m, i \notin \text{dom } f, \llbracket a_i \rrbracket \cap F' = \llbracket a_i \rrbracket \cap \bigcup_{a' \in A'} \llbracket a' \rrbracket = \bigcup_{a' \in A'} (\llbracket a_i \rrbracket \cap \llbracket a' \rrbracket) = \bigcup_{c_i \in a_i \wedge a'} \llbracket c_i \rrbracket$;
- and similarly, for each $j, 1 \leq j \leq n, j \notin \text{cod } f, F \cap \llbracket a'_j \rrbracket = \bigcup_{\substack{a \in A \\ c'_j \in a \wedge a'_j}} \llbracket c'_j \rrbracket$.

Finally, we notice that unions distribute over the \mid construction, meaning that $F \mid A \cup B, C_2, \dots, C_n$ is equal to the union of $F \mid A, C_2, \dots, C_n$ and $F \mid B, C_2, \dots, C_n$. By distributing all unions across the \mid construction, we obtain the indicated formula for \wedge' . \square

D Proofs of Results on Words with the Prefix Topology (Section 8)

Proposition 8.2 (recap). Let $X_1, X_2, \dots, X_n, \dots$ be countably many topological spaces. The sets of the form $\lceil F_1 F_2 \cdots F_n \rceil$, where each F_i is closed in X_i , form a subbase of closed sets for $\bigtriangleright_{n=1}^{+\infty} X_n$: these sets are closed, and every closed subset is an intersection of finite unions of such sets.

Proof. We first observe that the complement of $\lceil F_1 F_2 \cdots F_n \rceil$ is the open set $\lfloor \emptyset, X_1 \setminus F_1, X_1 X_2 \setminus F_1 F_2, \dots, X_1 X_2 \cdots X_m \setminus F_1 F_2 \cdots F_m, \dots, X_1 X_2 \cdots X_n \setminus F_1 F_2 \cdots F_n, X_1 X_2 \cdots X_n X_{n+1}, \dots, X_1 X_2 \cdots X_k, \dots \rceil = \lfloor \emptyset, U_1, U_1 X_2 \cup X_1 U_2, \dots, \bigcup_{i=1}^m X_1 \cdots X_{i-1} U_i X_{i+1} \cdots X_m, \dots, \bigcup_{i=1}^m X_1 \cdots X_{i-1} U_i X_{i+1} \cdots X_n, X_1 X_2 \cdots X_n X_{n+1}, \dots, X_1 X_2 \cdots X_k, \dots \rceil$, where U_i is the complement of F_i , $1 \leq i \leq n$.

Conversely, we claim that every open in the prefix topology is a union of subsets of the form ${}_m[V_1 V_2 \cdots V_m]_n$ ($0 \leq m \leq n$, V_i open in X_i for every i), where the latter denotes the set of all words w of length at least m such that either $|w| \geq n$ or the m -letter prefix of w is in $V_1 V_2 \cdots V_m$. Indeed, consider any wide telescope $\mathcal{U} = U_0, U_1, \dots, U_n, \dots$, where $U_k = \prod_{i=1}^k X_i$ for all $k \geq n$. \mathcal{U} certainly contains ${}_m[V_1 V_2 \cdots V_m]_n$ for any open rectangle $V_1 V_2 \cdots V_m$ contained in U_m : for any $w \in {}_m[V_1 V_2 \cdots V_m]_n$, either $|w| \geq n$, then $w \in \prod_{i=1}^{|w|} X_i = U_{|w|}$, or $m \leq |w| < n$, in which case w is in $V_1 V_2 \cdots V_m X_{m+1} \cdots X_{|w|}$, hence in $U_m X_{m+1} \cdots X_{|w|}$, and therefore in $U_{|w|}$ by the definition of telescopes. We claim that \mathcal{U} is equal to the union of all ${}_m[V_1 V_2 \cdots V_m]_n$, where $0 \leq m \leq n$, and $V_1 V_2 \cdots V_m$ ranges over the open rectangles contained in U_m . Indeed, given any word $w \in \lfloor \mathcal{U} \rceil$, either $|w| \geq n$ and we can take $m = n$, $V_1 = X_1, \dots, V_m = X_m$; or $|w| = m < n$, then w is in some open rectangle $V_1 V_2 \cdots V_m$ contained in U_m , by definition of the product topology on $\prod_{i=1}^m U_i$, whence $w \in {}_m[V_1 V_2 \cdots V_m]_n$.

Finally, we observe that the complement of ${}_m[V_1 V_2 \cdots V_m]_n$ is the set of words w such that either $|w| < m$, or $m \leq |w| < n$ and for some i , $1 \leq i \leq m$, the i th letter of w is not in V_i . When $m = 0$, this is empty. When $m = n$, the condition $m \leq |w| < n$ is always false, so the complement of ${}_m[V_1 V_2 \cdots V_m]_n$ is equal to $\lceil X^{m-1} \rceil$. Finally, when $m \neq 0$ and $m < n$, write F_i for the complement of V_i , then the complement of ${}_m[V_1 V_2 \cdots V_m]_n$ is equal to $\lceil X^{m-1} \rceil \cup \bigcup_{i=1}^m \lceil X^{i-1} F_i X^{n-i-1} \rceil$ if $m \geq 1$.

So the complement of any open of the prefix topology is an intersection of finite unions of the claimed subbasic closed sets, and we conclude. \square

We need the following, which also appears in [Goubault-Larrecq, 2013, Exercise 9.7.36].

Lemma D.1 *Let $X_1, X_2, \dots, X_n, \dots$ be countably many topological spaces. The map $i_n: X_1 \times \cdots \times X_n \rightarrow \bigtriangleright_{n=1}^{+\infty} X_n$ that sends each n -tuple (a_1, a_2, \dots, a_n) to the word $a_1 a_2 \cdots a_n$, and the map $cons: X_1 \times \bigtriangleright_{n=2}^{+\infty} X_n \rightarrow \bigtriangleright_{n=1}^{+\infty} X_n$ that sends $a_1, a_2 a_3 \cdots a_n$ to $a_1 a_2 a_3 \cdots a_n$ are both continuous.*

Proof. To show that i_n is continuous, we note that the inverse image of the open subset $\lfloor \mathcal{U} \rceil$, where \mathcal{U} is any telescope $U_0, U_1, \dots, U_n, \dots$, is the open subset U_n . To show that $cons$ is continuous, it is easier to show that the inverse image of a subbasic

closed set $\lceil F_1 F_2 \cdots F_n \rceil$ (see Proposition 8.2) is closed. Indeed, this inverse image is empty if $n = 0$, and equal to $F_1 \times \lceil F_2 \cdots F_n \rceil$ otherwise. \square

Lemma 8.3 (recap). Let $X_1, X_2, \dots, X_n, \dots$, be countably many topological spaces. The subsets of the form $\lceil C_1 C_2 \cdots C_n \rceil$, where C_i is irreducible closed in X_i for each i , $1 \leq i \leq n$, are irreducible closed in $\triangleright_{n=1}^{+\infty} X_n$.

Proof. $C_1 \times C_2 \times \cdots \times C_n$ is irreducible closed in $X_1 \times \cdots \times X_n$ by Lemma A.2. The map i_n is continuous, so $cl(i_n[C_1 \times C_2 \times \cdots \times C_n])$ is irreducible closed by Lemma A.1. We claim that the latter is exactly $\lceil C_1 C_2 \cdots C_n \rceil$, and this will prove the lemma. Indeed, $i_n[C_1 \times C_2 \times \cdots \times C_n]$ is contained in $\lceil C_1 C_2 \cdots C_n \rceil$, hence so does its closure. Conversely, any word in $\lceil C_1 C_2 \cdots C_n \rceil$ is also in the downward closure of $i_n[C_1 \times C_2 \times \cdots \times C_n]$ (with respect to \leq^p), hence in the set $cl(i_n[C_1 \times C_2 \times \cdots \times C_n])$, since the latter is closed hence downwards-closed. \square

Lemma 8.5 (recap). Let $X_1, X_2, \dots, X_n, \dots$, be countably many non-empty topological spaces. The whole space $\triangleright_{n=1}^{+\infty} X_n$ is irreducible closed in itself.

Proof. Passing to complements, it is equivalent to show that the intersection of two non-empty opens is again non-empty. Any two non-empty open subsets of $\triangleright_{n=1}^{+\infty} X_n$ are of the form $\lfloor \mathcal{U} \rfloor$ and $\lfloor \mathcal{U}' \rfloor$ for two wide telescopes $\mathcal{U} = (U_n)_{n \in \mathbb{N}}$ and $\mathcal{U}' = (U'_n)_{n \in \mathbb{N}}$. For n large enough, $U_n = U'_n = \prod_{i=1}^n X_i$, so any length n heterogeneous word is in $\lfloor \mathcal{U} \rfloor \cap \lfloor \mathcal{U}' \rfloor$. \square

To show that there is no other irreducible closed subset, we rest on the following general-purpose lemma.

Lemma D.2 *Let Y be a topological space, and \mathcal{B} be a subbase of closed sets of Y . Assume that any (possibly infinite) non-empty intersection of elements of \mathcal{B} can be written as a finite union of elements of \mathcal{B} . Then every irreducible closed subset of Y is a member of \mathcal{B} , or equals Y itself.*

Proof. Let C be an arbitrary irreducible closed subset of Y . As a closed set, C can be written as $\bigcap_{i \in I} \bigcup_{j \in J_i} F_{ij}$, where I is some index set, J_i is finite for every $i \in I$, and $F_{ij} \in \mathcal{B}$ for all i, j . If I is empty, then $C = Y$. Otherwise, for each $i \in I$, C is contained in $\bigcup_{j \in J_i} F_{ij}$, so C is contained in F_{ij_i} for some $j_i \in J_i$, since C is irreducible. Since F_{ij_i} is clearly contained in $\bigcup_{j \in J_i} F_{ij}$, $C = \bigcap_{i \in I} F_{ij_i}$. By assumption, C can be written as a finite union $\bigcup_{k=1}^n B_k$ of elements of \mathcal{B} . Since C is irreducible, again, C must equal some B_k , $1 \leq k \leq n$, whence $C \in \mathcal{B}$. \square

Lemma 8.6 (recap). Let $X_1, X_2, \dots, X_n, \dots$, be countably many non-empty topological spaces. The only irreducible closed subsets of $\triangleright_{n=1}^{+\infty} X_n$ are $\triangleright_{n=1}^{+\infty} X_n$ itself, and the subsets of the form $\lceil C_1 C_2 \cdots C_n \rceil$, where C_i is irreducible closed in X_i for each i , $1 \leq i \leq n$.

Proof. Let \mathcal{B} be the subbase of closed sets of the form $\lceil F_1 F_2 \cdots F_n \rceil$. Any non-empty intersection of such sets is again of this form. In fact, whenever I is non-empty, the intersection $\bigcap_{i \in I} \lceil F_{i1} F_{i2} \cdots F_{in_i} \rceil$ equals $\lceil \bigcap_{i \in I} F_{i1} \bigcap_{i \in I} F_{i2} \cdots \bigcap_{i \in I} F_{i \min_{j \in I} n_j} \rceil$. So Lemma D.2 applies: the only irreducible closed subsets of $\triangleright_{n=1}^{+\infty} X_n$ other than $\triangleright_{n=1}^{+\infty} X_n$ itself are in \mathcal{B} .

Now assume $\lceil F_1 F_2 \cdots F_n \rceil$ is irreducible. Without loss of generality, no F_i is empty: otherwise, letting k be the smallest index such that F_k is empty, one can rewrite

$\lceil F_1 F_2 \cdots F_n \rceil$ as $\lceil F_1 F_2 \cdots F_{k-1} \rceil$.

For each i , $1 \leq i \leq n$, since F_i is non-empty, fix an element x_i of F_i . We claim that F_i must be irreducible for each i , $1 \leq i \leq n$. Otherwise, there would be two closed subsets F' and F'' such that $F_i \subseteq F' \cup F''$, but F_i is contained neither in F' nor in F'' . In this case, let x' be an element of F_i outside F' , and x'' an element of F_i outside F'' . Then $x_1 x_2 \cdots x_{i-1} x'$ is in $\lceil F_1 F_2 \cdots F_i \cdots F_n \rceil$ but not in $\lceil F_1 F_2 \cdots F' \cdots F_n \rceil$ (where F' replaces F_i at position i), and $x_1 x_2 \cdots x_{i-1} x''$ is in $\lceil F_1 F_2 \cdots F_i \cdots F_n \rceil$ but not in $\lceil F_1 F_2 \cdots F'' \cdots F_n \rceil$ (where F'' replaces F_i at position i), although $\lceil F_1 F_2 \cdots F_i \cdots F_n \rceil$ is contained in the union $\lceil F_1 F_2 \cdots F' \cdots F_n \rceil \cup \lceil F_1 F_2 \cdots F'' \cdots F_n \rceil$. This would contradict the fact that $\lceil F_1 F_2 \cdots F_i \cdots F_n \rceil$ is irreducible. So each F_i is irreducible. \square

Proposition 8.7 (recap). Let $X_1, X_2, \dots, X_n, \dots$ be countably many non-empty topological spaces. The map $i: (\mathcal{D}_{n=1}^{+\infty} \mathcal{S}(X_n))^\top \rightarrow \mathcal{S}(\mathcal{D}_{n=1}^{+\infty} X_n)$ that sends \top to $\mathcal{D}_{n=1}^{+\infty} X_n$ and the word $C_1 C_2 \cdots C_n$ (where $C_i \in \mathcal{S}(X_i)$ for each i) to $\lceil C_1 C_2 \cdots C_n \rceil$ is an order-isomorphism and a homeomorphism.

Proof. First, i is well-defined, by Lemma 8.5 and Lemma 8.3. It is surjective by Lemma 8.6.

The specialization quasi-ordering on $(\mathcal{D}_{n=1}^{+\infty} \mathcal{S}(X_n))^\top$ is $\subseteq^{\triangleright \top}$.

Notice that $C_1 \cdots C_m \subseteq^{\triangleright \top} C'_1 \cdots C'_n$ iff $C_1 \cdots C_m \subseteq^{\triangleright} C'_1 \cdots C'_n$ iff $m \leq n$ and $C_1 \subseteq C'_1, \dots, C_m \subseteq C'_m$. We claim that this is equivalent to $\lceil C_1 \cdots C_m \rceil \subseteq \lceil C'_1 \cdots C'_n \rceil$. The only if direction is clear. In the if direction, pick $x_1 \in C_1, \dots, x_m \in C_m$: the word $x_1 \cdots x_m$ is in $\lceil C_1 \cdots C_m \rceil \subseteq \lceil C'_1 \cdots C'_n \rceil$, so $x_1 \in C'_1, \dots, x_m \in C'_m$. In particular, $m \leq n$; also, letting $x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_m$ remain fixed, but varying x_i in C_i , we obtain that $C_i \subseteq C'_i$.

Notice also that $\top \subseteq^{\triangleright \top} C'_1 \cdots C'_n$ never holds, and that $\mathcal{D}_{n=1}^{+\infty} X_n \subseteq \lceil C'_1 \cdots C'_n \rceil$ never holds either, since $\mathcal{D}_{n=1}^{+\infty} X_n$ contains words of arbitrary lengths.

Notice finally that $C_1 \cdots C_m \subseteq^{\triangleright \top} \top$ always holds, and correspondingly $\lceil C_1 \cdots C_m \rceil \subseteq \mathcal{D}_{n=1}^{+\infty} X_n$ always holds.

It follows that for every $w, w' \in (\mathcal{D}_{n=1}^{+\infty} \mathcal{S}(X_n))^\top$, $w \subseteq^{\triangleright \top} w'$ iff $i(w) \subseteq i(w')$. In particular, $i(w) = i(w')$ entails $w = w'$, so that i is injective. Since i is surjective, it is bijective. This also shows that i and its inverse are monotonic, so that i is an order-isomorphism.

For each closed subset F of a space X , write $\square F$ for the family of all irreducible closed subsets C of X such that $C \subseteq F$. Alternatively, $\square F$ is the complement of the open subset $\diamond U$ of $\mathcal{S}(X)$, for U the complement of F in X . In particular, $\square F$ is closed in $\mathcal{S}(X)$, and all closed subsets of $\mathcal{S}(X)$ are of this form. Since \diamond commutes with arbitrary unions and finite intersections, \square commutes with arbitrary intersections and finite unions, so given a subbase \mathcal{B} of closed subsets of X , the sets $\square F$ with $F \in \mathcal{B}$ form a subbase of the closed subsets of $\mathcal{S}(X)$. In particular, the sets $\square \lceil F_1, F_2, \dots, F_n \rceil$ with each F_i closed in X_i form a subbase of closed sets of $\mathcal{S}(\mathcal{D}_{n=1}^{+\infty} X_n)$. Their inverse image by i is the set of words $C_1 C_2 \cdots C_m$ in $\mathcal{D}_{n=1}^{+\infty} \mathcal{S}(X_n)$ such that $\lceil C_1, C_2, \dots, C_m \rceil \subseteq \lceil F_1, F_2, \dots, F_n \rceil$. This is equivalent to $C_1 \cdots C_m \subseteq^{\triangleright \top} F_1 F_2 \cdots F_n$, by an argument similar to one we have already seen at the beginning of the present proof. (The difference is that F_1, F_2, \dots, F_n are no

longer irreducible, contrarily to C'_1, \dots, C'_n —but we never used irreducibility there.) Therefore $i^{-1}(\square[F_1, F_2, \dots, F_n]) = \square[F_1, \square F_2, \dots, \square F_n]$. This shows that i is continuous.

Since i is bijective, let j be its inverse. We have just shown that $j^{-1}(\square[F_1, \square F_2, \dots, \square F_n]) = \square[F_1, F_2, \dots, F_n]$. The sets $\square F_i$, for F_i closed in X_i , span all the closed subsets of $\mathcal{S}(X_i)$, since their complements $\diamond U$ for U open span all the open subsets of $\mathcal{S}(X_i)$. Using Proposition 8.2, the sets $\square[F_1, \square F_2, \dots, \square F_n]$ with all F_i closed form a subbase of closed subsets of $\mathcal{D}_{n=1}^{+\infty} \mathcal{S}(X_n)$. Together with the whole set $(\mathcal{D}_{n=1}^{+\infty} \mathcal{S}(X_n))^\top$, they form a subbase of closed subsets of $(\mathcal{D}_{n=1}^{+\infty} \mathcal{S}(X_n))^\top$, whose inverse images by j are closed: either the closed set $\square[F_1, F_2, \dots, F_n]$, or the whole space $\mathcal{S}(\mathcal{D}_{n=1}^{+\infty} X_n)$. Therefore $j = i^{-1}$ is also continuous, hence i is a homeomorphism. \square

Lemma 8.8 (recap). Let X_1, X_2, \dots, X_n be non-empty topological spaces. The only irreducible closed subsets of $\mathcal{D}_{k=1}^n X_k$ are the subsets of the form $\square[C_1 C_2 \dots C_m]$, where C_i is irreducible closed in X_i for each i , $1 \leq i \leq m$, and $m \leq n$.

Proof. The proof of Lemma 8.6, with minor changes, shows that all irreducible closed subsets are of this form, except possibly for the whole space $\mathcal{D}_{k=1}^n X_k$. We show that the latter cannot be irreducible, unless it is itself of one of the above form.

Assume that $\mathcal{D}_{k=1}^n X_k$ is irreducible, that is, any two non-empty opens have a non-empty intersection. In particular, given any two non-empty open subsets U_k and V_k of X_k , the open subset $\langle \emptyset, \dots, \emptyset, X_1 \dots X_{k-1} U_k, X_1 \dots X_{k-1} U_k X_{k+1}, \dots, X_1 \dots X_{k-1} U_k X_{k+1} \dots X_n, \dots \rangle$ and the open subset $\langle \emptyset, \dots, \emptyset, X_1 \dots X_{k-1} V_k, X_1 \dots X_{k-1} V_k X_{k+1}, \dots, X_1 \dots X_{k-1} V_k X_{k+1} \dots X_n, \dots \rangle$ are both non-empty, and must have non-empty intersection. Any word in this intersection must be of length at least k , and its k th letter must be both in U_k and in V_k . So any two non-empty open subsets U_k and V_k of X_k must have non-empty intersection: X_k is irreducible. Then $\mathcal{D}_{k=1}^n X_k = \square[X_1 X_2 \dots X_n]$ is irreducible closed by Lemma 8.3, and we are done. \square

Proposition 8.9 (recap). Let X_1, X_2, \dots, X_n be non-empty topological spaces. The map $i: \mathcal{D}_{k=1}^n \mathcal{S}(X_k) \rightarrow \mathcal{S}(\mathcal{D}_{k=1}^n X_k)$ that sends the word $C_1 C_2 \dots C_k$ (where $k \leq n$ and $C_i \in \mathcal{S}(X_i)$ for each i) to $\square[C_1 C_2 \dots C_k]$ is an order-isomorphism and a homeomorphism.

Proof. The proof is as for Proposition 8.7, using now Lemma 8.8 instead of Lemma 8.6. \square

E Proofs of Results on Trees (Section 10)

It is sometimes convenient to be able to talk about subterms and positions p , together with the *subterm* $t|_p$ of t at position p . A *position* is a finite word over \mathbb{N} . The empty word ϵ is always a position in any term t , and $t|_\epsilon = t$. Whenever $t|_p$ is defined, and $t|_p$ is of the form $f(t_1, \dots, t_n)$, then p_i is a position in t for every i , $1 \leq i \leq n$, and $t|_{p_i} = t_i$. The *size* of a term is the number of its positions. We write $t[s]_p$ for the term t , except that the subterm at position p has been replaced by s .

The following generalizes the notion of simple tree expression: for U open in X and \mathcal{U} open in $\mathcal{T}(X)^*$, let $\diamond U \cdot \mathcal{U}$ be the set of all terms that have a subterm of the form

$f(\vec{t})$ with $f \in U$ and $\vec{t} \in \mathcal{U}$. We use them in proving the first part of Proposition 10.1:

Lemma E.1 *Let X be a topological space. Every finite intersection of simple tree expressions can be rewritten as a finite union of simple tree expressions. In particular, the simple tree expressions form a base of the tree topology.*

Proof. Let $Y = \mathcal{T}(X)$. The empty intersection is $\diamond X()$, and it remains to compute binary intersections. We do this in two steps. First, for all opens U and U' of X , for all opens \mathcal{U} and \mathcal{U}' of Y^* , we show that the intersection of $\diamond U \cdot \mathcal{U}$ and $\diamond U' \cdot \mathcal{U}'$ can be expressed as the union of simpler expressions of the form $\diamond U'' \cdot \mathcal{U}''$. Next, in the special case where $\diamond U \cdot \mathcal{U}$ is a simple tree expression $\diamond U(\pi_1 \mid \cdots \mid \pi_n)$, namely when \mathcal{U} is of the form $Y^* \pi_1 Y^* \cdots Y^* \pi_n Y^*$, and similarly for $\diamond U' \cdot \mathcal{U}'$, we show that each of the simpler expressions $\diamond U'' \cdot \mathcal{U}''$ obtained in step 1 can themselves be expressed as finite unions of simple tree expressions.

Step 1. The intersection of $\diamond U \cdot \mathcal{U}$ and $\diamond U' \cdot \mathcal{U}'$ is the union of:

1. $\diamond U \cdot (\mathcal{U} \cap (Y^*(\diamond U' \cdot \mathcal{U}')Y^*))$,
2. $\diamond U' \cdot (\mathcal{U}' \cap (Y^*(\diamond U \cdot \mathcal{U})Y^*))$,
3. $\diamond(U \cap U') \cdot (\mathcal{U} \cap \mathcal{U}')$,
4. $\diamond X \cdot (\diamond U \cdot \mathcal{U} \mid \diamond U' \cdot \mathcal{U}')$
5. $\diamond X \cdot (\diamond U' \cdot \mathcal{U}' \mid \diamond U \cdot \mathcal{U})$.

Indeed, the terms t of $\diamond U \cdot \mathcal{U} \cap \diamond U' \cdot \mathcal{U}'$ are those that have a subterm $t_{|p} = f(\vec{t})$ with $f \in U$ and $\vec{t} \in \mathcal{U}$ and that have a subterm $t_{|p'} = f'(\vec{t}')$ with $f' \in U'$ and $\vec{t}' \in \mathcal{U}'$, for some positions p and p' . If p is a proper prefix of p' (we say that $f(\vec{t})$ is *above* $f'(\vec{t}')$), then t is in $\diamond U \cdot (\mathcal{U} \cap (Y^*(\diamond U' \cdot \mathcal{U}')Y^*))$ (case 1 above); if p' instead is a proper prefix of p , then t is in case 2; if $p = p'$, then t is in case 3; if p and p' are incomparable, then t is in case 4 if $f(\vec{t})$ is to the left of $f'(\vec{t}')$ (i.e., the first element that differs in p and p' is less in p than in p'), and in case 5 if $f(\vec{t})$ is to the right of $f'(\vec{t}')$. Conversely, each of the opens 1–5 are clearly contained both in $\diamond U \cdot \mathcal{U}$ and in $\diamond U' \cdot \mathcal{U}'$.

Note. The operator $\diamond V \cdot$ commutes with finite unions, i.e., $\diamond V \cdot \bigcup_{i=1}^m \mathcal{V}_i = \bigcup_{i=1}^m \diamond V \cdot \mathcal{V}_i$. This is easy: both sides are the set of terms such that there is a subterm $f(\vec{t})$ with $f \in V$ and there is an i , $1 \leq i \leq m$, such that $\vec{t} \in \mathcal{V}_i$. We will use that freely below.

Step 2. Call an *elementary open* any open subset of Y^* of the form $Y^* \pi_1 Y^* \cdots Y^* \pi_n Y^*$, where π_1, \dots, π_n are simple tree expressions. We claim that, for all open subsets U and U' of X , for all elementary opens \mathcal{U} and \mathcal{U}' , $\diamond U \cdot \mathcal{U} \cap \diamond U' \cdot \mathcal{U}'$ is a finite union of (denotations of) simple tree expressions. We show this by induction over the size of the expressions $\diamond U \cdot \mathcal{U}$ and $\diamond U' \cdot \mathcal{U}'$. Write \mathcal{U} as $Y^* \pi_1 Y^* \cdots Y^* \pi_m Y^*$, and \mathcal{U}' as $Y^* \pi'_1 Y^* \cdots Y^* \pi'_n Y^*$. Then $\mathcal{U} \cap (Y^*(\diamond U' \cdot \mathcal{U}')Y^*)$ (case 1) is the union of all elementary opens of the form $Y^* \pi_1 Y^* \cdots Y^* \pi_{i-1} Y^* (\diamond U' \cdot \mathcal{U}') Y^* \pi_i Y^* \cdots Y^* \pi_m Y^*$, $1 \leq i \leq m+1$, plus all opens of the form $Y^* \pi_1 Y^* \cdots Y^* (\pi_i \cap \diamond U' \cdot \mathcal{U}') Y^* \cdots Y^* \pi_m Y^*$, $1 \leq i \leq m$. For each i , the latter is the (finite) union of the elementary opens $Y^* \pi_1 Y^* \cdots Y^* \pi'' Y^* \cdots Y^* \pi_m Y^*$, where π'' ranges over the (finitely many) simple

tree expressions given by the induction hypothesis, and whose union equals $\pi_i \cap \diamond U' \cdot U'$. So $\mathcal{U} \cap (Y^*(\diamond U' \cdot U')Y^*)$ is a (finite) union of elementary opens, say $\mathcal{U}_1, \dots, \mathcal{U}_k$: then the open $\diamond U \cdot (\mathcal{U} \cap (Y^*(\diamond U' \cdot U')Y^*))$ of case 1 is the (finite) union $\bigcup_{i=1}^k \diamond U \cdot \mathcal{U}_i$. Case 2 is symmetric.

Cases 4 and 5 are already in the form of simple tree expressions.

For case 3, we show that $\mathcal{U} \cap \mathcal{U}'$ is a finite union of elementary opens, by induction on $m + n$, using the formulae that we have already used in the proof of Lemma 6.1. When $m = 0$ or $n = 0$, this is clear. Otherwise, write $\mathcal{U} = Y^*\pi_1\mathcal{V}$, $\mathcal{U}' = Y^*\pi'_1\mathcal{V}'$, so $\mathcal{U} \cap \mathcal{U}'$ is the union of $Y^*\pi_1(\mathcal{V} \cap Y^*\pi'_1\mathcal{V}')$, of $Y^*\pi'_1(Y^*\pi_1\mathcal{V} \cap \mathcal{V}')$, and of $Y^*(\pi_1 \cap \pi'_1)(\mathcal{V} \cap \mathcal{V}')$. By induction hypothesis, $\mathcal{V} \cap Y^*\pi'_1\mathcal{V}'$ is a finite union of elementary opens, so $Y^*\pi_1(\mathcal{V} \cap Y^*\pi'_1\mathcal{V}')$ is, too, since unions distribute over concatenations. Similarly for $Y^*\pi'_1(Y^*\pi_1\mathcal{V} \cap \mathcal{V}')$. For $Y^*(\pi_1 \cap \pi'_1)(\mathcal{V} \cap \mathcal{V}')$, $\pi_1 \cap \pi'_1$ is a finite union of simple tree expressions by induction hypothesis (the first one, on π and π'), and $\mathcal{V} \cap \mathcal{V}'$ is a finite union of elementary opens by induction hypothesis (the second one, on \mathcal{U} and \mathcal{U}'). We then distribute unions over concatenations again to conclude. \square

Lemma E.2 *For every open subset U of X , for every open subset \mathcal{U} of $\mathcal{T}(X)^*$, $\diamond U \cdot \mathcal{U}$ is open in $\mathcal{T}(X)$.*

Proof. Let $Y = \mathcal{T}(X)$. Notice that \mathcal{U} can be written as a union of opens of the form $Y^*U_1Y^* \cdots Y^*U_nY^*$, where each U_i is open in Y , because those form a base of the word topology on Y^* (Lemma 6.1). By Lemma E.1, U_1, \dots, U_n can all be written as unions of simple tree expressions. Distributing unions over concatenations, \mathcal{U} is then a union of elementary opens \mathcal{U}_i of Y^* , $i \in I$ (in the sense already used in Lemma E.1). Then $\diamond U \cdot \mathcal{U} = \bigcup_{i \in I} \diamond U \cdot \mathcal{U}_i$ is a union of simple tree expressions, hence is open. \square

Lemma E.3 *Letting \leq be the specialization quasi-ordering of X , every open subset of $\mathcal{T}(X)$ is upwards-closed with respect to \leq .*

Proof. This is Exercise 9.7.43 of [Goubault-Larrecq, 2013]. For short, let Y denote $\mathcal{T}(X)$. We show that whenever $s \leq t$ and $s \in \pi$, then $t \in \pi$. This is by induction on the structure of π . Write π as $\diamond U(\pi_1 \mid \dots \mid \pi_n)$. There must be a subterm, say at position p , of s , of the form $f'(\vec{s}')$, with $f' \in U$ and $\vec{s}' \in Y^*\pi_1Y^* \cdots Y^*\pi_nY^*$. Let (*) be our induction hypothesis: whenever $u \leq v$ and $u \in \pi_i$ for some i , $1 \leq i \leq n$, then $v \in \pi_i$.

We show that $t \in \pi$ for any term t such that $s \leq t$, whenever s contains a subterm at some position p of the form $f'(\vec{s}')$, with $f' \in U$ and $\vec{s}' \in Y^*\pi_1Y^* \cdots Y^*\pi_nY^*$, by a secondary induction on the size of t . If $s \leq t$ by the first case of the definition, i.e., if $t = g(t_1, \dots, t_p)$ and $s \leq t_j$ for some j , $1 \leq j \leq p$, then by induction hypothesis $t_j \in \pi$, from which $t \in \pi$ follows immediately, by definition of (the denotation of) π .

So assume that $s \leq t$ by the second case of the definition, i.e., $s = f(\vec{s})$, $t = g(\vec{t})$, $f \leq g$, and $\vec{s} \leq \vec{t}$. Write \vec{s} as $s_1s_2 \dots s_m$, \vec{t} as $t_1t_2 \dots t_n$, so that there is an (injective) increasing map $h: \{1, 2, \dots, m\} \rightarrow \{1, 2, \dots, n\}$ with $s_1 \leq t_{h(1)}$, $s_2 \leq t_{h(2)}$, \dots , $s_m \leq t_{h(m)}$.

If $p = \varepsilon$, then $f' = f$ and $\vec{s}' = \vec{s}$, so $\vec{s} \in Y^*\pi_1Y^* \cdots Y^*\pi_nY^*$. Let $s_{i_1} \dots s_{i_n}$ be a subword of \vec{s} satisfying $s_{i_1} \in \pi_1, \dots, s_{i_n} \in \pi_n$. Then $t_{h(i_1)} \in \pi_1, \dots, t_{h(i_n)} \in \pi_n$

by $(*)$, and $t_{h(i_1)} \dots t_{h(i_n)}$ forms a subword of \vec{t} . So $\vec{t} \in Y^* \pi_1 Y^* \dots Y^* \pi_n Y^*$; since U is upwards-closed, $g \in U$; so $t \in \pi$.

Finally, if $p \neq \varepsilon$, then $f'(\vec{s}')$ must be a subterm of some s_i , $1 \leq i \leq m$. Since $s_i \preceq t_{h(i)}$, by induction hypothesis, $t_{h(i)}$ is in π , from which $t \in \pi$ follows immediately. \square

Lemma E.4 *For every open subset U of X , every open subset \mathcal{U} of $\mathcal{T}(X)^*$, and every open subset V of $\mathcal{T}(X)$, let $\diamond U \cdot \mathcal{U} // V$ be the set of all terms containing a subterm $f(\vec{t}) \in V$ with $f \in U$ and $\vec{t} \in \mathcal{U}$. Then $\diamond U \cdot \mathcal{U} // V$ is open in $\mathcal{T}(X)$.*

Proof. This is the first part of Exercise 9.7.44 of [Goubault-Larrecq, 2013]. Let $Y = \mathcal{T}(X)$. We first note that $\diamond U \cdot \mathcal{U} // V$ is open if V is (the denotation of) a simple tree expression $\pi = \diamond U'(\pi'_1 \mid \dots \mid \pi'_n)$: in that case $\diamond U \cdot \mathcal{U} // \pi = \diamond(U \cap U') \cdot (\mathcal{U} \cap Y^* \pi'_1 Y^* \dots Y^* \pi'_n Y^*) \cup \diamond U \cdot (\mathcal{U} \cap Y^* \pi Y^*)$. This is open by Lemma E.2. In the general case, by Lemma E.1, we can write V as a union of simple tree expressions π_i , $i \in I$. We conclude that $\diamond U \cdot \mathcal{U} // V = \bigcup_{i \in I} \diamond U \cdot \mathcal{U} // \pi_i$ is indeed open. \square

Lemma E.5 *For every closed subset F of X , and all closed subsets $\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_n$ of $Y = \mathcal{T}(X)$, let $F^{\vec{?}}(\mathcal{F}_1^{\vec{?}} \mathcal{F}_2^{\vec{?}} \dots \mathcal{F}_n^{\vec{?}})$ denote the union of $\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_n$ with the set of those terms $f(\vec{t})$ such that $f \in F$ and $\vec{t} \in \mathcal{F}_1^{\vec{?}} \mathcal{F}_2^{\vec{?}} \dots \mathcal{F}_n^{\vec{?}}$. (Recall the word products of Section 6.) Then $F^{\vec{?}}(\mathcal{F}_1^{\vec{?}} \mathcal{F}_2^{\vec{?}} \dots \mathcal{F}_n^{\vec{?}})$ is closed in Y : let $U = X \setminus F$, V be the complement of $\mathcal{F}_1 \cup \mathcal{F}_2 \cup \dots \cup \mathcal{F}_n$ in Y , \mathcal{U} be the complement of $\mathcal{F}_1^{\vec{?}} \mathcal{F}_2^{\vec{?}} \dots \mathcal{F}_n^{\vec{?}}$ in Y^* , then $F^{\vec{?}}(\mathcal{F}_1^{\vec{?}} \mathcal{F}_2^{\vec{?}} \dots \mathcal{F}_n^{\vec{?}})$ is the complement of $(\diamond X \cdot \mathcal{U} // V) \cup (\diamond U \cdot Y^* // V)$.*

Proof. This is the second part of Exercise 9.7.44 of [Goubault-Larrecq, 2013]. Let us characterize the complement of $F^{\vec{?}}(\mathcal{F}_1^{\vec{?}} \mathcal{F}_2^{\vec{?}} \dots \mathcal{F}_n^{\vec{?}})$. Note that U and V are open by definition, while \mathcal{U} is open by Corollary 6.6. Then $\diamond X \cdot \mathcal{U} // V$ and $\diamond U \cdot Y^* // V$ are open, as we have seen in Lemma E.4.

Let t be a term outside $F^{\vec{?}}(\mathcal{F}_1^{\vec{?}} \mathcal{F}_2^{\vec{?}} \dots \mathcal{F}_n^{\vec{?}})$. Since t is not in $F^{\vec{?}}(\mathcal{F}_1^{\vec{?}} \mathcal{F}_2^{\vec{?}} \dots \mathcal{F}_n^{\vec{?}})$, we first observe that t is in V , otherwise it would be in $\mathcal{F}_1 \cup \mathcal{F}_2 \cup \dots \cup \mathcal{F}_n$. Write t as $f(\vec{t})$. If $f \in F$, then \vec{t} cannot be in $\mathcal{F}_1^{\vec{?}} \mathcal{F}_2^{\vec{?}} \dots \mathcal{F}_n^{\vec{?}}$. Recall that \mathcal{U} is the complement of $\mathcal{F}_1^{\vec{?}} \mathcal{F}_2^{\vec{?}} \dots \mathcal{F}_n^{\vec{?}}$. So, if $f \in F$, then t is in $\diamond X \cdot \mathcal{U} // V$. If on the other hand $f \notin F$, then t is in $\diamond U \cdot Y^* // V$.

Conversely, consider any element t of $(\diamond X \cdot \mathcal{U} // V) \cup (\diamond U \cdot Y^* // V)$. We claim that t cannot be in $F^{\vec{?}}(\mathcal{F}_1^{\vec{?}} \mathcal{F}_2^{\vec{?}} \dots \mathcal{F}_n^{\vec{?}})$. Notice first that any subterm of a term in $F^{\vec{?}}(\mathcal{F}_1^{\vec{?}} \mathcal{F}_2^{\vec{?}} \dots \mathcal{F}_n^{\vec{?}})$ is again in $F^{\vec{?}}(\mathcal{F}_1^{\vec{?}} \mathcal{F}_2^{\vec{?}} \dots \mathcal{F}_n^{\vec{?}})$: this follows easily from the definition, and the fact that $\mathcal{F}_1 \cup \mathcal{F}_2 \cup \dots \cup \mathcal{F}_n$, being closed, is downwards-closed with respect to \preceq (as a consequence of Lemma E.3), hence is closed under taking subterms. Now let t be both in $(\diamond X \cdot \mathcal{U} // V) \cup (\diamond U \cdot Y^* // V)$ and in $F^{\vec{?}}(\mathcal{F}_1^{\vec{?}} \mathcal{F}_2^{\vec{?}} \dots \mathcal{F}_n^{\vec{?}})$. If t is in $\diamond X \cdot \mathcal{U} // V$, then t has a subterm $f(\vec{t}) \in V$ with $\vec{t} \in \mathcal{U}$. Since t is in $F^{\vec{?}}(\mathcal{F}_1^{\vec{?}} \mathcal{F}_2^{\vec{?}} \dots \mathcal{F}_n^{\vec{?}})$, its subterm $f(\vec{t})$ is in $F^{\vec{?}}(\mathcal{F}_1^{\vec{?}} \mathcal{F}_2^{\vec{?}} \dots \mathcal{F}_n^{\vec{?}})$, too. But since $\vec{t} \in \mathcal{U}$, $\vec{t} \notin \mathcal{F}_1^{\vec{?}} \mathcal{F}_2^{\vec{?}} \dots \mathcal{F}_n^{\vec{?}}$, so $f(\vec{t})$ must be in $\mathcal{F}_1 \cup \mathcal{F}_2 \cup \dots \cup \mathcal{F}_n$; but this would contradict the fact that $f(\vec{t}) \in V$. If t is instead in $\diamond U \cdot Y^* // V$, then t has a subterm $f(\vec{t}) \in V$ with $f \in U$, i.e., $f \notin F$. Again $f(\vec{t})$ is in $F^{\vec{?}}(\mathcal{F}_1^{\vec{?}} \mathcal{F}_2^{\vec{?}} \dots \mathcal{F}_n^{\vec{?}})$, and $f \notin F$ entails that $f(\vec{t})$ must be in $\mathcal{F}_1 \cup \mathcal{F}_2 \cup \dots \cup \mathcal{F}_n$, again contradicting $f(\vec{t}) \in V$.

So $F^{\bar{?}}(\mathcal{F}_1^? \mathcal{F}_2^? \dots \mathcal{F}_n^?)$ is the complement of $(\diamond X \cdot \mathcal{U} // V) \cup (\diamond U \cdot Y^* // V)$. Since the latter is open, the former is closed. \square

Proposition 10.1 (recap). Let X be a topological space. Every finite intersection of simple tree expressions can be rewritten as a finite union of simple tree expressions. In particular, the simple tree expressions form a base of the tree topology.

Letting \leq be the specialization quasi-ordering of X , the specialization quasi-ordering of $\mathcal{T}(X)$ is the embedding quasi-ordering \preceq_{\leq} .

Proof. The first part is Lemma E.1. For the second part, let \preceq denote temporarily the specialization quasi-ordering of $\mathcal{T}(X)$. Using Lemma E.3, $t \preceq_{\leq} t'$ implies $t \preceq t'$. Conversely, we show by structural induction on t' that its downward closure $\downarrow_{\preceq_{\leq}} t'$ in \preceq_{\leq} is closed: write t' as $f(t'_1, t'_2, \dots, t'_n)$, then $\downarrow_{\preceq_{\leq}} t'$ is equal to $(\downarrow_X f)^?((\downarrow_{\preceq_{\leq}} t'_1)^?)^? (\downarrow_{\preceq_{\leq}} t'_2)^? \dots (\downarrow_{\preceq_{\leq}} t'_n)^?)^?$, which is closed by Lemma E.5 and the induction hypothesis. So, if $t \preceq t'$ then t is in the closure of $\{t'\}$, hence in the closed set $\downarrow_{\preceq_{\leq}} t'$, whence $t \preceq_{\leq} t'$. \square

Proposition 10.2 (recap). Let X be a set quasi-ordered by \leq . The tree topology on $\mathcal{T}(X_a)$ is exactly the Alexandroff topology of \preceq_{\leq} on $\mathcal{T}(X)$.

Proof. Any upwards-closed subset A of $Y = \mathcal{T}(X)$ is the union of $\uparrow_Y s$, $s \in A$, where upward closure is taken relative to \preceq_{\leq} . We claim that $\uparrow_Y s$ is obtained recursively by $\uparrow_Y s = \diamond(\uparrow_X f)(\uparrow_Y s_1 \mid \dots \mid \uparrow_Y s_m)$, where $s = f(s_1, \dots, s_m)$. Indeed, let $t = g(t_1, \dots, t_n)$ be any term. We first show that if $t \in \uparrow_Y s$, i.e., if $s \preceq_{\leq} t$, then $t \in \diamond(\uparrow_X f)(\uparrow_Y s_1 \mid \dots \mid \uparrow_Y s_m)$: s clearly belongs to $\diamond(\uparrow_X f)(\uparrow_Y s_1 \mid \dots \mid \uparrow_Y s_m)$, and since open sets are upwards-closed in the specialization quasi-ordering, which is \preceq_{\leq} by Proposition 10.1, t is also in $\diamond(\uparrow_X f)(\uparrow_Y s_1 \mid \dots \mid \uparrow_Y s_m)$. Conversely, if $t \in \diamond(\uparrow_X f)(\uparrow_Y s_1 \mid \dots \mid \uparrow_Y s_m)$, then either $t = g(\vec{t})$ is itself the subterm such that $g \in \uparrow_X f$ and $\vec{t} \in Y^*(\uparrow_Y s_1)Y^* \dots Y^*(\uparrow_Y s_m)Y^*$, so $f \leq g$ and $\vec{s} \preceq_{\leq}^* \vec{t}$, in particular $s \preceq_{\leq} t$; or t_j is in $\diamond(\uparrow_X f)(\uparrow_Y s_1 \mid \dots \mid \uparrow_Y s_m)$ for some j , $1 \leq j \leq n$, so $s \preceq_{\leq} t_j$ by induction hypothesis, therefore $s \preceq_{\leq} t$.

It follows that $\uparrow_Y s$ is open in the tree topology for every $s \in A$, so $\uparrow_Y A$ is also open in the tree topology. Conversely, every open subset if upwards-closed in \preceq_{\leq} by Proposition 10.1. So the tree topology is the Alexandroff topology of \preceq_{\leq} . \square

E.1 Tree Steps

Lemma E.6 *The application map $@: X \times \mathcal{T}(X)^* \rightarrow \mathcal{T}(X)$, which sends (f, \vec{t}) to $f(\vec{t})$, is continuous.*

Proof. This is the first part of Exercise 9.7.47 of [Goubault-Larrecq, 2013]. Let $Y = \mathcal{T}(X)$, and $\pi = \diamond U(\pi_1, \dots, \pi_n)$ be a simple tree expression. We show that $@^{-1}(\pi)$ is open: $@^{-1}(\pi)$ is the union of the open $U \times (Y^* \pi_1 Y^* \dots Y^* \pi_n Y^*)$ with $X \times (Y^* \pi Y^*)$; indeed, $f(\vec{t}) \in \pi$ iff either $f \in U$ and $\vec{t} \in Y^* \pi_1 Y^* \dots Y^* \pi_n Y^*$ (case where the needed subterm of $f(\vec{t})$ that witnesses the fact that $f(\vec{t})$ is in π is $f(\vec{t})$ itself), or some element of the sequence \vec{t} is in π . So $@^{-1}(\pi)$ is open, and therefore $@$ is continuous. \square

Lemma 10.4 (recap). Let X be a topological space. For every closed subset F of

X , and every word-product \vec{P} on $\mathcal{T}(X)$, $\text{supp } \vec{P}$ and $F^{\vec{P}}$ are closed in $\mathcal{T}(X)$. If moreover $F = C$ is irreducible, then so is the tree step $C^{\vec{P}}$.

Proof. Let $Y = \mathcal{T}(X)$. The first part of the Lemma is a slight extension of Lemma E.5.

First, $\text{supp } \vec{P}$ is just $i^{-1}(\vec{P})$, where $i: Y \rightarrow Y^*$ is the continuous map such that $i(t)$ is the word with just one letter, t . By Corollary 6.6, \vec{P} is closed, so $\text{supp } \vec{P}$ is closed too.

Let t be a term outside $F^{\vec{P}}$. Let V be the complement of $\text{supp } \vec{P}$. We have just seen that V is open. Moreover, since t is not in $F^{\vec{P}}$, t is not in $\text{supp } \vec{P}$, so t is in V . Write t as $f(\vec{t})$. If $f \in F$, then \vec{t} cannot be in \vec{P} . Let \mathcal{U} be the complement of \vec{P} in $\mathcal{T}(X)^*$. This is open. So, if $f \in F$, then t is in $\diamond X \cdot \mathcal{U} // V$, which is open by Lemma E.4. If on the other hand $f \notin F$, then t is in $\diamond U \cdot Y^* // V$, where U is the complement of F in X ; therefore $t \in (\diamond X \cdot \mathcal{U} // V) \cup (\diamond U \cdot Y^* // V)$.

Conversely, consider any element t of $(\diamond X \cdot \mathcal{U} // V) \cup (\diamond U \cdot Y^* // V)$. We claim that t cannot be in $F^{\vec{P}}$. Notice that any subterm of a term in $F^{\vec{P}}$ is again in $F^{\vec{P}}$: this follows easily from the definition, and the fact that $\text{supp } \vec{P}$, being closed, is downwards-closed with respect to \preceq_{\leq} , hence is closed under taking subterms. So let t be both in $(\diamond X \cdot \mathcal{U} // V) \cup (\diamond U \cdot Y^* // V)$ and in $F^{\vec{P}}$. If t is in $\diamond X \cdot \mathcal{U} // V$, then t has a subterm $f(\vec{t}) \in V$ with $\vec{t} \in \mathcal{U}$. Since t is in $F^{\vec{P}}$, its subterm $f(\vec{t})$ is in $F^{\vec{P}}$, too. But since $\vec{t} \in \mathcal{U}$, $\vec{t} \notin \vec{P}$, so $f(\vec{t})$ must be in $\text{supp } \vec{P}$; this contradicts the fact that $f(\vec{t}) \in V$. If t is instead in $\diamond U \cdot Y^* // V$, then t has a subterm $f(\vec{t}) \in V$ with $f \in U$, i.e., $f \notin F$. Again $f(\vec{t})$ is in $F^{\vec{P}}$, and $f \notin F$ entails that $f(\vec{t})$ must be in $\text{supp } \vec{P}$, again contradicting $f(\vec{t}) \in V$.

So $F^{\vec{P}}$ is the complement of $(\diamond X \cdot \mathcal{U} // V) \cup (\diamond U \cdot Y^* // V)$. Since the latter is open, the former is closed.

Let us now assume that $F = C$ is irreducible.

By Lemma 6.7, the word-product \vec{P} over $\mathcal{T}(X)$ is irreducible closed in $\mathcal{T}(X)^*$. So $C \times \vec{P}$ is irreducible closed in $X \times \mathcal{T}(X)^*$, since the product of two irreducible closed subsets is irreducible closed (Lemma A.2). Since $@$ is continuous (Lemma E.6), one concludes that $cl(@[C \times \vec{P}])$ is irreducible closed in $\mathcal{T}(X)$ (Lemma A.1). Since $C^{\vec{P}}$ is closed, and clearly contains $@[C \times \vec{P}]$, it contains $cl(@[C \times \vec{P}])$. Conversely, the latter contains $\{f(\vec{t}) \mid f \in C, \vec{t} \in \vec{P}\}$, and is downwards-closed in \preceq_{\leq} , so is closed under taking subterms, whence $cl(@[C \times \vec{P}])$ contains $C^{\vec{P}}$. Therefore the latter is the closure $cl(@[C \times \vec{P}])$, and must then be irreducible. \square

E.2 Tree Iterators

We need the following lemma to show that tree iterators define closed sets. A relation R from a space X to a space Y is a subset of $X \times Y$. It is *lower semi-continuous* iff $\text{Pre}^{\exists} R(V) = \{x \in X \mid \exists y \in V \cdot x R y\}$ is open for every open subset V of Y . It is *upper semi-continuous* iff $\text{Pre}^{\forall} R(V) = \{x \in X \mid \forall y \cdot x R y \Rightarrow y \in V\}$ is open for every open subset V of Y . It is *continuous* if and only if it is both lower semi-continuous and upper semi-continuous.

Lemma E.7 Let Z be a topological space, \square be a hole outside Z , and *inst-of* be the relation from Z^* to $(Z + \{\square\})^*$ defined by w *inst-of* w' iff w is obtained from w' by replacing each occurrence of \square by (possibly distinct) elements from Z . Formally, iff w and w' have the same length and for every index i , the i th letter of w' is either \square or equal to the i th letter of w .

Then *inst-of* is continuous.

Proof. For short, let Y be $Z + \{\square\}$.

Lower semi-continuity. Consider any basic open $V = Y^*V_1Y^* \cdots Y^*V_nY^*$ of Y . Let $U_i = V_i \subseteq Z$ if $\square \notin V_i$, $U_i = Z$ otherwise. Then $\text{Pre}^{\exists} \text{inst-of}(V) = Z^*U_1Z^* \cdots Z^*U_nZ^*$ is open. Since every open subset of Y is a union of basic open sets, and since $\text{Pre}^{\exists} \text{inst-of}$ commutes with unions, $\text{Pre}^{\exists} \text{inst-of}(V)$ is open for every open subset V of Y^* .

Upper semi-continuity. We first observe that, for every word-product $P = e_1e_2 \cdots e_n$ on Y , $\text{Pre}^{\exists} \text{inst-of}(P)$ is a word-product on Z . Indeed, $\text{Pre}^{\exists} \text{inst-of}(P)$ is equal to $\text{Pre}^{\exists} \text{inst-of}(e_1)\text{Pre}^{\exists} \text{inst-of}(e_2) \cdots \text{Pre}^{\exists} \text{inst-of}(e_n)$, while $\text{Pre}^{\exists} \text{inst-of}(F^?)$ equals $Z^?$ if $\square \in F$ and $F^?$ otherwise, and $\text{Pre}^{\exists} \text{inst-of}(F^*)$ equals Z^* if $\square \in F$ and F^* otherwise. Call a *monotone Boolean combination* of word-products any finite union of finite intersections of word-products. Lemma 6.12 shows that any finite intersection of word-products can be rewritten as a finite union of word-products. So the monotone Boolean combinations of word-products are the finite unions of word-products $\bigcup_{i=1}^m P_i$. Now $\text{Pre}^{\exists} \text{inst-of}$ commutes with unions, so $\text{Pre}^{\exists} \text{inst-of}(F)$ is a finite union of word-products on Z (hence closed in Z^* by Corollary 6.6) for every monotone Boolean combination F of word-products on Y .

Using Lemma 6.2, the complement of any monotone Boolean combination U of basic opens of Y^* is a monotone Boolean combination of word-products. So $\text{Pre}^{\forall} \text{inst-of}(U)$, which is the complement of $\text{Pre}^{\exists} \text{inst-of}(F)$, assuming that F is the complement of U , is open in Z^* .

Consider now any open subset U of Y^* . U is a union of basic opens, hence a directed union $\bigcup_{i \in I} U_i$, where each U_i is a finite union (in particular, a monotone Boolean combination) of basic opens. We observe that $\text{Pre}^{\forall} \text{inst-of}$ commutes with directed unions. This is because each word w (say of length m) in Z^* only has finitely many images w_1, w_2, \dots, w_{2^m} , namely the 2^m words obtained from w by replacing each letter by \square , or not: if $w \in \text{Pre}^{\forall} \text{inst-of}(\bigcup_{i \in I} U_i)$, then for every j , $1 \leq j \leq 2^m$, there is an $i \in I$ such that $w_j \in U_i$; we may take the same i for every j , by directedness, whence $w \in \text{Pre}^{\forall} \text{inst-of}(U_i)$; the converse direction is obvious. So $\text{Pre}^{\forall} \text{inst-of}(U)$ is the directed union $\bigcup_{i \in I} \text{Pre}^{\forall} \text{inst-of}(U_i)$, and is therefore open: *inst-of* is upper semi-continuous. \square

Lemma E.8 Let X, Z be topological spaces. The relation $\text{id}_X \times \text{inst-of}$ that relates $(f, \vec{t}) \in X \times Z^*$ with $(f, \vec{u}) \in X \times (Z + \{\square\})^*$ if and only if \vec{t} *inst-of* \vec{u} is continuous.

Proof. Let $Y = Z + \{\square\}$, and fix an arbitrary open subset \mathcal{V} of $X \times (Z + \{\square\})^*$. Write \mathcal{V} as a union of open rectangles $\bigcup_{i \in I} U_i \times W_i$ (where every U_i and every W_i is open; this is the definition of the product topology).

Lower semi-continuity. $\text{Pre}^{\exists}(\text{id}_X \times \text{inst-of})(\mathcal{V})$ is equal to $\bigcup_{i \in I} U_i \times \text{Pre}^{\exists} \text{inst-of}(W_i)$, hence is open.

Upper semi-continuity. We must show that $\mathcal{U} = \text{Pre}^\forall(\text{id}_X \times \text{inst-of})(\mathcal{V})$ is open. For that, fix (f, \vec{t}) : it is enough to find an open rectangle containing (f, \vec{t}) and included in \mathcal{U} . As in Lemma E.7, note that there are only finitely many elements \vec{u} such that \vec{t} inst-of \vec{u} . List them as $\vec{u}_1, \dots, \vec{u}_m$. For each one, (f, \vec{u}_j) is in some $U_i \times W_i$: pick one such i and call it i_j . Our desired open rectangle is $U \times V$ where $U = \bigcap_{j=1}^m U_{i_j}$ and $V = \text{Pre}^\forall \text{inst-of}(\bigcup_{j=1}^m W_{i_j})$. By construction, (f, \vec{t}) is in $U \times V$. For every element (g, \vec{s}) of $U \times V$, by definition every element (g, \vec{u}) that is related to (g, \vec{s}) by $\text{id}_X \times \text{inst-of}$ is such that $g \in U$ and $\vec{u} \in \bigcup_{j=1}^m W_{i_j}$. Let j be such $\vec{u} \in W_{i_j}$. Since $g \in U$, g is in U_{i_j} , so (g, \vec{u}) is in $U_{i_j} \times W_{i_j} \subseteq \mathcal{U}$. \square

Lemma E.9 *Let $\mathfrak{C}^*.S$ be a tree iterator such that $\text{args } \mathfrak{C}$ is closed. Any subterm s of a term t in $\mathfrak{C}^*.S$ is again in $\mathfrak{C}^*.S$.*

Proof. This is proved by structural induction on t . If t is in S , then any subterm s of t is such that $s \preceq t$, hence $s \in S$. If $t \in \text{args } \mathfrak{C}$, we argue similarly, since \mathfrak{C} is closed (this is the first place where we need this assumption). Otherwise, either $s = t$ and the claim is obvious, or s is a proper subterm of t . In the latter case, there is an elementary context $c = f(u_1 u_2 \dots u_m) \in \mathfrak{C}$ such that t is obtained from c by replacing those u_j , $1 \leq j \leq m$, that equal \square by terms from $\mathfrak{C}^*.S$. I.e., t can be written $f(t_1, t_2, \dots, t_m)$, where $t_j = u_j$ if $u_j \neq \square$, $t_j \in \mathfrak{C}^*.S$ otherwise. For some j , s is a subterm of t_j . If $u_j = \square$, then $t_j \in \mathfrak{C}^*.S$, so that $s \in \mathfrak{C}^*.S$ by induction hypothesis. Otherwise, we claim that t_j is in $\text{args } \mathfrak{C}$. Indeed, $c = f(u_1 u_2 \dots u_m) \in \mathfrak{C}$, so the smaller $f(u_j)$ is in \mathfrak{C} , too, since \mathfrak{C} is closed in $X \times (\mathcal{T}(X) + \{\square\})^*$, hence downwards-closed. Since $u_j = t_j$, $f(t_j)$ is in \mathfrak{C} , so $t_j \in \text{args } \mathfrak{C}$. It follows that $t_j \in \mathfrak{C}^*.S$. Since s is a subterm of t_j , by induction hypothesis $s \in \mathfrak{C}^*.S$. \square

Lemma 10.9 (recap). Let X be a topological space, and \square be a hole outside $\mathcal{T}(X)$. Every tree iterator $\mathfrak{C}^*.S$ such that $\text{args } \mathfrak{C}$ is closed in $\mathcal{T}(X)$ denotes a closed subset of $\mathcal{T}(X)$.

Proof. Let V be the complement of $S \cup \text{args } \mathfrak{C}$ in $\mathcal{T}(X)$; this is open, since both S and $\text{args } \mathfrak{C}$ are closed. Let also \mathcal{V} be the open complement of $\mathfrak{C} \cap (X \times (\text{args } \mathfrak{C} + \{\square\})^*)$ in $X \times Y^*$, where we let Y abbreviate $\mathcal{T}(X) + \{\square\}$. Using Lemma E.8, the binary relation $\text{id}_X \times \text{inst-of}$, between $X \times \mathcal{T}(X)^*$ and $X \times Y^*$ is continuous, hence upper semi-continuous. Therefore $\mathcal{U} = \text{Pre}^\forall(\text{id}_X \times \text{inst-of})(\mathcal{V}) = \{(f, \vec{t}) \mid \forall \vec{u}. \vec{t} \text{ inst-of } \vec{u} \Rightarrow (f, \vec{u}) \in \mathcal{V}\}$ is open.

Write \mathcal{U} as a union of open rectangles $\bigcup_{i \in I} U_i \times W_i$. We claim that the complement of $\mathfrak{C}^*.S$ is $\bigcup_{i \in I} \diamond U_i \cdot W_i // V$. It will follow that $\mathfrak{C}^*.S$ is closed.

Let t be any term not in $\mathfrak{C}^*.S$. Consider a minimal subterm $f(\vec{t})$ of t that is not in $\mathfrak{C}^*.S$. By minimal, we mean that all its proper subterms are in $\mathfrak{C}^*.S$. Since $f(\vec{t})$ is not in $\mathfrak{C}^*.S$, it is in particular not in $S \cup \text{args } \mathfrak{C}$, hence it is in V . For any tuple $\vec{u} \in Y^*$ of which \vec{t} is an instance, i.e., such that \vec{t} inst-of \vec{u} , and such that the components of \vec{u} that are different from \square are in $\text{args } \mathfrak{C}$ (i.e., $\vec{u} \in (\text{args } \mathfrak{C} + \{\square\})^*$), (f, \vec{u}) cannot be in \mathfrak{C} : otherwise $f(\vec{t})$ would be obtained from $f(\vec{u})$ by replacing each occurrence of \square by some components of the tuple \vec{t} , which are all in $\mathfrak{C}^*.S$; so $f(\vec{t})$ would again be in $\mathfrak{C}^*.S$, which is impossible. Another way of stating this is that whenever (f, \vec{t}) ($\text{id}_X \times \text{inst-of}$) (f, \vec{u}) , then either \vec{u} is not in $(\text{args } \mathfrak{C} + \{\square\})^*$ or (f, \vec{u}) is not in \mathfrak{C} . That is, (f, \vec{t}) is in

\mathcal{U} . It follows that, for some $i \in I$, $f \in U_i$ and $\vec{t} \in W_i$. Recall that $f(\vec{t}) \in V$. So t is in $\diamond U_i \cdot W_i // V$.

Conversely, assume that $t \in \diamond U_i \cdot W_i // V$, for some $i \in I$. That is, t has a subterm $f(\vec{t})$ in V , with $f \in U_i$ and $\vec{t} \in W_i$. Assume, for the sake of contradiction, that t is in $\mathcal{E}^*.S$. By Lemma E.9, $f(\vec{t})$ is also in $\mathcal{E}^*.S$. Since $f(\vec{t}) \in V$, $f(\vec{t})$ is neither in S nor in $\text{args } \mathcal{E}$, so there is an elementary context $f(\vec{u})$ in \mathcal{E} such that \vec{t} is obtained from \vec{u} by replacing the \square elements in \vec{u} by some terms in $\mathcal{E}^*.S$. Since $f(\vec{u})$ is in \mathcal{E} , the components of \vec{u} that are different from \square are in $\text{args } \mathcal{E}$, by definition of the argument support. So \vec{u} is in $(\text{args } \mathcal{E} + \{\square\})^*$, hence $(f, \vec{u}) \in \mathcal{E} \cap (X \times (\text{args } \mathcal{E} + \{\square\})^*)$, the complement of \mathcal{V} . However, $(f, \vec{t}) \in (\text{id}_X \times \text{inst-of})(f, \vec{u})$, so (f, \vec{t}) cannot be in $\mathcal{U} = \text{Pre}^\forall(\text{id}_X \times \text{inst-of})(\mathcal{V})$. This contradicts the fact that (f, \vec{t}) is in $U_i \times W_i$.

This concludes our proof that the complement of $\mathcal{E}^*.S$ is $\bigcup_{i \in I'} \diamond U_i \cdot W_i // V$, so that $\mathcal{E}^*.S$ is closed. \square

Lemma 10.11 (recap). Let X be a topological space, and \square be a hole outside $\mathcal{T}(X)$, \mathcal{E} be a closed subset of $X \times (\mathcal{T}(X) + \{\square\})^*$, S be a closed subset of $\mathcal{T}(X)$, and assume that $\text{args } \mathcal{E}$ is closed. Then the tree iterator $\mathcal{E}^*.S$ is irreducible in the following cases:

- 1) if \mathcal{E} is non- \square -linear, and S is non-empty;
- 2) or if \mathcal{E} is \square -generated and \square -linear and S is irreducible;
- 3) or if \mathcal{E} is non-empty, \square -generated, and S is empty.

Proof. (1) \mathcal{E} is non- \square -linear, and S is non-empty. Since $S \subseteq \mathcal{E}^*.S$, $\mathcal{E}^*.S$ is non-empty. Since \mathcal{E} is non- \square -linear, in particular there is an elementary context $f(\vec{u}) \in \mathcal{E}$ such that \vec{u} has at least two occurrences of \square . More precisely, there is an element of the form $f(\vec{u}_1 \square \vec{u}_2 \square \vec{u}_3)$ in \mathcal{E} , i.e., one where \square occurs at least twice. If $\mathcal{E}^*.S$ is included in the union of two closed subsets S' and S'' , but not in S' and not in S'' , then pick t' in $\mathcal{E}^*.S$ outside S' , and t'' in $\mathcal{E}^*.S$ outside S'' . Pick some term t in $\mathcal{E}^*.S$ (for example, t' or t''), and let $\vec{t}_1, \vec{t}_2, \vec{t}_3$ be obtained from $\vec{u}_1, \vec{u}_2, \vec{u}_3$ respectively by replacing all occurrences of \square by t . Clearly $f(\vec{t}_1 t' \vec{t}_2 t'' \vec{t}_3)$ is in $\mathcal{E}^*.S$, hence in S' or in S'' . Assume without loss of generality that it is in S' . Then its subterm t' is in S' , contradiction. So $\mathcal{E}^*.S$ is irreducible.

(2) \mathcal{E} is \square -generated and \square -linear, and S is irreducible. Assume that $\mathcal{E}^*.S$ is included in the union of two closed subsets S' and S'' , but not in S' or in S'' . We claim that there is a context $c' \in \mathcal{E}^{\square*}$, with exactly one occurrence of \square , and a term $t' \in S \cup \text{args } \mathcal{E}$, such that $c'[t']$ is not in S' . Indeed, since \mathcal{E} is \square -linear, there is a context $c = c_1[c_2[\dots[c_k]\dots]]$, $k \in \mathbb{N}$, where each c_i is in \mathcal{E} , and such that one obtains a term outside S' by replacing the unique occurrence of \square (if any) in c by a term from $S \cup \text{args } \mathcal{E}$. If \square actually occurs (once) in c , let $c' = c$, and the term outside S' obtained above can be written $c'[t']$ for some $t' \in S \cup \text{args } \mathcal{E}$. Otherwise, some c_i does not contain an occurrence of \square . Pick i minimal: so \square occurs (once) in c_1, c_2, \dots, c_{i-1} , but not in c_i ; moreover, $c = c_1[c_2[\dots[c_{i-1}[c_i]\dots]]]$ is a term (i.e., where \square does not occur) outside S' . Write c_i as $f(\vec{u})$. Since \mathcal{E} is \square -generated, one can split \vec{u} as $\vec{u}_1 \vec{u}_2$ so that $f(\vec{u}_1 \square \vec{u}_2) \in \mathcal{E}$. Pick any term t' from S : this is easy since irreducible sets are non-empty. Let $c' = c_1[c_2[\dots[c_{i-1}[f(\vec{u}_1 \square \vec{u}_2)]]\dots]]$. Then

$c_i = f(\vec{u}_1 \vec{u}_2) \preceq_{\leq} f(\vec{u}_1 \square \vec{u}_2)[t'] = f(\vec{u}_1 t' \vec{u}_2)$, so $c \preceq_{\leq} c'[t']$. Since c is not in S' , $c'[t']$ is not in S' either.

In any case, there is a context $c' \in \mathfrak{C}^{\square*}$, with exactly one occurrence of \square , and a term $t' \in S \cup \text{args } \mathfrak{C}$, such that $c'[t']$ is not in S' . Similarly, there is a context $c'' \in \mathfrak{C}^{\square*}$, with exactly one occurrence of \square , and a term $t'' \in S \cup \text{args } \mathfrak{C}$, such that $c''[t'']$ is not in S'' . Note that both $c'[t']$ and $c''[t'']$ are in $\mathfrak{C}^*.S$.

Examine the case where t' or t'' is in $\text{args } \mathfrak{C}$, say t' by symmetry. So $f(t') \in \mathfrak{C}$ and \square does not occur in t' , for some $f \in X$. Since \mathfrak{C} is \square -generated, $f(t' \square)$ or $f(\square t')$ is in \mathfrak{C} , too, say $f(t' \square)$. The term $c'[f(t' \square)[c''[t'']]]$ is then in $\mathfrak{C}^*.S$, hence in S' or in S'' , say S' . However, the terms $c'[t']$ and $c''[t'']$ are below the latter term in the \preceq_{\leq} ordering, since \square occurs in c' . So they are both in S' , since S' is closed hence downwards-closed. But precisely, $c'[t']$ is not in S' , contradiction.

Then examine the case where t' and t'' are both in S . We recall from Lemma E.6 that $\textcircled{\ast}$ is continuous, and from Lemma B.1 that $i: Y \rightarrow Y^*$ and $\text{cat}: Y^* \times Y^* \rightarrow Y^*$ are continuous. By a simple induction on c' , the function that maps each term t to $c'[t]$ is then continuous. Similarly, the function that maps t to $c''[t]$ is continuous. Consider the map f that sends each term $t \in \mathcal{T}(X)$ to $c'[c''[t]] = c'[c'']$: f is continuous. Since $c'[c'']$ is in $\mathfrak{C}^{\square*}$, every $t \in S$ is such that $c'[c''[t]]$ is in $\mathfrak{C}^*.S$, hence in $S' \cup S''$. So S is included in $f^{-1}(S' \cup S'') = f^{-1}(S') \cup f^{-1}(S'')$. The latter is a union of two closed sets, since f is continuous. Since S is irreducible, S is included in $f^{-1}(S')$ or in $f^{-1}(S'')$. If $S \subseteq f^{-1}(S')$, then in particular $t' \in f^{-1}(S')$, that is, $c'[c''[t']] \in S'$. However, $c'[t'] \preceq_{\leq} c'[c''[t']]$, since \square occurs in c'' . So $c'[t']$ is in S' , a contradiction. Similarly, $S \subseteq f^{-1}(S'')$ also leads to a contradiction.

So $\mathfrak{C}^*.S$ is in fact included in S' or in S'' . We conclude that $\mathfrak{C}^*.S$ is irreducible.

(3) \mathfrak{C} is non-empty, \square -generated, and S is empty. Since \mathfrak{C} is non-empty, it contains an elementary context $f(\vec{u})$. Since \mathfrak{C} is closed, hence downwards-closed in $\leq \times \preceq_{\leq}$, $f()$ is also in \mathfrak{C} , so f is in $\mathfrak{C}^*.S$. Let S' be the closure of f in $\mathcal{T}(X)$. Since $\mathfrak{C}^*.S$ is closed, it contains S' . So $\mathfrak{C}^*.S \supseteq \mathfrak{C}^*.S'$. The converse inclusion follows since $S = \emptyset$, whence $\mathfrak{C}^*.S = \mathfrak{C}^*.S'$. By construction, S' is irreducible closed. If \mathfrak{C} is \square -linear, then $\mathfrak{C}^*.S = \mathfrak{C}^*.S'$ is irreducible by case (2). If \mathfrak{C} is not \square -linear, then $\mathfrak{C}^*.S = \mathfrak{C}^*.S'$ is irreducible by case (1). \square

Lemma 10.12 (recap). Let X be a topological space. The complement $\mathfrak{C}\pi$ of the open subset denoted by the simple tree expression $\pi = \diamond U(\pi_1 \mid \pi_2 \mid \cdots \mid \pi_n)$ is given by structural induction on π by:

- $\mathfrak{C}\pi = ((F \times \{\square\}^*) \cup (X \times (\mathfrak{C}\pi_1^* \{\square\}^? \mathfrak{C}\pi_2^* \{\square\}^? \cdots \{\square\}^? \mathfrak{C}\pi_n^*)))^*. \emptyset$ if $n \geq 1$, where F is the complement of U in X ;
- if $n = 0$, then $\mathfrak{C}\pi = (F \times \{\square\}^*)^*. \emptyset$.

Proof. We first deal with the case $n = 0$. The terms t that are not in $\diamond U()$ are those such that no function symbol occurring in t is in U . So they are the terms whose function symbols are all in F , i.e., the terms in $(F \times \{\square\}^*)^*. \emptyset$.

Next, we deal with the case $n \geq 1$. Let $\pi = \diamond U(\pi_1 \mid \pi_2 \mid \cdots \mid \pi_n)$. Let us explain the notation first. Notice that $\{\square\}$ is irreducible closed in $\mathcal{T}(X) + \{\square\}$. So $\{\square\}^*$ and $\mathfrak{C}\pi_1^* \{\square\}^? \mathfrak{C}\pi_2^* \{\square\}^? \cdots \{\square\}^? \mathfrak{C}\pi_n^*$ are word-products on $\mathcal{T}(X) + \{\square\}$, hence are closed in $(\mathcal{T}(X) + \{\square\})^*$, by Corollary 6.6. Write $\mathfrak{C} = (F \times \{\square\}^*) \cup$

$(X \times (\mathbb{C}\pi_1^*\{\square\}^?\mathbb{C}\pi_2^*\{\square\}^? \dots \{\square\}^?\mathbb{C}\pi_n^*))$. We must show that $\mathbb{C}\pi = \mathcal{E}^*.\emptyset$. Notice that $\text{args } \mathcal{E} = \mathbb{C}\pi_1 \cup \mathbb{C}\pi_2 \cup \dots \cup \mathbb{C}\pi_n$, so $\text{args } \mathcal{E}$ is closed.

For every term $t \in \mathcal{T}(X)$, we show, in one direction, that if t is not in π , then t is in $\mathcal{E}^*.\emptyset$, by structural induction on t . Write t as $f(\vec{t})$, where $\vec{t} = t_1 t_2 \dots t_m$. Necessarily, t_1, t_2, \dots, t_m are outside π as well. So t_1, t_2, \dots, t_m are in $\mathcal{E}^*.\emptyset$. Moreover, $f \notin U$ or $\vec{t} \notin Y^*\pi_1 Y^* \dots Y^*\pi_n Y^*$, where $Y = \mathcal{T}(X)$. If $f \notin U$, then $f \in F$, so $t = f(\vec{t})$ is obtained from the context $f(\square^m)$, in $F \times \{\square\}^*$, by replacing the holes by terms from $\mathcal{E}^*.\emptyset$. Therefore t is itself in $\mathcal{E}^*.\emptyset$. Otherwise, \vec{t} is in $\mathbb{C}\pi_1^* Y^? \mathbb{C}\pi_2^* Y^? \dots Y^? \mathbb{C}\pi_n^*$, by Lemma 6.2. (Recall that $n \geq 1$.) So one can write \vec{t} as a sequence $\vec{t}_1 \in \mathbb{C}\pi_1^*$, followed by zero or one term s_1 , followed by a sequence $\vec{t}_2 \in \mathbb{C}\pi_2^*$, followed by zero or one term s_2, \dots , followed by zero or one term s_{n-1} , followed by a sequence $\vec{t}_n \in \mathbb{C}\pi_n^*$. When there is indeed a term s_i between \vec{t}_i and \vec{t}_{i+1} , say that s_i exists. Note that those terms among s_1, s_2, \dots, s_{n-1} that do exist are in $\mathcal{E}^*.\emptyset$, since they form a subsequence of \vec{t} (use the induction hypothesis). Let \vec{u} be the sequence obtained by concatenating \vec{t}_1, \square if s_1 exists (and nothing otherwise), \vec{t}_2, \square if s_2 exists, $\vec{t}_3, \dots, \square$ if s_{n-1} exists, and \vec{t}_n . One obtains t by replacing the occurrences of \square in $f(\vec{u})$ by terms in $\mathcal{E}^*.\emptyset$, and $f(\vec{u})$ is in $X \times (\mathbb{C}\pi_1^*\{\square\}^?\mathbb{C}\pi_2^*\{\square\}^? \dots \{\square\}^?\mathbb{C}\pi_n^*)$ by construction, so $t \in \mathcal{E}^*.\emptyset$ again.

Conversely, we claim that no term in $\mathcal{E}^*.\emptyset$ can be in (the language of) π . We start by proving the following claim (a): for every $j, 1 \leq j \leq n$, no term in $\mathbb{C}\pi_j$ can be in π . Indeed, if $t \in \mathbb{C}\pi_j$ is in $\pi = \diamond U(\pi_1 \mid \pi_2 \mid \dots \mid \pi_n)$, then t has a subterm $s \in \pi_j$. Then $s \leq t$, to $t \in \pi_j$ since opens are upwards-closed: contradiction.

We then show that whenever $t \in \mathcal{E}^*.\emptyset$, then t cannot be in $\pi = \diamond U(\pi_1 \mid \pi_2 \mid \dots \mid \pi_n)$, by structural induction on t , following the definition of $\mathcal{E}^*.\emptyset$. Assume that $t \in \pi$: there is a subterm $s = g(\vec{s})$ of t such that $g \in U$ and \vec{s} is in $Y^*\pi_1 Y^* \dots Y^*\pi_n Y^*$. (Again, $Y = \mathcal{T}(X)$.) Note that s itself is in π . By Claim (a), t cannot be in $\mathbb{C}\pi_1 \cup \mathbb{C}\pi_2 \cup \dots \cup \mathbb{C}\pi_n = \text{args } \mathcal{E}$. It follows that $t = f(\vec{t})$ must be obtained from some elementary context $f(\vec{u})$ in \mathcal{E} by replacing the occurrences of the hole \square by terms, themselves from $\mathcal{E}^*.\emptyset$. Write \vec{t} as $t_1 t_2 \dots t_m$, \vec{u} as $u_1 u_2 \dots u_m$. There are two cases, corresponding to the definition of \mathcal{E} . If $(f, \vec{u}) \in F \times \{\square\}^*$, then $f \notin U$, so $s = g(\vec{s})$ must be different from t (since $g \in U$), hence s must be a subterm of some $t_j, 1 \leq j \leq m$. Moreover, u_j is a hole, so t_j is in $\mathcal{E}^*.\emptyset$. By induction hypothesis t_j cannot be in π , hence its subterm $s \leq t_j$ is not in π either. This is impossible since s is in π . The other case is when (f, \vec{u}) is in $X \times (\mathbb{C}\pi_1^*\{\square\}^?\mathbb{C}\pi_2^*\{\square\}^? \dots \{\square\}^?\mathbb{C}\pi_n^*)$. Then \vec{t} is in $\mathbb{C}\pi_1^* Y^? \mathbb{C}\pi_2^* Y^? \dots Y^? \mathbb{C}\pi_n^*$, i.e., not in $\mathcal{T}(X)^*\pi_1 \mathcal{T}(X)^* \dots \mathcal{T}(X)^*\pi_n \mathcal{T}(X)^*$, by Lemma 6.2 (recall that $n \geq 1$). So again s must be different from t , hence be a subterm of some $t_j, 1 \leq j \leq m$. Either t_j is in some $\mathbb{C}\pi_i, 1 \leq i \leq n$ (when $u_j \neq \square$), or $u_j = \square$ and t_j is in $\mathcal{E}^*.\emptyset$. However, $\mathbb{C}\pi_i$ is included in $\text{args } \mathcal{E}$, hence in $\mathcal{E}^*.\emptyset$. So in any case t_j is in $\mathcal{E}^*.\emptyset$. Since t_j contains $s = g(\vec{s})$ as a subterm, $s \leq t_j$ and therefore t_j is also in $\pi = \diamond U(\pi_1 \mid \pi_2 \mid \dots \mid \pi_n)$. This is impossible by induction hypothesis. Having reached a contradiction in each case, we conclude. \square

E.3 Checking Inclusion between Tree Steps

Lemma 10.14 (recap). Let X be a topological space, C and C' be two irreducible closed subsets of X , \vec{P} and \vec{P}' be two word-products over $\mathcal{T}(X)$. Then $C^{\vec{P}}(\vec{P}) \subseteq$

$C'^{\vec{\tau}}(\vec{P}')$ iff $C \subseteq C'$ and $\vec{P} \subseteq \vec{P}'$, or $C^{\vec{\tau}}(\vec{P}) \subseteq \text{supp } \vec{P}'$.

Proof. The if direction is obvious, noting that $\text{supp } \vec{P}' \subseteq C'^{\vec{\tau}}(\vec{P}')$. Conversely, assume $C^{\vec{\tau}}(\vec{P}) \subseteq C'^{\vec{\tau}}(\vec{P}')$.

For every pair $(f, \vec{t}) \in C \times \vec{P}$, since $f(\vec{t}) \in C^{\vec{\tau}}(\vec{P}) \subseteq C'^{\vec{\tau}}(\vec{P}')$, either $(f, \vec{t}) \in C' \times \vec{P}'$, or (f, \vec{t}) is in $S = \{(f, \vec{t}) \in X \times \mathcal{T}(X)^* \mid f(\vec{t}) \in \text{supp } \vec{P}'\}$. So $C \times \vec{P}$ is included in $(C' \times \vec{P}') \cup S$. Since @ is continuous (Lemma E.6) and $\text{supp } \vec{P}'$ is closed (Lemma 10.4), S is closed. By Lemma 6.7, \vec{P} is irreducible, so $C \times \vec{P}$ is irreducible. Also, $C' \times \vec{P}'$ is closed, since \vec{P}' is closed by Corollary 6.6. So $C \times \vec{P}$ is included in $C' \times \vec{P}'$ or in S .

If $C \times \vec{P} \subseteq S$, then $C^{\vec{\tau}}(\vec{P}) \subseteq \text{supp } \vec{P}'$. Indeed, all the terms $f(\vec{t})$ with $f \in C$ and $\vec{t} \in \vec{P}$ are in $\text{supp } \vec{P}'$, by the definition of S . And for every $t \in \text{supp } \vec{P}$, fix an arbitrary $f \in C$ (since C , being irreducible, is non-empty) to obtain that $f(t) \in C^{\vec{\tau}}(\vec{P})$ hence $f(t) \in \text{supp } \vec{P}'$; since $\text{supp } \vec{P}'$ is closed, hence downwards-closed in \leq , $t \in \text{supp } \vec{P}'$.

If $C \times \vec{P} \subseteq C' \times \vec{P}'$ on the other hand, then clearly $C \subseteq C'$ and $\vec{P} \subseteq \vec{P}'$, since neither C nor \vec{P} is empty. \square

E.4 Checking Inclusion between Tree Steps and Tree Iterators

Lemma E.10 *Let Z be a topological space, \square a hole outside Z , F a closed subset of Z . Let inst-of_F be the relation from Z^* to $(Z + \{\square\})^*$ defined by $w \text{ inst-of}_F w'$ iff w is obtained from w' by replacing each occurrence of \square by (possibly distinct) elements from F .*

Then inst-of_F is upper semi-continuous.

Proof. For short, let Y be $Z + \{\square\}$. For every word-product $P = e_1 e_2 \cdots e_n$ on Y , $\text{Pre}^{\exists} \text{inst-of}_F(P)$ is equal to $\text{Pre}^{\exists} \text{inst-of}_F(e_1) \text{Pre}^{\exists} \text{inst-of}_F(e_2) \cdots \text{Pre}^{\exists} \text{inst-of}_F(e_n)$, and for each atomic expression e_j , $\text{Pre}^{\exists} \text{inst-of}_F(e_j)$ is computed as follows: $\text{Pre}^{\exists} \text{inst-of}_F(F'^{?})$ is $((F' \setminus \{\square\}) \cup F)^?$ if $\square \in F'$ and $F'^{?}$ otherwise, $\text{Pre}^{\exists} \text{inst-of}_F(F'^{*})$ is $((F' \setminus \{\square\}) \cup F)^*$ if $\square \in F'$ and F'^{*} otherwise. (Notice that $F' \setminus \{\square\} = F' \cap Z$ is closed in Z .) The rest of the proof is as in Lemma E.7. \square

Lemma E.11 *Let X, Z be topological spaces, F be a closed subset of Z . The relation $\text{id}_X \times \text{inst-of}_F$ that relates $(f, \vec{t}) \in X \times Z^*$ with $(f, \vec{u}) \in X \times (Z + \{\square\})^*$ if and only if $\vec{t} \text{ inst-of}_F \vec{u}$ is upper semi-continuous.*

Proof. As for Lemma E.8, using Lemma E.10 instead of Lemma E.7. \square

Lemma E.12 *X be a topological space, \square a hole outside $\mathcal{T}(X)$, \mathfrak{C} be a closed subset of $X \times (\mathcal{T}(X) + \{\square\})^*$, and S be a closed subset of $\mathcal{T}(X)$. The set $\mathfrak{C}[S]$, defined as the set of all pairs (f, \vec{t}) where \vec{t} is obtained from \vec{u} by replacing each occurrence of \square by possibly different terms from S , for some \vec{u} such that $(f, \vec{u}) \in \mathfrak{C}$, is closed in $X \times \mathcal{T}(X)^*$.*

Proof. $\mathfrak{C}[S]$ is just $\text{Pre}^{\exists}(\text{id}_X \times \text{inst-of}_S)(\mathfrak{C})$; then use Lemma E.11. \square

Lemma 10.15 (recap). Let X be a topological space, C be an irreducible closed subset of X , \vec{P} be a word-product over $\mathcal{T}(X)$, \mathfrak{C} be a closed subset of $X \times (\mathcal{T}(X) + \{\square\})^*$ such that $\text{args } \mathfrak{C}$ is closed in $\mathcal{T}(X)$, where \square is a hole outside $\mathcal{T}(X)$, and S be a closed subset of $\mathcal{T}(X)$.

Then $C^{\vec{P}}(\vec{P}) \subseteq \mathfrak{C}^*.S$ iff $C \times \vec{P} \subseteq \mathfrak{C}[\mathfrak{C}^*.S]$ and $\text{supp } \vec{P} \subseteq \mathfrak{C}^*.S$, or $C^{\vec{P}}(\vec{P}) \subseteq \text{args } \mathfrak{C} \cup S$.

Proof. If $C \times \vec{P} \subseteq \mathfrak{C}[\mathfrak{C}^*.S]$ and $\text{supp } \vec{P} \subseteq \mathfrak{C}^*.S$, then we claim that every term $t = f(\vec{t})$ in $C^{\vec{P}}(\vec{P})$ is in $\mathfrak{C}^*.S$. Indeed, either $f \in C$ and $\vec{t} \in \vec{P}$, or $t \in \text{supp } \vec{P}$. In the first case, (f, \vec{t}) is obtained from some $(f, \vec{u}) \in \mathfrak{C}$ by replacing each occurrence of \square in \vec{u} by terms from $\mathfrak{C}^*.S$, so $f(\vec{t})$ is again in $\mathfrak{C}^*.S$. In the second case, where $t \in \text{supp } \vec{P}$, then $t \in \mathfrak{C}^*.S$ by assumption.

If $C^{\vec{P}}(\vec{P}) \subseteq \text{args } \mathfrak{C} \cup S$, then $C^{\vec{P}}(\vec{P})$ is trivially included in $\mathfrak{C}^*.S$.

Conversely, assume $C^{\vec{P}}(\vec{P}) \subseteq \mathfrak{C}^*.S$. For every $f \in C$ and $\vec{t} \in \vec{P}$, $f(\vec{t})$ is in $C^{\vec{P}}(\vec{P})$, hence in $\mathfrak{C}^*.S$. So either $f(\vec{t}) \in \text{args } \mathfrak{C} \cup S$, or there is an elementary context $f(\vec{u})$ in \mathfrak{C} such that \vec{t} is obtained from \vec{u} by replacing the occurrences of \square by terms from $\mathfrak{C}^*.S$. That is, $f(\vec{t})$ is in $\text{args } \mathfrak{C} \cup S$ or in $\mathfrak{C}[\mathfrak{C}^*.S]$. So $C \times \vec{P}$ is contained in the union of the set $@^{-1}(\text{args } \mathfrak{C} \cup S)$, which is closed since $@$ is continuous, and the set $\mathfrak{C}[\mathfrak{C}^*.S]$, which is closed by Lemma E.12 and Lemma 10.9. On the other hand, by Lemma 6.7, \vec{P} is irreducible, so $C \times \vec{P}$ is irreducible. So $C \times \vec{P}$ is included in $@^{-1}(\text{args } \mathfrak{C} \cup S)$ or in $\mathfrak{C}[\mathfrak{C}^*.S]$. If $C \times \vec{P} \subseteq @^{-1}(\text{args } \mathfrak{C} \cup S)$, then every term $f(\vec{t})$ with $f \in C$ and $\vec{t} \in \vec{P}$ is in $\text{args } \mathfrak{C} \cup S$; since $\text{args } \mathfrak{C} \cup S$ is closed hence downwards-closed, $\text{supp } \vec{P}$ is also included in $\text{args } \mathfrak{C} \cup S$, and therefore $C^{\vec{P}}(\vec{P})$ as well. Otherwise, $C \times \vec{P} \subseteq \mathfrak{C}[\mathfrak{C}^*.S]$. Moreover, $\text{supp } \vec{P} \subseteq C^{\vec{P}}(\vec{P}) \subseteq \mathfrak{C}^*.S$. \square

E.5 Checking Inclusion between Tree Iterators

Lemma 10.21 (recap). Let X be a topological space, \mathfrak{C} and \mathfrak{C}' be two closed subsets of $X \times (\mathcal{T}(X) + \{\square\})^*$ such that $\text{args } \mathfrak{C}$ and $\text{args } \mathfrak{C}'$ are closed in $\mathcal{T}(X)$, where \square is a hole outside $\mathcal{T}(X)$, and let S, S' be two closed subsets of $\mathcal{T}(X)$.

Then $\mathfrak{C}^*.S \subseteq \mathfrak{C}'^*.S'$ iff $\mathfrak{C}[\mathfrak{C}^*.S] \subseteq @^{-1}(\text{args } \mathfrak{C}' \cup S') \cup \mathfrak{C}'[\mathcal{T}^\square(X)]$ and $\text{args } \mathfrak{C} \cup S \subseteq \mathfrak{C}'^*.S'$.

Proof. A warning, first. Although we have used the notation $f(\vec{u})$ for elementary contexts, we must recall that this is an abbreviation for a pair (f, \vec{u}) . One obtains the term (or context) $f(\vec{u})$ from (f, \vec{u}) by applying $@$, hence the use of $@$ in the statement of the lemma.

If $\mathfrak{C}^*.S \subseteq \mathfrak{C}'^*.S'$, then in particular $\text{args } \mathfrak{C} \cup S \subseteq \mathfrak{C}^*.S \subseteq \mathfrak{C}'^*.S'$. Moreover, for every $(f, \vec{t}) \in \mathfrak{C}[\mathfrak{C}^*.S]$, $f(\vec{t})$ is in $\mathfrak{C}^*.S$. If $f(\vec{t})$ is not in $\text{args } \mathfrak{C}' \cup S'$, then (f, \vec{t}) is obtained from some elementary context $f(\vec{u}) = (f, \vec{u})$ in \mathfrak{C}' by replacing all occurrences of the hole \square by terms (in $\mathfrak{C}^*.S$, but this is irrelevant). In any case $(f, \vec{t}) \in @^{-1}(\text{args } \mathfrak{C}' \cup S') \cup \mathfrak{C}'[\mathcal{T}(X)]$, hence in $@^{-1}(\text{args } \mathfrak{C}' \cup S') \cup \mathfrak{C}'[\mathcal{T}^\square(X)]$. It follows that $\mathfrak{C}[\mathfrak{C}^*.S] \subseteq @^{-1}(\text{args } \mathfrak{C}' \cup S') \cup \mathfrak{C}'[\mathcal{T}^\square(X)]$.

Conversely, assume that $\mathfrak{C}[\mathfrak{C}^*.S] \subseteq @^{-1}(\text{args } \mathfrak{C}' \cup S') \cup \mathfrak{C}'[\mathcal{T}^\square(X)]$ and $\text{args } \mathfrak{C} \cup S \subseteq \mathfrak{C}'^*.S'$. Consider any term $t = f(\vec{t})$ in $\mathfrak{C}^*.S$. We show by induction on the definition of $\mathfrak{C}^*.S$ that t is in $\mathfrak{C}'^*.S'$. If $t \in \text{args } \mathfrak{C} \cup S$ (base case), then $t \in \mathfrak{C}'^*.S'$

by assumption. Otherwise, (f, \vec{t}) is obtained from some $(f, \vec{u}) \in \mathcal{C}$ by replacing each occurrence of \square in \vec{u} by elements of $\mathcal{C}^{\bar{*}}.S$. Let us make this clear. Write \vec{t} as $t_1 t_2 \cdots t_n$, \vec{u} as $u_1 u_2 \cdots u_n$. For each j , $1 \leq j \leq n$, either $t_j = u_j$ or $u_j = \square$ and $t_j \in \mathcal{C}^{\bar{*}}.S$. When $u_j \neq \square$, observe that $t_j = u_j$ is in $\text{args } \mathcal{C} \subseteq \mathcal{C}^{\bar{*}}.S$. Therefore, in any case, $t_j \in \mathcal{C}^{\bar{*}}.S$ for every j , $1 \leq j \leq n$. By induction hypothesis, $t_j \in \mathcal{C}'^{\bar{*}}.S'$ for every j , $1 \leq j \leq n$. On the other hand, the existence of (f, \vec{u}) , as specified, means that (f, \vec{t}) is in $\mathcal{C}[\mathcal{C}^{\bar{*}}.S]$. By assumption, (f, \vec{t}) is then in $@^{-1}(\text{args } \mathcal{C}' \cup S')$, or in $\mathcal{C}'[\mathcal{T}^{\square}(X)]$. In the first case, $f(\vec{t}) \in \text{args } \mathcal{C}' \cup S' \subseteq \mathcal{C}'^{\bar{*}}.S'$. In the second case, there is a pair $(f, \vec{v}) \in \mathcal{C}'$ such that \vec{t} is obtained from $\vec{v} = v_1 v_2 \cdots v_n$ by replacing each v_j that equals \square by the element t_j from \vec{t} . We have noticed that all such elements were in $\mathcal{C}'^{\bar{*}}.S'$. So $f(\vec{t})$ is in $\mathcal{C}'^{\bar{*}}.S'$. \square

Lemma 10.22 (recap). Let X be a topological space, \mathcal{C} and \mathcal{C}' be two closed subsets of $X \times (\mathcal{T}(X) + \{\square\})^*$, where \square is a hole outside $\mathcal{T}(X)$, and let S, S' be two closed subsets of $\mathcal{T}(X)$. Assume also that \mathcal{C} is of the form $\bigcup_{i=1}^m C_i \times Q_i$, and that \mathcal{C}' is of the form $\bigcup_{j=1}^n C'_j \times Q'_j$, where each C_i and each C'_j is irreducible closed in X , and Q_i and Q'_j are word-products over $\mathcal{T}(X) + \{\square\}$ for each i , $1 \leq i \leq m$, and each j , $1 \leq j \leq n$. Assume finally that $\mathcal{C}^{\bar{*}}.S$ is irreducible, and that $\square \in Q_i$ for every i , $1 \leq i \leq m$.

Then $\mathcal{C}^{\bar{*}}.S \subseteq \mathcal{C}'^{\bar{*}}.S'$ iff:

- either $\mathcal{C}^{\bar{*}}.S \subseteq \text{args } \mathcal{C}' \cup S'$,
- or $\text{args } \mathcal{C} \cup S \subseteq \mathcal{C}'^{\bar{*}}.S'$, and for every i , $1 \leq i \leq m$, there is a j , $1 \leq j \leq n$, such that $C_i \subseteq C'_j$ and $Q_i[\mathcal{T}^{\square}(X)] \subseteq Q'_j[\mathcal{T}^{\square}(X)]$.

Proof. Consider the following statements:

- (i) $\mathcal{C}^{\bar{*}}.S \subseteq \mathcal{C}'^{\bar{*}}.S'$;
- (ii) $\text{args } \mathcal{C} \cup S \subseteq \mathcal{C}'^{\bar{*}}.S'$, and for every i , $1 \leq i \leq m$, either $C_i^{\bar{*}}(Q_i[\mathcal{C}^{\bar{*}}.S]) \subseteq \text{args } \mathcal{C}' \cup S'$ or for some j , $1 \leq j \leq n$, $C_i \subseteq C'_j$ and $Q_i[\mathcal{C}^{\bar{*}}.S] \subseteq Q'_j[\mathcal{T}^{\square}(X)]$.
- (iii) either $\mathcal{C}^{\bar{*}}.S \subseteq \text{args } \mathcal{C}' \cup S'$, or $\text{args } \mathcal{C} \cup S \subseteq \mathcal{C}'^{\bar{*}}.S'$ and for every i , $1 \leq i \leq m$, there is a j , $1 \leq j \leq n$, such that $C_i \subseteq C'_j$ and $Q_i[\mathcal{T}^{\square}(X)] \subseteq Q'_j[\mathcal{T}^{\square}(X)]$.

The Lemma claims that (i) is equivalent to (iii). We shall show this by proving that (i) implies (ii) implies (iii) implies (i).

The differences between (ii) and (iii) are: first, there is an additional disjunct $\mathcal{C}^{\bar{*}}.S \subseteq \text{args } \mathcal{C}' \cup S'$ in (iii); second, (iii) dispenses with the disjunct $C_i^{\bar{*}}(Q_i[\mathcal{C}^{\bar{*}}.S]) \subseteq \text{args } \mathcal{C}' \cup S'$ that occurs in (ii); finally, we use $Q_i[\mathcal{C}^{\bar{*}}.S]$ versus $Q_i[\mathcal{T}^{\square}(X)]$ in the last inclusion.

Before we start, note that, using Lemma 10.8, $\text{args } \mathcal{C} = \bigcup_{i=1}^m \text{supp } Q_i \cap \mathcal{T}(X)$ is closed, and similarly, $\text{args } \mathcal{C}' = \bigcup_{j=1}^n \text{supp } Q'_j \cap \mathcal{T}(X)$ is closed: so Lemma 10.21 applies.

(i) \Rightarrow (ii). By Lemma 10.21, $\text{args } \mathcal{C} \cup S \subseteq \mathcal{C}'^{\bar{*}}.S'$ and $\mathcal{C}[\mathcal{C}^{\bar{*}}.S] = \bigcup_{i=1}^m C_i \times Q_i[\mathcal{C}^{\bar{*}}.S] \subseteq @^{-1}(\text{args } \mathcal{C}' \cup S') \cup \mathcal{C}'[\mathcal{T}^{\square}(X)] = @^{-1}(\text{args } \mathcal{C}' \cup S') \cup \bigcup_{j=1}^n (C'_j \times Q'_j[\mathcal{T}^{\square}(X)])$. Therefore, for every i , $1 \leq i \leq m$, $C_i \times Q_i[\mathcal{C}^{\bar{*}}.S] \subseteq @^{-1}(\text{args } \mathcal{C}' \cup S') \cup$

$\bigcup_{j=1}^n (C'_j \times Q'_j[\mathcal{T}^\square(X)])$. Since $\mathfrak{E}^\square.S$ is irreducible, and because $Q_i[\mathfrak{E}^\square.S]$ is obtained by syntactically replacing occurrences of \square by $\mathfrak{E}^\square.S$ (Lemma 10.18), $Q_i[\mathfrak{E}^\square.S]$ is a word-product. So $Q_i[\mathfrak{E}^\square.S]$ is irreducible by Lemma 6.7. It follows that $C_i \times Q_i[\mathfrak{E}^\square.S]$ is irreducible. So $C_i \times Q_i[\mathfrak{E}^\square.S] \subseteq @^{-1}(\text{args } \mathfrak{E}' \cup S')$ or, for some $1 \leq j \leq n$, $C_i \times Q_i[\mathfrak{E}^\square.S] \subseteq C'_j \times Q'_j[\mathcal{T}^\square(X)]$. In the first case, $C_i^{\overline{?}}(Q_i[\mathfrak{E}^\square.S]) \subseteq \text{args } \mathfrak{E}' \cup S'$, and we conclude.

(iii) \Rightarrow (i). If $\mathfrak{E}^\square.S \subseteq \text{args } \mathfrak{E}' \cup S'$, then (i) holds trivially. Otherwise, since $Q_i[\mathfrak{E}^\square.S] \subseteq Q_i[\mathcal{T}^\square(X)]$, we obtain that for every i , there is a j such that $C_i \times Q_i[\mathfrak{E}^\square.S] \subseteq C'_j \times Q'_j[\mathcal{T}^\square(X)]$. Therefore $\mathfrak{E}[\mathfrak{E}^\square.S] = \bigcup_{i=1}^m C_i \times Q_i[\mathfrak{E}^\square.S] \subseteq \bigcup_{j=1}^n (C'_j \times Q'_j[\mathcal{T}^\square(X)]) = \mathfrak{E}'[\mathcal{T}^\square(X)]$. Also, $\text{args } \mathfrak{E} \cup S \subseteq \mathfrak{E}^\square.S'$ by assumption, so by Lemma 10.21 $\mathfrak{E}^\square.S \subseteq \mathfrak{E}^\square.S'$.

(ii) \Rightarrow (iii). If $\mathfrak{E}^\square.S \subseteq \text{args } \mathfrak{E}' \cup S'$, then (iii) is clear. So let us also assume $\mathfrak{E}^\square.S \not\subseteq \text{args } \mathfrak{E}' \cup S'$.

To show (iii) under these assumptions, and in the view of what (ii) states, it is enough to show that $C_i^{\overline{?}}(Q_i[\mathfrak{E}^\square.S]) \subseteq \text{args } \mathfrak{E}' \cup S'$ is impossible, and that for every i and j , if $Q_i[\mathfrak{E}^\square.S] \subseteq Q'_j[\mathcal{T}^\square(X)]$, then the stronger inclusion $Q_i[\mathcal{T}^\square(X)] \subseteq Q'_j[\mathcal{T}^\square(X)]$ holds.

We start with the former. Since $\square \in Q_i$, $\mathfrak{E}^\square.S$ is included in $C_i^{\overline{?}}(Q_i[\mathfrak{E}^\square.S])$: indeed, for every $t \in \mathfrak{E}^\square.S$, $f(t) \in C_i^{\overline{?}}(Q_i[\mathfrak{E}^\square.S])$ for some (arbitrary) $f \in C_i$, so $t \preceq f(t)$ is also in $C_i^{\overline{?}}(Q_i[\mathfrak{E}^\square.S])$, which is closed hence downwards-closed. Therefore $C_i^{\overline{?}}(Q_i[\mathfrak{E}^\square.S])$ cannot be included in $\text{args } \mathfrak{E}' \cup S'$, since we assumed that $\mathfrak{E}^\square.S \not\subseteq \text{args } \mathfrak{E}' \cup S'$.

We proceed with the other claim. Write Q_i as the product of atomic expressions $e_1 e_2 \cdots e_m$ over $\mathcal{T}(X) + \{\square\}$, and similarly Q'_j as $e'_1 e'_2 \cdots e'_n$. Let e_i be written as F_i^* or $F_i^?$, $1 \leq i \leq m$, where F_i is closed in $\mathcal{T}(X) + \{\square\}$, and also irreducible in case e_i is written $F_i^?$. Similarly, write e'_j as $F_j'^*$ or $F_j'^?$.

The core of the argument is that: (*) when $\square \in e_i$, i.e., when e_i is of the form $\{\square\}^?$ or F_i^* with $\square \in F_i$, then $e_i[\mathfrak{E}^\square.S]$ is not included in any $e'_j[\mathcal{T}^\square(X)]$, $1 \leq j \leq n$, unless $\square \in e'_j$ as well. Indeed, assume that \square is not in e'_j . So $e'_j[\mathcal{T}^\square(X)] = e'_j$ is included in $\text{args } \mathfrak{E}'$. If $e_i[\mathfrak{E}^\square.S]$ were included in $e'_j[\mathcal{T}^\square(X)]$, then $\mathfrak{E}^\square.S$, which is included in $e_i[\mathfrak{E}^\square.S]$ since $\square \in e_i$, would be included in e'_j , hence in $\text{args } \mathfrak{E}'$. This would contradict the fact that $\mathfrak{E}^\square.S \not\subseteq \text{args } \mathfrak{E}' \cup S'$.

It follows that: (**) $e_i[\mathfrak{E}^\square.S] \subseteq e'_j[\mathcal{T}^\square(X)]$ iff $e_i[\mathcal{T}^\square(X)] \subseteq e'_j[\mathcal{T}^\square(X)]$. The if direction is obvious. In the only if direction, we distinguish four cases. If $\square \notin e_i$, then $e_i[\mathfrak{E}^\square.S] = e_i = e_i[\mathcal{T}^\square(X)]$, and the claim is clear. If $e'_j = \{\square\}^?$, then the assumption that $e_i[\mathfrak{E}^\square.S]$ is included in $e'_j[\mathcal{T}^\square(X)]$ means that $e_i[\mathfrak{E}^\square.S]$ is a collection of sequences of terms of length at most 1; this is then certainly also the case for e_i , hence of $e_i[\mathcal{T}^\square(X)]$, so $e_i[\mathcal{T}^\square(X)] \subseteq e'_j[\mathcal{T}^\square(X)]$. If $e'_j = F_j'^*$ where $\square \in F_j'$, then $e'_j[\mathcal{T}^\square(X)]$ is just $\mathcal{T}^\square(X)^*$ (see Lemma 10.18), and the claim is obvious. Otherwise, \square is in e_i , and not in e'_j , so the claim follows from (*).

The algorithmic characterization of inclusion of word-products given in Lemma 6.10 now allows us to conclude that $Q_i[\mathfrak{E}^\square.S] \subseteq Q'_j[\mathcal{T}^\square(X)]$ if and only if $Q_i[\mathcal{T}^\square(X)] \subseteq Q'_j[\mathcal{T}^\square(X)]$. Concretely, this algorithmic characterization only depends on the answers

to queries of the form $e_i[\mathfrak{C}^*.S] \subseteq e'_j[\mathcal{T}^\square(X)]$ in the first case, and on answers to the corresponding queries $e_i[\mathcal{T}^\square(X)] \subseteq e'_j[\mathcal{T}^\square(X)]$ in the second case. By (**) these answers must be the same. \square

E.6 Intersections of Tree-Products

Lemma 10.23 (recap). Let X be a topological space. The intersection of two tree steps $P = C^{\vec{?}}(\vec{P})$ and $P' = C'^{\vec{?}}(\vec{P}')$ is equal to $\bigcup_{j=1}^n (C \cap C')^{\vec{?}}(\vec{P}''_j) \cup (\text{supp } \vec{P} \cap P') \cup (P \cap \text{supp } \vec{P}')$, where $\vec{P} \cap \vec{P}'$ is expressed as a finite union $\bigcup_{j=1}^n \vec{P}''_j$ of word-products on $\mathcal{T}(X)$. If $C \cap C'$ can be written as the union of finitely many irreducible closed subsets C_i , $1 \leq i \leq m$, then $P \cap P'$ is also equal to the union of the tree steps $C_i^{\vec{?}}(\vec{P}''_j)$ ($1 \leq i \leq n$, $1 \leq j \leq m$), of $\text{supp } \vec{P} \cap P'$, and of $P \cap \text{supp } \vec{P}'$.

Proof. Let $t = f(\vec{t})$ be any term in $P \cap P'$. Since $t \in P$, t is in $\text{supp } \vec{P}$, or $f \in C$ and $\vec{t} \in \vec{P}$. In the first case, t is in $\text{supp } \vec{P} \cap P'$. Similarly, the claim that t' is in P' splits in two cases. The first one gives $t' \in P \cap \text{supp } \vec{P}'$. There remains the case where $f \in C$, $\vec{t} \in \vec{P}$, and $f \in C'$, $\vec{t} \in \vec{P}'$. Then f is in $C \cap C'$ (resp., in some C_i , if $C \cap C'$ can be written as a finite union of irreducible closed subsets C_i), and \vec{t} is in some \vec{P}''_j , so t is in $(C \cap C')^{\vec{?}}(\vec{P}''_j)$ (resp., in $C_i^{\vec{?}}(\vec{P}''_j)$).

Conversely, $\text{supp } \vec{P} \cap P'$ and $P \cap \text{supp } \vec{P}'$ are included in $P \cap P'$. It remains to show that $(C \cap C')^{\vec{?}}(\vec{P}''_j)$ (resp., $C_i^{\vec{?}}(\vec{P}''_j)$) is included in $P \cap P'$, namely in both P and P' . We only deal with the first case. For every term t in $(C \cap C')^{\vec{?}}(\vec{P}''_j)$ (resp., $C_i^{\vec{?}}(\vec{P}''_j)$), either t is in $\text{supp } \vec{P}''_j$ or $t = f(\vec{t})$ with $f \in C \cap C'$ (resp., $f \in C_i$) and $\vec{t} \in \vec{P}''_j$. In the first case, the one-element word t is in \vec{P}''_j , hence in \vec{P} , so t is in $\text{supp } \vec{P}$ and therefore in $P = C^{\vec{?}}(\vec{P})$. In the second case, f is in C , \vec{P} is in \vec{P} , so $t = f(\vec{t})$ is in $P = C^{\vec{?}}(\vec{P})$. \square

Let us write $\text{supp } Q[S]$ for $\text{supp}(Q[S])$.

Lemma E.13 Let X be a Noetherian space, Q be a word-product over $\mathcal{T}(X) + \{\square\}$, and S be a closed subset of $\mathcal{T}(X)$. Then $\text{supp } Q[S] \subseteq \text{supp } Q \cup S$. (See Lemma 10.18 for $Q[S]$.)

Proof. We first claim that for every atomic expression e , $\text{supp } e[S] \subseteq \text{supp } e \cup S$. If $e = \{\square\}^?$, then $\text{supp } e[S] = \text{supp } S^? = S$. If $e = I^?$ where I is irreducible closed in $\mathcal{T}(X)$, then $\text{supp } e[S] = \text{supp } I^? = I = \text{supp } e$. If $e = \mathcal{F}^*$ and $\square \in \mathcal{F}$, then $\text{supp } e[S] = \text{supp } ((\mathcal{F} \setminus \{\square\}) \cup S)^* = (\mathcal{F} \setminus \{\square\}) \cup S \subseteq \mathcal{F} \cup S = \text{supp } e \cup S$. If $e = \mathcal{F}^*$ and $\square \notin \mathcal{F}$, then $\text{supp } e[S] = \text{supp } \mathcal{F}^* = \mathcal{F} = \text{supp } e$.

Write Q as $e_1 e_2 \cdots e_n$, where each e_i is an atomic expression. Then $\text{supp } Q[S] = \bigcup_{i=1}^n \text{supp } e_i[S] \subseteq \bigcup_{i=1}^n (\text{supp } e_i \cup S) \subseteq \bigcup_{i=1}^n \text{supp } e_i \cup S = \text{supp } Q \cup S$. \square

Lemma 10.24 (recap). Let X be a Noetherian space, and S be a closed subset of $\mathcal{T}(X)$. Let $C^{\vec{?}}(\vec{P})$ be a tree step, \mathfrak{C} be a closed subset of $X \times (\mathcal{T}(X) + \{\square\})^*$ of the form $\bigcup_{j=1}^n C_j \times Q_j$, where each C_j is irreducible closed in X and each Q_j is a word-product over $\mathcal{T}(X) + \{\square\}$.

The intersection of the tree step $P = C^{\vec{\tau}}(\vec{P})$ and of the tree iterator $P' = \mathfrak{C}^*.S$ is the union of $\text{supp } \vec{P} \cap P'$, of $P \cap (S \cup \text{args } \mathfrak{C})$, and of $(C \cap C_j)^{\vec{\tau}}(\vec{P} \cap Q_j[P'])$, $1 \leq j \leq n$.

If, for each j , one can write $C \cap C_j$ as the union of finitely many irreducible subsets C_{ij} , $1 \leq i \leq m_j$, and if $\vec{P} \cap Q_j[P']$ can be expressed as the union of finitely many word-products $\vec{P}_{\ell j}$, $1 \leq \ell \leq q_j$, then $P \cap P'$ is also equal to the union of $\text{supp } \vec{P} \cap P'$, of $P \cap (S \cup \text{args } \mathfrak{C})$, and of $C_{ij}^{\vec{\tau}}(\vec{P}_{\ell j})$, $1 \leq j \leq n$, $1 \leq i \leq m_j$, $1 \leq \ell \leq q_j$.

Proof. Let $t = f(\vec{t})$ be any term in $P \cap P'$. Since $t \in P$, t is in $\text{supp } \vec{P}$, or $f \in C$ and $\vec{t} \in \vec{P}$. In the first case, t is in $\text{supp } \vec{P} \cap P'$. Similarly, since $t \in P'$, t is in $S \cup \text{args } \mathfrak{C}$, or there is an elementary context $c \in \mathfrak{C}$ such that t is in $c[P']$. In the first of these cases, t is in $P \cap (S \cup \text{args } \mathfrak{C})$. There remains the case where $f \in C$, $\vec{t} \in \vec{P}$, and t is in $c[P']$ for some elementary context c —necessarily of the form $f(\vec{u})$ —in \mathfrak{C} . In that case, \vec{t} is obtained from \vec{u} by replacing each occurrence of \square by possibly different terms from P' , in other words, (f, \vec{t}) is in $\mathfrak{C}[P']$. By Lemma 10.18, f is in C_j and \vec{t} is in $Q_j[P']$ for some j . It follows that f is in $C \cap C_j$ and \vec{t} is in $\vec{P} \cap Q_j[P']$, so $t = f(\vec{t})$ is in $(C \cap C_j)^{\vec{\tau}}(\vec{P} \cap Q_j[P'])$. (Additionally, if $C \cap C_j$ can be written as $\bigcup_{i=1}^{m_j} C_{ij}$ and $\vec{P} \cap Q_j[P']$ can be written as $\bigcup_{\ell=1}^{q_j} \vec{P}_{\ell j}$, then t is in $C_{ij}^{\vec{\tau}}(\vec{P}_{\ell j})$ for some j , i , and ℓ , too.)

For the converse inclusions, we check that:

- $\text{supp } \vec{P} \cap P' \subseteq P \cap P'$: every term in $\text{supp } \vec{P} \cap P'$ is in $\text{supp } \vec{P}$ hence in $P = C^{\vec{\tau}}(\vec{P})$, and is also in P' ;
- $P \cap (S \cup \text{args } \mathfrak{C}) \subseteq P \cap P'$: every term in $P \cap (S \cup \text{args } \mathfrak{C})$ is in $S \cup \text{args } \mathfrak{C}$, hence in $P' = \mathfrak{C}^*.S$, and also in P ;
- $(C \cap C_j)^{\vec{\tau}}(\vec{P} \cap Q_j[P'])$ (resp., $C_{ij}^{\vec{\tau}}(\vec{P}_{\ell j})$) is included in $P \cap P'$.

Every term t in $\text{supp}(\vec{P} \cap Q_j[P'])$ (resp., $\text{supp } \vec{P}_{\ell j}$) is such that the one-element word t is in $\vec{P} \cap Q_j[P']$ (resp., $\vec{P}_{\ell j}$, hence also in $\vec{P} \cap Q_j[P']$). Since that one-element word is in \vec{P} , t is in $\text{supp } \vec{P}$ hence in $P = C^{\vec{\tau}}(\vec{P})$, and since it is also in $Q_j[P']$, t is in $\text{supp } Q_j[P']$, hence in $\text{supp } Q_j \cup P'$ by Lemma E.13, hence in $\text{args } \mathfrak{C}' \cup P' \subseteq P'$.

Next, let t be any term of the form $f(\vec{t})$ with $f \in C \cap C_j$ and $\vec{t} \in \vec{P} \cap Q_j[P']$. (If instead $f \in C_{ij}$ and $\vec{t} \in \vec{P}_{\ell j}$, then f is in $C \cap C_j$ and \vec{t} is in $P \cap Q_j[P']$.) Since f is in C and \vec{t} is in \vec{P} , $t = f(\vec{t})$ is in $P = C^{\vec{\tau}}(\vec{P})$. Since f is in C_j and \vec{t} is in $Q_j[P']$, by Lemma 10.18 (f, \vec{t}) is in $\mathfrak{C}[P']$, so $t = f(\vec{t})$ is in $P' = \mathfrak{C}^*.S$. \square

Lemma 10.26 (recap). Let X be a Noetherian space, S and S' be closed subsets of $\mathcal{T}(X)$. Let also \mathfrak{C} (resp., \mathfrak{C}') be a closed subset of $X \times (\mathcal{T}(X) + \{\square\})^*$ of the form $\bigcup_{i=1}^m C_i \times Q_i$ (resp., $\bigcup_{j=1}^n C'_j \times Q'_j$), where each C_i and each C'_j is irreducible closed in X and each Q_i and each Q'_j is a normalized word-product over $\mathcal{T}(X) + \{\square\}$. For all i, j , write $C_i \cap C_j$ as $\bigcup_{k=1}^{p_{ij}} C''_{ijk}$ where each C''_{ijk} is irreducible closed in X , and let $Q''_{ij\ell}$, $1 \leq \ell \leq q_{ij}$ enumerate the elements of $\text{Meet}^{\mathcal{E}}(Q_i, Q'_j)$, where the oracle \mathcal{E} is defined in Lemma 10.25.

Then the intersection of the tree iterators $P = \mathfrak{C}^{\bar{*}}.S$ and $P' = \mathfrak{C}'^{\bar{*}}.S'$ is the tree iterator $\mathfrak{C}''^{\bar{*}}.S''$, where $\mathfrak{C}'' = \bigcup_{i,j,k,\ell} C''_{ijk} \times Q''_{ij\ell}$ and where S'' is the union of $P \cap (\text{args } \mathfrak{C}' \cup S')$ and of $(\text{args } \mathfrak{C} \cup S) \cap P'$.

Proof. Let $t = f(\vec{t})$ be in $P \cap P'$, where $\vec{t} = t_1 \cdots t_N$. We show that t is in $\mathfrak{C}''^{\bar{*}}.S''$ by induction on the size of t . If t is in $\text{args } \mathfrak{C} \cup S$, then t is in $(\text{args } \mathfrak{C} \cup S) \cap P'$, hence in S'' , hence in $\mathfrak{C}''^{\bar{*}}.S''$. Similarly if t is in $\text{args } \mathfrak{C}' \cup S'$. In the remaining case, there is an elementary context (f, \vec{u}) in \mathfrak{C} such that \vec{t} is obtained by replacing the occurrences of \square in \vec{u} by possibly different terms from P . Hence there is an index i , $1 \leq i \leq m$, such that f is in C_i and \vec{u} is in Q_i , so \vec{t} is in $Q_i[P]$. Similarly, there is an index j , $1 \leq j \leq n$, such that $f \in C'_j$ and \vec{t} is in $Q'_j[P']$. Since f is in $C_i \cap C'_j$, f is in C''_{ijk} for some k , $1 \leq k \leq p_{ij}$. By Lemma 10.25, \vec{t} is in some element of $\text{Meet}^{\mathcal{E}}(Q_i, Q'_j)[P \cap P']$, hence in $Q''_{ij\ell}[P \cap P']$ for some ℓ , $1 \leq \ell \leq q_{ij}$. Since every term t_k in the list \vec{t} is strictly smaller than t , every t_k that is in $P \cap P'$ is also in $\mathfrak{C}''^{\bar{*}}.S''$, by induction hypothesis. It follows that \vec{t} is in $Q''_{ij\ell}[\mathfrak{C}''^{\bar{*}}.S'']$. Therefore $(f, \vec{t}) \in C''_{ijk} \times Q''_{ij\ell}[\mathfrak{C}''^{\bar{*}}.S''] \subseteq \mathfrak{C}''[\mathfrak{C}''^{\bar{*}}.S''] \subseteq \mathfrak{C}''^{\bar{*}}.S''$.

This shows that $P \cap P' \subseteq \mathfrak{C}''^{\bar{*}}.S''$. Conversely, let $t = f(\vec{t})$ be any term in $\mathfrak{C}''^{\bar{*}}.S''$. We show that t is in $P \cap P'$ by induction on the size of t .

If t is in $P \cap (\text{args } \mathfrak{C}' \cup S')$, then t is in P , and in $\text{args } \mathfrak{C}' \cup S' \subseteq \mathfrak{C}'^{\bar{*}}.S' = P'$. If t is in $(\text{args } \mathfrak{C} \cup S) \cap P'$, then t is in $P \cap P'$ by a symmetric argument. This shows that if t is in S'' , then t is in $P \cap P'$.

If t is in $\text{args } \mathfrak{C}''$, then by Lemma 10.8, $\text{args } \mathfrak{C}'' = \bigcup_{i,j,k,\ell} \text{supp } Q''_{ij\ell} \cap \mathcal{T}(X)$, so t is in $\text{supp } Q''_{ij\ell}$ for some i, j and ℓ . By the last part of Lemma 10.25, t is in $\text{supp } Q_i \cap \text{supp } Q'_j$, or in $\text{supp } Q_i \cap P'$, or in $P \cap \text{supp } Q'_j$ (or in $\{\square\}$, but that is impossible since $t \in \mathcal{T}(X)$). Hence, using Lemma 10.8 again, t is in $\text{args } \mathfrak{C} \cap \text{args } \mathfrak{C}'$, or in $\text{args } \mathfrak{C} \cap P'$, or in $P \cap \text{args } \mathfrak{C}'$. In any case, t is in $P \cap P'$.

We have proved that if t is in $\text{args } \mathfrak{C}'' \cup S''$, then t is in $P \cap P'$. The other possibility for t to be in $\mathfrak{C}''^{\bar{*}}.S''$ is for f to be in C''_{ijk} and \vec{t} to be obtained from some elementary context \vec{u} in $Q''_{ij\ell}$ (for some i, j, k, ℓ) by replacing the occurrences of \square by possibly different terms from $\mathfrak{C}''^{\bar{*}}.S''$ —hence from $P \cap P'$, by induction hypothesis. Then f is in both C_i and C'_j , while \vec{t} is in $Q''_{ij\ell}[P \cap P']$, hence in $Q_i[P] \cap Q'_j[P']$ by Lemma 10.25. This shows that $t = f(\vec{t})$ is both in $\mathfrak{C}[P] = \mathfrak{C}[\mathfrak{C}^{\bar{*}}.S] \subseteq \mathfrak{C}^{\bar{*}}.S = P$ and in $\mathfrak{C}'[P'] \subseteq P'$. \square

Index

Symbols

η^{Γ}	embedding inside ideal completion 8	$\dagger E$	points with a point way-below it in E 7
η_X^S	embedding inside sobrification 12	A_{apo}	associative path ordering 46
γ	representation map of a (W)ADL	$[C_1 C_2 \dots C_n]$	irreducible closed subsets, prefix topology 39
ω	top code, heterogeneous words 41	$[F_1 F_2 \dots F_n]$	subbasic closed sets of the prefix topology 39
$\prod_{i \in I} X_i$	product of the spaces $X_i, i \in I$	$[\mathcal{U}]$	subbasic open of the prefix topology 38
Σ^*	words over Σ 6, 17	A^*	words with letters in A 28
\top	top element 13, 40	$A \mid u$	m-product notation 36
τ	set of codes representing the whole space 18	$\text{args } \mathcal{C}$	argument support of \mathcal{C} , in the ranked tree case 43, 52
(r)	ideal generated by r 24	$\text{args } \mathcal{C}$	syntactic argument support of \mathcal{C} , in the unranked tree case 62
\sqrt{I}	radical of ideal I 24	$\mathcal{C}^{\bar{x}}.S$	tree iterator 42, 60
$+$	sum of STREs 42, 60	$\mathcal{C}[\mathcal{T}^{\square}(X)]$	set of contexts obtained by replacing some holes in a context of \mathcal{C} by terms 56
\rightarrow_1	simplification rule for STREs 46	$\mathcal{C}[S]$	language substitution 55
\wedge	intersection operation on codes 18	$\mathcal{C}[S]$	context application 52
0	empty STRE 42, 60	$c[S]$	applying context c to set S of trees 43
\emptyset	empty multiset 34	$c^{\bar{r}}[S]$	optionally applying context c to set S of trees 43
\uplus	multiset union 34	$\mathcal{C}[t]$	context application 52
\square	hole 42, 52, 60	cat	concatenation map 76
$\square C$	subbasic closed set, powerset 22	$cl(A)$	closure of A 8
$\diamond U$	opens of $\mathcal{S}(X)$ 11	$cons$	adding a letter, prefix topology 83
$\diamond U$	subbasic open, Hoare powerdomain 12	$\mathbb{C}\pi$	complement of π 54
$\diamond U$	subbasic open, powerset 22	$\mathcal{C}^{\bar{r}}(\vec{P})$	tree step 60
$\diamond U(\pi_1 \mid \dots \mid \pi_n)$	simple tree expression 50	D^*	words over D 17, 33
$\diamond U \cdot \mathcal{U}$	specific open in $\mathcal{T}(X)$ 86	D^{\otimes}	finite multisets over D . . . 17, 36
$\diamond U \cdot \mathcal{U} // V$	specific open in $\mathcal{T}(X)$ 89	$F^{\bar{r}}(\vec{P})$	denotation of tree steps 51
$\llbracket u \rrbracket$	irreducible closed subset represented by code u 18	$f[E]$	image of E by f 10
$<$	strict part of \leq 6	F_I	closed subset in Zariski topology 24
\ll	way below relation 7	$F \mid C_1, \dots, C_n$	m-product 35
\triangleleft	quasi-ordering 6	$\bar{f}^{\bar{r}}(P_1, \dots, P_r)$	tree step 42
\triangleleft^*	specialization quasi-ordering . 10	$\mathcal{H}_V(X)$	Hoare powerdomain . . 12, 17, 21
\triangleleft^*	embedding quasi-ordering on words 28	$\mathcal{H}_V(X)_{\perp}$	lifted Hoare powerdomain . . 12, 17, 21
\triangleleft^{\otimes}	sub-multiset quasi-ordering . . 34	i	letter to word conversion 76
\approx	equivalence relation of \leq 6	$\mathbf{I}(X)$	ideal completion of X 8
\sqsupseteq	quasi-ordering on atomic expressions of word-products 31	i_n	conversion from tuples to words, prefix topology 83
\sqsupseteq^{\top}	quasi-ordering on Y^{\top} 40	inst-of	instance of relation 92
$\sqsupseteq^b, \sqsupseteq^{mul}, \sqsupseteq^b$	Hoare quasi-ordering 18, 21	inst-of_F	constrained instance of relation 97
\sqsupseteq^{mul}	multiset extension of \leq 34	$K[X_1, \dots, X_k]$	ring of polynomials with coefficients in K 24
\sqsupseteq	homeomorphic embedding . . 41, 42	$\text{Meet}^{\mathcal{E}}(P, P')$	intersection of word-products modulo oracle \mathcal{E} 31
\succ	opposite quasi-ordering 6	$\text{meet}[\wedge]$	lifting \wedge to words 79
$\Downarrow E$	points way-below some point in E 7		
$\downarrow E$	downward closure of E 6		
$\downarrow_X E$	downward closure of E in X . . 6		
$\uparrow E$	upward closure of E 6		
$\uparrow_X E$	upward closure of E in X . . . 6		

$\{x_1, \dots, x_n\}$	multiset containing x_1, \dots, x_n
\mathbb{N}_ω	completion of \mathbb{N} 4, 7, 8
$\mathbb{P}^*(X)$	non-empty powerset of X . . . 17,
	22, 23
$\mathbb{P}(X)$	powerset of X 7, 17, 22, 23
$\mathbb{P}_{\text{fin}}(X)$	finite powerset of X 7
$\text{Post}^*(E)$	iterated successors of E 3
$\text{Post}(E)$	one-step successors of E 3
$\text{Pre}^*(E)$	iterated predecessors of E 3
$\text{Pre}(E)$	one-step predecessors of E 2
$\text{Pre}^k(E)$	k -step predecessors of E 3
$\text{Pre}^{\leq k}(E)$	at-most- k -step predecessors of E
	3
$Q[S]$	result of replacing holes in Q by
	terms from S 57
$\mathbb{Q}[X_1, \dots, X_k]$	ring of polynomials with rational
	coefficients 25
$\mathcal{S}(X)$	sobrification of X 11, 17, 21
S°	translation of S-representations
	from ranked to unranked trees 50
$\text{Spec}(R)$	spectrum of ring R 17, 24, 25, 26
$\text{supp } \vec{P}$	support of \vec{P} 51
$\text{supp } \bar{P}$	syntactic support of \bar{P} 62
$\mathcal{T}^\square(X)$	terms plus an extra top element
	\square 56
$\mathcal{T}(D)$	finite trees over D 17, 50, 66
$\mathcal{T}(D, \{\square\})$	finite trees with one variable \square 50
$t[s]_p$	replacing subterm of t at position
	p by s 86
U/U	removing a U from the front of
	U 75
X^*	space of finite words over X . 27
X^\otimes	space of finite multisets over X
	34
X_a	X with the Alexandroff topology
	9
\widehat{X}	completion of X 3, 12, 17
X_σ	X with the Scott topology 9
X_u	X with the upper topology 10
$X \times Y$	product of X and Y 10
Y^\top	Y with new top \top 40
$\mathcal{Z}(u)$	common rational zeros of the poly-
	nomials of u 26

A

above	87
adequate domain of limits	4, 13, 14
ADL	4, 13, 14
Alexandroff topology	9
algebraic	7
almost open	10
antichain	6
argument support	43, 52
arity	42
ascending chain condition	11
atomic expression	29

B

backwards	3
base	9
basic opens	9
basis	3, 8
big end	39
\square -linear	46

C

closed set	8
closure	8
coarser	9
coarsest	9
codes	18
codomain	37, 81
cofinite topology	11
compact	10
completion	4
component	29
context	52
contexts	42
continuous	8, 9, 91
cover	3
coverability	3

D

dcpo	7
Dickson's Lemma	6
directed family	7
directed-complete partial order	7
discrete topology	9
domain	37, 81
domination quasi-ordering	18
downward closure	6
downwards-closed	6

E

effective WADL	14
elementary contexts	52
elementary open	28, 87
embedding quasi-ordering	50
empty	47

F

finer	9
finitary closed	10
finite	7
finite basis	6
free dcpo	8
free sober space	12
function symbol	42

G

\square -generated	47, 54
generated by	9
Gröbner base	25

H	
heterogeneous words	37, 38
Higman's Lemma	6
Hoare powerdomain	12
Hoare quasi-ordering	18
Hoffmann's Theorem	12
hole	42, 52
homeomorphism	9
I	
ideal	24
interior	8
interpolation	8
intersection map	18
irreducible	11
iterator	43
K	
Kaplansky's Theorem	24
Kruskal's Tree Theorem	6, 50
L	
Lasker-Noether Theorem	26
least upper bound	7
length	37
lifted Hoare powerdomain	12
limit points	14
limits	13
\square -linear	54
lower semi-continuous	91
lower Vietoris topology	12
M	
m-product	35
m-product notations	36, 81
m-SRE	36
matching	37, 81
monotone Boolean combination	92
monotonic	2, 6
N	
Noetherian	11, 24
normalized word-product	59
O	
open	8
open cover	10
order embeddings	6
order isomorphism	6
order quotient	6
ordering	6
P	
Parikh mapping	34
poset	6
position	86
prefix topology	38
primary ideal	26
prime ideal	24
product	10
product topology	10
Q	
quasi-order embedding	6
quasi-ordering	6
quotient map	10
quotient space	10
quotient topology	10
R	
Rabinowitch trick	25
radical	24
recognized at a state	44
representable	16
representable subset	14
representation map	14, 16
S	
S-representation	18
saturated	10
Scott closed	8
Scott open	8
Scott topology	8
Scott-continuous	7
signature	42
simple regular expressions	29
simple tree expression	50
simple tree regular expression	42
size	86
small end	39
sober	11
specialization quasi-ordering	10
spectrum	24
STRE	42
sub-multiset quasi-ordering	34
subbase	9
subbasic closed subset	9
subbasic open	9
subspace	10
subspace topology	10
subterm	50
support	51
syntactic argument support	62
syntactic support	62
syntactically \square -generated	61
syntactically \square -linear	61
T	
T_0	10
telescope	37
terms	42
topological embedding	10
topological WADL	16
topology	8

tree iterators	52
tree pre-products	42
tree step	51
tree steps	51
tree topology	50
tree-product	46
tree-products	42

U

upper bound	7
upper semi-continuous	91
upper topology	10
upward closure	6
upwards-closed	2, 6

W

WADL	13, 14
way below	7
weak adequate domain of limits	13, 14
well	6
well quasi-ordering	6
well-founded	6
well-ordered	7
wide telescope	38
wildcard	56
word language generators	37
word topology	27
word-product	29
word-product notations	79
word-SRE	29
wqo	6
WSTS	2

Z

Zariski topology	24, 27
------------------	--------