

# Possible polyphase periglaciation and glaciation adjacent to the Moreux impact-crater, Mars

R. J. Soare, Susan J. Conway, J.-P. Williams, A. J. Hepburn

## ► To cite this version:

R. J. Soare, Susan J. Conway, J.-P. Williams, A. J. Hepburn. Possible polyphase periglaciation and glaciation adjacent to the Moreux impact-crater, Mars. Icarus, 2021, 362, pp.114401. 10.1016/j.icarus.2021.114401. hal-03186645

# HAL Id: hal-03186645 https://hal.science/hal-03186645

Submitted on 31 Mar 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1	
2	
3	Possible polyphase periglaciation and glaciation
4	adjacent to the
5	Moreux impact-crater, Mars
6	
7	R.J. Soare, <sup>1</sup> S.J. Conway, <sup>2</sup> J-P Williams, <sup>3</sup> A.J. Hepburn <sup>4</sup>
8 9 10	<sup>1</sup> Geography Department, Dawson College, Montreal, Qc, Canada, H3Z 1A4 (rsoare@dawsoncollege.qc.ca)
11	<sup>2</sup> UMR, CNRS 6112, 2 rue de la Houssinière, 44322, Nantes, Cedex 3, France
12	<sup>3</sup> Earth, Planetary and Space Sciences, University of California, Los Angeles, CA, USA
13	<sup>4</sup> Department of Geography and Earth Sciences, Aberystwyth University, Aberystwyth, UK
14	
15	
16	
17	
18	
19	
20	
21	<b>Pages:</b> 50
22	Figures: 20
23	Keywords: Mars, atmosphere, climate, surface
24	
25	
26	
27	
28	

29 Abstract

The cyclicity and temporal succession of glacial-periglacial periods or epochs are keynotes of cold-climate geology on Earth. Relatively recent work within the Mars community has begun to dissect the mid- to higher-latitudinal terrain of Mars for analogical evidence of similar coldclimate cyclicity and succession.

Here, we carry on with this work by focusing on the terrain immediately to the north of the Moreux impact-crater (40-44<sup>0</sup> N, 43-47<sup>0</sup> E). The crater is located in northern Arabia Terra, to the south of Protonilus Mensae. It lies astride of and postdates Mars' crustal-dichotomy. The latter is a global geological-boundary that separates the ancient southern-highlands from the relatively younger northern-lowland plains.

Using cross-cutting relationships, relative stratigraphy and crater-size frequency 39 distributions (CSFDs) we identify three glacial and two periglacial periods that are temporally 40 intertwined and differentiated by a suite of features unique to each of these periods. For example, 41 we report and discuss clusters of pingo-like mounds amidst ridge and trough terrain or "brain 42 terrain". On Earth, the former are the work of freeze-thaw cycling; on Mars, the latter are thought 43 to be glacial remnants. In turn, the brain terrain is underlain by small-sized polygons possibly 44 45 formed by thermal contraction cracking and with margins underlain by degraded ice-wedges. Age estimates derived of CSFDs suggest that the polygonised terrain could as much as ~100 Ma, 46 whereas the brain terrain and pingo-like mounds are thought to be ~1 - ~10 Ma. Possible terminal-47 48 moraines that intercept brain-terrain fragments point to an even more recent period of glaciation

If the *CSFD* age-estimates are valid, then the polygons that underlie the brain terrain and incise the basin floors of our study zone could be an order of magnitude older than most of the age estimates associated with polygonised terrain at other locations on Mars. The fact that there are 52 two distinct periods of polygonization and periglacial activity with a wide offset of time within 53 one relatively small study zone also highlights the extent to which the freeze-thaw cycling of water 54 might be rooted as iteratively and as deeply in Mars' geological history as is its glaciation.

55 **1. Introduction** 

56 Our study zone is immediately to the north of the Moreux impact-crater (40-44<sup>0</sup> N, 43-47<sup>0</sup> 57 E) in northern Arabia Terra (**Fig. 1**). The latter lies astride of and postdates Mars' crustal-58 dichotomy, a global geological-boundary that separates the ancient southern highlands (McGill 59 and Dimitriou, 1990; Frey et al., 2002) from the relatively younger northern lowlands (Head et al., 50 2002; Tanaka et al, 2005).

Here, as elsewhere at similar and higher latitudes within both hemispheres, landscape assemblages, landforms and surface textures pointing to the work of: glaciation, i.e. the net and perennial accumulation (aggradation) or degradation of atmospherically-precipitated (water) ice; and, periglaciation, i.e. environments dominated by frost action and often, but not necessarily, the freeze-thaw cycling of water.

The long-term cyclicity and temporal succession of glacial-periglacial periods or epochs are keynotes of cold-climate geology on Earth. Relatively recent work within the Mars community has begun to dissect the mid- to higher-latitudinal terrain of Mars for analogical evidence of similar cold-climate cyclicity and succession (e.g. Dickson et al., 2008; Levy et al., 2009; Baker et al., 2015; Hubbard et al., 2011; Souness and Hubbard, 2013; Sinha and Murty, 2015; Hepburn et al., 2020).

Below, we carry on with and extend the range of this work to our study zone. In so doingwe have three principal aims:

- Integrate cross-cutting relationships and relative stratigraphy to posit a possible
   geochronological ordering of the disparate glacial and periglacial assemblages,
   landforms and surface textures.
- 77 2) Use crater-size frequency distribution to frame the proposed geochronological ordering
  78 in absolute terms.
- 3) Estimate the temporal distance between the latest and earliest onset of glaciation orperiglaciation.

### 81 **2.** Methods

Two *HiRISE* images (High Resolution Imaging Science Experiment, Mars Reconnaissance Orbiter, *MRO*, McEwen et al., 2007) (ESP\_045349\_2235, 43.405<sup>o</sup> N; 44.108<sup>o</sup> E; and, ESP\_042105\_2235, 43.163<sup>o</sup> N; 43.983<sup>o</sup> E) and one *CTX* image (Context Camera, Malin et al., 2007) (J01\_045349\_2239\_XN\_ 43N316W, 43.94<sup>o</sup> N; 43.99<sup>o</sup> E) frame our study region geographically and were used to evaluate cross-cutting relationships and relative stratigraphy therein.

Crater counts were conducted using the *Cratertools* plug-in for the *ERSI ArcGIS* to measure crater diameters (Kneissl et al., 2011). Crater size-frequency distributions (*CSFDs*) were compared with modeled crater-retention age isochrons from Hartmann (2005) using *Craterstats 2* (Michael and Neukum, 2010; Michael et al., 2016).

### 92 **3.** Observations

93 Small-sized (~5-15 m in diameter) and clastically non-sorted polygons (**Type-1**) are 94 ubiquitous in their coverage of the two basins in our study region (**Fig. 2a-c**). They also incise 95 some of the terrain near the eastern margin of the northernmost basin and are observed upslope as 96 well on its flanks (**Figs. 2a, c**). The polygons are shaped irregularly. Some of the polygons show 97 centres elevated slightly above the polygon margins, or high-centred polygons [*HCPs*]. Polygon
98 margins are metres to sub-metres in width and may or may not comprise sub-metre deep troughs

A slightly larger and morphologically-different type of polygon (**Type-2**) also is observed within the basins (**Figs. 3a-b**). The polygons tend to be four-sided, not irregular or asymmetrical, show margins that are etched slightly deeper and wider than the basin-surface polygons ,and are sub-divided by secondary cracks. Some of the polygons centres are knobby; many of them are relatively flat. Mirroring the spatially-constrained distribution of the basin-surface polygons, none of the etched polygons are observed beyond the basin borders.

105 The etched polygons occur amidst terrain superposed on the polygonised basin-surface. 106 The terrain, in turn, comprises outcrops, ridges and troughs, and small-sized mounds, described 107 below, each of which may or may not be contiguous. Outliers, distal from the principal areas of 108 distribution, are commonplace.

One of the basins in our study region exhibits wave-like and (mostly) equator-facing scarps 109 that span its multiple-kilometer reach (Fig. 4). The scarps display metres to decametres of 110 111 horizontal separation and, individually, metres of vertical drop. On a smaller scale the polygonised surface-basins also show uneven topography. The latter comprises metres-scale, serialized and 112 sub-parallel undulations (Figs. 5a-b). The undulations are shallow, perhaps comprising a few 113 metres of depth, exhibit kilometres<sup>2</sup> of reach and display a NE/SW vector; their minor axes are 114 scaled to the diameter of individual polygons. Elsewhere, circular to sub-circular and rimless 115 116 depressions are distributed in two basin-wide arcs (Fig. 5c). The depressions seemingly scale to the same shallow depth as the undulations noted above. 117

118 Coverage of the polygonised basins and overprinting by ridge and trough structures and 119 assemblages is widespread (**Figs. 6, 7a**). Individual ridges and troughs are metres in elevation and metres to decametres in width (**Fig. 7c**). Some of the ridge-trough assemblages are open; others are closed (**Fig. 7c**). Long axes often trend longitudinally (**Fig. 7c**). Numerous basins show a gradual decrease in the mass, height and surface coverage of the ridges and troughs as they edge further into the basin midsts and away from the margins (**Figs. 7a-b**), **also Fig. 6**). Spatial outliers also are observed throughout the basin surfaces (**Fig. 7b**).

As noted above, etched mounds populate the terrain superposed on the polygonised basinsurfaces. (Figs. 3a-b, 8a-d). Typically, long axes are sub-kilometre, elevation is scaled in metres, and shape is circular to sub-circular. Numerous mounds exhibit morphological asymmetry or irregularities at and within their margins, possibly by way or erosion or ablation (Figs. 8a-c). Many but not all of the mounds are clustered and contiguous with the larger-scale distributions of etched terrain on the polygonised basins (Fig. 8b). No mounds or mound remnants are observed on the elevated terrain that flank the basins.

The eastern flank (HiRISE image ESP\_045349\_2235) shows multiple closely-set and bow-132 shaped ridges (Fig. 9). The long-axial distribution of these ridges roughly follows the long axis of 133 134 the polygonised basin and exhibits kilometres of reach (Fig. 9). Most of the ridges comprise decametre to metre-scale boulders distributed continuously or discontinuously (Fig. 10a). At one 135 136 location, as many as nine serialised ridges are observed (Fig. 10a). Some of the ridges are subtle, being largely absent of boulders and observable only by means of a gentle decametres-wide and 137 metres-high change of (basin-surface) elevation (Fig. 10a). A much denser distribution of boulders 138 139 is observed at the topographical transition between the basin surface and the rockier flankingterrain immediately to the east (Figs. 10a-b). Where the ridges intercept the ridge-trough terrain 140 141 the amplitude and mass of the former increases perceptibly, compared to the ridge segments where 142 no interception occurs (Fig. 11).

Parallel albeit discontinuous lineations normal to these ridges intercept the latter and extend beyond them infrequently (**Figs. 10b-c**). The lineations show a strong variance in length, from multiple to hundreds of metres, and incise the same type of polygonised terrain that covers the basin surface elsewhere (**Fig. 10d**). Some of the lineations comprise metre-wide grooves with metre-to decametre-scale boulders nested therein; boulder size, most often, appears to vary randomly (**Fig. 10c**); others, comprise surface boulders whose size also seems to vary randomly. A few lineations exhibit smaller boulders on the lee side of larger ones (**Fig. 10c**).

Two other items are of note. First, much like the distribution of the serialized ridges, the density of boulder alignments and grooves increase with proximity to the basin's edge and the rockier terrain adjacent to the edge (**Figs. 10a-c**). Second, at some locations the grooved surfacelineations at/near basin edges extend through the positively sloped basin-adjacent terrain (**Fig. 9**, **top right-hand corner**).

### 155 4. Periglacial landscapes on Earth

### 156 *4.1 Thermal-contraction polygons*

157 Small-sized polygons are commonplace features of permafrost landscapes on Earth (e.g. Lachenbruch, 1962; Mackay, 1974; 1999; Czudek and Demek 1970; Washburn, 1973; Rampton 158 159 and Bouchard 1975; Rampton, 1988; French, 2007) (Figs. 12a-b). Typically ≤25 m in diameter, the polygons are produced by the tensile-induced fracturing of frozen sediment. This occurs when 160 the latter undergoes a sharp drop of sub-zero (Celsius) temperatures (de Leffingwell, 1915; 161 162 Lachenbruch, 1962). Fracturing, or *thermal-contraction* cracking, opens up shallow, narrow and vertical veins (Lachenbruch, 1962). In-filling prevents the ground from relaxing and returning to 163 its pre-cracked state as temperatures rise. Subsequent cycles of cracking and relaxation increase 164 165 the depth and the width of the veins, eventually forming wedges.

The fill type depends on the ambient boundary-conditions and the availability of: 1) 166 167 meltwater derived of thawed snow or ice; 2) winter hoarfrost; or, 3) windblown sand, mineral-soil, or a mixture of the two (e.g. de Leffingwell 1915; Péwé 1959; Lachenbruch 1962; Washburn 1973; 168 169 Sletten et al. 2003; Hallet et al. 2011) (Figs. 12b-d). The diurnal and seasonal iteration of cracking and filling grows the sub- to centimetre-scale veins into wedges with metre-scale depth and width 170 171 (Lachenbruch, 1962). As the cracks intercept one another they form individual polygons and, eventually, consolidated polygon networks (Figs. 12a-b). The latter may comprise hundreds of 172 square kilometres of continuous or discontinuous coverage and represent hundreds of cycles (e.g. 173 174 Black 1954; Lachenbruch 1962; Washburn 1973; Mackay 1974).

As ice- and/or sand-wedges aggrade by means of seasonal/annual supplements of fill, the 175 wedges and their sedimentary overburdens may rise above the elevation of the polygon centres; 176 177 this forms low-centred polygons [LCPs] (Péwé, 1959; Washburn 1973; Harris et al., 1988; Rampton, 1988; French, 2007). Ice or sand-wedge degradation, by thaw in the case of the former, 178 or aeolian erosion in the case of the latter, degrades and depletes wedge mass. If and as the wedges 179 180 fall below the elevation of the polygon centres the polygons begin to show high-centres [HCPs] (Péwé, 1959; Washburn 1973; Harris et al., 1988; Rampton, 1988; French 2007) (Fig. 12a). Some 181 182 ice/sand-wedge polygons show neither elevated nor collapsed margins. This is indicative of morphological nascency, evolved insufficiently to uplift marginal overburdens, or of a transitional 183 and topographically-neutral stage between aggradation and degradation. 184

Ice and sand-wedge polygons show a similar range of morphologies, depending on their stage of aggradation, degradation or neutrality. However, the respective presence or absence of sand-wedge polygons within a permafrost landscape points to climatic, hydrological and aeolian regimes that differ markedly from the regimes associated with ice-wedge polygons. Although the terrain fractured by sand-wedge polygons does require some ice-cementation (Lachenbruch, 1962), the boundary conditions associated with these polygons tend to be arid and extremely cold (Péwé, 1959; Lachenbruch, 1962; French and Guglielmin, 2000; Murton et al., 2000; Marchant et al., 2002; Sletten et al., 2003; Hallet et al. 2011; Wolfe et al., 2018). Permafrost landscapes where surface water is bountiful and freeze-thaw cycling is commonplace exhibit greater populations of ice-wedge than sand-wedge polygons (Péwé, 1959; Lachenbruch, 1962; French and Guglielmin, 2000; Murton et al., 2000; Sletten et al., 2003; Hallet et al. 2011; Wolfe et al., 2018).

The vulnerability of ice-wedge polygons to thaw means that the concurrent observation of low- and high-centred ice wedge polygons, or even a landscape dominated by high-centred polygons and thaw pits, would be expected when ice-rich landscapes undergo thermal stress and disequilibrium (Péwé, 1959; Washburn 1973; Harris et al., 1988; Rampton, 1988; French, 2007; Farquharson et al., 2019).

By comparison, sand-wedge polygons are insensitive to thermal stress. Aggradation and degradation are based uniquely on a gain of mass by aeolian deposition or mass loss by aeolian erosion. As aeolian processes would not be expected to vary substantially on a highly-localised scale, the closely set presence of low- and high-centred sand-wedge polygons in a permafrost landscape would be unusual (e.g. Péwé, 1959; French and Guglielmin, 2000; Marchant et al., 2002; Sletten et al., 2003; Hallett et al., 2011).

### 207 *4.2 Thermokarst and ice-rich permafrost*

*Thermokarst* refers to terrain in permafrost regions of excess ice which, depending upon
the currency of warming or cooling trends, could undergo frost heave or settlement as the excess
ice aggrades or degrades *in situ* (Taber, 1930; Penner, 1959; Hussey, 1966; Hughes, 1974;
Rampton, 1988) (Fig. 12d). *Excess ice* equals the volume of ice in the ground that exceeds the

total pore-volume that the ground would have under natural unfrozen conditions (Harris et al.,
1988; also, see Taber, 1930; Penner, 1959; Rampton and Mackay, 1971; Washburn, 1973;
Rampton, 1988; French, 2007).

Excess ice may comprise lenses, veins or larger masses of consolidated ice such as tabular 215 ice. All of these ice types originate and develop by the work of *ice segregation*. Ice segregation, 216 in turn, is the result of cryosuction pulling pore water to a freezing front where the ice consolidates 217 into lenses and, over time, into more substantial bodies of ice (e.g Taber, 1930; Black, 1954; 218 Penner, 1959; Rampton and Mackay, 1971; Rampton, 1988; French, 2007). As this occurs, the 219 220 overlying terrain undergoes frost heave proportional to the volumetric growth of ice-segregated lenses; frost settlement occurs when these lenses thaw or degrade and the newly voided or de-iced 221 soil settles under its own weight (e.g. Harris et al., 1988; Osterkamp et al., 2009; Farquharson et 222 223 al., 2019).

Fine to medium-grained soils such as silts or silty clays are particularly adept at hosting *segregation ice* because they have relatively-small interstices (e.g. Taber, 1930; Black, 1954; Penner, 1959; Rampton and Mackay, 1971; Rampton, 1988; French, 2007). Were the grain-size too fine, the migration of pore-water to the freezing front would be overly constricted and inadequate to the requirements of segregation-ice formation; opposingly, were the grain-size coarse, the migration of pore-water would be overly expeditious and flush out of the system before segregation ice could form.

The oscillation of regional mean-temperatures is one of the principal drivers of frost heave and settlement (e.g. Péwe, 1954; Czudek and Demek, 1970; Murton, 2001; Grosse et al., 2007; Osterkamp et al., 2009; Schirrmeister et al., 2013; Wetterich et al., 2014). This is exemplified, in part, by the hummocky sediments and rolling topography of the Tuktoyaktuk Coastlands of northern Canada (e.g. Rampton and Mackay, 1971; Rampton, 1988; Murton, 2001) and
northeastern Siberia (e.g. Czudek and Demek, 1970; Grosse et al., 2007; Schirrmeister et al.,
2013).

The time-frame of excess-ice aggradation and degradation, or of ice-induced heave and 238 settlement, however, need not be synchronous (e.g. Rampton and Mackay, 1971; Rampton, 1988; 239 Farquharson et al., 2019). For example, the wide-ranging presence of thermokarst lakes and alases 240 (thermokarst-lake basins absent of water) throughout the Tuktovaktuk Coastlands is rooted in the 241 relatively recent Holocene Era (e.g. Rampton and Mackay, 1971, Rampton 1988;). Contrarily, the 242 radiocarbon dating of wood ensconced in segregation-ice lenses and beds that are metres to tens 243 of metres deep point to region-wide ice-enrichment having taken place thousands and possibly 244 tens of thousands of years ago during the middle to late Wisconsinian glacial stade (Rampton and 245 Bouchard, 1988). This means that the geochronological distance or offset of time between ice 246 enrichment and depletion can be substantial. 247

248 *4.3 Perennially ice-cored mounds* 

Cold-climate and non-glacial landscapes on Earth show disparate mound types, i.e. earth
hummocks (e.g. Pettapiece, 1974; Kokelj et al., 2007), frost blisters (e.g. van Everdingen, 1982;
Pollard and French, 1985), and pingos, hydrostatic (e.g. Mackay, 1998) or hydraulic (e.g. Müller,
1963). Of these four types, only the *pingos* scale to the height and width of the mounds in our
study region and are perennial.

254 *4.3.1 Closed-system pingos* 

Hydrostatic, or closed-system, pingos [*CSPs*] are uniquely tied to ice-rich periglacial
landscapes (e.g. Müller, 1963; Washburn, 1973; Mackay, 1998; French, 2007). They originate and
develop in response to the loss of thermokarst lake-water by drainage or evaporation (Figs. 13a-

**d**). Lake-water loss triggers the propagation of a freezing front from the newly-exposed basin sides or floor. Pore water undergoes expulsion ahead of the freezing front into an increasingly compressed area of unfrozen ground, usually at or near the topographical low(s) of the basin. As the hydrostatic pressure of the expelled pore-water increases, the basin floor begins to deform and uplift. This creates a dome-like structure or mound. If and when the underlying water freezes, an ice core forms (Mackay, 1998, 1999) (**Fig. 13b**).

Mound shape, usually circular to sub-circular, is determined by the morphology or 264 bathymetry of the lake basins in which the mound is nested (e.g. Washburn, 1973; Mackay, 1998, 265 266 1999; French, 2007). Mound height (metres to decametres) and long-axis diameters (metres to hundreds of metres) are a function of four variables: 1) the surface area of the basin; 2) the volume 267 of pore water surrounding the basin that is exposed to permafrost aggradation; 3) the stage of 268 269 development, i.e. nascent, adolescent or growing, and mature or stagnant; and, 4) current (freezingfront) boundary conditions remaining relatively constant (Mackay 1998). Were thaw temperatures 270 to be lost from this cycling, arresting the loss of lake water and the further exposure of the 271 272 thermokarst lake-basin pore water, the shape, size and height of the pingo would be fixed. OSPs are less sensitive to this type of temperature-based fixation. Neither the migration of 273 274 topographically-driven or of deeply seated/geothermal water to the site of mound formation is contingent upon freeze-thaw boundary conditions at/or near the surface. 275

Metres-wide and deep fissures - dilation cracks - are not unusual amongst larger closedsystem pingos (**Fig. 13c**). These cracks propagate and trend from the mound summit as the pingo grows and tensile stresses increase (e.g. Washburn, 1973; Mackay, 1998, 1999; French, 2007). Mirroring the degradational pathway of the *OSPs* (see the discussion below), the summit cracks of *CSPs* may evolve into depressions as the thermal integrity of the overburden and the underlying ice-core dissipates (Fig. 13c). Subsequent stages of mound degradation possibly comprise fans,
slumps and collapse, leaving slightly elevated and irregular ramparts encircling the collapse basin
in the wake (Fig. 13d).

284 *4.3.2 Open-system pingos* 

Hydraulic or open-system pingos *[OSPs]* are circular to sub-circular in some instances (e.g. Bennike, 1983; Worsley and Gurney, 1996; Scholz and Baumann, 1997; Kelly, 2001) and linear, oblong or irregular in others (Cruickshank and Calhoun, 1965; O'Brien, 1971; Worsley and Gurney, 1996; Kelly, 2001). Mound height ranges from metres to decametres and long axes may reach hundreds of metres (Cruickshank and Colhoun, 1965; Allen, 1976; Bennike, 1983; Sholz and Baumann, 1997; Kelly, 2001) (**Figs. 14a-d**).

Fractures are observed at/or radiating from some mound summits or crests (Fig. 14b); they 291 form, as is the case with CSPs, in response to the tensional stresses within the mound overburden 292 as it undergoes uplift and sedimentary stretch (O'Brien, 1971; Allen, 1976; Worsley and Gurney, 293 1996; Scholz and Baumann, 1997). Further growth may translate these fractures into summit 294 295 depressions or craters, with the overburden becoming increasingly thin (O'Brien, 1971; Allen, 1976; Donner, 1978; Worsley and Gurney, 1996; Scholz and Baumann, 1997; Bennike, 1998; 296 Kelly, 2001) (Figs. 14a-b). Overburden thickness scales to the thermal integrity of the mound and 297 its ice core. Once this integrity is compromised by excessive stretching, ice core thaw could be 298 engendered. Mound collapse is the end-stage of pingo evolution (O'Brien, 1971; Allen, 1976; 299 Donner, 1978; Yoshikawa et al., 1994; Worsley and Gurney, 1996; Scholz and Baumann, 1997; 300 Kelly, 2001). 301

Three geological-pathways typically are associated with the origin and development of the
 *OSPs*, derived largely from field observations in Greenland:

Meltwater enters the local hydrological system at points of higher elevation, i.e.
 mountain slopes or valley walls, and migrates to lower elevations by means of sub-or
 intra-permafrost channels. The meltwater emerges where the permafrost is sufficiently
 thin or the hydraulic pressure is sufficiently high to uplift the near-surface/surface
 terrain into a mound (Müller, 1963; Bennike, 1998). An ice core forms as the water's
 near-surface exposure to perennial freezing evolves (Müller, 1963); or,

2) Meltwater forms basally under the snout of a surging or a receding glacier and migrates down-valley to the mound site(s) by means of near-surface faults, fractures or sub/intrapermafrost channels. Mound formation occurs where the permafrost is relatively thin and most susceptible to the hydraulic pressure generated by the valley-midst topography linking glaciers to mound sites (e.g. Müller, 1963; Yoshikawa et al., 1994; Yoshikawa and Harada, 1995); or,

# 316 3) Deeply-seated and possibly geothermally-driven water migrates to the surface through local or regional faults. The pressure is sufficient to uplift near-surface sediments, forming a mound and an underlying ice core as freezing iterates itself annually (e.g. O'Brien, 1971; Allen, 1976; Yoshikawa and Harada, 1995; Worsley and Gurney, 1996; Scholz and Baumann, 1997).

321 **5.** Glacial landscapes on Earth

### 322 *5.1 Glaciers*

*Glaciers* comprise perennial and net accumulations of water-ice precipitated episodically from the atmosphere and are classified according to their size and geographical context (e.g. Barry and Gann, 2011) (**Figs. 15a-c**). Valley glaciers, for example, are relatively small in scale compared to ice sheets and caps and their flow is constrained by the local topography (e.g. Sugden and John, 327 1976). Flow movement is the result of gravity and the internal deformation of ice (e.g. Barry and328 Gann, 2011).

329 *5.2 Moraines* 

*Glacial moraines* are composite landforms, formed and modified by a range of processes operative at glacial margins; these include glacio-tectonism, bulldozing/pushing, squeezing, freeze-on, melt-out, glaciofluvial, and gravity-driven processes (e.g. Benn and Evans, 2010) (**Figs. 15a, c**). Irrespective of their genesis, moraines may comprise unsorted glacial-till whose grain-size ranges from silts to boulder-sized clasts (**Fig. 15b**) and whose uneven distribution may engender topographical irregularity (e.g. Shakesby, 1989) (**Figs. 15 a, c**). The till is formed by the erosion and entrainment of sediment by glacial processes (van der Meer et al., 2003).

Moraines are referenced according to context and composition (e.g. Hambrey, 1994; Easterbrook, 1999) (**Figs. 15a, c**). Terminal or ice-marginal moraines are the outermost ridges which mark the maximum horizontal extent of a glacier (e.g. Hambrey, 1994; Summerfield, 1991). Depending on the orientation of moraines to ice flow, terminal and recessional moraines may be further divided into lateral or frontal moraines, or latero-frontal moraines where the entire tongue is demarcated by ridges (Benn and Evans, 2010). Serialised recessional-moraines are not unusual when multiple retreats and pauses of glaciers occur (e.g. Easterbrook, 1999) (**Fig. 15a, c**).

344 *5.3 Lineations (small-large scales)* 

Centimetre-scale *striations*, sub-metre-to metres-wide grooves and polished bedrock are the product of glacial abrasion at varying scales of width and depth (e.g. Hoppe and Schytt, 1953; Witkind, 1978; Lliboutry, 1994; Fjellander et al., 2006) (**Fig. 16**). Individual striations are gouged into the bedrock by single and relatively small clasts embedded and entrained in basal ice as it moves downstream (Påsse, 2004). *Grooves*, formed similarly, are the work of larger-sized clasts, cobbles and boulders. The clasts, cobbles and boulders are quarried from nearby rock walls and, as rock contact wears out and blunts their erosive competences, the distance covered by their gouging is proportional to the size of the clasts and cobbles (Lliboutry, 1994). Polished bedrock is the result of a much more comprehensive and evenly distributed type of wear, albeit by finegrained debris or clean ice, across the entire face of the exposed bedrock (Påsse, 2004).

Fluted ridges or flutes are linear (constructional) bedforms comprised of sub-glacial 355 sediments that have undergone post-depositional deformation (Boulton, 1976) (Figs. 17a-b). 356 Bracketed by shallow furrows or grooves, the flutes are low, typically  $\leq 5m$  in relative elevation 357 above the surrounding sediments (Hoppe and Schytt, 1953; Baranowksi, 1970; Glasser and 358 Hambrey, 2001; Benn and Evans, 2010), decimetres to metres wide, and tens of metres to hundreds 359 of metres long (Hoppe and Schytt, 1953; Roberson et al., 2001). When observed in groups, 360 individual flutes often exhibit metres-to multi-metres of separation (Hoppe and Schytt, 1953); the 361 groups also trend normal to the glacier's terminus and in line with the flow of the glacier (Roberson 362 et al., 2001). Although not the case in all instances, some flutes exhibit cobbles and boulders whose 363 364 long-axes also point in the direction of the ice movement (Hoppe and Schytt, 1953; Boulton, 1976).

Flutes develop beneath temperate glaciers and their warm-based ice, as till is pressured into 365 366 lee-side cavities of sub-glacially transported boulders and cobbles (e.g. Hoppe and Schytt, 1953; Hart, 1998; Glasser and Hambrey, 2001; Benn and Evans, 2010). Flute width is consistent 367 throughout the long axes and proportional to the boulders and cobbles that reside upslope of them 368 369 (Glasser and Hambrey, 2001). Bracketing furrows form adjacent to the ridges as the result of till migration into the lee-side boulder and cobble cavities (Boulton, 1976; also, Baranowski, 1970). 370 Since flute/furrow genesis is uniquely sub-glacial, assemblages thereof rarely are observed beyond 371 372 current or past glacial forelands (Glasser and Hambrey, 2001).

373 Boulder trains are linear clusters of erratic boulders that track the flow lines of former glaciers (Augustinus et al., 1997; Evenson et al., 2009; Hall and Phillips, 2006; Darvill et al., 2015) 374 (Figs. 18a-b). One of the two principal formation-hypotheses proposes that the constituent 375 boulders are plucked from rock outcrops, cliffs or tors and, subsequently, are dragged and 376 deposited linearly (albeit proximally) by basal ice as it flows downstream (Hall and Phillips, 2006; 377 Phillips et al., 2006; Darvill et al., 2015). According to this hypothesis, the distal deposition of the 378 379 boulders is neither expected nor observed (Augustinus et al., 1997; Hall and Phillips, 2006; Phillips et al., 2006; Darvill et al., 2015). Closely-set and cascadingly-smaller boulders downslope of the 380 381 larger lead boulder upslope is not infrequent, since the lead boulder would protect the lesser-sized boulders from further drag, and would be consistent with this hypothesis (Augustinus et al, 1997; 382 Hall and Phillips, 2006). 383

An alternate formation-hypothesis suggests that boulder trains are vestigial to rock avalanches, comprise markers of supraglacial transportation and are spread linearly by ice flow onto moraines (Evenson et al., 2009; Darvill et al., 2015). Distal deposition is not inconsistent with this hypothesis (Evenson et al., 2009). However, trains composed of cascadingly-smaller boulders downslope of the lead boulder upslope are explained less easily.

*Mega-scale glacial lineations* comprise a suite of parallel grooves and ridges that are aligned with the antecedent flow of ice (Clark et al., 2003; Storrar and Stokes, 2007; Benn and Evans, 2010; Fu et al., 2012) (**Fig. 19**). They exhibit up to tens of metres of height, hundreds of metres of width and as much as 100 km of length (Hättestrand and Clark, 2006; Fowler, 2010; Fu et al., 2012). The leading formation-hypothesis assumes that entrained boulders or rocks in basal ice act as "keels" that plough through soft sediments, carve elongate grooves and squeeze sediment into the intervening ridges (Clark et al., 2003).

### 396 6. Landscape interpretation

### 397 6.1 Basin-surface and ridge/trough nested polygons

The morphology and metres-scale of the basin-surface polygons and the ridge and trough 398 nested polygons in our study region are consistent with polygons formed by thermal-contraction 399 in permafrost on Earth (e.g. Lachenbruch, 1962; Oehler et al., 2016) and, possibly, elsewhere on 400 Mars (e.g. Pechmann, 1980; Costard and Kargel, 1995; Seibert and Kargel, 2001; Soare et al., 401 2008; Levy et al., 2009a,b; Oehler et al., 2016). However, where only HCPs or polygons that 402 exhibit no topographical variance between centres and margins are observed, as is the case with 403 404 the basin-surface polygons, identifying marginal fill is neither simple nor straightforward. Planview observations derived of HiRISE imagery, even at the best resolution and magnification, are 405 equivocal; the aggradational and degradational morphologies of sand- or ice-wedge polygons are 406 relatively congruent when observed remotely. 407

On Earth, ground truth invariably underlies the discrimination of the two polygon types.
On Mars, by contrast, a landscape-scale evaluation and conciliation of individual features (e.g.
Baker, 2003; Hauber et al., 2011) may comprise the most effective pathway for deducing marginal
fill and polygon origin.

### 412 6.2 Polygonised depressions

The polygonised terrain is our study region is punctuated at some locations by shallow circular to sub-circular depressions (**Fig. 5c**), lineated and parallel to sub-parallel variances of topography (**Figs. 5a-b**) and, in one kilometres-scale basin, cascading equator-ward scarps (**Fig.** 416 4). In permafrost regions on Earth polygonised terrain with similar variances of topography typically occur where/when ice-rich terrain undergoes thermal degradation, deflation or subsidence, and volumetric loss. These losses are the work of evaporation and/or meltwater
migration. Thaw-derived run-off channels often facilitate the latter.

On Mars, contrarily, none of the thermokarst-like landscapes exhibit run-off channels. If and when the atmospheric vapour-pressure at the mid-latitudes is low (relative to the vapour pressure of near-surface ice), and the boundary conditions are well below the triple point of water, the most plausible agent of ice loss would be sublimation (e.g. Morgenstern et al., 2007; Lefort et al., 2009, 2010; Ulrich et al., 2010; Séjourné et al., 2011; Dundas et al., 2015; Dundas, 2017).

This having been said, hypotheses concerning ice enrichment and ice depletion need not be univocal in identifying plausible environmental triggers. There is no conceptual or geological inconsistency in proposing ice enrichment by the freeze-thaw cycling of water and ice-depletion by sublimation. Inconsistencies are reared, however, if and when ice enrichment is explained by means of sublimation and adsorption cycles.

For example, the heterogenous distribution of excess ice in terrestrial landscapes is the result of highly-localised differences in sedimentary permeability, pore space and tortuosity. In turn, this facilitates or impedes the formation of segregation ice or ice lenses and the heaved deformation of terrain surfaces. Hypotheses favouring the origin and development of excess ice by adsorption-diffusion cycles implicitly assume a latitudinal homogeneity in the composition of near-surface regolith and of its ice richness (Mellon and Jakosky, 1993; also, Dundas et al, 2015). These hypotheses are weakened by two constraints.

First, and notwithstanding questions concerning the plausibility of regolith homogeneity on the scale assumed here, diffusion-adsorption cycles are self-limiting. Once the pore space of near-surface regolith has become saturated with adsorbed water ice, transport to a sub-adsorbed ice depth is choked off (Clifford, 1993; Mellon and Jakosky, 1993). As such, the decametres of possible thermokarst depth observed in one of our study-region basins, and elsewhere on Mars
(e.g. Morgenstern et al., 2007; Lefort et al., 2008; Séjourné et al., 2011), lie beyond the explanatory
reach of the adsorption-diffusion hypothesis. The highly iterative freeze-thaw cycling of water
would not be proscribed similarly.

445 *6.3 Ridge and trough terrain* 

Overlying portions of the polygonised surface of the basins in our study region is terrain 446 comprised of interconnected ridges and troughs, also referenced as *brain terrain* in the literature 447 (e.g. Levy et al., 2009a, b) (Figs. 6, 7a-c). They are metres in elevation and metres to decametres 448 449 in width. Basin-ward flanks of the ridge and trough assemblages are less dense and thick than the 450 assemblages closer to the basin margins. Outcrops of these assemblages, possibly remnants of basin coverage by them that was more extensive than today, also are observed (Fig. 7b). The ridges 451 452 and troughs could be the ablated remnants of an icy mantle (e.g. Levy et al., 2009a, b) precipitated atmospherically and accumulated episodically in the Late Amazonian Epoch (e.g. Head et al., 453 2003; Milliken et al., 2003; Madeleine et al., 2009; 2014; Shorghofer and Forget, 2012). 454

455 *6.4 Mounds* 

Decametre-wide, metres-high and circular to sub-circular mounds are observed throughout the polygonised basins in our study region (**Figs. 3b, 5, 8a-d**). Mound summits show fracturing and possible depletion. Mound surfaces, as noted above, exhibit polygonization. Mound distribution, typically, is clustered. Isolated outliers are infrequent. No mounds are observed on the terrain adjacent to the basin.

461 Nearby, i.e. the Moreux impact-crater region (Soare et al, in press), and elsewhere, i.e. the
462 mid-latitudes of Utopia Planitia (Soare et al., 2005; 2013; 2020; Dundas et al., 2008; de Pablo and
463 Komatsu, 2009), equatorial Athabasca Valles (Burr et al., 2005; Page and Murray, 2006; Balme

464 and Gallagher, 2009), and in the Argyre impact-crater region (Soare et al., 2014), pingo-like mounds that show similarities of size, shape, polygonization and possible degradation have been 465 observed. Mound origin is deduced from and categorized by mound location. For example, if the 466 mounds are nested in plains, basins and thermokarst-like depressions then a hydrostatic origin is 467 hypothesized (e.g. Soare et al., 2005; 2013; 2020; Burr et al., 2005; Balme and Gallagher, 2009). 468 A slope-side location in the Argyre region, adjacent to graben-like cavities and possible paleo-469 discharges of water, highlights the one location where a hydraulic origin for the pingo-like mounds 470 has been hypothesized (Soare et al., 2014). 471

472 6.5 Bow-like ridges and boulder/cobble lineations

Unlike periglacial landscape features such as small-sized polygons and pingo-like mounds 473 whose form does not necessarily constrain one and only one possible origin, the connexion 474 between process and form is less equivocal with regard to the bow-like ridges that populate the 475 (inner) eastern margin of the two main basins in our study zone. For example, the ridges are 476 topographically-irregular through their fronts and flanks, are sometimes serialized and are 477 478 distributed proximally to possible valley-constrained flows (Figs. 9, 10a). These traits are 479 commonplace amongst glacial moraines on Earth (Figs. 15a, c). On Mars, the moraine-like ridges are thought to be no less diagnostic of glaciation, especially when they are observed amidst other 480 glacier-like forms such as lineated valley fill and sinuous esker-like channels (e.g. Arftstrom and 481 Hartmann, 2005; Banks et al., 2008; Dickson et al., 2008; Head et al., 2010; Hubbard et al., 2011, 482 483 2014; Souness et al., 2012; Brough et al., 2016; Hepburn et al., 2020).

484 Speculation about the origin of boulder-alignments on Mars is wide-ranging and stretches 485 from periglacially induced self-sorting through to mass wasting or englacial flow (e.g. Banks et 486 al., 2008; Hubbard et al., 2011, 2014). A number of variables weigh against (periglacial) self487 sorting. First, decametre-scale boulders are not observed in self-sorted landscapes on Earth; convectional forces and frost heave substantially greater than those expressed terrestrially would 488 be required for stripes comprised of these large boulders to form. Second, some of the boulder 489 alignments and spatially-associated grooves are contiguous with grooves in the basin-adjacent 490 491 terrain. This would be more consistent with an erosive origin, proximally, and to mass movement or entrainment than with freeze-thaw cycling and the work of periglaciation. Third, the density of 492 boulder distribution and of boulder alignments decreases as the distance to the basin's edge 493 increases. This too points to the spatial proximity of source material and away from *in situ* self-494 495 sorting. With regard to mass wasting, it is not clear that the gravity-induced flow or slide of boulder material in and of itself could engender the tightly distributed lines of boulders and grooves 496 observed here. 497

By way of scale, the basin-margin grooves are too narrow and shallow to fit the terrestrial 498 definition of mega glacial-lineations, as well as too wide and deep to be deemed striations (Fig. 499 10b). Setting aside these scale-based definitions, the grooves could have been gouged by glacially-500 501 entrained boulders whose size befits the actual width and depth of the former. This hypothesis also would be consistent with the contiguous gouging observed upslope of the grooves in the basin-502 503 adjacent terrain. Where the mega-lineation explanation falls short, however, is with regard to the presence of groove-nested or -adjacent boulders at various basin-margin locations throughout our 504 study region. This suggests deposition, not erosion, and the possibility that the lineated grooves 505 506 and boulder lineations are the result of two separate and distinct glacial regimes.

507 As noted above, some boulder alignments do exhibit smaller boulders downslope of larger 508 ones and, in as much the distribution and density of boulder alignments increases with its proximity 509 to an upslope source on the basin-adjacent terrain, they could be boulder trains (**Figs. 10b-c**). At the same time, the entrainment of boulders and the gouging of erosive grooves or furrows is not a keynote characteristic of boulder trains. Once again, a separate and distinct glacial regime would have to be inferred as a complement to the boulder-train hypothesis for its plausibility to be warranted.

Of the glacially-associated explanations of boulder alignment, the glacial-flute hypothesis befits the geological context of our observations most closely For example, the hypothesis concurrently encompasses the formation of small-scale grooves and of adjacent terrain that is slightly elevated and bouldered, as seen in **Fig. 10b**. The adjacency of boulder sources proximal to the possible flutes (**Fig. 10d**) offers tertiary support of the hypothesis, although this would be equally supportive of the boulder-train hypothesis.

The ridge and trough terrain is thought to comprise the sublimated remnants of surface ice on Mars that is precipitated atmospherically (e.g. Levy et al., 2009a, b); as such, it could be considered artefactual to glaciation. However, this terrain has no observed analogue on Earth and questions concerning the origin of its distinctive and unique morphology remain largely unanswered.

### 525 7. Age-estimates and the temporal-ordering of local periglaciation and glaciation

### 526 7.1 Glacial Stade I (inferred not observed)

The clastic composition of the terrain incised by the small-sized polygons and the thermokarst-like depressions is unknown. However, for ice enrichment and wedging to occur the terrain would have needed to incorporate relatively fine-grained material. Then, as now, the removal, transport and deposition of regolith by the work of wind is a key player in Mars' dynamic surface-environment. Volcanic ash or loess, for example, could have been brought to the region by aeolian activity sufficient to form a thick surface to depth horizon or sets of horizons (e.g. Greeley and Williams, 1994; Bridges et al., 2010; Séjourné et al., 2012; Bridges and Muhs, 2012;
Skinner et al., 2012; Soare et al., 2014; Smalley et al., 2019).

The possibility that variances of obliquity and eccentricity have facilitated and induced the 535 atmospheric precipitation of dusty snow/ice and its surface accumulation at the northern mid-536 latitudes throughout the Late Amazonian Epoch is well documented in the literature (e.g. Jakosky 537 et al., 1995, Head et al., 2003, Laskar et al., 2004; Madeleine et al., 2009, 2014; Shorghofer and 538 Forget, 2012). Were this dusty/ice snow to have undergone thaw, with meltwater migrating into 539 regolith composed of volcanic ash or loess, this would have framed an ideal set of boundary 540 conditions for the freeze-thaw cycling of that meltwater (Soare et al., 2014). We define this 541 hypothesized period of glaciation and deglaciation as Glacial Stade-I, and suggest that it preceded 542 the **Type-1** polygonization of the currently-exposed basin surfaces in our study region. 543

### 544 7.2 Periglacial Period I

Age estimates of these **Type-1** polygonised basins derived from the evaluation of *CSFD* raise the possibility that the polygons are as old as 100 Ma (see Appendix A, Figs. 20a-e). We define this formation period as **Periglacial Period-I** (**PP-I**). The 100 Ma age estimate, if valid, is roughly an order of magnitude greater than the relatively recent age estimate of the fragmented or degraded terrain, possibly icy in origin, that overlies the **Type-1** polygons at some locations (**Fig. 7c**).

### 551 7.3 Glacial Stade II (possible remnants observed); Periglacial Period II

Fragmented-terrain members include: networked ridge and trough structures (i.e. brain terrain) pingo-like mounds and mound remnants; and, etched **Type-2** polygons that incise the mounds, mound remnants and adjacent terrain (**Figs. 7a-c, 8a-d**). Based on this relative stratigraphy and the *CFSD* of the brain terrain, we suggest that the **Type-2** polygons, possibly the mounds as well, regardless of whether they are open or closed, formed during a second and more recent periglacial period (**PP-II**), i.e. ~1 - ~10 Ma, than **PP-I**. Moreover, if brain terrain comprises glacial residue whose distribution varies between dense to sparse throughout the **Type-1** polygonised basins, then the antecedent presence of an icy mass that covers the basins completely (**Glacial Stade-II**) does not seem implausible. Left open is the question of whether the thermokarstic depressions formed during **PP-I** or **PP-II**.

All of the observed thermokarst-like depressions in the valley basins are incised by **Type**-562 1 polygons, many of which show high-centres. We suggest that the devolatilized depressions 563 564 formed prior to their polygonization during **PP-I**. Whether devolatilization was the work of thaw or sublimation is moot, as long as one differentiates between periods of ice enrichment, which 565 require water to undergo freeze-thaw cycling, and periods of ice loss, which do not. Arguably, 566 567 devolatilization could have postdated GS-II, been concurrent with PP-II and not involved the freeze-thaw cycling of water. Similarly large offsets of time between enrichment and depletion are 568 commonplace, for example. amidst late Quaternary Epoch and Holocene era permafrost 569 570 landscapes on Earth (e.g. Rampton, 1973, 1988). On the other hand, devolatisation of metres-deep ice-rich terrain could have occurred within a relatively short period of time during **PP-II**, as has 571 572 happened throughout the very late Holocene era (e.g. Farquaharson et al., 2019), when and if conditions of water meta-stability arose in our study region. 573

574 *7.4 Glacial Stade III* 

575 On the eastern margin of the southernmost **Type-1** polygonised basin in our study zone 576 some of the bouldery and debris-covered ridges, possibly moraines, intercept ridge and trough 577 remnants. The amplitude and mass of the bouldery ridges is greater at these contacts than where 578 the ridge and trough remnants are absent. This apparent push-morphology suggests that the remnants predate the bouldery ridges (Fig. 11). As such, the moraine-like structures could be
artefacts of a separate and distinctly successive period of glaciation, i.e. Glacial Stade III (GSIII).

To the east and upslope of the bouldery ridges, lineations reminiscent of glacial flutes and boulder-trains are observed (**Figs. 10a-b**). Some of them extend almost to the bouldery and moraine-like ridges themselves (**Fig. 10c**). They too could be features associated with **GS-III**. Interestingly, the possible flutes and boulder trains superpose polygons whose shape, size, texture and networked distribution matches the **Type-1** polygons that populate the adjacent basin to the west. At some locations, the two sets of polygons intersect and we interpret the sub-flute polygons to be formationally concurrent with the **Type-1** polygons that incise the basin polygons (**Fig. 2c**).

589 8. Conclusion

Using cross-cutting relationships, relative stratigraphy and crater-size frequency distributions (*CSFDs*) we have identified landscape features within our study region that point to the existence of three disparate albeit temporally intertwined glacial stades and two periglacial periods. Age estimates of the earliest polygons, **Type-1**, and of the polygonally-incised thermokarst-like depressions in our basin-based landscapes suggest an origin that could be 100 Ma. This is millions of years older and almost an order of age magnitude beyond that of the more youthful **Type-2** polygons.

Heretofore, much of the discussion about periglaciation on Mars has focused on its relative recentness and the glaciation that preceded it (e.g. Mustard et al., 2001; Head et al., 2003; Levy et al., 2009a, b; Séjourné et al., 2011; Schon et al., 2012; Sinha and Murty, 2015). The older estimation of periglaciation in our study region and the intertwined chronology of two (possible) periglacial periods and three glacial stades documented here suggests that the cycling of the two 602 environment types could be much more Earth-like and extend much more deeply into Mars'603 history than had been thought hitherto.

604 Appendix

605 Crater counts provide constraints on the exposure and crater retention ages of land surfaces. 606 Craters measured on the *CTX* image (**Fig 20a.**), below, across all surfaces generates a broad crater-607 retention age for our study region immediately to the north of the Moreux impact-crater. The *CSFD* 608 for crater diameters larger than 200 m are well approximated by a 1.5 Ga model isochron, assuming 609 the Hartmann (2005) model. This is broadly similar to the crater-count age estimate derived by 610 Sinha and Murty (2015).

Crater counts also were conducted on the polygonised basin-surface (**area #1**) and ridgetrough terrain (**area #2**) in *HiRISE* image ESP\_045349\_2235 (**Fig. 20b**) and the polygonised basin-surface (**area #3**) in *HiRISE* image ESP\_042105\_2235 (**Fig. 20c**). The *CSFDs* of these areas do not conform to a production function and the shallower power-law slope is indicative of a loss of smaller diameter craters from the populations. This is consistent with and similar to the age estimates derived of the polygonised basin surface on the eastern margin of the Moreux impactcrater (Soare et al., 2020, in press).

Various processes may alter a crater population on Mars: burial, erosion and exhumation.
Each of these processes preferentially remove smaller-diameter craters from the landscape. A
shallower power-law slope results from the size-dependent deficit of smaller craters (see reviews
by Williams et al., 2018 and Rubanenko et al. 2020 and references therein for further discussion).
We note that the larger craters are shallow, possibly infilled, and polygonized consistent
with older, more heavily modified craters compared with the smaller diameter craters which have

624 more pristine, bowl-shaped morphologies due to the relatively rapid loss of craters at these 625 diameters.

The CSFDs of the polygonised terrains in areas #1 and #3 (Figs. 20d-e) extend across 626 627 isochrons ~1 Ma at the small diameters and extend to >10 Ma at the large diameters with the largest craters in area #3 overlapping the 500 Ma isochron. This suggests that the polygonised terrains 628 are older than 10 Ma, possibly as old as ~100 Ma or more. The CSFD in area #2, composed of 629 ridges and troughs, extends from ~1 Ma to ~10 Ma. Areas #1 and #3 generally contain larger 630 diameter craters than area #2. This points to the polygonised terrain being older than the latter, 631 possibly by 10s to 100s Ma, and is consistent with the observed overprinting of the polygonised 632 basin-surface by the ridge and trough terrain in HiRISE image ESP\_045349\_2235. The similar 633 density of the more pristine, bowl-shaped craters on all three areas at the smaller sizes indicates 634 that the characteristic time of the retention of these smaller craters is more uniform across terrains. 635

```
636 References
```

Allen, C.R., O'Brien, M.G., Sheppard, S.M.F. 1976. The chemical and isotopic characteristics of
 some northeast Greenland surface and pingo waters, Arctic and Alpine Research, 8, 3, 297 317.

640 Arfstrom, J., Hartmann, W.H. 2005. Martian flow features, moraine-like ridges, and gullies:

- 641 Terrestrial analogs and interrelationships. Icarus 174, 321-335, doi:10.1016/j.icarus.2004.
  642 05.026.
- Augustinus, P.C., Gorp, D.B., Leishman, M.R., Zwartz, D. 1997. Reconstruction of ice flow across
  the Bunger Hills, East Antarctica. Antarctic Science 9, 3, 347-354.
- Baker, D.M.H., Head, J.W. 2015. Extensive Middle Amazonian mantling of debris aprons and

646	plains in Deuteronilus Mensae, Mars: Implications for the record of mid-latitude glaciation.
647	Icarus 260, 269-288, doi.org/10.1016/j.icarus.2015.06.036.
648	Baker, V.R. 2003. Icy Martian mysteries. Nature 426, 779-780, doi.org/10.1038/426779a.
649	Balme, M.R, Gallagher, C. 2009. An equatorial periglacial landscape on Mars. Earth and
650	Planetary Science Letters 285, 1-15, doi.10.1016/j.epsl.2009.05.031.
651	Banks, M.E., et al. 2008. High Resolution Imaging Science Experiment (HiRISE) observations of
652	glacial and periglacial morphologies in the circum-Argyre Planitia highlands, Mars.
653	Journal of Geophysical Research 113, E12015, doi:10.1029/2007JE002994,
654	Baranowski, S. 1970. The Origin of Fluted Moraine at the Fronts of Contemporary Glaciers.
655	Geografiska Annaler. Series A, Physical Geography 52, 1, 68-75.
656	Barry, R.G, Gan, T-Y. 2011. The global cryosphere: past, present and future. Cambridge:
657	Cambridge University Press.
658	Benn, D., Evans, D. 2010. Glaciers and glaciation, London UK, Hooder Education.
659	Bennike, O. 1983. Pingos in Peary Land, North Greenland. Bulletin of the Geological Society of
660	Denmark 32, 97-99.
661	Bennike, O. 1998. Pingos at Nioghalvfjerdsfjorden, eastern North Greenland. Geology of
662	Greenland Survey Bulletin 180, 159-162.
663	Bjarnadóttir, V. 2015. Flutes in the forefield of Nordenskiöldbreen, Svalbard. BSc Thesis. Faculty
664	of Earth Sciences School of Engineering and Natural Sciences University of Iceland,
665	Reykjavík, 35 p.
666	Black, R.F. 1954. Permafrost - a review. Bulletin of the Geological Society of America 85, 839-
667	856.
668	Boulton, G.S. 1976. The origin of glacially fluted surface observations and theory. Journal of

669 Glaciology. 17, 76, 287-309.

- Bridges, N.T., et al. 2010. Eolian bedforms, yardangs, and indurated surfaces in the Tharsis Montes
  as seen by the HiRISE camera: evidence for dust aggregates. Icarus 205:165-182, doi.org
- 672 /10.1016/ j.icarus.2009.05.017.
- Bridges, N.T., Muhs, D.R. 2012. Duststones on Mars: source, transport, deposition and erosion.
  Sedimentary Geology of Mars, SEPM special issue 112, 169-182.
- Brough, S., Hubbard, B., Hubbard, A. 2016. Former extent of glacier-like forms on Mars. Icarus
  274, 37-49, doi.org/10.1016/j.icarus.2016.03.006.
- Burr, D.M., Soare, R.J., Wan Bun Tseung, J.M., Emery, J. 2005. Young (late Amazonian), near
- surface, ground ice features near the equator, Athabasca Valles, Mars. Icarus 178, 56-73.
- 679 Clark, C.D., Tulczyk, S.M., Stokes, C.R., Canals, M. 2003. A groove-ploughing theory for the
- 680 production of mega-scale glacial lineations, and implications for ice-stream mechanics.
- 681 Journal of Glaciology 49, 165, 240-256.
- 682 Costard, F.M., Kargel, J.S. 1995. Outwash plains and thermokarst on Mars. Icarus 114, 1, 93-
- 683 112, doi.10.1006/icar.1995.1046.
- 684 Cruickshank, J.G., Colhoun., E.A. 1965. Observations on Pingos and Other LandForms in
  685 Schuchertdal, Northeast Greenland. Geografiska Annaler. Series A, Physical Geography
  686 47, 4, 224-236.
- 687 Czudek, T., Demek, J. 1970. Thermokarst in Siberia and its influence on the development of
  688 lowland relief. Quaternary Research 1, 103-120.
- De K. Leffingwell, E. 1915. The dominant form of ground-ice on the north coast of Alaska. The
  Journal of Geology 23, 7, 635-654.
- 691 De Pablo, M.A., Komatsu, G. 2009. Possible pingo fields in the Utopia basin, Mars: Geological

- and climatical implications. Icarus 49-74, doi:10.1016/j.icarus.2008.09.007.
- 693 Dickson, J.L, Head, J.W., Marchant, D.R. 2008. Late Amazonian glaciation at the dichotomy
- 694 boundary on Mars: Evidence for glacial thickness maxima and multiple glacial phases.
- 695 Geology 36, 5, 411–414; doi:10.1130/G24382A.
- Donner, J. 1978. Holocene history of the west coast of Disko, central west Greenland.
- 697 Geografiska Annaler. Series A, Physical Geography 60, 12, 63-72.
- Dundas, C.M., Mellon, M.T., McEwen, A.S., Lefort, A., Keszthelyi, L.P., Thomas, N., 2008.
- 699 HiRISE observations of fractured mounds: possible martian pingos. Geophysical Research
- 700 Letters 35, L04201, doi.org/10.1029/2007GL031798.
- Dundas, C.M., McEwen, A.S. 2010. An assessment of evidence for pingos on Mars using HiRISE.
  Icarus 205, 1, 244-258, doi.org/10.1016/j.icarus.2009.02.020.
- Dundas, C.M., Byrne, S., McEwen, A.S. 2015. Modeling the development of Martian sublimation
  thermokarst landforms. Icarus 262, 154-169, doi.org/10.1016/j.icarus.2015.07.033.
- 705 Dundas, C.M. 2017. Effects of varying obliquity on Martian sublimation thermokarst landforms.
- 706 Icarus 281, 1, 115-120, doi.org/10.1016/j.icarus.2016.08.031.
- 707 Easterbrook, D.J. 1991. Surface processes and landforms. New Jersey: Prentice Hall.
- Evenson, E.B., et al. 2009. Enigmatic boulder trains, supraglacial rock avalanches, and the origin
  of "Darwin's boulders," Tierra del Fuego. *GSA* Today 19, 12, doi:10.1130/GSATG72A.1.
- 710 Farquharson, L.M., Romanovsky, V.E., Cable, W.L., Walker, D.A., Kokelj, S.V., Nicolsky, D.
- 711 2019. Climate Change Drives Widespread and Rapid Thermokarst Development in Very
- 712 Cold Permafrost in the Canadian High Arctic. Geophysical Research Letters 6681-6689,
- 713 doi.org/10.1029/2019GL082187.
- Fjellanger, J., Sørbel, L., Linge, H., Brook, E.J., Raisbeck, G.M., Yiou, F. 2006. Glacial survival

715	of blockfields on the Varanger Peninsula, northern Norway. Geomorphology 82, 255-272.
716	Fowler, A.C. 2010. The formation of subglacial streams and mega-scale glacial lineations.
717	Proceedings of the Royal Society 466, 181-3201, doi:10.1098/rspa.2010.0009.
718	French, H.M. 2007. The periglacial environment, 3 <sup>rd</sup> ed., J. Wiley & Sons, West Sussex, England,
719	458 p.
720	French, H.M., Guglielmin, M. 2000. Frozen ground phenomena in the vicinity of Terra Nova Bay,
721	northern Victoria Land, Antarctica: a preliminary report. Geografiska Annaler 82, A, 513-
722	526.
723	Frey, H.V., Roark, J.H., Shockey, K.M., Frey, E.L., Sakimoto, S.E.H. 2002. Ancient lowlands on
724	Mars, Geophysical Research Letters 29, 10, 1384,10.1029/2001GL013832.
725	Fu, P., Heyman, J., Hättestrand, C., Stroeven, A.P., Harbo, J.M. Glacial geomorphology of the
726	Shaluli Shan area, southeastern Tibetan Plateau. Journal of Maps 8:1, 48-55, doi:10.1080/
727	17445647.2012.668762.
728	Glasser, N.F., Hambrey, M.J. 2001. Styles of sedimentation beneath Svalbard valley glaciers under
729	changing dynamic and thermal regimes. Journal of the Geological Society, London 158,
730	697-707.
731	Greeley, R., Williams, S.H. 1994. Dust deposits on Mars: the "Parna" analog. Icarus 110, 165-
732	177.
733	Grosse, G., Schirrmeister, L., Siegert, C., Kunitsky, V.K., Slagoda, E.A., Andreev, A.A.,
734	Dereviagyn, Y. 2007. Geological and geomorphological evolution of a sedimentary
735	periglacial landscape in northeast Siberia during the late Quaternary. Geomorphology 86,
736	25-51, doi.10.1016/j.geomorph.2006.08.005.
737	Hall, A.M., Phillips, W.M. 2006. Glacial modification of granite tors in the Cairngorms, Scotland.

- 738 Journal of Quaternary Science 21, 8, 811-830, doi:10.1002/jqs.1003.
- Hallet, B., Sletten, R., Whilden, K. 2011. Micro-relief development in polygonal patterned ground
- in the Dry Valleys of Antarctica. Quaternary Research 75, 347-355, doi.org/10.1016/j.
  yqres.2010.12.009.
- 742 Hambrey, M.J. 1994. Glacial environments. UBC Press.
- 743 Harris, S.A., French, H.M., Heginbottom, J.A., Johnston, G.H., Ladanyi, B., Sego, D.C., van
- Everdingen, R.O. (eds.), 1988. Glossary of permafrost and related ground-ice terms.
  Technical Memorandum 142, Permafrost Subcommittee, National Research Council of
  Canada, 154 p.
- Hartmann, W.K. 2005. Martian cratering 8: Isochron refinement and the chronology of Mars.
  Icarus 174, 294-320, doi.org/10.1016/j.icarus.2004.11.023.
- 749 Hart, J.K. 1998. The deforming bed/debris basal ice continuum and its implications for the
- formation of glacial landforms (flutes) and sediments (melt-out till). Quaternary Science
  Reviews 7, 737-754.
- Hättestrand, C., Clark, C.D. 2006. The glacial geomorphology of Kola Peninsula and adjacent
- areas in the Murmansk Region, Russia, Journal of Maps, 2:1, 30-42, doi:10.4113/
  jom.2006.41.
- Hauber et al. 2011. Periglacial landscapes on Svalbard: Terrestrial analogs for cold-climate
  landforms on Mars. The Geological Society of America Special Paper 483, 177-201, doi:
  10.1130/2011.2483(12).
- Head, J.W., Kreslavsky, M.A., Pratt, S. 2002. Northern lowlands of Mars: Evidence for
- videspread volcanic flooding and tectonic deformation in the Hesperian Period. Journal of
- 760 Geophysical Research doi.org/10.1029/2000JE001445.

761	Head, J.W., Mustard, J.F., Kreslavsky, M.A., Milliken, R.E., Marchant, D.R. 2003. Recent ice
762	ages on Mars. Nature 426, 797-802, doi.10.1038/nature02114.
763	Head, J.W., Marchant, D.R., Dickson, J.L., Kress, A.M., Baker, D.M. 2010. Northern mid-latitude
764	glaciation in the Late Amazonian period of Mars: criteria for the recognition of debris-
765	covered glacier and valley glacier land-system deposits. Earth and Planetary Science
766	Letters 294. 306-320, doi.10.1016/j.epsl.2009.06.041.
767	Hepburn, A. J., Ng, F., Livingstone, S. J., Holt, T., Hubbard, B. 2020. Polyphase mid-latitude
768	glaciation on Mars: Chronology of the formation of superposed glacier-like forms from
769	crater-count dating. Journal of Geophysical Research 125, e2019JE006102, doi.org/10.
770	1029/2019JE006102.
771	Hoppe, G., Schytt, V. 1953. Some Observations on Fluted Moraine Surfaces Geografiska
772	Annaler 35, 2, 105-115.
773	Hubbard, B., Milliken, R.E., Kargel, J.S., Limaye, A., Souness, C. 2011. Characterisation and
774	interpretation of a mid-latitude glacier-like form: Hellas Planitia, Mars. Icarus 211, 330-
775	346, doi.10.1016/j.icarus.2010.10.021.
776	Hubbard, B., Souness, C., Brough, S. 2014. Glacier-like forms on Mars. The Cryosphere 8, 2047-
777	2061, doi:10.5194/tc-8-2047-2014.
778	Hughes, O.L. 1974. Geology and permafrost in relation to hydrology and geophysics, in
779	Proceedings of Workshop Seminar, 1974: Ottawa, Canadian National Committee,
780	International Hydrological Decade, 21-30.
781	Hussey, K.M., Michelson, R.W. 1966. Tundra relief features near Point Barrow, Alaska. Arctic,
782	19, 2, 162-184.
783	Jakosky, B.M., Henderson, B.G., Mellon, M.T. 1995. Chaotic obliquity and the nature of the

784 martian climate. J. Geophys. Res. 100, 1579-1584.

- Kneissl, T., van Gasselt, S., Neukum, G. 2011. Map-projection-independent crater size-frequency
  determination in GIS environments new software tool for ArcGIS. Planetary and Space
  Science 59, 1243-1254, doi.org/10.1016/j.pss.2010.03.015.
- Kokelj, S.V., Burn, C.R., Tarnocai, C. 2007. The Structure and Dynamics of Earth Hummocks in
  the Subarctic Forest Near Inuvik, Northwest Territories, Canada, Arctic, Antarctic, and
  Alpine Research, 39,1, 99-109, doi.org/10.1657/1523-0430(2007)39[99:TSADOE]2.0.
  CO;2.
- Lachenbruch, A.H., 1962. Mechanics of Thermal Contraction Cracks and Ice-wedge Polygons in
   Permafrost. GSA Special Paper 70. vol. 69 Geological Society of America, New York.
- Laskar, J., Correia, A.C.M., Gastineau, M., Joutel, F., Levrard, B., Robutel, P. 2004. Long term
  evolution and chaotic diffusion of the insolation quantities of Mars. Icarus 170, 343-364.
- Lefort, A., Russell, P.S., McEwen, A.S., Dundas, C.M., Kirk, R.L. 2009. Observations of
- periglacial landforms in Utopia Planitia with the High Resolution Imaging Science
  Experiment (*HiRISE*). Journal of Geophysical Research 114, E04005, doi.10.1029/
  2008JE003264.
- Lefort, A., Russell, P.S., Thomas, N. 2010. Scalloped terrains in the Peneus and Amphitrites
  Paterae region of Mars as observed by HiRISE, Icarus, 205, 259-268, doi:10.1016/j.icarus.
  2009.06.005.
- Levy, J., Head, J., Marchant, D. 2009a. Concentric crater fill in Utopia Planitia: History and
- 804 interaction between glacial "brain terrain" and periglacial mantle processes. Icarus 202,
  805 462-476, doi.10.1016/j.icarus.2009.02.018.
- Levy, J., Head, J., Marchant, D. 2009b. Thermal contraction crack polygons on Mars:
- 807 Classification, distribution, and climate implications from HiRISE observations 114,
  808 E01007, doi:10. 1029/2008JE003273.
- Lliboutry, L.A. 1994. Monolithologic erosion of hard beds by temperate glaciers. Journal of
  Geology 40, 136, 433-450.
- Mackay, J.R. 1974. Ice wedge cracks, Garry Island, Northwest Territories. Canadian Journal of
  Earth Sciences 11, 1366-1383.
- Mackay, J.R. 1986. Growth of Ibyuk Pingo, Western Arctic Coast, Canada, and Some Implications
  for Environmental Reconstructions. Quaternary Research 26, 1, 68-80.
- 815 Mackay, J.R. 1998. Pingo growth and collapse, Tuktoyaktuk Peninsula areas, western arctic
- 816 coast, Canada: a long-term field study. Géographie physique et Quaternaire 52, 3, 1-53.
- 817 Mackay, J.R. 1999. Periglacial features developed on the exposed lake bottoms of seven
- 818 lakes that drained rapidly after 1950, Tuktoyaktuk Peninsula Area, western arctic coast,
- 819 Canada. Permafrost and Periglacial Processes 10, 39-63.
- 820 Madeleine, J-B., Forget, F., Head, J.W., Levrard, B., Montmessin, F., Millour, E. 2009.
- Amazonian northern mid-latitude glaciation on Mars: a proposed climate scenario. Icarus
  203, 390-405, doi.org/10.1016/j.icarus.2009.04.037.
- 823 Madeleine, J-B. et al. 2014. Recent ice ages on Mars: the role of radiatively active clouds and
- 824 cloud microphysics. Geophysical Research Letters 41, 14, 4873-4879, doi.10.1002/2014G
  825 L059861.
- 826 Malin, M.C. et al. 2007. Context Camera Investigation on board the Mars Reconnaissance Orbiter.
- Journal of Geophysical Research 112, E05S04, doi.org/10.1029/2006JE002808.
- 828 Marchant, D.R. et al. 2002. Formation of patterned ground and sublimation till over Miocene

- glacier ice in Beacon Valley, southern Victoria Land, Antarctica. Geological Society of
  America Bulletin 114, 6, 718-730.
- 831 McEwen, A.S. et al. 2007. Mars Reconnaissance Orbiter's High-Resolution Imaging Science
- 832 Experiment (HiRISE). Journal of Geophysical Research 112, E05S02, doi.org/10.1029/
  833 2005JE002605.
- McGill, G.E., Dimitriou, A.M. 1990. Origin of the Martian global dichotomy by crustal thinning
  in the Late Noachian or Early Hesperian. Journal of Geophysical Research doi.org/10.
  1029/JB095iB08p12595.
- Mellon, M.T., Jakosky, B.M. 1993. Geographic Variations in the Thermal and Diffusive Stability
  of Ground Ice on Mars. Journal of Geophysical Research 98, E2, 3345-3364.
- Michael, G.G., Neukum, G. 2010. Planetary surface dating from crater size-frequency distribution
  measurements: partial resurfacing events and statistical age uncertainty. Earth and
  Planetary Science Letters 294, 223-229. doi.org/10.1016/j.epsl.2009.12.041.
- 842 Michael G.G., Kneissl T., Neesemann, A. 2016. Planetary surface dating from crater size-
- 843 frequency distribution measurements: Poisson timing analysis. Icarus 277, 279-285,
  844 doi.org/10.1016/j.icarus.2016.05.019.
- 845 Milliken, R.E., Mustard, J.F., Goldsby, D.L. 2003. Viscous flow features on the surface of Mars:
- 846 Observations from high-resolution Mars Orbiter Camera (MOC) images. Journal of
  847 Geophysical Research 108, E6, 5057, doi.10.1029/2002JE002005.
- 848 Morgan, G.A., Head, J.W., Marchant, D.R. 2009. Lineated valley fill (LVF) and lobate debris
- aprons (LDA) in the Deuteronilus Mensae northern dichotomy boundary region, Mars:
- 850 Constraints on the extent, age and episodicity of Amazonian glacial events 202, 1, 22-38,
- doi.org/10.1016/j.icarus.2009.02.017.

852	Morgenstern, A., Hauber, E., Reiss, D., van Gasselt, S., Grosse, G., Schirrmeister, L. 2007.
853	Deposition and degradation of a volatile-rich layer in Utopia Planitia, and implications for
854	climate history on Mars. Journal of Geophysical Research 112, E06010, doi.10.1029/
855	2006JE002869.
856	Müller, F. 1963. Observations on Pingos. National Research Council of Canada, Ottawa, Canada,
857	Technical Translation 1073, pp, 119.
858	Murton, J.B. 2001. Thermokarst sediments and sedimentary structures, Tuktoyaktuk Coastlands,
859	western Arctic Canada. Global and Planetary Change 28, 175-192.
860	Murton, J.B., Worsley, P., Gozdzik, J. 2000. Sand veins and wedges in cold aeolian environments.
861	Quaternary Science Reviews, 19, 899-922.

- Mustard, J.F., Cooper, C.D., Rifkin, M.R. 2001. Evidence for recent climate change on Mars
  from the identification of youthful near-surface ground ice. Nature 412, 411-414, doi.10.
  1038/35086515.
- 0'Brien, R. 1971. Observations on pingos and permafrost hydrology in Schuchert Dal,
- 866 Northeast Greenland. Meddelelser om Grønland 195, 1, 20 pp.
- 867 Oehler, D., Mangold, N., Hallet, B., Fairén, .G., Le Deit, L., Williams, A.J., Sletten, R.S.,

Martínez-Frías, J. 2016. Origin and significance of decameter-scale polygons in the lower
Peace Vallis fan of Gale crater, Mars. Icarus 277, 56-72,
doi.org/10.1016/j.icarus.2016.04.038.

- 871 Osterkamp, T.E., Jorgenson, M.T., Schuur, E.A.G., Shur, Y.L., Kanevskiy, M.A., Vogel, J.G.,
- 872 Tumskov, V.E. 2009. Physical and Ecological Changes Associated with Warming
- 873 Permafrost and Thermokarst in Interior Alaska. Permafrost and Periglacial Processes 20,
- 874 235, 256, doi:10.1002/ppp656.

875	Page, D.P., Murray, J.B. 2006. Stratigraphical and morphological evidence for pingo genesis in
876	the Cerberus plains. Icarus 183, 46–54, doi.10.1016/j.icarus.2006.01.017.

- Påsse, T. 2004. The amount of glacial erosion of the bedrock. Sveriges Geologiska Undersökning.
  Technical Report TR-04-25, 39 p.
- Pechmann, J.C. 1980. The origin of polygonal troughs on the northern plains of Mars. Icarus 42,
  185-210.
- Penner, E. 1959. The mechanism of frost heaving in soils. Highway Research Board, Bulletin
  225, 1-22.
- Pettapiece, W.W. 1974. A hummocky permafrost soil from the subarctic of northwestern Canada
  and some influences of fire. Canadian Journal of Soil Science 54, 4, 343-355.
- Péwé, T. 1959. Sand-wedge polygons (tessellations) in the McMurdo Sound region, Antarcticaa progress report. American Journal of Science 257, 545-552.
- Phillips, W.M., Hall, A.M., Mottram, R., Fifield, L.K., Sugden, D.E. 2006. Cosmogenic <sup>10</sup>Be
- and <sup>26</sup>Al exposure ages of tors and erratics, Cairngorm Mountains, Scotland: Timescales
- for the development of a classic landscape of selective linear glacial erosion.
  Geomorphology 73, 222-245, doi:10.1016/j.geomorph.2005.06.009.
- Pollard, W.H., French, H.M. 1985. The internal structure and ice crystallography of seasonal frost
  mounds. Journal of Glaciology 31, 108, 157-162.
- 893 Rampton, V.N. 1973. The influence of ground ice and thermokarst upon the geomorphology of
- the Mackenzie Beaufort region, in Research in polar and alpine geomorphology,
- 895 Proceedings, 3<sup>rd</sup> Guelph Symposium on Geomorphology, edited by B.D. Fahey and R.D.
- Thompson, Geo. Abstracts, Norwich, UK, 43-59.
- 897 Rampton, V.N. 1988. Quaternary geology of the Tuktoyaktuk Coastlands, Northwest Territories,

898

Geological Survey of Canada, Memoir 423, 98 p.

- 899 Rampton, V.N. Mackay, J.R. 1971. Massive ice and icy sediments throughout the Tuktoyaktuk
- 900 Peninsula, Richards Island, and nearby areas, District of Mackenzie, Geological Survey of
- 901 Canada, Paper 71-21, 16 p.
- Rampton, V.N., Bouchard, M. 1975. Surficial geology of Tuktoyaktuk, District of Mackenzie,
  Geological Survey of Canada, Paper 74-53, 16 p.
- 904 Reznichenko, N.V., Davies, T.R., Winkler, S. 2016. Revised palaeoclimatic significance of
- Mueller Glacier moraines, Southern Alps, New Zealand. Earth Surface Processes and
  Landforms 41, 2, 196-207, doi.org/10.1002/esp.3848.
- 907 Roberson, S., Hubbard, B., Coulson, H., Boomer, I. 2001. Physical properties and formation of
- 908 flutes at a polythermal valley glacier: Midre Lovénbreen, Svalbard. Geografika Annaler:
  909 Series A, Physical Geography, 93, 71-88. doi:10.1111/j.1468-0459.2011.00420.x.
- 910 Schirrmeister, L. Froese, D., Tumskoy, V., Grosse, G., Wetterich, S. 2013. Yedoma: late
- Pleistocene ice-rich syngenetic permafrost of Beringia. eds. S.A. Elias & C.J. Mock,
  Encyclopedia of Quaternary Science, 2<sup>nd</sup> ed., 3, 542-552.
- Scholz, H., Baumann, M. 1997. An "open system" pingo near Kangerlussuaq (Søndre Strømfjord),
  West Greenland, Geology of Greenland Bulletin 104-108.
- Schon, S.C., Head, J.W., Fassett, C.I. 2012. Recent high-latitude resurfacing by a climate-related
  latitude-dependent mantle: Constraining age of emplacement from counts of small craters.
- Planetary and Space Science 69, 49-61, doi.org/10.1016/j.pss.2012.03.015.
- Schorghofer, N., Forget, F. 2012. History and anatomy of subsurface ice on Mars. Icarus 220,
- 919 1112-1120, doi.10.1016/j.icarus.2012.07.003.
- 920 Seibert, N.M., Kargel, J.S. 2001. Small scale martian polygonal terrain: implications for liquid

- 921 surface water. Geophysical Research Letters 28, 5, 899-902.
- 922 Séjourné, A., Costard, F., Gargani, J., Soare, R.J., Fedorov, A., Marmo, C. 2011. Scalloped
- depressions and small-sized polygons in western Utopia Planitia: a new formation
  hypothesis. Planetary and Space Science 59, 412-422, doi:10.1016/j.pss.2011.01.007.
- 925 Séjourné, A., Costard, F., Gargani, J., Soare, R.J., Marmo, C. 2012. Evidence of an eolian ice-rich
- and stratified permafrost in Utopia Planitia, Mars. Planetary and Space Science 60, 248254, doi:10.1016/j.pss.2011.09.004.
- 928 Shakesby, R.A., 1989. Variability in Neoglacial moraine morphology and composition, Storbreen,
- Jotunheimen, Norway: within-moraine patterns and their implications. Geografiska
  Annaler: Series A, Physical Geography 71, 1-2, 17-29.
- Sinha, R.K., Murty, S.V.S. 2015. Amazonian modification of Moreux crater: Record of recent
  and episodic glaciation in the Protonilus Mensae region of Mars. Icarus 245, 122-144,
- 933 doi.org/10.1016/j.icarus.2014.09.028.
- 934 Skinner, J.A., Tanaka, K.L., Platz, T. 2012. Widespread loess-like deposit in the Martian northern
  935 lowlands identifies Middle Amazonian climate change. Geology 40, 12, 1127-1130,
  936 doi:10.1130/G33513.1
- 937 Sletten, R.S., Hallet, B., Fletcher, R.C. 2003. Resurfacing time of terrestrial surfaces by the
- formation and maturation of polygonal patterned ground. Journal of Geophysical Research
  108, E4, 8044, dx.doi.org/10.1029/2002JE001914.
- 940 Smalley, I., Marshall, J., Fitzsimmons, K., Whalley, W.B., Ngambi, S. 2019. Desert loess: a
- 941 selection of relevant topics. Geologos 25, 1, 191-102, doi.10.2478/logos-2019-0007.
- 942 Soare, R.J., Kargel, J.S., Osinski, G.R., Costard, F. 2007. Thermokarst processes and the origin
- 943 of crater-rim gullies in Utopia and western Elysium Planitia. Icarus 191, 1, 95-112, doi.

- 944 org/10.1016/j.icarus.2007.04.018.
- Soare, R.J., Osinski, G.R., Roehm, C.L. 2008. Thermokarst lakes and ponds on Mars in the very
  recent (late Amazonian) past. Earth and Planetary Science Letters 272, 1-2, 382-393, doi.
  10.1016/j.epsl.2008.05.10.
- 948 Soare, R.J., Conway, S.J, Pearce, G., Dohm, J.M., Grindrod, P.M. 2013. Possible crater-based
- 949 pingos, paleolakes and periglacial landscapes in the high latitudes of Utopia Planitia, Mars.
  950 Icarus, 225, 2, 971-981, doi.org/10.1016/j.icarus.2012.08.041.
- 951 Soare, R.J., Conway, S.J., Dohm, J.M., El-Maarry, M.R. 2014. Possible open-system pingos in
- and around the Argyre impact basin, Mars. Earth and Planetary Science Letters 398, 2536, doi.10. 1016/j.epsl.2014.04.044.
- Soare, R.J., Conway, S.J., Williams, J-P., Gallagher, C., McKeown, L.E. 2020. Possible (closed
  system) pingo-thermokarst complexes at the mid-latitudes of Utopia Planitia, Mars. Icarus,
  doi.org/10.1016/j.icarus.2019.03.010.
- 957 Soare, R.J., Williams, J-P, Conway, S.J., El-Maarry, M.R. 2021. Pingo-like mounds and possible
- polyphase periglaciation/glaciation at/adjacent to the Moreux impact crater, in MarsGeological Enigmas, from the deep past through to the present. Elsevier, in press.
- Souness, C.J., Hubbard, B., Milliken, R.E., Quincey, D. 2012. An inventory and population-scale
  analysis of martian glacier-like forms. Icarus 217, 243-255, doi.10.1016/j.icarus.2011.10.
  020.
- Souness, C.J., Hubbard, B. 2013. An alternative interpretation of late Amazonian ice flow:
- 964 Protonilus Mensae, Mars. Icarus 225, 495-505, doi.org/10.1016/j.icarus.2013.03.030.
- 965 Storrar, R., Stokes, C.R. 2007. A Glacial Geomorphological Map of Victoria Island, Canadian
- 966 Arctic, Journal of Maps, 3, 1, 191-210, doi:10.1080/jom.2007.9710838

967	Sugden, D.E., John, B.S. 1976. Glaciers and landscape: a geomorphological approach. E. Arnold.
968	Summmerfield, M.A. 1999. Global geomorphology. Harlow, England, Pearson Education.
969	Taber, S. 1930. The mechanics of frost heaving, 9-26, in Historical perspectives in frost heave
970	research: the early works of S. Taber and G. Beskow. Special report 91-23, U.S. Army
971	Corp of Engineers, (eds.) P.B Black and M.J. Hardenburg, 1991, 159 p.
972	Tanaka, K.L., Skinner, J.A., Hare, T.M. 2005. Geological map of the northern plains of Mars, map
973	2888. Flagstaff, USGS.
974	Ulrich, M., Morgenstern, A., Günther, F., Reiss, D., Bauch, K.E., Hauber, E., Rössler, S,
975	Schirrmeister, L. 2010. Thermokarst in Siberian ice-rich permafrost: comparison to
976	asymmetric scalloped depressions on Mars. Journal of Geophysical Research 115, E10009,
977	doi:10.1029/2010JE003640.
978	van der Meer, J.J., Menzies, J., Rose, J. 2003. Subglacial till: the deforming glacier bed.
979	Quaternary Science Reviews, 22, 15-17, 1659-1685, doi:10.1016/S0277-3791(03)00141-
980	0.
981	van Everdingen, R.O. 1982. Frost Blisters of the Bear Rock Spring Area near Fort Norman, NWT.
982	Arctic 35,2 243-265.
983	Washburn, A.L. 1973. Periglacial Processes and Environment. St Martin's Press, New York, NY,
984	p. 320.
985	Williams, JP., van der Bogert, C.H., Pathare, A.V., Michael, G.G., Kirchoff, M.R., Hiesinger, H.
986	2018. Dating very young planetary surfaces from crater statistics: A review of issues and
987	challenges. Meteoritics and Planetary Science 53, 4, 554-582, doi.org/10.1111/maps.
988	12924.
989	Witkind, I.J. 1978. Giant glacial grooves at the north end of the mission range, northwest Montana.

990

Journal of Research, U.S. Geological Survey 6, 4, 425-433.

- 991 Wolfe, S.A., Morse, P.D., Neudorf, C.M., Kokelj, S.V., Lian, O.B., O'Neill, B. 2018.
- 992 Contemporary sand-wedge development in seasonally frozen ground and
  993 paleoenvironmental implications. Geomorphology 215-229, doi.org/10.1016/j.geomorph.
  994 2018.02.015.
- 995 Worsley, P., Gurney, S.D. 1996. Geomorphology and hydrogeological significance of the
- Holocene pingos in the Karup Valley area, Traill Island, northern east Greenland. Journalof Quaternary Science 11, 3, 249-262.
- 998 Yoshikawa, K., Harada, K. 1995. Observations on nearshore pingo growth, Adventalen,
- 999 Spitsbergen. Permafrost and Periglacial Processes, 6, 4, 361-372.
- 1000 Yoshikawa, K., Nakamura, T., Igarashi, Y. 1996. Growth and Collapse History of Pingos,

1001 Kuganguaq, Disko Island, Greenland. Polarforschung 64, 3, 109-113.

1002 Figures

Fig. 1. (a) Geographical context and topographical location of the study site north of the Moreux 1003 1004 impact-crater, Mars. Top panel shows a colour-keyed digital elevation model using Mars Orbiter Laser Altimeter (MOLA) data overlain on the Thermal Emission Imaging System 1005 1006 (THEMIS) daytime IR controlled-image mosaic. The inset gives the location of the map within the global view of Mars rendered as a hill-shaded relief version of the MOLA data. 1007 (b) The study area in the *THEMIS* daytime *IR* controlled-image mosaic. The two (red) 1008 1009 boxes showing the location of the *HiRISE* images used within the fretted terrain north of Moreux crater's rim. (c-d) All of the figures and their footprints framed within Mars 1010 Reconnaissance Orbiter Context Camera (CTX) image J01\_045349\_2239\_XN\_43N316W. 1011 1012 Image credit: Malin Space Science Systems.

1013 Fig. 2. (a) Context image of Type-1 polygon distribution in one of the two basins studied by us.

1014 Note the overprinting of ridge and trough terrain (bottom left hand corner) and of possible moraines (top right-hand corner) located respectively at the southwestern and northeastern 1015 1016 margins of the basin. (b) Random example of high-centred Type-1 polygons. (c-d) Example of polygonised terrain whose relative elevation and morphology are sufficiently 1017 similar to the terrain highlighted in panel (b), despite lying amidst landscape features that 1018 could be glacial in origin, for a (roughly) similar age to be ascribed (HiRISE image 1019  $43.405^{\circ}$ N:  $44.108^{\circ}$ ESP 045349 2235. E. 25 cm/pixel). Image 1020 credit: 1021 NASA/JPL/University of Arizona.

Fig. 3. (a) Etched polygons incise terrain fragments and (b) isolated mounds which, antecedently,
 could have been contiguous with icy coverage of the surrounding basin (*HiRISE* image
 ESP\_045349\_2235, 43.405<sup>0</sup> N; 44.108<sup>0</sup> E, 25 cm/pixel). Image credit:
 *NASA/JPL/*University of Arizona.

## Fig. 4. Basin-covering scarps, possibly thermokarstic in origin, with equator-ward orientation. The loss of elevation trends to the south (*CTX* image J01\_045349\_2239\_XN\_43N316W, 43.94<sup>0</sup> N: 43.99<sup>0</sup> E, 5.94 m/pixel). Image credit: Malin Space Science Systems.

Fig. 5. (a) Context image of undulating, polygonised terrain, comprised of high-centred polygons.
 Note the longitudinal trend and the sub-parallel alignment of the undulations (b)
 Magnification of same. (c) Arc-like distribution of circular and sub-circular depressions
 within the basin-surface polygons (*HiRISE* image ESP\_042105\_2235, 43.163<sup>0</sup> N; 43.983<sup>0</sup>
 E, 50 cm/pixel). Image credit: NASA/JPL/University of Arizona.

**Fig. 6.** Landscape-scale image (*CTX* J01\_045349\_2239\_XN, 43.94<sup>0</sup> N; 43.99<sup>0</sup> E, 5.94 m/pixel) of

1035our study region. Black rectangles identify the surface coverage of the two *HiRISE* images1036highlighted by us therein. Black arrows point to the areas of the basin where the dense1037distribution of ridge and trough terrain begins to dissipate. Image credit: Malin Space1038Science Systems. Image credit: Malin Space Science Systems.

- Fig. 7. (a) Partial basin-coverage by ridge and trough terrain. Note the dense distribution of etched 1039 terrain and mounds in the latter's midst. (b) Eastward transition of ridge and trough 1040 distribution: dense and continuous on the left; discontinuous, with less amplitude and mass, 1041 on the centre-left; and, possible remnants to the right, top corner (also see Fig. 8d). (c) 1042 1043 Overprinting of the basin-surface polygons by the ridges and troughs (*HiRISE* image ESP 045349 2235,  $43.405^{\circ}$ N:  $44.108^{\circ}$ E. 25 cm/pixel). 1044 Image credit: NASA/JPL/University of Arizona. 1045
- Fig. 8. Mounds and etched (Type-2) polygons. (a-c) Clustered distribution of mounds amidst the 1046 ridge and trough terrain. Note the truncation/erosion and dissection of the mounds by the 1047 ridge and trough structures, as well as the mound incision by the **Type-2** polygons. (d) 1048 1049 Density of ridge/trough distribution decreases with distance from the western flank of the basin; note outlying fragment (see Fig. 9 for context) at the far right of this tile. 1050 Collectively, these observations hint at the possibility that the basin coverage by the 1051 ridge/trough terrain could have been much more extensive than is apparent today (HiRISE 1052 image ESP\_045349\_2235, 43.405<sup>0</sup> N; 44.108<sup>0</sup> E, 25 cm/pixel). Image credit: 1053 1054 NASA/JPL/University of Arizona.

1055 Fig. 9. Kilometre-scale, longitudinal distribution of serialized raised-ridges (possible moraines),

1056 with kilometres of reach onto the basin surface at some locations (black rectangle). Note1057 the horseshoe-shaped and etched mounds as well as the small patch of etched terrain within

the rectangle, possibly outlying remnants of the etched terrain to the west (*HiRISE* image
ESP 045349\_2235, 43.405<sup>0</sup> N; 44.108<sup>0</sup> E, 25 cm/pixel). Image credit: *NASA/JPL/*University of Arizona.

1061 Fig. 10. (a) ~Nine serialized ridges with varying degrees of amplitude and mass. Note the increased density of boulders and bouldery terrain with proximity to the basin's edge and the rockier 1062 elevated-terrain on the right flank of the image. (b) Small-scale lineations/grooves with 1063 adjacent (aligned) boulders (black arrows), consistent with presumed flow of glacier. 1064 Suggestive of Earth-like flutes. (c) Possible boulder trains (black rectangles). (d) Larger-1065 1066 scale view of lineated train, possibly comprised of glacial flutes and/or boulder trains, linking a possible rocky source upslope with boulder-strewn fields of polygons downslope. 1067 (*HiRISE* image ESP 045349 2235, 43.405<sup>0</sup> N; 44.108<sup>0</sup> E, 25 cm/pixel). Image credit: 1068 NASA/JPL/University of Arizona. 1069

Fig. 11. Apparent push-contact and interception of the ridge and trough terrain by moraine-like
ridges (white arrows highlight contact; hollow black arrow points to one of the lobate
structures). Note the loss of ridge mass and amplitude where there is no contact (top-left
of figure, above the white arrows) (*CTX* image J01\_045349\_2239\_XN, 43.94<sup>o</sup> N; 43.99<sup>o</sup>
E, 5.94 m/pixel). Image credit: Malin Space Science Systems.

Fig. 12. Polygonal terrain in the Tuktoyaktuk Coastlands, Northwest Territories, Canada. (a) Icewedge polygons (~10-20 m in diameter) with meltwater-filled marginal troughs slightly
lower in elevation than the polygon centers. The high-centred morphology of the polygons
is the result of ice-wedge degradation on the margins. Photo credits: R. Soare. (b-c).
Incision of ice wedge by sand wedge and vice versa (Murton et al., 2000). (d) Panoramic
and cross-sectional view of ice wedges (white arrows) in the midst of massive-ice outcrops.

1081 The small, surface depressions above the ice-wedge in (c) and the ice-wedge to the right, 1082 here, mark the location of bilateral polygonal troughs associated with a field of high-1083 centered polygons. The depressions, along with the hummocky terrain above and 1084 surrounding the massive-ice, comprise thermokarst. Photo credit: R. Soare.

1085 Fig. 13. Closed-system pingos in the Tuktoyaktuk Coastlands. (a) Thermokarst-lake drainage in 1086 ice-rich, polygonised terrain and a nascent CSP (plan-view) in the midst. (b) Adolescent *CSP* developing adjacent to the village of Tuktovaktvuk ( $69.4454^{\circ}$  N,  $133.0342^{\circ}$  W). (c) 1087 Ibyuk Pingo (late winter), thought to ~1300 years old (+/-200 years) (Mackay, 1986), as 1088 seen from the summit of Split Pingo. Ibyuk Pingo stands at ~49m above sea-level and is 1089 1090 the 2nd highest pingo on Earth. Note the radial fracturing of Ibyuk Pingo as well as the irregular topography of the summits at both Split and Ibyuk Pingos. Photo credits: R. Soare. 1091 1092 (d) Metre-scale (elevation) pingo rampart, the end-stage of pingo evolution. Photo credit: D.M. Burr. 1093

Fig. 14. Open-system pingos in Greenland and Canada. (a) Two open-system pingos (~5-10 m 1094 1095 high; 20-50 m diameters) on glacial-valley floor, Mellemfjord, Disko, Central West Greenland. Braided glacio-fluvial river system dissects the valley Photo credit: 1096 1097 Christiansen, 1995. Left-most mound displays irregularly-shaped summit depression. (b) Partially-collapsed, open-system pingo in outwash plain underlain by continuous albeit 1098 thin permafrost, Niohalvfjerdsfjorden, eastern North Greenland. Photo credit: Ole Bennike, 1099 1100 GEUS. (c) View of Müller ice-cap terminus from summit of open-system pingo, Axel Heiberg Island, Nunavut, Canada. Small-sized thermal-contraction polygons incise terrain 1101 1102 separating the pingo from ice-cap terminus (not clearly identifiable at this oblique angle).

(d) Summit of Axel Heiberg pingo; note seasonal icings (off-centred from the summit apex) and glacio-fluvial river system (to left of mound). Photo credits, c-d: R. Soare.

Fig. 15. Landforms and deposition associated with glaciation on Earth. (a) Aerial view of Mueller 1105 1106 Glacier, New Zealand, highlighted by pro-glacial lake and (marginal) moraines. White arrows indicate the crests of a large 140 m high Little Ice-Age latero-frontal and that of a 1107 1108 smaller outermost terminal moraine. (b) Basal till deposited during the younger Dryas glaciation, c. 18,000 years ago, on the north shore of Loch Torridon, Scotland. Note the 1109 angular and poorly-sorted mixture of boulders and cobbles. Photo credit: M.J. Hambrey; 1110 1111 reproduced with permission of Glaciers Online (https://www.swisseduc.ch/glaciers/). (c) Recessional-moraine assemblage at the terminus of Mueller glacier; moraines indicated by 1112 white arrows in a) are marked by orange lines. At least three moraines in sequence are 1113 identifiable. Aerial imagery in c) from the *LINZ* database (https://data.linz.govt.nz/). 1114

1115 Fig. 16. Parallel scratches in bedrock are glacial striations (western Manitoulin Island, Ontario,

1116 Canada). Scale provided by a Canadian dollar coin ("Loonie") in the centre of the photo.

- 1117 Location: Mississagi Lighthouse Campground, Photo credit: A. Fyon (<u>https://www.</u>
- 1118 <u>ontariobeneathourfeet.com/glacial-striation</u>).

Fig. 17. (a) Glacial flute in Iceland, 75 m from the margin of Múlajökull glacier. The flute has
three boulders at its head, all aligned with the flute's long axis. The farthest up-glacier
boulder is ~0.37 m tall. This flute is 20 m long, 0.21 m high and 0.86 m wide nearest its
boulders. (b) Up-glacier view of multiple flutes. Meltwater flows in the depressed areas on
either side of the flutes. Photo credits: L. Ives.

1124 Fig. 18. (a) Clach Bun Rudhtair, eastern Cairngorms mountains, Scotland: The summit of the

upper tower has been removed by ice flowing in the direction of the lineated train of
boulders; the summit of the third tower is unmodified and carries weathering pits over 1 m
deep (Hall and Phillips, 2006). (b) Boulder train highlighted by aligned and closely-set
members, cascading clast-size lee of the largest boulder and in the direction of glacial flow.
Photo credit, V. Bjarnadóttir.

1130 Fig. 19. Glacial grooves, vestigial to and carved by the Laurentide Ice sheet Glacial Grooves

State Monument, Kelley's Island, Ohio. Photo credit: Hill/National Snow and Ice Data
Center, University of Colorado, Boulder (https://nsidc.org/cryosphere/glaciers/
gallery/grooves.html).

Fig. 20. (a) CTX image J01\_045349\_2239\_XN, craters marked with red circles. The black boxes 1134 are the locations of the two *HiRISE* images in (b) ESP\_045349\_2235 and (c) 1135 ESP 042105 2235 with count areas outlined in black with color shading. Count areas 1 1136 and 3 represent polygonised terrain and count area 2 is comprised of ridge and trough 1137 terrain. North is up and illumination is from the west. (d) Cumulative and (e) differential 1138 1139 *CSFDs* of counts with model isochrons of Hartmann (2005) (gray curves) for 1 Ma, 10 Ma, 100 Ma, 500 Ma, and 3 Ga and the model-age estimate of  $1.5\pm0.2$  Ga for the CTX counts 1140 1141 derived by Poisson timing analysis (Michael et al., 2016). CSFDs extend across model isochrons >10 Ma suggesting terrains are older than 10 Ma. The crater diameters extend to 1142 larger sizes in areas 1 and 3 consistent with the polygonised terrain being older than the 1143 1144 ridge and trough terrain. Image credit: Malin Space Science Systems, NASA/JPL/University of Arizona. 1145

1146











































□ 10 km

□ 2 km



Red 'X' are bowl shaped craters