

Ultrasound-induced microbubble coalescence by parametric instability

Michiel Postema, Philippe Marmottant, Charles T Lancée, Sascha Hilgenfeldt, Nico de Jong

► To cite this version:

Michiel Postema, Philippe Marmottant, Charles T Lancée, Sascha Hilgenfeldt, Nico de Jong. Ultrasound-induced microbubble coalescence by parametric instability. Proceedings of the 10th Dutch Annual Conference on Biomedical Engineering, 2003, Enschede, Netherlands. pp.177, 10.5281/zen-odo.4841069. hal-03189256

HAL Id: hal-03189256 https://hal.science/hal-03189256

Submitted on 25 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.







Ultrasound-induced microbubble coalescence by

parametric instability

Michiel Postema,^{1,2*} Philippe Marmottant,³ Charles T. Lancée,¹ Sascha Hilgenfeldt,³ Nico de Jong^{1,2,3}

¹Department of Experimental Echocardiography, Thoraxcentre, Erasmus MC, Rotterdam;

²Interuniversity Cardiology Institute of the Netherlands (ICIN), Utrecht;

³Physics of Fluids, Department of Applied Physics, Faculty of Science and Technology, University of Twente, Enschede.

Micrometer-sized gas bubbles in a liquid will oscillate upon insonification by ultrasound. During the expansion phase, bubble coalescence is frequently observed. We give a description of the microbubble coalescence mechanism, based on high-speed optical observations and theoretical modeling.

Ultrasound-induced microbubble coalescence is the fusion of two or more microbubbles. As adjacent bubbles expand, the following stages can be distinctly observed: flattening of the bubble surfaces prior to contact, drainage of the interposed liquid film toward a critical thickness, rupture of the liquid film, and formation of a single bubble. Locally the critical thickness is more quickly reached than might be explained from film drainage alone, due to thickness perturbations induced by secondary radiation forces and amplified by a parametric instability. The coalescence process is thus sped up to durations of less than 1 μ s for ultrasound contrast agent microbubbles in the micron diameter range.





* To whom correspondence should be addressed. E-mail: m.postema@erasmusmc.n