



HAL
open science

Removal of Metal–Metal Bonding in a Dimetallic Paddlewheel Complex: Molecular and Electronic Structure of Bis(phenyl) Dirhodium(III) Carboxamidate Compounds

Joffrey Wolf, Rinaldo Poli, Jian-Hua Xie, Jason Nichols, Bin Xi, Peter Zavalij, Michael P Doyle

► To cite this version:

Joffrey Wolf, Rinaldo Poli, Jian-Hua Xie, Jason Nichols, Bin Xi, et al.. Removal of Metal–Metal Bonding in a Dimetallic Paddlewheel Complex: Molecular and Electronic Structure of Bis(phenyl) Dirhodium(III) Carboxamidate Compounds. *Organometallics*, 2008, 27 (22), pp.5836-5845. 10.1021/om800631b . hal-03191444

HAL Id: hal-03191444

<https://hal.science/hal-03191444v1>

Submitted on 7 Apr 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Removal of Metal-Metal Bonding in a Dimetallic Paddlewheel Complex – Molecular and Electronic Structure of Bis(phenyl) Dirhodium(III) Carboxamidate Compounds

Joffrey Wolf,[†] Rinaldo Poli,^{†§} Jian-Hua Xie,[‡] Jason Nichols,[‡] Bin Xi,⁺ Peter Zavalij[‡] and Michael P.
Doyle^{‡*}*

Department of Chemistry and Biochemistry, University of Maryland, College Park, MD 20742 USA;

Laboratoire de Chimie de Coordination, UPR CNRS 8241 liée par convention à l'Université Paul Sabatier et à l'Institut National Polytechnique de Toulouse, 205 route de Narbonne, 31077 Toulouse Cedex 4, France; and Institut Universitaire de France, 103, bd Saint-Michel, 75005 Paris, France; and

Department of Chemistry, Purdue University, West Lafayette, Indiana 47907 USA

[†] Laboratoire de Chimie de Coordination, UPR CNRS 8241 liée par convention à l'Université Paul Sabatier et à l'Institut National Polytechnique de Toulouse, 205 route de Narbonne, 31077 Toulouse Cedex 4, France. [§] Institut Universitaire de France, 103, bd Saint-Michel, 75005 Paris, France. [‡] Department of Chemistry and Biochemistry, University of Maryland, College Park, MD 20742 USA. ⁺ Department of Chemistry, Purdue University, West Lafayette, Indiana 47907.


rinaldo.poli@lcc-toulouse.fr

RECEIVED DATE (to be automatically inserted after your manuscript is accepted if required according to the journal that you are submitting your paper to)

Abstract : The unique structural features and chemical stabilities of bis(phenyl) tetracarboxamidatodirhodium(III) are reported, and their electronic structures are mapped through XPS, electrochemical, and computational methods. Comparison with the structures of dirhodium(II,II) and dirhodium(II,III) oxidative precursors portrays the diphenyl dirhodium(III) compounds as two square-pyramidal rhodium units that have undergone conrotatory motion in order to optimize metal-ligand bonding. Axial phenyl ligands are severely distorted from their expected Rh-Rh-C linear array. XPS data for this series of dirhodium compounds are consistent with the absence of a rhodium-rhodium bond for the diphenyl dirhodium(III) compounds, and electrochemical measurement shows a single reversible $\text{Rh}_2^{6+}/\text{Rh}_2^{7+}$ redox couple. Notably, they exhibit high thermal stability, and Brønsted acid removal of carboxamidate ligands precedes the formation of benzene. The ability of a phenyl group to impart unusual stability to rhodium(III) compounds is explained by theoretical analysis of the electronic structure of bis- σ -(phenyl)-tetrakis- μ -(carboxamido)dirhodium(III), and comparison is made with previously reported dinitrosyl dirhodium(III) complexes. Formally, two phenyl radicals in combination with a rhodium-rhodium bond are transformed into two phenyl-rhodium bonds. The severe distortion of phenyl rings from linearity is suggested to result from a long-range interaction between the electron-deficient rhodium and distal oxygen atoms.

Introduction

The recent definitive treatise on multiple bonds between metal atoms introduced oxidation states with the statement that “the most common oxidation states for the M_2^{n+} units in paddlewheel complexes corresponds to values of n of 4, 5, and 6”.¹ Examples of the $n = 6$ systems are less common. Our report last year of remarkably stable bis(phenyl)dirhodium(III) caprolactamate (Scheme 1, left)² was only the second example for Rh_2^{6+} compounds, $Rh_2(O_2CR)_4(NO)_4$ ($R = Me, Et, iPr$) (Scheme 1, center) having been previously reported by the Lippard group exhibiting the same structural type and an essentially identical Rh-Rh separation.³ Although no specific oxidation state assignment was mentioned in the Lippard report, the bent geometry observed for the NO ligands justifies the assignment of a formal oxidation state of III to the Rh centers, and this was later supported by computational studies.⁴⁻⁶ One of the surprising features of these Rh_2^{6+} compounds is the apparent absence of a rhodium-rhodium bond. In contrast, a paramagnetic paddlewheel Ir_2^{6+} compound, $Ir_2(hpp)_4Cl_2$ (Scheme 1, right), also from group 9, has been reported to have a doubly-bonded dimetallic core with two unpaired electrons.⁷ Isoelectronic Ru_2^{4+} complexes are reported to have the $\sigma^2\pi^4\delta^2\delta^*2\pi^*2$ electronic configuration^{1, 8} implying a Ru-Ru double bond and a spin triplet ground state, whereas Ru_2^{6+} compounds have the $\pi^4\delta^2\pi^*4$ electronic configuration with a net δ single bond.^{1, 9, 10}




Scheme 1. Comparison of two isoelectronic Group 9 paddlewheel dimers.

Homo-dinuclear σ -alkynyl complexes of both ruthenium and rhodium are known,¹¹ although with rhodium only in the (II,III)-oxidation state.¹² Electron donation from the axial carbon ligand to rhodium increases the Rh-Rh bond distance, but only a fraction of that found with **1**, and their electronic properties imply retention of a metal-metal bond.

The ability of paddlewheel complexes to hold together two metals without a metal-metal bond is well-established,^{13, 14} and the structural and electronic influences that result have substantial theoretical and practical implications.¹⁵ In this contribution, we report the molecular structures and chemical stabilities of the diphenyl tetracarboxamidatodirrhodium(III) system and their study by XPS spectroscopy, electrochemistry, and by computational methods.

Results and Discussion

Synthesis and Characterization. In order to identify the unique structural features of bis(phenyl)dirrhodium(III) carboxamidates we have prepared the dirrhodium(II) pyrrolidinate (pyr), valerolactamate (val), and caprolactamate (cap) bis-acetonitrile complexes (**1**),¹⁶ oxidized these compounds to the corresponding paramagnetic Rh(II)Rh(III) derivatives (**2**, eq 1), and also prepared their diamagnetic bis(phenyl)dirrhodium(III) analogues (**3**, eq 2). The synthesis of dirrhodium(II,III) complexes was achieved by oxidation of the dirrhodium(II) carboxamidate using copper(II) triflate, nitrosonium salts (eq 1),^{17, 18} or with the use of phenyldiazonium tetrafluoroborate; oxidations with nitrosonium salts were the most versatile and gave dirrhodium(II,III) product without byproducts in solution. As mentioned in the Introduction, the synthesis (using sodium tetraphenylborate) and characterization of bis(phenyl)dirrhodium(III) caprolactamate (**3c**) has been described in a preliminary communication.² Since this first report, a new methodology for convenient access to diaryl analogues directly from dirrhodium(II) carboxamidates and arylboronic acids catalyzed by copper sulfate (eq 2) has been communicated,¹⁹ and the spectral data of these products have been described.



The crystal structures of all nine compounds have been obtained. Those of the pyrrolidinate systems $\text{Rh}_2(\text{pyr})_4(\text{MeCN})_2$ (**1a**), $[\text{Rh}_2(\text{pyr})_4(\text{PhCN})_2]\text{BF}_4$ (**2a'**), and $\text{PhRh}(\text{pyr})_4\text{RhPh}$ (**3a**) are presented in Figure 1. Unpublished data for the other 6 compounds are provided as supporting information. A selection of bond lengths and angles for all nine compounds is reported in **Table 1** and **Table 2**. As can be seen from these data, the diphenyl-dirhodium(III) compounds exhibit severe structural distortions relative to their dirhodium(II,II) and dirhodium(II,III) analogues. They can best be evaluated by comparing a few key bond angles [the values are averages over the three structures within each family and listed in the order **1, 2, 3**]: $L_{\text{axial}}\text{-Rh-Rh}$ [$175(2)$, $174(4)$, $159(4)^\circ$]; Rh-O-C [$118(2)$, $118(2)$, $126(1)^\circ$]; Rh-N-C [$122(1)$, $122(1)$, $118(1)^\circ$]. These distortions take the form of a severe bending of the Rh-Rh-Ph bond axis from 180° , a major opening of the Rh-O-C angle and a less pronounced closing of the Rh-N-C angle from the ideal 120° . The positioning of the rhodium units in the diphenyl-dirhodium structures appears as if two square-pyramidal units have undergone conrotatory motion so as to minimize repulsions and metal-metal interactions (Figure 2). In addition to these geometrical distortions, there is a significant lengthening of the Rh-Rh bond on going from **2** to **3**, whereas the bond shortens on going from **1** to **2**.




Figure 1. Structures of dirhodium pyrrolidates (a) **1a**, (b) **2a'**, and (c) **3a**. Ellipsoids are shown at the 30% probability level. Hydrogen atoms, BF_4^- (for **2a'**) and solvent molecules are omitted for clarity.

Table 1. Selected bond lengths of dirhodium carboxamidates in the (II,II), (II,III), and (III,III) oxidation states.

		Bond lengths of dirhodium complexes, Å		
	bond	1a	2a'^a	3a
From $\text{Rh}_2(\text{pyr})_4$	Rh1–Rh2	2.4540(5)	2.4432(4)	2.5738(12)
	Rh2–O1	2.077(2)	2.030(6)	2.093(6)
	Rh2–O2	2.080(2)	2.034(5)	2.095(5)
	Rh1–N1	2.003(3)	1.973(6)	1.993(7)
	Rh1–N2	2.011(3)	1.994(8)	2.012(6)
	Rh–L _{ax}	2.241(2)	2.248(8)	2.037(7)
	bond	1b	2b'^b	3b
From $\text{Rh}_2(\text{val})_4$	Rh1–Rh2	2.4187(5)	2.3872(3)	2.5162(5)
	Rh2–O1	2.061(2)	2.0143(16)	2.105(3)
	Rh2–O2	2.073(2)	2.0211(16)	2.061(2)
	Rh1–N1	2.023(3)	2.0078(19)	1.997(3)

	Rh1–N2	2.034(3)	2.016(2)	2.027(3)
	Rh–L _{ax}	2.254(3)	2.3071(16)	1.996(3)
	bond	1c	2c	3c
From Rh ₂ (cap) ₄	Rh1–Rh2	2.4221(4)	2.3840(6)	2.5188(3)
	Rh2–O1	2.050(2)	2.023(2)	2.0825(16)
	Rh2–O2	2.053(2)	2.030(2)	2.0777(16)
	Rh1–N1	2.038(3)	1.988(3)	2.012(2)
	Rh1–N2	2.045(3)	1.996(3)	2.0078(19)
	Rh–L _{axial}	2.336(3)	2.288(2)	2.010(2)

^a The bis-benzonitrile complex was prepared after failed attempts to prepare the bis-acetonitrile complex. ^b The bis-aquo complex was formed and crystallized, even though reaction was performed in acetonitrile.

Table 2. Selected bond angles for dirhodium carboxamidates in the (II,II), (II,III), and (III,III) oxidation states.

Bond angles of dirhodium complexes, °				
	angle	1a	2a'^a	3a
From Rh ₂ (pyr) ₄	Rh1–Rh2–O1	90.04(6)	89.66(16)	82.43(15)
	Rh1–Rh2–O2	90.01(6)	89.94(15)	83.87(14)
	Rh2–Rh1–N1	85.30(7)	85.9(2)	89.32(18)
	Rh2–Rh1–N2	85.35(7)	86.15(18)	90.61(18)
	Rh2–O1–C1	115.16(19)	115.8(5)	121.1(4)
	Rh2–O2–C2	115.9(2)	117.7(5)	122.4(5)
	Rh1–N1–C1	123.40(2)	122.4(5)	118.0(5)
	Rh1–N2–C2	124.00(2)	123.7(5)	118.5(5)
	Rh1–Rh2–L _{axial}	176.90(7)	178.22(16)	164.20(20)
	angle	1b	2b'^b	3b
From Rh ₂ (val) ₄	Rh1–Rh2–O1	88.03(7)	89.61(5)	74.58(7)
	Rh1–Rh2–O2	88.85(7)	89.66(4)	85.05(7)
	Rh2–Rh1–N1	87.08(8)	87.11(5)	99.10(8)

	Rh2–Rh1–N2	87.89(8)	87.23(5)	88.47(8)
	Rh2–O1–C1	118.8(2)	118.89(14)	133.0(2)
	Rh2–O2–C2	119.3(2)	119.08(14)	122.1(2)
	Rh1–N1–C1	121.5(2)	121.10(16)	121.8(4)
	Rh1–N2–C2	121.8(2)	122.07(16)	119.8(2)
	Rh1–Rh2–L _{axial}	173.84(8)	172.27(5)	157.34(11)
	angle	1c	2c	3c
From Rh ₂ (cap) ₄	Rh1–Rh2–O1	89.62(6)	88.58(7)	79.29(4)
	Rh1–Rh2–O2	89.37(6)	88.50(7)	78.39(5)
	Rh2–Rh1–N1	86.70(7)	88.42(8)	94.46(5)
	Rh2–Rh1–N2	86.81(7)	88.60(8)	95.32(5)
	Rh2–O1–C1	118.3(2)	119.8(2)	128.11(15)
	Rh2–O2–C2	118.96(19)	119.5(2)	128.97(15)
	Rh1–N1–C1	121.5(2)	120.8(2)	114.45(16)
	Rh1–N2–C2	121.7(2)	121.2(2)	113.51(15)
	Rh1–Rh2–L _{axial}	174.42(7)	171.28(7)	156.19(7)

^a The bis-benzonitrile complex was prepared after failed attempts to prepare the bis-acetonitrile complex. ^b The bis-aquo complex was formed and crystallized, even though reaction was performed in acetonitrile.




Figure 2. ORTEP views of (a) PhRh(pyr)₄RhPh (**3a**), (b) PhRh(val)₄RhPh (**3b**), and (c) PhRh(cap)₄RhPh (**3c**) showing comparable structural distortions. Ellipsoids are shown at the 30% probability level. Hydrogen atoms and solvent molecules are omitted for clarity. The corresponding Rh–Rh–C bond angles are 164.2° (**3a**), 157.3° (**3b**), and 156.2° (**3c**).

X-ray photoelectron spectroscopy (XPS) data for these nine compounds is given in Table 3. As evidenced by this direct measurement of the binding energy of *d*-electrons that constitute the dirhodium core, there is a significant change in binding energy upon oxidation of dirhodium(II,II) to dirhodium(II,III) (~ 0.8 eV) but an insignificant change upon oxidation of dirhodium(II,III) to dirhodium(III,III) (< 0.2 eV).

Table 3. XPS data for dirhodium carboxamidates in the (II,II), (II,III), and (III,III) oxidation states.^a

Compound	Rh-3d _{5/2} , eV	Rh-3d _{3/2} , eV
Rh ₂ (pyr) ₄ (MeCN) ₂ (1a)	308.000	312.563
Rh ₂ (val) ₄ (MeCN) ₂ (1b)	308.129	312.843
Rh ₂ (cap) ₄ (MeCN) ₂ (1c)	308.080	312.812
Rh ₂ (pyr) ₄ (PhCN) ₂ BF ₄ (2a')	308.905	313.514
Rh ₂ (val) ₄ (H ₂ O) ₂ BF ₄ (2b')	308.841	313.567
Rh ₂ (cap) ₄ (MeCN) ₂ BF ₄ (2c)	309.094	313.718
PhRh(pyr) ₄ RhPh (3a)	309.045	313.739
PhRh(val) ₄ RhPh (3b)	309.011	313.723
PhRh(cap) ₄ RhPh (3c)	309.105	313.725

^a Calibrated to carbon 1s at 284.6 eV.

The relative electron richness of the diphenyldirhodium(III) carboxamidate complexes is also suggested by the results of an electrochemical investigation. Cyclic (CV) and differential pulse voltammograms (DPV) of **3a**, **3b**, and **3c** were measured in CH₂Cl₂, and their half-wave potentials ($E_{1/2}$) are reported in Table 4, along with their electronic absorption maxima (λ_{max}). Figure 3 shows the CV and DPV traces for PhRh(pyr)₄RhPh (**3a**). For each of the complexes, a single, reversible Rh₂⁶⁺/Rh₂⁷⁺ redox couple (A) was observed. The oxidation potential of **3** increases with decreasing number of methylene units in the bridging cyclic amide, which follows that of the Rh₂^{4+/5+} couple of the parent complexes.²⁰ For complex **3a**, an additional irreversible redox couple (B) near the reduction limit of the

solvent was observed and is assigned to the $\text{Rh}_2^{5+/6+}$ redox couple. Attempts to isolate and characterize the oxidized $\text{PhRh}(\text{pyr})_4\text{RhPh}$ or any of its derivatives have thus far been unsuccessful.

Table 4. Electrode potentials (V) from CV measurements and λ_{max} of **3**.

Compound	A (V) ^a	B (V) ^b	λ_{max} (nm) ^c
3a	0.91	-1.36	442
3b	0.81	-	429
3c	0.78	-	430

^aReversible anodic wave reported as $E_{1/2}$, measured with platinum wire/glassy carbon working electrodes against a Ag/AgCl reference electrode in CH_2Cl_2 with tetra-*n*-butylammonium hexafluorophosphate as the supporting electrolyte. ^bIrreversible cathodic reduction reported as cathodic peak potential (E_{pc}) in parentheses. ^cMeasured in CH_2Cl_2 .





Figure 3. Cyclic (CV) and differential pulse voltammetry (DPV) measurements of **3a** recorded in 0.2M CH_2Cl_2 solution of Bu_4NPF_6 at a scan rate of 0.10 V/s.

The stabilities of bis(phenyl)dirhodium(III) carboxamidates are noteworthy. They are air stable and thermally stable to temperatures below 130°C. They do not react with nucleophiles such as cyanide or azide, but they are sensitive to electrophilic attack. Treatment of compound *bis-σ*-(6-methoxynaphthalen-2-yl)-*tetrakis-μ*-(caprolactamato)dirhodium(III)¹⁹ with trifluoroacetic acid, for example, resulted in quantitative conversion to 2-methoxynaphthalene (eq 3) and caprolactam (capH). Product analysis as a function of time with *bis-σ*-(phenyl)-*tetrakis-μ*-(caprolactamato)dirhodium(III) demonstrated the stability of the phenyl-rhodium(III) bond (Figure 4); caprolactam was released from the complex prior to release of benzene, but attempts to isolate the anticipated diphenyl derivative of dirhodium(III) trifluoroacetate were unsuccessful. Although *bis-σ*-(phenyl)-*tetrakis-μ*-(caprolactamato)dirhodium(III) was stable as a solid to temperatures above 200° C with a measured decomposition point at 256° C, when this compound was heated in refluxing chlorobenzene over long periods of time, biphenyl as well as mono- and di-chlorobiphenyl are formed, suggestive of homolytic cleavage of the rhodium-phenyl bond.







Figure 4. Reaction of PhRh(cap)₄RhPh (3.7 μmol) with (a) 10 and (b) 15 molar equiv. of trifluoroacetic acid in CDCl₃ at room temperature.

Computational study. The structural novelty of the bis(aryl)dirhodium(III) family and their peculiar properties (metal-metal bond length, stability), in comparison with isoelectronic systems (see Introduction) warrant a more detailed analysis of their electronic structure. DFT methods have proven quite successful when applied to dinuclear metal-metal bonded compounds.^{21, 22} The B3LYP functional provides the best agreement between experimental and calculated structures,¹ and it has therefore been adopted in this study. The study used the standard LANL2DZ basis set in order to provide a qualitative description of the electronic structure. A model system was constructed with the amidato ligands being replaced with simple formamidato ligands, [OCCHNH]⁻.

(a) [Rh₂(amidato)₄], [Rh₂(amidato)₄(MeCN)₂] and [Rh₂(amidato)₄(MeCN)₂]⁺ systems.

Before analyzing the more interesting bis(aryl)dirhodium(III,III) derivatives, it is necessary to start with the dirhodium(II,II) and dirhodium(II,III) precursors. These families of compounds have already been extensively studied, thus our analysis is limited to a comparison of the optimized and experimental geometries, and to a comparison of the basic electronic structural features with those previously presented for other members of the same families²³⁻³⁰ (fuller details are available as Supporting Information). Geometry optimizations were carried out for the simple tetrakis(amidato)dirhodium(II,II) system (**I** in Scheme 2), for its bis(acetonitrile) adduct (**II**), and for the one-electron oxidation product of

the latter (**III**). All these calculations were only carried out with the experimentally observed head-head-tail-tail (*cis*-2,2)²⁰ relative orientation of the four bridging amidato ligands (see Scheme 2). The salient features of the optimized geometries are shown in Table 5. There is excellent agreement between optimized and experimental parameters, attesting to the suitability of the computational method and of the chosen model. The Rh-Rh distance trend is perfectly reproduced by the calculation. The very minor elongation (0.045 Å) computed on going from **I** to **II** agrees with the notion that this bond is rather insensitive to the presence of axial ligands, an experimental example being Rh₂(O₂C-C₆H₂iPr₃-2,4,6)₄ and its bis(acetone) adduct (distances of 2.350(1) and 2.370(1), respectively).^{31, 32}



Scheme 2. Tetraformamidato systems used for the DFT calculations.

Table 5. Selected DFT-optimized geometric parameters (distances in Å, angles in deg) for the model compounds **I-III** and comparison with the experimental geometries.

Parameter	System I		System II		System III	
		Opt.	Exp. ^a	Opt.	Opt.	Exp. ^b
(a) Distances						
Rh-Rh	2.454	2.497	2.43(2)	2.481	2.40(3)	
Rh-N(amidato)	2.047	2.039	2.03(2)	2.011	2.00(2)	
Rh-O	2.085	2.112	2.066(4)	2.064	2.025(7)	
Rh-N(acetonitrile)	-	2.286	2.28(5)	2.285	2.28(3)	
(b) Angles						
Rh-Rh-N(amidato)	87.15	86.36	87(1)	86.95	87(1)	
Rh-Rh-O	89.25	88.80	89.3(7)	88.32	89.3(2)	
Rh-Rh-N(acetonitrile)	-	176.61	175(2)	177.54	174(4)	

^aAverages over all chemically equivalent parameters of compounds **1a**, **1b** and **1c**. ^bAverages over all chemically equivalent parameters of compounds **2a'**, **2b'** and **2c**.

The specific Rh-Rh σ , π , and δ^* interactions are symmetry-mixed with symmetry-adapted linear combinations (SALC) of the Rh-O and Rh-N bonds. The effect of this mixing is that each of these interactions is mainly distributed in two molecular orbitals. Within each pair of orbitals, the one carrying the greater Rh-Rh character is that at lower energy for σ and π and that at higher energy for δ^* . From the energetic point of view, the major Rh-Rh σ MO is found at higher energy than the major Rh-Rh π MO, and the major Rh-Rh δ^* MO is higher than the Rh-Rh π^* MO. Upon MeCN axial ligation, the σ competition between the Rh-NCMe and Rh-Rh interactions has the effect of raising the Rh-Rh σ level. Weak axial ligands (e.g., H₂O) establish weak interactions with the Rh atoms and weakly affect the Rh-Rh σ MO energy,²³ whereas stronger ligands (e.g., phosphines) have a more pronounced effect and greatly raise the energy of the Rh-Rh σ MO to the point that this may even become the HOMO.²⁵ In the present case, the interaction is relatively weak but sufficient to push the Rh-Rh σ MO higher than the Rh-Rh δ MO but lower than the Rh-Rh π^* MOs. Upon MeCN ligation, the energy of all orbitals is raised by ca. 0.4 eV on average, reflecting the additional electron density provided to the Rh atoms by the axial ligands. On the other hand, upon one-electron oxidation, all orbitals drop by ca. 1.5 eV on average as a result of the increased effective positive charge, but the MO energy ordering remains approximately unchanged (the σ level remains between the δ and π^* levels). For all three systems, the HOMO (doubly occupied for **I** and **II**, singly occupied for **III**) is the Rh-Rh δ^* MO.

(b) The [Rh₂(amidato)₄Ph₂] system.

A geometry optimization was carried out on the model system Rh₂(OCHNH)₄(C₆H₅)₂ with the same (*cis*-2,2) arrangement as in the **I-III** congeners (**IV**, see Scheme 2). The input geometry was constructed directly from the experimentally observed geometry of compound **3c**. Since this geometry is very close to an ideal *C*_{2h} symmetry, that symmetry was imposed in the initial geometry optimization. The converged geometry (Figure 5, *i*) remained very close to the initial one, but the frequency analysis

revealed one imaginary frequency whose motion showed the tendency for the two Rh-Ph axes to move in opposite directions away from the symmetry plane. Therefore, a free optimization in C_1 symmetry was subsequently carried out. The resulting geometry (Figure 5, *ii*) was essentially C_i -symmetric, with the expected 3N-6 positive vibrational frequencies, and with a co-planar arrangement for the two Ph groups. The energy of the global minimum is just 0.2 kcal/mol lower than the C_{2h} structure. The ideal geometry where the Ph-Rh-Rh-Ph moiety was forced to be coplanar and linear (Figure 5, *iii*), on the other hand, turned out to be energetically less favored by 8.2 kcal/mol. These combined results indicate that the origin of the experimentally observed distortion of the Rh-Ph bonds is not in the steric interaction between the Ph groups and the amidato ligands, but rather in an electronic effect that forces the three-bond Ph-Rh-Rh-Ph sequence out of co-linearity. Of the two different and nearly isoenergetic distortion modes (*i*) and (*ii*), the first one is adopted in the crystal structure for possible packing reasons.




Figure 5. Two mutually perpendicular views of the geometries, and relative energies, of system **IV** obtained under three different conditions: (*i*) constrained optimization in C_{2h} symmetry; (*ii*) full optimization without symmetry constraints; (*iii*) idealized geometry with linear Ph-Rh-Rh-Ph moiety.

Selected optimized geometric parameters for system **IV** are shown in Table 6. Like for systems **II** and **III**, the optimized structure of **IV** (C_{2h} symmetry) agrees very well with the observed structures. Except for the Rh-Rh distance (ca. 0.1 Å longer in the optimized geometry), all distances are in very close

agreement with the experimentally determined structure. A significant difference is observed in the Rh-Rh-Ph angle (ca. 10° smaller in the X-ray structures), presumably because of crystal packing effects. The distortion in the lower-energy C_i structure is different. It causes a slight asymmetry in the Rh coordination geometry yielding alternating long-short Rh-N and Rh-O distances, which is confirmed experimentally. The longer distances are associated to the smaller Rh-Rh-X (X = N, O) angles.

Table 6. Selected DFT-optimized geometric parameters (distances in Å, angles in deg) for the model compound **IV** and comparison with the experimental geometries.

Parameter	System IV (no symmetry)	System IV (C_{2h} symmetry)	Exp. ^a
(a) Distances			
Rh-Rh	2.644	2.639	2.54(3)
Rh-N	2.025, 2.053	2.022	2.01(2)
Rh-O	2.095, 2.133	2.138	2.09(2)
Rh-C	1.999	1.999	2.01(2)
(b) Angles			
Rh-Rh-N	90.22, 81.83	90.73	93(1)
Rh-Rh-O	89.75, 81.50	81.13	81(1)
Rh-Rh-C	166.92	166.44	159(4)

^aAverages over all chemically equivalent parameters of compounds **3a**, **3b** and **3c**.

The electronic structure analysis helps rationalize the molecular constitution and geometry of the novel dinuclear Rh^{III} complexes. The essential features are reported in Figure 6, while fuller details are presented in the Supporting Information. The relevant MO's for **IV** derive from the full geometry optimization, while those of the Rh₂(OCHNH)₄ fragment (**V**) and the Ph radicals are calculated for the corresponding fixed geometries as found in **IV**. The frontier orbitals of interest for the Ph radical are the

filled, familiar Hückel-type π_1 , π_2 and π_3 orbitals, plus the σ orbital pointing out from the unsaturated carbon in a radial direction. The latter is expected to be used for the formation of the Rh^{III}-Ph σ bond.




Figure 6. Simplified interaction diagram for system IV, built from the Rh₂(OCHNH)₄ fragment (V) and two phenyl radicals.

Upon interaction between V and the two Ph radicals, we observe that those molecular orbitals that are not involved in a significant interaction change their energy only slightly. More specifically, there is a slight energy increase for the phenyl orbitals 17 (π_1) and 20 (π_3) and a slight energy decrease for V orbitals 51 and 52 (Rh-Rh π), 59 (Rh-Rh δ), and 63 (Rh-Rh δ^*). Inspection of the contour plots of the related orbitals in system IV (see Supporting Information) shows that they do not involve a significant mixing of components from the different fragments.

The greater energy change experienced by the Ph π_2 orbitals is caused by the combination with the appropriately oriented Rh-Rh π and π^* orbitals of V, resulting in Rh-Ph π -type interaction. However,

since these are filled-filled interactions (both the in-phase and out-of-phase combinations are filled) there is no net Rh-Ph π bonding. The interaction is stronger (greater mixing and energy changes) for the Rh-Rh π^* orbital because of a better energy match. The in-phase and out-of-phase combinations are found in orbitals 94 and 103, while the corresponding combinations with the Rh-Rh π MO are found in orbitals 93 and 102.

The most important interaction between **V** and the two Ph radicals results from the combination of the σ -type orbitals. There are four such orbitals in the frontier region, two for Rh₂(OCHNH)₄ (the filled Rh-Rh σ orbital, mixed with a nonbonding ligand-based combination in MO's 54 and 55, *vide supra*, and the empty Rh-Rh σ^* orbital, MO 64), and two for the two Ph radicals (the half-filled frontier σ orbital, MO 21). This gives rise to two filled MO's (86 and 91) and to two empty MO's (105 and 108) in **IV**, all of which are shown in Figure 7 (a more extensive sample of MO contour diagrams for **IV** is presented in the Supporting Information). The two filled orbitals have mostly Rh-Ph σ character but also feature other contributions: the former has substantial Rh-Rh and C-C/C-H σ contributions, whereas the latter has Rh-Rh σ^* character. Thus, very little Rh-Rh σ bonding is present in the molecule. Most of the Rh-Rh σ interaction is found in the empty MO 105, which also has significant Rh-Ph σ^* character and is therefore raised above the energy of the Rh-Rh δ^* orbital. Finally, MO 108 has both Rh-Rh and Rh-Ph σ^* character (Figure 7). The Rh-Rh interaction in this molecule can therefore be formally described as $\pi^4\delta^2\pi^*\delta^*2$, leading to a formal bond order of zero.

Using a basic, qualitative valence bond approach, we can describe the molecule as two interpenetrating octahedra, using the familiar d^2sp^3 hybrids for bonding. This would lead to the establishment of all three σ interactions (two Rh-Ph bonds and the Rh-Rh bond) by implication of the higher energy 5s and 5p_z orbitals for the two rhodium atoms. Obviously, this implication is not extensive enough to keep the third bonding orbital sufficiently low in energy to be occupied. We note that the involvement of the higher energy s orbital has been invoked³³ to rationalize the observed short distance of paddle-wheel d^3-d^3 W₂(O₂CET)₄R'₂ compounds (E = Et and R' = CH₂Ph or CH₂tBu),³⁴ which are able

to maintain their W-W σ interaction as in the parent d^4 - d^4 $W_2(O_2CEt)_4$ compound. The involvement of the higher energy orbitals seems a more common feature for 5d elements, as a result of relativistic effects. The effect of the σ interaction scheme on the electronic structure is such that the HOMO of **IV** is the same orbital, the δ^* orbital, as the HOMO of **I**. The two are essentially identical because the Ph groups do not significantly contribute. This agrees with the observed reversible electrochemical behavior upon oxidation: the electron is pulled off the essentially nonbonding δ^* orbital, thus the Rh_2^{7+} oxidation product is predicted to have essentially no Rh-Rh bonding interaction and to display a structure closely related to that of its Rh_2^{6+} precursor.

One remaining point of interest is the origin of the non linear arrangement of the Ph-Rh-Rh-Ph moiety. One possible cause, related to the Ph-Rh-Rh-Ph σ bonding scheme, is a second order Jahn-Teller effect involving the filled Rh-Ph and the empty Rh-Rh σ orbitals. However, this idea is contradicted by the higher energy of the bonding orbitals 86 and 91 in the optimized minimum relative to the idealized linear geometry (*iii* in Figure 5, orbital energies available in SI). This suggests that the Ph-Rh-Rh-Ph σ bonding framework prefers the linear configuration. The relative weakness of this interaction and its symmetry-mixing with other bonding features, however, do not allow its clear identification in the molecular orbitals. Therefore, the reason for the molecular distortion of **3** out of linearity must be attributed to other factors. A possible clue comes from the observation of very narrow Rh-Rh'-O bond angles in **3** (less than 90°), suggesting the presence of an attractive interaction between Rh and the O atom bonded to Rh'. The through space crystallographic distances between rhodium and its distal oxygens are 3.00 and 3.24 Å for **3a**, 2.82 and 3.11 Å for **3b**, and 2.90 and 2.97 Å for **3c**. Given the absence of direct Rh-Rh bonding, the Rh atoms are in fact electronically unsaturated (16-electron configuration), and therefore seek electron density from the distal O lone pairs. However, we note that an alkylrhodium(III) compound with a nearly identical coordination geometry, the O_2N_2 coordination environment being provided by a Schiff base ligand, maintains a mononuclear 5-coordination geometry with an open coordination site.³⁵




Figure 7. MOLDEN contour plots of relevant orbitals for system IV.

(c) The $[Rh_2(O_2CCH_3)_4(NO)_2]$ system.

Compounds $Rh_2(O_2CR)_4(NO)_2$ ($R = Me, Et, iPr$), featuring the same paddlewheel structure as compounds **3** with axial NO ligation and a very similar Rh-Rh distance, have been recently described.³ The electronic structure of these compounds (with $R = H$ and CH_3), has also been reported.⁴⁻⁶ Here, we wish to compare the electronic structures of system $Rh_2(O_2CCH_3)_4(NO)$ (**VI**), and **IV**. For this purpose, the geometry of **VI** has been optimized at the same computational level as **IV**. The orbital interaction diagram for **VI** with respect to its parent $Rh_2(O_2CCH_3)_4$ and NO fragments is shown in Figure 8, while fuller details of the electronic structure of **VI** are presented in the Supporting Information.

The axial donor, in this case, mostly interacts with the dimetal system via the N lone pair (which is mixed with the O lone pair in MO 7). However, because of the Rh-N-O bending, the NO $\pi_{||}$ and $\pi_{||}^*$ orbitals can also contribute to the Rh-Rh and Rh-N σ bonding scheme. The perpendicular N-O π orbitals (π_{\perp} and π_{\perp}^*), have the correct symmetry to mix with the Rh-Rh π and π^* components of the $Rh_2(O_2CCH_3)_4$ fragment, much like what happens in structure **IV** for the phenyl π_2 orbitals (*vide supra*).




Figure 8. Simplified interaction diagram for system VI, built from $\text{Rh}_2(\text{O}_2\text{CCH}_3)_4$ and two nitrosyl radicals.

Going from the fragments to the full system, as with the formamidato system discussed above, the metal-metal π , δ , π^* and δ^* orbitals are not significantly affected. The N-O π_{\perp} orbitals can potentially establish Rh-N π interactions but, because of their relatively deep energy, do not significantly mix with the metal orbitals and therefore are also little effected by the axial coordination. As for the amidato system, the most important change concerns the σ -type orbitals. Relative to the diphenyl system, the situation is complicated by the greater number of participating orbitals [the Rh-Rh σ and σ^* orbitals from fragment V, plus the N lone pair and the N-O $\pi_{||}$ and $\pi_{||}^*$ orbitals from each of the two NO ligands, for a total of 8 resulting MO's]. The four lowest energy combinations (orbitals 51, 52, 61 and 65 in

Figure 8) have their major contribution from the Rh-N σ and N-O $\pi_{||}$ bonds. Contour diagrams of the other four orbitals are shown in Figure 9 (other relevant orbitals are shown in the Supporting Information). They are qualitatively similar to those of system IV, including their occupation. Thus, as with the diphenyl formamidato system, the formation of the two Rh-NO bonds essentially destroys the Rh-Rh interaction, since the Rh-Rh σ^* component that was empty in the dinuclear Rh^{II} fragment becomes now engaged in the axial σ -bonding system. That there is no N-Rh-Rh-N distortion in the nitrosyl complex is likely due to the symmetric nature of the 3-atom bridges; since oxygen atoms constitute both sides, the molecule remains symmetric.




Figure 9. Contour plots of relevant orbitals for system VI.

(d) Comparison of electronic structures

A comparison between the two MO diagrams of IV in Figure 6 and VI in Figure 8 shows that those orbitals with the largest Rh contribution, and not involved in the σ interactions (namely the π , π^* , δ and

δ^* orbitals), have lower energies for system **VI** than **IV**. This is rationalized by the more electronegative nature of the carboxylato ligands relative to the amidato ligands. The most important difference between the two systems, however, is in the σ manifold. For **IV**, the Rh-Rh(σ^*) of the Rh₂(OCHNH)₄ fragment interacts strongly with the Ph donor orbitals, yielding a stabilized bonding orbital and leaving the Rh-Rh(δ^*) orbital as the HOMO. The unexpectedly low observed increase in binding energy for the *d*-electrons (XPS data in Table 3) upon going from compounds **2** to compounds **3** is consistent with the relatively small change of electronic structure in the frontier region and with the reversible oxidative electrochemistry (*vide supra*).

On the other hand, the interaction between the Rh-Rh(σ^*) orbital of the Rh₂(O₂CCH₃)₄ fragment and the orbitals of the NO axial donors is not nearly as strong and yields a MO at relatively high energy, the HOMO of system **VI**. The nature of this orbital is Rh-Rh(σ^*) and Rh-N(σ), see Figure 9, thus oxidation of this compound may be expected to lead to rupture of the Rh-NO interactions and to the establishment of a Rh-Rh bond. Thus, although systems **IV** and **VI** have a closely related electronic structure and very similar bonding parameters, their chemical behavior toward oxidation should be drastically different.

It is also interesting to compare the electronic structure of these two systems with that of the isoelectronic Ir₂⁶⁺ complex Ir₂(hpp)₄Cl₂ (shown on the right in Scheme 1). No computational study on this system has been presented as yet, to the best of our knowledge. There is little doubt, however, that the general features of the electronic structure will parallel those presented above for systems **IV** and **VI**. The question of importance is the strength of the σ interaction between the axial ligands and the metal-metal framework. For the electronegative and spherically symmetrical Cl atom, the 4 resulting MO (analogous to 86, 91, 105 and 108 for system **IV** or to 87, 94, 96 and 100 for system **VI**) would certainly end up at lower energy relative to the block of metal-metal orbitals of π and δ symmetry. The stronger relativistic effect for the 5d Ir element should also contribute to a stabilization of the orbitals of M-M σ symmetry through a greater involvement of the metal *s* orbital. In case the second highest of these four orbitals (having M-M σ and M-L_{ax} σ^* contributions) is stabilized to the point of falling below the

energy of the HOMO, then the molecule will indeed have a formal metal-metal double bond, because the two electrons needed to fill the M-M σ orbital originate from a previously filled antibonding orbital of π^* or δ^* type. If δ^* is located below π^* , the molecule will display two unpaired electrons as is indeed observed for compound $\text{Ir}_2(\text{hpp})_4\text{Cl}_2$.

Conclusion

The ability of a phenyl group to impart unusual stability to rhodium(III) compounds is explained by theoretical analysis of the electronic structure of bis- σ -(phenyl)-tetrakis- μ -(carboxamido)dirhodium(III). Formally, two phenyl radicals in combination with a rhodium-rhodium bond are transformed into two phenyl-rhodium bonds. The absence of a rhodium-rhodium σ bonding interaction is suggested in the lengthening of the rhodium-rhodium distances from those of dirhodium(II,II) and dirhodium(II,III) complexes, as well as from XRD data, and is confirmed by DFT analysis. The stability of the phenyl-rhodium bonds is seen in the preferential removal of equatorial carboxamidate ligands by trifluoroacetic acid and in the homolytic loss of phenyl radical from **3** at high reaction temperatures. The observed severe distortion of phenyl rings from linearity is suggested to have its origin in long-range interactions between the electron deficient rhodium and the distal oxygen atoms. These structural characteristics are unique among M_2^{n+} compounds, and they suggest intriguing potential for future structural revelations.

Experimental Section

General: All oxidation reactions were performed in air. NaBPh_4 and CuOTf were purchased and used as received. ^1H NMR (400 MHz) and ^{13}C NMR (100 MHz) spectra were obtained from solutions in CDCl_3 . Chemical shifts are reported in parts per million (ppm, δ) downfield from Me_4Si (TMS); coupling constants are reported in Hertz (Hz). High resolution mass spectra were done on an ESI instrument. UV-Visible spectra were obtained using a xenon flash lamp. Both cyclic and differential pulse voltammograms were recorded in 0.2 M (*n*-Bu) $_4\text{NPF}_6$ solution (CH_2Cl_2 , N_2 -degassed) on a voltammetric analyzer with a glassy carbon working electrode, a platinum-wire auxiliary electrode, and

a Ag/AgCl reference electrode. The ferrocenium/ferrocene couple was observed at 0.42 V (vs. Ag/AgCl) under experimental conditions. All mass spectra were recorded on JEOL AccuTOF-CS from CH₂Cl₂ solution using MeCN/H₂O or MeOH/H₂O (9:1) as solvent in presence of 1% of triflic acid. Elemental analysis were carried out by the Service d'Analyse du Laboratoire de Chimie de Coordination in Toulouse. Dirhodium(II) acetate, carboxamidate ligands, and reagents described in this report were obtained commercially.

Tetrakis- μ -(pyrrolidinato)bis(acetonitrile)dirhodium(II) (1a).¹⁶ In a dry flask under N₂ were mixed Rh₂(OAc)₄ (500 mg, 1.31 mmol), 2-pyrrolidinone (2.3 g, 27.0 mmol) and anhydrous chlorobenzene (40 mL). The flask was equipped with a Soxhlet apparatus with a thimble containing a 2:1 NaHCO₃/sand mixture (5 g). The solution was kept under N₂ and stirred at reflux for 24 hours, after which the solution was cooled and solvent was removed under reduced pressure. The residue was washed with diethyl ether (2 x 50 mL) and pentane (50 mL). The purple solid obtained was heated at 80°C in acetonitrile (15 mL), and some methanol was added to dissolve the entire solid. The mixture was crystallized at 20°C to give a pink-red solid. Yield: 565 mg (80%). Suitable x-ray quality crystals (pink-red prisms) were obtained by slow evaporation of diethyl ether from an acetonitrile solution. ¹H NMR (400 MHz, CD₃CN + CD₃OD): δ 3.55 (br, N-CH₂), 3.44 (m, N-CH₂), 2.34 (t, ³J = 7.6 Hz, 8H, CH₂-C), 1.98 (q, ³J = 7.6 Hz, 8H, N-CH₂-CH₂). ¹³C NMR (100 MHz, CD₃CN + CD₃OD) : δ 186.89 (s, NCO), 55.41 (s, N-CH₂), 33.12 (s, CH₂-C), 21.91 (s, N-CH₂-CH₂). MS (ESI), m/z, (%): 584.01, (100) [Rh₂C₁₈H₂₈N₅O₄]⁺ = M(MeCN)⁺; 625.04, (28) [Rh₂C₂₀H₃₁N₆O₄] = M(MeCN)₂⁺. UV/visible spectrum (MeCN + MeOH, one drop), λ (ϵ , M⁻¹cm⁻¹) : 350 nm (224); 505 nm (235). XPS: Rh-3d a 5/2 308.000 eV; Rh-3d b 5/2 309.082 eV; Rh-3d a 3/2 312.563 eV; Rh-3d b 3/2 313.648 eV.

Tetrakis- μ -(pyrrolidinato)bis(acetonitrile)dirhodium(II,III) Tetrafluoroborate (2a). A solution of NOBF₄ (22.3 mg, 191 μ mol) in anhydrous acetonitrile (9 mL) was slowly added (over 3 min) to a suspension of **1a** (108.5 mg, 174 μ mol) in CH₂Cl₂ (25 mL). The suspension rapidly turned from purple to dark red-purple. After complete addition the mixture was stirred for an additional 30 min at room temperature. Solvents were removed under reduced pressure, and the residue was dissolved in CH₂Cl₂ (5

mL), then filtered. The filtrate was concentrated (*ca.* 2-3 mL), precipitated in diethyl ether (50 mL), filtered and dried under vacuum. A dark red-violet solid was obtained (117.2 mg, 94.8%), but the bis-acetonitrile complex did not form crystals suitable for X-ray analysis. Suitable X-ray crystals (black plate) of the bis(benzonitrile) adduct, **2a'**, were obtained after an axial ligand exchange (by heating the product 5 min at 60°C in a small amount of benzonitrile) and a layer diffusion of diethyl ether in a 2:1 benzonitrile/CH₂Cl₂ solution. A satisfactory elemental analysis for this compound could not be obtained, presumably because of the facile axial ligand exchange. NMR silent; HRMS (ESI) calcd. for Rh₂C₁₈H₂₇N₅O₄ (M(MeCN)⁺): 583.01731; found: 583.01516. MS (ESI), *m/z*, (%): 574.1, (100) [Rh₂C₁₇H₂₈N₄O₅]⁺ = M(MeOH)⁺; 542.0, (24) [Rh₂C₁₆H₂₄N₄O₄]⁺ = M⁺; 87.0, (100) [BF₄]⁻. UV/Visible spectrum (CH₂Cl₂), λ (ε, M⁻¹cm⁻¹) : 489 nm (4150); 530 nm (-); 1030 nm (1440). XPS : Rh-3d a 5/2 308.905 eV; Rh-3d b 5/2 310.722 eV; Rh-3d a 3/2 313.514 eV; Rh-3d b 3/2 315.069 eV.

Bis-σ(phenyl)-tetrakis-μ-(pyrrolidinato)dirhodium(III) (3a). To a solution MeOH:CH₂Cl₂ solution (5 mL, 1:1) of **2a** (30.5 mg, 42.9 μmol) were added NaBPh₄ (73.4 mg, 214.5 μmol) and CuOTf (1.5 mg, 7.1 μmol). The initial purple solution rapidly turned to yellow-green, and the mixture was stirred at room temperature for 18 hours. During this period a yellow-green solid precipitated. Solvents were removed under reduced pressure, MeOH (10 mL) was added to the residue, and the mixture was stirred 5 min to obtain a homogenous suspension. The mixture was filtered, washed with MeOH (2 x 5 mL) and Et₂O (10 mL), and dried under vacuum to give a yellow-green product. Yield: 20.6 mg (69%). Suitable crystals (yellow prism) for x-ray analysis were obtained by slow evaporation of dichloromethane from a CH₂Cl₂ solution. Calcd. for C₂₈H₃₄N₄O₄Rh₂ (Mw = 696.41): C 48.29, H 4.92, N 8.05%. Found: C 48.10, H 4.93, N 7.29%. ¹H NMR (400 MHz, CDCl₃): δ 7.33 (m, 4H, *o*-Ph), 7.1 (m, 6H, *m,p*-Ph), 3.64 (m, 4H, N-CH₂), 3.01 (m, 4H, N-CH₂), 2.52 (m, 4H, CH₂-C), 2.41 (m, 4H, CH₂-C), 1.94 (m, 4H, N-CH₂-CH₂), 1.82 (m, 4H, N-CH₂-CH₂). ¹³C NMR (100.6 MHz, CDCl₃): δ 183.00 (s, NCO), 141.18 (d, ¹J_{Rh-C} = 34 Hz), 136.60 (s, *o*-Ph), 127.50 (s, *m*-Ph), 125.21 (s, *p*-Ph), 53.46 (s, N-CH₂), 34.12 (s, CH₂-C), 21.82 (s, N-CH₂-CH₂). UV/Visible spectrum (CH₂Cl₂) 442 nm (ε = 4492 M⁻¹cm⁻¹).

$^1\text{cm}^{-1}$). HRMS (ESI) calcd for $\text{Rh}_2\text{C}_{28}\text{H}_{35}\text{N}_4\text{O}_4$ 697.07684, found 697.07580 (MH^+). XPS : Rh-3d a 5/2: 309.045 eV; Rh-3d a 3/2: 313.739 eV.

Compounds 1b, 2b, 2b', 3b, 1c, 2c, and 3c. These compounds were synthesized by the same general procedures described above for compounds **1a, 2a, 2a'**, and **3a**. The details are given as Supporting Information.

Reaction of *Bis- σ -(6-methoxynaphthalen-2-yl)-tetrakis- μ -(caprolactamato)dirhodium(III) with Trifluoroacetic Acid.* A solution of trifluoroacetic acid (114 mg, 1.0 mmol) in CH_2Cl_2 (1 mL) was slowly added (over 5 min) to a solution of *bis- σ -(6-methoxynaphthalen-2-yl)-tetrakis- μ -(caprolactamato)dirhodium(III)* (52 mg, 53.7 μmol) in anhydrous CH_2Cl_2 (6 mL). The reaction solution rapidly turned from green to orange. After completed addition, the mixture was stirred at room temperature for 24 hrs. Solvents were then removed under reduced pressure, and the residue was chromatographed on silica gel using Hexanes/EtOAc (10:1) as eluent. The product 2-methoxynaphthalene was obtained as white solid (15.5 mg, 91% yield). Mp: 71–73 $^\circ\text{C}$; lit.³⁶ 71–73 $^\circ\text{C}$. ^1H NMR (400 MHz, CDCl_3): δ 7.76–7.74 (m, 3H), 7.44–7.39 (m, 1H), 7.34–7.29 (m, 1H), 7.14–7.10 (m, 2H), 3.91 (s, 3H).

Reaction of 3c with Trifluoroacetic Acid. In a dry NMR tube were added 3.0 mg (3.7 μmol) bis(phenyl)dirhodium(III) caprolactamate (**3c**) and 0.5 mL CDCl_3 . After the solid was dissolved and a clear green solution was obtained, 13 μL (18.5 μmol , 5 equiv.) or 26 μL (37.0 μmol , 10 equiv.) or 39 μL (55.5 μmol , 15 equiv.) trifluoroacetic acid solution in CDCl_3 (1.35 mmol/mL) was then added. The mixture was rapidly oscillated for a minute and then submitted to ^1H NMR to follow reaction progress. Spectra were taken every 5 minutes for 1 h and then every hour, and data were obtained by using CHCl_3 in CDCl_3 as an internal standard.

Computational details. All geometry optimizations were performed using the B3LYP three-parameter hybrid density functional method of Becke,³⁷ as implemented in the Gaussian03 suite of programs.³⁸ The basis functions consisted of the standard LANL2DZ for all atoms, which included the

Hay and Wadt effective core potentials (ECP)³⁹ for Rh. Unless otherwise stated, the geometry optimizations were carried out without any symmetry constraint and the final geometries were characterized as local minima of the potential energy surface (PES) by verifying that all second derivatives of the energy were positive. The unrestricted formulation was used for the open-shell, mixed valence Rh₂⁵⁺ system. The value of $\langle S^2 \rangle$ at convergence was 0.7579, very close to the expected value of 3/4, indicating minor spin contamination.

Acknowledgement. We are grateful to the National Science Foundation (CHE-0456911) for their generous support. We also thank the Fond Social Européen for funding Joffrey Wolf and Bristol-Meyers-Squibb for supporting Jason Nichols with an American Chemical Society Organic Division Fellowship. RP thanks the CICT (Project CALMIP) for granting free computer time.

Supporting Information Available. Synthesis and characterization of **2** and **3**. Crystallographic data for eight compounds (**1-3**) in CIF format are available as supporting information. Cartesian coordinates and relative energies of all optimized structures. Energies, composition and contour diagrams of the relevant molecular orbitals for geometry optimized systems **I**, **II**, **III**, **IV** and **VI**.

References

- ¹ Cotton, F. A.; Murillo, C. A.; Walton, R. A., *Multiple Bonds Between Metal Atoms*. 3rd ed.; Springer Science: New York, 2005; 'Vol.' p.
- ² Nichols, J. M.; Wolf, J.; Zavalij, P.; Varughese, B.; Doyle, M. P., *J. Am. Chem. Soc.* **2007**, 129, 3504-3505.
- ³ Hilderbrand, S. A.; Lim, M. H.; Lippard, S. J., *J. Am. Chem. Soc.* **2004**, 126, 4972-4978.
- ⁴ Sizova, O. V., *J. Mol. Struct. Theochem.* **2006**, 760, 183-187.
- ⁵ Sizova, O. V.; Skripnikov, L. V.; Sokolov, A. Y.; Ivanova, N. V., *Russ. J. Coord. Chem.* **2007**, 33, 588-593.
- ⁶ Sizova, O. V.; Sokolov, A. Y.; Skripnikov, L. V.; Baranovski, V. I., *Polyhedron* **2007**, 26, 4680-4690.
- ⁷ Cotton, F. A.; Murillo, C. A.; Timmons, D. J., *Chem. Commun.* **1999**, 1427-1428.
- ⁸ Cotton, F. A.; Ren, T.; Eglin, J. L., *Inorg. Chem.* **1991**, 30, 2552-2558.
- ⁹ Ren, T., *Organometallics* **2002**, 21, 732-738.
- ¹⁰ Bear, J. L.; Li, Y. L.; Han, B. C.; Kadish, K. M., *Inorg. Chem.* **1996**, 35, 1395-1398.
- ¹¹ Hurst, S. K.; Ren, T., *J. Organomet. Chem.* **2003**, 670, 188-197.
- ¹² Yao, C. L.; Park, K. H.; Khokhar, A. R.; Jun, M. J.; Bear, J. L., *Inorg. Chem.* **1990**, 29, 4033-4039.

13 Mendiratta, A.; Cummins, C. C.; Cotton, F. A.; Ibragimov, S. A.; Murillo, C. A.; Villagran, D.,
Inorg. Chem. **2006**, 45, 4328-4330.

14 Aakeroy, C. B.; Schultheiss, N.; Desper, J., *Dalton Trans.* **2006**, 1627-1635.

15 Liu, C. S.; Wang, J. J.; Yan, L. F.; Chang, Z.; Bu, X. H.; Sanudo, E. C.; Ribas, J., *Inorg. Chem.*
2007, 46, 6299-6310.

16 Rh₂(pyr)₄(Hpyr)₂ and Rh₂(val)₄(Hval)₂ have been prepared and their crystal structures reported:
Bear, J. L.; Lifsey, R. S.; Chau, K. L.; Ahsan, M. Q.; Korp, J. D.; Chavan, M.; Kadish, K. M. *J.*
Chem. Soc. Dalton Trans. **1989**, 93-100.

17 Wang, Y.; Wolf, J.; Zavalij, P.; Doyle, M. P., *Angew. Chem., Int. Ed.* **2008**, 47, 1439-1442.

18 For uses of nitrosonium salts as oxidants, see: Connelly, N. G.; Geiger, W. E. *Chem Rev.* **1996**,
96, 877-910.

19 Xie, J.-H.; Nichols, J. M.; Lubek, C.; Doyle, M. P., *Chem. Commun.* **2008**, 2671-2673.

20 Doyle, M. P.; Ren, T., *Prog. Inorg. Chem.* **2001**, 49, 113-168.

21 Cotton, F. A.; Feng, X., *J. Am. Chem. Soc.* **1997**, 119, 7514-7520.

22 Cotton, F. A.; Feng, X. J., *J. Am. Chem. Soc.* **1998**, 120, 3387-3397.

23 Norman, J. G., Jr.; Kolari, H. J., *J. Am. Chem. Soc.* **1978**, 100, 791-799.

24 Norman, J. G., Jr.; Renzoni, G. E.; Case, D. A., *J. Am. Chem. Soc.* **1979**, 101, 5256-67.

25 Bursten, B. E.; Cotton, F. A., *Inorg. Chem.* **1981**, 20, 3042-8.

26 Cotton, F. A.; Feng, X. J., *Inorg. Chem.* **1989**, 28, 1180-1183.

27 Kawamura, T.; Maeda, M.; Miyamoto, M.; Usami, H.; Imaeda, K.; Ebihara, M., *J. Am. Chem.*
Soc. **1998**, 120, 8136-8142.

28 Mougnot, P.; Demuynck, J.; Benard, M., *Chem. Phys. Lett.* **1987**, 136, 279-282.

29 Nakatsuji, H.; Ushio, J.; Kanda, K.; Onishi, Y.; Kawamura, T.; Yonezawa, T., *Chem. Phys. Lett.*
1981, 79, 299-304.

30 Nakatsuji, H.; Onishi, Y.; Ushio, J.; Yonezawa, T., *Inorg. Chem.* **1983**, 22, 1623-1630.

31 Cotton, F. A.; Hillard, E. A.; Murillo, C. A., *J. Am. Chem. Soc.* **2002**, 124, 5658-5660.

32 Cotton, F. A.; Hillard, E. A.; Liu, C. Y.; Murillo, C. A.; Wang, W. N.; Wang, X. P., *Inorg. Chim.*
Acta **2002**, 337, 233-246.

33 Chisholm, M. H.; Clark, D. L.; Huffman, J. C.; Van Der Sluys, W. G.; Kober, E. M.;
Lichtenberger, D. L.; Bursten, B. E., *J. Am. Chem. Soc.* **1987**, 109, 6796-6816.

34 Chisholm, M. H.; Hoffman, D. M.; Huffman, J. C.; Vandersluys, W. G.; Russo, S., *J. Am. Chem.*
Soc. **1984**, 106, 5386-5388.

35 Anderson, D. J.; Eisenberg, R., *Inorg. Chem.* **1994**, 33, 5378-5379.

36 Arsenijevic, L.; Arsenijevic, V.; Horeau, A.; Jacques, J., *Org. Syn.* **1973**, 53, 5-8.

37 Becke, A. D., *J. Chem. Phys.* **1993**, 98, 5648-5652.


38 Frisch, M. J., *et al.*, *Gaussian 03, Revision B.04.* ed.; Gaussian, Inc.: Wallingford CT, 2003.

39 Hay, P. J.; Wadt, W. R., *J. Chem. Phys.* **1985**, 82, 270-283.

TOC Graphics

Removal of Metal-Metal Bonding in a Dimetallic Paddlewheel Complex – Molecular and Electronic Structure of Bis(phenyl) Dirhodium(III) Carboxamidate Compounds

Joffrey Wolf,[†] Rinaldo Poli,^{†§*} Jian-Hua Xie,[‡] Jason Nichols,[‡] Bin Xi,⁺ Peter Zavalij[‡] and Michael P. Doyle^{‡*}



TOC Text

The molecular and electronic structures of bis(phenyl) tetracarboxamidatodirhodium(III) show that these thermally stable compounds do not have any metal-metal bonding. They have hydrolytically stable Rh(III)-Ph bonds and can be reversibly oxidized to Rh₂⁷⁺ complexes on the cyclic voltammetric timescale.