



**HAL**  
open science

# Reactivity of Phosphane–Imidazolium Salts Towards [Ir(COD)Cl] 2: Preparation of New Hydrido-iridium(III) Complexes Bearing Abnormal Carbenes

Joffrey Wolf, Agnès Labande, Jean-Claude Daran, Rinaldo Poli

► **To cite this version:**

Joffrey Wolf, Agnès Labande, Jean-Claude Daran, Rinaldo Poli. Reactivity of Phosphane–Imidazolium Salts Towards [Ir(COD)Cl] 2: Preparation of New Hydrido-iridium(III) Complexes Bearing Abnormal Carbenes. *European Journal of Inorganic Chemistry*, 2008, 2008 (19), pp.3024-3030. 10.1002/ejic.200800270 . hal-03192568

**HAL Id: hal-03192568**

**<https://hal.science/hal-03192568>**

Submitted on 8 Apr 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Reactivity of phosphane-imidazolium salts toward [Ir(COD)Cl]<sub>2</sub>: preparation of new hydridoiridium(III) complexes bearing abnormal carbenes.<sup>§</sup>**

1 Joffrey Wolf, Agnès Labande,\* Jean-Claude Daran and Rinaldo Poli\*

2  
3 *Laboratoire de Chimie de Coordination, UPR CNRS 8241 (lié par convention à l'Université*  
4 *Paul Sabatier et à l'Institut National Polytechnique de Toulouse), 205 route de Narbonne,*  
5 *31077 Toulouse Cedex 4, France. E-mail : [agnes.labande@lcc-toulouse.fr](mailto:agnes.labande@lcc-toulouse.fr)*

6  
7  
8  
9 **Keywords:** Carbenes / Phosphane ligands / Iridium / Hydrides / C-H activation

10  
11  
12  
13 **Corresponding author:**

14 Agnès Labande

15 Tél +33-561333158

16 Fax +33-561553003

17 **E-mail:** [Agnès.Labande@lcc-toulouse.fr](mailto:Agnès.Labande@lcc-toulouse.fr)

18  
19  
20  
21  
22  
23  
24  
25  
26  
27 <sup>§</sup> Dedicated to Prof. W. A. Herrmann on the occasion of his 60<sup>th</sup> birthday.

1     **Abstract**

2

3     The unusual reactivity of chelating phosphine-imidazolium salts MesImEtPPh<sub>2</sub><sup>+</sup>Br<sup>-</sup>, DIPP-  
4 ImEtPPh<sub>2</sub><sup>+</sup>Br<sup>-</sup> and MesImEtPPh<sub>2</sub><sup>+</sup>BF<sub>4</sub><sup>-</sup> toward low oxidation state iridium complex [Ir(COD)(μ-  
5 Cl)]<sub>2</sub> was studied. In the absence of a base, the C-H insertion at the C5 position of the  
6 imidazolium ring is the only reaction occurring, with no normal NHC observed, and lead to  
7 iridium(III) hydride complexes. This reactivity is independent of the nature of the  
8 imidazolium counteranion and of the substitution pattern of the aryl group.


9

## 1 Introduction

2  
3 N-heterocyclic carbene (NHC) complexes are now widely used for catalytic applications<sup>[1]</sup>.  
4 This interest is particularly due to the production of more robust catalysts by coordination of  
5 NHCs on transition metals such as palladium, nickel, rhodium, ruthenium and iridium<sup>[2]</sup>. Over  
6 the last fifteen years, a number of bidentate as well as polydentate ligands incorporating  
7 NHCs<sup>[3]</sup> and other coordinating atoms<sup>[4-6],[7,8]</sup> have been developed and used successfully in  
8 catalysis.

9 More recently a different binding mode for NHCs has been described by Crabtree and  
10 co-workers:<sup>[9]</sup> instead of the normal C2 coordination, the metal can be bound to the C4 or  
11 C5 atom of an imidazolium salt. Since this discovery, only few examples of “abnormal”  
12 carbenes have been described, for example with iridium,<sup>[10-12]</sup> ruthenium,<sup>[13]</sup> osmium<sup>[14]</sup>  
13 and several other transition metals.<sup>[15]</sup> In the case of Crabtree’s work with iridium  
14 precursor IrH<sub>5</sub>(PPh<sub>3</sub>)<sub>2</sub>, it has been shown that the formation of classical or abnormal  
15 NHCs was anion-dependent: imidazolium salts bearing a halogen anion led preferentially  
16 to classical NHCs, whereas non-coordinating anions BF<sub>4</sub><sup>-</sup> or PF<sub>6</sub><sup>-</sup> gave mostly abnormal  
17 NHCs.<sup>[12]</sup> Monodentate ligands have then been used instead of bidentate NHCs, with  
18 similar results, showing no chelate effect.<sup>[11]</sup> Very recently, the group of Li has reported  
19 iridium<sup>III</sup> hydride complexes bearing chelating phosphine-(abnormal)NHC ligands.<sup>[16]</sup>  
20 They did not observe any influence of the counter-anion on the product selectivity.

21 We are interested in the preparation of bifunctional imidazolium salts containing a  
22 coordinating heteroatom such as phosphorus or sulfur, and their use as ligands or ligand  
23 precursors for transition metals.<sup>[4-6]</sup> More particularly, we have recently described the use  
24 of phosphine-imidazolium salts to prepare zwitterionic Ni<sup>II</sup> complexes and shown their  
25 excellent activities for Kumada-Tamao-Corriu cross coupling reactions.<sup>[4]</sup> We now report  
26 the unusual reactivity of these salts with [IrCl(COD)]<sub>2</sub>. These results were obtained during  
27 attempts to synthesise new Ir<sup>I</sup> pre-catalysts, following the same procedure that allowed us  
28 to access Ni<sup>II</sup> and Pd<sup>II</sup> analogues.<sup>[4,5]</sup>




**Scheme 1.** Synthesis of phosphine-imidazolium salts. Mes = mesityl or 2,4,6-trimethylphenyl; DIPP = 2,6-diisopropylphenyl.

## 1 Results and discussion

2

3 The phosphine-imidazolium salts were prepared according to previously reported  
4 procedures (Scheme 1).<sup>[4]</sup> The reaction of 1-(2-diphenylphosphinoethyl)-3-(2,4,6-  
5 trimethylphenyl) imidazolium bromide salt MesImC<sub>2</sub>H<sub>4</sub>PPh<sub>2</sub><sup>+</sup>Br<sup>-</sup> **1** with [Ir(COD)(μ-Cl)]<sub>2</sub>  
6 yields a mixture of two hydride complexes (**4a** and **4b**) in a 1:1 ratio (Scheme 2). The two  
7 complexes exhibit two almost equivalent <sup>31</sup>P NMR signals at δ -1.1 and -2.4 and two  
8 equivalent <sup>1</sup>H NMR hydride doublet resonances at δ -15.22 and -14.55, see figure 1a.




9

10 **Scheme 2.** Mixture of isomers **4a+4b** from the reaction of ligand **1**, or isomers **6a+6b**  
11 from the reaction of ligand **2**, with [Ir(COD)(μ-Cl)]<sub>2</sub>.


12

13 We attribute this behavior to halogen exchange, induced by the bromide anion of **1**. This  
14 hypothesis is confirmed by the observation that using MesImC<sub>2</sub>H<sub>4</sub>PPh<sub>2</sub><sup>+</sup>BF<sub>4</sub><sup>-</sup> **3** in place of  
15 **1** leads to a single product **5**, whose <sup>31</sup>P and <sup>1</sup>H (hydride) resonances match with those of  
16 **4a** (Scheme 3, figure 1b). Thus, both products **4a** and **5** contain a chloride ligand and a  
17 different counterion (Br<sup>-</sup> for **4a** and BF<sub>4</sub><sup>-</sup> for **5**), whereas the chloride and bromide ions are  
18 exchanged in **4b**.




19

1 **Figure 1.**  $^1\text{H}$  NMR spectra (500 MHz, 20°C) of complexes a) **4a/4b** ( $\text{CDCl}_3$ ), b) **5**  
2 ( $\text{CD}_2\text{Cl}_2$ ), c) **6a/6b** ( $\text{CD}_2\text{Cl}_2$ ) (hydride resonances).




**Scheme 3.** Synthesis of complex **5** via an iridium(I) intermediate.

The  $^1\text{H}$  NMR spectrum of the **4a/4b** mixture indicates  $C_1$  symmetry and reveals three signals corresponding to the imidazole ring protons at  $\delta$  9.74, 6.24 and 6.18, with integration values of 1: 0.5: 0.5, whereas that of compound **5** exhibits two 1:1 resonances at  $\delta$  8.45 and 6.40, see Figure 2. Whereas the upfield imidazole ring  $^1\text{H}$  resonances of **5** approximately matches with that of **4a**, the downfield one is quite anion dependent. Furthermore, 1:1  $^{13}\text{C}$  resonances for quaternary carbons are observed at  $\delta$  119.9 and 121.1, which is a typical region for aryl substituents and not for NHC ligands. These results indicate that the products are  $\text{Ir}^{\text{III}}$  complexes coordinated by a bidentate phosphine-abnormal carbene ligand (scheme 2). The shift of the downfield  $^1\text{H}$  resonance is attributed to differences in H-bonding between the acidic imidazolium proton and the different counterion. Related iridium(III) hydride complexes, also featuring an abnormal carbene coordination, have been recently reported by Li *et al.* and show similar NMR features (complex **A**, figure 3).<sup>[16]</sup> Danopoulos *et al.* also obtained and crystallographically characterised a similar complex,  $\text{Ir}(\text{COD})(\text{Br})\{\kappa^2\text{-}P,C5\text{-Ph}_2\text{PCH}_2\text{CH}_2(\text{N}_2\text{C}_3\text{HMe}_s)\}$  (**B** in figure 3), from the reaction of **1** with  $[\text{Ir}(\text{COD})(\text{HCl})(\mu\text{-Cl})_2]_2$ . It is related to **4b** by elimination of  $\text{HCl}$ .<sup>[8]</sup>




**Figure 2.**  $^1\text{H}$  NMR spectra (500 MHz,  $20^\circ\text{C}$ ) of complexes a) **4a/4b** ( $\text{CDCl}_3$ ), b) **5** ( $\text{CD}_2\text{Cl}_2$ ), c) **6a/6b** ( $\text{CD}_2\text{Cl}_2$ ) (imidazolium resonances).



**Figure 3.** Related iridium complexes reported by Li *et al.* (**A**)<sup>[16]</sup> and Danopoulos *et al.* (**B**)<sup>[8]</sup>.

Complex **5** has also been characterised by X-ray crystallography (figure 4, table 1). The Ir-C(2) distance [2.068(5) Å] and the Ir(1)-Cl(1) distance are in agreement with the data given by Li *et al.* for (**A**) (table 2).<sup>[16]</sup> The Ir(1)-P(1) distance, however, appears much shorter than in Li's compound, but is very close to the distance measured in Danopoulos' compound (**B**),<sup>[8]</sup> where the N(2) atom is also substituted with a bulky aryl group. The C(2)-Ir(1)-P(1) bite angle is also very close to that of Li's complex, with a large value of  $92.70(13)^\circ$ . The structure confirms the binding of the imidazole moiety through C5 (labeled as {C2} in the structure). It also shows the presence of a hydride *trans* to the chloride, with a Ir(1)-H(100) bond length of  $1.586(19)$  Å, and a relative angle to the Cl atom of  $164(2)^\circ$ . The coordination is distorted octahedral at the iridium(III) center. All other characterisations (elemental analysis and mass spectrometry) confirmed the nature of these products.



1  
 2 **Figure 4.** An ORTEP view of compound **5**. Ellipsoids are represented at the 30%  
 3 probability level. Hydrogen atoms, except H100, cocrystallised solvent molecules and  
 4  $\text{BF}_4^-$  are omitted for clarity.

5

6 **Table 1.** Crystallographic data for compound **5**.

<b>5</b>	
Empirical formula	$\text{C}_{36}\text{H}_{44}\text{BCl}_5\text{F}_4\text{IrN}_2\text{P}$
Formula weight	991.96
Temperature (K)	180(2)
Crystal system	triclinic
Space group	P -1
$a$ (Å)	10.071(5)
$b$ (Å)	15.126(5)
$c$ (Å)	15.988(5)
$\alpha$ (°)	68.528(5)
$\beta$ (°)	74.532(5)
$\gamma$ (°)	78.030(5)
Volume (Å <sup>3</sup> )	2168.0(15)
$Z$	2
Density	1.520
Absorption coefficient. (mm <sup>-1</sup> )	3.468
$F(0\ 0\ 0)$	984
$\theta$ range for data collection	$30.5 > \theta > 1.4$
Reflections collected	57426
Independent reflections	13227 [ $R(\text{int}) = 0.0612$ ]
Goodness-of-fit on $F^2$	1.087
Final $R$ indices [ $I > 2\sigma(I)$ ]	$R_1 = 0.0481, wR_2 = 0.1217$
$R$ indices for all data	$R_1 = 0.0628, wR_2 = 0.1267$

7

8

9



1 **Table 2.** Comparison of the main structural parameters of **5** and of the iridium complexes  
 2 described by Li *et al.* (A)<sup>[16]</sup> and Danopoulos *et al.* (B)<sup>[8]</sup>.

	<b>5</b>	(A)	(B)
Bond distances (Å)			
Ir(1)–C(2) <sup>a</sup>	2.068(5)	2.065(6)	2.018(9)
Ir(1)–P(1)	2.2972(16)	2.585(1)	2.283(2)
Ir(1)–Cl(1)	2.4845(15)	2.5097(9)	–
Ir(1)–Br(1)	–	–	2.6865(11)
Ir(1)–H(100)	1.586(19)	–	–
Bond angles (°)			
C(2)–Ir(1)–P(1)	92.70(13)	92.24(16)	88.9(3)
C(2)–Ir(1)–Cl(1)	82.75(15)	85.50(16)	–
P(1)–Ir(1)–Cl(1)	86.81(5)	90.83(5)	–
C(2)–Ir(1)–H(100)	82(2)	–	–
P(1)–Ir(1)–H(100)	92(2)	–	–
Cl(1)–Ir(1)–H(100)	164(2)	–	–

3 <sup>a</sup> Labelled as C(9) in Li's complex (A).<sup>[16]</sup>

4  
 5 A <sup>31</sup>P NMR monitoring of the [Ir(COD)(μ-Cl)]<sub>2</sub>/**3** reaction at room temperature  
 6 revealed additional features. After five minutes, the ligand was completely consumed and  
 7 two products were visible: the final product at 1.1 ppm, and an intermediate compound at  
 8 19.4 ppm which quickly disappeared (Scheme 3). Our hypothesis is that, in a first step, the  
 9 phosphine coordinates to the iridium center, splitting the dimer. Subsequently, the CH  
 10 insertion occurs at the C5 position of the imidazolium ring, possibly assisted by a  
 11 chelating effect. This hypothesis is supported by the recent results obtained by Li *et al.*<sup>[16]</sup>

12 However, our observed reactivity is at variance with the results described by  
 13 Danopoulos with a similar ligand, where the mesityl substituent is replaced by DIPP (2,6-  
 14 diisopropylphenyl, compound **2**).<sup>[8]</sup> That contribution indicated that the interaction of  
 15 [Ir(COD)(μ-Cl)]<sub>2</sub> with **2**, under unspecified reaction conditions, affords the simple  
 16 addition product where the chloride ligand has however been exchanged with bromide,  
 17 [Ir(COD)(Br)(PPh<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>Im<sup>DIPP</sup>)], similar to the proposed intermediate of our reaction  
 18 leading to **5**. In the interest of checking whether a simple change of aryl group would  
 19 enforce such a big reactivity change, we decided to repeat the same reaction under  
 20 conditions similar to those leading to **4a/4b**. The reactions were carried out in two

1 different solvents (CH<sub>2</sub>Cl<sub>2</sub> or THF), always leading to complete conversion in 20 minutes  
2 at room temperature and leading to iridium complex bearing an abnormal carbene  
3 (mixture of isomers, **6a**+**6b**, Scheme 2). These compounds have been fully characterised  
4 by NMR and mass spectrometry, leaving no doubt about their nature. Indeed, the NMR  
5 spectra present many similar features to those of complexes **4a** and **4b**: the <sup>1</sup>H NMR  
6 spectrum shows four signals corresponding to the imidazole ring protons at δ 9.73, 9.71,  
7 6.35 and 6.28, with integration values of 0.5 each, for the mixture of **6a** and **6b**. We can  
8 also observe two equivalent <sup>1</sup>H NMR hydride doublets at δ -15.07 and -14.41 (figure 1c)  
9 and two signals at δ -2.04 and -3.08 in the <sup>31</sup>P NMR spectrum, corresponding to the two  
10 isomers. Moreover, the mass spectrum clearly shows two peaks at *m/z* 777.9 and 821.8,  
11 corresponding respectively to cations **6a**<sup>+</sup> and **6b**<sup>+</sup>.

12

### 13 **Conclusions**

14

15 In summary, we have described an unusual reactivity of chelating phosphine-  
16 imidazolium salts toward low oxidation state iridium complexes. In the absence of a base,  
17 we have shown that the C-H insertion at the C5 position of the imidazolium ring is the  
18 only occurring reaction, with no normal NHC observed. Contrary to related iridium  
19 chemistry previously reported by the Crabtree group,<sup>[12]</sup> this reactivity is independent of  
20 the nature of the imidazolium counteranion. It is also independent of the substitution  
21 pattern of the aryl group. In fact, the recent paper by Li *et al.*<sup>[16]</sup> shows identical chemistry  
22 for ligands bearing alkyl substituents (Me, *i*Pr) on the 3-N atom of the imidazolium ring.  
23 Thus, this reactivity pattern appears to be quite general.

24

## 1 **Experimental Section**

2  
3 All reactions were carried out under a dry argon atmosphere using Schlenk glassware and  
4 vacuum line techniques. Solvents for syntheses were dried and degassed by standard methods  
5 before use. Elemental analyses were carried out by the analytical service of the “Laboratoire  
6 de Chimie de Coordination” in Toulouse.  $^1\text{H}$  data were recorded on a Bruker AV-500  
7 spectrometer, operating at 500 MHz.  $^{13}\text{C}\{\text{H}, \text{P}\}$  and  $^{31}\text{P}\{\text{H}\}$  NMR data were recorded on a  
8 Bruker AV-500 instrument, operating at 125.8 and 202.5 MHz, respectively.  $^{19}\text{F}$  NMR data  
9 were recorded on a Bruker AC-200 instrument, operating at 188 MHz. The spectra were  
10 referenced internally using the signal from the residual protiosolvent ( $^1\text{H}$ ) or the solvent  
11 signals ( $^{13}\text{C}$ ), the signal of  $\text{CF}_3\text{CO}_2\text{H}$  (10%) in  $\text{C}_6\text{D}_6$  ( $^{19}\text{F}$ ) and externally using 85%  $\text{H}_3\text{PO}_4$  for  
12  $^{31}\text{P}$ . Mass spectra were obtained from acetonitrile or methanol solutions on a TSQ7000  
13 instrument from ThermoElectron (electrospray ionisation), and from DMSO or DMF  
14 solutions on a Nermag R10-10 instrument (FAB).  $[\text{Ir}(\text{COD})(\mu\text{-Cl})_2]$  was from Strem and used  
15 as received.  $\text{MesImEtPPh}_2^+\text{Br}^-$  **1**,  $\text{DIPP-ImEtPPh}_2^+\text{Br}^-$  **2** and  $\text{MesImEtPPh}_2^+\text{BF}_4^-$  **3** were  
16 prepared as previously described.<sup>[4]</sup>

### 17 18 **General procedure for the synthesis of Ir<sup>III</sup> complexes**

19 In a Schlenk tube,  $\text{CH}_2\text{Cl}_2$  was added to a phosphine-imidazolium salt and  $[\text{Ir}(\text{COD})\text{Cl}]_2$   
20 mixture and stirred for 30 minutes at room temperature. The solvent was removed under  
21 vacuum, the residue washed with diethyl ether until the ethereal phase remains colourless,  
22 then dried under vacuum.

23 These reactions can also be done in THF without any change in the results.

### 24 25 **Iridium chloro-1,4-cyclooctadienyl-1-(2-(diphenylphosphinoethyl) -3-(2,4,6-trimethylphe** 26 **nyl) imidazol-5-ylidene bromide (4a+4b)**

27  $\text{CH}_2\text{Cl}_2$  (3 mL),  $\text{MesImEtPPh}_2^+\text{Br}^-$  **1** (60 mg, 125.4  $\mu\text{mol}$ ) and  $[\text{Ir}(\text{COD})\text{Cl}]_2$  (46 mg, 68.5  
28  $\mu\text{mol}$ ). A pale yellow, air-sensitive product was obtained. Yield 94 mg (92%).

29  $\text{C}_{34}\text{H}_{40}\text{BrClIrN}_2\text{P}$  (815.25) Calcd for  $\text{C}_{34}\text{H}_{40}\text{BrClIrN}_2\text{P}\cdot\text{CHCl}_3$ : C 44.98, H 4.42, N 3.00; found  
30 C 45.01, H 4.35, N 2.80.

31 *First isomer:*

32  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ , 20°C):  $\delta$  = 9.74 (s, 1 H, NCHN), 7.88-7.98 (m, 2 H, *o*-CH( $\text{Ph}^2$ )),  
33 7.54-7.64 (m, 3 H, *m,p*-CH( $\text{Ph}^2$ )), 7.32-7.51 (m, 5 H, *o,m,p*-CH( $\text{Ph}^1$ )), 6.92 (br s, 2 H,

1  $CH(\text{Mes})$ ), 6.18 (d,  $^3J = 1.4$  Hz, 1 H,  $\text{MesNCH=}$ ), 5.74-5.83 (m, 1 H,  $\text{NCH}_2$ ), 5.29-5.33 (m, 1  
2 H,  $CH(\text{COD})$ ), 5.17-5.23 (m, 1 H,  $CH(\text{COD})$ ), 4.61-4.69 (m, 1 H,  $CH(\text{COD})$ ), 3.83-3.91 (m, 1  
3 H,  $\text{NCH}_2$ ), 3.73-3.81 (m, 1 H,  $CH(\text{COD})$ ), 3.37-3.55 (m, 1 H,  $\text{CH}_2\text{P}$ ), 3.01-3.12 (m, 1 H,  
4  $\text{CH}_2(\text{COD})$ ), 2.65-2.80 (m, 4 H,  $\text{CH}_2(\text{COD})$ ;  $\text{CH}_2\text{P}$ ), 2.40-2.60 (m, 2 H,  $\text{CH}_2(\text{COD})$ ), 2.25-  
5 2.40 (m + s, 4 H,  $\text{CH}_2(\text{COD})$ ;  $p\text{-CH}_3$ ), 1.90-2.12 (m + s, 7 H,  $\text{CH}_2(\text{COD})$ ;  $o\text{-CH}_3$ ), -14.55 (d,  
6  $^2J_{\text{P-H}} = 8.1$  Hz, 1 H, Ir- $H$ ).  $^{13}\text{C}$  NMR (125.8 MHz,  $\text{CDCl}_3$ , 20°C):  $\delta = 140.2$  (s,  $p\text{-C}(\text{Mes})$ ),  
7 137.1 (s,  $\text{NCHN}$ ), 135.0; 134.1 (s,  $o\text{-C}(\text{Mes})$ ), 134.5 (d,  $^2J_{\text{C-P}} = 10.4$  Hz,  $o\text{-CH}(\text{Ph}^2)$ ), 132.9 (d,  
8  $^2J_{\text{C-P}} = 7.8$  Hz,  $o\text{-CH}(\text{Ph}^1)$ ), 132.6 (d,  $^4J_{\text{C-P}} = 2$  Hz,  $p\text{-CH}(\text{Ph}^2)$ ), 131.7 (d,  $^4J_{\text{C-P}} = 2$  Hz,  $p\text{-}$   
9  $\text{CH}(\text{Ph}^1)$ ), 131.5 (s,  $\text{NC}(\text{Mes})$ ), 129.5 (d,  $^3J_{\text{C-P}} = 15$  Hz,  $m\text{-CH}(\text{Ph}^2)$ ), 129.4; 129.3 (s,  
10  $\text{CH}(\text{Mes})$ ), 128.6 (d,  $ipso\text{-C}(\text{Ph}^2)$ ), 128.1 (d,  $^3J_{\text{C-P}} = 10.6$  Hz,  $m\text{-CH}(\text{Ph}^1)$ ), 128.0 (d,  $ipso\text{-}$   
11  $\text{C}(\text{Ph}^1)$ ), 123.8 (s,  $\text{MesNCH=}$ ), 119.9 (d,  $^2J_{\text{C-P}} = 7.9$  Hz, Ir- $C$ ), 96.7 (d,  $^2J_{\text{C-P}} = 16$  Hz,  
12  $\text{CH}(\text{COD})$ ), 93.3 (d,  $^2J_{\text{C-P}} = 9.5$  Hz,  $\text{CH}(\text{COD})$ ), 91.7; 82.4 (s,  $\text{CH}(\text{COD})$ ), 45.3 (s,  $\text{NCH}_2$ ),  
13 34.1; (d,  $^3J_{\text{C-P}} = 2$  Hz,  $\text{CH}_2(\text{COD})$ ), 31.6; 30.0 (s,  $\text{CH}_2(\text{COD})$ ), 29.7 (d,  $^3J_{\text{C-P}} = 2$  Hz,  
14  $\text{CH}_2(\text{COD})$ ), 25.0 (d,  $^1J_{\text{C-P}} = 40.1$  Hz,  $\text{CH}_2\text{P}$ ), 21.1 (s,  $p\text{-CH}_3(\text{Mes})$ ), 17.9; 17.7 (s,  $o\text{-}$   
15  $\text{CH}_3(\text{Mes})$ ).  $^{31}\text{P}$  NMR (202.5 MHz,  $\text{CDCl}_3$ , 20°C):  $\delta = -2.39$  (s,  $\text{PPh}_2$ ).

16 *Second isomer:*

17  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ , 20°C):  $\delta = 9.74$  (s, 1 H,  $\text{NCHN}$ ), 7.88-7.98 (m, 2 H,  $o\text{-CH}(\text{Ph}^2)$ ),  
18 7.54-7.64 (m, 3 H,  $m,p\text{-CH}(\text{Ph}^2)$ ), 7.32-7.51 (m, 5 H,  $o,m,p\text{-CH}(\text{Ph}^1)$ ), 6.94 (br s, 2 H,  
19  $\text{CH}(\text{Mes})$ ), 6.24 (d,  $^3J = 1.4$  Hz, 1 H,  $\text{MesNCH=}$ ), 5.74-5.83 (m, 1 H,  $\text{NCH}_2$ ), 5.35-5.41 (m, 1  
20 H,  $CH(\text{COD})$ ), 5.06-5.11 (m, 1 H,  $CH(\text{COD})$ ), 4.61-4.69 (m, 1 H,  $CH(\text{COD})$ ), 4.07-4.17 (m, 1  
21 H,  $\text{NCH}_2$ ), 3.63-3.69 (m, 1 H,  $CH(\text{COD})$ ), 3.37-3.55 (m, 1 H,  $\text{CH}_2\text{P}$ ), 3.01-3.12 (m, 1 H,  
22  $\text{CH}_2(\text{COD})$ ), 2.65-2.80 (m, 4 H,  $\text{CH}_2(\text{COD})$ ;  $\text{CH}_2\text{P}$ ), 2.40-2.60 (m, 2 H,  $\text{CH}_2(\text{COD})$ ), 2.25-  
23 2.40 (m + s, 4 H,  $\text{CH}_2(\text{COD})$ ;  $p\text{-CH}_3$ ), 1.90-2.12 (m + s, 7 H,  $\text{CH}_2(\text{COD})$ ;  $o\text{-CH}_3$ ), -15.22 (d,  
24  $^2J_{\text{P-H}} = 8.3$  Hz, 1 H, Ir- $H$ ).  $^{13}\text{C}$  NMR (125.8 MHz,  $\text{CDCl}_3$ , 20°C):  $\delta = 140.2$  (s,  $p\text{-C}(\text{Mes})$ ),  
25 137.4 (s,  $\text{NCHN}$ ), 134.8; 134.2 (s,  $o\text{-C}(\text{Mes})$ ), 134.4 (d,  $^2J_{\text{C-P}} = 10.4$  Hz,  $o\text{-CH}(\text{Ph}^2)$ ), 132.8 (d,  
26  $^2J_{\text{C-P}} = 7.9$  Hz,  $o\text{-CH}(\text{Ph}^1)$ ), 133.3 (d,  $^4J_{\text{C-P}} = 11$  Hz,  $p\text{-CH}(\text{Ph}^2)$ ), 131.6 (d,  $^4J_{\text{C-P}} = 2$  Hz,  $p\text{-}$   
27  $\text{CH}(\text{Ph}^1)$ ), 131.5 (s,  $\text{NC}(\text{Mes})$ ), 129.6 (d,  $^3J_{\text{C-P}} = 15$  Hz,  $m\text{-CH}(\text{Ph}^2)$ ), 129.4; 129.3 (s,  
28  $\text{CH}(\text{Mes})$ ), 128.6 (d,  $ipso\text{-C}(\text{Ph}^2)$ ), 128.2 (d,  $^3J_{\text{C-P}} = 10.4$  Hz,  $m\text{-CH}(\text{Ph}^1)$ ), 128.0 (d,  $ipso\text{-}$   
29  $\text{C}(\text{Ph}^1)$ ), 123.5 (s,  $\text{MesNCH=}$ ), 121.1 (d,  $^2J_{\text{C-P}} = 8.1$  Hz, Ir- $C$ ), 98.6 (d,  $^2J_{\text{C-P}} = 16.2$  Hz,  
30  $\text{CH}(\text{COD})$ ), 93.8 (d,  $^2J_{\text{C-P}} = 9.1$  Hz,  $\text{CH}(\text{COD})$ ), 93.9; 83.7 (s,  $\text{CH}(\text{COD})$ ), 45.6 (s,  $\text{NCH}_2$ ),  
31 35.5; 31.0 (s,  $\text{CH}_2(\text{COD})$ ), 29.6 (d,  $^3J_{\text{C-P}} = 2$  Hz,  $\text{CH}_2(\text{COD})$ ), 28.2; (d,  $^3J_{\text{C-P}} = 2$  Hz,  
32  $\text{CH}_2(\text{COD})$ ), 24.9 (d,  $^1J_{\text{C-P}} = 39.5$  Hz,  $\text{CH}_2\text{P}$ ), 21.1 (s,  $p\text{-CH}_3(\text{Mes})$ ), 17.8; 17.6 (s,  $o\text{-}$   
33  $\text{CH}_3(\text{Mes})$ ).  $^{31}\text{P}$  NMR (202.5 MHz,  $\text{CDCl}_3$ , 20°C):  $\delta = -1.14$  (s,  $\text{PPh}_2$ ).

1 MS (FAB, mNBA matrix):  $m/z$  (%) = 627 (100) [ $C_{26}H_{28}ClIrN_2P^+$ ], 671 (92)  
2 [ $C_{26}H_{26}BrIrN_2P^+$ ], 735 (34) [ $C_{34}H_{40}ClIrN_2P^+$ ], 779 (36) [ $C_{34}H_{40}BrN_2IrP^+$ ].

3

4 **Iridium chloro-1,4-cyclooctadienyl-1-(2-(diphenylphosphinoethyl) -3-(2,4,6- trimethylph**  
5 **enyl) imidazol-5-ylidene tetrafluoroborate (5)**

6  $CH_2Cl_2$  (3 mL), MesImEtPPh<sub>2</sub><sup>+</sup>BF<sub>4</sub><sup>-</sup> **3** (60 mg, 123  $\mu$ mol) and [Ir(COD)Cl]<sub>2</sub> (43 mg, 63  
7  $\mu$ mol). A cream-white, air-sensitive product was obtained (87 mg, 86%). Suitable X-ray  
8 crystals were obtained by layer diffusion of pentane into a  $CH_2Cl_2$  solution at -20°C.

9  $C_{34}H_{40}BClF_4IrN_2P$  (822.15) Calcd for  $C_{34}H_{40}BClF_4IrN_2P \cdot 0,5 CH_2Cl_2$ : C 47.93, H 4.78, N  
10 3.24; found C 47.53, H 4.90, N 3.36.

11 <sup>1</sup>H NMR (500 MHz,  $CD_2Cl_2$ , 20°C):  $\delta$  = 8.45 (s, 1 H, NCHN), 7.99 (dd, <sup>3</sup> $J$  = 7 Hz; <sup>3</sup> $J_{P-H}$  =  
12 11.5 Hz, 2 H, *o*-CH(Ph<sup>2</sup>)), 7.63-7.72 (m, 3 H, *m,p*-CH(Ph<sup>2</sup>)), 7.43-7.57 (m, 5 H, *o,m,p*-  
13 CH(Ph<sup>1</sup>)), 7.04 (br s, 2 H, CH(Mes)), 6.40 (d, <sup>4</sup> $J$  = 1 Hz, 1 H, MesNCH=), 5.47-5.54 (m, 1 H,  
14 CH(COD)), 5.14 (dddd, <sup>2,3</sup> $J$  = 2.3; 7.6; 14.3 Hz; <sup>3</sup> $J_{P-H}$  = 27.4 Hz, 1 H, NCH<sub>2</sub>), 5.05-5.12 (m, 1  
15 H, CH(COD)), 4.73-4.77 (m, 1 H, CH(COD)), 4.12 (ddd, <sup>2,3</sup> $J$  = 11; 12 Hz; <sup>3</sup> $J_{P-H}$  = 19.6 Hz, 1  
16 H, NCH<sub>2</sub>), 3.62-3.69 (m, 1 H, CH(COD)), 3.49-3.58 (m, 1 H, CH<sub>2</sub>P), 3.02-3.12 (m, 1 H,  
17 CH<sub>2</sub>(COD)), 2.92-3.00 (m, 1 H, CH<sub>2</sub>(COD)), 2.66-2.82 (m, 3 H, CH<sub>2</sub>(COD); CH<sub>2</sub>P), 2.47-  
18 2.58 (m, 2 H, CH<sub>2</sub>(COD)), 2.32-2.42 (m + s, 4 H, CH<sub>2</sub>(COD); *p*-CH<sub>3</sub>), 2.00-2.15 (m + s, 7 H,  
19 CH<sub>2</sub>(COD); *o*-CH<sub>3</sub>), -15.12 (d, <sup>2</sup> $J_{P-H}$  = 8.3 Hz, 1 H, Ir-*H*). <sup>13</sup>C NMR (125.8 MHz,  $CD_2Cl_2$ ,  
20 20°C):  $\delta$  = 140.6 (s, *p*-C(Mes)), 135.9 (s, NCHN), 134.8; 134.4 (s, *o*-C(Mes)), 134.3 (d, <sup>2</sup> $J_{C-P}$   
21 = 7.3 Hz, *o*-CH(Ph<sup>2</sup>)), 132.8 (d, <sup>2</sup> $J_{C-P}$  = 8 Hz, *o*-CH(Ph<sup>1</sup>)), 132.6 (d, <sup>4</sup> $J_{C-P}$  = 2.3 Hz, *p*-  
22 CH(Ph<sup>2</sup>)), 131.4 (d, <sup>4</sup> $J_{C-P}$  = 2.6 Hz, *p*-CH(Ph<sup>1</sup>)), 131.3 (s, NC(Mes)), 129.5 (d, <sup>3</sup> $J_{C-P}$  = 10.6 Hz,  
23 *m*-CH(Ph<sup>2</sup>)), 129.4 (br s, CH(Mes)), 128.8 (d, <sup>1</sup> $J_{C-P}$  = 58.7 Hz, *ipso*-C(Ph<sup>1</sup>)), 128.6 (d, <sup>1</sup> $J_{C-P}$  =  
24 58.7 Hz, *ipso*-C(Ph<sup>2</sup>)), 128.1 (d, <sup>3</sup> $J_{C-P}$  = 10.5 Hz, *m*-CH(Ph<sup>1</sup>)), 124.5 (s, MesNCH=), 121.4 (d,  
25 <sup>2</sup> $J_{C-P}$  = 9.2 Hz, Ir-C), 98.7 (d, <sup>2</sup> $J_{C-P}$  = 15.6 Hz, CH(COD)), 94.6 (d, <sup>3</sup> $J_{C-P}$  = 9.4 Hz, CH(COD)),  
26 94.1; 84.4 (s, CH(COD)), 45.7 (s, NCH<sub>2</sub>), 35.6; 30.7 (s, CH<sub>2</sub>(COD)), 29.6 (d, <sup>3</sup> $J_{C-P}$  = 2.1 Hz,  
27 CH<sub>2</sub>(COD)), 28.0 (d, <sup>3</sup> $J_{C-P}$  = 3.8 Hz, CH<sub>2</sub>(COD)), 25.1 (d, <sup>1</sup> $J_{C-P}$  = 39.7 Hz, CH<sub>2</sub>P), 20.8 (s, *p*-  
28 CH<sub>3</sub>(Mes)), 17.1; 17.0 (s, *o*-CH<sub>3</sub>(Mes)). <sup>31</sup>P NMR (202.5 MHz,  $CD_2Cl_2$ , 20°C):  $\delta$  = 2.42 (s,  
29 PPh<sub>2</sub>). <sup>19</sup>F NMR (188 MHz,  $CD_2Cl_2$ , 20°C):  $\delta$  = -76.2 (s, BF<sub>4</sub>). MS (ESI, positive mode)  $m/z$   
30 (%) = 627.1 (10) [ $C_{26}H_{28}ClIrN_2P^+$ ], 735.1 (100) [ $C_{34}H_{40}ClIrN_2P^+$ ]. MS (ESI, negative mode)  
31  $m/z$  (%) = 87 (100) [BF<sub>4</sub><sup>-</sup>].

32

33 **Iridium chloro-1,4-cyclooctadienyl-1-(2-(diphenylphosphinoethyl) -3-(2,6-diisopropylphe**  
34 **nyl) imidazol-5-ylidene bromide (6a+6b)**

1 CH<sub>2</sub>Cl<sub>2</sub> (3 mL), DIPP-ImEtPPh<sub>2</sub><sup>+</sup>Br<sup>-</sup> **2** (82 mg, 157 μmol) and [Ir(COD)Cl]<sub>2</sub> (53 mg, 79  
2 μmol). A pale yellow, air-sensitive product was obtained (118 mg, 87%).

3 C<sub>37</sub>H<sub>46</sub>BrClIrN<sub>2</sub>P (857.33). Calcd for C<sub>37</sub>H<sub>46</sub>BrClIrN<sub>2</sub>P·CH<sub>2</sub>Cl<sub>2</sub>: C 48.44, H 5.13, N 2.97;  
4 found C 48.41, H 5.13, N 2.92.

5 *First isomer:*

6 <sup>1</sup>H NMR (500 MHz, CD<sub>2</sub>Cl<sub>2</sub>, 20°C): δ = 9.73 (d, <sup>4</sup>J = 1.1 Hz, 1 H, NCHN), 8.02-8.05 (m, 2  
7 H, *o*-CH(Ph<sup>2</sup>)), 7.63-7.69 (m, 3 H, *m,p*-CH(Ph<sup>2</sup>)), 7.39-7.55 (m, 5 H, *o,m,p*-CH(Ph<sup>1</sup>)), 7.32 (d,  
8 <sup>3</sup>J = 6.5 Hz, 1 H, *m*-CH(DIPP)), 7.31 (d, <sup>3</sup>J = 6.6 Hz, 1 H, *m*-CH(DIPP)), 6.28 (d, <sup>3</sup>J = 1.6 Hz,  
9 1 H, (DIPP)NCH=), 5.70-5.78 (m, 1 H, NCH<sub>2</sub>), 5.31-5.36 (m, 1 H, CH(COD)), 5.16-5.20 (m,  
10 1 H, CH(COD)), 4.67-4.75 (m, 1 H, CH(COD)), 3.89-3.97 (m, 1 H, NCH<sub>2</sub>), 3.66-3.74 (m, 1 H,  
11 CH(COD)), 3.54-3.61 (m, 1 H, CH<sub>2</sub>P), 3.00-3.09 (m, 1 H, CH<sub>2</sub>(COD)), 2.66-2.91 (m, 3 H,  
12 CH<sub>2</sub>(COD); CH<sub>2</sub>P), 2.54-2.63 (m, 1 H, CH<sub>2</sub>(COD)), 2.32-2.52 (m, 4 H, CH<sub>2</sub>(COD);  
13 CH(CH<sub>3</sub>)<sub>2</sub>), 1.90-2.12 (m, 1 H, CH<sub>2</sub>(COD)), 1.16-1.26 (m, 12 H, CH(CH<sub>3</sub>)<sub>2</sub>), -14.41 (d, <sup>2</sup>J<sub>P-H</sub> =  
14 8.2 Hz, 1 H, Ir-H). <sup>13</sup>C NMR (125.8 MHz, CD<sub>2</sub>Cl<sub>2</sub>, 20°C): δ = 146.0; 145.3 (s, *o*-C(DIPP)),  
15 137.4 (s, NCHN), 134.6 (d, <sup>2</sup>J<sub>P-C</sub> = 10.5 Hz, *o*-CH(PPh<sub>2</sub>)), 133.0 (s, <sup>2</sup>J<sub>P-C</sub> = 6.7 Hz, *o*-  
16 CH(PPh<sub>2</sub>)), 132.5 (d, <sup>4</sup>J<sub>P-C</sub> = 2.6 Hz, *p*-CH(PPh<sub>2</sub>)), 131.4 (d, <sup>4</sup>J<sub>P-C</sub> = 2.7 Hz, *p*-CH(PPh<sub>2</sub>)),  
17 131.2 (s, NC(Mes)), 131.0 (s, *p*-CH(DIPP)), 129.4 (d, <sup>3</sup>J<sub>P-C</sub> = 10.5 Hz, *m*-CH(PPh<sub>2</sub>)), 127.9 (s  
18 d, <sup>3</sup>J<sub>P-C</sub> = 10.5 Hz, *m*-CH(PPh<sub>2</sub>)), 125.5 (d, <sup>3</sup>J<sub>C-P</sub> = 8.5 Hz, (DIPP)NCH=), 124.3; 124.1 (s, *m*-  
19 CH(DIPP)), 120.1 (d, <sup>2</sup>J<sub>C-P</sub> = 8.4 Hz, Ir-C), 96.8 (d, <sup>2</sup>J<sub>P-C</sub> = 16.2 Hz, CH(COD)), 93.8 (d, <sup>2</sup>J<sub>P-C</sub>  
20 = 9.0 Hz, CH(COD)), 91.7; 82.7 (s, CH(COD)), 45.4 (s, NCH<sub>2</sub>), 33.7 (d, <sup>3</sup>J<sub>P-C</sub> = 2.1 Hz,  
21 CH<sub>2</sub>(COD)), 31.9 (s, CH<sub>2</sub>(COD)), 29.9 (d, <sup>3</sup>J<sub>P-C</sub> = 4.3 Hz, CH<sub>2</sub>(COD)), 29.7 (d, <sup>3</sup>J<sub>P-C</sub> = 2.2 Hz,  
22 CH<sub>2</sub>(COD)), 28.5; 28.4 (s, CH(CH<sub>3</sub>)<sub>2</sub>), 25.0 (d, <sup>1</sup>J<sub>C-P</sub> = 40.3 Hz, CH<sub>2</sub>P), 24.3; 24.2; 24.1; 24.0  
23 (s, CH(CH<sub>3</sub>)<sub>2</sub>). <sup>31</sup>P NMR (202.5 MHz, CD<sub>2</sub>Cl<sub>2</sub>, 20°C): δ = -3.08 (s, PPh<sub>2</sub>).

24 *Second isomer:*

25 <sup>1</sup>H NMR (500 MHz, CD<sub>2</sub>Cl<sub>2</sub>, 20°C): δ = 9.71 (d, <sup>4</sup>J = 1.2 Hz, 1 H, NCHN), 7.98-8.01 (m, 2  
26 H, *o*-CH(Ph<sup>2</sup>)), 7.63-7.69 (m, 3 H, *m,p*-CH(Ph<sup>2</sup>)), 7.39-7.55 (m, 5 H, *o,m,p*-CH(Ph<sup>1</sup>)), 7.36 (d,  
27 <sup>3</sup>J = 7.9 Hz, 1 H, *m*-CH(DIPP)), 7.35 (d, <sup>3</sup>J = 7.9 Hz, 1 H, *m*-CH(DIPP)), 6.35 (d, <sup>3</sup>J = 1.5 Hz,  
28 1 H, (DIPP)NCH=), 5.70-5.78 (m, 1 H, NCH<sub>2</sub>), 5.37-5.43 (m, 1 H, CH(COD)), 5.10-5.17 (m,  
29 1 H, CH(COD)), 4.67-4.75 (m, 1 H, CH(COD)), 4.20-4.27 (m, 1 H, NCH<sub>2</sub>), 3.76-3.83 (m, 1 H,  
30 CH(COD)), 3.38-3.43 (m, 1 H, CH<sub>2</sub>P), 3.00-3.09 (m, 1 H, CH<sub>2</sub>(COD)), 2.66-2.91 (m, 3 H,  
31 CH<sub>2</sub>(COD); CH<sub>2</sub>P), 2.54-2.63 (m, 1 H, CH<sub>2</sub>(COD)), 2.32-2.52 (m, 4 H, CH<sub>2</sub>(COD);  
32 CH(CH<sub>3</sub>)<sub>2</sub>), 1.90-2.12 (m, 1 H, CH<sub>2</sub>(COD)), 1.16-1.26 (m, 12 H, CH(CH<sub>3</sub>)<sub>2</sub>), -15.07 (d, <sup>2</sup>J<sub>P-H</sub> =  
33 8.1 Hz, 1 H, Ir-H). <sup>13</sup>C NMR (125.8 MHz, CD<sub>2</sub>Cl<sub>2</sub>, 20°C): δ = 145.9; 145.4 (s, *o*-C(DIPP)),  
34 137.6 (s, NCHN), 134.4 (d, <sup>2</sup>J<sub>P-C</sub> = 10.5 Hz, *o*-CH(PPh<sub>2</sub>)), 132.9 (s, <sup>2</sup>J<sub>P-C</sub> = 8.0 Hz, *o*-

1 CH(PPh<sub>2</sub>), 132.5 (d, <sup>4</sup>J<sub>P-C</sub> = 2.4 Hz, *p*-CH(PPh<sub>2</sub>)), 131.4 (d, <sup>4</sup>J<sub>P-C</sub> = 2.5 Hz, *p*-CH(PPh<sub>2</sub>)),  
2 131.1 (s, NC(Mes)), 131.0 (s, *p*-CH(DIPP)), 129.4 (d, <sup>3</sup>J<sub>P-C</sub> = 10.5 Hz, *m*-CH(PPh<sub>2</sub>)), 128.1 (s  
3 d, <sup>3</sup>J<sub>P-C</sub> = 10.6 Hz, *m*-CH(PPh<sub>2</sub>)), 125.3 (d, <sup>3</sup>J<sub>C-P</sub> = 8.5 Hz, (DIPP)NCH=), 124.3; 124.2 (s, *m*-  
4 CH(DIPP)), 120.2 (d, <sup>2</sup>J<sub>C-P</sub> = 8.6 Hz, Ir-C), 98.7 (d, <sup>2</sup>J<sub>P-C</sub> = 15.7 Hz, CH(COD)), 94.2 (d, <sup>2</sup>J<sub>P-C</sub>  
5 = 9.3 Hz, CH(COD)), 93.9; 84.2 (s, CH(COD)), 45.8 (s, NCH<sub>2</sub>), 35.2 (d, <sup>3</sup>J<sub>P-C</sub> = 2.1 Hz,  
6 CH<sub>2</sub>(COD)), 32.2 (s, CH<sub>2</sub>(COD)), 29.4 (d, <sup>3</sup>J<sub>P-C</sub> = 2.1 Hz, CH<sub>2</sub>(COD)), 28.6; 28.5 (s,  
7 CH(CH<sub>3</sub>)<sub>2</sub>), 28.3 (d, <sup>3</sup>J<sub>P-C</sub> = 3.9 Hz, CH<sub>2</sub>(COD)), 25.1 (d, <sup>1</sup>J<sub>C-P</sub> = 39.6 Hz, CH<sub>2</sub>P), 24.3; 24.2;  
8 24.1; 24.0 (s, CH(CH<sub>3</sub>)<sub>2</sub>). <sup>31</sup>P NMR (202.5 MHz, CD<sub>2</sub>Cl<sub>2</sub>, 20°C): δ = -2.04 (s, PPh<sub>2</sub>).  
9 MS (ESI, positive mode) *m/z* (%) = 777.9 (100) [C<sub>37</sub>H<sub>46</sub>ClIrN<sub>2</sub>P<sup>+</sup>], 821.8 (33)  
10 [C<sub>37</sub>H<sub>46</sub>BrIrN<sub>2</sub>P<sup>+</sup>].  
11

## 12 Crystallographic Data

13 A single crystal was mounted under inert perfluoropolyether on the tip of a glass fibre and  
14 cooled in the cryostream of an Bruker APEX2 diffractometer. Data were collected using the  
15 monochromatic Mo Kα radiation (λ = 0.71073). The final unit cells parameters were obtained  
16 by the least-squares refinement of a large number of selected reflections. The structures were  
17 solved by direct methods (SIR97<sup>[17]</sup>) and refined by least-squares procedures on *F*<sup>2</sup> with the  
18 SHELXL-97 program<sup>[18]</sup> using the integrated system WINGX(1.63).<sup>[19]</sup> Hydrogen atoms  
19 attached to carbon atoms were introduced at calculated positions and treated as riding on their  
20 parent atoms [d(CH) = 0.96–0.98 Å] with a displacement parameter equal to 1.2 (C<sub>6</sub>H<sub>5</sub>, CH<sub>2</sub>)  
21 or 1.5 (CH<sub>3</sub>) times that of the parent atom. The hydride atom was located in difference Fourier  
22 syntheses and its coordinates were refined using a Ir-H restraint of 1.60(2) Å. The BF<sub>4</sub> anion  
23 was partially disordered by rotation around one B-F axis, this disorder was treated using the  
24 tools available in SHELX-97<sup>[17]</sup>. Some residual electron densities were difficult to modelise  
25 and therefore, the SQUEEZE function of PLATON<sup>[20]</sup> was used to eliminate the contribution  
26 of the electron density in the solvent region from the intensity data, and the solvent-free  
27 model was employed for the final refinement. There are two cavities of 127 Å<sup>3</sup> per unit cell.  
28 PLATON estimated that each cavity contains 62 electrons which may correspond to a mixture  
29 of dichloromethane and pentane solvent molecules. The molecular views were realised with  
30 the help of ORTEP-III.<sup>[21],[22]</sup>  
31 CCDC 681170 contains the supplementary crystallographic data for this paper. These data can  
32 be obtained free of charge from The Cambridge Crystallographic Data Centre via  
33 [www.ccdc.cam.ac.uk/datarequest/cif](http://www.ccdc.cam.ac.uk/datarequest/cif).

34

## 1 Acknowledgements

2 We thank the Centre National de la Recherche Scientifique (C.N.R.S.) for support of this  
3 work, the Fonds Social Européen (F.S.E.) for a Ph.D. fellowship to Joffrey Wolf, and Yannick  
4 Coppel for N.M.R. analysis of the iridium complexes.

5

## 6 References

7

- 8 [1] D. Bourissou, O. Guerret, F. P. Gabbaï, G. Bertrand, *Chem. Rev.* **2000**, *100*, 39-91; W.  
9 A. Herrmann, *Angew. Chem. Int. Ed.* **2002**, *41*, 1290-1309; F. Glorius, *N-Heterocyclic*  
10 *Carbenes in Transition Metal Catalysis, Vol. 21*, Springer-Verlag New York, LLC,  
11 New York, **2007**; S. P. Nolan, *N-Heterocyclic Carbenes in Synthesis*, Wiley-VCH,  
12 **2006**.
- 13 [2] A. C. Hillier, H. M. Lee, E. D. Stevens, S. P. Nolan, *Organometallics* **2001**, *20*, 4246-  
14 4252; N. M. Scott, R. Dorta, E. D. Stevens, A. Correa, L. Cavallo, S. P. Nolan, *J. Am.*  
15 *Chem. Soc.* **2005**, *127*, 3516-3526; N. M. Scott, V. Pons, E. D. Stevens, D. M.  
16 Heinekey, S. P. Nolan, *Angew. Chem. Int. Ed.* **2005**, *44*, 2512-2515; R. A. Kelly, III,  
17 H. Clavier, S. Giudice, N. M. Scott, E. D. Stevens, J. Bordner, I. Samardjiev, C. D.  
18 Hoff, L. Cavallo, S. P. Nolan, *Organometallics* **2008**, *27*, 202-210.
- 19 [3] J. A. Mata, M. Poyatos, E. Peris, *Coord. Chem. Rev.* **2007**, *251*, 841-859.
- 20 [4] J. Wolf, A. Labande, J.-C. Daran, R. Poli, *J. Organomet. Chem.* **2006**, *691*, 433-443;  
21 J. Wolf, A. Labande, M. Natella, J.-C. Daran, R. Poli, *J. Mol. Catal. A: Chem.* **2006**,  
22 *259*, 205-212.
- 23 [5] J. Wolf, A. Labande, J.-C. Daran, R. Poli, *Eur. J. Inorg. Chem.* **2007**, 5069-5079.
- 24 [6] A. Labande, J.-C. Daran, E. Manoury, R. Poli, *Eur. J. Inorg. Chem.* **2007**, 1205-1209.
- 25 [7] W. A. Herrmann, C. Köcher, L. J. Gooßen, G. R. J. Artus, *Chem. Eur. J.* **1996**, *2*,  
26 1627-1636; N. Tsoureas, A. A. Danopoulos, A. A. D. Tulloch, M. E. Light,  
27 *Organometallics* **2003**, *22*, 4750-4758; H. Seo, H.-J. Park, B. Y. Kim, J. H. Lee, S. U.  
28 Son, Y. K. Chung, *Organometallics* **2003**, *22*, 618-620; T. Focken, G. Raabe, C.  
29 Bolm, *Tetrahedron: Asymmetry* **2004**, *15*, 1693-1706; H. M. Lee, P. L. Chiu, J. Y.  
30 Zeng, *Inorg. Chim. Acta* **2004**, *357*, 4313-4321; P. L. Chiu, H. M. Lee,  
31 *Organometallics* **2005**, *24*, 1692-1702; L. D. Field, B. A. Messerle, K. Q. Vuong, P.  
32 Turner, *Organometallics* **2005**, *24*, 4241-4250; M. Poyatos, A. Maise-François, S.  
33 Bellemin-Laponnaz, L. H. Gade, *Organometallics* **2006**, *25*, 2634-2641; P. L. Arnold,  
34 C. Wilson, *Inorg. Chim. Acta* **2007**, *360*, 190-196; L. H. Gade, S. Bellemin-Laponnaz,  
35 *Coord. Chem. Rev.* **2007**, *251*, 718-725; O. Kuehl, *Chem. Soc. Rev.* **2007**, *36*, 592-  
36 607; S. K. Yen, L. L. Koh, H. V. Huynh, T. S. A. Hor, *Dalton Trans.* **2007**, 3952-  
37 3958; M. Merzouk, T. Moore, N. A. Williams, *Tetrahedron Lett.* **2007**, *48*, 8914-  
38 8917; I. Özdemir, N. Gürbüz, Y. Gök, B. Çetinkaya, *Heteroat. Chem.* **2008**, *19*, 82-86;  
39 M. V. Jiménez, J. J. Pérez-Torrente, M. I. Bartolomé, V. Gierz, F. J. Lahoz, L. A. Oro,  
40 *Organometallics* **2008**, *27*, 224-234.
- 41 [8] N. Stylianides, A. A. Danopoulos, N. Tsoureas, *J. Organomet. Chem.* **2005**, *690*,  
42 5948-5958.
- 43 [9] S. Gründemann, A. Kovacevic, M. Albrecht, J. W. Faller, R. H. Crabtree, *Chem.*  
44 *Commun.* **2001**, 2274-2275.
- 45 [10] S. Gründemann, A. Kovacevic, M. Albrecht, J. W. Faller, R. H. Crabtree, *J. Am.*  
46 *Chem. Soc.* **2002**, *124*, 10473-10481.




- 1 [11] A. R. Chianese, A. Kovacevic, B. M. Zeglis, J. W. Faller, R. H. Crabtree,  
2 *Organometallics* **2004**, *23*, 2461-2468.
- 3 [12] L. N. Appelhans, D. Zuccaccia, A. Kovacevic, A. R. Chianese, H. R. Miecznikowski,  
4 A. Macchioni, E. Clot, O. Einsenstein, R. H. Crabtree, *J. Am. Chem. Soc.* **2005**, *127*,  
5 16299-16311.
- 6 [13] C. E. Ellul, M. F. Mahon, O. Saker, M. K. Whittlesey, *Angew. Chem. Int. Ed.* **2007**,  
7 *46*, 6343-6345.
- 8 [14] M. Baya, B. Eguillor, M. A. Esteruelas, M. Olivan, E. Onate, *Organometallics* **2007**,  
9 *26*, 6556-6563; B. Eguillor, M. A. Esteruelas, M. Olivan, M. Puerta, *Organometallics*  
10 **2008**, *27*, 445-450.
- 11 [15] P. L. Arnold, S. Pearson, *Coord. Chem. Rev.* **2007**, *251*, 596-609; X. Hu, I. Castro-  
12 Rodriguez, K. Meyer, *Organometallics* **2003**, *22*, 3016-3018; A. A. Danopoulos, N.  
13 Tsoureas, J. A. Wright, M. E. Light, *Organometallics* **2004**, *23*, 166-168; H. Lebel, M.  
14 K. Janes, A. B. Charette, S. P. Nolan, *J. Am. Chem. Soc.* **2004**, *126*, 5046-5047.
- 15 [16] G. Song, X. Wang, Y. Li, X. Li, *Organometallics* **2008**, *ASAP*.
- 16 [17] A. Altomare, M. C. Burla, M. Camalli, G. L. Cascarano, C. Giacovazzo, A.  
17 Guagliardi, A. G. G. Moliterni, G. Polidori, R. Spagn, *J. Appl. Cryst.* **1999**, *32*, 115-  
18 119.
- 19 [18] G. M. Sheldrick, SHELXL97, Göttingen, Germany, 1998.
- 20 [19] L. J. Farrugia, *J. Appl. Cryst.* **1999**, *32*, 837-838.
- 21 [20] A. L. Spek, *J. Appl. Cryst.* **2003**, *36*, 7-13.
- 22 [21] M. N. Burnett, C. K. Johnson, ORTEP-III\_Report ORNL-6895, Oak Ridge National  
23 Laboratory, Oak Ridge, Tennessee, 1996.
- 24 [22] L. J. Farrugia, *J. Appl. Cryst.* **1997**, *30*, 565.
- 25  
26

## 1 Graphical abstract

Phosphine-imidazolium salts add to iridium complex  $[\text{Ir}(\text{COD})(\mu\text{-Cl})_2]$  without the need of a base and lead to iridium(III) hydride complexes containing “abnormal” carbene ligands, through C-H insertion at the C5 position of the imidazolium ring.

**Keywords:** Carbenes / Phosphane ligands / Iridium / Hydrides / C-H activation

**Key Topic:** Iridium complexes of abnormal carbenes



2