



**HAL**  
open science

# Kuhn's Equivalence Theorem for Games in Product Form

Benjamin Heymann, Michel de Lara, Jean-Philippe Chancelier

► **To cite this version:**

Benjamin Heymann, Michel de Lara, Jean-Philippe Chancelier. Kuhn's Equivalence Theorem for Games in Product Form. 2021. hal-03193448v1

**HAL Id: hal-03193448**

**<https://hal.science/hal-03193448v1>**

Preprint submitted on 9 Apr 2021 (v1), last revised 12 Jul 2022 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Kuhn’s Equivalence Theorem for Games in Product Form

Benjamin Heymann\*, Michel De Lara<sup>†</sup>, Jean-Philippe Chancelier<sup>†</sup>

April 9, 2021

## Abstract

We propose an alternative to the tree representation of extensive form games. Games in product form represent information with  $\sigma$ -fields over a product set, and do not require an explicit description of the play temporality, as opposed to extensive form games on trees. This representation encompasses games with a continuum of actions, randomness and players, as well as games for which the play order cannot be determined in advance. We adapt and prove Kuhn’s theorem — regarding equivalence between mixed and behavioral strategies under perfect recall — for games in product form with continuous action sets.

**Keywords.** Games with information, Kuhn’s equivalence theorem, perfect recall, Witsenhausen intrinsic model.

## 1 Introduction

From the origin, games in extensive form have been formulated on a tree. In his seminal paper *Extensive Games and the Problem of Information* [10], Kuhn claimed that “The use of a geometrical model (...) clarifies the delicate problem of information”. The proper handling of information was thus a strong motivation for Kuhn’s extensive games. On the game tree, moves are those vertices that possess alternatives, then moves are partitioned into players moves, themselves partitioned into information sets (with the constraint that no two moves in an information set can be on the same play). Kuhn mentions agents, one agent per information set, to “personalize the interpretation” but the notion is not central (to the point that his definition of perfect recall “obviates the use of agents”). Then (pure) strategies of a player are defined as mappings from player moves to alternatives, with the property of being constant on every information set.

---

\*Criteo, Paris, France

<sup>†</sup>CERMICS, Ecole des Ponts, Marne-la-Vallée, France

By contrast, agents play a central role in the so-called Witsenhausen’s intrinsic model [15, 16]. The novelty introduced by Witsenhausen is the notion of information field, that we summarize as follows: (i) each agent is equipped with a measurable action space (set and  $\sigma$ -algebra); (ii) the product of those measurable spaces, called the hybrid space, serves as a unique domain for all the strategies (or policies in a control theoretic wording); (iii) the hybrid product  $\sigma$ -algebra hosts the agents’ information subfields, and the (pure) strategies of an agent are required to be measurable with respect to the agent’s information field.

Witsenhausen’s intrinsic model was elaborated in the control theory setting, to handle how information is distributed among agents and how it impacts their strategies. Although not explicitly designed for games, Witsenhausen’s intrinsic model had, from the start, the potential to be adapted to games. Indeed, in [15] Witsenhausen placed his own model in the context of game theory, as he made references to von Neuman and Morgenstern [14], Kuhn [10] and Aumann [3].

In this paper, we<sup>1</sup> introduce a new representation of games that we call *games in product form*. Game representations play a key role in their analysis (see the illuminating introduction of the book [2]). In the philosophy of the tree-based extensive form (Kuhn’s view), the temporal ordering is hardcoded in the tree structure: one goes from the root to the leaves, making decisions at the moves, contingent on information, chance and strategies. For Kuhn, the chronology (tree) comes first; information comes second (partition of the move vertices). By contrast, for Witsenhausen, information comes first; the chronology comes (possibly) second, under a so-called *causality* assumption contingent on the information structure [15].

Trees are perfect to follow step by step how a game is played as any strategy profile induces a unique play: one goes from the root to the leaves, passing from one node to the next by an edge that depends on the strategy profile. On the other hand, the notion of product games (W-games) does not require an explicit description of the play temporality, and the product form replaces the tree structure with a product structure, that unlocks a new set of concepts and mathematical machinery for the study of games.

We claim that games in product form display the following appealing features. Not having a hardcoded temporal ordering makes mathematical representations less constrained, hence more general. Beliefs and transition probabilities can be introduced in a unified framework, and extended to the ambiguity setting and beyond. Having a product structure enables the possibility of decomposition, agent by agent. As a first step in a broader research program (sketched in the Conclusion section), we illustrate our claim with a proof of the celebrated Kuhn’s equivalence theorem for games in product form: we show that, for W-games, perfect recall implies the equivalence between mixed and behavioral strategies; we also show the reverse implication.

The paper is organized as follows. In Sect. 2, we present a slightly extended version of Witsenhausen’s intrinsic model. Then, in Sect. 3, we propose a formal definition of games in product form (W-games), and define mixed and behavioral strategies. Finally, we derive an

---

<sup>1</sup>The paper uses the convention that the pronoun “we” refers to the authors, or the authors and the reader in the formal statements.

equivalent of Kuhn’s equivalence theorem for games in product form in Sect. 4. The proofs are relegated in Sect. 5.

In the paper, we adopt (except for the Alice and Bob models) the convention that a player is female (hence using “she” and “her”), whereas an agent is male (“he”, “his”).

## 2 Witsenhausen’s intrinsic model

In this paper, we tackle the issue of information in the context of games. For this purpose, we now present the so-called intrinsic model of Witsenhausen [16, 6]. In §2.1, we present an extended version of Witsenhausen’s intrinsic model, where we highlight the role of the configuration field that contains the information subfields of all agents. In §2.2, we illustrate, on a few examples, the ease with which one can model information in strategic contexts, using subfields of the configuration field. Finally, we present in §2.3 the notion of playability (solvability).

### 2.1 Witsenhausen’s intrinsic model (W-model)

We present an extended version of Witsenhausen’s intrinsic model, introduced some five decades ago in the control community [15, 16].

Recall that a  $\sigma$ -field (or  $\sigma$ -algebra or, shortly, field) over the set  $\mathbb{D}$  is a subset  $\mathcal{D} \subset 2^{\mathbb{D}}$ , containing  $\mathbb{D}$ , and which is closed under complementation and countable union. The trivial field over the set  $\mathbb{D}$  is the field  $\{\emptyset, \mathbb{D}\}$ . The complete field over the set  $\mathbb{D}$  is the power set  $2^{\mathbb{D}}$ . If  $\mathcal{D}$  is a  $\sigma$ -field over the set  $\mathbb{D}$  and if  $\mathbb{D}' \subset \mathbb{D}$ , then  $\mathbb{D}' \cap \mathcal{D} = \{\mathbb{D}' \cap \mathbb{D}'' \mid \mathbb{D}'' \in \mathcal{D}\}$  is a  $\sigma$ -field over the set  $\mathbb{D}'$ , called *trace field*. Consider two fields  $\mathcal{D}$  and  $\mathcal{D}'$  over the finite set  $\mathbb{D}$ . We say that the field  $\mathcal{D}$  is *finer* than the field  $\mathcal{D}'$  if  $\mathcal{D} \supset \mathcal{D}'$  (notice the reverse inclusion); we also say that  $\mathcal{D}'$  is a *subfield* of  $\mathcal{D}$ . As an illustration, the complete field is finer than any field or, equivalently, any field is a subfield of the complete field. The *least upper bound* of two fields  $\mathcal{D}$  and  $\mathcal{D}'$ , denoted by  $\mathcal{D} \vee \mathcal{D}'$ , is the smallest field that contains  $\mathcal{D}$  and  $\mathcal{D}'$ . The least upper bound of two fields is finer than any of the two. Consider a family  $(\mathcal{D}_i)_{i \in I}$ , where  $\mathcal{D}_i$  is a field over the set  $\mathbb{D}_i$ , for all  $i \in I$ . The *product field*  $\bigotimes_{i \in I} \mathcal{D}_i$  is the smallest field, over the product set  $\prod_{i \in I} \mathbb{D}_i$ , that contains all the cylinders.

**Definition 1.** (adapted from [15, 16])

A W-model is a collection  $(\mathbb{A}, (\Omega, \mathcal{F}), (\mathbb{U}_a, \mathcal{U}_a)_{a \in \mathbb{A}}, (\mathbb{J}_a)_{a \in \mathbb{A}})$ , where

- $\mathbb{A}$  is a set, whose elements are called agents;
- $\Omega$  is a set which represents “chance” or “Nature”; any  $\omega \in \Omega$  is called a state of Nature;  $\mathcal{F}$  is a  $\sigma$ -field over  $\Omega$ ;
- for any  $a \in \mathbb{A}$ ,  $\mathbb{U}_a$  is a set, the set of actions for agent  $a$ ;  $\mathcal{U}_a$  is a  $\sigma$ -field over  $\mathbb{U}_a$ ;

- for any  $a \in \mathbb{A}$ ,  $\mathcal{J}_a$  is a subfield of the following product field

$$\mathcal{J}_a \subset \mathcal{F} \otimes \bigotimes_{b \in \mathbb{A}} \mathcal{U}_b, \quad \forall a \in \mathbb{A} \quad (1)$$

and is called the information field of the agent  $a$ .

In [15, 16], the set  $\mathbb{A}$  of agents is supposed to be finite, but we have relaxed this assumption. Indeed, there is no formal difficulty in handling a general set of agents, which makes the W-model possibly relevant for differential or nonatomic games. A *finite W-model* is a W-model for which the sets  $\mathbb{A}$ ,  $\Omega$  and  $\mathcal{U}_a$ , for all  $a \in \mathbb{A}$ , are finite, and the  $\sigma$ -fields  $\mathcal{F}$  and  $\mathcal{U}_a$ , for all  $a \in \mathbb{A}$ , are the power sets (that is, the complete fields).

The *configuration space* is the product space (called *hybrid space* by Witsenhausen, hence the  $\mathbb{H}$  notation)

$$\mathbb{H} = \Omega \times \prod_{a \in \mathbb{A}} \mathcal{U}_a \quad (2a)$$

equipped with the product *configuration field*

$$\mathcal{H} = \mathcal{F} \otimes \bigotimes_{a \in \mathbb{A}} \mathcal{U}_a. \quad (2b)$$

A *configuration*  $h \in \mathbb{H}$  is denoted by

$$h = (\omega, (u_a)_{a \in \mathbb{A}}) \iff h_\emptyset = \omega \text{ and } h_a = u_a, \quad \forall a \in \mathbb{A}. \quad (2c)$$

Now, we introduce the notion of pure W-strategy.

**Definition 2.** ([15, 16]) A pure W-strategy of agent  $a \in \mathbb{A}$  is a mapping

$$\lambda_a : (\mathbb{H}, \mathcal{H}) \rightarrow (\mathcal{U}_a, \mathcal{U}_a) \quad (3a)$$

from configurations to actions, which is measurable with respect to the information field  $\mathcal{J}_a$  of agent  $a$ , that is,

$$\lambda_a^{-1}(\mathcal{U}_a) \subset \mathcal{J}_a. \quad (3b)$$

We denote by  $\Lambda_a$  the set of all pure W-strategies of agent  $a \in \mathbb{A}$ . A pure W-strategies profile  $\lambda$  is a family

$$\lambda = (\lambda_a)_{a \in \mathbb{A}} \in \prod_{a \in \mathbb{A}} \Lambda_a \quad (4a)$$

of pure W-strategies, one per agent  $a \in \mathbb{A}$ . The set of pure W-strategies profiles is

$$\Lambda = \prod_{a \in \mathbb{A}} \Lambda_a. \quad (4b)$$

Condition (3b) expresses the property that any (pure)  $W$ -strategy of agent  $a$  may only depend upon the information  $\mathcal{J}_a$  available to  $a$ .

The following self-explanatory notations (for  $B \subset \mathbb{A}$ ) will be useful:

$$\mathbb{U}_B = \prod_{b \in B} \mathbb{U}_b, \quad (5a)$$

$$\mathcal{U}_B = \bigotimes_{b \in B} \mathcal{U}_b \otimes \bigotimes_{a \notin B} \{\emptyset, \mathbb{U}_a\} \subset \bigotimes_{a \in \mathbb{A}} \mathcal{U}_a, \quad (5b)$$

$$\mathcal{H}_B = \mathcal{F} \otimes \mathcal{U}_B = \mathcal{F} \otimes \bigotimes_{b \in B} \mathcal{U}_b \otimes \bigotimes_{a \notin B} \{\emptyset, \mathbb{U}_a\} \subset \mathcal{H}, \quad (5c)$$

$$\text{(when } B \neq \emptyset) \quad h_B = (h_b)_{b \in B} \in \prod_{b \in B} \mathbb{U}_b, \quad \forall h \in \mathbb{H}, \quad (5d)$$

$$\text{(when } B \neq \emptyset) \quad \pi_B : \mathbb{H} \rightarrow \prod_{b \in B} \mathbb{U}_b, \quad h \mapsto h_B, \quad (5e)$$

$$\text{(when } B \neq \emptyset) \quad \lambda_B = (\lambda_b)_{b \in B} \in \prod_{b \in B} \Lambda_b, \quad \forall \lambda \in \Lambda. \quad (5f)$$

In (5b), when  $B = \{a\}$  is a singleton, we will sometimes (abusively) identify  $\mathcal{U}_{\{a\}} = \mathcal{U}_a \otimes \bigotimes_{b \neq a} \{\emptyset, \mathbb{U}_b\} \subset \mathcal{U}_{\mathbb{A}}$  with  $\mathcal{U}_a$ .

## 2.2 Examples

We illustrate, on a few examples, the ease with which one can model information in strategic contexts, using subfields of the configuration field.

**Alice and Bob models.** To illustrate the  $W$ -formalism presented above in §2.1, we give here three examples with two agents, Alice and Bob (who can belong either to the same player or to two different players)<sup>2</sup>: first, acting simultaneously (Figure 1i); second, one acting after another (Figure 1ii) ; third acting after the Nature's move (Figure 1iii).

*Alice and Bob as unordered agents (trivial information, Figures 1i and 2).* In the simplest model, we consider two agents  $a$  (Alice) and  $b$  (Bob) having two possible actions each (top  $T$  and bottom  $B$  for Alice  $a$ , left  $L$  and right  $R$  for Bob  $b$ ), that is,

$$\mathbb{U}_a = \{u_T, u_B\}, \quad \mathbb{U}_b = \{u_L, u_R\}. \quad (6a)$$

We also suppose that Alice and Bob have no information about each other's actions — see Figure 2 where are represented the (here trivial) atoms (that is, the minimal elements for the inclusion order) of the finite  $\sigma$ -fields  $\mathcal{J}_a$  and  $\mathcal{J}_b$  — that is,  $\mathcal{J}_a = \mathcal{J}_b = \{\emptyset, \mathbb{U}_a\} \otimes \{\emptyset, \mathbb{U}_b\}$ ,

---

<sup>2</sup>For the Alice and Bob models, we do not follow the convention that a player is female, whereas an agent is male.




Figure 1: Alice and Bob examples in the tree model

which can be interpreted as Alice and Bob acting simultaneously. As Nature is absent, the configuration space consists of four elements

$$\mathbb{H} = \mathbb{U}_a \times \mathbb{U}_b = \{u_T, u_B\} \times \{u_L, u_R\}, \quad (6b)$$

hence the square in Figure 2.




Figure 2: Atoms of the information fields of the agents  $a$  and  $b$  acting simultaneously (case of Figure 1i)

*Alice and Bob as ordered agents (without Nature, Figures 1ii and 3).*

As in the previous example, Nature is absent, and there are two agents  $a$  (Alice) and  $b$  (Bob), having two possible actions each (see (6a)), so that the configuration space consists of four elements (see (6b)). Suppose that Bob's information field is trivial (Bob knows nothing of Alice's actions), that is,

$$\mathcal{J}_b = \{\emptyset, \{u_T, u_B\}\} \otimes \{\emptyset, \{u_L, u_R\}\},$$

and that Alice knows what Bob does (Alice can distinguish between  $u_L$  and  $u_R$ )

$$\mathcal{J}_a = \{\emptyset, \{u_T, u_B\}\} \otimes \{\emptyset, \{u_L\}, \{u_R\}, \{u_L, u_R\}\}.$$

In this example, the agents are naturally ordered: Bob plays first, Alice plays second. Had the order been inverted, then there would have been a sort of paradox – Alice would play first, before Bob, and would know Bob's action that has not been yet taken by him.




Figure 3: Atoms of the information fields of the ordered agents  $a$  and  $b$ , without Nature (case of Figure 1ii)

*Alice and Bob as ordered agents (with Nature, Figures 1iii and 4).*

In this example, there are two agents  $a$  (Alice) and  $b$  (Bob) and two states of Nature  $\Omega = \{\omega^+, \omega^-\}$  (say, heads or tails). As in the previous examples, agents have two possible actions each (see (6a)). Thus, the configuration space consists of eight elements

$$\mathbb{H} = \{\omega^+, \omega^-\} \times \{u_T, u_B\} \times \{u_L, u_R\},$$

hence the cube in Figure 4. We consider the following information structure:

$$\mathcal{J}_b = \overbrace{\{\emptyset, \{\omega^+\}, \{\omega^-\}, \{\omega^+, \omega^-\}\}}^{\text{Bob knows Nature's move}} \otimes \overbrace{\{\emptyset, \{u_T, u_B\}\}}^{\text{Bob does not know what Alice does}} \otimes \{\emptyset, \mathbb{U}_b\}, \quad (7a)$$

$$\mathcal{J}_a = \overbrace{\{\emptyset, \{\omega^+\}, \{\omega^-\}, \{\omega^+, \omega^-\}\}}^{\text{Alice knows Nature's move}} \otimes \{\emptyset, \mathbb{U}_a\} \otimes \overbrace{\{\emptyset, \{u_L\}, \{u_R\}, \{u_L, u_R\}\}}^{\text{Alice knows what Bob does}}. \quad (7b)$$

Again, here agents are naturally ordered: Bob plays first, Alice plays second.




Figure 4: Atoms of the information fields of the ordered agents  $a$  and  $b$ , with Nature (case of Figure 1iii)



**Sequential decision-making.** Suppose a player makes her actions (say, an element of  $\mathbb{R}^n$ ) at every discrete time step in the set<sup>3</sup>  $\llbracket 0, T-1 \rrbracket$ , where  $T \geq 1$  is an integer. The situation will be modeled with (possibly) Nature set and field  $(\Omega, \mathcal{F})$ , and with  $T$  agents in  $\mathbb{A} = \llbracket 0, T-1 \rrbracket$ , and their corresponding sets,  $\mathbb{U}_t = \mathbb{R}^n$ , and fields,  $\mathcal{U}_t = \mathcal{B}_{\mathbb{R}^n}$  (the Borel  $\sigma$ -field of  $\mathbb{R}^n$ ), for  $t \in \mathbb{A}$ . Then, one builds up the product set  $\mathbb{H} = \Omega \times \prod_{t=0}^{T-1} \mathbb{U}_t$  and the product field  $\mathcal{H} = \mathcal{F} \otimes \bigotimes_{t=0}^{T-1} \mathcal{U}_t$ . Every agent  $t \in \llbracket 0, T-1 \rrbracket$  is equipped with an information field  $\mathcal{J}_t \subset \mathcal{H}$ . Then, we show how we can express four information patterns: sequentiality, memory of past information, memory of past actions, perfect recall. Following the notation (5b), we set  $\mathcal{U}_{\{0, \dots, t-1\}} = \bigotimes_{s=0}^{t-1} \mathcal{U}_s \otimes \bigotimes_{s=t}^{T-1} \{\emptyset, \mathbb{U}_s\}$  for  $t \in \llbracket 1, T \rrbracket$ . The inclusions  $\mathcal{J}_t \subset \mathcal{H}_{\{0, \dots, t-1\}} = \mathcal{F} \otimes \mathcal{U}_{\{0, \dots, t-1\}}$ , for  $t \in \llbracket 0, T-1 \rrbracket$ , express that every agent can remember no more than his past actions (sequentiality); memory of past information is represented by the inclusions  $\mathcal{J}_{t-1} \subset \mathcal{J}_t$ , for  $t \in \llbracket 1, T-1 \rrbracket$ ; memory of past actions is represented by the inclusions  $\{\emptyset, \Omega\} \otimes \mathcal{U}_{\{0, \dots, t-1\}} \subset \mathcal{J}_t$ , for  $t \in \llbracket 1, T-1 \rrbracket$ ; perfect recall is represented by the inclusions  $\mathcal{J}_{t-1} \vee (\{\emptyset, \Omega\} \otimes \mathcal{U}_{\{0, \dots, t-1\}}) \subset \mathcal{J}_t$ , for  $t \in \llbracket 1, T-1 \rrbracket$ .

To represent  $N$  players — where each player  $p$  makes a sequence of actions, one for each period  $t \in \llbracket 0, T^p-1 \rrbracket$  — we use  $\prod_{p=1}^N T^p$  agents, labeled by  $(p, t) \in \bigcup_{q=1}^N (\{q\} \times \llbracket 0, T^q-1 \rrbracket)$ . With obvious notations, the inclusions  $\mathcal{J}_{(p, t-1)} \subset \mathcal{J}_{(p, t)}$  express memory of one's own past information, whereas (with obvious notation) the inclusions  $\bigvee_{q=1}^N \{\emptyset, \Omega\} \otimes \bigotimes_{s=0}^{t-1} \mathcal{U}_s \otimes \bigotimes_{s=t}^{T^q-1} \{\emptyset, \mathbb{U}_s^q\} \subset \mathcal{J}_{(p, t)}$  express memory of all players past actions.

**Principal-Agent models.** A branch of Economics studies so-called *Principal-Agent* models with two decision-makers (agents) — the Principal  $\text{Pr}$  (*leader*) who takes actions  $u_{\text{Pr}} \in \mathbb{U}_{\text{Pr}}$ , where the set  $\mathbb{U}_{\text{Pr}}$  is equipped with a  $\sigma$ -field  $\mathcal{U}_{\text{Pr}}$ , and the Agent  $\text{Ag}$  (*follower*) who takes actions  $u_{\text{Ag}} \in \mathbb{U}_{\text{Ag}}$ , where the set  $\mathbb{U}_{\text{Ag}}$  is equipped with a  $\sigma$ -field  $\mathcal{U}_{\text{Ag}}$  — and with Nature, corresponding to *private information (or type)* of the Agent  $\text{Ag}$ , taking values in a set  $\Omega$ , equipped with a  $\sigma$ -field  $\mathcal{F}$ .

*Hidden type* (leading to adverse selection or to signaling) is represented by any information structure with the property that, on the one hand,

$$\mathcal{J}_{\text{Pr}} \subset \underbrace{\{\emptyset, \Omega\}}_{\text{Ag type not observed}} \otimes \underbrace{\{\emptyset, \mathbb{U}_{\text{Pr}}\}}_{\text{Ag's action possibly observed}} \otimes \mathcal{U}_{\text{Ag}}, \quad (8)$$

that is, the Principal  $\text{Pr}$  does not know the Agent  $\text{Ag}$  type, but can possibly observe the Agent  $\text{Ag}$  action, and, on the other hand, that

$$\underbrace{\mathcal{F}}_{\text{known type}} \otimes \{\emptyset, \mathbb{U}_{\text{Pr}}\} \otimes \{\emptyset, \mathbb{U}_{\text{Ag}}\} \subset \mathcal{J}_{\text{Ag}}, \quad (9)$$

that is, the Agent  $\text{Ag}$  knows the state of nature (his type).

*Hidden action* (leading to moral hazard) is represented by any information structure with the property that, on the one hand,

$$\mathcal{J}_{\text{Pr}} \subset \underbrace{\mathcal{F}}_{\text{possibly knows Ag type}} \otimes \{\emptyset, \mathbb{U}_{\text{Pr}}\} \otimes \underbrace{\{\emptyset, \mathbb{U}_{\text{Ag}}\}}_{\text{cannot observe Ag's action}}, \quad (10)$$

---

<sup>3</sup>For any integers  $a \leq b$ ,  $\llbracket a, b \rrbracket$  denotes the subset  $\{a, a+1, \dots, b-1, b\}$ .

that is, the Principal **Pr** does not know the Agent **Ag** action, but can possibly observe the Agent **Ag** type and, on the other hand, that the inclusion (9) holds true, that is, the agent **Ag** knows the state of nature (his type).

**Stackelberg leadership model.** In Stackelberg games, the leader **Pr** takes an action  $u_{\text{Pr}} \in \mathbb{U}_{\text{Pr}}$  — based at most upon the (full or partial) observation of the state  $\omega \in \Omega$  of Nature — and the the follower **Ag** takes an action  $u_{\text{Ag}} \in \mathbb{U}_{\text{Ag}}$  — based at most upon the (full or partial) observation of the state of Nature  $\omega \in \Omega$ , and upon the leader action  $u_{\text{Pr}} \in \mathbb{U}_{\text{Pr}}$ . This kind of information structure is expressed with the following inclusions of fields:

$$\mathcal{J}_{\text{Pr}} \subset \mathcal{F} \otimes \{\emptyset, \mathbb{U}_{\text{Pr}}\} \otimes \{\emptyset, \mathbb{U}_{\text{Ag}}\} \text{ and } \mathcal{J}_{\text{Ag}} \subset \mathcal{F} \otimes \mathbb{U}_{\text{Pr}} \otimes \{\emptyset, \mathbb{U}_{\text{Ag}}\}. \quad (11)$$

Even if the players are called leader and follower, there is no explicit chronology in the model, but just the information structure (11). It is the information structure that reveals the chronology. Indeed, if we label the leader **Pr** as  $t = 0$  (first player) and the follower **Ag** as  $t = 1$  (second player), the inclusions (11) become the inclusions  $\mathcal{J}_0 \subset \mathcal{F} \otimes \{\emptyset, \mathbb{U}_0\} \otimes \{\emptyset, \mathbb{U}_1\}$ , and  $\mathcal{J}_1 \subset \mathcal{F} \otimes \mathbb{U}_0 \otimes \{\emptyset, \mathbb{U}_1\}$ : the sequence  $\mathcal{J}_0, \mathcal{J}_1$  of information fields is “adapted” to the filtration  $\mathcal{F} \otimes \{\emptyset, \mathbb{U}_0\} \otimes \{\emptyset, \mathbb{U}_1\} \subset \mathcal{F} \otimes \mathbb{U}_0 \otimes \{\emptyset, \mathbb{U}_1\}$ . But if we label the leader **Pr** as  $t = 1$  and the follower **Ag** as  $t = 0$ , the new sequence of information fields would not be “adapted” to the new filtration. It is the information structure (11) that prevents the follower to play first, but that makes possible the leader to play first and the follower to play second.

## 2.3 Playability (solvability)

Regarding the Kuhn tree formulation, Witsenhausen says that “For any combination of policies one can find the corresponding outcome by following the tree along selected branches, and this is an explicit procedure” [15]. In the Witsenhausen product formulation, there is no such explicit procedure as, for any combination of policies, there may be none, one or many solutions to the closed-loop equations; these equations express the action of one agent as the output of his strategy, supplied with Nature outcome and with all agents actions. This is why Witsenhausen needs a property that he called *solvability* in [15], whereas Kuhn does not need it as it is hardcoded in the tree structure. From now on, we will no longer use the terminology of Witsenhausen and we will use *playability* and *playable*, where he used solvability and solvable. We indeed think that such vocabulary is more telling to a game theory audience.

### 2.3.1 Playability

With any given pure W-strategies profile  $\lambda = (\lambda_a)_{a \in \mathbb{A}} \in \prod_{a \in \mathbb{A}} \Lambda_a$  we associate the set-valued mapping (correspondence)

$$\mathcal{M}_\lambda : \Omega \rightrightarrows \prod_{b \in \mathbb{A}} \mathbb{U}_b, \quad \omega \mapsto \left\{ (u_b)_{b \in \mathbb{A}} \in \prod_{b \in \mathbb{A}} \mathbb{U}_b \mid u_a = \lambda_a(\omega, (u_b)_{b \in \mathbb{A}}), \quad \forall a \in \mathbb{A} \right\}, \quad (12a)$$

as well as the *solution correspondence*  $\mathcal{S}_\lambda = (I_\Omega, \mathcal{M}_\lambda)$  given by (see notation (2c) for  $h_\emptyset$ )

$$\mathcal{S}_\lambda : \Omega \rightrightarrows \mathbb{H} , \quad \omega \mapsto \left\{ h \in \mathbb{H} \mid h_\emptyset = \omega \text{ and } h_a = \lambda_a(h_\emptyset, (h_b)_{b \in \mathbb{A}}) = \lambda_a(h) , \quad \forall a \in \mathbb{A} \right\} . \quad (12b)$$

With this definition, we slightly reformulate below how Witsenhausen introduced the property of playability.

**Definition 3.** ([15, 16]) *The playability property holds true for the W-model of Definition 1 when, for any pure W-strategies profile  $\lambda = (\lambda_a)_{a \in \mathbb{A}} \in \prod_{a \in \mathbb{A}} \Lambda_a$ , the set-valued mapping  $\mathcal{M}_\lambda$  in (12a) is a mapping whose domain is  $\Omega$ , that is, the cardinal of  $\mathcal{M}_\lambda(\omega)$  is equal to one, for any state of nature  $\omega \in \Omega$ .*

*Thus, under the playability property, for any state of nature  $\omega \in \Omega$ , there exists one, and only one, action profile  $(u_b)_{b \in \mathbb{A}} \in \prod_{b \in \mathbb{A}} \mathbb{U}_b$  which is a solution of the closed-loop equations*

$$u_a = \lambda_a(\omega, (u_b)_{b \in \mathbb{A}}) , \quad \forall a \in \mathbb{A} . \quad (13)$$

*In this case, we define the solution map*

$$S_\lambda : \Omega \rightarrow \mathbb{H} \quad (14a)$$

*as the unique element contained in the set  $\mathcal{S}_\lambda(\omega)$  in (12b), that is,  $\mathcal{S}_\lambda(\omega) = \{S_\lambda(\omega)\}$  or, equivalently, by, for all  $h \in \mathbb{H}$  and using notation (2c),*

$$S_\lambda(\omega) = h \iff \begin{cases} h_\emptyset &= \omega \\ h_a &= \lambda_a(h) , \quad \forall a \in \mathbb{A} . \end{cases} \quad (14b)$$

We now present some useful properties of playable W-models. Let a W-model be playable, let  $\lambda = (\lambda_a)_{a \in \mathbb{A}} \in \prod_{a \in \mathbb{A}} \Lambda_a$  be a pure W-strategies profile like in (4a), and let  $B \subset \mathbb{A}$  be a nonempty subset of agents. From (14b), we readily get that

$$\pi_B(S_\lambda(\omega)) = \lambda_B(S_\lambda(\omega)) , \quad \forall \omega \in \Omega , \quad (15)$$

where the projection  $\pi_B$  is defined in Equation (5e) and  $\lambda_B$  is defined in Equation (5f). Now, we examine what happens when we replace some of the W-strategies  $\lambda_a$  by constant ones. For this purpose, for any subset  $B \subset \mathbb{A}$  of agents, we introduce the *partial solution map*  $\widehat{S}_{\lambda_{-B}}^B$ , defined by

$$\widehat{S}_{\lambda_{-B}}^B(\omega, u_B) = S_{u_B, \lambda_{-B}}(\omega) , \quad \forall \omega \in \Omega , \quad \forall u_B \in \mathbb{U}_B , \quad (16)$$

where  $(u_B, \lambda_{-B})$  has to be understood as the pure W-strategies profile made of two subprofiles, like in (5f), namely constant subprofile with values  $u_B$  and subprofile  $\lambda_{-B} = (\lambda_c)_{c \notin B} \in \prod_{c \notin B} \Lambda_c$ .

We obtain the following result, as a straightforward application of (14)–(15)–(16).

**Proposition 4.** *Let a W-model be playable, as in Definition 3. For any subset  $B \subset \mathbb{A}$  of agents, the solution map  $S_\lambda$  in (14) and the partial solution map  $\widehat{S}_{\lambda_{-B}}^B$  in (16) are related as follows:*

$$S_\lambda(\omega) = S_{\lambda_B, \lambda_{-B}}(\omega) = \widehat{S}_{\lambda_{-B}}^B(\omega, \pi_B(S_\lambda(\omega))) = \widehat{S}_{\lambda_{-B}}^B(\omega, \lambda_B(S_\lambda(\omega))), \quad \forall \omega \in \Omega. \quad (17)$$

*As a consequence, for any two pure W-strategies profile  $\lambda$  and  $\lambda'$  which are such that  $\lambda_{-B} = \lambda'_{-B}$ , we have that  $\widehat{S}_{\lambda_{-B}}^B = \widehat{S}_{\lambda'_{-B}}^B$  and that*

$$\pi_B(S_\lambda(\omega)) = \pi_B(S_{\lambda'}(\omega)) \implies S_\lambda(\omega) = S_{\lambda'}(\omega), \quad \forall \omega \in \Omega. \quad (18)$$

Here is a nice application of property (17), that will be useful in the proof of Kuhn's equivalence Theorem (Lemma 21).

**Proposition 5.** *Let a W-model be playable, as in Definition 3. Let  $a \in \mathbb{A}$  be an agent, and  $Z : (\mathbb{H}, \mathcal{J}_a) \rightarrow (\mathbb{Z}, \mathcal{Z})$  be a measurable mapping, where  $\mathbb{Z}$  is a set (not to be taken in the sense of relative integers) and where the  $\sigma$ -field  $\mathcal{Z}$  contains the singletons. Then, for any pair  $\lambda = (\lambda_b)_{b \in \mathbb{A}}$  and  $\lambda' = (\lambda'_b)_{b \in \mathbb{A}}$  of W-strategy profiles such that  $b \neq a \implies \lambda_b = \lambda'_b$ , we have that  $Z \circ S_\lambda = Z \circ S_{\lambda'}$ .*

*Proof.* The proof is by contradiction. Let  $\lambda = (\lambda_b)_{b \in \mathbb{A}}$  and  $\lambda' = (\lambda'_b)_{b \in \mathbb{A}}$  be a pair of W-strategy profiles such that  $b \neq a \implies \lambda_b = \lambda'_b$ , and suppose that there exists  $\omega \in \Omega$  such that  $Z(S_\lambda(\omega)) \neq Z(S_{\lambda'}(\omega))$ .

Consider  $H = Z^{-1}(Z(S_\lambda(\omega))) \subset \mathbb{H}$ . By definition of the subset  $H$  and by property of  $\omega \in \Omega$ , we get that  $S_\lambda(\omega) \in H$  and  $S_{\lambda'}(\omega) \notin H$ . Moreover,  $H \in \mathcal{J}_a$  since  $Z : (\mathbb{H}, \mathcal{J}_a) \rightarrow (\mathbb{Z}, \mathcal{Z})$  is a measurable mapping and the  $\sigma$ -field  $\mathcal{Z}$  contains the singletons. We define a new W-strategy  $\lambda''_a$  for agent  $a$  as follows:

$$\forall h'' \in \mathbb{H}, \quad \lambda''_a(h'') = \begin{cases} \pi_a(S_\lambda(\omega)) & \text{if } h'' \notin H, \\ \pi_a(S_{\lambda'}(\omega)) & \text{if } h'' \in H. \end{cases} \quad (19)$$

Thus defined, the mapping  $\lambda''_a$  indeed is a W-strategy because, as  $H \in \mathcal{J}_a$ , the mapping  $\lambda''_a : (\mathbb{H}, \mathcal{J}_a) \rightarrow (\mathbb{U}_a, \mathcal{U}_a)$  is measurable. We define the W-strategies profile  $\lambda'' = (\lambda''_b)_{b \in \mathbb{A}}$  by completing  $\lambda''_a$  with  $\lambda''_b = \lambda_b = \lambda'_b$  when  $b \neq a$ .

We prove that solvability fails for the W-strategy profile  $\lambda''$  (hence the contradiction). For this purpose, we consider the following only two possibilities for  $S_{\lambda''}(\omega)$ , depending whether it belongs or not to  $H$ .

First, we assume that  $S_{\lambda''}(\omega) \notin H$ . Then, we have that

$$\begin{aligned} \pi_a(S_{\lambda''}(\omega)) &= \lambda''_a(S_{\lambda''}(\omega)) && \text{(by (15))} \\ &= \pi_a(S_\lambda(\omega)). && \text{(by the first case of (19) as } S_{\lambda''}(\omega) \notin H \text{ by assumption)} \end{aligned}$$

Using Implication (18) with the W-strategies profiles  $\lambda''$  and  $\lambda$  and with the subset  $B = \{a\}$ , we get that  $S_{\lambda''}(\omega) = S_\lambda(\omega)$ . Therefore, as  $S_\lambda(\omega) \in H$ , we deduce that  $S_{\lambda''}(\omega) \in H$ , which contradicts the assumption that  $S_{\lambda''}(\omega) \notin H$ .

Second, we assume that  $S_{\lambda''}(\omega) \in H$ . Then, we have that

$$\begin{aligned} \pi_a(S_{\lambda''}(\omega)) &= \lambda''_a(S_{\lambda''}(\omega)) && \text{(by (15))} \\ &= \pi_a(S_{\lambda'}(\omega)) . && \text{(by the second case of (19) as } S_{\lambda''}(\omega) \in H \text{ by assumption)} \end{aligned}$$

Using Implication (18) with the W-strategies profiles  $\lambda''$  and  $\lambda'$  and with the subset  $B = \{a\}$ , we get that  $S_{\lambda''}(\omega) = S_{\lambda'}(\omega)$ . Therefore, as  $S_{\lambda'}(\omega) \notin H$ , we deduce that  $S_{\lambda''}(\omega) \notin H$ , which contradicts the assumption that  $S_{\lambda''}(\omega) \in H$ .

We obtain a contradiction and conclude that  $Z(S_{\lambda}(\omega)) = Z(S_{\lambda'}(\omega))$ .

This ends the proof.  $\square$

Witsenhausen introduced the notion of solvable measurable property in [15] when the solution map is measurable. We will need a stronger definition.

**Definition 6.** *Let  $B \subset \mathbb{A}$  be a nonempty subset of agents. We say that a playable W-model is  $B$ -strongly measurable if, for any subset  $B' \subset B$ , the partial solution map in (16) is a measurable mapping  $\widehat{S}_{\lambda_{-B'}}^{B'} : (\Omega \times \mathbb{U}_{B'}, \mathcal{F} \otimes \mathcal{U}_{B'}) \rightarrow (\mathbb{H}, \mathcal{H})$ , for any pure W-strategies profile  $\lambda = (\lambda_a)_{a \in \mathbb{A}} \in \prod_{a \in \mathbb{A}} \Lambda_a$  like in (4a) .*

Of course, a playable finite W-model is always  $B$ -strongly measurable, for any nonempty subset  $B \subset \mathbb{A}$  of agents.

### 2.3.2 An example of a playable game that cannot be framed on a tree

Witsenhausen defines the notion of causality and proves in [15] that causality implies playability. The reverse, however, is not true. In [15, Theorem 2], Witsenhausen exhibits an example of noncausal W-model that is playable. The construction relies on three agents with binary action sets –hence  $\mathbb{A} = \{a, b, c\}$ ,  $\mathbb{U}_a = \mathbb{U}_b = \mathbb{U}_c = \{0, 1\}$ – and Nature does not play any role – so that  $\mathbb{H} = \{0, 1\}^3$ . The example (see Figure 5) relies on a choice of information fields so that (i) no information fields is trivial — which means that there is no first agent — , (ii) the W-model is playable though. The triplet of information fields (where  $\sigma$  denotes here the  $\sigma$ -field generated by a measurable mapping)  $\mathcal{J}_a = \sigma(u_b(1 - u_c))$  ,  $\mathcal{J}_b = \sigma(u_c(1 - u_a))$  ,  $\mathcal{J}_c = \sigma(u_a(1 - u_b))$  clearly satisfies (i). Let us show that playability holds. First we observe that the W-strategies can be written as

$$\lambda_a(u_a, u_b, u_c) = \widetilde{\lambda}_a(u_b(1 - u_c)), \quad \lambda_b(u_a, u_b, u_c) = \widetilde{\lambda}_b(u_c(1 - u_a)), \quad \lambda_c(u_a, u_b, u_c) = \widetilde{\lambda}_c(u_a(1 - u_b)),$$

where  $\widetilde{\lambda} : \{0, 1\} \rightarrow \{0, 1\}$ , hence  $(\widetilde{\lambda}_a, \widetilde{\lambda}_b, \widetilde{\lambda}_c) \in \{\text{Id}, 1 - \text{Id}\}^3$  (Id denotes the identity mapping). From there, we check that playability holds true, with the (constant) solution map given by

$$S_{(\text{Id}, \text{Id}, \text{Id})} = (0, 0, 0), \quad S_{(1 - \text{Id}, \text{Id}, \text{Id})} = (1, 0, 1), \quad S_{(1 - \text{Id}, 1 - \text{Id}, \text{Id})} = (0, 1, 0), \quad S_{(1 - \text{Id}, 1 - \text{Id}, 1 - \text{Id})} = (1, 1, 1).$$

Hence the W-model is noncausal (because there is no first agent) but playable.

Witsenhausen's intrinsic model deals with agents, information and strategies, but not with players and preferences. We now turn to extending the Witsenhausen's intrinsic model to games.




Figure 5: Noncausal playable W-model: information partitions of the three agents

### 3 Games in product form

We are now ready to embed Witsenhausen's intrinsic model into game theory. In §3.1, we introduce a formal definition of a game in product form (W-game). In §3.2, we define mixed and behavioral strategies in the spirit of Aumann [3].

#### 3.1 Definition of a game in product form (W-game)

We introduce a formal definition of a game in product form (W-game).

**Definition 7.** A W-game  $\left( ((\mathbb{A}^p)_{p \in P}, (\Omega, \mathcal{F}), (\mathbb{U}_a, \mathcal{U}_a, \mathcal{J}_a)_{a \in \mathbb{A}}), (\preceq^p)_{p \in P} \right)$ , or a game in product form, is made of

- a set  $\mathbb{A}$  of agents with a partition  $(\mathbb{A}^p)_{p \in P}$ , where  $P$  is the set of players; each subset  $\mathbb{A}^p$  is interpreted as the subset of executive agents of the player  $p \in P$ ;
- a W-model (called underlying W-model)  $(\mathbb{A}, (\Omega, \mathcal{F}), (\mathbb{U}_a, \mathcal{U}_a, \mathcal{J}_a)_{a \in \mathbb{A}})$ , as in Definition 1;
- for each player  $p \in P$ , a preference relation  $\preceq^p$  on the set of measurable mappings  $(\Omega, \mathcal{F}) \rightarrow \Delta(\mathbb{U}_{\mathbb{A}}, \mathcal{U}_{\mathbb{A}})$  (where  $\Delta(\mathbb{U}_{\mathbb{A}}, \mathcal{U}_{\mathbb{A}})$  is the set of probability distributions over  $(\mathbb{U}_{\mathbb{A}}, \mathcal{U}_{\mathbb{A}})$ ).

Let  $p \in P$  be a player. A W-game is said to be playable (resp.  $p$ -strongly measurable) if the underlying W-model is playable as in Definition 3 (resp.  $\mathbb{A}^p$ -strongly measurable as in Definition 6).

A finite W-game is a W-game whose underlying W-model is finite. In a W-game, the family  $(\mathbb{A}^p)_{p \in P}$  consists of pairwise disjoint nonempty sets whose union is  $\mathbb{A} = \bigcup_{p \in P} \mathbb{A}^p$ . When we focus on a specific player  $p \in P$ , we denote  $\mathbb{A}^{-p} = \bigcup_{q \in P \setminus \{p\}} \mathbb{A}^q$ . In what follows, agents appear as lower indices and (most of the time) players as upper indices.

Our proposal for the preference relation  $\preceq^p$  is one among possible others (as a fact, we do not need a preference relation for the results in this paper). With it, the above definition covers (like in [5]) the most traditional preference relation  $\preceq^p$ , which is the numerical *expected utility* preference. In this latter, each player  $p \in P$  is endowed, on the

one hand, with a *criterion* (payoff, objective function), that is, a measurable function  $J^p : (\mathbb{H}, \mathcal{H}) \rightarrow [-\infty, +\infty[$  (we include  $-\infty$  in the codomain of the criterion as a way to handle constraints) which is bounded above, and, on the other hand, with a *belief*, that is, a probability distribution  $\nu^p : \mathcal{F} \rightarrow [0, 1]$  over the states of Nature  $(\Omega, \mathcal{F})$ . Then, given two measurable mappings  $S_i : (\Omega, \mathcal{F}) \rightarrow \Delta(\mathbb{U}_{\mathbb{A}}, \mathcal{U}_{\mathbb{A}})$ ,  $i = 1, 2$ , one says that  $S_1 \preceq^p S_2$  if  $\int_{\Omega} \nu^p(d\omega) \int_{\mathbb{U}_{\mathbb{A}}} J^p(\omega, u) S_1(\omega, du) \leq \int_{\Omega} \nu^p(d\omega) \int_{\mathbb{U}_{\mathbb{A}}} J^p(\omega, u) S_2(\omega, du)$  where both integrals are well defined in  $[-\infty, +\infty[$  because the function  $J^p$  is supposed to be bounded above.

Note also that Definition 7 includes Bayesian games, by specifying a product structure  $\Omega = \prod_{n \in \mathcal{N}} \Omega_n$  — where one factor may represent chance, and the other may represent types of players — and a probability on  $\Omega$ .

### 3.2 Mixed and behavioral strategies

We define mixed and behavioral strategies in the spirit of Aumann in [3]. For this purpose, for any agent  $a \in \mathbb{A}$ , we denote by  $(\mathbb{W}_a, \mathcal{W}_a)$  a copy of the Borel space  $([0, 1], \mathcal{B}_{[0,1]})$ , by  $\ell_a$  a copy of the Lebesgue measure on  $(\mathbb{W}_a, \mathcal{W}_a) = ([0, 1], \mathcal{B}_{[0,1]})$ , and we also set

$$\mathbb{W}^p = \prod_{a \in \mathbb{A}^p} \mathbb{W}_a, \quad \mathcal{W}^p = \bigotimes_{a \in \mathbb{A}^p} \mathcal{W}_a, \quad \ell^p = \bigotimes_{a \in \mathbb{A}^p} \ell_a, \quad \forall p \in P \quad (20a)$$

and

$$\mathbb{W} = \prod_{p \in P} \mathbb{W}^p, \quad \mathcal{W} = \bigotimes_{p \in P} \mathcal{W}^p, \quad \ell = \bigotimes_{p \in P} \ell^p. \quad (20b)$$

The existence of a product probability space  $(\mathbb{W}, \mathcal{W}, \ell)$ , that is the existence of a product space  $\mathbb{W}$  equipped with a product  $\sigma$ -algebra  $\mathcal{W}$  and a probability measure  $\ell$  with  $\ell_a$  as marginal probability for each agent  $a \in \mathbb{A}$  is developed in [1, §15.6] and is, in the case we consider, a consequence of the Kolmogorov extension theorem.

**Definition 8.** For the player  $p \in P$ , an  $A$ -mixed strategy is a family  $m^p = (m_a^p)_{a \in \mathbb{A}^p}$  of measurable mappings

$$m_a^p : \left( \prod_{b \in \mathbb{A}^p} \mathbb{W}_b \times \mathbb{H}, \bigotimes_{b \in \mathbb{A}^p} \mathcal{W}_b \otimes \mathcal{J}_a \right) \rightarrow (\mathbb{U}_a, \mathcal{U}_a), \quad \forall a \in \mathbb{A}^p, \quad (21a)$$

an  $A$ -behavioral strategy is an  $A$ -mixed strategy  $m^p = (m_a^p)_{a \in \mathbb{A}^p}$  with the property that

$$(m_a^p)^{-1}(\mathcal{U}_a) \subset \left( \mathbb{W}_a \otimes \bigotimes_{b \in \mathbb{A}^p \setminus \{a\}} \{\emptyset, \mathbb{W}_b\} \right) \otimes \mathcal{J}_a, \quad \forall a \in \mathbb{A}^p, \quad (21b)$$

and an  $A$ -pure strategy is an  $A$ -mixed strategy  $m^p = (m_a^p)_{a \in \mathbb{A}^p}$  with the property that

$$(m_a^p)^{-1}(\mathcal{U}_a) \subset \bigotimes_{b \in \mathbb{A}^p} \{\emptyset, \mathbb{W}_b\} \otimes \mathcal{J}_a, \quad \forall a \in \mathbb{A}^p. \quad (21c)$$

An  $A$ -mixed strategy profile is a family  $m = (m^p)_{p \in P}$  of  $A$ -mixed strategies.

By definition, A-behavioral strategies form a subset of A-mixed strategies. Equation (21b) means that, for any agent  $a$  and any fixed configuration  $h \in \mathbb{H}$ ,  $m_a^p(w^p, h)$  only depends on the randomizing component  $w_a$ . Thus, under the product probability distribution  $\ell^p = \bigotimes_{a \in \mathbb{A}^p} \ell_a$  in (20), the random variables  $(m_a^p(\cdot, h))_{a \in \mathbb{A}^p}$  are independent. In other words, an A-behavioral strategy is an A-mixed strategy in which the randomization is made independently, agent by agent, for each fixed configuration  $h \in \mathbb{H}$ . An A-pure strategy is an A-mixed strategy in which there is no randomization, hence can be identified with a W-pure strategy as in Definition 2.

The connection between A-mixed strategies profiles and pure W-strategies profiles, as in (4a), is as follows: if  $m^p = (m_a^p)_{a \in \mathbb{A}^p}$  is an A-mixed strategy (21a), then every mapping

$$m_a^p(w^p, \cdot) : (\mathbb{H}, \mathcal{J}_a) \rightarrow (\mathbb{U}_a, \mathcal{U}_a), \quad h \mapsto m_a^p(w^p, h), \quad \forall w^p = (w_b)_{b \in \mathbb{A}^p} \in \mathbb{W}^p = \prod_{b \in \mathbb{A}^p} \mathbb{W}_b$$

belongs to  $\Lambda_a$  (see (3)), for  $a \in \mathbb{A}^p$ , and thus  $(m_a^p(w^p, \cdot))_{a \in \mathbb{A}^p} \in \Lambda^p = \prod_{a \in \mathbb{A}^p} \Lambda_a$ . In the same way, an A-mixed strategy profile  $m = (m^p)_{p \in P}$  induces, for any  $w \in \mathbb{W}$ , a mapping  $m(w, \cdot) \in \Lambda = \prod_{a \in \mathbb{A}} \Lambda_a$  in (4b).

Consider a playable W-model (see Definition 3), and a profile  $m = (m^p)_{p \in P}$  of A-mixed strategies. For any  $w \in \mathbb{W}$ ,  $m(w, \cdot)$  is a pure strategy and  $S_{m(w, \cdot)}(\omega)$  is well defined by playability. We use the following compact notation for the solution map as in (14):

$$T_m^\omega(w) = S_{m(w, \cdot)}(\omega), \quad \forall \omega \in \Omega, \quad \forall w \in \mathbb{W}. \quad (22)$$

As we introduce A-mixed strategies, we need to adapt the definition of solvable measurable (SM) property in [15]. To stress the difference, the notion below is for W-games (to distinguish it from a possible definition for W-models inspired by the SM property in [15]).

**Definition 9.** *We say that a W-game is playable and measurable if, for any profile  $m = (m^p)_{p \in P}$  of A-mixed strategies, the following mapping is measurable*

$$T_m : (\mathbb{W} \times \Omega, \mathcal{W} \otimes \mathcal{F}) \rightarrow (\mathbb{H}, \mathcal{H}), \quad (w, \omega) \mapsto T_m^\omega(w), \quad (23)$$

where  $T_m^\omega(w)$  is defined in Equation (22). In that case, for any probability  $\nu$  on  $(\Omega, \mathcal{F})$ , we denote by

$$\mathbb{Q}_m^\nu = \mathbb{Q}_{(m^p)_{p \in P}}^\nu = \left( \left( \bigotimes_{p \in P} \ell^p \right) \otimes \nu \right) \circ (T_{(m^p)_{p \in P}})^{-1} = (\ell \otimes \nu) \circ (T_m)^{-1} \quad (24)$$

the pushforward probability, on the space  $(\mathbb{H}, \mathcal{H})$ , of the product probability distribution  $\ell \otimes \nu = \left( \bigotimes_{p \in P} \ell^p \right) \otimes \nu$  on  $\mathbb{W} \times \Omega = \left( \prod_{p \in P} \mathbb{W}^p \right) \times \Omega$  by the mapping  $T_m$  in (22).

Of course, a playable finite W-game is always playable and measurable.

## 4 Kuhn's equivalence theorem

Now, we are equipped to give, for games in product form, a statement and a proof of the celebrated Kuhn's equivalence theorem: when a player satisfies perfect recall, for any mixed W-strategy, there is an equivalent behavioral strategy (and the reciprocal).



## 4.1 Perfect recall in W-games

For any agent  $a \in \mathbb{A}$ , we define the *choice field*  $\mathcal{C}_a \subset \mathcal{H}$  by<sup>4</sup>

$$\mathcal{C}_a = \mathcal{U}_a \bigvee \mathcal{J}_a, \quad \forall a \in \mathbb{A}. \quad (25)$$

Thus defined, the choice field of agent  $a$  contains both what the agent did ( $\mathcal{U}_a$  identified with  $\mathcal{U}_{\{a\}}$ ) and what he knew ( $\mathcal{J}_a$ ) when making the action. Thus doing, our definition is close to the notion of choice in [2, Definition 4.1].

We consider a focus player  $p \in P$  and we suppose that  $\mathbb{A}^p$  is a finite set with cardinality  $|\mathbb{A}^p|$ . For any  $k \in \llbracket 1, |\mathbb{A}^p| \rrbracket$ , let  $\Sigma_k^p$  denote the set of *k-orderings of player p*, that is, injective mappings from  $\llbracket 1, k \rrbracket$  to  $\mathbb{A}^p$ :

$$\Sigma_k^p = \{ \kappa : \llbracket 1, k \rrbracket \rightarrow \mathbb{A}^p; \kappa \text{ is an injection} \}. \quad (26a)$$

We define the *set of orderings of player p*, shortly *set of p-orderings*, by

$$\Sigma^p = \bigcup_{k=1}^{|\mathbb{A}^p|} \Sigma_k^p. \quad (26b)$$

The set  $\Sigma_{|\mathbb{A}^p|}^p$  is the set of *total orderings of player p*, shortly *total p-orderings*, of agents in  $\mathbb{A}^p$ , that is, bijective mappings from  $\llbracket 1, |\mathbb{A}^p| \rrbracket$  to  $\mathbb{A}^p$  (in contrast with *p-partial orderings* in  $\Sigma_k^p$  for  $k < |\mathbb{A}^p|$ ). For any  $k \in \llbracket 1, |\mathbb{A}^p| \rrbracket$ , any  $p$ -ordering  $\kappa \in \Sigma_k^p$ , and any  $i \in \llbracket 1, k \rrbracket$ ,  $\kappa|_{\llbracket 1, i \rrbracket} \in \Sigma_i^p$  is the restriction of the  $p$ -ordering  $\kappa$  to the first  $i$  integers. For any  $k \in \llbracket 1, |\mathbb{A}^p| \rrbracket$ , there is a natural mapping

$$\psi_k : \Sigma_{|\mathbb{A}^p|}^p \rightarrow \Sigma_k^p, \quad \rho \mapsto \rho|_{\llbracket 1, k \rrbracket}, \quad (27)$$

which is the restriction of any (total)  $p$ -ordering of  $\mathbb{A}^p$  to  $\llbracket 1, k \rrbracket$ . For any  $k \in \llbracket 1, |\mathbb{A}^p| \rrbracket$ , we define the *range*  $\|\kappa\|$  of the  $p$ -ordering  $\kappa \in \Sigma_k^p$  as the subset of agents

$$\|\kappa\| = \{ \kappa(1), \dots, \kappa(k) \} \subset \mathbb{A}^p, \quad \forall \kappa \in \Sigma_k^p, \quad (28a)$$

the *cardinality*  $|\kappa|$  of the  $p$ -ordering  $\kappa \in \Sigma_k^p$  as the integer

$$|\kappa| = k \in \llbracket 1, |\mathbb{A}^p| \rrbracket, \quad \forall \kappa \in \Sigma_k^p, \quad (28b)$$

the *last element*  $\kappa_\star$  of the  $p$ -ordering  $\kappa \in \Sigma_k^p$  as the agent

$$\kappa_\star = \kappa(k) \in \mathbb{A}^p, \quad \forall \kappa \in \Sigma_k^p, \quad (28c)$$

the *first elements*  $\kappa_-$  as the restriction of the  $p$ -ordering  $\kappa \in \Sigma_k^p$  to the first  $k-1$  elements

$$\kappa_- = \kappa|_{\llbracket 1, k-1 \rrbracket} \in \Sigma_{k-1}^p, \quad \forall \kappa \in \Sigma_k^p, \quad (28d)$$

---

<sup>4</sup>As indicated after the definition (5b), we (abusively) identify  $\mathcal{U}_{\{a\}} = \mathcal{U}_a \otimes \bigotimes_{b \neq a} \{ \emptyset, \mathbb{U}_b \} \subset \mathcal{U}_{\mathbb{A}}$  with  $\mathcal{U}_a$ .

with the convention that  $\kappa_- = \emptyset \in \Sigma_0^p = \{\emptyset\}$  when  $\kappa \in \Sigma_1^p$ . With obvious notation, any  $p$ -ordering  $\kappa \in \Sigma^p$  can be written as  $\kappa = (\kappa_-, \kappa_*)$ , with the convention that  $\kappa = (\kappa_*)$  when  $\kappa \in \Sigma_1^p$ .

The following notion of configuration-ordering is adapted from [15, Property C, p. 153].

**Definition 10.** *A  $p$ -configuration-ordering is a mapping  $\varphi : \mathbb{H} \rightarrow \Sigma_{|A^p|}^p$  from configurations towards total  $p$ -orderings. With any  $p$ -configuration-ordering  $\varphi$ , and any  $p$ -ordering  $\kappa \in \Sigma^p$ , we associate the subset  $\mathbb{H}_\kappa^\varphi \subset \mathbb{H}$  of configurations defined by*

$$\mathbb{H}_\kappa^\varphi = \{h \in \mathbb{H}; \psi_{|\kappa|}(\varphi(h)) = \kappa\}, \quad \forall \kappa \in \Sigma^p \cup \Sigma_0^p. \quad (29)$$

By convention, we put  $\mathbb{H}_\emptyset^\varphi = \mathbb{H}$ .

The following definition of perfect recall is new.

**Definition 11.** *We say that a player  $p \in P$  in a  $W$ -model satisfies perfect recall if there exists a  $p$ -configuration-ordering  $\varphi : \mathbb{H} \rightarrow \Sigma_{|A^p|}^p$  such that<sup>5</sup>*

$$\mathbb{H}_\kappa^\varphi \cap H \in \mathcal{J}_{\kappa_*}, \quad \forall H \in \mathcal{C}_{\|\kappa_-\|}, \quad \forall \kappa \in \Sigma^p, \quad (30a)$$

where the subset  $\mathbb{H}_\kappa^\varphi \subset \mathbb{H}$  of configurations has been defined in (29), the last agent  $\kappa_*$  in (28c), the  $p$ -ordering  $\kappa_-$  in (28d), the set  $\Sigma^p$  in (26b), and where<sup>6</sup>

$$\mathcal{C}_{\|\kappa_-\|} = \bigvee_{a \in \|\kappa_-\|} \mathcal{C}_a = \bigvee_{a \in \|\kappa_-\|} \mathcal{U}_a \vee \mathcal{J}_a \subset \mathcal{H}. \quad (30b)$$

We will use the property that  $\mathbb{H}_\kappa^\varphi \in \mathcal{J}_{\kappa_*}$ , by (30) with  $H = \mathbb{H}$ .

We interpret the above definition as follows. A player satisfies perfect recall when any of her executive agents — when called to play as the last one in an ordering — knows at least what did ( $\mathcal{U}_a$  identified with  $\mathcal{U}_{\{a\}}$ ) and knew ( $\mathcal{J}_a$ ) those of the executive agents that are both his predecessors (in the range of the ordering under consideration) and that are executive agents of the player.

While this definition departs from the standard one [10, 12] because it makes no reference to probabilities, it is very close in spirit to the definitions proposed in [11, Definition 203.3], [3] and [13], that rely on “recording” or “recall” functions (whereas (30) involves  $\sigma$ -fields). To illustrate the definition, let us revisit Alice and Bob examples in §2.2. If we consider that Alice and Bob are agents of the same player, then perfect recall is satisfied in the second case (one acting after another as in Figures 1ii and 3) and third case (acting after the Nature’s move as in Figures 1iii and 4), but not in the first case (acting simultaneously as in Figures 1i and 2) because neither Alice nor Bob knows which action the other made.

We are going to show, in Proposition 14 to come, that perfect recall implies the existence of a temporal order on the agents of the focus player. For this purpose, we introduce

<sup>5</sup>When  $\kappa \in \Sigma_1^p$ , the statement (30a) is void.

<sup>6</sup>See Footnote 4 for the abuse of notation for  $\mathcal{U}_a$

the following definition of partial causality, inspired by the property of causality in [15, Property C, p. 153] (and slightly generalized in [16, p. 324]). For any player  $p \in P$ , we set

$$\mathcal{H}_B^p = \mathcal{F} \otimes \bigotimes_{b \in B} \mathcal{U}_b \otimes \bigotimes_{a \in \mathbb{A}^p \setminus B} \{\emptyset, \mathcal{U}_a\} \otimes \bigotimes_{c \notin \mathbb{A}^p} \mathcal{U}_c \subset \mathcal{H}, \quad \forall B \subset \mathbb{A}^p. \quad (31)$$

**Definition 12.** We say that a player  $p \in P$  in a  $W$ -model satisfies partial causality if there exists a  $p$ -configuration-ordering  $\varphi : \mathbb{H} \rightarrow \Sigma_{|\mathbb{A}^p|}^p$  such that

$$\mathbb{H}_\kappa^\varphi \cap H \in \mathcal{H}_{\|\kappa_-\|}^p, \quad \forall H \in \mathcal{J}_{\kappa_*}, \quad \forall \kappa \in \Sigma^p, \quad (32)$$

where the subset  $\mathbb{H}_\kappa^\varphi \subset \mathbb{H}$  of configurations has been defined in (29), the last agent  $\kappa_*$  in (28c), the  $p$ -ordering  $\kappa_-$  in (28d), the set  $\Sigma^p$  in (26b), and  $\mathcal{H}_{\|\kappa_-\|}^p$  in (31). When  $\kappa \in \Sigma_1^p$ ,  $\mathcal{H}_{\|\kappa_-\|}^p = \mathcal{H}_\emptyset^p = \mathcal{F} \otimes \bigotimes_{a \in \mathbb{A}^p} \{\emptyset, \mathcal{U}_a\} \otimes \bigotimes_{c \notin \mathbb{A}^p} \mathcal{U}_c = \mathcal{F} \otimes \mathcal{U}_{\mathbb{A}^p}$ .

The following Lemma 13 will be instrumental in the coming proofs.

**Lemma 13.** Suppose that player  $p \in P$  satisfies partial causality with  $p$ -configuration-ordering  $\varphi : \mathbb{H} \rightarrow \Sigma^p$ . Let  $\kappa \in \Sigma^p$  be a  $p$ -ordering. Then, for any integer  $j \in \llbracket 1, |\kappa| \rrbracket$  and for any  $\mathcal{J}_{\kappa(j)}$ -measurable mapping  $Z : (\mathbb{H}, \mathcal{H}) \rightarrow (\mathbb{Z}, \mathcal{Z})$  — where  $\mathbb{Z}$  is a set (not to be taken in the sense of relative integers) and where the  $\sigma$ -field  $\mathcal{Z}$  contains the singletons — we have the property that

$$\begin{aligned} h' \in \mathbb{H}, \quad h \in \mathbb{H}_{\kappa(1), \dots, \kappa(j-1)}^\varphi, \quad (h_\emptyset, h_{\mathbb{A}^p}, h_{\kappa(1)}, \dots, h_{\kappa(j-1)}) &= (h'_\emptyset, h'_{\mathbb{A}^p}, h'_{\kappa(1)}, \dots, h'_{\kappa(j-1)}) \\ \implies h' \in \mathbb{H}_{\kappa(1), \dots, \kappa(j-1)}^\varphi \quad \text{and} \quad Z(h') &= Z(h), \end{aligned} \quad (33a)$$

which we shortly denote by

$$Z(h) = Z(h_\emptyset, h_{\mathbb{A}^p}, h_{\kappa(1)}, \dots, h_{\kappa(j-1)}), \quad \forall h \in \mathbb{H}_{\kappa(1), \dots, \kappa(j-1)}^\varphi, \quad (33b)$$

where the right hand side means the common value  $Z(h_\emptyset, h_{\mathbb{A}^p}, h_{\kappa(1)}, \dots, h_{\kappa(j-1)}, h'_{\mathbb{A}^p \setminus \{\kappa(1), \dots, \kappa(j-1)\}})$  for any  $h'_{\mathbb{A}^p \setminus \{\kappa(1), \dots, \kappa(j-1)\}}$ .

*Proof.* Suppose that player  $p$  satisfies partial causality with  $p$ -configuration-ordering  $\varphi : \mathbb{H} \rightarrow \Sigma^p$ . Let  $\kappa \in \Sigma^p$ ,  $j \in \llbracket 1, |\kappa| \rrbracket$  and  $Z : (\mathbb{H}, \mathcal{H}) \rightarrow (\mathbb{Z}, \mathcal{Z})$  be a  $\mathcal{J}_{\kappa(j)}$ -measurable mapping. For any configuration  $h \in \mathbb{H}_{\kappa(1), \dots, \kappa(j-1)}^\varphi$ , the set  $Z^{-1}(Z(h))$  contains  $h$  and belongs to  $\mathcal{J}_{\kappa(j)}$ , by the measurability assumption on the mapping  $Z$  and the assumption that the  $\sigma$ -field  $\mathcal{Z}$  contains the singletons. By partial causality (32), we get that  $\mathbb{H}_{\kappa(1), \dots, \kappa(j-1)}^\varphi \cap Z^{-1}(Z(h)) \in \mathcal{H}_{\kappa(1), \dots, \kappa(j-1)}^p$ . By definition (31) of this latter field, the set  $\mathbb{H}_{\kappa(1), \dots, \kappa(j-1)}^\varphi \cap Z^{-1}(Z(h))$  is a cylinder such that, if  $h' \in \mathbb{H}$  and  $(h_\emptyset, h_{\mathbb{A}^p}, h_{\kappa(1)}, \dots, h_{\kappa(j-1)}) = (h'_\emptyset, h'_{\mathbb{A}^p}, h'_{\kappa(1)}, \dots, h'_{\kappa(j-1)})$ , then  $h' \in \mathbb{H}_{\kappa(1), \dots, \kappa(j-1)}^\varphi \cap Z^{-1}(Z(h))$ . Therefore, we have gotten (33a).  $\square$

Now, we show that perfect recall implies the existence of a temporal order on the agents of the focus player.

**Proposition 14.** *In a playable W-model, if a player satisfies perfect recall with some configuration-ordering, then she satisfies partial causality with the same configuration-ordering.*

*Proof.* The proof is by contradiction. We will show that there necessarily exists an agent  $b \in \mathbb{A}^p$  such that  $J_b \not\subset \mathcal{H}_{\mathbb{A} \setminus \{b\}}$  (see Equation (5c)). Now, Witsenhausen proves in [16] that, in a playable W-model, all agents satisfy what he calls *absence of self-information*, namely any agent  $a \in \mathbb{A}$  is such that  $J_a \subset \mathcal{H}_{\mathbb{A} \setminus \{a\}}$ . Therefore, we will obtain a contradiction as it is assumed that the W-model is playable.

We now give the details. Using Definition 11 of perfect recall, there exists a configuration-ordering  $\varphi : \mathbb{H} \rightarrow \Sigma^p$  such that (30) holds true. We suppose that player  $p$  is not partially causal for this very configuration-ordering  $\varphi$ . Then, it follows from Equation (32) that there exists  $\kappa \in \Sigma^p$  and  $H \in \mathcal{J}_{\kappa_\star}$  such that  $\mathbb{H}_\kappa^\varphi \cap H \notin \mathcal{H}_{\|\kappa_-\|}^p$ . Now, by definitions (31) and (5c), we have that  $\mathcal{H}_{\|\kappa_-\|}^p = \bigcap_{b \in \mathbb{A}^p \setminus \|\kappa_-\|} \mathcal{H}_{\mathbb{A} \setminus \{b\}}$ , where the set  $\mathbb{A}^p \setminus \|\kappa_-\|$  is not empty as it contains  $\kappa_\star$ . As a consequence, there exists  $b \in \mathbb{A}^p \setminus \|\kappa_-\|$  such that  $\mathbb{H}_\kappa^\varphi \cap H \notin \mathcal{H}_{\mathbb{A} \setminus \{b\}}$ . By absence of self-information, itself a consequence of the W-model being playable, we have that  $J_{\kappa_\star} \subset \mathcal{H}_{\mathbb{A} \setminus \{\kappa_\star\}}$ , hence that  $\mathbb{H}_\kappa^\varphi \cap H \in J_{\kappa_\star} \subset \mathcal{H}_{\mathbb{A} \setminus \{\kappa_\star\}}$ . As  $\mathbb{H}_\kappa^\varphi \cap H \notin \mathcal{H}_{\mathbb{A} \setminus \{b\}}$ , we deduce that  $b \neq \kappa_\star$ . Then, we denote by  $\Sigma_b^p$  the subset of  $\Sigma^p$  of all  $p$ -orderings  $\kappa' \in \Sigma^p$  such that  $|\kappa'| > |\kappa|$  and  $\psi_{|\kappa|}(\kappa') = \kappa$ , where  $\psi_{|\kappa|}$  has been defined in (27), and such that  $\kappa'_\star = b$ . As  $b \in \mathbb{A}^p \setminus \|\kappa_-\|$ , we get that  $b \notin \|\kappa_-\|$ . Therefore, it readily follows from the definition (26b) of  $\Sigma^p$  that

$$\bigcup_{\kappa' \in \Sigma_b^p} \mathbb{H}_{\kappa'}^\varphi = \mathbb{H}_\kappa^\varphi, \quad (34)$$

as, with any  $h \in \mathbb{H}_\rho^\varphi$ , we associate the total  $p$ -ordering  $\rho = \varphi(h) \in \Sigma_{|\mathbb{A}^p|}^p$  and that  $b \in \{\rho(|\kappa|+1), \dots, \rho(|\mathbb{A}^p|)\}$ , because  $b \in \mathbb{A}^p \setminus \|\kappa_-\|$  and  $b \neq \kappa_\star$ . From there, we get that

$$\begin{aligned} \mathbb{H}_\kappa^\varphi \cap H &= \left( \bigcup_{\kappa' \in \Sigma_b^p} \mathbb{H}_{\kappa'}^\varphi \right) \cap H && \text{(by (34))} \\ &= \bigcup_{\kappa' \in \Sigma_b^p} (\mathbb{H}_{\kappa'}^\varphi \cap H) && \text{(by developing)} \\ &= \bigcup_{\kappa' \in \Sigma^p} \underbrace{(\mathbb{H}_{\kappa'}^\varphi \cap H)}_{\in J_b} \end{aligned}$$

by perfect recall property (30) of agent  $b$  for the subset  $H \in \mathcal{J}_{\kappa_\star} \subset \mathcal{C}_{\|\kappa'_-\|} = \bigvee_{a \in \|\kappa'_-\|} \mathcal{U}_a \vee J_a$ ,

where the last inclusion comes from  $\psi_{|\kappa|}(\kappa') = \kappa$ ,  $|\kappa'| > |\kappa|$  and  $\kappa'_\star = b \neq \kappa_\star$  which imply that  $\kappa_\star \in \|\kappa_-\| \subset \|\kappa'_-\| \setminus \{b\} = \|\kappa'_-\| \setminus \{\kappa'_\star\} = \|\kappa'_-\|$

$$\in J_b. \quad \text{(as the set } \Sigma_b^p \text{ is finite)}$$

As a conclusion, we have therefore obtained that  $\mathbb{H}_\kappa^\varphi \cap H \in J_b$  and  $\mathbb{H}_\kappa^\varphi \cap H \notin \mathcal{H}_{\mathbb{A} \setminus \{b\}}$  and therefore  $J_b \not\subset \mathcal{H}_{\mathbb{A} \setminus \{b\}}$ . Now, this contradicts the absence of self information for agent  $b$ , hence contradicts playability.

This ends the proof. □

The statement of Proposition 14 resembles the one by Ritzenger in [12] on the fact “that present past and future have an unambiguous meaning” when the player satisfies perfect recall.

## 4.2 Main results

We can now state the main results of the paper. The proofs are provided in Sect. 5.

### 4.2.1 Sufficiency of perfect recall for behavioral strategies to be as powerful as mixed strategies

It happens that, for the proof of the first main theorem, we resort to regular conditional distributions, and that these objects display nice properties when defined on Borel spaces, and when the conditioning is with respect to measurable mappings (and not general  $\sigma$ -fields). This is why we introduce the following notion that informations fields are generated by Borel measurable mappings.

**Definition 15.** *We say that player  $p \in P$  in a  $W$ -game satisfies the Borel measurable functional information assumption if there exists a family  $((\mathbb{Z}_a, \mathcal{Z}_a))_{a \in \mathbb{A}^p}$  of Borel spaces and a family  $(Z_a)_{a \in \mathbb{A}^p}$  of measurable mappings  $Z_a : (\mathbb{H}, \mathcal{H}) \rightarrow (\mathbb{Z}_a, \mathcal{Z}_a)$  such that  $Z_a^{-1}(\mathcal{Z}_a) = \mathcal{I}_a$ , for all  $a \in \mathbb{A}^p$ .*

Of course, a player in a finite  $W$ -game always satisfies the Borel measurable functional information assumption.

We now state the first main theorem, namely sufficiency of perfect recall for behavioral strategies to be as powerful as mixed strategies.

**Theorem 16** (Kuhn’s theorem). *We consider a playable and measurable  $W$ -game (see Definition 9). Let  $p \in P$  be a given player. We suppose that the  $W$ -game is  $p$ -strongly measurable (see Definition 7), that player  $p$  satisfies the Borel measurable functional information assumption (see Definition 15), that  $\mathbb{A}^p$  is a finite set, that  $(\mathbb{U}_a, \mathcal{U}_a)$  is a Borel space, for all  $a \in \mathbb{A}^p$ , and that  $(\Omega, \mathcal{F})$  is a Borel space.*

*Suppose that the player  $p$  satisfies perfect recall, as in Definition 11. Then, for any probability  $\nu$  on  $(\Omega, \mathcal{F})$ , for any  $A$ -mixed strategy  $m^{-p} = (m_a^{-p})_{a \in \mathbb{A}^{-p}}$  of the other players and for any  $A$ -mixed strategy  $m^p = (m_a^p)_{a \in \mathbb{A}^p}$ , of the player  $p$ , there exists an  $A$ -behavioral strategy  $m'^p = (m'_a{}^p)_{a \in \mathbb{A}^p}$  of the player  $p$  such that*

$$\mathbb{Q}_{(m^{-p}, m^p)}^\nu = \mathbb{Q}_{(m^{-p}, m'^p)}^\nu, \quad (35)$$

where the pushforward probability  $\mathbb{Q}_{(m^{-p}, m^p)}^\nu$  has been defined in (24).

Theorem 16 applies to the case where the focus player (the one satisfying perfect recall) makes her actions in finite sets, so that we cover the original result in [10]. Regarding the case where the focus player makes her actions in infinite (Borel) sets, the only result that we know of is [3] (to the best of our knowledge, see the discussion at the end of §6.4 in [2,

p. 159]). We emphasize proximities and differences. In [3], the focus player make her actions in Borel sets, and plays a countable number of times where the order of actions is fixed in advance. In our result, the focus player also makes her actions in Borel sets and the order of actions is not fixed in advance, but she plays a finite number of times.

#### 4.2.2 Necessity of perfect recall for behavioral strategies to be as powerful as mixed strategies

After stating the second main theorem, namely necessity of perfect recall for behavioral strategies to be as powerful as mixed strategies, we will comment on our formulation.

**Theorem 17.** *We consider a playable and measurable W-game (see Definition 9). Let  $p \in P$  be a given player. We suppose that player  $p$  satisfies the Borel measurable functional information assumption (see Definition 15) and partial causality (see Definition 12), that  $\mathbb{A}^p$  is a finite set, and that  $\mathbb{U}_a$  contains at least two distinct elements, for all  $a \in \mathbb{A}^p$ .*

*Suppose that, for the  $p$ -configuration-ordering  $\varphi : \mathbb{H} \rightarrow \Sigma^p$  given by partial causality, there exists a  $p$ -ordering  $\kappa \in \Sigma^p$  such that*

$$\exists h^+, h^- \in \mathbb{H}_\kappa^\varphi, \quad Z_{\kappa_*}(h^+) = Z_{\kappa_*}(h^-), \quad (Z_a(h^+), h_a^+)_{a \in \|\kappa_{-}\|} \neq (Z_a(h^-), h_a^-)_{a \in \|\kappa_{-}\|}. \quad (36)$$

*Then, there exists an A-mixed strategy  $m^{-p} = (m_a^{-p})_{a \in \mathbb{A}^{-p}}$  of the other players, an A-mixed strategy  $m^p = (m_a^p)_{a \in \mathbb{A}^p}$  of the player  $p$ , and a probability distribution  $\nu$  on  $\Omega$  such that, for any A-behavioral strategy  $m'^p = (m'_a{}^p)_{a \in \mathbb{A}^p}$  of the player  $p$ , we have that  $\mathbb{Q}_{(m^{-p}, m^p)}^\nu \neq \mathbb{Q}_{(m^{-p}, m'^p)}^\nu$  where the pushforward probability  $\mathbb{Q}_{(m^{-p}, m^p)}^\nu$  has been defined in (24).*

In case of a finite W-game, the condition (36) is the negation of the perfect recall property (30) (characterize the condition (30) in terms of atoms, and then express the negation using the property that the mappings  $Z_a$  are constant on suitable atoms). For more general W-games, we could formally define a weaker notion of perfect recall than (30): a functional version of perfect recall would replace the  $\sigma$ -fields inclusions in (30) by functional constraints of the form<sup>7</sup>  $(Z_a(h), h_a)_{a \in \|\kappa_{-}\|} = \phi^\kappa(Z_{\kappa_*}(h))$ , for all  $h \in \mathbb{H}_\kappa^\varphi$ , where the mappings  $\phi^\kappa$  would not be supposed to be measurable. We do not pursue this formal path and we prefer to recognize that there is a technical difficulty in negating a  $\sigma$ -fields inclusion — or, equivalently, by Doob functional theorem [7, Chap. 1, p. 18], in negating the existence of a measurable functional constraint. Thus doing, we follow [13] who also had to negate a weaker version of perfect recall and who had to invoke the weaker notion of R-games to prove the necessity of perfect recall.

Theorem 17 applies to the case where the focus player makes her actions in finite sets, so that we cover the original result in [10]. Regarding the case where the focus player makes her actions in infinite (Borel) sets, the only result that we know of is [13] (to the best of our knowledge, see the discussion at the end of §6.4 in [2, p. 159]). We emphasize proximities and differences. In [13], the focus player make her actions in Borel sets, and plays a countable

---

<sup>7</sup>The mappings  $\phi^\kappa$  correspond to the “recall” functions in [3, 13].

number of times where the order of actions is fixed in advance. In our result, the focus player also makes her actions in any measurable set with at least two elements, and the order of actions is not fixed in advance, but she plays a finite number of times.

## 5 Proofs of the main results

We give the proof of Theorem 16 in §5.1 and of Theorem 17 in §5.2.

### 5.1 Proof of Theorem 16

We will need the notion of stochastic kernel. Let  $(\mathbb{X}, \mathcal{X})$  and  $(\mathbb{Y}, \mathcal{Y})$  be two measurable spaces. A *stochastic kernel* from  $(\mathbb{X}, \mathcal{X})$  to  $(\mathbb{Y}, \mathcal{Y})$  is a mapping  $\Gamma : \mathbb{X} \times \mathcal{Y} \rightarrow [0, 1]$  such that for any  $Y \in \mathcal{Y}$ ,  $\Gamma(\cdot, Y) : \mathbb{X} \rightarrow [0, 1]$  is  $\mathcal{X}$ -measurable and, for any  $x \in \mathbb{X}$ ,  $\Gamma(x, \cdot) : \mathcal{Y} \rightarrow [0, 1]$  is a probability measure on  $\mathcal{Y}$ .

The proof of Theorem 16 is decomposed into four lemmas and a final proof. The overall logic is as follows:

1. in Lemma 18, we obtain key technical disintegration formulas for stochastic kernels on the action spaces,
2. in Lemma 19, we identify the candidate behavioral strategy,
3. in Lemma 20, we show that one step substitution (ordered agent by ordered agent) between behavioral and mixed strategies is possible,
4. we apply the substitution procedure between the first and last agent of the player and obtain, in the substitution Lemma 21, a kind of Kuhn's Theorem, but on the randomizing device space  $\mathbb{W}$  instead of the configuration space  $\mathbb{H}$ ,
5. we conclude the proof of Kuhn's Theorem 16 (sufficiency) on the configuration space  $\mathbb{H}$ , by enabling the use of Lemma 21 with the pushforward probability formula (24).

We start with the technical Lemma 18 on stochastic kernels on the action spaces.

**Lemma 18** (Disintegration). *Suppose that the assumptions of Theorem 16 are satisfied, and that the player  $p \in P$  satisfies perfect recall, as in Definition 11. We consider a probability  $\nu$  on  $(\Omega, \mathcal{F})$ , an  $A$ -mixed strategy  $m^p = (m_a^p)_{a \in \mathbb{A}^p}$ , of the player  $p$  and an  $A$ -mixed strategy  $m^{-p} = (m_a^{-p})_{a \in \mathbb{A}^{-p}}$  of the other players.*

*As  $(\mathbb{W} \times \Omega, \mathcal{W} \otimes \mathcal{F})$  is a Borel space, as the mapping  $Z_a$  is measurable by the Borel measurable functional information assumption (see Definition 15), and as the mapping  $T_m$  is measurable by assumption that the  $W$ -game is playable and measurable, we denote by  $(\ell \otimes \nu)^{|Z_a \circ T_m} [d\omega d\omega | z]$  the regular conditional distribution on the probability space  $(\mathbb{W} \times \Omega, \mathcal{W} \otimes \mathcal{F}, \ell \otimes \nu)$  knowing the random variable  $Z_a \circ T_m : (\mathbb{W} \times \Omega, \mathcal{W} \otimes \mathcal{F}) \rightarrow (\mathbb{Z}_a, \mathcal{Z}_a)$ .*

*Then, there exists*

- a family  $(\Gamma_\kappa)_{\kappa \in \Sigma^p}$  of stochastic kernels, where  $\Gamma_\kappa : \mathcal{U}_{\|\kappa\|} \times \mathbb{H}_\kappa^\varphi \rightarrow [0, 1]$  is an  $(\mathbb{H}_\kappa^\varphi \cap \mathcal{J}_{\kappa_\star})$ -measurable stochastic kernel, such that

$$\Gamma_\kappa[du_\kappa | h] = ((\ell \otimes \nu)^{|Z_{\kappa_\star} \circ T_m}[\cdot | Z_{\kappa_\star}(h)] \circ m_\kappa^p(\cdot, h)^{-1})(du_\kappa), \quad \forall h \in \mathbb{H}_\kappa^\varphi, \quad (37)$$

where we use the shorthand notation  $m_\kappa^p = (m_a^p)_{a \in \|\kappa\|}$ , and that

$$\Gamma_\kappa[du_\kappa | h] = \mathbf{1}_{\{u_{\|\kappa\|} = h_{\|\kappa\|}\}} \Gamma_\kappa[du_\kappa | h] = \mathbf{1}_{\{u_{\|\kappa_- \|} = h_{\|\kappa_- \|}\}} \Gamma_\kappa[du_\kappa | h], \quad \forall h \in \mathbb{H}_\kappa^\varphi, \quad (38)$$

- a family  $(\Gamma_\kappa^{\kappa_-})_{\kappa \in \Sigma^p}$  of stochastic kernels where  $\Gamma_\kappa^{\kappa_-} : \mathcal{U}_{\|\kappa_- \|} \times \mathbb{H}_\kappa^\varphi \rightarrow [0, 1]$ , such that

$$\Gamma_\kappa^{\kappa_-}[du_{\kappa_-} | h] = \left( \ell^{|Z_{\kappa_\star} \circ T_m^\omega}[\cdot | Z_{\kappa_\star}(h)] \circ m_{\kappa_-}^p(\cdot, h)^{-1} \right)(du_{\kappa_-}), \quad \forall h \in \mathbb{H}_\kappa^\varphi, \quad (39)$$

- a family  $(\Gamma_\kappa^{\kappa_\star})_{\kappa \in \Sigma^p}$  of stochastic kernels, where<sup>8</sup>  $\Gamma_\kappa^{\kappa_\star} : \mathcal{U}_{\kappa_\star} \times (\mathbb{U}_{\|\kappa_- \|} \times \mathbb{H}_\kappa^\varphi) \rightarrow [0, 1]$  is an  $\mathcal{U}_{\|\kappa_- \|} \otimes (\mathbb{H}_\kappa^\varphi \cap \mathcal{J}_{\kappa_\star})$ -measurable stochastic kernel, such that

$$\Gamma_\kappa[du_\kappa | h] = \Gamma_\kappa[du_{\kappa_-} du_{\kappa_\star} | h] = \Gamma_\kappa^{\kappa_\star}[du_{\kappa_\star} | u_{\kappa_-}, h] \otimes \Gamma_\kappa^{\kappa_-}[du_{\kappa_-} | h], \quad \forall h \in \mathbb{H}_\kappa^\varphi. \quad (40)$$

*Proof.* We consider a  $p$ -ordering  $\kappa \in \Sigma^p$ . We are going to prove the following preliminary result: the mapping<sup>9</sup>  $m_\kappa^p = (m_a^p)_{a \in \|\kappa\|} : (\mathbb{W} \times \mathbb{H}_\kappa^\varphi, \mathcal{W} \otimes (\mathbb{H}_\kappa^\varphi \cap \mathcal{J}_{\kappa_\star})) \rightarrow (\mathbb{U}_{\|\kappa\|}, \mathcal{U}_{\|\kappa\|})$  is measurable, by studying each component  $m_a^p : (\mathbb{W} \times \mathbb{H}_\kappa^\varphi, \mathcal{W} \otimes (\mathbb{H}_\kappa^\varphi \cap \mathcal{J}_{\kappa_\star})) \rightarrow (\mathbb{U}_a, \mathcal{U}_a)$  for  $a \in \|\kappa\|$ . Indeed, on the one hand, as the mapping  $m_{\kappa_\star}^p$  is  $\mathcal{W} \otimes \mathcal{J}_{\kappa_\star}$ -measurable by definition (21a) of an A-mixed strategy, we deduce that the (restriction) mapping  $m_{\kappa_\star}^p : (\mathbb{W} \times \mathbb{H}_\kappa^\varphi, \mathcal{W} \otimes (\mathbb{H}_\kappa^\varphi \cap \mathcal{J}_{\kappa_\star})) \rightarrow (\mathbb{U}_{\kappa_\star}, \mathcal{U}_{\kappa_\star})$  is measurable (by definition of the trace field  $\mathbb{H}_\kappa^\varphi \cap \mathcal{J}_{\kappa_\star}$ ). On the other hand, for any  $a \in \|\kappa_- \|$ , the mapping  $m_a^p$  is  $\mathcal{W} \otimes \mathcal{J}_a$ -measurable by definition (21a) of an A-mixed strategy, where  $\mathbb{H}_\kappa^\varphi \cap \mathcal{J}_a \subset \mathcal{J}_{\kappa_\star}$  by perfect recall (30); we deduce that the (restriction) mapping  $m_a^p : (\mathbb{W} \times \mathbb{H}_\kappa^\varphi, \mathcal{W} \otimes (\mathbb{H}_\kappa^\varphi \cap \mathcal{J}_{\kappa_\star})) \rightarrow (\mathbb{U}_a, \mathcal{U}_a)$  is measurable.

We define  $\Gamma_\kappa$  by (37), that is, for any  $U_{\|\kappa\|} \in \mathcal{U}_{\|\kappa\|}$  and  $h \in \mathbb{H}_\kappa^\varphi$ :

$$\Gamma_\kappa[U_{\|\kappa\|} | h] = \int_{\mathbb{W} \times \Omega} (\ell \otimes \nu)^{|Z_{\kappa_\star} \circ T_m}[\cdot | Z_{\kappa_\star}(h)] \mathbf{1}_{\{m_\kappa^p(w, h) \in U_{\|\kappa\|}\}} \cdot$$

The function  $\mathbb{H}_\kappa^\varphi \ni h \mapsto \Gamma_\kappa[U_{\|\kappa\|} | h]$  is  $(\mathbb{H}_\kappa^\varphi \cap \mathcal{J}_{\kappa_\star})$ -measurable because the stochastic kernel  $(\ell \otimes \nu)^{|Z_{\kappa_\star} \circ T_m}$  is  $\mathcal{J}_{\kappa_\star}$ -measurable by its very definition, and the function  $\mathbb{H}_\kappa^\varphi \ni h \mapsto \mathbf{1}_{\{m_\kappa^p(w, h) \in U_{\|\kappa\|}\}}$  is measurable, from our preliminary result. As a consequence,  $\Gamma_\kappa : \mathcal{U}_{\|\kappa\|} \times \mathbb{H}_\kappa^\varphi \rightarrow [0, 1]$  is an  $(\mathbb{H}_\kappa^\varphi \cap \mathcal{J}_{\kappa_\star})$ -measurable stochastic kernel. As  $\mathbf{1}_{\{m_\kappa^p(w, T_m^\omega(w)) = \pi_{\|\kappa\|}(T_m^\omega(w))\}} = \mathbf{1}_{\{m_{\kappa_-}^p(w, T_m^\omega(w)) = \pi_{\|\kappa_- \|}(T_m^\omega(w))\}} = 1$  by playability property (15) and by definition (22) of  $T_m^\omega(w)$ , we get (38).

By parametric disintegration [4, p. 135] — which holds true because  $(\mathbb{U}_a, \mathcal{U}_a)$  is a Borel space, for all  $a \in \mathbb{A}^p$ , by assumption of Theorem 16 — there exists a stochastic kernel

<sup>8</sup>If  $\|\kappa_- \| = \emptyset$ ,  $(\mathbb{U}_{\|\kappa_- \|} \times \mathbb{H}_\kappa^\varphi) = \mathbb{H}_\kappa^\varphi$  and  $\mathcal{U}_{\|\kappa_- \|} \otimes (\mathbb{H}_\kappa^\varphi \cap \mathcal{J}_a) = (\mathbb{H}_\kappa^\varphi \cap \mathcal{J}_a)$ .

<sup>9</sup>By abuse of notation, we use the same symbol to denote a mapping and the restriction of this mapping to a subset of the domain.



$\Gamma_{\kappa^*}^{\kappa^*} : \mathcal{U}_{\kappa^*} \times (\mathbb{U}_{\|\kappa_{-}\|} \times \mathbb{H}_{\kappa}^{\varphi}) \rightarrow [0, 1]$ , which is  $\mathcal{U}_{\|\kappa_{-}\|} \otimes (\mathbb{H}_{\kappa}^{\varphi} \cap \mathcal{J}_{\kappa^*})$ -measurable, and a stochastic kernel  $\Gamma_{\kappa_{-}}^{\kappa} : \mathcal{U}_{\|\kappa_{-}\|} \times \mathbb{H}_{\kappa}^{\varphi} \rightarrow [0, 1]$ , which is  $(\mathbb{H}_{\kappa}^{\varphi} \cap \mathcal{J}_{\kappa^*})$ -measurable, such that (40) holds true. By taking marginal distributions, we get (39).

This ends the proof.  $\square$

Lemma 18 is particularly useful to prove the next result, which provides us with a candidate behavioral strategy.

**Lemma 19** (Candidate behavioral strategy for equivalence). *Suppose that the assumptions of Theorem 16 are satisfied, and that the player  $p \in P$  satisfies perfect recall, as in Definition 11. We consider a probability  $\nu$  on  $(\Omega, \mathcal{F})$ , an  $A$ -mixed strategy  $m^p = (m_a^p)_{a \in \mathbb{A}^p}$ , of the player  $p$  and an  $A$ -mixed strategy  $m^{-p} = (m_a^{-p})_{a \in \mathbb{A}^{-p}}$  of the other players.*

*Then, there exists an  $A$ -behavioral strategy  $m^p = (m_a^p)_{a \in \mathbb{A}^p}$  of the player  $p$  such that, for any agent  $a \in \mathbb{A}^p$ , and any  $p$ -ordering  $\kappa \in \Sigma^p$ , we have that*

$$\kappa_{\star} = a \implies (\ell_a \circ m_a^p(\cdot, h)^{-1})(du_a) = \Gamma_{\kappa}^a[du_a \mid h_{\|\kappa_{-}\|}, h] = \Gamma_{\kappa^*}^{\kappa^*}[du_{\kappa^*} \mid h_{\|\kappa_{-}\|}, h], \quad (41)$$

$$\forall h \in \mathbb{H}_{\kappa}^{\varphi}.$$

*Proof.* We consider an agent  $a \in \mathbb{A}^p$  and we define, for any  $p$ -ordering  $\kappa \in \Sigma^p$  such that  $\kappa_{\star} = a$ ,

$$\beta_a^{\kappa}[U_a \mid h] = \Gamma_{\kappa}^a[U_a \mid h_{\|\kappa_{-}\|}, h], \quad \forall U_a \in \mathcal{U}_a, \quad \forall h \in \mathbb{H}_{\kappa}^{\varphi}.$$

Thus defined, the function  $\beta_a^{\kappa} : \mathcal{U}_a \times \mathbb{H}_{\kappa}^{\varphi} \rightarrow [0, 1]$  is an  $\mathbb{H}_{\kappa}^{\varphi} \cap \mathcal{J}_a$ -measurable stochastic kernel because, for any  $U_a \in \mathcal{U}_a$ , the function  $h \mapsto \beta_a^{\kappa}[U_a \mid h]$  is obtained by composition

$$(\mathbb{H}_{\kappa}^{\varphi}, \mathbb{H}_{\kappa}^{\varphi} \cap \mathcal{J}_a) \rightarrow (\mathbb{U}_{\|\kappa_{-}\|} \times \mathbb{H}_{\kappa}^{\varphi}, \mathcal{U}_{\|\kappa_{-}\|} \otimes (\mathbb{H}_{\kappa}^{\varphi} \cap \mathcal{J}_a)) \rightarrow [0, 1]$$

$$h \mapsto (h_{\|\kappa_{-}\|}, h) \mapsto \Gamma_{\kappa}^a[U_a \mid h_{\|\kappa_{-}\|}, h].$$

In this composition, the second mapping is measurable since  $\Gamma_{\kappa}^a$  is a  $(\mathbb{H}_{\kappa}^{\varphi} \cap \mathcal{J}_a)$ -measurable stochastic kernel by Lemma 18, and since the first mapping  $h \mapsto h_{\|\kappa_{-}\|}$  is  $(\mathbb{H}_{\kappa}^{\varphi} \cap \mathcal{U}_{\|\kappa_{-}\|})$ -measurable, hence  $(\mathbb{H}_{\kappa}^{\varphi} \cap \mathcal{J}_a)$ -measurable by perfect recall (30).

The family  $(\mathbb{H}_{\kappa}^{\varphi})_{\kappa \in \Sigma^p, \kappa_{\star} = a}$  consists of pairwise disjoint (possibly empty) sets whose union is  $\mathbb{H}$ . Indeed, for any  $h \in \mathbb{H}$ , we consider the total  $p$ -ordering  $\rho = \varphi(h)$ , we denote by  $k \in \mathbb{N}^*$  the index such that  $\rho(k) = a$ , we set the restriction  $\kappa = \psi_k(\rho) \in \Sigma^p$ , where  $\psi_k$  has been defined in (27) for  $k \in \llbracket 1, |\mathbb{A}^p| \rrbracket$ , and we get  $h \in \mathbb{H}_{\kappa}^{\varphi}$  with  $\kappa_{\star} = a$ . What is more, for every subset of the family, we have that  $\mathbb{H}_{\kappa}^{\varphi} \in \mathcal{J}_{\kappa^*} = \mathcal{J}_a$ , by (30a) with  $H = \mathbb{H}$ . Then, for any  $U_a \in \mathcal{U}_a$ , we define  $\beta_a[U_a \mid h] = \sum_{\kappa \in \Sigma^p, \kappa_{\star} = a} \mathbf{1}_{\mathbb{H}_{\kappa}^{\varphi}}(h) \beta_a^{\kappa}[U_a \mid h]$ , for any  $h \in \mathbb{H}$ . As we have established that the function  $h \mapsto \beta_a^{\kappa}[U_a \mid h]$  is  $\mathcal{J}_a$ -measurable and that the  $\mathbb{H}_{\kappa}^{\varphi}$  belong to  $\mathcal{J}_a$ , we conclude that the function  $\beta_a : \mathcal{U}_a \times \mathbb{H} \rightarrow [0, 1]$  is an  $\mathcal{J}_a$ -measurable stochastic kernel.

By [9, Lemma 3.22] (realization lemma), the  $\mathcal{J}_a$ -measurable stochastic kernel  $\beta_a$  can be *realized* as the pushforward of the Lebesgue measure  $\ell_a$  by a measurable random variable  $m_a''(\cdot, h)$ ,  $\mathcal{J}_a$ -measurably in  $h$ . More precisely, there exists a measurable mapping  $m_a'' : (\mathbb{W}_a \times \mathbb{H}, \mathcal{W}_a \otimes \mathcal{J}_a) \rightarrow (\mathbb{U}_a, \mathcal{U}_a)$  such that

$$(\ell_a \circ m_a''(\cdot, h)^{-1})(du_a) = \beta_a[du_a \mid h].$$

We easily extend the mapping  $m_a''$  from the domain  $\mathbb{W}_a$  to the domain  $\mathbb{W}$  in (20), by setting  $m_a'^p : (\prod_{b \in \mathbb{A}^p} \mathbb{W}_b \times \mathbb{H}, \mathcal{W}_a \otimes \mathcal{J}_a) \rightarrow (\mathbb{U}_a, \mathcal{U}_a)$  defined by  $m_a'^p((w_b)_{b \in \mathbb{A}^p}) = m_a''(w_a)$ . Thus, we get (41).

This ends the proof.  $\square$

The next Lemma 20 concentrates much of the technical difficulty. It provides us with a way to replace the A-mixed strategy  $m^p$  by the A-behavioral strategy  $m'^p$  in an integral expression, which gives us a clear path toward Kuhn's theorem. It combines Lemma 19 with results from probability theory, in particular Doob functional theorem and properties of regular conditional distributions.

**Lemma 20** (One step mixed/behavioral substitution). *Suppose that the assumptions of Theorem 16 are satisfied, and that the player  $p \in P$  satisfies perfect recall, as in Definition 11. We consider a probability  $\nu$  on  $(\Omega, \mathcal{F})$ , an A-mixed strategy  $m^p = (m_a^p)_{a \in \mathbb{A}^p}$ , of the player  $p$  and an A-mixed strategy  $m^{-p} = (m_a^{-p})_{a \in \mathbb{A}^{-p}}$  of the other players.*

*Then, the A-behavioral strategy  $m'^p = (m_a'^p)_{a \in \mathbb{A}^p}$  of Lemma 19 has the property that, for any  $p$ -ordering  $\kappa \in \Sigma^p$  and for any bounded measurable function  $\Phi : \mathbb{U}_{\|\kappa\|} \rightarrow \mathbb{R}$ , we have that*

$$\begin{aligned} & \int_{\mathbb{W}} \ell(dw) \mathbf{1}_{\mathbb{H}_\kappa^\varphi}(T_m^\omega(w)) \Phi(m_\kappa^p(w^p, T_m^\omega(w))) \\ &= \int_{\mathbb{W}} \ell(dw) \mathbf{1}_{\mathbb{H}_\kappa^\varphi}(T_m^\omega(w)) \int_{\mathbb{W}_{\kappa_\star}} \ell_{\kappa_\star}(dw'_{\kappa_\star}) \Phi\left(m_{\kappa_-}^p(w^p, T_m^\omega(w)), m_{\kappa_\star}^p(w'_{\kappa_\star}, T_m^\omega(w))\right), \end{aligned} \quad (42)$$

where we use the shorthand notation  $m_\kappa^p = (m_a^p)_{a \in \|\kappa\|}$ .

*Proof.* Let  $\kappa \in \Sigma^p$  and  $\Phi : \mathbb{U}_{\|\kappa\|} \rightarrow \mathbb{R}$  be a bounded measurable function. As a preliminary result, we show that there exists a measurable function  $\Psi : (\mathbb{W} \times \mathbb{Z}_{\kappa_\star}, \mathcal{W} \otimes \mathcal{Z}_{\kappa_\star}) \rightarrow (\mathbb{R}, \mathcal{B}_{\mathbb{R}})$  such that

$$\Psi(w, Z_{\kappa_\star}(h)) = \mathbf{1}_{\mathbb{H}_\kappa^\varphi}(h) \Phi(m_\kappa^p(w^p, h)), \quad \forall h \in \mathbb{H}. \quad (43)$$

Indeed, the function  $\mathbb{W} \times \mathbb{H} \ni (w, h) \mapsto \mathbf{1}_{\mathbb{H}_\kappa^\varphi}(h) \Phi(m_\kappa^p(w^p, h))$  is measurable with respect to  $\mathcal{W} \otimes (\mathbb{H}_\kappa^\varphi \cap (\bigvee_{a \in \|\kappa\|} \mathcal{J}_a))$  by definition (21a) of an A-mixed strategy and by definition of the trace field  $\mathbb{H}_\kappa^\varphi \cap (\bigvee_{a \in \|\kappa\|} \mathcal{J}_a)$ , hence with respect to  $\mathcal{W} \otimes (\mathbb{H}_\kappa^\varphi \cap (\mathcal{C}_{\|\kappa-\|} \vee \mathcal{J}_{\kappa_\star}))$  by definition (30b) of  $\mathcal{C}_{\|\kappa-\|}$ , hence with respect to  $\mathcal{W} \otimes (\mathcal{J}_{\kappa_\star} \vee (\mathbb{H}_\kappa^\varphi \cap \mathcal{J}_{\kappa_\star}))$  by perfect recall (30), hence with respect to  $\mathcal{W} \otimes \mathcal{J}_{\kappa_\star}$  as  $\mathbb{H}_\kappa^\varphi \in \mathcal{J}_{\kappa_\star}$  by (30) with  $H = \mathbb{H}$ . As a consequence, as  $Z_{\kappa_\star}^{-1}(\mathcal{Z}_{\kappa_\star}) = \mathcal{J}_{\kappa_\star}$  by assumption, by Doob functional theorem [7, Chap. 1, p. 18], there exists a measurable function  $\Psi : (\mathbb{W} \times \mathbb{Z}_{\kappa_\star}, \mathcal{W} \otimes \mathcal{Z}_{\kappa_\star}) \rightarrow (\mathbb{R}, \mathcal{B}_{\mathbb{R}})$  such that (43) holds true, because  $(\mathbb{W} \times \mathbb{Z}_{\kappa_\star}, \mathcal{W} \otimes \mathcal{Z}_{\kappa_\star})$  is a product of Borel spaces, hence is a Borel space.

We have that

$$\begin{aligned}
& \int_{\mathbb{W} \times \Omega} \ell(dw) \nu(d\omega) \mathbf{1}_{\mathbb{H}_\kappa^\varphi}(T_m^\omega(w)) \Phi(m_\kappa^p(w^p, T_m^\omega(w))) && \text{(where } w = (w^p, w^{-p})\text{)} \\
&= \int_{\mathbb{W} \times \Omega} \ell(dw) \nu(d\omega) \Psi(w, Z_{\kappa_\star}(T_m^\omega(w))) && \text{(by property (43) of the function } \Psi\text{)} \\
&= \int_{\mathbb{W} \times \Omega} \ell(dw) \nu(d\omega) \left[ \int_{\mathbb{W} \times \Omega} (\ell \otimes \nu)^{|Z_{\kappa_\star} \circ T_m}[dw' d\omega' | z] \Psi(w', z) \right]_{|z=Z_{\kappa_\star} \circ T_m^\omega(w)}
\end{aligned}$$

by property of regular conditional distributions [9, Th. 6.4]

$$\begin{aligned}
&= \int_{\mathbb{W} \times \Omega} \ell(dw) \nu(d\omega) \left[ \int_{\mathbb{W} \times \Omega} (\ell \otimes \nu)^{|Z_{\kappa_\star} \circ T_m}[dw' d\omega' | Z_{\kappa_\star}(h)] \Psi(w', Z_{\kappa_\star}(h)) \right]_{|h=T_m^\omega(w)} \\
&\quad \text{(by the change of variables } z = Z_{\kappa_\star}(h), h = T_m^\omega(w)\text{)} \\
&= \int_{\mathbb{W} \times \Omega} \ell(dw) \nu(d\omega) \left[ \int_{\mathbb{W} \times \Omega} (\ell \otimes \nu)^{|Z_{\kappa_\star} \circ T_m}[dw' d\omega' | Z_{\kappa_\star}(h)] \right. \\
&\quad \left. \mathbf{1}_{\mathbb{H}_\kappa^\varphi}(h) \Phi(m_\kappa^p(w^p, h)) \right]_{|h=T_m^\omega(w)} \\
&\quad \text{(by property (43) of the function } \Psi\text{)} \\
&= \int_{\mathbb{W} \times \Omega} \ell(dw) \nu(d\omega) \left[ \mathbf{1}_{\mathbb{H}_\kappa^\varphi}(h) \int_{\mathbb{W} \times \Omega} (\ell \otimes \nu)^{|Z_{\kappa_\star} \circ T_m}[dw' d\omega' | Z_{\kappa_\star}(h)] \Phi(m_\kappa^p(w^p, h)) \right]_{|h=T_m^\omega(w)}
\end{aligned}$$

where the inner integral (the last one inside the brackets) is given by

$$\begin{aligned}
& \int_{\mathbb{W} \times \Omega} (\ell \otimes \nu)^{|Z_{\kappa_\star} \circ T_m}[dw' d\omega' | Z_{\kappa_\star}(h)] \Phi(m_\kappa^p(w^p, h)) \\
&= \int_{\mathbb{U}_{\|\kappa\|}} \Phi(u_\kappa) \Gamma_\kappa[du_\kappa | h] && \text{(by definition (37) of the stochastic kernel } \Gamma_\kappa\text{)} \\
&= \int_{\mathbb{U}_{\|\kappa\|}} \Phi(u_{\kappa_-}, u_{\kappa_\star}) \mathbf{1}_{\{u_{\kappa_-} = h_{\|\kappa_-}\| \}} \Gamma_\kappa^{\kappa_\star}[du_{\kappa_\star} | u_{\kappa_-}, h] \otimes \Gamma_\kappa^{\kappa_-}[du_{\kappa_-} | h]
\end{aligned}$$

by change of variables  $u_\kappa = (u_{\kappa_-}, u_{\kappa_\star})$ , by property (38) and by disintegration formula (40) for the stochastic kernel  $\Gamma_\kappa$

$$= \int_{\mathbb{U}_{\|\kappa_-}\|} \Gamma_\kappa^{\kappa_-}[du_{\kappa_-} | h] \mathbf{1}_{\{u_{\kappa_-} = h_{\|\kappa_-}\| \}} \int_{\mathbb{U}_{\kappa_\star}} \Phi(u_{\kappa_-}, u_{\kappa_\star}) \Gamma_\kappa^{\kappa_\star}[du_{\kappa_\star} | h_{\|\kappa_-}\|, h]$$

by Fubini Theorem and by substitution  $u_{\kappa_-} = h_{\|\kappa_-\|}$  in the term  $\Gamma_{\kappa^*}^{\kappa^*}[du_{\kappa^*} | u_{\kappa_-}, h]$

$$\begin{aligned}
&= \int_{\mathbb{U}_{\|\kappa\|}} \Gamma_{\kappa}^{\kappa_-}[du_{\kappa_-} | h] \int_{\mathbb{U}_{\|\kappa\|}} \Phi(u_{\kappa_-}, u_{\kappa^*}) \Gamma_{\kappa}^{\kappa^*}[du_{\kappa^*} | h_{\|\kappa_-\|}, h] \\
&\hspace{15em} \text{(by property (38) for the stochastic kernel } \Gamma_{\kappa} \text{)} \\
&= \int_{\mathbb{U}_{\|\kappa\|}} \Gamma_{\kappa}^{\kappa_-}[du_{\kappa_-} | h] \int_{\mathbb{W}_{\kappa^*}} \ell_{\kappa^*}(dw'_{\kappa^*}) \Phi(u_{\kappa_-}, m_{\kappa^*}^{p'}(w'_{\kappa^*}, h)) \\
&\hspace{15em} \text{(by property (41) of the mapping } m_{\kappa^*}^{p'} \text{)} \\
&= \int_{\mathbb{W} \times \Omega} (\ell \otimes \nu)^{|Z_{\kappa^*} \circ T_m}[dw'' d\omega'' | Z_{\kappa^*}(h)] \int_{\mathbb{W}_{\kappa^*}} \ell_{\kappa^*}(dw'_{\kappa^*}) \Phi(m_{\kappa_-}^p(w''^p, h), m_{\kappa^*}^{p'}(w'_{\kappa^*}, h)) . \\
&\hspace{15em} \text{(by property (39) for the stochastic kernel } \Gamma_{\kappa} \text{)}
\end{aligned}$$

Now, we show that there exists a measurable function  $\Psi' : (\mathbb{W} \times \mathbb{Z}_{\kappa^*}, \mathcal{W} \otimes \mathcal{Z}_{\kappa^*}) \rightarrow (\mathbb{R}, \mathcal{B}_{\mathbb{R}})$  such that

$$\begin{aligned}
\Psi'(w'', Z_{\kappa^*}(h)) &= \mathbf{1}_{\mathbb{H}_{\kappa}^{\varphi}}(h) \int_{\mathbb{W}_{\kappa^*}} \ell_{\kappa^*}(dw'_{\kappa^*}) \Phi(m_{\kappa_-}^p(w''^p, h), m_{\kappa^*}^{p'}(w'_{\kappa^*}, h)) , \\
&\hspace{15em} \forall (w'', h) \in \mathbb{W} \times \mathbb{H} .
\end{aligned} \tag{44}$$

Indeed, the function  $\mathbb{W}^p \times \mathbb{W}_{\kappa^*} \times \mathbb{H} \ni (w'', w'_{\kappa^*}, h) \mapsto \mathbf{1}_{\mathbb{H}_{\kappa}^{\varphi}}(h) \Phi(m_{\kappa_-}^p(w''^p, h), m_{\kappa^*}^{p'}(w'_{\kappa^*}, h))$  is measurable with respect to  $\mathcal{W}^p \otimes \mathcal{W}_{\kappa^*} \otimes (\mathbb{H}_{\kappa}^{\varphi} \cap (\bigvee_{a \in \|\kappa\|} \mathcal{J}_a))$  by definition (21a) of an A-mixed strategy and by definition of the trace field  $\mathbb{H}_{\kappa}^{\varphi} \cap (\bigvee_{a \in \|\kappa\|} \mathcal{J}_a)$ , hence with respect to  $\mathcal{W}^p \otimes \mathcal{W}_{\kappa^*} \otimes (\mathbb{H}_{\kappa}^{\varphi} \cap (\mathcal{C}_{\|\kappa_-\|} \vee \mathcal{J}_{\kappa^*}))$  by definition (30b) of  $\mathcal{C}_{\|\kappa_-\|}$ , hence with respect to  $\mathcal{W}^p \otimes \mathcal{W}_{\kappa^*} \otimes (\mathcal{J}_{\kappa^*} \vee (\mathbb{H}_{\kappa}^{\varphi} \cap \mathcal{J}_{\kappa^*}))$  by perfect recall (30), hence to  $\mathcal{W}^p \otimes \mathcal{W}_{\kappa^*} \otimes \mathcal{J}_{\kappa^*}$  as  $\mathbb{H}_{\kappa}^{\varphi} \in \mathcal{J}_{\kappa^*}$  by (30) with  $H = \mathbb{H}$ . By Fubini Theorem, we deduce that the function  $\mathbb{W} \times \mathbb{H} \ni (w, h) \mapsto \mathbf{1}_{\mathbb{H}_{\kappa}^{\varphi}}(h) \int_{\mathbb{W}_{\kappa^*}} \ell_{\kappa^*}(dw'_{\kappa^*}) \Phi(m_{\kappa_-}^p(w''^p, h), m_{\kappa^*}^{p'}(w'_{\kappa^*}, h))$  is measurable with respect to  $\mathcal{W} \otimes \mathcal{J}_{\kappa^*}$ . As a consequence, as  $Z_{\kappa^*}^{-1}(\mathcal{Z}_{\kappa^*}) = \mathcal{J}_{\kappa^*}$  by the Borel measurable functional information assumption (see Definition 15), by Doob functional theorem [7, Chap. 1, p. 18], there exists a measurable function  $\Psi' : (\mathbb{W} \times \mathbb{Z}_{\kappa^*}, \mathcal{W} \otimes \mathcal{Z}_{\kappa^*}) \rightarrow (\mathbb{R}, \mathcal{B}_{\mathbb{R}})$  such that (44) holds true, because  $(\mathbb{W} \times \mathbb{Z}_{\kappa^*}, \mathcal{W} \otimes \mathcal{Z}_{\kappa^*})$  is a product of Borel spaces, hence is a Borel space.

We conclude that

$$\begin{aligned}
&\int_{\mathbb{W} \times \Omega} \ell(dw) \nu(d\omega) \mathbf{1}_{\mathbb{H}_{\kappa}^{\varphi}}(T_m^{\omega}(w)) \Phi(m_{\kappa}^p(w^p, T_m^{\omega}(w))) \\
&= \int_{\mathbb{W} \times \Omega} \ell(dw) \nu(d\omega) \left[ \int_{\mathbb{W} \times \Omega} \mathbf{1}_{\mathbb{H}_{\kappa}^{\varphi}}(h) (\ell \otimes \nu)^{|Z_{\kappa^*} \circ T_m^p}[dw'' d\omega'' | Z_{\kappa^*}(h)] \right. \\
&\hspace{15em} \left. \int_{\mathbb{W}_{\kappa^*}} \ell_{\kappa^*}(dw'_{\kappa^*}) \Phi(m_{\kappa_-}^p(w''^p, h), m_{\kappa^*}^{p'}(w'_{\kappa^*}, h)) \right]_{|h=T_m^{\omega}(w)} \\
&\hspace{15em} \text{(by substitution of the inner integral expression)} \\
&= \int_{\mathbb{W} \times \Omega} \ell(dw) \nu(d\omega) \left[ \int_{\mathbb{W} \times \Omega} (\ell \otimes \nu)^{|Z_{\kappa^*} \circ T_m}[dw'' d\omega'' | z] \Psi'(w'', z) \right]_{|z=Z_{\kappa^*} \circ T_m^{\omega}(w)}
\end{aligned}$$

by property (44) of the function  $\Psi'$

$$= \int_{\mathbb{W} \times \Omega} \ell(dw) \nu(dw) \Psi'(w, Z_{\kappa_*} \circ T_m^\omega(w))$$

by property of regular conditional distributions [9, Th. 6.4]

$$= \int_{\mathbb{W} \times \Omega} \ell(dw) \nu(dw) \mathbf{1}_{\mathbb{H}_\kappa^\varphi}(T_m^\omega(w)) \\ \int_{\mathbb{W}_{\kappa_*}} \ell_{\kappa_*}(dw'_{\kappa_*}) \Phi(w^p, m_{\kappa_-}^p(w^p, T_m^\omega(w)), m'_{\kappa_*}(w'_{\kappa_*}, T_m^\omega(w)))$$

by property (44) of the function  $\Psi'$ .

This ends the proof.  $\square$

The next Lemma 21 is a kind of Kuhn's Theorem, but on the randomizing device space  $\mathbb{W}$  instead of the configuration space  $\mathbb{H}$ . The proof combines the previous Lemma 20 with the playability property of the solution map and an induction.

**Lemma 21** (Equivalence on  $\mathbb{W}^p$ ). *Suppose that the assumptions of Theorem 16 are satisfied, and that the player  $p \in P$  satisfies perfect recall, as in Definition 11. We consider a probability  $\nu$  on  $(\Omega, \mathcal{F})$ , an  $A$ -mixed strategy  $m^p = (m_a^p)_{a \in \mathbb{A}^p}$ , of the player  $p$  and an  $A$ -mixed strategy  $m^{-p} = (m_a^{-p})_{a \in \mathbb{A}^{-p}}$  of the other players. We let  $m'^p = (m'_a{}^p)_{a \in \mathbb{A}^p}$  denote the  $A$ -behavioral strategy of the player  $p$  given by Lemma 19.*

*Then, for any total  $p$ -ordering  $\rho \in \Sigma_{|\mathbb{A}^p|}^p$ , for any bounded measurable function  $J : (\mathbb{H}, \mathcal{H}) \rightarrow (\mathbb{R}, \mathcal{B}_{\mathbb{R}})$ , for any  $\omega \in \Omega$  and  $w^{-p} \in \mathbb{W}^{-p}$ , we have that*

$$\int_{\mathbb{W}^p} \ell^p(dw^p) (\mathbf{1}_{\mathbb{H}_\rho^\varphi} J)(T_{(m^{-p}, m^p)}^\omega(w^{-p}, w^p)) = \int_{\mathbb{W}^p} \ell^p(dw^p) (\mathbf{1}_{\mathbb{H}_\rho^\varphi} J)(T_{(m^{-p}, m'^p)}^\omega(w^{-p}, w^p)). \quad (45)$$

*Proof.* For any total  $p$ -ordering  $\rho \in \Sigma_{|\mathbb{A}^p|}^p$  and any  $p$ -ordering  $\kappa \in \Sigma^p$ , we say that  $\kappa \subset \rho$  if  $\kappa = \psi_{|\kappa|} \rho$  where  $\psi_{|\kappa|}$  has been defined in (27). When  $\kappa \subset \rho$ , we introduce the tail ordering  $\rho \setminus \kappa = (\rho(i))_{i=|\kappa|+1, \dots, |\rho|}$  so that  $\kappa \subset \rho \implies \rho = (\kappa_-, \kappa_*, \rho \setminus \kappa)$ . We also denote  $w_\kappa = (w_a)_{a \in \|\kappa\|}$ ,  $w_{\rho \setminus \kappa} = (w_a)_{a \in \|\rho \setminus \kappa\|}$  and  $\mathbb{W}_{\rho \setminus \kappa} = \prod_{a \in \|\rho \setminus \kappa\|} \mathbb{W}_a$ .

Let  $\omega \in \Omega$  and  $w^{-p} \in \mathbb{W}^{-p}$  be fixed. Let  $\rho \in \Sigma_{|\mathbb{A}^p|}^p$  be a total  $p$ -ordering of the agents in  $\mathbb{A}^p$ . As, by assumption, the  $W$ -game is playable and  $p$ -strongly measurable, for any  $\kappa \subset \rho$  and  $w'^p \in \mathbb{W}^p$ , we get by (17) the existence of a measurable mapping  $\widehat{S}_{(m^{-p}(w^{-p}, \cdot), m_{\rho \setminus \kappa}^p(w^p, \cdot))}^{\|\kappa\|}$  such that

$$S_{(m^{-p}(w^{-p}, \cdot), m_{\rho \setminus \kappa}^p(w^p, \cdot), m'_{\rho \setminus \kappa}(w^p, \cdot))}(\omega) \\ = \widehat{S}_{(m^{-p}(w^{-p}, \cdot), m'_{\rho \setminus \kappa}(w^p, \cdot))}^{\|\kappa\|} \left( \omega, m_{\rho \setminus \kappa}^p(w^p, S_{(m^{-p}(w^{-p}, \cdot), m_{\rho \setminus \kappa}^p(w^p, \cdot), m'_{\rho \setminus \kappa}(w^p, \cdot))}(\omega)) \right),$$

where we have used the shorthand notation  $m_{\kappa}^p = (m_a^p)_{a \in \|\kappa\|}$  and  $m_{\rho/\kappa}^p = (m^p)_{a \in \rho/\kappa}$ .

As  $m^p$  is an A-behavioral strategy, Equation (21b) implies that  $m_{\rho/\kappa}^p(w^p, \cdot)$  only depends on the randomizing component  $w'_{\rho/\kappa} \in \mathbb{W}_{\rho/\kappa}$  and, going back to the original definition (16) we can denote  $\widehat{S}_{(m^{-p}(w^{-p}, \cdot), m_{\kappa}^p(w^p, \cdot), m_{\rho/\kappa}^p(w'_{\rho/\kappa}, \cdot))}^{\|\kappa\|} = \widehat{S}_{(m^{-p}(w^{-p}, \cdot), m_{\rho/\kappa}^p(w'_{\rho/\kappa}, \cdot))}^{\|\kappa\|}$ , obtaining thus that

$$\begin{aligned} & S_{(m^{-p}(w^{-p}, \cdot), m_{\kappa}^p(w^p, \cdot), m_{\rho/\kappa}^p(w'_{\rho/\kappa}, \cdot))}(\omega) \\ &= \widehat{S}_{(m^{-p}(w^{-p}, \cdot), m_{\rho/\kappa}^p(w'_{\rho/\kappa}, \cdot))}^{\|\kappa\|} \left( \omega, m_{\kappa}^p(w^p, S_{(m^{-p}(w^{-p}, \cdot), m_{\kappa}^p(w^p, \cdot), m_{\rho/\kappa}^p(w'_{\rho/\kappa}, \cdot))}(\omega)) \right). \end{aligned} \quad (46)$$

For any  $p$ -ordering  $\kappa \in \Sigma^p$  such that  $\kappa \subset \rho$ , we prove that the following quantity

$$\theta(\kappa) = \int_{\mathbb{W}_{\rho/\kappa}} \ell_{\rho/\kappa}(dw'_{\rho/\kappa}) \int_{\mathbb{W}^p} \ell^p(dw^p) (\mathbf{1}_{\mathbb{H}_{\rho}^{\varphi}} J) (S_{(m^{-p}(w^{-p}, \cdot), m_{\kappa}^p(w^p, \cdot), m_{\rho/\kappa}^p(w'_{\rho/\kappa}, \cdot))}(\omega)) \quad (47)$$

is equal to  $\theta(\kappa_-)$ . This proves the desired result as

$$\begin{aligned} \theta(\rho) &= \int_{\mathbb{W}^p} \ell^p(dw^p) (\mathbf{1}_{\mathbb{H}_{\rho}^{\varphi}} J) (T_{(m^{-p}, m^p)}^{\omega}(w^{-p}, w^p)), \\ \theta(\emptyset) &= \int_{\mathbb{W}^p} \ell^p(dw^p) (\mathbf{1}_{\mathbb{H}_{\rho}^{\varphi}} J) (T_{(m^{-p}, m^p)}^{\omega}(w^{-p}, w^p)), \end{aligned}$$

where the notation  $\emptyset$  in  $\theta(\emptyset)$  refers to the convention that  $\kappa_- = \emptyset \in \Sigma_0^p = \{\emptyset\}$  when  $\kappa \in \Sigma_1^p$ .

First, we focus on the inner integral in (47): for fixed  $w'_{\rho/\kappa} \in \mathbb{W}_{\rho/\kappa}$ , we have that

$$\begin{aligned} & \int_{\mathbb{W}^p} \ell^p(dw^p) (\mathbf{1}_{\mathbb{H}_{\rho}^{\varphi}} J) (S_{(m^{-p}(w^{-p}, \cdot), m_{\kappa}^p(w^p, \cdot), m_{\rho/\kappa}^p(w'_{\rho/\kappa}, \cdot))}(\omega)) \\ &= \int_{\mathbb{W}^p} \ell^p(dw^p) \left[ (\mathbf{1}_{\mathbb{H}_{\rho}^{\varphi}} J) \left( \widehat{S}_{(m^{-p}(w^{-p}, \cdot), m_{\rho/\kappa}^p(w'_{\rho/\kappa}, \cdot))}^{\|\kappa\|} (\omega, m_{\kappa}^p(w^p, h)) \right) \right]_{h=} \\ & \quad S_{(m^{-p}(w^{-p}, \cdot), m_{\kappa}^p(w^p, \cdot), m_{\rho/\kappa}^p(w'_{\rho/\kappa}, \cdot))}(\omega) \quad (\text{by (46)}) \\ &= \int_{\mathbb{W}^p} \ell^p(dw^p) \left[ (\mathbf{1}_{\mathbb{H}_{\rho}^{\varphi}} J) \left( \widehat{S}_{(m^{-p}(w^{-p}, \cdot), m_{\rho/\kappa}^p(w'_{\rho/\kappa}, \cdot))}^{\|\kappa\|} (\omega, m_{\kappa_-}^p(w^p, h), m_{\kappa_{\star}}^p(w^p, h)) \right) \right]_{h=} \\ & \quad S_{(m^{-p}(w^{-p}, \cdot), m_{\kappa_-}^p(w^p, \cdot), m_{\kappa_{\star}}^p(w^p, \cdot), m_{\rho/\kappa}^p(w'_{\rho/\kappa}, \cdot))}(\omega) \\ & \quad (\text{by using the decomposition } m_{\kappa}^p = (m_{\kappa_-}^p, m_{\kappa_{\star}}^p)) \\ &= \int_{\mathbb{W}^p} \ell^p(dw^p) \left[ \mathbf{1}_{\mathbb{H}_{\kappa}^{\varphi}}(h) \Phi(m_{\kappa_-}^p(w^p, h), m_{\kappa_{\star}}^p(w^p, h)) \right]_{h=} \\ & \quad S_{(m^{-p}(w^{-p}, \cdot), m_{\kappa_-}^p(w^p, \cdot), m_{\kappa_{\star}}^p(w^p, \cdot), m_{\rho/\kappa}^p(w'_{\rho/\kappa}, \cdot))}(\omega) \end{aligned}$$

where we have used the property  $\mathbf{1}_{\mathbb{H}_\kappa^\varphi} \mathbf{1}_{\mathbb{H}_\rho^\varphi} = \mathbf{1}_{\mathbb{H}_\rho^\varphi}$  since  $\mathbb{H}_\rho^\varphi \subset \mathbb{H}_\kappa^\varphi$  as  $\kappa \subset \rho$ , and where we have dropped the variables  $\omega, w^{-p}, w', w'_{\rho\kappa}$  that do not contribute to the integration (to the difference of  $w^p$ ) inside the notation

$$\Phi(u_{\kappa_-}, u_{\kappa_*}) = (\mathbf{1}_{\mathbb{H}_\rho^\varphi} J) \left( \widehat{S}_{(m^{-p}(w^{-p}, \cdot), m'_{\rho\kappa}(w'_{\rho\kappa}, \cdot))}^{\|\kappa\|}(\omega, (u_{\kappa_-}, u_{\kappa_*})) \right),$$

where the function  $\Phi : \mathbb{U}_{\|\kappa\|} \rightarrow \mathbb{R}$  is bounded measurable — as  $\mathbf{1}_{\mathbb{H}_\rho^\varphi}$  is measurable by (30a), as the function  $J$  is measurable by assumption, and as the mapping  $\widehat{S}_{(m^{-p}(w^{-p}, \cdot), m'_{\rho\kappa}(w'_{\rho\kappa}, \cdot))}^{\|\kappa\|}$  is measurable by assumption that the W-game is playable and  $p$ -strongly measurable

$$\begin{aligned} &= \int_{\mathbb{W}^p} \ell^p(dw^p) \int_{\mathbb{W}_{\kappa_*}} \ell_{\kappa_*}(dw'_{\kappa_*}) \\ &\quad \left[ \mathbf{1}_{\mathbb{H}_\kappa^\varphi}(h) \Phi(m_{\kappa_-}^p(w^p, h), m'_{\kappa_*}(w'_{\kappa_*}, h)) \right]_{h=} \\ &\quad S_{(m^{-p}(w^{-p}, \cdot), m_{\kappa_-}^p(w^p, \cdot), m'_{\kappa_*}(w'_{\kappa_*}, \cdot), m'_{\rho\kappa}(w'_{\rho\kappa}, \cdot))}(\omega) \end{aligned}$$

by using Lemma 20 making possible the substitution (42) where the term  $m_{\kappa_*}^p(w^p, h)$  has been replaced by  $m'_{\kappa_*}(w'_{\kappa_*}, h)$  inside a new integral

$$\begin{aligned} &= \int_{\mathbb{W}^p} \ell^p(dw^p) \int_{\mathbb{W}_{\kappa_*}} \ell_{\kappa_*}(dw'_{\kappa_*}) \\ &\quad \left[ \mathbf{1}_{\mathbb{H}_\kappa^\varphi}(h) \Phi(m_{\kappa_-}^p(w^p, h), m'_{\kappa_*}(w'_{\kappa_*}, h)) \right]_{h=} \\ &\quad S_{(m^{-p}(w^{-p}, \cdot), m_{\kappa_-}^p(w^p, \cdot), m'_{\kappa_*}(w'_{\kappa_*}, \cdot), m'_{\rho\kappa}(w'_{\rho\kappa}, \cdot))}(\omega) \end{aligned}$$

where, in the expression  $h = S_{(m^{-p}(w^{-p}, \cdot), m_{\kappa_-}^p(w^p, \cdot), m_{\kappa_*}^p(w'_{\kappa_*}, \cdot), m_{\rho_{\kappa_*}}^p(w'_{\rho_{\kappa_*}}, \cdot))}(\omega)$ , the term  $m_{\kappa_*}^p(w^p, \cdot)$  has been substituted for  $m_{\kappa_*}^p(w'_{\kappa_*}, \cdot)$  by Proposition 5 because the function  $\mathbb{H} \ni h \mapsto \mathbf{1}_{\mathbb{H}_\kappa^\varphi}(h) \Phi(m_{\kappa_-}^p(w^p, h), m_{\kappa_*}^p(w'_{\kappa_*}, h))$  is  $\mathcal{J}_{\kappa_*}$ -measurable; indeed, the function is measurable with respect to  $\mathbb{H}_\kappa^\varphi \cap (\bigvee_{a \in \|\kappa\|} \mathcal{J}_a)$  by definition (21a) of an A-mixed strategy (recall that  $m_{\kappa_*}^p(w'_{\kappa_*}, \cdot)$  is  $\mathcal{J}_{\kappa_*}$ -measurable by Lemma 19) and by definition of the trace field  $\mathbb{H}_\kappa^\varphi \cap (\bigvee_{a \in \|\kappa\|} \mathcal{J}_a)$ , hence with respect to  $\mathbb{H}_\kappa^\varphi \cap (\mathcal{C}_{\|\kappa_-\|} \vee \mathcal{J}_{\kappa_*})$  by definition (30b) of  $\mathcal{C}_{\|\kappa_-\|}$ , hence with respect to  $\mathcal{J}_{\kappa_*} \vee (\mathbb{H}_\kappa^\varphi \cap \mathcal{J}_{\kappa_*})$  by perfect recall (30), hence with respect to  $\mathcal{J}_{\kappa_*}$  as  $\mathbb{H}_\kappa^\varphi \in \mathcal{J}_{\kappa_*}$  by (30) with  $H = \mathbb{H}$

$$\begin{aligned}
&= \int_{\mathbb{W}^p} \ell^p(dw^p) \int_{\mathbb{W}_{\kappa_*}} \ell_{\kappa_*}(dw'_{\kappa_*}) \left[ (\mathbf{1}_{\mathbb{H}_\rho^\varphi} J) \right. \\
&\quad \left. \left( \widehat{S}_{(m^{-p}(w^{-p}, \cdot), m_{\rho_{\kappa_*}}^p(w'_{\rho_{\kappa_*}}, \cdot))}(\omega, m_{\kappa_-}^p(w^p, h), m_{\kappa_*}^p(w'_{\kappa_*}, h)) \right) \right]_{h=} \\
&\quad S_{(m^{-p}(w^{-p}, \cdot), m_{\kappa_-}^p(w^p, \cdot), m_{\kappa_*}^p(w'_{\kappa_*}, \cdot), m_{\rho_{\kappa_*}}^p(w'_{\rho_{\kappa_*}}, \cdot))}(\omega) \\
&\quad \text{(by definition of the function } \Phi) \\
&= \int_{\mathbb{W}^p} \ell^p(dw^p) \int_{\mathbb{W}_{\kappa_*}} \ell_{\kappa_*}(dw'_{\kappa_*}) (\mathbf{1}_{\mathbb{H}_\rho^\varphi} J) \left( S_{(m^{-p}(w^{-p}, \cdot), m_{\kappa_-}^p(w^p, \cdot), m_{\kappa_*}^p(w'_{\kappa_*}, \cdot), m_{\rho_{\kappa_*}}^p(w'_{\rho_{\kappa_*}}, \cdot))}(\omega) \right)
\end{aligned}$$

by formula (46), but with  $(m^{-p}(w^{-p}, \cdot), m_{\kappa_-}^p(w^p, \cdot), m_{\rho_{\kappa_*}}^p(w'_{\rho_{\kappa_*}}, \cdot))$  replaced by  $(m^{-p}(w^{-p}, \cdot), m_{\kappa_-}^p(w^p, \cdot), m_{\kappa_*}^p(w'_{\kappa_*}, \cdot), m_{\rho_{\kappa_*}}^p(w'_{\rho_{\kappa_*}}, \cdot))$ .

Thus, inserting this last expression in the right hand side of Equation (47), we conclude that

$$\begin{aligned}
\theta(\kappa) &= \int_{\mathbb{W}_{\rho_{\kappa_*}}} \ell_{\rho_{\kappa_*}}(dw'_{\rho_{\kappa_*}}) \int_{\mathbb{W}^p} \ell^p(dw^p) \\
&\quad \int_{\mathbb{W}_{\kappa_*}} \ell_{\kappa_*}(dw'_{\kappa_*}) (\mathbf{1}_{\mathbb{H}_\rho^\varphi} J) \left( S_{(m^{-p}(w^{-p}, \cdot), m_{\kappa_-}^p(w^p, \cdot), m_{\kappa_*}^p(w'_{\kappa_*}, \cdot), m_{\rho_{\kappa_*}}^p(w'_{\rho_{\kappa_*}}, \cdot))}(\omega) \right) \\
&= \int_{\mathbb{W}_{\kappa_*}} \ell_{\kappa_*}(dw'_{\kappa_*}) \int_{\mathbb{W}_{\rho_{\kappa_*}}} \ell_{\rho_{\kappa_*}}(dw'_{\rho_{\kappa_*}}) \int_{\mathbb{W}^p} \ell^p(dw^p) \\
&\quad (\mathbf{1}_{\mathbb{H}_\rho^\varphi} J) \left( S_{(m^{-p}(w^{-p}, \cdot), m_{\kappa_-}^p(w^p, \cdot), m_{\kappa_*}^p(w'_{\kappa_*}, \cdot), m_{\rho_{\kappa_*}}^p(w'_{\rho_{\kappa_*}}, \cdot))}(\omega) \right) \\
&\quad \text{(by Fubini Theorem)} \\
&= \int_{\mathbb{W}_{\kappa_*} \times \mathbb{W}_{\rho_{\kappa_*}}} (\ell_{\kappa_*} \otimes \ell_{\rho_{\kappa_*}})(dw'_{\kappa_*} dw'_{\rho_{\kappa_*}}) \int_{\mathbb{W}^p} \ell^p(dw^p) \\
&\quad (\mathbf{1}_{\mathbb{H}_\rho^\varphi} J) \left( S_{(m^{-p}(w^{-p}, \cdot), m_{\kappa_-}^p(w^p, \cdot), m_{\kappa_*}^p(w'_{\kappa_*}, \cdot), m_{\rho_{\kappa_*}}^p(w'_{\rho_{\kappa_*}}, \cdot))}(\omega) \right)
\end{aligned}$$

by Fubini Theorem and by definition of the product probability  $\ell_{\kappa_*} \otimes \ell_{\rho_{\kappa_*}}$

$$= \int_{\mathbb{W}_{\rho_{\kappa_*}}} \ell_{\rho_{\kappa_*}}(dw'_{\rho_{\kappa_*}}) \int_{\mathbb{W}^p} \ell^p(dw^p) (\mathbf{1}_{\mathbb{H}_\rho^\varphi} J) \left( S_{(m^{-p}(w^{-p}, \cdot), m_{\kappa_-}^p(w^p, \cdot), m_{\rho_{\kappa_*}}^p(w'_{\rho_{\kappa_*}}, \cdot))}(\omega) \right)$$



by changes of variables  $(w'_{\kappa_\star}, w'_{\rho\kappa_-}) = w'_{\rho\kappa_-}$  and  $\rho \setminus \kappa_- = (\kappa_\star, \rho \setminus \kappa)$

$$= \theta(\kappa_-).$$

This ends the proof. □

**Proof of Theorem 16.**

*Proof.* To prove (35), we consider a bounded measurable function  $J : (\mathbb{H}, \mathcal{H}) \rightarrow (\mathbb{R}, \mathcal{B}_{\mathbb{R}})$ , and we proceed with

$$\begin{aligned} & \int_{\mathbb{H}} J(h) \mathbb{Q}_{(m^{-p}, m^p)}^\nu(dh) \\ &= \int_{\Omega} d\nu(\omega) \int_{\mathbb{W}^{-p} \times \mathbb{W}^p} \ell^{-p}(dw^{-p}) \otimes \ell^p(dw^p) J(T_{(m^{-p}, m^p)}^\omega(w^{-p}, w^p)) \end{aligned}$$

by the pushforward probability formula (24) and by detailing the product structures of  $\mathbb{W}$  and  $\ell$  in (20)

$$\begin{aligned} &= \int_{\Omega} d\nu(\omega) \int_{\mathbb{W}^{-p}} \ell^{-p}(dw^{-p}) \left[ \int_{\mathbb{W}^p} \ell^p(dw^p) J(T_{(m^{-p}, m^p)}^\omega(w^{-p}, w^p)) \right] \quad (\text{by Fubini Theorem}) \\ &= \int_{\Omega} d\nu(\omega) \int_{\mathbb{W}^{-p}} \ell^{-p}(dw^{-p}) \sum_{\rho \in \Sigma_{|\mathbb{A}^p|}^p} \left[ \int_{\mathbb{W}^p} \ell^p(dw^p) (\mathbf{1}_{\mathbb{H}_\rho^\varphi} J)(T_{(m^{-p}, m^p)}^\omega(w^{-p}, w^p)) \right] \end{aligned}$$

since the subsets  $\mathbb{H}_\rho^\varphi$  in (29) are pairwise disjoint when the total ordering  $\rho$  varies in  $\Sigma_{|\mathbb{A}^p|}^p$ , and their union is  $\mathbb{H}$

$$\begin{aligned}
&= \int_{\Omega} d\nu(\omega) \int_{\mathbb{W}^{-p}} \ell^{-p}(dw^{-p}) \sum_{\rho \in \Sigma_{|\mathbb{A}^p|}^p} \left[ \int_{\mathbb{W}^p} \ell^p(dw'^p) \int_{\mathbb{W}^p} \ell^p(dw^p) (\mathbf{1}_{\mathbb{H}_\rho^\varphi} J)(T_{(m^{-p}, m'^p)}^\omega(w^{-p}, w'^p)) \right] \\
&\hspace{25em} \text{(by (45) in the substitution Lemma 21)} \\
&= \int_{\Omega} d\nu(\omega) \int_{\mathbb{W}^p} \ell^p(dw'^p) \int_{\mathbb{W}^{-p}} \ell^{-p}(dw^{-p}) \sum_{\rho \in \Sigma_{|\mathbb{A}^p|}^p} \left[ \int_{\mathbb{W}^p} \ell^p(dw^p) (\mathbf{1}_{\mathbb{H}_\rho^\varphi} J)(T_{(m^{-p}, m'^p)}^\omega(w^{-p}, w'^p)) \right] \\
&\hspace{25em} \text{(by Fubini Theorem)} \\
&= \int_{\Omega} d\nu(\omega) \int_{\mathbb{W}^p} \ell^p(dw'^p) \int_{\mathbb{W}^{-p}} \ell^{-p}(dw^{-p}) \int_{\mathbb{W}^p} \ell^p(dw^p) J(T_{(m^{-p}, m'^p)}^\omega(w^{-p}, w'^p)) \\
&= \int_{\Omega} d\nu(\omega) \int_{\mathbb{W}^p} \ell^p(dw'^p) \int_{\mathbb{W}^{-p}} \ell^{-p}(dw^{-p}) J(T_{(m^{-p}, m'^p)}^\omega(w^{-p}, w'^p)) \\
&= \int_{\Omega} d\nu(\omega) \int_{\mathbb{W}^{-p} \times \mathbb{W}^p} \ell^{-p}(dw^{-p}) \otimes \ell^p(dw'^p) J(T_{(m^{-p}, m'^p)}^\omega(w^{-p}, w'^p)) \quad \text{(by Fubini Theorem)} \\
&= \int_{\Omega} d\nu(\omega) \int_{\mathbb{W}} d\ell(w) J(T_{(m^{-p}, m'^p)}^\omega(w)) \quad \text{(as } \ell = \ell^{-p} \otimes \ell^p \text{ and } \mathbb{W} = \mathbb{W}^{-p} \times \mathbb{W}^p \text{ by (20))} \\
&= \int_{\mathbb{H}} J(h) \mathbb{Q}'_{(m^{-p}, m'^p)}(dh). \quad \text{(by the pushforward probability formula (24))}
\end{aligned}$$

This ends the proof.  $\square$

## 5.2 Proof of Theorem 17

We start with Lemma 22, gives constraints on the marginals of the pushforward probability induced by any A-behavioral strategy  $m^p$  of the player  $p$  satisfying Equation (35).

**Lemma 22.** *We consider a playable and measurable  $W$ -game (see Definition 7). We focus on the player  $p \in P$  and we suppose that  $\mathbb{A}^p$  is a finite set. Let be given a probability  $\nu$  on  $(\Omega, \mathcal{F})$ , an A-mixed strategy  $m^{-p} = (m_a)_{a \in \mathbb{A}^{-p}}$  of the other players, an A-mixed strategy  $m^p = (m_a^p)_{a \in \mathbb{A}^p}$ , of the player  $p$ , and an A-behavioral strategy  $m'^p = (m_a'^p)_{a \in \mathbb{A}^p}$  of the player  $p$ . We set*

$$\mathbb{W}'_a[h] = \{w_a \in \mathbb{W}_a \mid m_a'^p(w_a, h) = h_a\}, \quad \forall a \in \mathbb{A}^p, \quad \forall h \in \mathbb{H}. \quad (49)$$

Then, we have the following implication, for any  $h \in \mathbb{H}$ ,

$$\mathbb{Q}'_{(m^{-p}, m^p)} = \mathbb{Q}'_{(m^{-p}, m'^p)} \quad \text{and} \quad \mathbb{Q}'_{(m^{-p}, m^p)}(h) > 0 \implies \ell^a(\mathbb{W}'_a[h]) > 0, \quad \forall a \in \mathbb{A}^p. \quad (50)$$

*Proof.* Let a configuration  $h \in \mathbb{H}$  be given. Then, we have that

$$\mathbb{Q}'_{(m^{-p}, m^p)}(h) = (\ell^{-p} \otimes \bigotimes_{a \in \mathbb{A}^p} \ell_a \otimes \nu) \left\{ (w, \omega) \in \mathbb{W}^{-p} \times \prod_{a \in \mathbb{A}^p} \mathbb{W}_a \times \Omega \mid T_{m^p}^\omega(w) = h \right\}$$

by definition (24) of the pushforward probability  $\mathbb{Q}'_{(m^{-p}, m^p)}$  and by (20)

$$= (\ell^{-p} \otimes \bigotimes_{a \in \mathbb{A}^p} \ell_a \otimes \nu) \left\{ (w, \omega) \in \mathbb{W}^{-p} \times \prod_{a \in \mathbb{A}^p} \mathbb{W}_a \times \Omega \mid \omega = h_\emptyset, \right. \\ \left. m_{\mathbb{A}^q}(w^q, h) = h_{\mathbb{A}^q}, \forall q \in P \setminus \{p\}, m_a^p(w_a, h) = h_a, \forall a \in \mathbb{A}^p \right\}$$

by solution map property (14b) and by definition (22) of  $T_{m^p}^\omega(w)$

$$= \nu\{h_\emptyset\} \times \prod_{q \in P \setminus \{p\}} \ell^q \{w^q \in \mathbb{W}^q \mid m_{\mathbb{A}^q}(w^q, h) = h_{\mathbb{A}^q}\} \\ \times \prod_{a \in \mathbb{A}^p} \ell_a \{w_a \in \mathbb{W}_a \mid m_a^p(w_a, h) = h_a\} \\ \text{(by definition of a product probability)} \\ = \nu\{h_\emptyset\} \times \prod_{q \in P \setminus \{p\}} \ell^q \{w^q \in \mathbb{W}^q \mid m_{\mathbb{A}^q}(w^q, h) = h_{\mathbb{A}^q}\} \times \prod_{a \in \mathbb{A}^p} \ell_a(\mathbb{W}'_a[h]) . \\ \text{(by definition of } \mathbb{W}'_a[h] \text{ in (49))}$$

As a consequence, if  $\mathbb{Q}'_{(m^{-p}, m^p)} = \mathbb{Q}'_{(m^{-p}, m^p)}$  and  $\mathbb{Q}'_{(m^{-p}, m^p)}(h) > 0$ , we deduce that the nonnegative quantity  $\ell_a(\mathbb{W}'_a[h])$  must be positive for all  $a \in \mathbb{A}^p$ .

We have proven (50) and this ends the proof.  $\square$

### Proof of Theorem 17.

*Proof.* We consider a playable and measurable W-game (see Definition 7). We focus on the player  $p \in P$  and we suppose that she satisfies the Borel measurable functional information assumption (see Definition 15) and partial causality (see Definition 12), that  $\mathbb{A}^p$  is a finite set, and that  $\mathbb{U}_a$  contains at least two distinct elements, for all  $a \in \mathbb{A}^p$ .

By assumption (see Equation (36)), we have that, for the  $p$ -configuration-ordering  $\varphi : \mathbb{H} \rightarrow \Sigma^p$  given by Definition 12, there exists a  $p$ -ordering  $\kappa \in \Sigma^p$  such that

$$\exists h^+, h^- \in \mathbb{H}_\kappa^\varphi, Z_{\kappa_\star}(h^+) = Z_{\kappa_\star}(h^-), (Z_a(h^+), h_a^+)_{a \in \|\kappa_-\|} \neq (Z_a(h^-), h_a^-)_{a \in \|\kappa_-\|} .$$

Therefore, setting  $j_c = |\kappa| \geq 2$  (because the case  $|\kappa| = 1$  is void) and  $c = \kappa(j_c) = \kappa_\star$ , we deduce that one of the following two exclusive cases holds true:

1. either there exists  $h^+, h^- \in \mathbb{H}_\kappa^\varphi$  such that  $Z_c(h^+) = Z_c(h^-)$ , and there exists an agent  $b \in \|\kappa_-\|$  such that  $h_b^+ \neq h_b^-$ ,
2. or  $Z_c(h) = Z_c(h') \implies h_a = h'_a$ , for all  $h, h' \in \mathbb{H}_\kappa^\varphi$  and for all  $a \in \|\kappa_-\|$ , and there exists  $h^+, h^- \in \mathbb{H}_\kappa^\varphi$  such that  $Z_c(h^+) = Z_c(h^-)$ , and there exists an agent  $b \in \|\kappa_-\|$  such that  $Z_b(h^+) \neq Z_b(h^-)$ .

In both cases, we denote  $h^+ = (\omega^+, u^+)$  and  $h^- = (\omega^-, u^-)$ . For any mixed strategy  $m_c^p$  of the agent  $c$ , we have that  $m_c^p(w_c, h^+) = m_c^p(w_c, h^-)$  since the mapping  $\mathbb{H} \ni h \mapsto m_c^p(w_c, h)$  is  $\mathcal{I}_c$ -measurable, since  $Z_c^{-1}(\mathcal{Z}_c) = \mathcal{I}_c$  and  $Z_c(h^+) = Z_c(h^-)$ . Without loss of generality, we can suppose that  $h_c^+ = u_c^+ \neq u_c^- = h_c^-$ . Indeed, as the player  $p$  satisfies partial causality, we have that  $\mathbb{H}_\kappa^p \cap \{h \in \mathbb{H} \mid Z_c(h) = Z_c(h^-)\} \in \mathcal{H}_{\|\kappa_-\|}^p$  by (32), so that  $Z_c((h_c^-, u_c)) = Z_c(h^-)$  for any  $u_c \in \mathbb{U}_c$ , and we choose  $u_c \neq u_c^+$ .

Now for the proof, by contradiction. We set  $\tilde{P} = P \setminus \{p\}$  and, in both cases above, we consider the same A-mixed strategy  $m^{-p} = (m^q)_{q \in \tilde{P}}$  for the other players than player  $p$ . For this purpose, we introduce, for any player  $q \in \tilde{P}$ , a partition  $\mathbb{W}_q^+$  and  $\mathbb{W}_q^-$  of  $\mathbb{W}^q$  with  $\ell^q(\mathbb{W}_q^+) = \ell^q(\mathbb{W}_q^-) = 1/2$ , and we define the A-mixed strategy  $m^q = (m_a^q)_{a \in \mathbb{A}^q}$  by

$$m_a^q(w^q, h) = h_a^\epsilon, \quad \forall q \in \tilde{P}, \quad \forall a \in \mathbb{A}^q, \quad \forall \epsilon \in \{+, -\}, \quad \forall w^q \in \mathbb{W}_q^\epsilon, \quad \forall h \in \mathbb{H}. \quad (51)$$

Notice that the above definition is valid even if  $h_a^+ = h_a^-$ , and that, for any fixed  $w^q \in \mathbb{W}_q^\epsilon$ , the pure strategy profile  $m^q(w^q, \cdot)$  is a constant mapping with value  $(h_a^\epsilon)_{a \in \mathbb{A}^q}$ .

We consider the first case where there exists an agent  $b \in \|\kappa_-\|$  such that  $h_b^+ \neq h_b^-$ . We can always suppose that  $b = \kappa(j_b)$  where  $j_b = \inf\{j \in \llbracket 1, j_c - 1 \rrbracket \mid h_{\kappa(j)}^+ \neq h_{\kappa(j)}^-\}$  so that  $h_{\kappa(j)}^+ = h_{\kappa(j)}^-$ , for all  $j \in \llbracket 1, j_b - 1 \rrbracket$  (the empty set if  $j_b = 1$ ). We define the A-mixed strategy  $m^p = (m_a^p)_{a \in \mathbb{A}^p}$  of player  $p$  in the same way than for the other players: we introduce a partition  $\mathbb{W}_p^+$  and  $\mathbb{W}_p^-$  of  $\mathbb{W}^p$  with  $\ell^p(\mathbb{W}_p^+) = \ell^p(\mathbb{W}_p^-) = 1/2$ , and we define

$$m_a^p(w^p, h) = h_a^\epsilon, \quad \forall a \in \mathbb{A}^p, \quad \forall \epsilon \in \{+, -\}, \quad \forall w^p \in \mathbb{W}_p^\epsilon, \quad \forall h \in \mathbb{H}. \quad (52)$$

Then, we consider any probability distribution  $\nu$  on  $\Omega$  such that  $\nu\{\omega^+\} > 0$ ,  $\nu\{\omega^-\} > 0$  and  $\nu\{\omega^+, \omega^-\} = 1$ , thus covering both cases where  $\omega^+ = \omega^-$  or  $\omega^+ \neq \omega^-$ .

On the one hand, as, for any player  $q \in P$  and for any  $w^q \in \mathbb{W}_q^+$  (resp.  $w^q \in \mathbb{W}_q^-$ ) the pure strategy profile  $m^q(w^q, \cdot)$  takes the constant value  $(h_a^+)_{a \in \mathbb{A}^q}$  (resp.  $(h_a^-)_{a \in \mathbb{A}^q}$ ) by (51)–(52), we readily get — by definition (22) of  $T_{(m^{-p}, m^p)}^{\omega^\epsilon}(w)$  and by characterization (14b) of the solution map — that

$$\begin{aligned} w \in \prod_{q \in P} \mathbb{W}_q^+ &\implies T_{(m^{-p}, m^p)}^{\omega^+}(w) = (\omega^+, u^+) = h^+, \\ w \in \prod_{q \in P} \mathbb{W}_q^- &\implies T_{(m^{-p}, m^p)}^{\omega^-}(w) = (\omega^-, u^-) = h^-, \end{aligned}$$

hence, as  $\nu\{\omega^\epsilon\} > 0$  and  $\prod_{q \in P} \ell^q(\mathbb{W}_q^\epsilon) = 1/2^{|P|} > 0$  for  $\epsilon \in \{+, -\}$ , that

$$\mathbb{Q}_{(m^{-p}, m^p)}^\nu(h^+) > 0 \quad \text{and} \quad \mathbb{Q}_{(m^{-p}, m^p)}^\nu(h^-) > 0. \quad (53a)$$

On the other hand, we also readily get, in the same way but focusing on (52), that

$$\mathbb{Q}_{(m^{-p}, m^p)}^\nu(h) > 0 \implies \text{either } h_{\mathbb{A}^p} = h_{\mathbb{A}^p}^+ \text{ or } h_{\mathbb{A}^p} = h_{\mathbb{A}^p}^-. \quad (53b)$$

The proof is by contradiction and we suppose that there exists an A-behavioral strategy  $m'^p = (m'_a{}^p)_{a \in \mathbb{A}^p}$  of the player  $p$  such that  $\mathbb{Q}'_{(m^{-p}, m'^p)} = \mathbb{Q}_{(m^{-p}, m'^p)}$ . Applying Lemma 22 to  $h^+$  and  $h^-$ , we obtain that  $\ell_a(\mathbb{W}'_a[h^+]) > 0$  and  $\ell_a(\mathbb{W}'_a[h^-]) > 0$ ,  $\forall a \in \mathbb{A}^p$ . As a consequence, the following set

$$\mathbb{W}^\pm = \prod_{q \in \tilde{P}} \mathbb{W}_q^+ \times \prod_{a \in \mathbb{A}^p \setminus \{c\}} \mathbb{W}'_a[h^+] \times \mathbb{W}'_c[h^-] \quad (54)$$

has positive probability and, for any  $w \in \mathbb{W}^\pm$ , we are going to show that the configuration  $h = T_{(m^{-p}, m'^p)}^{\omega^+}(w)$  contradicts (53b). First, we observe that the configuration  $h$  is such that  $h_{\mathbb{A}^{-p}} = h_{\mathbb{A}^{-p}}^+$  because, for any player  $q \in \tilde{P}$ , the pure strategy profile  $m^q(w^q, \cdot)$  takes the constant value  $(h_a^+)_{a \in \mathbb{A}^q}$  when  $w \in \mathbb{W}^\pm$  by definition (54) of  $\mathbb{W}^\pm$ . Second, we prove by induction on  $j \in \llbracket 1, j_c - 1 \rrbracket$  (where  $j_c = |\kappa| \geq 2$ , hence  $j_c - 1 \geq 1$ ) that  $h_{\kappa(j)} = h_{\kappa(j)}^+$  and that  $h \in \mathbb{H}_{\kappa(1), \dots, \kappa(j-1)}^\varphi$ .

We suppose that  $j \geq 1$  and that  $h_{\kappa(i)} = h_{\kappa(i)}^+$  for all  $i \in \llbracket 1, j - 1 \rrbracket$  and  $h \in \mathbb{H}_{\kappa(1), \dots, \kappa(j-1)}^\varphi$  (the special case  $j = 1$  corresponds to the initialization part of the proof by induction that we cover too). Then, we have that

$$\begin{aligned} h_{\kappa(j)} &= m_{\kappa(j)}'^p(w_{\kappa(j)}, h) && \text{(by solution map property (14b) of } h = T_{(m^{-p}, m'^p)}^{\omega^+}(w)) \\ &= m_{\kappa(j)}'^p(w_{\kappa(j)}, (\omega^+, h_{\mathbb{A}^{-p}}, h_{\kappa(1)}, \dots, h_{\kappa(j-1)})) \end{aligned}$$

by partial causality property (33a) and short notation (33b), using Lemma 13 as  $h_\emptyset = \omega^+$  and  $h \in \mathbb{H}_{\kappa(1), \dots, \kappa(j-1)}^\varphi$  by induction assumption (remaining true in the special case  $j = 1$  because  $h_{\mathbb{A}^{-p}} = h_{\mathbb{A}^{-p}}^+$  and  $h \in \mathbb{H}_\emptyset^\varphi = \mathbb{H}$ )

$$= m_{\kappa(j)}'^p(w_{\kappa(j)}, (\omega^+, h_{\mathbb{A}^{-p}}^+, h_{\kappa(1)}^+, \dots, h_{\kappa(j-1)}^+))$$

as we have seen that  $h_{\mathbb{A}^{-p}} = h_{\mathbb{A}^{-p}}^+$ , and as  $(h_{\kappa(1)}, \dots, h_{\kappa(j-1)}) = (h_{\kappa(1)}^+, \dots, h_{\kappa(j-1)}^+)$  by the induction assumption

$$\begin{aligned} &= m_{\kappa(j)}'^p(w_{\kappa(j)}, h^+) && \text{(by using again partial causality, but with } h^+ \text{ this time)} \\ &= h_{\kappa(j)}^+ \end{aligned}$$

as  $w \in \mathbb{W}^\pm$ , hence  $w_{\kappa(j)} \in \mathbb{W}'_{\kappa(j)}[h^+]$  by definition (54) of the set  $\mathbb{W}^\pm$ , and by definition (49) of the set  $\mathbb{W}'_{\kappa(j)}[h^+]$ . From  $h \in \mathbb{H}_{\kappa(1), \dots, \kappa(j-1)}^\varphi$ ,  $h_{\mathbb{A}^{-p}} = h_{\mathbb{A}^{-p}}^+$  and  $(h_{\kappa(1)}, \dots, h_{\kappa(j)}) = (h_{\kappa(1)}^+, \dots, h_{\kappa(j)}^+)$ , we deduce that  $h \in \mathbb{H}_{\kappa(1), \dots, \kappa(j)}^\varphi$  by partial causality property (33a), using Lemma 13 as  $h^+ \in \mathbb{H}_\kappa^\varphi \subset \mathbb{H}_{\kappa(1), \dots, \kappa(j)}^\varphi$  by definition (29) of  $\mathbb{H}_\kappa^\varphi$ . Thus, the induction is completed and we obtain that  $h_{\|\kappa-\|} = (h_{\kappa(1)}, \dots, h_{\kappa(j_c-1)}) = (h_{\kappa(1)}^+, \dots, h_{\kappa(j_c-1)}^+) = h_{\|\kappa-\|}^+$ , and that  $h \in \mathbb{H}_{\kappa-}^\varphi = \mathbb{H}_{\kappa(1), \dots, \kappa(j_c-1)}^\varphi$ .

Third, we compute

$$\begin{aligned} h_c &= m_c'^p(w_c, h) && \text{(by solution map property (14b) of } h = T_{(m^{-p}, m'^p)}^{\omega^+}(w)) \\ &= m_c'^p(w_c, (\omega^+, h_{\mathbb{A}^{-p}}, h_{\kappa(1)}, \dots, h_{\kappa(j_c-1)})) \end{aligned}$$

by partial causality property (33a), and short notation (33b), using Lemma 13 as  $h^+ \in \mathbb{H}_\kappa^\varphi \subset \mathbb{H}_{\kappa_-}^\varphi$  by definition (29) of  $\mathbb{H}_\kappa^\varphi$ , and as  $c = \kappa(j_c) = \kappa_*$

$$= m_c^{lp}(w_c, (\omega^+, h_{\mathbb{A}^{-p}}^+, h_{\kappa(1)}^+, \dots, h_{\kappa(j_c-1)}^+))$$

as  $h_{\mathbb{A}^{-p}} = h_{\mathbb{A}^{-p}}^+$  and as  $(h_{\kappa(1)}, \dots, h_{\kappa(j_c-1)}) = (h_{\kappa(1)}^+, \dots, h_{\kappa(j_c-1)}^+)$  as proved above by induction

$$\begin{aligned} &= m_c^{lp}(w_c, h^+) \quad (\text{by using again partial causality, but with } h^+ \in \mathbb{H}_\kappa^\varphi \text{ this time}) \\ &= h_c^- \end{aligned}$$

as  $w \in \mathbb{W}^\pm$ , hence  $w_c \in \mathbb{W}'_c[h^-]$  by definition (54) of the set  $\mathbb{W}^\pm$ , and by definition (49) of the set  $\mathbb{W}'_c[h^-]$ . As the set  $\mathbb{W}^\pm$  has positive probability, we conclude that

$$\mathbb{Q}'_{(m^{-p}, m^p)} \{h \in \mathbb{H} \mid h_b = h_b^+, h_c = h_c^-\} > 0.$$

Since  $\mathbb{Q}'_{(m^{-p}, m^p)} = \mathbb{Q}_{(m^{-p}, m^p)}$  by assumption, we deduce that

$$\mathbb{Q}'_{(m^{-p}, m^p)} \{h \in \mathbb{H} \mid h_b = h_b^+, h_c = h_c^-\} > 0.$$

But this contradicts (53b) because  $h_b^+ \neq h_b^-$  and  $h_c^+ \neq h_c^-$ .

We consider the second case where  $Z_c(h) = Z_c(h') \implies h_a = h'_a$ , for all  $h, h' \in \mathbb{H}_\kappa^\varphi$  and for all  $a \in \|\kappa_-\|$ , and there exists  $h^+, h^- \in \mathbb{H}_\kappa^\varphi$  such that  $Z_c(h^+) = Z_c(h^-)$ , and there exists an agent  $b \in \|\kappa_-\|$  such that  $Z_b(h^+) \neq Z_b(h^-)$ . Thus, from  $Z_c(h^+) = Z_c(h^-)$ , we deduce that  $h_a^+ = h_a^-$ , for all  $a \in \|\kappa_-\|$ , that is,  $h_{\|\kappa_-\|}^+ = h_{\|\kappa_-\|}^-$ . There exists an element  $\bar{h}_b \neq h_b^+$  by assumption (action sets have at least two distinct elements). We introduce a partition  $\mathbb{W}_p^+$  and  $\mathbb{W}_p^-$  of  $\mathbb{W}^p$  with  $\ell^p(\mathbb{W}_p^+) = \ell^p(\mathbb{W}_p^-) = 1/2$ , and we define the A-mixed strategy  $m^p = (m_a^p)_{a \in \mathbb{A}^p}$  by

$$m_a^p(w^p, h) = h_a^+, \quad \forall a \in \mathbb{A}^p \setminus \{b, c\}, \quad \forall w^p \in \mathbb{W}^p, \quad \forall h \in \mathbb{H}, \quad (55a)$$

$$m_b^p(w^p, h) = \begin{cases} h_b^+ & \text{if } Z_b(h) = Z_b(h^+) \text{ and } w^p \in \mathbb{W}_p^+, \\ \bar{h}_b & \text{if } Z_b(h) \neq Z_b(h^+) \text{ and } w^p \in \mathbb{W}_p^+, \\ \bar{h}_b & \text{if } Z_b(h) = Z_b(h^+) \text{ and } w^p \in \mathbb{W}_p^-, \\ h_b^+ & \text{if } Z_b(h) \neq Z_b(h^+) \text{ and } w^p \in \mathbb{W}_p^-, \end{cases} \quad (55b)$$

and finally (beware of the interversion in the following expression)

$$m_c^p(w^p, h) = \begin{cases} h_c^- & \text{if } Z_c(h) = Z_c(h^+) \text{ and } w^p \in \mathbb{W}_p^-, \\ h_c^+ & \text{else} \end{cases} \quad (55c)$$

For any agent  $a \in \mathbb{A}^p \setminus \{b, c\}$ , the mapping  $m_a^p(w^p, \cdot)$  is  $\mathcal{J}_a$ -measurable as it is constant by (55a). The mapping  $m_b^p(w^p, \cdot)$  is  $\mathcal{J}_b$ -measurable as it measurably expressed in (55b) as a function of the  $\mathcal{J}_b$ -measurable mapping  $Z_b$ . The same holds true for  $m_c^p(w^p, \cdot)$  in (55c).

Then, we consider any probability distribution  $\nu$  on  $\Omega$  such that  $\nu\{\omega^+\} > 0$ ,  $\nu\{\omega^-\} > 0$  and  $\nu\{\omega^+, \omega^-\} = 1$ , thus covering both cases where  $\omega^+ = \omega^-$  or  $\omega^+ \neq \omega^-$ . As a preliminary result, we prove that

$$\mathbb{Q}_{(m^{-p}, m^p)}^\nu \{h \in \mathbb{H} \mid h \in \mathbb{H}_\kappa^\varphi, Z_b(h) = Z_b(h^+), h_c = h_c^-\} = 0. \quad (56)$$

Indeed, by (55c), any  $h = T_{(m^{-p}, m^p)}^\omega(w) \in \mathbb{H}_\kappa^\varphi$  such that  $h_c = h_c^-$  must be such that both  $Z_c(h) = Z_c(h^+)$  and  $w^p \in \mathbb{W}_p^-$ . But, as  $Z_c(h') = Z_c(h'') \implies h'_a = h''_a$ , for all  $h', h'' \in \mathbb{H}_\kappa^\varphi$  and for all  $a \in \|\kappa_-\|$ , we deduce that  $h_b = h_b^+$ . As  $w^p \in \mathbb{W}_p^-$ , we get by (55b) that necessarily  $Z_b(h) \neq Z_b(h^+)$ . Thus, we have proven (56), and we will now show that any A-behavioral strategy contradicts (56).

First, we get that

$$w \in \prod_{q \in \tilde{P}} \mathbb{W}_q^+ \times \mathbb{W}_p^+ \implies T_{(m^{-p}, m^p)}^{\omega^+}(w) = (\omega^+, h_{\mathbb{A}^{-p}}^+, h_{\mathbb{A}^p \setminus \{b, c\}}^+, h_b^+, h_c^+) = h^+$$

because, for any player  $q \in \tilde{P}$  and for any  $w^q \in \mathbb{W}_q^+$  the pure strategy profile  $m^q(w^q, \cdot)$  takes the constant value  $(h_a^+)_{a \in \mathbb{A}^q}$ , and by the expressions (55a)–(55b)–(55c) of  $m^p(w^p, \cdot)$  when  $w^p \in \mathbb{W}_p^+$ . Now, we have that  $\nu\{\omega^+\} > 0$  and  $\prod_{q \in \tilde{P}} \ell^q(\mathbb{W}_q^+) \times \ell^p(\mathbb{W}_p^+) = 1/2^{|\tilde{P}|} > 0$ . Thus, we get that  $\mathbb{Q}_{(m^{-p}, m^p)}^\nu(h^+) > 0$  and, using Lemma 22 as in the first case, we obtain that  $\ell_a(\mathbb{W}'_a[h^+]) > 0$ , for any  $a \in \mathbb{A}^p$ .

Second, we set

$$h^\mp = (\omega^-, h_{\mathbb{A}^{-p}}^-, h_{\mathbb{A}^p \setminus \{b, c\}}^+, h_b^+, h_c^-), \quad (57a)$$

and we show that  $\mathbb{Q}_{(m^{-p}, m^p)}^\nu(h^\mp) > 0$ .

For this purpose, we first establish that

$$Z_b(h^\mp) = Z_b(\omega^-, h_{\mathbb{A}^{-p}}^\mp, h_{\kappa(1)}^\mp, \dots, h_{\kappa(j_b-1)}^\mp)$$

by partial causality property (33a), and short notation (33b), using Lemma 13 as  $h^\mp \in \mathbb{H}_{\kappa(1), \dots, \kappa(j_b-1)}^\varphi$  since  $h_{\mathbb{A}^{-p}}^\mp = h_{\mathbb{A}^{-p}}^-$  and  $h_{\|\kappa_-\|}^\mp = h_{\|\kappa_-\|}^-$  — by definition (57a) of  $h^\mp$ , using that  $h_{\|\kappa_-\|}^+ = h_{\|\kappa_-\|}^-$  — and as  $h^- \in \mathbb{H}_\kappa^\varphi \subset \mathbb{H}_{\kappa(1), \dots, \kappa(j_b-1)}^\varphi$  by definition (29) of  $\mathbb{H}_\kappa^\varphi$

$$\begin{aligned} &= Z_b(\omega^-, h_{\mathbb{A}^{-p}}^-, h_{\kappa(1)}^+, \dots, h_{\kappa(j_b-1)}^+) && \text{(by definition (57a) of } h^\mp) \\ &= Z_b(\omega^-, h_{\mathbb{A}^{-p}}^-, h_{\kappa(1)}^-, \dots, h_{\kappa(j_b-1)}^-) \\ &\quad \text{(as } h_a^+ = h_a^-, \text{ for all } a \in \|\kappa_-\| \supset \{\kappa(1), \dots, \kappa(j_b-1)\}) \\ &= Z_b(\omega^-, h_{\mathbb{A}^{-p}}^-, h_{\mathbb{A}^p}^-) \end{aligned}$$

again by partial causality property (33a), but with  $h^- \in \mathbb{H}_\kappa^\varphi$  this time, and as  $b = \kappa(j_b)$

$$= Z_b(h^-). \quad \text{(as } h^- = (\omega^-, h_{\mathbb{A}^{-p}}^-, h_{\mathbb{A}^p}^-))$$

Then, we get that

$$w \in \prod_{q \in \tilde{P}} \mathbb{W}_q^- \times \mathbb{W}_p^- \implies T_{(m^{-p}, m^p)}^{\omega^-}(w) = (\omega^-, h_{\mathbb{A}^{-p}}^-, h_{\mathbb{A}^p \setminus \{b, c\}}^+, h_b^+, h_c^-) = h^\mp,$$

because, for any player  $q \in \tilde{P}$  and for any  $w^q \in \mathbb{W}_q^-$  the pure strategy profile  $m^q(w^q, \cdot)$  takes the constant value  $(h_a^-)_{a \in \mathbb{A}^q}$ , and by the expressions (55a)–(55b)–(55c) of  $m^p(w^p, \cdot)$  when  $w^p \in \mathbb{W}_p^+$  using that  $Z_b(h^\mp) = Z_b(h^-) \neq Z_b(h^+)$ . Now, as  $\nu\{\omega^-\} > 0$  and  $\prod_{q \in P} \ell^q(\mathbb{W}_q^-) = 1/2^{|P|} > 0$  we obtain that  $\mathbb{Q}'_{(m^{-p}, m^p)}(h^\mp) > 0$ .

Third, using Lemma 22, we deduce that  $\ell_a(\mathbb{W}'_a[h^\mp]) > 0$  for any  $a \in \mathbb{A}^p$  hence, in particular, that  $\ell_c(\mathbb{W}'_c[h^\mp]) > 0$ . Now, we prove that  $\ell_c(\mathbb{W}'_c[h^-]) > 0$ , where the set  $\mathbb{W}'_c[h^-]$  has been defined in (49), by showing that  $\mathbb{W}'_c[h^\mp] \subset \mathbb{W}'_c[h^-]$ . Indeed, for  $w_c \in \mathbb{W}'_c[h^\mp]$ , we have that

$$\begin{aligned} m_c^p(w_c, h^-) &= m_c^p(w_c, (\omega^-, h_{\mathbb{A}^{-p}}^-, h_{\kappa(1)}^-, \dots, h_{\kappa(j_c-1)}^-)) && \text{(by partial causality)} \\ &= m_c^p(w_c, (\omega^-, h_{\mathbb{A}^{-p}}^-, h_{\kappa(1)}^+, \dots, h_{\kappa(j_c-1)}^+)) \end{aligned}$$

as  $h_a^+ = h_a^-$ , for all  $a \in \|\kappa_-\| \supset \kappa(1), \dots, \kappa(j_c-1)$

$$\begin{aligned} &= m_c^p(w_c, (\omega^-, h_{\mathbb{A}^{-p}}^\mp, h_{\kappa(1)}^\mp, \dots, h_{\kappa(j_c-1)}^\mp)) && \text{(by definition (57a) of } h^\mp) \\ &= m_c^p(w_c, h^\mp) && \text{(by partial causality)} \\ &= h_c^\mp && \text{(by definition of } \mathbb{W}'_c[h^\mp] \text{ in (49))} \\ &= h_c^- . && \text{(by definition (57a) of } h^\mp) \end{aligned}$$

We have shown that  $\mathbb{W}'_c[h^\mp] \subset \mathbb{W}'_c[h^-]$ , hence we deduce that  $\ell_c(\mathbb{W}'_c[h^-]) \geq \ell_c(\mathbb{W}'_c[h^\mp]) > 0$ . Thus, the set  $\mathbb{W}^\pm$  in (54) has positive probability and, for any  $w \in \mathbb{W}^\pm$ , we are going to show that the configuration  $h = T_{(m^{-p}, m^p)}^{\omega^\pm}(w)$  contradicts (56). Indeed, the configuration  $h$  is such that  $h_{\mathbb{A}^{-p}} = h_{\mathbb{A}^{-p}}^+$  because, for any player  $q \in \tilde{P}$ , the pure strategy profile  $m^q(w^q, \cdot)$  takes the constant value  $(h_a^+)_{a \in \mathbb{A}^q}$  when  $w \in \mathbb{W}^\pm$  by definition (54) of  $\mathbb{W}^\pm$ . Then, we get that

$$Z_b(h^+) = Z_b(\omega^+, h_{\mathbb{A}^{-p}}^+, h_{\kappa(1)}^+, \dots, h_{\kappa(j_b-1)}^+)$$

by partial causality property (33a), and short notation (33b), using Lemma 13 as  $h^+ \in \mathbb{H}_\kappa^\varphi$ , and as  $b = \kappa(j_b)$

$$= Z_b(\omega^+, h_{\mathbb{A}^{-p}}, h_{\kappa(1)}, \dots, h_{\kappa(j_b-1)})$$

as we have just established that  $h_{\mathbb{A}^{-p}} = h_{\mathbb{A}^{-p}}^+$ , and as  $(h_{\kappa(1)}, \dots, h_{\kappa(j_b-1)}) = (h_{\kappa(1)}^+, \dots, h_{\kappa(j_b-1)}^+)$  by definition (55a) of  $m_a^p(w^p, h) = h_a^+$  for any  $a \in \mathbb{A}^q \setminus \{b, c\}$

$$= Z_b(h)$$



by partial causality property (33a), and short notation (33b), using Lemma 13 as  $h^+ \in \mathbb{H}_\kappa^\varphi$ ,  $h_{\mathbb{A}^{-p}} = h_{\mathbb{A}^{-p}}^+$ ,  $(h_{\kappa(1)}, \dots, h_{\kappa(j_b-1)}) = (h_{\kappa(1)}^+, \dots, h_{\kappa(j_b-1)}^+)$  and  $b = \kappa(j_b)$ . Now, by definition (54) of  $\mathbb{W}^\pm$ , we have that  $h_c = h_c^-$ . As the set  $\mathbb{W}^\pm$  has positive probability, we conclude that

$$\mathbb{Q}_{(m^{-p}, m^p)}^\nu \{h \in \mathbb{H} \mid h \in \mathbb{H}_\kappa^\varphi, Z_b(h) = Z_b(h^+), h_c = h_c^-\} > 0,$$

but this contradicts (56) since  $\mathbb{Q}_{(m^{-p}, m^p)}^\nu = \mathbb{Q}_{(m^{-p}, m^p)}^\nu$  by assumption.

This ends the proof. □

## 6 Discussion

In this paper, we have introduced the notion of games in product form. For this, we have adapted Witsenhausen’s intrinsic model — where information is represented by  $\sigma$ -fields — to games. In contrast to games in extensive form formulated on a tree, product games do not require an explicit description of the play temporality. Moreover, the product representation is more general than the tree-based ones as, for instance, it allows to describe noncausal situations.

As part of a larger research program, we have focused here on Kuhn’s equivalence theorem. We have also been working on the embedding of tree-based games in extensive form into W-games (by a mapping that associates each information set with an agent), and on the restricted class of W-games that can be embedded in tree-based games [8]. Further research includes

- extensions to infinite number of agents or players, and connections with differential games and mechanism design;
- development of theoretical and numerical decomposition tools (subsystems, subgames, subgame perfect equilibrium, backward induction, approximations, etc.);
- applications (games such as poker, games on graphs, etc.).

### Acknowledgments.

We thank Dietmar Berwanger and Tristan Tomala for fruitful discussions, and for their valuable comments on a first version of this paper. We thank Danil Kadnikov for the pictures of trees and W-models. We thank the editor and a reviewer for their insightful comments. This research benefited from the support of the FMJH Program PGM0 and from the support to this program from EDF.

## References

- [1] C. D. Aliprantis and K. C. Border. *Infinite dimensional analysis*. Springer, Berlin, third edition, 2006. A hitchhiker’s guide.

- [2] C. Alós-Ferrer and K. Ritzberger. *The theory of extensive form games*. Springer Series in Game Theory. Springer-Verlag, Berlin, 2016.
- [3] R. Aumann. Mixed and behavior strategies in infinite extensive games. In M. Dresher, L. S. Shapley, and A. W. Tucker, editors, *Advances in Game Theory*, volume 52, pages 627–650. Princeton University Press, 1964.
- [4] D. P. Bertsekas and S. E. Shreve. *Stochastic Optimal Control: The Discrete-Time Case*. Athena Scientific, Belmont, Massachusetts, 1996.
- [5] L. Blume, A. Brandenburger, and E. Dekel. Lexicographic probabilities and choice under uncertainty. *Econometrica*, 59(1):61–79, 1991.
- [6] P. Carpentier, J.-P. Chancelier, G. Cohen, and M. De Lara. *Stochastic Multi-Stage Optimization. At the Crossroads between Discrete Time Stochastic Control and Stochastic Programming*. Springer-Verlag, Berlin, 2015.
- [7] C. Dellacherie and P. Meyer. *Probabilités et potentiel*. Hermann, Paris, 1975.
- [8] D. Kadnikov. *Game Theory with Information. Games in Intrinsic Witsenhausen Form*. PhD thesis, Université Paris-Est, 2020.
- [9] O. Kallenberg. *Foundations of Modern Probability*. Springer-Verlag, New York, second edition, 2002.
- [10] H. W. Kuhn. Extensive games and the problem of information. In H. W. Kuhn and A. W. Tucker, editors, *Contributions to the Theory of Games*, volume 2, pages 193–216. Princeton University Press, Princeton, 1953.
- [11] M. J. Osborne and A. Rubinstein. *A course in game theory*. MIT press, 1994.
- [12] K. Ritzberger. Recall in extensive form games. *International Journal of Game Theory*, 28(1):69–87, 1999.
- [13] G. Schwarz. Ways of randomizing and the problem of their equivalence. *Israel Journal of Mathematics*, 17:1–10, 1974.
- [14] J. von Neuman and O. Morgenstern. *Theory of games and economic behaviour*. Princeton University Press, Princeton, second edition, 1947.
- [15] H. S. Witsenhausen. On information structures, feedback and causality. *SIAM J. Control*, 9(2):149–160, May 1971.
- [16] H. S. Witsenhausen. The intrinsic model for discrete stochastic control: Some open problems. In A. Bensoussan and J. L. Lions, editors, *Control Theory, Numerical Methods and Computer Systems Modelling*, volume 107 of *Lecture Notes in Economics and Mathematical Systems*, pages 322–335. Springer-Verlag, 1975.