



HAL
open science

Fluctuating temperature regime impairs growth in giant gourami (*Osphronemus goramy*) larvae

Vitas Atmadi Prakoso, Simon Pouil, Wahyulia Cahyanti, Sri Sundari, Otong Zenal Arifin, Jojo Subagja, Anang Hari Kristanto, Jacques Slembrouck

► To cite this version:

Vitas Atmadi Prakoso, Simon Pouil, Wahyulia Cahyanti, Sri Sundari, Otong Zenal Arifin, et al.. Fluctuating temperature regime impairs growth in giant gourami (*Osphronemus goramy*) larvae. *Aquaculture*, 2021, 539, 10.1016/j.aquaculture.2021.736606 . hal-03196727

HAL Id: hal-03196727

<https://hal.science/hal-03196727v1>

Submitted on 5 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Fluctuating temperature regime impairs growth in giant gourami**
2 *(Osphronemus goramy)* larvae
3
4 Vitas Atmadi Prakoso^{a*}, Simon Pouil^{b†}, Wahyulia Cahyanti^a, Sri Sundari^a, Otong
5 Zenal Arifin^a, Jojo Subagja^a, Anang Hari Kristanto^a, Jacques Slembrouck^c
6
7 Affiliations:
8 ^a Research Institute for Freshwater Aquaculture and Fisheries Extension
9 (RIFAFE), Bogor, Indonesia
10 ^b Environmental Sciences Division, Oak Ridge National Laboratory, Oak Ridge,
11 TN, USA
12 ^c ISEM, IRD, Université de Montpellier, CNRS, EPHE, Montpellier, France
13
14 *Corresponding author: vitas.atmadi@gmail.com
15 Research Institute for Freshwater Aquaculture and Fisheries Extension (RIFAFE),
16 Jl. Sempur No. 1, Bogor, West Java, Indonesia 16129
17 Phone: + 62 251 8313200
18
19 [†] Present address: Université Paris-Saclay, INRAE, AgroParisTech, GABI, 78350
20 Jouy-en-Josas, France

21 **Abstract**

22 Growth and physiology were compared in giant gourami (*Osphronemus goramy*)
23 larvae reared in constant (25, 30, and 35°C) and daily fluctuating (25-35°C)
24 temperature regimes. Larvae aged eight days post-hatching were reared indoors
25 for 21 days in four closed recirculating systems (four aquaria per system) in the
26 different temperature regimes. Samples of larvae were collected every week from
27 each temperature regime for growth measurements. At the end of the experiment,
28 cortisol and glucose concentrations, as well as oxygen consumption and
29 ventilation rates, were determined in fish from each temperature regime. Results
30 showed that the fluctuating temperature regime did not affect survival or
31 abnormality, but significantly impaired growth in giant gourami larvae.
32 Physiological responses, assessed using cortisol and glucose levels and oxygen
33 consumption, corroborate previous findings, suggesting that a stable temperature
34 of 30°C is optimal for rearing giant gourami larvae. Our experimental approach
35 demonstrated that temperature variations observed in larval rearing structures on
36 small-scale fish farms are likely to impair the production of giant gourami fry.

37

38 **Keywords:** Growth, Larval rearing, Oxygen consumption, Physiology, Thermal
39 fluctuation

40

41 **1. Introduction**

42 In aquatic environments, the thermal tolerance range of ectothermic organisms is
43 largely determined by the amplitude of thermal variation in their natural habitat.
44 Accordingly, the range of thermal tolerance is generally wider in temperate
45 species than in those that inhabit colder, warmer, and more stable water
46 temperatures (Pörtner, 2002; Souchon and Tissot, 2012). Fish growth usually
47 increases with increasing temperature up to an optimal temperature, and then
48 decreases at temperatures higher than this optimum (Árnason et al., 2019).
49 Temperature changes that exceed the thermal limit for long durations may
50 compromise fish survival (Whitney et al., 2016; Colchen et al., 2017; Mueller et
51 al., 2017; Réalis-Doyelle et al., 2018). Thus, knowledge on thermal plasticity is
52 important to understand how thermal regimes affect fish aquaculture production
53 (Mommsen et al., 1999; Whitney et al., 2016).

54 The effects of constant and fluctuating thermal regimes on fish have been assessed
55 in previous studies. Several studies performed on temperate species have revealed
56 that exposure to fluctuating temperatures improves fish tolerance to high
57 temperatures (e.g. Carveth et al., 2007; Eldridge et al., 2015; Pisano et al., 2019).
58 In contrast, fluctuating temperatures confer no benefits compared with a constant
59 temperature regime in Lahontan cutthroat trout *Oncorhynchus clarki henshawi*
60 (Dickerson and Vinyard, 1999) or smallmouth bass *Micropterus dolomieu*
61 (Eldridge et al., 2015).

62 The effects of temperature fluctuations on fish growth may be diverse. Although
63 growth can be boosted by daily fluctuating temperatures (Konstantinov et al.,
64 1990; Carveth et al., 2007; Azaza et al., 2010), other studies, mostly involving

65 temperate species, have shown slower growth than at constant thermal regimes
66 (Meeuwig et al., 2004, Hevrøy et al., 2012; Eldridge et al., 2015; Tromp et al.,
67 2016). Still other species show no significant differences, such as blue tilapia
68 *Oreochromis aureus* (Baras et al., 2000) and Japanese medaka *Oryzias latipes*
69 (Dhillon and Fox, 2007). These various studies highlight that the effects of
70 temperature fluctuations are species-dependent and that there are few data on
71 tropical species.

72 Fish metabolism and physiology are influenced by fluctuating temperatures,
73 affecting in turn growth, health, and survival (Pérez-Casanova et al., 2008a,
74 2008b; Singh et al., 2013; Tromp et al., 2016; Fu et al., 2018). In general,
75 individuals respond directly to acute exposure to stressors by increasing cortisol
76 and glucose levels to cope with stress (Iwama et al., 2005; Tromp et al., 2018).
77 Oxygen consumption in fish also depends on temperatures and increases with
78 increasing temperatures (Claësson et al., 2016; Remen et al., 2015). Such changes
79 in physiology may affect growth and survival.

80 There are still gaps in knowledge on several aspects of the biology of some
81 aquaculture tropical species. Among them, giant gourami (*Osphronemus goramy*
82 Lacepède, 1801) is the main species being farmed in Indonesia and is in great
83 demand for the food aquaculture industry (Amornsakun et al., 2014; Budi and
84 Suprayudi, 2015). Currently, with an estimated production of more than 173,000 t
85 in 2018, Indonesia is the world's highest giant gourami producer (>98% of world
86 production; FAO, 2020). Giant gourami aquaculture activities in Indonesia are
87 mainly driven by micro and small-scale farms, which are concentrated on Java
88 Island (Maskur et al., 2013). Although the giant gourami is an interesting species

89 in regard to current Indonesian aquaculture guidance, its production has slowed
90 for various reasons. One of the main impediments in giant gourami aquaculture is
91 ensuring a regular supply of fry (Slembrouck et al., 2019). A recent survey on
92 small-scale giant gourami farms in West Java identified irregular temperatures as
93 the main limiting factor for egg and fry production (Kristanto et al., 2020).
94 Current practices for raising giant gourami larvae still involve their rearing in
95 plastic basins and being moved to small outdoor ponds (Kristanto et al., 2020).
96 However, the Indonesian National Standard recommends indoor larval rearing to
97 avoid exposure to the large variations encountered in environmental conditions
98 (SNI, 2000). Temperatures fluctuate widely in outdoor rearing systems with
99 observed variations ranging from 27 to 40 °C, and massive mortality events are
100 frequently reported by farmers (Kristanto et al., 2020).
101 In this context, the objectives of this study were (1) to assess the growth and
102 survival of giant gourami (*O. goramy*) larvae reared in closed recirculating
103 systems at four rearing temperature regimes (fluctuating: 25-35 °C and constant:
104 25, 30, and 35 °C) and (2) to determine the effects of temperature fluctuation on
105 larval physiology.

106

107 **2. Materials and methods**

108 2.1. Source of larvae

109 Giant gourami *O. goramy* larvae used in this experiment came from natural
110 spawns of the same giant gourami broodfish pair. The broodfish, belonging to the
111 “Galunggung” strain, were approximately 4-5 years old and spawned at the
112 Tasikmalaya Center for the Development of Giant Gourami Culture (BPPSIGN)

113 where temperatures in broodfish ponds usually range from 25 to 32 °C (Kristanto
114 et al., 2020). After spawning, eggs were collected from the nest and then
115 transported to the Germplasm Freshwater Aquaculture Research Station
116 (RIFAFE, Cijeruk, Bogor, West Java, Indonesia). Eggs were then placed in an
117 incubation basin (20 L) inside an experiment room with a daily water change, a
118 temperature of 29.5 ± 0.7 °C, and light:dark cycle of 12:12 h, until the eggs
119 hatched. After hatching, larvae were kept unfed in the incubation basin until they
120 reached 8 days post-hatching (dph) for the experiment.

121

122 2.2. Live prey

123 Giant gourami larvae were fed with live tubifex worms (*Tubifex tubifex*) during
124 the experiment based on current practices (SNI, 2000) for optimal survival and
125 growth. Live tubifex worms, purchased regularly, were stored in a 100-L
126 aquarium in the experiment room with a daily water change, temperature of $29.5 \pm$
127 0.7 °C, and a light:dark cycle of 12:12 h; they were kept unfed.

128

129 2.3. Temperature regime experiment

130 2.3.1. Experimental design

131 Zootechnical and physiological performances of giant gourami larvae were
132 evaluated in four temperature treatments: one fluctuating temperature regime (25-
133 35 °C) and three constant temperature regimes (25, 30, and 35 °C). The
134 experiment was carried out in four identical indoor recirculating systems (RAS;
135 one system for each temperature treatment) under a light:dark cycle of 12:12 h
136 (daylight intensity: 60-4500 lux) including a sump with mechanical and biological

137 filtration and a buffer tank. PVC pipes connected each RAS to the experimental
138 tanks (i.e. 30 L glass aquaria $40 \times 30 \times 30$ cm) placed on a bench. At 8 dph, larvae
139 (mean body mass: 12.1 ± 1.2 mg; mean total length: 9.4 ± 0.3 mm) were randomly
140 distributed into the experimental tanks ($n = 4$ tanks per temperature treatment;
141 stocking density of 72 larvae per tanks or 2.4 larvae L^{-1}) in each RAS. During the
142 experiment, water inlet flow into rearing tanks was maintained at $33 L h^{-1}$ for the
143 first 5 days, and then increased to $78 L h^{-1}$ until the end of the experiment.
144 Prior to the experiment, giant gourami larvae in each treatment were kept in their
145 respective experimental tanks at an initial temperature of 28 °C. Then, larvae were
146 gradually acclimated to target temperatures (± 1 °C h^{-1}). Temperatures were kept
147 stable at target temperatures, except for the fluctuating temperature treatment
148 (Figure 1). Submersible heaters and chillers (when appropriate) regulated by
149 thermostats and placed in the sump of each RAS were used to control temperature
150 during the experiment. Six weeks prior to the experiment, preliminary tests were
151 performed to evaluate the temperature in the RASs. Temperature was monitored
152 regularly, especially in the fluctuating regime during the ascent and descent
153 phases to ensure that the variations were gradual. The experiment started several
154 days after all the temperature treatments were running properly and temperature
155 was then checked as described Section 2.3.2. To prevent heat loss and temperature
156 variations, the outside of the experimental tanks was insulated with plastic and
157 extruded polystyrene foam and covered with a transparent polycarbonate sheet.
158 Each experimental tank was equipped with an aeration system. The experiment
159 was carried out for 21 days, the typical rearing time to reach market size.
160

161 2.3.2. Feeding protocol and water quality monitoring

162 Giant gourami larvae were fed live tubifex worms every day, except on the
163 sampling days. To ensure continuous and non-limiting feeding, live tubifex
164 worms were placed on the bottom of each aquarium twice a day (at 8:00 and
165 17:00), the quantities were given in excess and in an equivalent amount for each
166 treatment during the whole experiment. Feeding in excess allows accurate
167 estimation of feed intake by weighing the uneaten tubifex worms remaining on the
168 bottom of each aquarium (Prakoso et al., 2019). Prior to their distribution, live
169 tubifex worms were collected, washed, and drained on a 50- μ m mesh and weighed
170 (to the nearest 0.1 g). Uneaten tubifex worms were removed from each aquarium
171 before giving the new ration and weighed to calculate feed intake. The
172 temperature in each experimental tank was observed three times a day (at 08:00,
173 12:00, and 16:00). Meanwhile, water quality was directly measured for pH,
174 dissolved oxygen (DO), and conductivity once a week using a portable
175 multiparameter meter (HI 9829 Hanna Instruments). Additionally, total ammonia
176 nitrogen (TAN), NH_3 , NH_4^+ , and NO_2^- were measured using spectrophotometry (n
177 = 12 per treatment). The results of water quality measurements are given in Table
178 1.

179

180 2.4. Observations and measurements on larvae

181 Larvae (n = 22 for each aquarium, i.e. ~30% of the total number of larvae) were
182 sampled every seven days. Prior to measurement, larvae were transferred to a
183 plastic basin containing water from their respective aquaria to keep them in the
184 same water temperature conditions. Larvae were anesthetized by using eugenol

185 and measured (total body length, TL, mm) under a stereomicroscope with a
186 micrometer (accuracy ranging from 0.05 to 0.1 mm, depending on fish size and
187 magnification). Meanwhile, body mass (BM, mg) was measured using a digital
188 weighing scale (accuracy of 0.1 mg). After measurement, fish were returned to
189 their respective experimental tanks.

190 At the end of the experiment, the living larvae in each aquarium were counted to
191 determine the survival rate and were visually observed for anatomic abnormalities
192 (i.e. morphological deformities). Additionally, cortisol and glucose levels were
193 recorded from each treatment following Prakoso et al. (2019). Whole fish were
194 collected, anesthetized, and ground (n = 4 pools of 10 fish). Subsequently, the
195 samples were vortex-mixed (30 s) and centrifuged (10,000 rpm for 10 min at 25
196 °C) to collect the supernatant before being frozen. Cortisol levels were determined
197 using the Cortisol ELISA kit (DRG International Inc., Germany) and a microplate
198 photometer (Biosan HiPo MPP-96, Latvia) according to the manufacturer's
199 instructions. Inter-plate and intra-plate coefficients of variation were always less
200 than 5%. Meanwhile, blood glucose samples were collected as described below (n
201 = 4 pools of 10 fish). One drop of blood was collected from each anesthetized fish
202 from each treatment (n = 10 fish) and was read on a blood glucose meter (Accu-
203 Chek Active, Roche, Germany).

204

205 2.5. Oxygen consumption, ventilation rate, and Q_{10} measurements

206 At the end of the rearing period, oxygen consumption was measured in giant
207 gourami fry using a closed respirometer (1.4 L) connected to a dissolved oxygen
208 meter (Trans Instrument HD3030, Singapore). After a 10 min acclimation period,

209 measurements were read for the three constant temperature treatments (25, 30, and
210 35 °C). The fluctuating temperature treatment (25-35 °C) was not measured due to
211 technical constraints. Twenty-one fish were combined for each measurement with
212 three replicates for each constant temperature treatment. Dissolved oxygen
213 concentrations were recorded every five minutes. Ventilation rates were also
214 determined by counting opercular movements for 1 min with nine replications for
215 each treatment. Furthermore, the measured oxygen consumption data were used to
216 calculate Q_{10} (e.g. Hölker, 2006; see Section 2.6 for details).

217

218 2.6. Physiological and zootechnical parameters

219 Temperature effects on survival, growth, and feed intake were determined by
220 calculating the following parameters for each experimental treatment.

221 Survival rates (SR), expressed as a percentage, were calculated by comparing the
222 final number (N_f) with the initial number of larvae (N_i): $SR (\%) = (N_f/N_i) \times 100$.

223 Abnormal fish rates in % were calculated by comparing the number of deformed
224 fish (N_{ab}) with the final number of live fish (N_f): $(N_{ab}/N_f) \times 100$.

225 Specific growth in body mass (SGR_{BM} , %) was calculated according to the
226 following equation: $SGR_{BM} = (\ln BM_f - \ln BM_i)/ED \times 100$, where BM_i and BM_f
227 are the initial and final body mass of larvae (mg), respectively, and ED is the
228 duration of the experiment in days.

229 Specific growth in total length (SGR_{TL} , %) was calculated using the same
230 calculation by substituting BM for TL: $SGR_{TL} = (\ln TL_f - \ln TL_i)/ED \times 100$.

231 Heterogeneity of larval size (in body mass or total length) was assessed using the
232 coefficient of variation (CV, %) calculated as $CV_{BM} = SD_{BM}/BM$ and $CV_{TL} =$

233 SD_{TL}/TL , where SD is the standard deviation and BM is the average body mass
234 (mg) and TL, the average body length (mm).

235 Fulton's condition factor (K) was calculated according to the relationship
236 $K = BM_f/TL_f^3$. The equation was multiplied by 100 to bring the value close to
237 one.

238 Fish biomass gain (BG, g) was calculated following the equation: $BG = (N_f BM_f -$
239 $N_i BM_i)/1000$, where BM_i and BM_f are the initial and the final average body mass
240 (mg), respectively.

241 The feed conversion ratio (FCR) was calculated using the following equation:
242 $FCR = F/(N_f BM_f - N_i BM_i)$, where F is the total quantity of food intake in wet
243 weight during the whole rearing period. F was determined as the total amount of
244 uneaten food subtracted from the total amount of food provided (g).

245 Oxygen consumption was calculated by the following equation: $OC = (DO_0 -$
246 $DO_t) \times (V/(t \times W))$, where OC is the oxygen consumption expressed as $mg\ kg^{-1}\ h^{-1}$
247 ¹, DO_0 and DO_t are the dissolved oxygen concentrations at initial time and after t
248 hours ($mg\ L^{-1}$), t is the time elapsed for fish consuming the oxygen (h), V is water
249 volume on respirometer (L), and W is the total weight of fish (kg).

250 Q_{10} values were calculated using the following equation:

$$Q_{10} = \left(\frac{R_2}{R_1}\right)^{\left(\frac{10}{t_2-t_1}\right)}$$

251 where R_1 is the oxygen consumption rate at temperature t_1 , and R_2 , the oxygen
252 consumption rate at temperature t_2 .

253

254 2.7. Data analysis

255 Statistical comparisons were done for each parameter described in Section 2.6.
256 Data were first assessed to confirm normality (Shapiro's test) and homogeneity of
257 variances (Levene's test) and, where necessary, data were arcsine-square-root-
258 transformed prior to analysis. One-way ANOVA followed by Tukey's test was
259 used to determine significant differences among temperature treatments. When
260 assumptions of normality and homogeneity of variances were not met, data were
261 analyzed using the Kruskal-Wallis and Dunn's non-parametric tests.

262 The growth kinetics of the larvae, both BM and total length TL, reared at the five
263 different temperatures were best fitted using an exponential model: BM or $TL =$
264 BM_i or $TF_i \times e^{at}$ where a is the growth rate (d^{-1}), and t , time (d). Model constants
265 were estimated by iterative adjustment using the least squares regression method.
266 Differences in growth (BM, TL) and body condition (K) over time were assessed
267 using a repeated-measures ANOVA designed as a linear-mixed-effect model.
268 "Temperature" and "Time" were used as fixed factors and "Tank" was used as a
269 random factor to take into account a possible "tank effects" (Sswat et al., 2018).
270 The models were fitted using means of body mass and total length per tank and
271 sampling time (Sswat et al., 2018). The level of significance for statistical
272 analyses was always set to $\alpha = 0.05$. All statistics were performed using R
273 freeware version 3.6.3 (R Development Core Team, 2020).

274

275 **3. Results**

276 3.1. Water quality parameters

277 The targeted temperature regimes were met throughout the experiment (Table 1)
278 and significant differences were confirmed between each treatment ($F = 381.93$, p

279 < 0.001, Table 1). Regular monitoring indicated that pH was similar between
280 treatments, but some differences were observed for conductivity ($\chi^2 = 57.7$, $p <$
281 0.001, Table 1). Dissolved oxygen concentration remained above 6.50 mg L⁻¹ in
282 all treatments. TAN was not significantly different between the four experimental
283 treatments, and the NO₂⁻ and NH₃ concentrations were lower than 0.1 and 0.05 mg
284 L⁻¹, respectively.

285

286 3.2. Survival and abnormality

287 The larval survival rate measured at the end of the experiment ranged from 56.6 ±
288 9.2% (35 °C treatment) to 96.2 ± 1.3% (30 °C treatment), with significant
289 differences in survival between treatments ($F = 37.34$, $p < 0.001$) except between
290 the fluctuating temperature regime (i.e. 25-35 °C treatment) and the two lowest
291 constant temperatures (i.e. 25 and 30 °C; Fig. 2A). The proportion of
292 abnormalities, mainly malformation of the spinal column, in living larvae varied
293 from 0 (30 °C treatment) to 11.8 ± 9.7% (35 °C treatment), but was not
294 significantly different between experimental treatments (Fig. 2B).

295

296 3.3. Growth and size heterogeneity

297 Larval growth (BM and TL) increased over time and varied significantly between
298 the temperature treatments ($F = 64.58-72.46$, $p < 0.001$) with a significant
299 interaction between temperature and time ($F = 70.10-146.849$, $p < 0.001$),
300 indicating an accentuation of the differences as the experiment progressed (Fig. 3).

301 At the end of the 21-day experiment, average BM and TL of larvae ranged from
302 147.1 ± 30.6 mg and 21.2 ± 1.6 mm and 276.7 ± 47.5 mg and 26.1 ± 1.4 mm

303 when they were reared at 25 and 30 °C, respectively (Table 2). Growth
304 significantly increased with increasing temperature up to 30 °C, with the lowest
305 growth observed for larvae reared at 25 °C (Fig. 3 and Table 2), and a slight, but
306 non-significant, decrease in growth was observed at the highest tested temperature
307 (35 °C, Fig. 3 and Table 2). Intermediate growth was observed in fish reared in
308 the fluctuating temperature regime (25-35 °C). This trend was confirmed by the
309 specific growth rate: significant increases in SGR_{BM} ($F = 37.10$, $p < 0.001$) and
310 SRG_{TL} ($F = 45.70$, $p < 0.001$) were observed with increasing temperatures up to
311 30 °C, with values varying from $12.1 \pm 0.6\%$ for SGR_{BM} and $3.9 \pm 0.2\%$ for
312 SRG_{TL} at 25 °C to $15.1 \pm 0.2\%$ for SGR_{BM} and $4.9 \pm 0.1\%$ SRG_{TL} at 30 °C (Table
313 2). No significant differences were observed in SGR between 30 and 35 °C for
314 either length or weight (Table 2).

315 Size heterogeneity in larvae as a function of temperature was assessed based on
316 calculation of the coefficients of variation for body mass (CV_{BM}) and total length
317 (CV_{TL}) at the end of the experiment. CV_{BM} ranged from $17.1 \pm 2.7\%$ (30 °C
318 treatment) to $26.4 \pm 7.9\%$ (35 °C treatment) and CV_{TL} ranged from $5.5 \pm 0.8\%$
319 (30 °C treatment) to $9.3 \pm 3.2\%$ (35 °C treatment). Although no significant change
320 was observed for CV_{TL} , a slight change was observed for CV_{BM} ($F = 3.59$, $p =$
321 0.046 , Table 2) with significant differences between the 30 °C and 35°C
322 treatments. There was a significant change in K for the different temperature
323 regimes ($\chi^2 = 195$, $p < 0.001$), with final K ranging from 1.48 ± 0.18 and $1.67 \pm$
324 0.28 for larvae reared at the fluctuant temperature regime (25-35 °C) and at the
325 highest constant temperature (i.e. 35 °C, Table 2), respectively.

326

327 3.4. Feed efficiency

328 FCR was affected by the temperature regime ($F = 10.73$, $p = 0.001$), with the
329 lowest FCR observed at 30 °C (4.75 ± 0.92) and the highest FCR observed at 35
330 °C (10.58 ± 2.27 , Table 2). There were no significant differences between the
331 fluctuating temperature regime (25-35 °C) and the lowest constant temperature
332 treatments (25 and 30 °C).

333

334 3.5. Cortisol and blood glucose concentrations

335 Cortisol and blood glucose concentrations were measured in giant gourami at the
336 end of the 21-day experiment for each temperature treatment. Cortisol
337 concentrations were not significantly different among treatments except between
338 30 °C and 35 °C ($F = 4.98$, $p = 0.031$), with values ranging from 6.22 ± 6.21 ng
339 mL^{-1} at 30 °C to 28.46 ± 2.72 ng mL^{-1} at 35 °C (Fig. 4). Intermediate cortisol
340 values were observed for the fluctuating temperature regime (25-35 °C) and the
341 lowest constant temperature regime (25 °C) with 13.81 ± 5.64 ng mL^{-1} and 12.56
342 ± 11.63 ng mL^{-1} , respectively (Fig. 4). At the end of the experiment, no
343 differences were observed in glucose concentrations in giant gourami reared at the
344 different temperature regimes, with values ranging from 37.83 ± 10.07 mg dL^{-1} to
345 54.17 ± 9.87 mg dL^{-1} at 25 °C and 35 °C, respectively (Fig. 4).

346

347 3.6. Oxygen consumption, ventilation rates, and Q_{10}

348 Oxygen consumption and ventilation rates were not assessed for the fluctuating
349 temperature regime (25-35 °C) due to technical constraints. Among the constant
350 temperature treatments, oxygen consumption increased with temperature ($F =$

351 62.20, $p < 0.001$), with values ranging from $422 \pm 70 \text{ mg kg}^{-1} \text{ h}^{-1}$ at 25 °C to $780 \pm$
352 $90 \text{ mg kg}^{-1} \text{ h}^{-1}$ at 35 °C (Fig. 5). Overall, the same trend was observed for
353 ventilation rates ($\chi^2 = 22.8$, $p < 0.001$), with values ranging from 163 ± 20 times
354 min^{-1} at 25 °C to 215 ± 8 times min^{-1} at 35 °C (Fig. 5), although there were no
355 significant differences between the two highest constant temperature regimes (i.e.
356 30 and 35 °C). The temperature coefficient (Q_{10}) of giant gourami larvae between
357 25 and 30 °C (the highest) was 1.98, between 25 and 35 °C, it was 1.92, and
358 between 30 and 35 °C, it was 1.91.

359

360 **4. Discussion**

361 4.1. Survival and abnormalities at the different temperature regimes

362 In the present study, daily temperature fluctuations (25-35 °C) did not
363 significantly affect the mortality of giant gourami larvae compared with the
364 constant temperature regimes at 25 and 30 °C. However, mortality increased
365 drastically at 35 °C. Overall, our results corroborate those of previous studies on
366 the same species (Sarah et al., 2009, Verawati et al. 2015, Arifin et al. 2019).
367 Although higher TAN values, explained by NH_4^+ (approx. 1 mg L^{-1}), were
368 observed at 25 °C than for the other temperature treatments, the NH_3 values
369 remained within a non-toxic range for tropical fish (i.e. $< 0.05 \text{ mg L}^{-1}$; Boyd and
370 Tucker, 2012). Thus, we assumed that TAN did not explain the mortality
371 observed at 25 °C. The increase of mortality for the low temperature treatment
372 was also observed by Prakoso et al. (2019). In the present study, there were no
373 significant differences in the fish anatomic abnormality rate between the different
374 temperature regimes or between fluctuating and constant temperatures.

375 Nevertheless, at the highest temperature treatment (35 °C), heterogeneity in the
376 abnormality rate per aquarium was high, with abnormalities ranging from 0 to
377 22%. Dead larvae were not checked for abnormalities, but our observations
378 suggest that taking them into account could have increased the actual abnormality
379 rate for the 35 °C treatment. Our findings showed once again that a constant 30 °C
380 regime is optimal for survival in giant gourami larval rearing, as previously
381 suggested (Prakoso et al. 2019); this temperature minimizes the fish abnormality
382 rate. The high mortality found in this study (i.e. 30-51% at 35 °C) clearly shows
383 that 35 °C is higher than the tolerance limit of giant gourami larvae. Our
384 experimental approach demonstrated that daily thermal fluctuations do not impair
385 survival.

386

387 4.2. Effects of fluctuating temperatures on growth and feed conversion

388 The fluctuating temperature regime had a negative effect on giant gourami growth.
389 Our results also showed that fluctuating temperatures tend to increase
390 heterogeneity in size slightly. Interestingly, the final average body mass increased
391 with average temperatures during the experiment: 147.1 mg at constant 25 °C
392 regime (24.8 ± 0.1 °C), 178.0 mg at fluctuating regime (29.0 ± 3.4 °C), 276.7 mg
393 at constant 30 °C regime (29.9 ± 0.1 °C), and 262.9 mg at constant 35 °C regime
394 (34.8 ± 0.1 °C). In terms of growth, the difference between fluctuating
395 temperatures and constant 30 °C was higher than between the fluctuating
396 temperature regime and the lowest constant temperature (25°C), suggesting that
397 minimizing fluctuations in temperature is a key factor in maximizing larval
398 growth. Surprisingly, the effects of a fluctuating temperature regime have been

399 poorly investigated in the literature. Overall, fluctuating temperatures negatively
400 affect fish growth. For example, at comparable developmental stages, spikedace
401 (*Meda fulgida*) held at fluctuations of 24–34 °C and 30–34 °C show significantly
402 lower growth than the constant temperature control (25 °C) (Carveth et al., 2007).
403 Furthermore, although small juveniles of the Nile tilapia *Oreochromis niloticus*
404 grow slightly faster when raised under fluctuating temperatures than under
405 constant temperatures, thermal fluctuations become increasingly unsuitable in fish
406 of increasing size (Azaza et al., 2010). A similar conclusion was drawn in a study
407 on blue tilapia (Baras et al., 2000). In the present study, at constant temperatures,
408 there was an overall trend of increased giant gourami larval growth with
409 temperatures of up to 30 °C. From 30 to 35 °C, this trend was reversed. At
410 comparable temperatures and stocking densities, growth performances were
411 similar to those reported in recent studies on giant gourami larval rearing (Arifin
412 et al., 2019; Sarah et al., 2009).

413 In regard to growth, the fluctuating temperature treatment showed significantly
414 higher FCR values compared with the constant 30 °C treatment, but remained
415 lower than the two extreme temperatures (i.e. 25 and 35 °C). These results are the
416 consequence of the combined effects of temperature in feed consumption, food
417 passage time, and enzyme activity in fish (Michalsen et al., 1998), and corroborate
418 previous findings on giant gourami (Prakoso et al., 2019) and other tropical fish
419 species (e.g. Degani et al., 1989; Watanabe et al., 1993). FCRs are generally high
420 in giant gourami (Arifin et al., 2019; Prakoso et al., 2019), and our findings
421 emphasize that controlling — and especially stabilizing — temperature during

422 larval rearing is the key to improving feed efficiency and minimizing production
423 costs related to diet (Kristanto et al., 2020).

424

425 4.3. Physiological responses to temperature fluctuation

426 Temperature changes are expected to stress aquatic organisms due to their
427 influence on all biological processes (Iwama et al., 2005; Pörtner et al., 2006;
428 Killen et al., 2013; Verhille et al., 2016; Tromp et al., 2018), which can be
429 predicted by measuring glucose and whole-body cortisol as the main indicators of
430 stress in fish larvae (Rudneva, 2013). Overall, there were no significant
431 differences between fluctuating and constant temperatures regarding cortisol and
432 glucose concentrations, except for cortisol between the constant temperature
433 treatments of 30 and 35 °C. Most previous studies on tropical fish species report
434 effects of temperatures on cortisol and glucose levels when temperature deviates
435 from the tolerance limits specific to each species (Chen et al., 2002; Zaragoza et
436 al., 2008; Long et al., 2012). Prakoso et al. (2019) found changes in cortisol levels
437 over time (but not detected at the beginning of the experiment) in giant gourami
438 10-dph larvae exposed to different temperatures for several weeks. The same
439 authors reported limited effects of temperature on glucose concentrations (initial
440 levels ranging from 30 to 40 mg dL⁻¹) measured at the end of the experiment,
441 except a decrease in glucose levels at the extreme temperatures tested (22.5 and
442 32.5 °C), but cortisol levels were highest (i.e. up to 78 ng mL⁻¹) at 25 and 27.5 °C.
443 Our findings corroborate those of Ebrahimi et al. (2011), who did not show
444 differences in glucose levels among giant gourami juveniles reared at
445 temperatures of 28, 30 and 32 °C. In this study, cortisol and glucose levels were

446 measured only at the end of the experiment; therefore, we cannot highlight any
447 changes during development. However, on larvae of the same age exposed for
448 three weeks to different temperatures, we found that cortisol levels were the
449 highest at 35 °C, indicating the occurrence of stress for larvae reared at this
450 temperature regime as shown by Hanke et al. (2019) in juvenile milkfish *Chanos*
451 *chanos* exposed to a gradual temperature increase of 1 °C per day in the range
452 from 26 °C to 33 °C. This study on milkfish warns that plasma cortisol values may
453 be biased by the fish-handling procedures. As described above, in the present
454 study, whole fish were collected, anesthetized, and ground to analyze cortisol
455 levels. Because all the fish were treated in the same way, the cortisol
456 concentrations we measured are comparable between treatments. Based on these
457 findings, we can reasonably assume that the highest constant temperature tested in
458 this study negatively affected the physiological response of giant gourami.
459 Interestingly, the fluctuating temperature regime tested in this study (25-35 °C)
460 was less harmful for giant gourami larvae compared with the constant 35 °C
461 rearing temperature. Nevertheless, as in Prakoso et al. (2019), we used larvae
462 from the same pair of brooders from a production site located at 480 m above the
463 sea level where the range of water temperature is restricted. Thus, further
464 investigations based on other gourami strains are needed to confirm our results on
465 a broader scale.

466

467 4.4. Changes in oxygen consumption with temperature

468 Fish metabolism is influenced by water temperature, which is also related to
469 health, aerobic swimming performance, stress response, growth, and survival

470 (Pérez-Casanova et al., 2008a, 2008b; McKenzie and Claireaux, 2010; Singh et al.,
471 2013; Tromp et al., 2016; Fu et al., 2018). Giant gourami larvae responded to the
472 increase in temperatures (i.e. 25, 30, and 35 °C) by increasing oxygen
473 consumption and ventilation rates. The oxygen consumption pattern we observed
474 is similar to previous reports on other freshwater tropical fish such as climbing
475 perch *Anabas testudineus* (117, 126, and 198 mg O₂ kg⁻¹ h⁻¹ at 25, 30, and 35 °C,
476 respectively; Sarma et al., 2010), common carp *Cyprinus carpio* (64, 71, and 99
477 mg O₂ kg⁻¹ h⁻¹ at 25, 30, and 35 °C, respectively; Chatterjee et al., 2004),
478 yellowtail catfish *Pangasius pangasius* (from 111, 232, 236 mg O₂ kg⁻¹ h⁻¹ at 30,
479 34, and 38 °C, respectively; Debnath et al., 2006), and Nile tilapia (from 22 to 95
480 mg O₂ kg⁻¹ h⁻¹ at 19 and 31 °C, respectively; Turker, 2011). Moreover, the pattern
481 of ventilation rates in giant gourami fry we measured is consistent with previous
482 observations on other fish species such as the barred loach *Nemacheilus fasciatus*
483 (115 to 265 times min⁻¹ at 21.5 and 31.5 °C, respectively; Prakoso and Kurniawan,
484 2020), the pikeperch *Sander lucioperca* (from 8 to 31 times min⁻¹ at 13 and 28 °C,
485 respectively; Frisk et al., 2012), and the exquisite sand-goby *Favonigobius*
486 *exquisitus* (from 31 to 160 times min⁻¹ at 15 and 35 °C, respectively; Ford et al.,
487 2004). Each fish species has a specific range of temperature preferences, resulting
488 in various oxygen consumption patterns for specific metabolic processes (Das et
489 al., 2005; Majhi and Das, 2013). Here, we found that higher temperatures lead to
490 higher metabolic rates for giant gourami larvae reared at temperatures of up to 35
491 °C.

492 The lowest temperature coefficient (Q₁₀) in giant gourami larvae was estimated
493 between 30 and 35 °C (1.91). This result indicates that within this temperature

494 range, this species maintains homeostasis more efficiently (González et al. 2010).
495 According to previous studies (e.g. Das et al., 2018; Prakoso and Kurniawan,
496 2020), the value at which Q_{10} drops among temperature treatments indicates the
497 ideal thermal range for growth. Such results corroborate a previous study showing
498 that the optimal larval rearing temperature for giant gourami is 30 °C (Prakoso et
499 al., 2019). All larvae we used have the same genetic background; similar studies
500 on other strains/populations/geographic locations are needed to generalize our
501 findings.

502

503 **5. Conclusion**

504 The present study demonstrated that a fluctuating temperature regime impairs
505 growth in giant gourami larvae. Accordingly, we observed a >60% reduction in
506 growth between larvae reared in optimal conditions (i.e. stable temperature, at
507 30 °C) and larvae that undergo daily fluctuations in temperature (i.e. 25-35 °C).
508 Additional, albeit less significant, impacts were observed for survival and
509 physiological responses. Our experimental approach confirmed that daily
510 temperature variations observed in larval rearing structures in small-scale fish
511 farms (Kristanto et al. 2020) are likely to be an impediment in the production of
512 giant gourami fry.

513

514 **Acknowledgements**

515 The authors acknowledge Imam Akhir Abdullah, technician at the Cijeruk
516 research station, and Ahmad Sihabuddin, technician at IRD Indonesia, for their

517 technical assistance throughout the experiment. This is publication ISEM XXXX-
518 XXX.

519

520 **References**

521 Amornsakun, T., Kullai, S., Hassan, A., 2014. Some aspects in early life stage of
522 giant gourami, *Osphronemus goramy* (Lacepède) larvae. Songklanakarin J.
523 Sci. Tech. 36, 493-498.

524 Arifin, O.Z., Prakoso, V.A., Kristanto, A.H., Pouil, S., Slembrouck, J., 2019.
525 Effect of stocking density on growth, food intake and survival of giant
526 gourami (*Osphronemus goramy*, Lacepède) larvae. Aquaculture 509, 159-
527 166.

528 Árnason, T., Gunnarsson, Á., Steinarsson, A., Daníelsdóttir, A.K., Björnsson,
529 B.T., 2019. Impact of temperature and growth hormone on growth
530 physiology of juvenile Atlantic wolffish (*Anarhichas*
531 *lupus*). Aquaculture 504, 404-413.

532 Azaza, M.S., Legendre, M., Kraiem, M.M., Baras, E., 2010. Size- dependent
533 effects of daily thermal fluctuations on the growth and size heterogeneity
534 of Nile tilapia *Oreochromis niloticus*. J. Fish Biol. 76(3), 669-683.

535 Baras, E., Prignon, C., Gohoungo, G., Méalard, C., 2000. Phenotypic sex
536 differentiation of blue tilapia under constant and fluctuating thermal
537 regimes and its adaptive and evolutionary implications. J. Fish Biol. 57(1),
538 210-223.

- 539 Boyd, C.E., Tucker, C.S., 2012. Pond aquaculture water quality management.
540 Springer Science + Business Media, NY, USA.
- 541 Budi, D.S., Suprayudi, M.A., 2015. Growth response and feed utilization of giant
542 gourami (*Osphronemus goramy*) juvenile feeding different protein levels
543 of the diets supplemented with recombinant growth hormone. Hayati 22(1),
544 12-19.
- 545 Carveth, C.J., Widmer, A.M., Bonaar, S.A., Simms, J.R. 2007. An examination of
546 the effects of chronic static and fluctuating temperature on the growth and
547 survival of spinedace, *Meda fulgida*, with implications for management. J.
548 Therm. Biol. 32, 102-108.
- 549 Chatterjee, N., Pal, A.K., Manush, S.M., Das, T., Mukherjee, S.C., 2004. Thermal
550 tolerance and oxygen consumption of *Labeo rohita* and *Cyprinus carpio*
551 early fingerlings acclimated to three different temperatures. J. Therm.
552 Biol. 29(6), 265-270.
- 553 Chen, W.H., Sun, L.T., Tsai, C.L., Song, Y.L., Chang, C.F., 2002. Cold-stress
554 induced the modulation of catecholamines, cortisol, immunoglobulin M,
555 and leukocyte phagocytosis in tilapia. Gen. Comp. Endocr. 126(1), 90-100.
- 556 Claësson, D., Wang, T., & Malte, H. (2016). Maximal oxygen consumption
557 increases with temperature in the European eel (*Anguilla anguilla*) through
558 increased heart rate and arteriovenous extraction. Conserv. Physiol. 4(1),
559 cow027.

- 560 Colchen, T., Teletchea, F., Fontaine, P., Pasquet, A., 2017. Temperature modifies
561 activity, inter-individual relationships and group structure in a fish. *Curr.*
562 *Zool.* 63(2), 175-183.
- 563 Das, S.K., Noor, N.M., Kai, K.S., Juan, Q.Z., Iskandar, N.S.M., De, M., 2018.
564 Effects of temperature on the growth, gastric emptying time, and oxygen
565 consumption rate of mahseer (*Tor tambroides*) under laboratory
566 conditions. *Aquac. Rep.* 12, 20-24.
- 567 Das, T., Pal, A.K., Chakraborty, S.K., Manush, S.M., Sahu, N.P., Mukherjee, S.C.,
568 2005. Thermal tolerance, growth and oxygen consumption of *Labeo rohita*
569 fry (Hamilton, 1822) acclimated to four temperatures. *J. Therm.*
570 *Biol.* 30(5), 378-383.
- 571 Debnath, D., Pal, A.K., Sahu, N.P., Baruah, K., Yengkokpam, S., Das, T., Manush,
572 S.M., 2006. Thermal tolerance and metabolic activity of yellowtail catfish
573 *Pangasius pangasius* (Hamilton) advanced fingerlings with emphasis on
574 their culture potential. *Aquaculture* 258(1-4), 606-610.
- 575 Degani, G., Ben-Zvi, Y., Levanon, D., 1989. The effect of different protein levels
576 and temperatures on feed utilization, growth and body composition of
577 *Clarias gariepinus* (Burchell 1822). *Aquaculture* 76(3-4), 293-301.
- 578 Dhillon, R.S., Fox, M.G., 2007. Growth- independent effects of a fluctuating
579 thermal regime on the life- history traits of the Japanese medaka (*Oryzias*
580 *latipes*). *Ecol. Freshw. Fish* 16(3), 425-431.

- 581 Dickerson, B.R., Vinyard, G.L., 1999. Effects of high chronic temperatures and
582 diel temperature cycles on the survival and growth of Lahontan cutthroat
583 trout. *Trans. Am. Fish. Soc.* 128(3), 516-521.
- 584 Ebrahimi, M.H., Imanpour, M.R., Adlo, M.N., 2011. The effects of temperature
585 on growth, survival and some hematological parameters of giant gourami
586 (*Osphronemus goramy* Lacepède, 1801). *Iranian Journal of Biology* 24(5),
587 648-654.
- 588 Eldridge, W.H., Sweeney, B.W., Law, J.M., 2015. Fish growth, physiological
589 stress, and tissue condition in response to rate of temperature change
590 during cool or warm diel thermal cycles. *Can. J. Fish. Aquat. Sci.* 72(10),
591 1527-1537.
- 592 FAO, 2020. Fishery and aquaculture statistics. Global aquaculture production
593 1950-2018 (FishstatJ). FAO Fisheries and Aquaculture Department, Rome,
594 Italy.
- 595 Ford, J., Tibbetts, I., Carseldine, L., 2004. Ventilation rate and behavioural
596 responses of two species of intertidal goby (Pisces: Gobiidae) at extremes
597 of environmental temperature. *Hydrobiologia* 528(1-3), 63-73.
- 598 Frisk, M., Skov, P.V., Steffensen, J.F., 2012. Thermal optimum for pikeperch
599 (*Sander lucioperca*) and the use of ventilation frequency as a predictor of
600 metabolic rate. *Aquaculture* 324, 151-157.
- 601 Fu, K.K., Fu, C., Qin, Y.L., Bai, Y., Fu, S.J., 2018. The thermal acclimation rate
602 varied among physiological functions and temperature regimes in a
603 common cyprinid fish. *Aquaculture* 495, 393-401.

- 604 González, R.A., Díaz, F., Licea, A., Re, A.D., Sánchez, L.N., García-Esquivel, Z.,
605 2010. Thermal preference, tolerance and oxygen consumption of adult
606 white shrimp *Litopenaeus vannamei* (Boone) exposed to different
607 acclimation temperatures. J. Therm. Biol. 35(5), 218-224.
- 608 Hanke, I., Ampe, B., Kunzmann, A., Gärdes, A., Aerts, J., 2019. Thermal stress
609 response of juvenile milkfish (*Chanos chanos*) quantified by ontogenetic
610 and regenerated scale cortisol. Aquaculture 500, 24-30.
- 611 Hevrøy, E.M., Waagbø, R., Torstensen, B.E., Takle, H., Stubhaug, I., Jørgensen,
612 S.M., Torgersen, T., Tvenning, L., Susort, S., Breck, O., Hansen, T., 2012.
613 Ghrelin is involved in voluntary anorexia in Atlantic salmon raised at
614 elevated sea temperatures. General and Comparative
615 Endocrinology 175(1), 118-134.
- 616 Hölker, F., 2006. Effects of body size and temperature on metabolism of bream
617 compared to sympatric roach. Anim. Biol. 56(1), 23-37.
- 618 Iwama, G., Afonso, L.O.B., Vijayan, M., Claiborne, J., 2005. Stress in fishes, in:
619 Evans, D.H., Claiborne, J.B. (Eds.), The Physiology of Fishes. CRC Press,
620 Boca Raton, Florida, pp. 319-342.
- 621 Killen, S.S., Marras, S., Metcalfe, N.B., McKenzie, D.J., Domenici, P., 2013.
622 Environmental stressors alter relationships between physiology and
623 behaviour. Trends Ecol. Evol., 28(11), 651-658.
- 624 Konstantinov, A.S., Zdanovich, V.V., Tikhomirov, D.G., 1990. Effect of
625 temperature oscillation on the metabolic rate and energetics of juvenile
626 fish. J. Ichthyol. 30, 38-47.

- 627 Kristanto, A.H., Slembrouck, J., Subagja, J., Pouil, S., Arifin, O.Z., Prakoso, V.A.,
628 Legendre, M., 2020. Survey on egg and fry production of giant gourami
629 (*Osphronemus goramy*): Current rearing practices and recommendations
630 for future research. J. World Aquac. Soc. 51(1), 119-138.
- 631 Long, Y., Li, L., Li, Q., He, X., Cui, Z., 2012. Transcriptomic characterization of
632 temperature stress responses in larval zebrafish. PLoS one 7(5), e37209.
- 633 Majhi, S.K., Das, S.K., 2013. Thermal tolerance, oxygen consumption and stress
634 response in *Danio dangila* and *Brachydanio rerio* (Hamilton, 1822)
635 acclimated to four temperatures. Turk. J. Fish Aquat. Sc. 13(2), 359-365.
- 636 Maskur, M., Rina, R., Hamid, M.A., 2013. Small-scale freshwater aquaculture
637 extension development in Indonesia.
638 [https://enaca.org/?id=189&title=small-scale-aquaculture-extension-in-](https://enaca.org/?id=189&title=small-scale-aquaculture-extension-in-indonesia)
639 [indonesia](https://enaca.org/?id=189&title=small-scale-aquaculture-extension-in-indonesia), Accessed date: 29 November 2018.
- 640 McKenzie, D., Claireaux, G., 2010. The effects of environmental factors on the
641 physiology of aerobic exercise, in: Domenici, P., Kapoor, B.G.
642 (Eds.), Fish Locomotion: An Eco-ethological Perspective. Science
643 Publishers, Enfield, New Hampshire, USA, pp. 296-332.
- 644 Meeuwig, M.H., Dunham, J.B., Hayes, J.P., Vinyard, G.L., 2004. Effects of
645 constant and cyclical thermal regimes on growth and feeding of juvenile
646 cutthroat trout of variable sizes. Ecol. Freshw. Fish 13(3), 208-216.
- 647 Michalsen, K., Ottersen, G., Nakken, O., 1998. Growth of North-east Arctic cod
648 (*Gadus morhua* L.) in relation to ambient temperature. ICES J. Mar.
649 Sci. 55(5), 863-877.

650 Mommsen, T.P., Vijayan, M.M., Moon, T.W., 1999. Cortisol in teleosts:
651 Dynamics, mechanisms of action, and metabolic regulation. *Rev. Fish Biol.*
652 *Fish.* 9(3), 211-268.

653 Mueller, J.S., Grabowski, T.B., Brewer, S.K., Worthington, T.A., 2017. Effects of
654 temperature, total dissolved solids, and total suspended solids on survival
655 and development rate of larval Arkansas River Shiner. *J. Fish Wildl.*
656 *Manag.* 8(1), 79-88.

657 Pérez-Casanova, J.C., Afonso, L.O.B., Johnson, S.C., Currie, S., Gamperl, A.K.,
658 2008a. The stress and metabolic responses of juvenile Atlantic cod *Gadus*
659 *morhua* L. to an acute thermal challenge. *J. Fish Biol.* 72, 899-916.

660 Pérez-Casanova, J.C., Rise, M.L., Dixon, B., Afonso, L.O.B., Hall, J.R., Johnson,
661 S.C., Gamperl, A.K., 2008b. The immune and stress responses of Atlantic
662 cod to long-term increases in water temperature. *Fish Shellfish Immunol.*
663 24, 600-609.

664 Pisano, O.M., Kuparinen, A., Hutchings, J.A., 2019. Cyclical and stochastic
665 thermal variability affects survival and growth in brook trout. *J. Therm.*
666 *Biol.* 84, 221-227.

667 Pörtner, H.O., 2002. Climate variations and the physiological basis of temperature
668 dependent biogeography: Systemic to molecular hierarchy of thermal
669 tolerance in animals. *Comp. Biochem. Physiol. Part A Mol. Integr. Physiol.*
670 132, 739-761.

671 Pörtner, H.O., Bennett, A.F., Bozinovic, F., Clarke, A., Lardies, M.A., Lucassen,
672 M., Pelster, B., Schiemer, F., Stillman, J.H., 2006. Trade-off s in thermal

673 adaptation: The need for a molecular to ecological integration. *Physiol.*
674 *Biochem. Zool.* 79, 295-313.

675 Prakoso, V.A., Kurniawan, 2020. Oxygen consumption of barred
676 loach *Nemacheilus fasciatus* (Valenciennes, 1846) on different
677 temperatures. *IOP Conf. Ser.: Earth Environ. Sci.* 457, 012065.

678 Prakoso, V.A., Pouil, S., Prabowo, M.N.I., Sundari, S., Arifin, O.Z., Subagja, J.,
679 Affandi, R., Kristanto, A.H., Slembrouck, J., 2019. Effects of temperature
680 on the zootechnical performances and physiology of giant gourami
681 (*Osphronemus goramy*) larvae. *Aquaculture* 510, 160-168.

682 R Development Core Team, 2020. R: A language and environment for statistical
683 computing. R Foundation for Statistical Computing, Vienna, Austria.

684 Réalis-Doyelle, E., Gisbert, E., Alcaraz, C., Teletchea, F., Pasquet, A., 2018.
685 Temperature affects growth allometry and development patterns in brown
686 trout (*Salmo trutta*) fry: A multitrait approach. *Can. J. Fish. Aquat.*
687 *Sci.*, 75(5), 714-722.

688 Remen, M., Nederlof, M.A., Folkedal, O., Thorsheim, G., Sitjà-Bobadilla, A.,
689 Pérez-Sánchez, J., Oppedal, F., Olsen, R.E., 2015. Effect of temperature
690 on the metabolism, behaviour and oxygen requirements of *Sparus aurata*.
691 *Aquac. Environ. Interact.* 7(2), 115-123.

692 Rudneva, I., 2013. Biomarkers for Stress in Fish Embryos and Larvae. CRC Press,
693 Boca Raton, Florida.

- 694 Sarah, S., Widanarni, Sudrajat, A.O., 2009. Pengaruh padat penebaran terhadap
695 pertumbuhan dan kelangsungan hidup benih ikan gurame (*Osphronemus*
696 *goramy* Lac.). Jurnal Akuakultur Indonesia, 8(2), 199-207 (in Indonesian).
- 697 Sarma, K., Pal, A.K., Ayyappan, S., Das, T., Manush, S.M., Debnath, D., Baruah,
698 K., 2010. Acclimation of *Anabas testudineus* (Bloch) to three test
699 temperatures influences thermal tolerance and oxygen consumption. Fish
700 Physiol. Biochem. 36(1), 85-90.
- 701 Singh, S.P., Sharma, J.G., Ahmad, T., Chakrabarti, R., 2013. Effect of water
702 temperature on the physiological responses of Asian catfish *Clarias*
703 *batrachus* (Linnaeus 1758). Asian Fish. Sci. 26, 26-38.
- 704 Slembrouck, J., Arifin, O.Z., Pouil, S., Subagja, J., Yani, A., Kristanto, A.H.,
705 Legendre, M., 2019. Gender identification in farmed giant gourami
706 (*Osphronemus goramy*): A methodology for better broodstock
707 management. Aquaculture 498, 388-395.
- 708 SNI, 2000. SNI: 01-6485.3-2000: Produksi benih ikan gurame (*Osphronemus*
709 *goramy*, Lac) kelas benih sebar. Badan Standardisasi Nasional (BSN),
710 Jakarta, Indonesia. (in Indonesian)
- 711 Souchon, Y., Tissot, L., 2012. Synthesis of thermal tolerances of the common
712 freshwater fish species in large Western Europe rivers. Knowl. Manag.
713 Aquat. Ecosyst. 405, 03.
- 714 Sswat, M., Stiasny, M.H., Jutfelt, F., Riebesell, U., Clemmesen, C., 2018. Growth
715 performance and survival of larval Atlantic herring, under the combined
716 effects of elevated temperatures and CO₂. PLoS One, 13(1), e0191947.

- 717 Tromp, J.J., Jones, P.L., Brown, M.S., Donald, J.A., Biro, P.A., Afonso, L.O.,
718 2018. Chronic exposure to increased water temperature reveals few
719 impacts on stress physiology and growth responses in juvenile Atlantic
720 salmon. *Aquaculture* 495, 196-204.
- 721 Tromp, J.J., Jones, P.L., Symonds, J.E., Walker, S.P., Pope, A., Pether, S.M.,
722 Afonso, L.O., 2016. Effects of commercial diets and temperature on the
723 growth performance and stress response of hapuku (*Polyprion*
724 *oxygeneios*). *Aquaculture* 452, 128-133.
- 725 Turker, H., 2011. The effect of water temperature on standard and routine
726 metabolic rate in two different sizes of Nile tilapia. *Kafkas Univ. Vet. Fak.*
727 *Derg.* 17(4), 575-80.
- 728 Verawati, Y., Muarif, Mumpuni, F.Y., 2015. Pengaruh perbedaan padat penebaran
729 terhadap pertumbuhan dan kelangsungan hidup benih ikan gurami
730 (*Osphronemus gouramy*) pada sistem resirkulasi. *Jurnal Mina Sains*, 1(1),
731 6-12 (in Indonesian).
- 732 Verhille, C.E., English, K.K., Cocherell, D.E., Farrell, A.P., Fangué, N.A., 2016.
733 High thermal tolerance of a rainbow trout population near its southern
734 range limit suggests local thermal adjustment. *Conserv. Physiol.* 4(1),
735 cow057.
- 736 Watanabe, W.O., Ernst, D.H., Chasar, M.P., Wicklund, R.I., Olla, B.L., 1993. The
737 effects of temperature and salinity on growth and feed utilization of
738 juvenile, sex-reversed male Florida red tilapia cultured in a recirculating
739 system. *Aquaculture* 112(4), 309-320.

740 Whitney, J.E., Al-Chokhachy, R., Bunnell, D.B., Caldwell, C.A., Cooke, S.J.,
741 Eliason, E.J., Rogers, M., Lynch, A.J., Paukert, C.P., 2016. Physiological
742 basis of climate change impacts on North American inland fishes.
743 Fisheries 41, 332-345.

744 Zaragoza, O.D.R., Rodríguez, M.H., Bückle Ramirez, L.F., 2008. Thermal stress
745 effect on tilapia *Oreochromis mossambicus* (Pisces: Cichlidae) blood
746 parameters. Mar. Freshw. Behav. Physiol. 41(2), 79-89.

747 **Captions to figures**

748

749 **Figure 1.** Illustration of the experimental protocol used to test for the effect of the
750 daily temperature fluctuations on the growth performance and physiology of giant
751 gourami *Osphronemus goramy* (indoor recirculating systems, 12L:12D). The
752 continuous lines show the daily fluctuating (25-35 °C) and three constant (25, 30
753 and 35°C) temperature regimes. T_{opt} is the optimal temperature for giant gourami
754 larval rearing (Prakoso et al., 2020).

755

756 **Figure 2.** Percentages of (A) mortality and (B) abnormality observed in giant
757 gourami *Osphronemus goramy* larvae (n = 4) reared at one daily fluctuating (25-
758 35 °C) and three constant (25, 30 and 35 °C) temperature regimes for 21 days.
759 Values are means ± SD. Asterisks denote significant differences: *(p < 0.05), ***
760 (p < 0.001), **** (p < 0.0001) between treatments.

761

762 **Figure 3.** (A) Body mass (mg) and (B) total length (mm) of giant gourami
763 *Osphronemus goramy* larvae (n = 4 pools of 22 larvae) reared at four temperature
764 regimes from 8 days post-hatching. Values are means ± SD. Different letters
765 denote significant differences (p < 0.05) between temperature regimes.


766

767 **Figure 4.** Concentration of cortisol and glucose (n = 4 pools of 10 fish)
768 concentrations in the blood of giant gourami *Osphronemus goramy* larvae reared
769 at one daily fluctuating temperature (25-35 °C) and three constant (25, 30 and


770 35 °C) temperature regimes for 21 days. Values are means \pm SD. Asterisks denote
771 significant differences: * ($p < 0.05$) between regimes.

772



773 **Figure 5.** Oxygen consumption and ventilation rates ($n = 3$ batches of 21 larvae)
774 concentrations in the blood of giant gourami *Osphronemus goramy* larvae reared
775 at one daily fluctuating (25-35 °C) and three constant (25, 30, and 35 °C)
776 temperature regimes for 21 days. Values are means \pm SD. Asterisks denote
777 significant differences: * ($p < 0.05$), *** ($p < 0.001$), **** ($p < 0.0001$) between
778 treatments.




779 **Figure 1**




780 **Figure 2**



781 **Figure 3**



782 **Figure 4**



783 **Figure 5**

784 Table 1. Summary of water quality parameters measured for the one fluctuating
 785 (25-35°C) and the three constant (25 °C, 30 °C and 35 °C) temperature regimes
 786 during the experiment. Values are means \pm SD. For each parameter, values with
 787 different letters are significantly different ($p < 0.05$) between temperature
 788 treatments.

Parameter	n	Temperature treatment			
		Fluctuating (25-35 °C)	Constant (25 °C)	Constant (30 °C)	Constant (35 °C)
Temperature (°C)	66	29.0 \pm 3.4 ^a	24.8 \pm 0.1 ^b	29.9 \pm 0.1 ^c	34.8 \pm 0.1 ^d
Temperature range (°C)	66	24.7-34.4 ^a	24.6-25.2 ^b	29.6-30.2 ^c	34.4-35.2 ^d
DO (mg L ⁻¹)	16	7.81 \pm 0.25 ^a	7.77 \pm 0.18 ^a	6.98 \pm 0.16 ^b	6.59 \pm 0.11 ^b
pH	16	7.32 \pm 0.46 ^a	7.47 \pm 0.38 ^a	7.67 \pm 0.17 ^a	7.66 \pm 0.51 ^a
Conductivity (μ S cm ⁻¹)	16	158.3 \pm 8.1 ^a	172.9 \pm 3.6 ^{ab}	201.3 \pm 7.8 ^c	186.5 \pm 5.6 ^{bc}
TAN (mg L ⁻¹)	12	0.24 \pm 0.08 ^a	0.95 \pm 0.97 ^a	0.24 \pm 0.11 ^a	0.25 \pm 0.09 ^a
NH ₃ (mg L ⁻¹)	12	0.006 \pm 0.007 ^a	0.013 \pm 0.010 ^a	0.007 \pm 0.004 ^a	0.012 \pm 0.001 ^a
NH ₄ ⁺ (mg L ⁻¹)	12	0.23 \pm 0.08 ^a	0.93 \pm 0.96 ^a	0.23 \pm 0.11 ^a	0.23 \pm 0.08 ^a
NO ₂ ⁻ (mg L ⁻¹)	12	0.022 \pm 0.017 ^a	0.029 \pm 0.01 ^a	0.024 \pm 0.013 ^a	0.033 \pm 0.025 ^a

789 DO: dissolved oxygen; TAN: total ammonia nitrogen

790 Table 2. Growth performances of giant gourami *Osphronemus goramy* larvae
 791 reared at one fluctuating (25-35 °C) and three constant temperature (25 °C, 30 °C,
 792 and 35 °C) for 21 days from 8 days post-hatching. See Materials and Methods for
 793 details on the parameters. Values are means \pm SD. For each parameter, values
 794 with different letters are significantly different ($p < 0.05$) between temperature
 795 treatments.

Parameter	n	Temperature treatment			
		Fluctuating (25-35 °C)	Constant (25 °C)	Constant (30 °C)	Constant (35 °C)
BM _i (mg)	4*	12.0 \pm 1.1 ^a	11.6 \pm 1.1 ^a	11.7 \pm 1.1 ^a	13.0 \pm 1.4 ^a
BM _f (mg)	4*	178.0 \pm 38.1 ^a	147.1 \pm 30.6 ^b	276.7 \pm 47.5 ^c	262.9 \pm 72.2 ^c
TL _i (mm)	4*	9.3 \pm 0.3 ^a	9.3 \pm 0.3 ^a	9.4 \pm 0.2 ^a	9.4 \pm 0.2 ^a
TL _f (mm)	4*	22.8 \pm 1.7 ^a	21.2 \pm 1.6 ^b	26.1 \pm 1.4 ^c	24.9 \pm 2.4 ^d
K (mg mm ⁻³ x 100)	4*	1.48 \pm 0.18 ^a	1.53 \pm 0.18 ^{ab}	1.54 \pm 0.09 ^b	1.67 \pm 0.28 ^c
SGR _{BM} (% d ⁻¹)	4	12.8 \pm 0.4 ^a	12.1 \pm 0.5 ^a	15.1 \pm 0.2 ^b	14.3 \pm 0.5 ^b
SGR _{TL} (% d ⁻¹)	4	4.3 \pm 0.1 ^a	3.9 \pm 0.2 ^b	4.9 \pm 0.1 ^c	4.6 \pm 0.2 ^c
CV _{BM} (%)	4	19.8 \pm 3.3 ^{ab}	18.1 \pm 1.2 ^{ab}	17.1 \pm 2.7 ^a	26.4 \pm 7.9 ^b
CV _{TL} (%)	4	7.1 \pm 1.9 ^a	6.6 \pm 1.5 ^a	5.5 \pm 0.8 ^a	9.3 \pm 3.2 ^a
BG (g)	4	11.1 \pm 0.7 ^{ab}	8.5 \pm 1.6 ^a	20.7 \pm 5.2 ^b	8.4 \pm 2.4 ^a
FCR	4	7.33 \pm 0.35 ^{ab}	8.47 \pm 1.64 ^{ac}	4.75 \pm 0.92 ^b	10.58 \pm 2.27 ^c

796 BG: biomass gain, BM: body mass, BM_i: initial body mass, BM_f: final body mass,
 797 CV: coefficient of variation, FCR: feed conversion ratio, K: Fulton's condition
 798 factor, TL: total length, SGR: specific growth rate, TL_i: initial total length, TL_f:
 799 final total length.

800 * For BM, TF and K, 22 larvae were sampled from each of the four tanks.

801 Table 3. Parameters (Mean \pm SE, n = 20 at each time) of exponential growth in
 802 body mass (BM, mg) and total length (TL, mm) of giant gourami (*Osphronemus*
 803 *goramy*) larvae reared in four different temperature treatments. Model parameters:
 804 a: growth rate (d^{-1}). R^2 : determination coefficient. *** Probability of the model
 805 adjustment: $p < 0.001$

Temperature treatment	a \pm SE	R^2
Body mass (BM)		
25-35 °C	0.128 \pm 0.001***	0.95
25 °C	0.121 \pm 0.001***	0.96
30 °C	0.150 \pm 0.001***	0.97
35 °C	0.143 \pm 0.001***	0.93
Total length (TL)		
25-35 °C	0.042 \pm 0.001***	0.97
25 °C	0.039 \pm 0.001***	0.98
30 °C	0.049 \pm 0.001***	0.99
35 °C	0.047 \pm 0.001***	0.96

806