



**HAL**  
open science

# Diophantine Gaussian excursions and random walks

Raphaël Lachièze-Rey

► **To cite this version:**

| Raphaël Lachièze-Rey. Diophantine Gaussian excursions and random walks. 2021. hal-03197690v2

**HAL Id: hal-03197690**

**<https://hal.science/hal-03197690v2>**

Preprint submitted on 27 May 2021 (v2), last revised 21 Sep 2022 (v4)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Diophantine Gaussian excursions and random walks

Raphaël Lachieze-Rey \*

**Abstract** We investigate the asymptotic variance of Gaussian nodal excursions in the Euclidean space, focusing on the case where the spectral measure has incommensurable atoms. This study requires to establish fine recurrence properties around 0 for the associated irrational random walk on the torus. We show in particular that the recurrence magnitude depends strongly on the diophantine properties of the atoms, and the same goes for the variance asymptotics of nodal excursions.

More specifically, if the spectral measures contains atoms which ratios are well approximable by rationals, the variance is likely to have large fluctuations as the observation window grows, whereas the variance is bounded by the  $(d-1)$ -dimensional measure of the window boundary if these ratio are badly approximable. We also show that, given any reasonable variance asymptotic behaviour, there are uncountably many sets of parameters that realise it.

The general formula we derive for the variance asymptotics of Gaussian excursions is also used in a different context, to show a variance cancellation phenomenon for Gaussian random waves in any dimension.

**Keywords:** Gaussian fields, nodal excursion, random walk, diophantine approximation, hyperuniformity, Gaussian random waves, variance cancellation.

**MSC Classification:** 60G15, 60G50, 11J13, 34L20

## 1 Introduction and examples

The primary motivation of this article is to study the variance of the excursion volume for Gaussian fields, and exhibit a parametric *diophantine* model that realises any prescribed asymptotic variance behavior. It turns out that this work is strongly related to the random walk which increment measure is the spectral measure of the Gaussian field, and which behaviour depends on the diophantine properties of the atoms.

To conduct this program, we hence establish two unrelated results which are of independent interest. The first one, Theorem 6, contains general bounds for the variance of Euclidean Gaussian fields; its versatility is illustrated by the application to fields of a fundamentally different nature, the Gaussian random waves (see Section 1.5). The second one, Theorem 5, contains bounds for random walks with irrational increments. These results can be projected on the torus and adds some uniform estimates to the existing literature (see Section 1.6).

### 1.1 Gaussian excursions

Properties of excursions and level sets of continuous random Gaussian functions have been studied under many different instances. The zero set of a one-dimensional Gaus-

---

\*Laboratoire MAP5, Université de Paris, 45 Rue des Saints-Pères, raphael.lachieze-rey@parisdescartes.fr

sian stationary process is the subject of an almost century long line of research, starting with the seminal works of Kac & Rice [17], or Cramer & Leadbetter [8], and followed by many other authors [9, 30, 18], mainly interested by second order behaviour. In higher dimensions, zeros of Gaussian entire functions [14, 25] and nodal sets of high energy Gaussian harmonics on a compact manifold [19, 36, 22] (and their Euclidean counterpart the Random Wave Model [24]) have attracted a lot of attention from both physicists and mathematicians. Random trigonometric Gaussian polynomials, i.e. independent Gaussian coefficients multiplied by trigonometric monomials based on a fundamental frequency, have also been studied in the asymptotics of the large degree, see for instance [37] and references therein. We propose here a crucial perturbation of the spectral measure support: instead of taking frequencies in a proportionnal relation, we choose finitely many frequencies which are incommensurable; this specificity allows for instance to reach all possible behaviours for the variance asymptotics.

A different approach to our results is through the lens of hyperuniform models. In the last decades, physicist have put in evidence states of matter intermediate between crystals and liquids, where the medium exhibits apparent disorder at all scales, but fluctuations are suppressed at large scales. This denotes in some sense an internal compensation of the medium behaviour, and is considered by physicists a *new state of matter*, see the works by S. Torquato and his co-authors [34, 35] that introduce the topic and expose the main tools and discoveries. Even though the focus was primarily on atomic measures, i.e. point configurations, this concept has been then generalised to other random measures, in particular bi-phased random media [33, 32]. Such heterogeneous materials abound in nature and synthetic situations. Examples include composite and porous media, metamaterials, biological media (e.g., plant and animal tissue), foams, polymer blends, suspensions, granular media, cellular solids, colloids.

The Gaussian world provides models for many types of phenomena, and the present work is an attempt to produce Gaussian hyperuniform random sets. The model we present here shares some similarities with perturbed lattices, in the sense that the long range correlations are very strong, but its disorder state is also one step above as one cannot write it as the (perturbed) repetition of a given pattern. It shares with quasi-crystals the property of *almost periodicity*, defined below, and exhibits a spectrum reminiscent of quasi-crystals, see Fig 1. Any asymptotic variance can be achieved, yielding in particular hyperuniform models. According to the typology established in [34, 6.1.2], the model is type-I hyperuniform for almost all choice of parameters; but uncountably many choices of the parameters will yield type-II or actually any type of hyperuniformity. We also give randomised versions of the model not involving diophantine parameters which exhibit different types of hyperuniformity.

As it turns out, it seems that a non-isotropic model is necessary to obtain a hyperuniform behaviour. We use the general variance formula of Theorem 6 to study Gaussian random waves in any dimension and prove a variance cancellation phenomenon, computations seem to indicate that isotropic Gaussian excursions might never be hyperuniform.

The main actors of this article are real Gaussian random fields  $\{X(\mathbf{t}); \mathbf{t} \in \mathbb{R}^d\}$ , which law is invariant under translations of  $\mathbb{R}^d$ . It is known that such fields are completely characterised by their reduced covariance function

$$C(\mathbf{t}) = \mathbb{E}(X(0)X(\mathbf{t})), \mathbf{t} \in \mathbb{R}^d,$$

or by their spectral measure, i.e. the unique finite symmetric measure  $\boldsymbol{\mu}$  on  $\mathbb{R}^d$  such that

$C$  admits the representation

$$C(\mathbf{t}) = \int_{\mathbb{R}^d} e^{i\mathbf{t}\cdot\mathbf{x}} \boldsymbol{\mu}(d\mathbf{x}), \mathbf{t} \in \mathbb{R}^d, \quad (1)$$

where  $\cdot$  denotes the standard scalar product. See the monograph [1] for a comprehensive expositions of main properties and results about Gaussian fields and their geometry.

Excursions of Gaussian processes on the real line have often been studied through their number of crossings with the axis [17, 8, 9, 30, 18]. Elementary considerations yield that the average number of crossings on an interval is proportionnal to the length of the interval. Furthermore, if  $\mu$  contains more than 1 (symmetrised) atom, the variance of the number of crossings is quadratic [21]. On  $\mathbb{R}^d$ , we rather focus on the Lebesgue measure of the nodal excursions

$$\{X > 0\} = \{\mathbf{t} \in \mathbb{R}^d : X(\mathbf{t}) > 0\}.$$

Here again, the field centering yields that the expectation is proportionnal to the volume:

$$\mathbb{E}(\mathcal{L}^d(A \cap \{X > 0\})) = \frac{\mathcal{L}^d(A)}{2}, A \subset \mathbb{R}^d.$$

We give in Section 4 general upper and lower bounds for the variance of the excursion volume. The need for models that yield any prescribed variance asymptotics is explained in [6], along with another such procedure based on the Fourier transform.

We consider models where the spectral measure is finite and observe that the excursion variance can have any prescribed behaviour. They fall within the class of *almost periodic fields*, and their properties seem to put them between ergodic models, and crystalline arrangements. We show that, depending on the diophantine properties of the atoms, the variance can either be minimal, i.e. of *surface-scaling order* in  $T^{d-1}$ , or maximal, i.e. of *quadratic order* in  $T^{2d}$ , and achieves any reasonable intermediate behaviour. For sublinear variances (below  $T^d$ ), the excursions are hence hyperuniform, contributing to the already large research body on the subject. We also consider Gaussian random waves to illustrate the generality of the variance formula established.

## 1.2 Variance asymptotics for diophantine measures

We consider symmetric spectral measures whose support contains incommensurate atoms. A model that offers a lot of diversity in  $\mathbb{R}^d$  is parametrized by a  $d$ -tuple  $\boldsymbol{\omega}$  where each entry belongs to  $\mathbb{R}^m$  for some  $m \geq 1$ :  $\boldsymbol{\omega} = (\omega_{[k],i})_{\substack{1 \leq k \leq d \\ 1 \leq i \leq m}} \in (\mathbb{R}^m)^d$  and the measure

$$\boldsymbol{\mu} = \frac{1}{d(m+1)} \sum_{k=1}^d (\bar{\delta}_{\mathbf{e}_k} + \sum_{i=1}^m \bar{\delta}_{\omega_{[k],i} \mathbf{e}_k}), \text{ with } \bar{\delta}_u = \frac{1}{2}(\delta_u + \delta_{-u}), u \in \mathbb{R}^d, \quad (2)$$

where  $(\mathbf{e}_k)_{1 \leq k \leq d}$  is some basis of  $\mathbb{R}^d$ .

Denote by  $X_{\boldsymbol{\omega}}$  a centered stationary Gaussian random field which spectral measure is  $\boldsymbol{\mu}$ . The excursion volume variance is denoted by

$$V_{\boldsymbol{\omega}}(T) := \mathbf{Var}(\mathcal{L}^d(\{X_{\boldsymbol{\omega}} > 0\} \cap B_d(0, T)))$$

where  $B_d(0, T)$  is the centred ball of  $\mathbb{R}^d$  with radius  $T$ .

To describe the asymptotic behaviour, introduce for  $\psi : \mathbb{N}_* \rightarrow [0, 1]$  the set of  $\psi$ -BA (Badly Approximable) numbers, which are the  $\omega \in \mathbb{R}^m$  such that for  $p \in \mathbb{Z}, q \in \mathbb{Z}^m$  with  $|q|$  sufficiently large,

$$|p - \omega \cdot q| \geq 2\psi(|q|),$$

where  $|q|$  is the Euclidean norm of  $q$ , the typical example being  $\psi(|q|) = |q|^{-\tau}$ ,  $\tau > 0$ . On the other hand, a vector  $\omega \in \mathbb{R}^m$  is  $\psi$ -WA<sup>(\*)</sup> (Well Approximable) if for some  $c_W > 0$  there are infinitely many couples  $p \in \mathbb{Z}$ ,  $q \in \mathbb{Z}^m$  such that  $(p, q) \equiv 1$  and

$$|p - \omega \cdot q| \leq c_W \psi(|q|).$$

where here and in all the paper, for  $p \geq 1$  and  $\mathbf{x} = (x_i)_{1 \leq i \leq p} \in \mathbb{Z}^p$  we write  $\mathbf{x} \equiv 1$  if  $\sum_{i=1}^p x_i$  is an odd number.

**Definition 1.** Say that  $\psi$  is *regular* (or  $\tau$ -regular) if it is of the form  $\psi(q) = q^{-\tau} L(q)$ ,  $q \in \mathbb{N}^*$ , where  $\tau > 0$  and  $L$  does not vanish and is slowly varying in the sense that  $|L(q) - L(q+1)| = o(L(q)q^{-1})$  as  $q \rightarrow \infty$ .

This property is motivated by Proposition 7 and yields that  $\ln(L(q))/\ln(q) \rightarrow 0$  as  $q \rightarrow \infty$ , hence  $L$  is dominated by any power of  $q$  and  $\tau$  is uniquely defined. We introduce the pseudo-inverse  $\psi^{-1} : (0, 1] \mapsto \mathbb{N}$  by (5). We can show that for every finite  $r > 0$  there are finite  $c_i > 0$  such that  $c_1 \psi(q) \leq \psi(rq) \leq c_2 \psi(q)$  and  $c_3 \psi^{-1}(\varepsilon) \leq \psi^{-1}(r\varepsilon) \leq c_4 \psi^{-1}(\varepsilon)$  on their domains of definition. Remark that  $q\psi(q)$  is strictly non-increasing for sufficiently large  $q$  if  $\tau > 1$ .

Given two functions  $F(T), G(T), T \geq 1$ , write  $F(T) \stackrel{\text{i.o.}}{\leq} G(T)$  (infinitely often) if  $F(T_k) \leq G(T_k)$  for some sequence  $T_k \rightarrow \infty$ . Our most precise and general result concerns the case where frequencies of  $\boldsymbol{\mu}$  are the same in all  $d$  directions, it implies that diophantine Gaussian nodal excursions provide stationary random sets in  $\mathbb{R}^d$  with any reasonable asymptotic prescribed variance behaviour.

**Theorem 1.** Let  $\tau > 0$ ,  $\psi : \mathbb{N}_* \rightarrow (0, 1]$   $\tau$ -regular and  $\omega \in \mathbb{R}^m$  that is  $\psi$ -WA<sup>(\*)</sup> and  $\psi$ -BA, and  $\boldsymbol{\omega} := (\omega, \omega, \dots, \omega) \in (\mathbb{R}^m)^d$ . Then with  $\tau^* = \frac{1+d(m+1)}{1+d}$ , as  $T \rightarrow \infty$ ,

$$\begin{aligned} V_{\boldsymbol{\omega}}(T) &\leq c_+ T^{d-1} && \text{if } \psi(q) > q^{-\tau^*} \ln(q) \\ V_{\boldsymbol{\omega}}(T) &\leq c_+ T^{d-1} \ln(T) && \text{if } \psi(q) = q^{-\tau^*} \\ \frac{c_- T^{2d}}{\psi^{-1}(T^{-1})^{1+d(m+1)}} &\stackrel{\text{i.o.}}{\leq} V_{\boldsymbol{\omega}}(T) \leq \frac{c_+ T^{2d}}{\psi^{-1}(T^{-1})^{1+d(m+1)}} = o(T^{2d}) && \text{if } \tau > \tau^* \end{aligned}$$

for some  $0 < c_- \leq c_+ < \infty$  depending on  $d, m, \psi$ . If  $m = 1$  and  $\tau > 1$ , there are uncountably many such  $\omega \in \mathbb{R}^1$ .

**Remark 1.** The presence of the term  $T^{d-1}$  on the RHS, proportionnal to the surface measure of the observation window, is natural as random stationary measures applied to a large window are usually not expected to have a variance behaviour lower than the boundary measure. No rigorous general result in this direction is known by the author, Beck [2] gives a formal proof in the case of point processes. See also [34], which classifies hyperuniform behaviours in three types: type I have asymptotic variance in  $T^{d-1}$ , type II in  $T^{d-1} \ln(T)$ , and type III gathers all other sublinear behaviours. The previous theorem shows that all types can be reached with the current model. Intermediate behaviours between  $T^{d-1}$  and  $T^{d-1} \ln(T)$  can likely be obtained by the same method.

**Remark 2.** This type of behaviour is really specific of nodal excursions. The volume variance for excursions  $\{X > u\} \cap B_d(0, T)$  always behave in  $T^{2d}$  if  $u \neq 0$ , see Theorem 6. The phenomenon of variance cancellation at  $u = 0$  is heavily documented for Gaussian random waves (see Section 3).

*Proof.* To prove this theorem, apply first Theorem 6 to the measure  $\boldsymbol{\mu}$  to have bounds on the variance in terms of the function  $\mathbf{J}_3$  (with  $\gamma$  as the unit ball indicator function,

see Example 1). Proposition 7 yields the upper bound. Then lower bounds for  $\mathbf{J}_3$  are derived in Theorem 5, noticing that  $\omega$  is  $\psi$ -BA and  $\psi$ -SWA<sup>(\*)</sup> thanks to Proposition 6.

It is a standard fact in diophantine approximation that if  $m = 1$  the set of  $\omega$  that are  $\psi$ -BA and  $\psi$ -WA is uncountable when  $q\psi(q)$  is non-increasing at infinity, see the seminal construction based on continued fractions by Jarník [16]. □

Prominent examples are provided by power functions, for which we introduce a special notation: say that  $\omega \in \mathbb{R}^m$  is  $(\tau)$ -BA (resp.  $(\tau)$ -WA<sup>(\*)</sup>) if it is  $cq^{-\tau}$ -BA (resp.  $cq^{-\tau}$ -WA<sup>(\*)</sup>) for some finite  $c > 0$ . These considerations are further developed and commented in Section 2 (see Groshev's theorem and Corollary 2), let us simply mention that for  $\eta > 0$ ,  $\mathcal{L}^m$ -a.a.  $\omega \in \mathbb{R}^m$  is  $(m + \eta)$ -BA and  $(m)$ -WA<sup>(\*)</sup>. For any  $\eta \geq 0$ , there are uncountably many  $\omega \in \mathbb{R}^m$  that are  $(m + \eta)$ -BA and  $(m + \eta)$ -WA<sup>(\*)</sup>. There are also uncountably many *Liouville numbers*, i.e.  $\omega \in \mathbb{R}$  that are  $(\tau)$ -WA<sup>(\*)</sup> for any  $\tau > 0$ .

**Remark 3.** The random excursion is hyperuniform if  $\omega$  is  $(\eta)$ -BA with  $\eta < \frac{1+d}{d}$  (see also Section 1.3 for characterisation of hyperuniformity via the structure factor), and strongly hyperuniform if  $\eta < 1 - \frac{m}{d+1}$ .

The following corollary is then a consequence of Theorem 1.

**Corollary 1.** For  $\omega \in \mathbb{R}^m$ , let  $\omega = (\omega, \dots, \omega) \in (\mathbb{R}^m)^d$ ,  $\tau^* = \frac{1+d(m+1)}{1+d}$ .

- (i) If  $d \geq m$ , for  $\tau \in (m, \tau^*)$ ,  $\mathcal{L}^m$ -a.a.  $\omega \in \mathbb{R}^m$  is  $(\tau)$ -BA, and the variance of  $V_\omega(T)$  is bounded by  $c_+T^{d-1}$  for some  $c_+ < \infty$  (strong hyperuniformity).
- (ii) If  $d < m$ , since for  $\mathcal{L}^m$ -a.a.  $\omega \in \mathbb{R}^m$ ,  $\omega$  is  $(m)$ -WA<sup>(\*)</sup> and  $(m + \eta)$ -BA for  $\eta > 0$ , we have  $c_-T^{d-\frac{1+d}{m}} \stackrel{\text{i.o.}}{\leq} V_\omega(T) \leq c_+T^{d-\frac{1+d}{m+\eta}}$  for some  $0 < c_-, c_+ < \infty$ .
- (iii) Let  $m = 1$ . For  $d-1 \leq \beta < 2d$ , the set of  $\omega \in \mathbb{R}$  such that for some  $0 < c_- \leq c_+ < \infty$

$$c_-T^\beta \stackrel{\text{i.o.}}{\leq} V_\omega(T) \leq c_+T^\beta, T \geq 1$$

is uncountable ( there is  $\tau \geq 1$  such that  $\beta = 2d - \frac{1+2d}{\tau}$ , and uncountably many  $\omega$  are  $(\tau)$ -WA<sup>(\*)</sup> and  $(\tau)$ -BA ).

- (iv) Let  $m = 1$ . For  $\omega$  a Liouville number, for every  $\varepsilon > 0$ , for some  $c_- > 0$ ,

$$c_-T^{2d-\varepsilon} \stackrel{\text{i.o.}}{\leq} V_\omega(T).$$

- (v) In all cases,  $V_\omega(T) = o(T^{2d})$ .

**Remark 4.** An interesting observation is that for such a model, the variance of the excursion indicator might be minimal (  $T^{d-1}$ , e.g. bounded in dimension  $d = 1$ ), while the zero-counting measure, its derivative in the distributional sense, has maximal variance, in  $T^{2d}$  (see [21, Theorem 2-(iii)]).

If the  $\omega_{[k]}$  differ along the directions  $1 \leq k \leq d$ , a similar application of Theorem 5, Proposition 7, and Theorem 6 yield that the upper bound corresponds to the worst upper bound among the  $\omega_{[k]}$ :

**Proposition 1.** Let  $\psi : \mathbb{N}_* \rightarrow (0, 1]$  regular. Assume  $\omega \in (\mathbb{R}^d)^m$  is such that each  $\omega_{[k]}$  is  $\psi$ -BA,  $1 \leq k \leq d$ . Then the same upper bounds as in Theorem 1 hold. In particular

- (i) if  $d \geq m$ , for  $\mathcal{L}^{md}$ -a.a.  $\omega \in (\mathbb{R}^m)^d$ ,  $V_\omega(T) \leq c_+T^{d-1}$  for some  $c_+ < \infty$ .

(ii) for every  $\omega \in (\mathbb{R}^m)^d$ ,  $V_\omega(T) = o(T^{2d})$

We need a further notion to state the lower bound. Say that  $\omega = (\omega_{[k]})_k$  is  $\psi$ -SWA (Simultaneously Well Approximable) if for some  $c_W > 0$ , for infinitely many  $q^j \in \mathbb{Z}^m$ ,  $j \geq 1$ , there exist  $p_{[k]}^j \in \mathbb{Z}$ ,  $1 \leq k \leq d$  such that

$$|p_{[k]}^j - \omega_{[k]} \cdot q^j| < c_W \psi(|q^j|), 1 \leq k \leq d,$$

and say that it is  $\psi$ -SWA<sup>(\*)</sup> if furthermore  $\sum_{k=1}^d (p_{[k]}^j + \sum_{i=1}^m q_i^j)$  is odd. The need to distinguish between  $\psi$ -SWA and  $\psi$ -SWA<sup>(\*)</sup> (or  $\psi$ -WA and  $\psi$ -WA<sup>(\*)</sup>) is discussed in Remark 7. Theorem 5-(iii) and Theorem 6 yield:

**Theorem 2.** *Assume that for some function  $\psi : \mathbb{N}^* \rightarrow (0, 1]$  converging to 0,  $\omega$  is  $\psi$ -SWA<sup>(\*)</sup>. Then for some  $c_+ < \infty$*

$$T^{2d} \psi^{-1}(T^{-1})^{-1-d(m+1)} \stackrel{\text{i.o.}}{\leq} c_+ V_\omega(T)$$

where  $\psi^{-1}$  denotes the pseudo-inverse of  $\psi$  (see (5)).

Thanks to Groshev's theorem (see Section 2), for  $\eta > 0$ ,  $\mathcal{L}^{md}$ -a.a.  $\omega \in (\mathbb{R}^m)^d$  is  $|q|^{-m/d}$ -SWA but not  $|q|^{-m/d-\eta}$ -SWA.

### 1.3 Structure factor, quasicrystals and almost periodicity

We have observed, for some values of the parameter  $\omega$ , the suppression of the variance at large scales, also called hyperuniformity phenomenon. A more general mathematical indicator of hyperuniformity is the structure factor, or more generally the behaviour around zero of the Fourier transform of the signal in some sense. Here we adopt the more general definition of the structure factor of the random excursion indicator function  $\mathbf{t} \mapsto \mathbf{1}_{\{X(\mathbf{t}) > 0\}}$ : if it exists, it is the measure  $\mathcal{S}$  satisfying

$$\int_{\mathbb{R}^d} \varphi(\mathbf{t}) \text{Cov}(\mathbf{1}_{\{X(0) > 0, X(\mathbf{t}) > 0\}}) d\mathbf{t} = \int_{\mathbb{R}^d} \hat{\varphi}(\mathbf{x}) \mathcal{S}(d\mathbf{x}),$$

for  $\varphi$  smooth with compact support. Integrable hyperuniform models are characterised by  $\mathcal{S}(d\mathbf{x}) \sim |\mathbf{x}|^\alpha d\mathbf{x}$  near 0 (for some  $\alpha > 0$ ), which integrates in

$$\mathcal{S}(B_d(0, \varepsilon)) \sim \varepsilon^{\alpha+d}, \varepsilon \rightarrow 0.$$

Following [27], we adopt the latter characterization as hyperuniformity with index  $\alpha > 0$ . Let us consider a diophantine Gaussian field with spectral measure  $\boldsymbol{\mu}$ , as described in the previous sections. We easily deduce from Lemma 5 that, with a computation similar to (23)


$$\mathcal{S} = \sum_{n \text{ odd}} \frac{\binom{n}{2n}}{4^n (2n+1)} \boldsymbol{\mu}^{*(2n+1)}$$

$$\mathcal{S}(B_d(0, \varepsilon)) \leq c \mathbf{J}_3(\varepsilon) \leq c' \varepsilon^{\frac{1+d(m+1)}{m+\eta}} \text{ using (9)}$$

where  $m \geq 1$  is the number of different frequencies and  $\eta \geq 0$  is the diophantine approximation parameter of the model (see the previous section), and a similar lower bound holds for infinitely many  $\varepsilon \rightarrow 0$ ; hence the model is indeed hyperuniform for  $\eta < \frac{1+d}{d}$ , as found at Corollary 1-(i), and strongly hyperuniform (i.e.  $\alpha > 1$ ) for  $\eta < 1 - \frac{m}{d+1}$ .

We give an approximate representation at figure 1 in a special case.

Figure 1: Structure factor for  $\boldsymbol{\mu} = \bar{\delta}_{\mathbf{e}_1} + \bar{\delta}_{\sqrt{2}\mathbf{e}_1} + \bar{\delta}_{\mathbf{e}_2} + \bar{\delta}_{\sqrt{2}\mathbf{e}_2}$



This kind of spectrum is reminiscent of Bragg peaks in quasi-crystals, and more generally of almost periodic fields, for which we give a definition here: a field  $X : \mathbb{R}^d \rightarrow \mathbb{R}$  is almost periodic if for any sequence of vectors  $\mathbf{t}_n \rightarrow \infty$ , there is a subsequence  $\mathbf{t}_{n'}$  such that  $\|X - X(\mathbf{t}_{n'} + \cdot)\|_\infty \rightarrow 0$ , see for instance the monograph [7]. Covariance functions and random Gaussian fields considered in this paper are  $\|\cdot\|_\infty$ -limits of trigonometric polynomials, and as such they are almost periodic. On the other hand, their excursions, seen as  $\{0, 1\}$ -valued functions, are not almost periodic in this sense, mainly because of the discontinuities at the set boundary. On the other hand, they are likely almost periodic for weaker norms, and could hence be seen as *almost periodic sets*, but this question is outside the scope of the current article.

#### 1.4 A randomised model

It is easy to build randomised models that exploit the metric properties of diophantine approximation to yield hyperuniform models.

**Proposition 2.** *Let  $\Omega$  be a real random variable whose law is continuous with respect to Lebesgue measure, and let  $a_k^i, i \geq 0, k \geq 1$  be independent and identically distributed standard Gaussian variables. Define*

$$X(\mathbf{t}) = \frac{1}{2d} \sum_{k=1}^d (a_k^0 \cos(t_{[k]}) + a_k^1 \sin(t_{[k]}) + a_k^2 \cos(\Omega t_{[k]}) + a_k^3 \sin(\Omega t_{[k]})), \mathbf{t} = (t_{[k]}) \in \mathbb{R}^d,$$

$M_T = \mathcal{L}^d(\{X > 0\} \cap B_d(0, T))$  and  $V(T) = \mathbf{Var}(M_T)$ . Then for some  $c_+ < \infty$ ,

$$V(T) \leq c_+ T^{d-1}.$$

*Proof.* Since the Gaussian field is centered, for any fixed  $\omega \in \mathbb{R}$ ,

$$\mathbb{E}(M_T \mid \Omega = \omega) = \mathcal{L}^d(B_d(0, T))/2$$

is deterministic. We also know that a.a.  $\omega \in \mathbb{R}$  is (2)-BA, and if we condition by  $\Omega = \omega$ ,  $X$  is the Gaussian field with reduced covariance  $\frac{1}{2d} \sum_{k=1}^d (\cos(t_{[k]}) + \cos(\omega t_{[k]}))$ . Hence the



conditional variance formula and Corollary 1 yield

$$\begin{aligned} V(T) &= \mathbb{E}(\mathbf{Var}(M_T | \Omega)) + \mathbf{Var}(\mathbb{E}(M_T | \Omega)) \\ &\leq c_+ T^{d-1}. \end{aligned}$$

□

The same arguments with  $1 \leq m < d$  yield the following:

**Proposition 3.** *Let  $(\Omega_0, \dots, \Omega_m)$  a random  $(m+1)$ -tuple of vectors with continuous law with respect to  $\mathcal{L}^{(m+1)d}$ , and*

$$X(\mathbf{t}) = \frac{1}{d(m+1)} \sum_{k=1}^d \sum_{i=0}^m (a_k^{2i} \cos(\Omega_i t_k) + a_k^{2i+1} \sin(\Omega_i t_k)).$$

*Then the variance is bounded by  $c_+ T^{d-1}$  if  $d > m$ .*

Along similar lines, exploiting Corollary 1-(iii) with  $m > d$  yields randomised models which variance is in  $T^\beta$  for  $d-1 < \beta < 2d$ .

**Remark 5.** Similar models in the context of random walks (Section 3) yield interesting examples of random walks in a random environment.

## 1.5 Variance cancellation for Gaussian random waves

Let us give another (and unrelated) application of Theorem 6 in the context of ergodic isotropic fields. Let  $d \geq 2$ ,  $\mathbb{S}^{d-1} = \{\mathbf{t} \in \mathbb{R}^d : \|\mathbf{t}\| = 1\}$  the  $d$ -dimensional unit sphere and  $\mu_d$  the Haar distribution on the sphere, i.e. the unique probability measure on  $\mathbb{S}^{d-1}$  invariant under rotations. Let  $X_d(\mathbf{t})$  be a centred Gaussian random field with spectral measure  $\mu_d$  and reduced covariance

$$C_d(\mathbf{t}) = \int_{\mathbb{S}^{d-1}} \exp(i\mathbf{t} \cdot \mathbf{x}) \mu_d(d\mathbf{x}) = c_d \frac{\mathbf{B}_{\frac{d}{2}-1}(\|\mathbf{t}\|)}{\|\mathbf{t}\|^{\frac{d}{2}-1}}, \mathbf{t} \in \mathbb{R}^d \quad (3)$$

for some  $c_d > 0$ , see [13, (21)] and Example 1 for the definition of the Bessel function of the first kind  $\mathbf{B}_a$ .

The field  $X_d$ , called *Berry's random wave model*, is of central importance as it is the unique stationary isotropic field satisfying  $\Delta X_d = -X_d$  a.s. [26]. It can be seen as a local approximation of random eigenfunctions of the Laplacian on compact  $d$ -dimensional smooth manifolds, of high interest in the physics literature. Since nodal statistics are local quantities, it makes sense to expect analogies between the behaviours of random waves on different manifolds as  $T \rightarrow \infty$ . Such random Gaussian harmonics have been recently heavily studied in dimension 2 in the mathematics literature, especially on the sphere or the torus [19, 23, 12, 22], and the Euclidean version  $X_d$  has also been investigated, through its percolation properties [24] or the statistical properties of their nodal lines [26, 12]. An interesting property of Gaussian random harmonics is the variance cancellation phenomenon, i.e. the very small asymptotic fluctuations of some statistic of the excursion set at the level  $u = 0$  in the high energy limit, compared to other levels  $u \neq 0$ . First conjectured by Berry [3] for the length of the excursion boundary (*nodal lines*) on the torus, it has then been observed and deeply analysed in several other instances [19, 22].

We prove here that a variance cancellation at the level  $u = 0$  also occurs for the nodal excursions of the Euclidean Gaussian random waves in any dimension. More specifically, while the excursion volume is overfluctuating for levels  $u \neq 0$  (i.e. the volume of large

windows is negligible with respect to the variance of the excursions of  $X_d$  restricted to this window), the fluctuations are linear for  $u = 0$ , as would be the case for fluctuations of short range random fields such as the Bargmann-Fock field (see e.g.[12] for another example). For  $T > 0$ , the rescaled version  $X_{d,T}(\mathbf{t}) = X_d(T\mathbf{t})$  satisfies  $\Delta X_{d,T} = -T^2 X_{d,T}$  and hence can be compared to random harmonics with same wavelength on compact manifolds.

**Theorem 3.** *Denote by  $V^u(T)$  the variance of  $\mathcal{L}^d(B_d(0, 1) \cap \{X_{d,T} > 0\})$ . For  $u \neq 0$ , there is  $c_u > 0$  such that*

$$c_u T^{1-d} \leq V^u(T), T > 0$$

and there is  $0 < c_{-,0} \leq c_{+,0} < \infty$  such that

$$c_{-,0} T^{-d} \leq V^0(T) \leq c_{+,0} T^{-d}, T > 0.$$

The proof is at Section 4.1. This result can be compared with similar results on the sphere, see the work of Marinucci and Wigman in dimension 2 [23], and then of Rossi [29] in dimension  $d \geq 2$ , who study the excursion volume of spherical Gaussian harmonics  $X$  satisfying  $\Delta_{\mathbb{S}^2} X = -\ell(\ell + d - 1)X$ ,  $\ell \in \mathbb{N}$ , where  $\Delta_{\mathbb{S}^2}$  is the Laplace-Beltrami operator on the sphere. They also obtain a variance of magnitude  $\ell^{1-d}$  at levels  $u \neq 0$  and  $\ell^{-d}$  at the level  $u = 0$ , echoing experimental results from Blum, Gnutzmann and Smilansky [4]. Hence the present results are consistent with those obtained on the sphere, and shows that Theorem 6 can also deal with such variance cancellation phenomena.

**Remark 6.** It is likely that the same phenomenon occurs for the excursions of any isotropic Gaussian field which spectral measure is supported by a lower dimensional set, e.g. linear combinations of independent Gaussian random waves at different energy levels, see Section 4.1. Computations also hint at the fact that excursions (at any level) of stationary isotropic Gaussian fields will never be hyperuniform, but more general arguments are required for a formal proof.

## 1.6 Diophantine random walk on the torus

Let  $\mu$  be a symmetric measure on  $\mathbb{R}^d$  of the form (2) and  $\bar{\mathbf{U}}_n$  the corresponding random walk on the torus

$$\bar{\mathbf{U}}_n = \left\{ \sum_{i=1}^n X_i \right\}$$

where the  $X_i$  are independent and identically distributed with law  $\mu$  and  $\{\mathbf{x}\} = (\{x_{[k]}\}) \in [0, 1]^d$  is the fractional part in  $\mathbb{R}^d$ . As before, let  $\omega$  be the tuple of vectors generating the support of  $\mu$  in (2), with incommensurable coordinates.

It is clear that if  $\omega$ 's coordinates are well approximable by rationals, the same goes for the increments of the random walk, hence it is likely to come back closer to 0 faster. The study of random walks on a group started on finite arithmetic groups with the works of Diaconis, Saloff-Coste, Rosenthal, Porod, (see references in [31]) and results for such irrational random walks in the continuous settings were then achieved by Diaconis [10], and finally Su [31], who gave the optimal speed of convergence of the law of  $\bar{\mathbf{U}}_n$  in an appropriate distance. Then Prescott and Su [28] extended the study in higher dimensional tori.

The novelty of our approach is to consider estimates as  $\varepsilon \rightarrow 0$  uniformly in  $n$ ; we show in Section 3 that for a given  $\varepsilon$ , irrelevant of the number of steps  $n$ , there is a probability

always smaller than  $c_+ \varepsilon^{\frac{m}{m+\eta}}$  that the walk on the torus ends up in  $B_d(0, \varepsilon)$  after  $n$  steps, where  $\eta$  is such that  $\omega$  is  $(m + \eta)$ -BA and  $c_+ < \infty$  does not depend on  $n$  or  $\varepsilon$ .

This value is actually very sensitive to the probability of vanishing coordinates  $\overline{\mathbf{U}}_{n,[k]}$  of  $\overline{\mathbf{U}}_n$ , in the sense that it decays slowly in  $\varepsilon$  because of the fast recurrence to 0 on the axes: for  $p < d$

$$\mathbb{P}(\overline{\mathbf{U}}_{n,[1]} = \overline{\mathbf{U}}_{n,[2]} = \dots = \overline{\mathbf{U}}_{n,[p]} = 0) \sim n^{-p/2}.$$

We don't give a formal proof but one can sense this result by noticing that the probability that the elementary random walk on  $\mathbb{Z}$  with  $\pm 1$  increments has a probability  $\sim n^{-1/2}$  to come back to 0 in  $2n$  steps, and the components are *almost* independent up to the parity relation  $n \equiv \sum_{k=1}^d \mathbf{U}_{n,[k]}$  (see Lemma 2). Hence only non-vanishing coordinates matter in the speed of decay as  $\varepsilon \rightarrow 0$ . Denote by  $\llbracket d \rrbracket$  the set  $\{1, 2, \dots, d\}$ . Define for  $K \subset \llbracket d \rrbracket$ ,  $K \neq \emptyset$ , the restricted ball  $B_K(\varepsilon) := B_d(0, \varepsilon) \cap H_K$  where

$$H_K := \{\mathbf{y} = (y_{[k]})_{1 \leq k \leq d} \in \mathbb{R}^d : y_{[k]} \neq 0, k \in K \text{ and } y_{[k]} = 0, k \notin K\}.$$

Then we have according to Theorem 5 for some  $c < \infty$ , uniformly on  $n, \varepsilon$ ,

$$\overline{p}_n^K(\varepsilon) \leq cn^{-\frac{(d-|K|)m}{2}} \varepsilon^{\frac{|K|m}{m+\eta}} \text{ where } \overline{p}_n^K(\varepsilon) = \mathbb{P}(\overline{\mathbf{U}}_n \in B_K(\varepsilon)).$$

Regarding the dependance in  $\varepsilon$ , the random walk hence comes back to 0 faster on subspaces with fewer coordinates equal to 0 (the dependence as  $n$  increases is opposite). The most interesting part of the convergence, i.e. where the magnitude is not dominated by coordinates equal to 0, seems to happen on the domain  $H_{\llbracket d \rrbracket}$  of points with non-vanishing coordinates. More precise results are derived in (7), in Section 3, dedicated to irrational random walks; the results are derived in particular in terms of the optimal function  $\psi$  such that  $\omega$  is  $\psi$ -BA.

Lower bounds are more erratic and difficult to prove, hence we consider the smoothed estimate, for  $\beta > 2$ ,

$$\mathbf{I}_\beta(\varepsilon) = \sum_{n \geq n_\varepsilon} n^{-\beta/2} \mathbb{P}(0 < |\overline{\mathbf{U}}_n| < \varepsilon) \quad (4)$$

where  $n_\varepsilon \geq 1$  grows sufficiently slowly. We prove also in Theorem 5 that  $\mathbf{I}_\beta(\varepsilon)$  behaves in  $\varepsilon^{\frac{\beta-2+dm}{m+\eta}}$  as  $\varepsilon \rightarrow 0$ , regardless of the number of vanishing coordinates.

In Section 3, similar results (but with different magnitudes in  $n$ ) are actually derived first for the random walk  $\mathbf{U}_n = \sum_{i=1}^n X_i$  itself, and passed on to  $\overline{\mathbf{U}}_n$  to yield the aforementioned results. A quantity  $\mathbf{J}_\beta(\varepsilon)$  related to  $\mathbf{U}_n$  and analogue to  $\mathbf{I}_\beta(\varepsilon)$  is estimated and used in Section 4 to determine the variance of the excursion of the Gaussian field which spectral measure is  $\mu$ , as discussed previously.

## 2 Diophantine approximation

The core of the paper is provided by results from diophantine approximation, we explain here basic principles and results, as well as the more advanced ones we will need. The definitions of  $\psi$ -BA,  $\psi$ -WA<sup>(\*)</sup>,  $\psi$ -SWA are given in the introduction. The quality of the approximation of a tuple  $\omega \in \mathbb{R}^m$  is measured by the numbers

$$d_q(\omega) = \inf_{p \in \mathbb{Z}} |p - q \cdot \omega|, q \in \mathbb{Z}^m.$$

The most basic, yet useful result is the Dirichlet principle:

**Proposition 4.** *Let  $m \geq 1$ . There is  $c_m < \infty$  such that for  $N \in \mathbb{N}^*$ , one can find  $q, q' \in B_N := (\mathbb{Z} \cap [-N, N])^m$  distinct such that for  $\omega \in \mathbb{R}^m$ ,*

$$d_{q-q'}(\omega) \leq N^{-m} \leq c_m \|q - q'\|^{-m},$$

*which yields that  $\omega$  is  $(m)$ -WA and if  $\omega$  is  $(m + \eta)$ -BA, then necessarily  $\eta \geq 0$ .*

*Proof.* Simply remark that if one divides  $[0, 1]$  in  $M := |B_N| - 1$  bins of size  $M^{-1}$ , out of the  $|B_N|$  values  $d_q(\omega), q \in B_N$ , at least two of them will end up in the same bin, yielding for some  $q, q' \in B_N$  distinct

$$d_{q-q'}(\omega) \leq |d_q(\omega) - d_{q'}(\omega)| \leq M^{-1} \leq N^{-m}.$$

The second inequality comes from  $\|q - q'\| \leq 2\sqrt{m}N \leq \sqrt{m}2^{1-m}M^{1/m}$ . □

Another fundamental but more technical result is the Khintchine-Groshev theorem, we do not include the proof here, see the latest improvement by Hussain and Yusupova [15].

**Theorem 4** (Khintchine-Groshev). *Let  $\psi : \mathbb{N} \rightarrow \mathbb{R}_+$  tending to 0 such that*

$$\sum_{q \in \mathbb{Z}^m} \psi(|q|)^d < \infty.$$

*Then the set of  $\omega \in (\mathbb{R}^m)^d$  that are  $\psi$ -SWA is  $\mathcal{L}^{md}$ -negligible. If on the other hand the sum diverges then  $\mathcal{L}^{md}$ -a.a.  $\omega \in (\mathbb{R}^m)^d$  is  $\psi$ -SWA, in the case  $m = d = 1$   $\psi$  needs furthermore to be monotonic.*

The theorem yields that  $\mathcal{L}^{md}$ -a.a.  $\omega$  has irrationality index  $\tau(\omega) = m/d$ , where the irrationality index of some  $\omega \in (\mathbb{R}^m)^d$  is defined by

$$\tau(\omega) := \inf\{\tau : \omega \text{ is not } \tau\text{-SWA}\} = \sup\{\tau : \omega \text{ is } \tau\text{-SWA}\}.$$

The following result can be retrieved from Dirichlet's principle and Groshev's theorem with  $d = 1$ . The fourth point goes back to [16], who built explicit examples based on the continued fraction expansion of a real number.

**Corollary 2.** *1. All  $\omega \in \mathbb{R}^m$  is  $(m)$ -WA*

*2.  $\mathcal{L}^m$ -a.a.  $\omega \in \mathbb{R}^m$  is not  $(m)$ -BA.*

*3. For  $\eta > 0$ ,  $\mathcal{L}^m$ -a.a.  $\omega$  is not  $(m + \eta)$ -WA and is  $(m + \eta)$ -BA*

*4. For  $m = 1$  and  $\psi$  regular, the set of  $\omega \in \mathbb{R}$  that are  $\psi$ -WA and  $\psi$ -BA has positive Hausdorff dimension and is in particular uncountable.*

This third point implies that for  $\mathcal{L}^m$ -a.a.  $\omega \in \mathbb{R}^m$ ,  $\omega$  is  $(m + \eta)$ -BA for each  $\eta > 0$ . The following result yields that the situation is essentially the same if WA is replaced by  $\text{WA}^{(*)}$ . Recall that for  $p \geq 1$  and  $\mathbf{x} = (x_i)_{1 \leq i \leq p} \in \mathbb{Z}^p$ ,  $\mathbf{x} \equiv 1$  means that  $\sum_{i=1}^p x_i$  is an odd number (otherwise  $\mathbf{x} \equiv 0$ ).

**Proposition 5.** *If  $\omega$  is  $\psi$ -WA and not  $\psi\text{-WA}^{(*)}$  for  $\psi$  non-increasing, then for some  $1 \leq i \leq m$ ,  $\omega - \mathbf{e}_i$  is  $\psi\text{-WA}^{(*)}$ .*

*Proof.* Since  $\omega$  is  $\psi$ -WA and not  $\psi$ -WA<sup>(\*)</sup>, there are infinitely many  $p^j \in \mathbb{Z}, q^j \in \mathbb{Z}^m$  such that  $(p^j, q^j) \equiv 0$  and  $|p^j - \omega \cdot q^j| < c_W \psi(|q^j|)$ . Let  $k_j \in \mathbb{N}$  be the largest natural integer such that  $2^{k_j}$  divides  $p^j$  and each  $q_i^j, 1 \leq i \leq m$ , let  $\tilde{p}^j = 2^{-k_j} p^j, \tilde{q}^j = 2^{-k_j} q^j$ . We have

$$|\tilde{p}^j - \omega \cdot \tilde{q}^j| < 2^{-k_j} c_W \psi(|q^j|) = 2^{-k_j} c_W \psi(2^{k_j} |\tilde{q}^j|) < c_W \psi(|\tilde{q}^j|)$$

hence either  $(\tilde{p}^j, \tilde{q}^j) \equiv 1$  for infinitely many  $j$  and the result is proved, or for infinitely many  $j$ ,  $(\tilde{p}^j, \tilde{q}^j) \equiv 0$ . In the latter case, since  $2^1$  does not divide each  $\tilde{q}_i^j$  and  $\tilde{p}^j$ , but  $(\tilde{p}^j, \tilde{q}^j) \equiv 0$ , necessarily for each  $j$ ,  $\tilde{q}_i^j \equiv 1$  for at least one  $i$ . By the drawer principle, for at least one  $i$  there is an infinity of  $j$  such that  $\tilde{q}_i^j$  is odd, denote it by  $i^*$ . Let then  $\omega' = \omega + \mathbf{e}_{i^*}, p'^j = \tilde{p}^j + \tilde{q}_{i^*}^j$ . We have

$$|p'^j - \omega' \cdot \tilde{q}^j| = |p'^j - \tilde{q}_{i^*}^j - \omega \cdot \tilde{q}^j| = |\tilde{p}^j - \omega \cdot \tilde{q}^j| < c_W \psi(|\tilde{q}^j|).$$

Since  $(p'^j, \tilde{q}^j) \equiv \tilde{q}_{i^*}^j \equiv 1$  for infinitely many  $j$ , the result is proved.  $\square$

The next result is useful for tensorizing variance estimates.

**Proposition 6.** *If  $\omega \in \mathbb{R}^m$  is  $\psi$ -WA<sup>(\*)</sup>,  $\boldsymbol{\omega} = (\omega, \dots, \omega) \in (\mathbb{R}^m)^d$  is  $\psi$ -SWA<sup>(\*)</sup>.*

*Proof.* Since  $\omega$  is assumed to be  $\psi$ -WA<sup>(\*)</sup>, there is a sequence  $(p^j, q^j)_j$  such that  $|p^j - \omega \cdot q^j| < c_W \psi(|q^j|)$  and  $(p^j, q^j) \equiv 1$ . Hence with  $q_{[k]}^j = q^j, p_{[k]}^j = p^j$ ,

$$|p_{[k]}^j - q_{[k]}^j \cdot \omega| = |p^j - q^j \cdot \omega| < c_W \psi(|q^j|)$$

but  $\sum_{k=1}^d (p_{[k]}^j + \sum_{i=1}^m q_{[k],i}^j) = d(p^j + \sum_i q_i^j)$  is odd only if  $d$  is odd. If  $d$  is even, choose instead  $p_{[1]}^j = 2p^j, q_{[1]}^j = 2q^j$ , so that  $|p_{[1]}^j - q_{[1]}^j \cdot \omega| < 2c_W \psi(|q^j|)$ , and  $\sum_{k=1}^d (p_{[k]}^j + \sum_i q_{[k],i}^j) = (2d+1)(p^j + q^j)$  is indeed odd. This sequence demonstrates that  $\boldsymbol{\omega}$  is  $\psi$ -SWA<sup>(\*)</sup>.  $\square$

### 3 Irrational random walks

Given  $\boldsymbol{\mu} = \sum_{i=1}^M \theta_i \bar{\delta}_{\boldsymbol{\omega}_i}$  a finite atomic symmetric probability measure on  $\mathbb{R}^d$ , we are interested in the associated random walk

$$\mathbf{U}_n := \sum_{k=1}^n X_k$$

where the  $X_k$  are independent and identically distributed with law  $\boldsymbol{\mu}$ , hence centred, and the  $\theta_i$  are in  $[0, \frac{1}{2}]$  and sum to 1 ( $\frac{1}{2}$  could be replaced by any element of  $(0, 1)$ , as long as it is fixed throughout the paper). The study of  $\mathbf{U}_n$  is related to the random walk on the torus

$$\bar{\mathbf{U}}_n = \mathbf{U}_n - [\mathbf{U}_n] \in [0, 1]^d,$$

which has been intensively studied, the consequences of the current results to the random walk on the torus are discussed at Section 1.6. To avoid degenerate behaviour, we assume that  $\boldsymbol{\mu}$  is  $\mathbb{Z}$ -free, i.e. there is no  $\mathbf{q} \in \mathbb{Z}^M \setminus \{0\}$  such that  $\sum_{i=1}^M \mathbf{q}_i \boldsymbol{\omega}_i = 0$ . In general we further assume that  $\boldsymbol{\omega} := (\boldsymbol{\omega}_i)$  is  $\psi$ -BA for some non-vanishing function  $\psi$ , which automatically implies that  $\boldsymbol{\mu}$  is  $\mathbb{Z}$ -free.

According to the Central Limit Theorem, the law of the renormalised sum  $n^{-1/2}\mathbf{U}_n$  weakly converges to a Gaussian measure (see also Lemma 2 for precise estimates), and the law  $\overline{\mu}_n$  of  $\overline{\mathbf{U}}_n$  is known to converge to Lebesgue measure on  $[0, 1]^d$  [31]. But it seems that if we zoom in further on this convergence, it becomes very irregular.

We estimate the following quantities below:

$$p_n^{\mathbf{x}}(\varepsilon) = \mathbb{P}(0 < |\mathbf{U}_n - \mathbf{x}| \leq \varepsilon), \varepsilon > 0, \mathbf{x} \in \mathbb{Z}^d,$$

$$\bar{p}_n(\varepsilon) = \sum_{\mathbf{x} \in \mathbb{Z}^d} p_n^{\mathbf{x}}(\varepsilon) = \mathbb{P}(0 < |\overline{\mathbf{U}}_n| \leq \varepsilon).$$

Remark that in general, if the sum  $n + \sum_{k=1}^d \mathbf{x}_{[k]}$  is even,  $\mathbb{P}(\mathbf{U}_n = \mathbf{x})$  is in  $n^{-\frac{M}{2}}$  and hence dominates  $p_n^{\mathbf{x}}(\varepsilon)$  for  $\varepsilon \rightarrow 0$ , which is why it is estimated separately. For odd values, since  $\mu$  is  $\mathbb{Z}$ -free,  $\mathbb{P}(\mathbf{U}_n = \mathbf{x}) = 0$ , hence  $p_n^{\mathbf{x}}(\varepsilon)$  is simply  $\mathbb{P}(|\mathbf{U}_n - \mathbf{x}| \leq \varepsilon)$ .

A fine analysis of the recurrence around 0 yields that the rate strongly depends on the number of coordinates equal to 0, expressed through

$$p_n^{K, \mathbf{x}}(\varepsilon) = \mathbb{P}(\mathbf{U}_n - \mathbf{x} \in B_K(0, \varepsilon)), \bar{p}_n^K(\varepsilon) = \mathbb{P}(\overline{\mathbf{U}}_n \in B_K(0, \varepsilon)).$$

We show below for instance that for some  $c < \infty$ , for  $K \subset \llbracket d \rrbracket$ , for  $\omega$  that is  $(m+\eta)$ -BA,  $\eta \geq 0$ ,

$$\bar{p}_n^K(\varepsilon) \leq cn^{-m \frac{d-|K|}{2}} \varepsilon^{\frac{m|K|}{m+\eta}}$$

so that it is really the number of vanishing coordinates that determines the recurrence probabilities. To avoid this technicality and obtain lower bounds, we consider the smoothed estimates for  $\beta > 0$ , with  $p_n^{\mathbf{0}} = p_n^{\llbracket d \rrbracket, \mathbf{0}}$

$$\mathbf{J}_\beta(\varepsilon) := \sum_{n \geq n_\varepsilon, n \in \mathbb{N}, n \text{ odd}} n^{-\beta/2} p_n^{\mathbf{0}}(\varepsilon)$$

analogue to  $\mathbf{I}_\beta(\varepsilon)$  defined in the introduction at (4), where  $n_\varepsilon$  does not grow too fast as  $\varepsilon \rightarrow 0$  and serves the purpose to show that it is the series tail that actually matters. Considering this statistic also allows to suppress the erratic behaviour in  $n$ , and we can prove that  $\mathbf{I}_\beta(\varepsilon), \mathbf{J}_\beta(\varepsilon)$  both behave in  $\varepsilon^{\frac{m}{m+\eta}}$  and find a matching lower bound. The summation over odd  $n$  in  $\mathbf{J}_\beta(\varepsilon)$  is adapted to estimating the volume variance of Gaussian nodal excursions (see Remark 7).

Assume there is  $m \in \mathbb{N}$ , and real numbers  $\omega_{[k], i}, 1 \leq k \leq d, 0 \leq i \leq m$  such that  $\mu$  is of the form (2) and denote by  $\omega_{[k]} = (\omega_{[k], 0}, \dots, \omega_{[k], m}), \boldsymbol{\omega} = (\omega_{[k]})_{1 \leq k \leq d}$ . For  $\psi : \mathbb{N}^* \rightarrow [0, 1]$  that converges to 0 let  $\psi^{-1}$  its pseudo-inverse defined by

$$\psi^{-1}(\varepsilon) = \min\{q \in \mathbb{N}^* : \psi(q) \leq \varepsilon\}, \varepsilon > 0. \quad (5)$$

**Theorem 5.** *Assume that  $\mu$  is under the form (2) for some  $\boldsymbol{\omega} = (\omega_{[k]})_{1 \leq k \leq d} \in (\mathbb{R}^m)^d$ . Let  $\psi$  be some mapping  $\mathbb{N}^* \rightarrow (0, 1]$  converging to 0. Let  $\beta > 0$ . There is  $0 < c < \infty$  depending on  $d, m, \psi, \beta$  such that the following holds:*

(i) *Assume each  $\omega_{[k]}$  is  $\psi$ -BA. We have for  $\mathbf{x} \in \mathbb{Z}^d, K \subset \llbracket d \rrbracket, 0 < \varepsilon < \frac{1}{2}, n \in \mathbb{N}^*$*

$$p_n^{K, \mathbf{x}}(\varepsilon) \leq cn^{-d/2} n^{-\frac{(d-|K|)m}{2}} \psi^{-1}(\varepsilon)^{-m|K|} \exp(-cn^{-1} \|\mathbf{x}\|^2) \quad (6)$$

$$\bar{p}_n^K(\varepsilon) \leq cn^{-\frac{(d-|K|)m}{2}} \psi^{-1}(\varepsilon)^{-m|K|} \quad (7)$$

$$\mathbf{I}_\beta(\varepsilon) \leq c\psi^{-1}(\varepsilon)^{-\beta-dm+2} \quad (8)$$

$$\mathbf{J}_\beta(\varepsilon) \leq c\psi^{-1}(\varepsilon)^{-\beta-d(m+1)+2} \quad (9)$$

(ii) Assume  $\omega$  is  $\psi$ -SWA. Then if  $n_\varepsilon \leq \psi^{-1}(\varepsilon)^2, \varepsilon > 0$ ,

$$\psi^{-1}(\varepsilon)^{-\beta-dm+2} \stackrel{\text{i.o.}}{\leq} c\mathbf{I}_\beta(\varepsilon), \varepsilon \rightarrow 0. \quad (10)$$

(iii) Assume  $\omega$  is  $\psi$ -SWA $^{(*)}$ . Then if  $n_\varepsilon \leq \psi^{-1}(\varepsilon)^2, \varepsilon > 0$ ,

$$\psi^{-1}(\varepsilon)^{-\beta-d(m+1)+2} \stackrel{\text{i.o.}}{\leq} c\mathbf{J}_\beta(\varepsilon), \varepsilon \rightarrow 0 \quad (11)$$

**Remark 7.** The reason why  $\omega$  has to be  $\psi$ -SWA $^{(*)}$  instead of  $\psi$ -SWA at point (iii) is because summands are odd in the definition of  $\mathbf{J}_\beta$ . As proved at Proposition 5, this subtlety does not influence final results about Gaussian excursions, hence one would like a general result from diophantine approximation that states that  $\psi$ -SWA tuples are also  $\psi$ -SWA $^{(*)}$ , but that is unlikely.

### 3.1 Proof of Theorem 5

**Notation.** We introduce the notation  $A = \Theta B$  to indicate that there are finite constants  $c, c' > 0$  depending on  $\mu, m, d, \psi$  and not (further) on  $\omega, \varepsilon, T, n$  such that  $A \leq cB, B \leq c'A$ .

Also, for a  $d$ -tuple of vectors of  $\mathbb{R}^{m+1}$ ,  $\bar{\mathbf{x}} = (\bar{x}_{[1]}, \dots, \bar{x}_{[d]}) \in (\mathbb{R}^{m+1})^d$  with  $\bar{x}_{[k]} = (x_{[k],0}, \dots, x_{[k],m}) \in \mathbb{R}^{m+1}$ , remove the bar when the 0-th component is removed from each vector:

$$x_{[k]} = (x_{[k],1}, \dots, x_{[k],m}), \quad \mathbf{x} = (x_{[1]} \dots, x_{[d]}).$$

Euclidean norms in  $\mathbb{R}^m$  are denoted by a single bar and in  $(\mathbb{R}^m)^d$  by two bars:

$$|x_{[k]}|^2 = \sum_{i=1}^m x_{i,[k]}^2, \quad \|\mathbf{x}\|^2 = \sum_{k=1}^d |x_{[k]}|^2.$$

**Lemma 1.** Let  $\omega \in \mathbb{R}^m$  that is  $\psi$ -BA. For  $1/2 \geq \varepsilon > 0$ , define

$$I_\varepsilon(\omega) := \{q \in \mathbb{Z}^m \setminus \{0\} : 0 < d_q(\omega) \leq \varepsilon\}.$$

Let  $q^{(N)}, N \in \mathbb{N}^*$ , the elements of  $I_\varepsilon$  ordered by increasing radius. Then

$$|q^{(N)}| \geq \Theta N^{\frac{1}{m}} \psi^{-1}(\varepsilon). \quad (12)$$

In particular, we have the following estimate:

$$\sum_{q \in I_\varepsilon} \exp(-\Theta n^{-1}|q|^2) \leq \sum_{N=1}^{\infty} \exp(-\Theta n^{-1} N^{\frac{2}{m}} \psi^{-1}(\varepsilon)^2) \leq \Theta n^{\frac{m}{2}} \psi^{-1}(\varepsilon)^{-m}. \quad (13)$$

*Proof.* The starting point is that for  $q \in I_\varepsilon$ , since  $\varepsilon > \varepsilon/2 \geq d_q(\omega)/2 \geq \psi(|q|)$ , we have  $|q| \geq \rho := \psi^{-1}(\varepsilon)$ . And the triangular inequality yields for  $q \neq q' \in I_\varepsilon$ ,

$$2\psi(q - q') \leq d_{q-q'}(\omega) \leq 2\varepsilon,$$

hence  $|q - q'| \geq \rho$  as well. It follows that all  $q \in I_\varepsilon$  are pairwise distant by more than  $\rho$ , and the balls  $B_m(q, \rho/2), q \in I_\varepsilon$  are disjoint. Hence for  $N_0 \in \mathbb{N}^*$ , the total  $\mathcal{L}^m$ -measure occupied by the  $B_m(q^{(N)}, \rho/2), N \leq N_0$  is larger than  $\Theta N_0 \rho^m$ . This volume is necessarily smaller than the volume of the ball with radius  $|q^{(N_0)}| + \rho/2 \leq 2|q^{(N_0)}|$ , hence

$$\Theta N_0 \rho^m \leq \Theta |q^{(N_0)}|^m$$

which yields (12). Finally (13) follows from

$$\sum_{N=1}^{\infty} \exp(-\Theta n^{-1} (N^{\frac{1}{m}} \psi^{-1}(\varepsilon))^2) \leq 2 \int_{1/2}^{\infty} \exp(-\Theta (n^{-\frac{m}{2}} \psi^{-1}(\varepsilon)^m y)^{\frac{2}{m}}) dy \leq \Theta n^{\frac{m}{2}} \psi^{-1}(\varepsilon)^{-m}.$$

□

*Proof of Theorem 5.* Let  $M = d(m+1)$ . The proof is based on the study of the symmetric random walk  $(S_n)_n$  on  $\mathbb{Z}^M$  with independent increments defined by  $S_0 = 0$  and

$$\mathbb{P}(S_{n+1} = S_n \pm \mathbf{e}_j) = \frac{1}{2M}, 1 \leq j \leq M,$$

where  $(\mathbf{e}_j)_j$  is the canonical basis. Following the notation introduced above, denote also  $\bar{\omega}_{[k]} = (1, \omega_{[k]})$  and  $\bar{\omega} = (\bar{\omega}_{[k]})_k$ .

For  $\bar{q}_{[k]} = (q_{[k],0}, q_{[k]}) \in \mathbb{Z}^{m+1}$ ,  $1 \leq k \leq d$ ,  $\bar{\mathbf{q}} = (\bar{q}_{[1]}, \dots, \bar{q}_{[d]}) \in (\mathbb{Z}^{m+1})^d \approx \mathbb{Z}^M$ , denote by  $q_{[k]} = (q_{[k],1}, \dots, q_{[k],m}) \in \mathbb{Z}^m$ ,  $\mathbf{q} = (q_{[k]})_k \in (\mathbb{Z}^m)^d$ . We define

$$\bar{\mathbf{q}} \otimes \bar{\omega} := (\bar{q}_{[k]} \cdot \bar{\omega}_{[k]})_{k=1, \dots, d} = \mathbf{q}_0 + \mathbf{q} \otimes \boldsymbol{\omega} \in \mathbb{R}^d$$

where  $\mathbf{q}_0 = (q_{[k],0})_{k=1}^d$ , so that we have the representation  $\mathbf{U}_n = S_n \otimes \bar{\omega}$ .

We use the notation, for  $\mathbf{x} = (x_{[1]}, \dots, x_{[d]}) \in \mathbb{Z}^d$ ,  $K \subset \llbracket d \rrbracket$ ,

$$\bar{\mathbf{I}}_{\varepsilon}^{\mathbf{x}, K} = \{\bar{\mathbf{q}} \in \mathbb{Z}^M : \bar{q}_{[k]} = 0, k \notin K \text{ and } 0 < |\bar{q}_{[k]} \cdot \bar{\omega}_{[k]} - x_{[k]}| \leq \varepsilon, k \in K\}.$$

For  $\varepsilon < 1/2$ , an element  $\bar{\mathbf{q}} \in \bar{\mathbf{I}}_{\varepsilon}^{\mathbf{x}, K}$  satisfies the following for  $1 \leq k \leq d$ :

$$|q_{[k],0} + q_{[k]} \cdot \omega_{[k]} - x_{[k]}| < \varepsilon,$$

hence since  $x_{[k]} \in \mathbb{Z}$  and  $q_{[k]} \in I_{\varepsilon}(\omega_{[k]})$ ,  $d_{q_{[k]}}(\omega_{[k]}) \leq \varepsilon$ . It follows that  $q_{[k],0}$  depends explicitly on other coordinates

$$\begin{aligned} q_{[k],0} = q_{[k],0}(x_{[k]}, q_{[k]}) &:= \operatorname{argmin}_{p \in \mathbb{Z}} |p + q_{[k]} \cdot \omega_{[k]} - x_{[k]}| \\ \mathbf{q}_0 = \mathbf{q}_0(\mathbf{x}, \mathbf{q}) &:= (q_{[k],0})_k. \end{aligned} \quad (14)$$

In particular,  $|x_{[k]}| \leq |q_{[k],0}| + |q_{[k]} \cdot \omega_{[k]}| + 1$ , and

$$\|\bar{\mathbf{q}}\|^2 = \|\mathbf{q}_0\|^2 + \|\mathbf{q}\|^2 \geq \max(\|\mathbf{q}\|^2, \|\mathbf{q}\|^2 + \Theta(\|\mathbf{x}\|^2 - 1)) \geq \Theta(\|\mathbf{q}\|^2 + \|\mathbf{x}\|^2). \quad (15)$$

We also have the one-to-one correspondance

$$\mathbf{I}_{\varepsilon}^{\mathbf{x}, K}(\boldsymbol{\omega}) := \left\{ \mathbf{q} \in \prod_k \mathbb{Z}^m : (\mathbf{q}_0(\mathbf{x}, \mathbf{q}), \mathbf{q}) \in \bar{\mathbf{I}}_{\varepsilon}^{\mathbf{x}, K}(\boldsymbol{\omega}), k \in K \right\} \equiv \{\mathbf{0}\}^{d-|K|} \times \prod_{k \in K} I_{\varepsilon}(\omega_{[k]}). \quad (16)$$



**Proof of (i):** By the Gaussian approximation Lemma 2 (below), and (15),

$$\begin{aligned}
p_n^{\mathbf{x},K}(\varepsilon) &= \sum_{\bar{\mathbf{q}} \in \bar{\mathbf{I}}_\varepsilon^{\mathbf{x},K}} \mathbb{P}(S_n = \bar{\mathbf{q}}) = \sum_{\mathbf{q} \in \mathbf{I}_\varepsilon^{\mathbf{x},K}} \mathbb{P}(S_n = (\mathbf{q}_0(\mathbf{x}, \mathbf{q}), \mathbf{q})) \\
&\leq \Theta \sum_{\mathbf{q} \in \mathbf{I}_\varepsilon^{K,\mathbf{x}}} n^{-\frac{M}{2}} \exp(-\Theta n^{-1}(\|\mathbf{q}\|^2 + \|\mathbf{q}_0\|^2)) \\
&\leq \Theta n^{-\frac{M}{2}} \sum_{\mathbf{q} \in \mathbf{I}_\varepsilon^{K,\mathbf{x}}} \exp(-\Theta n^{-1}\|\mathbf{q}\|^2) \exp(-\Theta n^{-1}\|\mathbf{x}\|^2) \\
&\leq \Theta n^{-\frac{M}{2}} \exp(-\Theta n^{-1}\|\mathbf{x}\|^2) \prod_{k \in K} \sum_{q_{[k]} \in I_\varepsilon(\omega_{[k]})} \exp(-\Theta n^{-1}|q_{[k]}|^2) \text{ by (16)} \quad (17) \\
&\leq \Theta n^{-\frac{d(m+1)}{2}} \exp(-\Theta n^{-1}\|\mathbf{x}\|^2) (n^{\frac{m}{2}} \psi^{-1}(\varepsilon)^{-m})^{|K|} \text{ with (13),} \\
&\leq \Theta n^{-d/2} n^{-\frac{(d-K)m}{2}} \psi^{-1}(\varepsilon)^{-m|K|} \exp(-\Theta n^{-1}\|\mathbf{x}\|^2)
\end{aligned}$$

and (6) is proved.

The bound (7) immediately stems from  $\bar{p}_n^K = \sum_{\mathbf{x} \in \mathbb{Z}^d} p_n^{\mathbf{x},K}$  and Lemma 3 (after summing over  $i \in \{0, 1\}$ ). Hence using (13), and (17) with  $\mathbf{x} = 0$

$$\begin{aligned}
\mathbf{I}_\beta^0(\varepsilon) &:= \sum_{n \geq n_\varepsilon} n^{-\beta/2} p_n^0(\varepsilon) \\
&= \sum_{n \geq n_\varepsilon} n^{-\beta/2} \sum_{K \neq \emptyset} p_n^{0,K}(\varepsilon) \\
&\leq \Theta \mathbf{K}_\beta(\varepsilon)
\end{aligned}$$

$$\begin{aligned}
\text{with } \mathbf{K}_\beta(\varepsilon) &:= \sum_{n \geq n_\varepsilon} n^{-\beta/2 - \frac{M}{2}} \sum_{K \subset [d], K \neq \emptyset} \prod_{k \in K} \sum_{N_k=1}^{\infty} \exp(-\Theta n^{-1} (N_k^{\frac{1}{m}} \psi^{-1}(\varepsilon))^2) \\
&\leq \sum_{n \geq n_\varepsilon} n^{-\beta/2 - \frac{M}{2}} \sum_{K \subset [d], K \neq \emptyset} \sum_{N_k \geq 1, k \in K}^{\infty} \exp(-\Theta n^{-1} \sum_{k \in K} (N_k^{\frac{1}{m}} \psi^{-1}(\varepsilon))^2) \\
&\leq \Theta \sum_{K \subset [d], K \neq \emptyset} \sum_{N_k \geq 1, k \in K} \sum_{n \geq n_\varepsilon} \int_n^{n+1/2} (z - 1/2)^{-\beta/2 - \frac{M}{2}} \exp(-\Theta z^{-1} \sum_{k \in K} \psi^{-1}(\varepsilon)^2 N_k^{2/m}) dz \\
&\leq \Theta \sum_{K \subset [d], K \neq \emptyset} \sum_{N_k \geq 1, k \in K} \left( \sum_{k \in K} \psi^{-1}(\varepsilon)^2 N_k^{2/m} \right)^{1-\beta/2 - \frac{M}{2}} \int_0^\infty y^{\beta/2 + \frac{M}{2} - 2} \exp(-\Theta y) dy \\
&\leq \Theta (\psi^{-1}(\varepsilon)^2)^{1-\beta/2 - \frac{M}{2}} \max_{K \subset [d], K \neq \emptyset} \int_{[1, \infty]^{|K|}} \left( \sum_{k \in K} x_k^{\frac{2}{m}} \right)^{1-\beta/2 - \frac{M}{2}} \prod_{k \in K} dx_k \\
&\leq \Theta \psi^{-1}(\varepsilon)^{2-M-\beta} \max_{1 \leq p \leq d} \int_{[1, \infty]^p} \left( \sum_{k=1}^p y_k \right)^{1-\beta/2 - \frac{M}{2}} \prod_{k=1}^p y_k^{\frac{m}{2} - 1} dy_k \\
&\leq \Theta \psi^{-1}(\varepsilon)^{2-d(m+1)-\beta} \max_{1 \leq p \leq d} \int_1^\infty (\Theta r)^{1-\beta/2 - \frac{(m+1)d}{2}} r^{mp/2 - p} r^{p-1} dr
\end{aligned}$$

and the integral converges if  $\beta/2 > 1 - d/2$ . Since there are less terms in  $\mathbf{J}_\beta(\varepsilon)$ , than in  $\mathbf{I}_\beta^0(\varepsilon)$ , the upper bound holds and (9) is proved.

With the same computations, using first (17), and then(13), and Lemma 3,

$$\begin{aligned} \mathbf{I}_\beta(\varepsilon) &= \sum_{n \geq n_\varepsilon} n^{-\beta/2} \sum_{\mathbf{x} \in \mathbb{Z}^d} \sum_{K \neq \emptyset} p_n^{\mathbf{x}, K}(\varepsilon) \\ &\leq \sum_{n \geq n_\varepsilon} n^{-\beta/2 - \frac{M}{2}} \sum_{\mathbf{x} \in \mathbb{Z}^d} \exp(-\Theta n^{-1} \mathbf{x}^2) \sum_{K \subset \llbracket d \rrbracket, K \neq \emptyset} \prod_{k \in K} \sum_{N_k=1}^{\infty} \exp(-\Theta n^{-1} (N_k^{\frac{1}{m}} \psi^{-1}(\varepsilon))^2) \\ &\leq \Theta \mathbf{K}_{\beta-d}(\varepsilon) \end{aligned}$$

provided  $\beta/2 > 1$ , which proves (8).

Let us conclude with the proof of (iii), the proof of (ii) is similar and easier. There are by hypothesis infinitely many  $q^j \in \mathbb{Z}^m, j \geq 1$  and  $p_{[k]}^j \in \mathbb{Z}, 1 \leq k \leq d$ , such that  $\bar{\mathbf{q}}^j := ((p_{[k]}^j, q^j))_k \equiv 1$  and

$$|p_{[k]}^j - \omega_{[k]} \cdot q^j| = d_{q^j}(\omega_{[k]}) \leq c_W \psi(|q^j|) =: c_W \varepsilon_j$$

(we have  $\varepsilon_j \rightarrow 0$  because  $\psi$  converges to 0 by hypothesis). We have in particular with Cauchy-Schwarz inequality

$$\|\bar{\mathbf{q}}^j\| \leq \sum_{k=1}^d (|p_{[k]}^j| + |q^j|) \leq \sum_{k=1}^d (|\omega_k| |q^j| + 1 + |q^j|) \leq \Theta |q^j|$$

and clearly the other inequality as well  $|q^j| \leq \|\bar{\mathbf{q}}^j\|$ .

Then, by Lemma 2, with  $\tilde{n}_j := c_{\inf}^{-1} \|\bar{\mathbf{q}}^j\| \vee n_{\varepsilon_j}$

$$\begin{aligned} \mathbf{J}_\beta(\varepsilon_j) &= \sum_{n \geq n_{\varepsilon_j}, n \text{ odd}} n^{-\beta/2} p_n^{\mathbf{0}}(\varepsilon_j) \geq \sum_{n \geq n_{\varepsilon_j}, n \text{ odd}} n^{-\beta/2} \mathbb{P}(S_n = \bar{\mathbf{q}}^j) \\ &\geq \Theta \sum_{n \geq \tilde{n}_j, n \equiv \bar{\mathbf{q}}^j \equiv 1} n^{-\beta/2} n^{-\frac{d(m+1)}{2}} \exp(-\Theta n^{-1} \|\bar{\mathbf{q}}^j\|^2) \\ &\geq \Theta \int_{[\tilde{n}_j/2]}^{\infty} y^{-\beta/2 - d\frac{m+1}{2}} \exp(-\Theta y^{-1} |q^j|^2) dy \\ &\geq \Theta |q^j|^{2-\beta-d(m+1)} \int_0^{\Theta |q^j|^2 \tilde{n}_j^{-1}} z^{\beta/2 + d\frac{m+1}{2} - 2} \exp(-\Theta z) dz \\ &\geq \Theta \psi^{-1}(\varepsilon_j)^{2-\beta-d(m+1)} \end{aligned}$$

provided  $\beta > 0$ , because  $|q^j|^2 |\bar{\mathbf{q}}^j|^{-1} \geq \Theta > 0$  and  $n_{\varepsilon_j} \leq \psi^{-1}(\varepsilon_j)^2$  yields (recalling  $\psi(|q^j|) = \varepsilon_j$ )

$$|q^j|^2 n_{\varepsilon_j}^{-1} \geq \psi^{-1}(\varepsilon_j)^2 \psi^{-1}(\varepsilon_j)^{-2} = 1,$$

hence (11) is proved. The proof of (10) is similar without the requirement that  $\bar{\mathbf{q}}^j \equiv 1$ , hence the sum is over all  $n \geq n_{\varepsilon_j}$  (even and odd).  $\square$

## 3.2 Gaussian approximation

The following lemma quantifies how much  $S_n$  is close to a Gaussian distribution.

**Lemma 2.** *Let  $p \in \mathbb{N}$  and  $S_n$  be the centred random walk on  $\mathbb{Z}^p$  with weights  $\theta_i \in (0, 1), 1 \leq i \leq p$ , summing to 1, i.e.*

$$\mathbb{P}(S_{n+1} = S_n \pm \mathbf{e}_i) = \frac{\theta_i}{2}, 1 \leq i \leq p, n \in \mathbb{N}.$$

For  $\mathbf{q} = (\mathbf{q}_i) \in \mathbb{Z}^p$ ,  $n \in \mathbb{N}$ , write  $\mathbf{q} \equiv n$  if  $\sum_{i=1}^p \mathbf{q}_i$  and  $n$  have the same parity, and remark that  $\mathbb{P}(S_n = \mathbf{q}) = 0$  if  $\mathbf{q} \not\equiv n$ . There is a constant  $c_{\text{inf}} > 0$  such that for  $\mathbf{q} \in \mathbb{Z}^p$

$$\mathbb{P}(S_n = \mathbf{q}) \leq \Theta n^{-\frac{p}{2}} \exp(-\Theta n^{-1} \|\mathbf{q}\|^2) \quad (18)$$

$$\mathbf{1}_{\{|\mathbf{q}| \leq c_{\text{inf}} n\}} \mathbb{P}(S_n = \mathbf{q}) \geq \Theta n^{-\frac{p}{2}} \exp(-\Theta n^{-1} \|\mathbf{q}\|^2) \mathbf{1}_{\{|\mathbf{q}| \leq c_{\text{inf}} n\}} \text{ for } \mathbf{q} \equiv n.$$

**Remark 8.** The constants involved in this result depend also on the  $\theta_i$ ,  $1 \leq i \leq p$ .

*Proof.* Let  $N_i$  be the number of times direction  $i$  has been chosen in the random walk, and let  $B_i \leq N_i$  be the number of  $+\mathbf{e}_i$  increments, hence  $N_i - B_i$  is the number of  $-\mathbf{e}_i$  increments. The  $i$ -th component of  $S_n$  is therefore  $S_{n,i} := N_i - 2B_i$ . We have  $N_i \sim \mathcal{B}(n, \theta_i)$ ,  $B_i \sim \mathcal{B}(N_i, 1/2)$ , and the  $B_i$  are independent conditionally on  $\mathbf{N} := (N_i)_i$ . Hence for  $|\varepsilon| \leq c_{\text{Bin}}$ , from Lemma 4

$$\mathbb{P}(B_i = \lfloor N_i(1/2 + \varepsilon) \rfloor \mid \mathbf{N}) = \Theta \exp(-\Theta N_i \varepsilon^2) N_i^{-1/2}.$$

Let  $\mathbf{q} = (\mathbf{q}_i) \in \mathbb{Z}^p$  such that for  $1 \leq i \leq p$ ,  $|\mathbf{q}_i| \leq c_{\text{Bin}} N_i$ , let  $\varepsilon_i = N_i^{-1} \mathbf{q}_i$ ,

$$\mathbb{P}(S_{n,i} = \mathbf{q}_i \mid \mathbf{N}) = \mathbb{P}(B_i = N_i/2 + \mathbf{q}_i/2 \mid \mathbf{N}) = \begin{cases} 0 & \text{if } N_i \not\equiv \mathbf{q}_i \\ \Theta N_i^{-1/2} \exp(-\Theta N_i \varepsilon_i^2) = \Theta N_i^{-1/2} \exp(-\Theta N_i^{-1} \mathbf{q}_i^2) & \text{otherwise.} \end{cases}$$

Let

$$c_{\text{inf}} := c_{\text{Bin}}(1 - c_{\text{Bin}}) \min_i \theta_i > 0.$$

If for all  $i$ ,  $N_i > \theta_i(1 - c_{\text{Bin}})n$  and  $|\mathbf{q}_i| < c_{\text{inf}} n$ , then  $|\mathbf{q}_i| < c_{\text{Bin}} N_i$  and we have the lower bound

$$\begin{aligned} \mathbb{P}(S_n = \mathbf{q}) &= \mathbb{E}(\mathbb{P}(S_n = \mathbf{q} \mid \mathbf{N})) \\ &= \mathbb{E}(\mathbf{1}_{\{\mathbf{q}_i \equiv N_i, \forall i\}} \mathbb{P}(S_n = \mathbf{q} \mid \mathbf{N})) \\ &\geq \mathbb{E}(\mathbf{1}_{\{\mathbf{q}_i \equiv N_i, N_i > \theta_i(1 - c_{\text{Bin}})n, \forall i\}} \mathbb{P}(S_n = \mathbf{q} \mid \mathbf{N})) \\ &\geq \mathbb{E}(\mathbf{1}_{\{\mathbf{q}_i \equiv N_i, N_i > \theta_i(1 - c_{\text{Bin}})n, \forall i\}} \Theta \prod_i N_i^{-\frac{1}{2}} \exp(-\Theta N_i^{-1} \mathbf{q}_i^2)) \\ &\geq \mathbb{E}(\mathbf{1}_{\{\mathbf{q}_i \equiv N_i, N_i > \theta_i(1 - c_{\text{Bin}})n, \forall i\}} \Theta n^{-\frac{p}{2}} \exp(-\Theta n^{-1} \mathbf{q}^2)) \\ &\geq \Theta n^{-\frac{p}{2}} \exp(-\Theta n^{-1} \mathbf{q}^2) \mathbb{P}(\mathbf{q}_i \equiv N_i, N_i > \theta_i(1 - c_{\text{Bin}})n, \forall i). \end{aligned}$$

If  $\mathbf{q} \equiv n$ , Asymptotically a fraction  $2^{-(p-1)}$  of tuples  $\mathbf{N} \in \llbracket n \rrbracket^p$  are such that  $\sum_i N_i \equiv n$  and  $N_i > \theta_i(1 - c_{\text{Bin}})n$  will be compatible with  $\mathbf{q}$ , hence  $\mathbb{P}(\mathbf{q}_i \equiv N_i, N_i > (\theta_i - c_{\text{Bin}})n, \forall i) = \Theta \mathbf{1}_{\{\mathbf{q} \equiv n\}} \mathbb{P}(N_i > (\theta_i - c_{\text{Bin}})n, \forall i)$  and the latter probability converges to 1 thanks to Lemma 4, hence the lower bound is proved.

The upper bound is a bit delicate. Let us start by the trivial bound, if  $|\mathbf{q}_i| > n$  for some  $i$ ,

$$0 = \mathbb{P}(S_{n,i} = \mathbf{q}_i \mid \mathbf{N}) = \mathbb{P}(S_n = \mathbf{q}) \leq \Theta n^{-1/2} \exp(-\Theta n^{-1} \mathbf{q}_i^2).$$

Assume henceforth that  $|\mathbf{q}_i| \leq n$  for all  $i$ . Let  $\Omega$  be the event that for some  $i$ ,  $N_i < \theta_i(1 - c_{\text{Bin}})n$ . On  $\Omega^c$ ,  $N_i = \Theta n$  for all  $i$ , hence

$$\mathbb{P}(S_{n,i} = \mathbf{q}_i \mid \Omega^c) \leq \Theta n^{-1/2} \exp(-\Theta n^{-1} \mathbf{q}_i^2).$$

Finally, in all cases,

$$\begin{aligned} \mathbb{P}(S_n = \mathbf{q}) &\leq \mathbb{E}(\mathbf{1}_{\{\Omega^c\}} \prod_i \mathbb{P}(S_{n,i} = \mathbf{q}_i \mid \mathbf{N})) + \mathbb{P}(\Omega) \\ &\leq \mathbb{E}(\mathbf{1}_{\{\Omega^c\}} \Theta \prod_i n^{-\frac{1}{2}} \exp(-\Theta n^{-1} \mathbf{q}_i^2)) + \mathbb{P}(\Omega) \\ &\leq n^{-p/2} \exp(-\Theta n^{-1} \mathbf{q}^2) + \mathbb{P}(\Omega). \end{aligned}$$

Then Lemma 4 with  $\varepsilon = -c_{\text{Bin}}$  yields, using the decreasing of binomial probabilities around the mean,

$$\begin{aligned}
\mathbb{P}(\Omega) &\leq \sum_i \sum_{k < [n\theta_i(1-c_{\text{Bin}})]} \mathbb{P}(N_i = k) \\
&\leq \sum_i \Theta n \mathbb{P}(N_i = n\theta_i(1+\varepsilon)) \\
&\leq \Theta n^{1/2} \exp(-\Theta n) \\
&\leq \Theta n^{-\frac{p}{2}} \exp(-\Theta n/2) \\
&\leq \Theta n^{-\frac{p}{2}} \exp(-\Theta n^{-1} \mathbf{q}^2),
\end{aligned}$$

using  $|\mathbf{q}| \leq n$ , which concludes the proof of (18).  $\square$

**Lemma 3.** For  $i \in \{0, 1\}$

$$\sum_{\mathbf{x} \in \mathbb{Z}^d, \mathbf{x} \equiv i} \exp(-\Theta n^{-1} \mathbf{x}^2) = \Theta n^{d/2}.$$

where  $\mathbf{x} \equiv i$  means that  $\sum_{k=1}^d \mathbf{x}_{[k]}$  has the same parity as  $i$ .

*Proof.* The lower bound stems from  $y^2 \geq \min_{\mathbf{x} \in \mathbb{Z}^d \cap B(y, 2), \mathbf{x} \equiv i \text{ or } \mathbf{x} = 0} \mathbf{x}^2$ ,  $y \in \mathbb{R}^d$ , and

$$\begin{aligned}
\Theta n^{d/2} &\leq \int_{\mathbb{R}^d} \exp(-\Theta n^{-1} y^2) dy \leq \int_{\mathbf{x} \in \mathbb{Z}^d \cap B(y, 2), \mathbf{x} \equiv i \text{ or } \mathbf{x} = 0} \max \exp(-\Theta n^{-1} \mathbf{x}^2) dy \\
&\leq 4^d \sum_{\mathbf{x} \in \mathbb{Z}^d, \mathbf{x} \equiv i \text{ or } \mathbf{x} = 0} \exp(-\Theta n^{-1} \mathbf{x}^2) \\
&\leq 4^d \left( \sum_{\mathbf{x} \in \mathbb{Z}^d, \mathbf{x} \equiv i} \exp(-\Theta n^{-1} \mathbf{x}^2) + 1 \right)
\end{aligned}$$

because at most a mass  $4^d$  of  $y$ 's are within distance 2 from some  $\mathbf{x} \in \mathbb{Z}^d$ . For the upper bound, for  $\mathbf{x} \in \mathbb{Z}^d \setminus \{0\}$ , there is at least one cube  $C_{\mathbf{x}}$  within the  $2^d$  cubes that touch  $\mathbf{x}$  such that for all  $y \in C_{\mathbf{x}}$ ,  $y^2 \leq \mathbf{x}^2$ . Hence

$$\begin{aligned}
\sum_{\mathbf{x} \equiv i} \exp(-\Theta n^{-1} \mathbf{x}^2) &\leq \sum_{\mathbf{x} \equiv i, \mathbf{x} \neq 0} \exp(-\Theta n^{-1} \mathbf{x}^2) + 1 \leq \sum_{\mathbf{x} \in \mathbb{Z}^d \setminus \{0\}} \int_{C_{\mathbf{x}}} \exp(-\Theta n^{-1} y^2) dy + 1 \\
&\leq 2^d \int_{\mathbb{R}^d} \exp(-\Theta n^{-1} y^2) dy + 1 \leq \Theta n^{d/2}.
\end{aligned}$$

$\square$

### 3.3 Binomial estimates

**Lemma 4.** Let  $\theta_0 < 1$ . There is a universal constant  $c_{\text{Bin}} \in (0, 1)$  such that for  $0 < \theta \leq \theta_0$ ,  $B \sim \mathcal{B}(m, \theta)$ , for  $-c_{\text{Bin}}\theta \leq \varepsilon_m = \varepsilon \leq c_{\text{Bin}}\theta$

$$\mathbb{P}(B = [m(\theta + \varepsilon)]) = \Theta m^{-1/2} \exp(-\Theta m \varepsilon^2)$$

where the constants involved in  $\Theta$  depend on  $\theta_0$ .

*Proof.* Let  $k = [m(\theta + \varepsilon)]$ . By Stirling's formula,

$$\begin{aligned}
\mathbb{P}(B = k) &= \Theta \frac{\sqrt{m}}{\sqrt{k}\sqrt{m-k}} \theta^k (1-\theta)^{m-k} \frac{m^m}{k^k (m-k)^{m-k}} \\
&= \Theta m^{-1/2} \frac{\theta^k (1-\theta)^{m-k}}{\sqrt{(\theta + \varepsilon)(1-\theta - \varepsilon)}} \frac{m^m}{(\theta m)^k (m(1-\theta))^{m-k} \left(\frac{k}{\theta m}\right)^k \left(\frac{m-k}{m(1-\theta)}\right)^{m-k}} \\
&= \Theta m^{-1/2} \frac{1}{\sqrt{\theta}} \left(1 + \frac{\varepsilon}{\theta}\right)^{-k} \left(1 - \frac{\varepsilon}{1-\theta}\right)^{k-m} \\
&= \Theta m^{-1/2} \theta^{-1/2} \exp(\gamma_{\varepsilon, \theta})
\end{aligned}$$

where for  $|\varepsilon| \leq \theta/2$

$$\begin{aligned}
\gamma_{\varepsilon, \theta} &= -m(\theta + \varepsilon) \left( \frac{\varepsilon}{\theta} - \frac{\varepsilon^2}{2\theta^2} + O(\varepsilon^3/\theta^3) \right) - m((1-\theta) - \varepsilon) \left( -\frac{\varepsilon}{1-\theta} + \frac{\varepsilon^2}{2(1-\theta)^2} + O(\varepsilon^3) \right) \\
&= m \frac{\varepsilon^2}{2\theta} - \frac{m\varepsilon^2}{\theta} + O(m\varepsilon^3\theta^{-2} + m\varepsilon^4\theta^{-3}) - \frac{m\varepsilon^2}{2(1-\theta)} - \frac{m\varepsilon^2}{1-\theta} + O(m\varepsilon^3 + m\varepsilon^4) \\
&= \frac{-m\varepsilon^2}{2\theta} - \frac{3m\varepsilon^2}{2(1-\theta)} + O(m\varepsilon^3\theta^{-2})
\end{aligned}$$

using  $\varepsilon \leq \theta$ , hence for some  $C$

$$\left| \gamma_{\varepsilon, \theta} - \left( \frac{-m\varepsilon^2}{2\theta} - \frac{3m\varepsilon^2}{2(1-\theta)} \right) \right| \leq Cm\varepsilon^3\theta^{-1}$$

and for every  $C' > 0$  there is  $c$  such that for  $|\varepsilon| < c$ , the right hand side is smaller than  $C'm\varepsilon^2\theta^{-1}$ , hence  $\gamma_{\varepsilon, \theta} = -\Theta m\varepsilon^2$  for  $c$  sufficiently small and  $|\varepsilon| \leq c$ .  $\square$

## 4 Variance estimates

Let  $X$  be some stationary Gaussian field on  $\mathbb{R}^d$  with spectral measure  $\boldsymbol{\mu}$  and reduced covariance function denoted by  $C$ . We study here the excursion set

$$E_u = \{\mathbf{t} \in \mathbb{R}^d : X(\mathbf{t}) \geq u\}, u \in \mathbb{R},$$

and the statistic

$$M_u^\gamma = \int_{\mathbb{R}^d} \mathbf{1}_{\{X(\mathbf{t}) > 0\}} \gamma(\mathbf{t}) d\mathbf{t}$$

where  $\gamma$  is some measurable *window function*, bounded with compact support. Define

$$\hat{\gamma}(\mathbf{x}) = \int_{\mathbb{R}^d} e^{i\mathbf{t} \cdot \mathbf{x}} \gamma(\mathbf{t}) d\mathbf{t}, \mathbf{x} \in \mathbb{R}^d,$$

and  $\gamma_T(\mathbf{t}) = \gamma(T^{-1}\mathbf{t})$ ,  $T > 0$ . The variance is

$$V_{\boldsymbol{\mu}}^{\gamma, u}(T) = \mathbf{Var}(\mathcal{L}^d(M_u^{\gamma_T})) \text{ with the notation } V_{\boldsymbol{\mu}}^\gamma := V_{\boldsymbol{\mu}}^{\gamma, 0}.$$

The most prominent example is the unit sphere indicator  $\gamma^d = \mathbf{1}_{\{B_d(0,1)\}}(\cdot)$ , and in this case  $\gamma^d$  is also implicit in the notation  $V_{\boldsymbol{\mu}}^{\gamma^d, u} = V_{\boldsymbol{\mu}}^u$ . Recall the notation

$$\mathbf{J}_3(\varepsilon) = \sum_{n \in \mathbb{N} \text{ odd}} n^{-\frac{3}{2}} \mathbb{P}(|\mathbf{U}_n| < \varepsilon)$$

where  $(\mathbf{U}_n)_n$  is the random walk on  $\mathbb{R}^d$  with increment law  $\boldsymbol{\mu}$ . Say that  $\boldsymbol{\mu}$  is  $\mathbb{Z}$ -free if  $\mathbb{P}(\mathbf{U}_n = 0) = 0$  for  $n \in \mathbb{N}$  odd.

**Theorem 6.** Assume  $\mu$  is  $\mathbb{Z}$ -free and  $\hat{\gamma}(\mathbf{t}) \geq c_1 > 0$ ,  $\mathbf{t} \in B_d(0, 4r)$  for some  $r > 0$ .

(i) Then for some  $c_2 > 0$ , as  $T \rightarrow \infty$ ,

$$c_2 T^{2d} \mathbf{J}_3(rT^{-1}) \stackrel{\text{i.o.}}{\leq} V_\mu^\gamma(T). \quad (19)$$

(ii) If in addition for some  $c_3 < \infty$ ,  $|\hat{\gamma}(\mathbf{x})| \leq c_3 \|\mathbf{x}\|^{-\frac{d+1}{2}}$  as  $\|\mathbf{x}\| \rightarrow \infty$ , then for some  $c_4 < \infty$ ,

$$V_\mu^\gamma(T) \leq c_4 (T^{2d} \mathbf{J}_3(rT^{-1}) + T^{d-1} \int_1^{(T/r)^{d+1}} \mathbf{J}_3(y^{-\frac{1}{d+1}}) dy). \quad (20)$$

(iii) If the hypotheses of (i) and (ii) and Proposition 7 below are satisfied, as  $T \rightarrow \infty$ ,

$$V_\mu^\gamma(T) = o(T^{2d}).$$

(iv) There is  $c_5 > 0$  such that for  $u \neq 0$ ;

$$V_\mu^{\gamma,u}(T) \geq c_5 T^{2d} \mathbb{P}(\|\mathbf{U}_2\| < rT^{-1}).$$

In particular if  $\mu$  has an atom at  $\mathbf{x}_0$ , we have for  $u \neq 0$

$$c_5 \mu(\{\mathbf{x}_0\})^2 T^{2d} \leq V_\mu^{\gamma,u}(T) \leq T^{2d}.$$

The bounds (i) and (ii) are sharp when applied to diophantine Gaussian fields, which have a purely atomic measure (see Section 1), and we show below that it is still the case in the context of more diffuse isotropic spectral measures; we are in particular able to capture the variance cancellation phenomenon for Gaussian random waves (see Theorem 3).

**Example 1.** For the unit sphere indicator, we have the classical formula ([13, Chap. 5])

$$\hat{\gamma}^d(\mathbf{x}) = \kappa_d |\mathbf{x}|^{-d/2} \mathbf{B}_{d/2}(|\mathbf{x}|)$$

where  $\kappa_d = \mathcal{L}^d(B_d(0, 1))$  and  $\mathbf{B}_a$  is the Bessel function of the first kind with parameter  $a$

$$\mathbf{B}_a(r) = \sum_{m=0}^{\infty} \frac{(-1)^m}{m! \Gamma(m+a+1)} \left(\frac{r}{2}\right)^{2m+a}, r \geq 0.$$

In particular,  $\hat{\gamma}^d(\mathbf{x}) \sim \kappa_d \Gamma(d/2+1)^{-1} > 0$  in 0 and

$$\hat{\gamma}^d(\mathbf{x}) \sim \kappa_d (2/\pi)^{1/2} |\mathbf{x}|^{-\frac{d+1}{2}} \cos(|\mathbf{x}| + \Delta_d)$$

as  $\mathbf{x} \rightarrow \infty$ , for some  $\Delta_d \in \mathbb{R}$ . It is known [11] that the first zero of  $\mathbf{B}_a$ ,  $a \geq 1/2$  is larger than the first zero of  $\mathbf{B}_{1/2}$ , which is  $\pi$ , hence we can take  $r = \frac{1}{2}$  in the result above.

The terms on the right hand side of (20) have similar magnitude in the regular diophantine case:

**Proposition 7.** Let  $\psi(q) = q^{-\tau} L(q)$  be regular (Definition 1) and assume  $\mu$  is of the form (2) with  $\omega$  that is  $\psi$ -BA. Let  $\tau^* = \frac{1+d(m+1)}{d+1}$ . Then as  $T \rightarrow \infty$ ,

$$\max \left\{ T^{2d} \mathbf{J}_3(T^{-1}), T^{d-1} \int_1^{T^{d+1}} \mathbf{J}_3(y^{-\frac{1}{d+1}}) dy \right\} \leq \begin{cases} \Theta T^{d-1} & \text{if } \psi(q) > q^{-\tau^*} \log(q)^{1/d} \\ \Theta T^{d-1} \ln(T) & \text{if } \psi(q) = q^{-\tau^*} \\ \Theta T^{2d} \psi^{-1}(T^{-1})^{-1-d(m+1)} & \text{if } \tau > \tau^* \end{cases}$$

**Remark 9.** It is likely that the upper bound in  $T^{d-1} \ln(T)$  is sharp if  $\psi(q) = q^{-\tau^*}$ , proving it rigourously would require a lower bound for  $v_T^{(2)}$  in the proof, which raises some technical difficulties because of the cosine term.

*Proof.* According to (9) in Theorem 5-(i),

$$\mathbf{J}_3(\varepsilon) \leq \Theta \psi^{-1}(\varepsilon)^{-1-d(m+1)}, \varepsilon > 0,$$

which yields that  $T^{2d} \mathbf{J}_3(T^{-1})$  admits an upper bound consistent with the claim.

To deal with the other term, assume without loss of generality that  $\psi$  is extended to a smooth strictly non-increasing function  $z^{-\tau} L(z) : [a, \infty) \rightarrow (0, 1]$  for some  $a \geq 1$ , such that  $L'(z) = o(z^{-1} L(z))$ . Make the change of variables  $z = \psi^{-1}(y^{-\frac{1}{d+1}})$ , i.e.  $\psi(z)^{-d-1} = y$ ,  $Z = \psi^{-1}(T^{-1}) \leq \Theta \psi^{-1}(T^{-1})$ ,

$$\int_a^{T^{d+1}} \mathbf{J}_3(y^{-\frac{1}{d+1}}) dy \leq \Theta \int_a^{T^{d+1}} \psi^{-1}(y^{-\frac{1}{d+1}})^{-1-d(m+1)} dy = \Theta \int_{\Theta}^Z z^{-1-d(m+1)} (\psi(z)^{-d-1})' dz.$$

The hypotheses on  $\psi$  yield

$$\begin{aligned} (\psi(z)^{-d-1})' &= (d+1)(\tau z^{-\tau-1} L(z) - z^{-\tau} L'(z)) \psi(z)^{-d-2} = (d+1)(\tau z^{-1} \psi(z) - z^{-\tau} o(z^{-1} L(z))) \psi(z)^{-d-2} \\ &\sim_{z \rightarrow \infty} (d+1) \tau z^{-1} \psi(z)^{-d-1}. \end{aligned} \quad (21)$$

In the case  $\tau \leq \tau^*$ , the previous two displays yield

$$\int_a^{T^{d+1}} \mathbf{J}_3(y^{-\frac{1}{d+1}}) dy \leq \Theta \int_{\Theta}^Z z^{-2-d(m+1)} \psi(z)^{-(d+1)} dz$$

and the integral converges if  $\psi(q) > q^{-\tau^*} \ln(q)^{1/d}$ , and if  $\psi(q) = q^{-\tau^*}$  it behaves in  $\ln(Z) = \Theta \ln(T)$ .

Let us turn to the case  $\tau > \tau^*$ . Let  $\tau' \in (\tau^*, \tau)$ , we have by (21) as  $z \rightarrow \infty$

$$\begin{aligned} (z^{-1-d(m+1)} \psi(z)^{-d-1})' &= z^{-1-d(m+1)} (\psi(z)^{-d-1})' - (1+d(m+1)) z^{-2-d(m+1)} \psi(z)^{-d-1} \\ &\geq z^{-1-d(m+1)} (\psi(z)^{-d-1})' - \frac{1+d(m+1)}{d+1} z^{-1-d(m+1)} (\psi(z)^{-d-1})' \\ &\geq z^{-1-d(m+1)} (\psi(z)^{-d-1})' (1 - \frac{\tau^*}{\tau'}) \end{aligned}$$

which results in

$$\int_a^{T^{d+1}} \mathbf{J}_3(y^{-\frac{1}{d+1}}) dy \leq \frac{\Theta}{1 - \tau^*/\tau'} [z^{-1-d(m+1)} \psi(z)^{-d-1}]_{\Theta}^Z = \Theta \psi^{-1}(T^{-1})^{-1-d(m+1)} T^{d+1}$$

which allows to conclude.  $\square$

*Proof of Theorem 6.* The starting point is the following lemma, available for instance partially in [5, Lemma 2].

**Lemma 5.** *We have for every  $u \in \mathbb{R}$  coefficients  $\alpha_{n,u} \geq 0, n \in \mathbb{N}$  such that for two standard gaussian variables  $X, Y$  with correlation  $\rho$*

$$\Gamma_u(\rho) := \text{Cov}(\mathbf{1}_{\{X>u\}}, \mathbf{1}_{\{Y>u\}}) = \sum_{n=1}^{\infty} \alpha_{n,u} \rho^n = \frac{1}{2\pi} \int_0^{\rho} \frac{1}{\sqrt{1-r^2}} \exp\left(-\frac{u^2}{1+r}\right) dr \quad (22)$$

*in particular,  $\Gamma_0(\rho) = \arcsin(\rho)$  with  $\alpha_{2n,0} = 0$  and*

$$\alpha_{2n+1} := \alpha_{2n+1,0} = \frac{\binom{n}{2n}}{4^n (2n+1)} = \Theta n^{-3/2}.$$

*We also have  $\alpha_{2,u} \neq 0$  for  $u \neq 0$ .*

*Proof.* The result is a consequence of the proof of [5, Lemma 2]. Denoting by  $\Phi^{(k)}$  the  $k$ -th order derivative of the standard Gaussian distribution function, it is shown there that (22) is true with

$$\alpha_{n,u} = \frac{\Phi^{(n)}(u)^2}{n!},$$

hence for instance  $\alpha_{2,u} > 0$  for  $u \neq 0$ . For  $u = 0$  we have the explicit decomposition of the arcsine function with

$$\Gamma_0(\rho) = \arcsin(\rho) = \sum_{n=0}^{\infty} \alpha_{2n+1} \rho^{2n+1} \text{ with } \alpha_{2n+1} = \frac{\binom{2n}{n}}{4^n(2n+1)} \sim_{n \rightarrow \infty} \frac{1}{\sqrt{\pi n}(2n+1)}.$$

□

Let  $\mathbf{U}_n = \sum_{i=1}^n X_i$  where the  $X_i$  are independent and identically distributed with law  $\boldsymbol{\mu}$ . Recall the Fourier inversion formula

$$\gamma(\mathbf{t}) = (2\pi)^{-d} \int \widehat{\gamma}(\mathbf{t}) e^{it \cdot \mathbf{x}} d\mathbf{x}, \mathbf{t} \in \mathbb{R}^d.$$

Denote by  $\gamma^{*2}$  the auto-convolution of  $\gamma$  with itself, and by  $\boldsymbol{\mu}^{*(n)}$  the  $n$ -fold convolution of the probability measure  $\boldsymbol{\mu}$  with itself, which is the law of  $\mathbf{U}_n$ . We have by Lemma 5

$$\begin{aligned} V_{\boldsymbol{\mu}}^u(T) &= \int_{(\mathbb{R}^d)^2} \Gamma_u(C(\mathbf{t} - \mathbf{s})) \gamma(\mathbf{t}/T) \gamma(\mathbf{s}/T) d\mathbf{t} d\mathbf{s} & (23) \\ &= \int_{(\mathbb{R}^d)^2} \Gamma_u(C(\mathbf{z})) \gamma\left(\frac{\mathbf{z} + \mathbf{w}}{2T}\right) \gamma\left(\frac{\mathbf{w} - \mathbf{z}}{2T}\right) d\mathbf{w} d\mathbf{z} \\ &= \int_{\mathbb{R}^d} \Gamma_u(C(\mathbf{z})) \gamma_{2T}^{*2}(2\mathbf{z}) d\mathbf{z} \\ &= \sum_{n \in \mathbb{N}} \alpha_{n,u} \int C(\mathbf{z})^n \gamma_{2T}^{*2}(2\mathbf{z}) d\mathbf{z} \\ &= \sum_{n \in \mathbb{N}} \alpha_{n,u} \int \boldsymbol{\mu}^{*(n)}(d\mathbf{z}) \hat{\gamma}_{2T}(2\mathbf{z})^2 d\mathbf{z} \text{ using (1)} \\ &= \sum_{n \in \mathbb{N}} \alpha_{n,u} \int \boldsymbol{\mu}^{*(n)}(d\mathbf{z}) (2T)^{2d} \hat{\gamma}(4T\mathbf{z})^2 d\mathbf{z} \\ &= (2T)^{2d} \sum_{n \in \mathbb{N}} \alpha_{n,u} \mathbb{E}(\hat{\gamma}(4T\mathbf{U}_n)^2) \\ &= 2^{2d} (v_T^{(1)} + v_T^{(2)}) \end{aligned}$$

where, with  $A_1 = [0, r], A_2 = ]r, \infty]$

$$v_T^{(i)} = T^{2d} \sum_{n \in \mathbb{N}} \alpha_{n,u} \mathbb{E}(\hat{\gamma}(4T\mathbf{U}_n)^2 \mathbf{1}_{\{\|\mathbf{U}_n\| \in A_i\}}).$$

For the case  $u \neq 0$ , we simply have according to Lemma 5

$$V_{\boldsymbol{\mu}}(T) \geq 2^{2d} v_T^{(1)} \geq 2^{2d} T^{2d} \alpha_{2,u} \mathbb{P}(\mathbf{U}_2 = 0) \gamma_d(0)^2 > 0.$$

Let us now focus on the case  $u = 0, \alpha_n = \alpha_{n,u} = \Theta n^{-3/2} \mathbf{1}_{\{n \text{ odd}\}}$ , and estimate  $v_T^{(1)}$ . Remark first that since  $\boldsymbol{\mu}$  is  $\mathbb{Z}$ -free,  $\mathbb{P}(\mathbf{U}_n = 0) = 0$  for  $n$  odd, and since by hypothesis  $C \geq \hat{\gamma}(\mathbf{t}) \geq c > 0$  on  $B_d(0, 4r)$  for some  $r \in (0, 1)$ ,

$$v_T^{(1)} = T^{2d} \sum_{n \text{ odd}} \Theta n^{-3/2} \mathbb{P}(\|\mathbf{U}_n\| < rT^{-1}) = \Theta T^{2d} \mathbf{J}_3(rT^{-1})$$



hence (19) is proved.

For the second point, the second term satisfies

$$\begin{aligned}
v_T^{(2)} &\leq T^{2d} \sum_{n \text{ odd}} \Theta n^{-3/2} \mathbb{E}(\|T\mathbf{U}_n\|^{-d-1} \mathbf{1}_{\{|T\mathbf{U}_n|>r\}}) \\
&\leq T^{2d} \sum_{n \text{ odd}} \Theta n^{-3/2} \int_0^{r^{-d-1}} \mathbb{P}((T\|\mathbf{U}_n\|)^{-d-1} > y) dy \\
&\leq \Theta T^{2d} T^{-d-1} \sum_{n \text{ odd}} n^{-3/2} \int_0^{(T/r)^{d+1}} \mathbb{P}(\|\mathbf{U}_n\| < y^{-\frac{1}{d+1}}) dy \\
&= \Theta T^{d-1} \int_1^{(T/r)^{d+1}} \mathbf{J}_3(y^{-\frac{1}{d+1}}) dy.
\end{aligned}$$

Point (iii) is a consequence of Proposition 7 and the fact that  $\psi^{-1}$  converges to 0 in 0.  $\square$

## 4.1 Proof of Theorem 3

We study the field at the original scale  $X_d$ , it is then straightforward to deduce the results for  $X_{d,T}, T > 0$ . We need to estimate  $\mathbb{P}(\|\mathbf{U}_n\| \leq \varepsilon)$  for  $n \geq 1$ , where  $\mathbf{U}_n$  is the random walk which increment measure is  $\boldsymbol{\mu}_d$ . Equation (3) and the universal bound  $|B_a(\mathbf{t})| \leq \Theta \|\mathbf{t}\|^{-1/2}, \mathbf{t} \in \mathbb{R}^d$  yield  $|C_d(\mathbf{t})| \leq \Theta(1 + \|\mathbf{t}\|)^{-\frac{d-1}{2}}$ . Then

$$\mathbb{P}(\|\mathbf{U}_n\| \leq \varepsilon) = \int \mathbf{1}_{\{B_\varepsilon\}}(\mathbf{z}) \boldsymbol{\mu}^{*n}(d\mathbf{z}) \leq \varepsilon^d \int \mathbf{1}_{\{B_1\}}(\mathbf{t}) \left| \int_{\mathbb{R}^d} C(\mathbf{t})^n e^{i\varepsilon \mathbf{t} \mathbf{z}} d\mathbf{z} \right| d\mathbf{t} \leq \Theta \varepsilon^d \int_{\mathbb{R}^d} (1 + \|\mathbf{t}\|)^{-n \frac{d-1}{2}} d\mathbf{t},$$

hence  $\mathbb{P}(\|\mathbf{U}_n\| \leq \varepsilon) \leq c_{5+} \varepsilon^d < \infty$  for  $n \geq 5$  (and  $n \geq 3$  if  $d \geq 4$ ). We still have to deal with  $1 \leq n \leq 4$ , and independently with the lower bounds. Let us analyse the self-convoluted measures  $\boldsymbol{\mu}_d^{*n}, n \geq 1$ . They are related by the recurrence relation, based on the isotropy of the measures  $\boldsymbol{\mu}^{*n}, n \geq 1$ ,

$$\begin{aligned}
\boldsymbol{\mu}_d^{*n+1}(B_d(0, r)) &= \int_{\mathbb{S}^{d-1} \times \mathbb{R}^d} \mathbf{1}_{\{s+x \in B_d(0, r)\}} \boldsymbol{\mu}_d^{*n}(ds) \boldsymbol{\mu}_d(dx) \\
&= \boldsymbol{\mu}_d(\mathbb{S}^{d-1}) \boldsymbol{\mu}_d^{*n}(\{x : x + e_1 \in B_d(0, r)\}) = \boldsymbol{\mu}_d^{*n}(B_d(e_1, r))
\end{aligned} \tag{24}$$

where  $e_1$  is some vector of  $\mathbb{S}^{d-1}$ , e.g.  $e_1 = (1, 0, \dots, 0)$ . Hence

$$\boldsymbol{\mu}^{*2}(B_d(0, r)) = \boldsymbol{\mu}_d(\mathbb{S}^{d-1} \cap B_d(e_1, r)) \tag{25}$$

which is equivalent to  $\mathcal{L}^{d-1}(B_{d-1}(0, r))$  as  $r \rightarrow 0$ . It implies that

$$\mathbf{J}_3(\varepsilon) \geq 2^{-3/2} \boldsymbol{\mu}^{*2}(B_d(0, r)) \geq c_{2,-} \varepsilon^{d-1}, \varepsilon > 0$$

for some  $c_{2,-} > 0$ . Theorem 6-(i) then implies the lower bound in the case  $u \neq 0$ .

Since  $\alpha_{2,0} = \alpha_{4,0} = 0$  (Lemma 5), to treat the case  $u = 0$  we must study  $\boldsymbol{\mu}^{*3}$  (only for  $d = 2$  and  $d = 3$ ). Using (24)-(25) easily yields  $0 < c_{3,-} \varepsilon^d \leq \boldsymbol{\mu}_d^{*3}(B_d(0, \varepsilon)) \leq c_{3,+} \varepsilon^d < \infty$  as  $\varepsilon \rightarrow 0$ . Hence

$$2^{-3/2} c_{3,-} \varepsilon^d \leq \mathbf{J}_3(\varepsilon) = \sum_{n \geq 3, n \text{ odd}} n^{-3/2} \mathbb{P}(\|\mathbf{U}_n\| \leq \varepsilon) \leq c_{3,+} \varepsilon^d + \sum_{n=5}^{\infty} n^{-3/2} c_{5+} \varepsilon^d$$

gives the desired upper and lower bounds for  $u = 0$  (using Theorem 6-(i),(ii)).

## Acknowledgements

I am indebted to M. A. Klatt, with whom I had many discussions about hyperuniformity and Gaussian excursions. I also wish to thank S. Torquato for comments on the final manuscript. I am particularly grateful to Yann Bugeaud for insights about diophantine approximation.

This study was supported by the IdEx Université de Paris, ANR-18-IDEX-0001.

## References

- [1] R. J. Adler and J. E. Taylor. *Random Fields and Geometry*. Springer, 2007.
- [2] J. Beck. Irregularities of distribution. i. *Acta Math.*, 159:1–49, 1987.
- [3] Michael V Berry. Statistics of nodal lines and points in chaotic quantum billiards: perimeter corrections, fluctuations, curvature. *Journal of Physics A: Mathematical and General*, 35(13):3025, 2002.
- [4] G. Blum, S. Gnutzmann, and U. Smilansky. Nodal domains statistics: A criterion for quantum chaos. *Phys. Rev. Lett.*, 88(114101), 2002.
- [5] A. Bulinski, E. Spodarev, and F. Timmermann. Central limit theorems for the excursion set volumes of weakly dependent random fields. *Bernoulli*, 18(1):100–118, 2012.
- [6] D. Chen and S. Torquato. Designing disordered hyperuniform two-phase materials with novel physical properties. *Acta Materialia*, 142:152–161, 2018.
- [7] C. Corduneanu, N. Gheorghiu, and V. Barbu. *Almost periodic functions*. Chelsea Publishing Company, 1989.
- [8] H. Cramér and M. R. Leadbetter. *Stationary and Related Stochastic Processes*. Wiley, 1967.
- [9] J. Cuzick. A central limit theorem for the number of zeros of a stationary Gaussian process. *Ann. Prob.*, 4(4):547–556, 1976.
- [10] P. Diaconis. Group representations in probability and statistics. *Inst. Math. Stat. Lect. Notes*, 11, 1988.
- [11] A. Elbert. Some recent results on the zeros of bessel functions and orthogonal polynomials. *J. Comp. Appl. Math.*, 133:65–83, 2001.
- [12] A. Estrade and J. R. Leon. A central limit theorem for the Euler characteristic of a Gaussian excursion set. *Ann. Prob.*, 44(6):3849–3878, 2016.
- [13] I. I. Gikhman and A. V. Skorokhod. *Introduction to the theory of random processes, Vol. I*. W. B. Saunders Company, 1965.
- [14] Ben Hough, Krishnapur, Peres, and Viràg. *Zeros of Gaussian Analytic Functions and Determinantal Point Processes*. University Lecture Series. Institute of Mathematical Statistics, 2009.
- [15] M. Hussain and T. Yusupova. A note on the weighted khintchine-groshev theorem. *J. de théorie des nombres de Bordeaux*, 26(2):385–397, 2014.

- [16] V. Jarník. über die konvexe hülle von  $n$  zufällig gewählten punkten iiber die simul-tanen diophantische approximationen. *Math. Z.*, 33:505–543, 1931.
- [17] M. Kac. On the average number of real roots of a random algebraic equation. *Bull. Amer. Math. Soc.*, 49:314–320, 1943.
- [18] M. Kratz and J. R. Leon. Hermite polynomial expansion for non-smooth functionals of stationary gaussian processes: Crossings and extremes. *Stoc. Proc. Appl.*, 66:237–252, 1997.
- [19] M. Krishnapur, P I. P. Kurlberg, and Wigman. Nodal length fluctuations for arith-metic random waves. *Annals of Mathematics*, 177(2):699–737, 2013.
- [20] R. Lachièze-Rey. Excursions of almost periodic gaussian processes and diophantine approximation. in preparation.
- [21] R. Lachièze-Rey. Variance linearity for real gaussian zeros. arXiv:2006.10341.
- [22] D. Marinucci, G. Peccati, M. Rossi, and I. Wigman. Non-universality of nodal length distribution for arithmetic random waves. *Geom. Funct. Anal.*, 26:926–960, 2016.
- [23] D. Marinucci and I. Wigman. The defect variance of random spherical harmonics. *J. Phys. A: Math. and Theor.*, 44(35):355206, 2011.
- [24] S. Muirhead, A. Rivera, and H. Vanneuville. The phase transition for planar gaussian percolation models without FKG. <https://arxiv.org/abs/2010.11770>, 2020.
- [25] F. Nazarov and M. Sodin. Fluctuations in random complex zeroes: Asymptotic normality revisited. *Int. Math. Res. Notic.*, 24:5720–5759, 2011.
- [26] I. Nourdin, G. Peccati, and M. Rossi. Nodal statistics of planar random waves. *Comm. Math. Phys.*, 369:99–151, 2019.
- [27] E. C. Oğuz, J. E. S. Socolar, P. J. Steinhardt, and S. Torquato. Hyperuniformity of quasicrystals. *Physical Review B*, 95(054119), 2017.
- [28] T. Prescott and F. E. Su. Random walks on the torus with several generators. *Rand. Struct. Alg.*, 25(3):336–345, 2004.
- [29] M. Rossi. The defect of random hyperspherical harmonics. *J. Theor. Prob.*, 32:2135–2165, 2019.
- [30] E. Slud. Multiple Wiener-Ito integral expansions for level-crossing-count functionals. *Prob. Th. Rel. Fields*, 87:349–364, 1991.
- [31] F. E. Su. Convergence of random walks on the circle generated by an irrational rotation. *Trans. AMS*, 350(9):3717–3741, 1998.
- [32] S. Torquato. Disordered hyperuniform heterogeneous materials. *J. Phys.: Condens. Matter*, 28(414012), 2016.
- [33] S. Torquato. Hyperuniformity and its generalizations. *Phys. Rev. E*, 94(022122), 2016.
- [34] S. Torquato. Hyperuniform states of matter. *Physics Reports*, 745:1–95, 2018.
- [35] S. Torquato and F. H. Stillinger. Local density fluctuations, hyperuniform systems, and order metrics. *Phys. Rev. E*, 68(041113):1–25, 2003.

- [36] I. Wigman. Fluctuations of the nodal length of random spherical harmonics. *Comm. Maths Phys.*, 298:787–831, 2010.
- [37] I. Wigman and A. Granville. The distribution of the zeros of random trigonometric polynomials. *Amer. J. Math.*, 133(2):295–357, 2011.