

Multiobjective Hypervolume Based ISOOMOO Algorithms Converge with At Least Sublinear Speed to the Entire Pareto Front

Eugénie Marescaux, Anne Auger

▶ To cite this version:

Eugénie Marescaux, Anne Auger. Multiobjective Hypervolume Based ISOOMOO Algorithms Converge with At Least Sublinear Speed to the Entire Pareto Front. 2021. hal-03198414v1

HAL Id: hal-03198414 https://hal.science/hal-03198414v1

Preprint submitted on 13 Jul 2021 (v1), last revised 22 Apr 2022 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1MULTIOBJECTIVE HYPERVOLUME BASED ISOOMOO2ALGORITHMS CONVERGE WITH AT LEAST SUBLINEAR SPEED TO3THE ENTIRE PARETO FRONT

EUGÉNIE MARESCAUX AND ANNE AUGER

Abstract. In multiobjective optimization, one is interested in finding a good approximation of the 56 Pareto set and the Pareto front, i.e the sets of best compromises in the decision and objective spaces, 7 respectively. In this context, we introduce a new algorithm framework, Incremental SingleObjective 8 Optimization for MultiObjective Optimization (ISOOMOO) for approximating the Pareto front with an increasing number of points. We focus on HV-ISOOMOO, its instanciation with the hypervolume 9 10 indicator, a set-quality indicator which is widely used for algorithms design and performance assessment. HV-ISOOMOO algorithms approximate the Pareto front by greedily maximizing the hypervolume. We 11 12study the convergence to the entire Pareto front of HV-ISOOMOO coupled with a perfect singleobjective 13 optimizer. The convergence is defined as the convergence of the hypervolume of the sets of all meta-14 iterations incumbents towards the hypervolume of the Pareto front. We prove tight lower bounds on the 15 convergence-speed for convex and bilipschitz Pareto fronts in $O(1/n^c)$ with n being the number of metaiterations and c = 1 and $c \leq 1$, respectively. For convex Pareto fronts, the convergence rate is exactly in 16 $\Theta(1/n)$, namely the highest convergence rate achievable by a biobjective optimization algorithm. These 17 are the first results on the convergence-speed of multiobjective optimization algorithms towards the entire 18 19Pareto front. We also analyze theoretically the asymptotic convergence behavior.

20 Key words. multiobjective optimization, convergence, hypervolume, pareto front

21 AMS subject classifications. 90C29, 90C30

4

22 1. Introduction. Real-world problems often involve the optimization of several conflicting objectives. The solution of such problems is the set of non-dominated decision 23 vectors, the Pareto set defined as the set of solutions that cannot be improved along one 24objective without degrading along another one. Its image in the objective-space is the 2526 Pareto front. A decision maker is then often involved to choose, based on its preferences, a single best compromise. The shape of the Pareto front informs on the trade-off between 27 objectives. Many algorithms such as evolutionary algorithms approximate the Pareto 28 front with a number of points fixed in the beginning. But some algorithms, in particular 29Direct Multi-Search Methods such as MultiGLODS [9], DMS [10] and D-Multi-MADS [6] 30 aim at approximating the entire Pareto set or Pareto front with as many well-distributed 31 32 points as possible. Ideally, the quality of the Pareto front approximation increases with time without stagnating and such algorithms can be considered as *anytime* algorithms. 33

The convergence speed towards a critical decision vector or a vector of the Pareto 34 front have been examined for many algorithms such as (1+1) evolutionary multiobjective 35algorithms [4] or Newton's method [13]. Convergence rates are typically similar to the 36 ones obtained for singleobjective optimization. Indeed, this is still a convergence towards 38 a single point. Its analysis is sometimes reduced to the study of the convergence of a singleobjective optimization algorithm. The convergence of anytime algorithms towards 39 the whole Pareto set or front is of a different kind because these are sets and not points. 40 It has already been theoretically investigated for some algorithms [9] and more abstract 41 frameworks [18], but analysis of the rate of convergence are missing. Additionally, empir-42 ical studies typically focus on determining which algorithm is faster and do not provide 43 orders of convergence or precise convergence rate. Yet, while largely overlooked, studying 44 convergence rates either theoretically or empirically is crucial. In this context, it has been 45

46 proven that convergence towards the whole Pareto front is always sublinear in the number 47 of function evaluations, at least when measuring convergence with the hypervolume indi-48 cator [15] or the multiplicative ϵ -indicator [7], and thus much slower than typical speeds 49 of convergence to a single point. The hypervolume is a set-quality indicator widely used 50 in multiobjective optimization, both to guide algorithms and for performance assessment. 51 It is at the core of all known stricly Pareto-compliant indicators [19].

In this paper, we introduce a new anytime algorithm framework, Incremental Sin-52gleObjective Optimization for MultiObjective Optimization (ISOOMOO). We focus on its instanciation with the hypervolume indicator, HV-ISOOMOO. HV-ISOOMOO al-54gorithms try to greedily maximize the hypervolume by adding points with the highest 56 hypervolume contribution. These points are obtained by running a singleobjective optimizer. Greediness is already used in the selection part of some multiobjective optimization evolutionary algorithms such as SMS-EMOA [5] to approximate the set of p points among 58 the population with the highest hypervolume. The hypervolume of such discrete greedy 5960 approximation is proven to be at least (e-1)/e times the one of the p-optimal distribution [16]. To our knowledge, we provide the first continuous equivalent of this result. We in-61 62 vestigate the rate of convergence of HV-ISOOMOO towards the whole Pareto front in the ideal case of perfect singleobjective optimization, measuring the convergence with the hy-63 pervolume. For convex and bilipschitz Pareto fronts, we prove that the convergence-speed 64 is in $O(1/n^c)$ with c = 1 and $c \leq 1$, respectively, with n being the number of singleob-65 jective optimization runs performed. For convex Pareto fronts, since it has already been 66 proven that no biobjective optimization algorithm can converge to the Pareto front faster 67 68 than in $\Omega(1/n)$ [15], the convergence rate is exactly in $\Theta(1/n)$. Additionally, we prove that for bilipschitz and smooth enough Pareto fronts doubling the number of points in 69 the approximation halves the optimality gap, asymptotically. In the proof process, we 70 obtain bounds on the normalized maximum hypervolume and a geometric interpretation 71 of optimality conditions. 72

The paper is organized as follows. In Section 2, we lay the foundations of the problem we investigate. In Section 3, we prove preliminary results later used to investigate convergence. In Section 4, we derive lower bounds on the convergence rate of the perfect version of HV-ISOOMOO for bilipschitz and convex Pareto fronts and an insight on the asymptotic convergence behavior.

Notations and conventions. For $a, b \in \mathbb{N}$, we note $[\![a; b]\!]$ the set $\{a, a+1, \ldots, b-1, b\}$. For a vector $u \in \mathbb{R}^2$, we note u_1 and u_2 respectively its first and its second coordinate. If the vector notation already contains an index, we separate the two indices with a comma. For simplicity sake, we often replace the set $\{u\}$ by u in the notations. We say that a function $f : \mathbb{R} \to \mathbb{R}$ is decreasing (respectively strictly decreasing) when for all x < y, we have $f(x) \ge f(y)$ (respectively f(x) > f(y)).

2. Background, algorithm framework and assumptions. Here, we lay the foundations of the problem we analyze. First, we recall some classic concepts of multiobjective optimization. Then, we introduce the ISOOMOO class of algorithms and its hypervolume based instanciation HV-ISOOMOO. We also formalize a mathematical abstraction of HV-ISOOMOO, the greedy set sequences. Finally, we examine our assumptions on the biobjective optimization problem. 2.1. Biobjective optimization problems, the Pareto front and the hyper volume indicator. We consider a biobjective minimization problem:

92 (2.1)
$$\min_{x \in \Omega \subset \mathbb{R}^d} F(x)$$

94 with $F: \Omega \subset \mathbb{R}^d \to \mathbb{R}^2 : x \mapsto (F_1(x), F_2(x))$. We define two dominance relations for 95 vectors in the objective space. We say that u weakly dominates v denoted by $u \preceq v$ if 96 $u_1 \leq v_1$ and $u_2 \leq v_2$ and that u dominates v denoted by $u \prec v$ if $u \preceq v$ and $u \neq v$. 97 A vector of the objective-space \mathbb{R}^2 is said *feasible* when it belongs to $F(\Omega)$. Solving the 98 optimization problem consists in finding a good approximation of the *Pareto front*, the 99 set of non-dominated feasible vectors, $\{F(X) : X \in \Omega, \forall Y \in \Omega, F(Y) \not\preceq F(X)\}$. We 100 restrict ourselves to Pareto fronts with an explicit representation:

101 (2.2)
$$PF_f = \{(x, f(x)) : x \in [x_{\min}, x_{\max}]\}$$

103 with $f : \mathbb{R} \to \mathbb{R}$ decreasing. We denote by $u_{\min} := (x_{\min}, f(x_{\min}))$ and $u_{\max} :=$ 104 $(x_{\max}, f(x_{\max}))$ the extreme vectors of the Pareto front. Likewise, we denote by $\tilde{u}_{\min,r} :=$ 105 $(\tilde{x}_{\min,r}, f(\tilde{x}_{\min,r}))$ and $\tilde{u}_{\max,r} := (\tilde{x}_{\max,r}, f(\tilde{x}_{\max,r}))$ the extremes vectors of the part of 106 the Pareto front dominating a reference point r, with $\tilde{x}_{\min,r} := \max(x_{\min}, f^{-1}(r_2))$ and 107 $\tilde{x}_{\max,r} := \min(x_{\max}, r_1)$. The vector $(x_{\max}, f(x_{\min}))$ is called the *nadir* point. All these 108 notations are illustrated in Figure 1.



Fig. 1: Illustration of notations. The extreme vectors u_{\min} and u_{\max} and the nadir point $(x_{\max}, f(x_{\min}))$ (leftmost); the extreme vectors relative to the reference point $r \tilde{u}_{\min,r}$ and $\tilde{u}_{\max,r}$ (left); three vectors u_1, u_2 and u_3 and the regions weakly dominated by them and dominating $r, \mathcal{D}_{u_1}^r \square, \mathcal{D}_{u_2}^r \square$ and $\mathcal{D}_{u_3}^r \square$ (right); the greedy set $S_3 = \{v_1, v_2, v_3\}$, its four gap regions \square and the associated reference points (rightmost).

108

The hypervolume with respect to a reference point r of an assessed set S of objective vectors, that we denote by $HV_r(S)$, is the Lebesgue measure of the region of the objectivespace dominated by S and strictly dominating the reference point r. When no vector of the Pareto front dominates the reference point r, the hypervolume with respect to r of any set of feasible points of the objective space is null. Since this particular case is not interesting, we only consider reference points dominated by at least one vector of the Pareto front from now on. We refer to such reference points as *valid*.

The region of the objective-space dominated by S and dominating r, see the righthand plot of Figure 1, is denoted by \mathcal{D}_S^r and formally defined as:

$$\mathcal{D}_S^r = \{ w \in \mathbb{R}^2 : \exists u \in S : u \preceq w \prec r \}.$$

120 The hypervolume of a set S relative to the reference point r equals $\lambda(\mathcal{D}_S^r)$ with $\lambda(.)$ being 121 the Lebesgue measure. In this paper, we only consider two-dimensional objective spaces 122 and refer to the Lebesgue measure of a set as its area. The set \mathcal{D}_S^r is the union of the \mathcal{D}_u^r 123 for $u \in S$, \mathcal{D}_u^r being the rectangle $[u_1, r_1] \times [u_2, r_2]$ when u dominates r and \emptyset otherwise, 124 see the righthand plot of Figure 1. Note that the \mathcal{D}_u^r are not disjoints.

We use the hypervolume to characterize the convergence of a set S of objective vectors to the entire Pareto front. For a fixed valid reference point r, a set S is said to converge to the Pareto front when the hypervolume difference $HV_r(PF_f) - HV_r(S)$ converges to 0. We define the *optimality gap* of S with respect to a valid reference point r as $HV_r(PF_f) - HV_r(S)$.

Another quantity of interest is how much adding a vector to a set affects its hypervolume. The hypervolume improvement with respect to r of the vector u to the set S is $HVI_r(u, S) = HV_r(S \cup \{u\}) - HV_r(S)$. We also use the term hypervolume improvement to refer to the hypervolume increase of an increasing sequence of sets. More precisely, the hypervolume improvement at iteration n of a monotone sequence $(\mathcal{S}_n)_{n \in \mathbb{N}^*}$ is $HV_r(\mathcal{S}_{n+1}) - HV_r(\mathcal{S}_n)$.

2.2. The ISOOMOO framework, its HV-ISOOMOO instanciation and the 136 associated greedy set sequences. The Incremental SingleObjective Optimization for 137MultiObjective Optimization (ISOOMOO) framework builds incrementally a monotone¹ 138 sequence $(\mathcal{I}_n)_{n \in \mathbb{N}^*}$ of sets of vectors of the objective space. The pseudo code of ISOOMOO 139is given in Algorithm 2.1, where the current value of \mathcal{I}_n is denoted by \mathcal{I} . At each so-140 called meta-iteration, a generic singleobjective maximization algorithm SOOPTIMIZER 141(line 3 in Algorithm 2.1) is run on the criterion $X \in \Omega \subset \mathbb{R}^d \mapsto J(\mathcal{I}, X)$ and the resulting 142 solution is added to \mathcal{I} (line 4 in Algorithm 2.1). We use the term meta-iteration to separate 143between the (meta-)iterations of ISOOMOO and the iterations of SOOPTIMIZER. Since 144 the set \mathcal{I} is composed of the final objective incumbents of previous runs of SOOPTIMIZER 145and (ideally) provides an approximation of the Pareto front, we call it *final incumbents* 146 147 Pareto front approximation.

The singleobjective optimization procedure may vary between meta-iterations. More precisely, the run of SOOPTIMIZER depends on data about precedent runs stored in D (line 4 in Algorithm 2.1). This allows to alternate between various singleobjective optimization algorithms with different features, but also to adapt the initialization. This could be done by storing in D an iteration index or the final search-space incumbents of SOOPTIMIZER runs.

Algorithm 2.1 Incremental SingleObjective Optimization for MultiObjective Optimization (ISOOMOO)

1: while not stopping criterion do

- 2: $Y, d \leftarrow \text{SOOPTIMIZER}(X \mapsto J(\mathcal{I}, X), D)$
- 3: $\mathcal{I} \leftarrow \mathcal{I} \cup \{F(Y)\}$ # update of the approximation of the Pareto front
- 4: $D \leftarrow D \cup \{d\} \#$ update of the data collected
- 5: end while

In this paper, we study HV-ISOOMOO, an instanciation of ISOOMOO for which the criterion $J(\mathcal{I}, .)$ relates to the hypervolume improvement to \mathcal{I} . Formally, HV-ISOOMOO

¹A sequence of set $\{A_n, n \ge 0\}$ is monotone if the following inclusions $A_0 \subset A_1, \ldots \subset A_n \subset \ldots$ hold.

is a class of algorithms derived from ISOOMOO for which the maximization of the criterion J is *compliant* with the maximization of the hypervolume improvement as defined below.

159 Assumption 2.1. (Compliance to hypervolume improvement maximization) The max-160 imization of a criterion J as in ISOOMOO is *compliant* with the maximization of the 161 hypervolume improvement if for any set \mathcal{I} of objective vectors, maximizing $J(\mathcal{I}, .)$ is 162 equivalent to maximizing the hypervolume improvement with respect to a valid reference 163 point r to the set \mathcal{I} :

164 (2.4)
$$\operatorname{argmax}_{X \in \mathbb{R}^d} J(\mathcal{I}, X) = \operatorname{argmax}_{X \in \mathbb{R}^d} \operatorname{HVI}_r(F(X), \mathcal{I})$$

165 We simply refer to an algorithm built from the HV-ISOOMOO framework as an HV-166 ISOOMOO algorithm.

167 DEFINITION 2.2 (HV-ISOOMOO). We define an HV-ISOOMOO algorithm as an 168 ISOOMOO algorithm as described in Algorithm 2.1 where the criterion J satisfies As-169 sumption 2.1.

At each meta-iteration n, an HV-ISOOMOO algorithm seeks a feasible vector maximizing the hypervolume improvement to the final incumbents Pareto front approximation \mathcal{I}_n . Ideally, when n goes to infinity, the non-dominated subset of $(\mathcal{I}_n)_{n\in\mathbb{N}^*}$ converges to the (entire) Pareto front, namely the non-dominated set of feasible objective vectors, which maximizes the hypervolume. In other words, HV-ISOOMOO algorithms try to approximate the Pareto front with a greedy approach.

176 DEFINITION 2.3. We define the convergence of HV-ISOOMOO as the convergence of 177 $HV_r(\mathcal{I}_n)$ towards $HV_r(PF_f)$.

The performance of a specific HV-ISOOMOO algorithm depends crucially on the 178 choice of the criterion J. In this respect, the hypervolume improvement to \mathcal{I} is not a 179good candidate for $J(\mathcal{I}, .)$. Indeed, it is constant equal to zero in the region dominated 180 by \mathcal{I} , which makes it difficult to optimize. A criterion whose maximization is compliant 181 with the maximization of the hypervolume improvement and designed to be easier to 182optimize has already been introduced in [17] under the name uncrowded hypervolume 183 improvement (UHVI). In the non-zero region of the hypervolume improvement to a set \mathcal{I} , 184 namely for F(X) not dominated by \mathcal{I} , the UHVI is simply the hypervolume improvement. 185Otherwise, in the region where the hypervolume improvement is null, the UHVI is negative 186 and equals minus the distance to the empirical non-dominated front of the set \mathcal{I} relative 187to r. It is easy to see that the UHVI satisfies (2.4). 188

189 The choice of SOOPTIMIZER also plays a key role in the performance of an HV-190 ISOOMOO algorithm. In this paper, we analyze HV-ISOOMOO under the assumption 191 of perfect singleobjective optimization formalized below.

192 Assumption 2.4 (Perfect Singleobjective Optimization). At every meta-iteration n, 193 for any final incumbents Pareto front approximation \mathcal{I}_n , the run of SOOPTIMIZER (line 2 194 in Algorithm 2.1) returns $Y \in \operatorname{argmax}_{X \in \Omega} J(\mathcal{I}_n, F(X))$.

The assumption of perfect singleobjective optimization is reminiscent to the assumption of perfect line search which is common in the analysis of gradient based methods [11]. Under this assumption, all choices of criterions verifying Assumption 2.1 are equivalent. The convergence of HV-ISOOMOO under perfect singleobjective optimization is a necessary condition for the soundness of the approach. Additionally, we could obtain a lower bound on the convergence-speed of a real instanciation of HV-ISOOMOO by combining
such lower bounds on the convergence-speed of HV-ISOOMOO under Assumption 2.4
with existing lower bounds on the convergence speed of singleobjective optimization algorithms.

We introduce below the notions of greedy sequence and greedy set sequence, which are mathematical abstractions of HV-ISOOMOO under Assumption 2.4 of perfect singleobjective optimization.

DEFINITION 2.5 (Greedy sequence and greedy set sequence). Given a valid reference point r, we define as greedy sequence, a sequence $\{v_n, n \ge 1\}$ satisfying

209 (2.5)
$$v_1 \in \arg \max_{v \in V(v)} HV_r(v)$$
 and

210 (2.6) $v_{n+1} \in \arg \max_{v \in F(\Omega)} HV_r(\{v_1, \cdots, v_n, v\}) \text{ for all } n \ge 1$.

The greedy set sequence $(S_n)_{n \in \mathbb{N}^*}$ associated to the greedy sequence $\{v_n, n \ge 1\}$ is composed of the greedy sets $S_n := \{v_k, k \le n\}$.

There is a bijection between greedy sequences and greedy set sequences. The *n*-th element of the greedy sequence $\{v_n, n \ge 1\}$ associated to a greedy set sequence $(S_n)_{n \in \mathbb{N}^*}$ is simply the unique element of $S_n \setminus S_{n-1}$ if n > 1 and of S_1 if n = 1.

The recurrence relation of the greedy sequence (2.6) is equivalent to v_{n+1} belonging to arg max_{$v \in F(\Omega)$} HVI_{$r}(<math>v, S_n$) for all $n \ge 1$. It is immediate to see that under Assumption 2.4, the final incumbents generated by HV-ISOOMOO constitute *a* greedy sequence while the final incumbents Pareto front approximations form the associated greedy set sequence (\mathcal{I}_n)_{$n \in \mathbb{N}^*$}. The indices *n* of both greedy and greedy set sequences iterations correspond to HV-ISOOMOO meta-iterations. In this paper, we derive convergence results for greedy set sequences, which transfer to HV-ISOOMOO under Assumption 2.4.</sub>

As we will see in Subsection 3.1, the problem of maximizing the hypervolume improvement to a fixed set can be rewritten as the maximum of a finite number of hypervolume maximization problems. Therefore, we can infer from [3, Theorem 1] that as soon as the Pareto front is lower semi-continuous, there exists a greedy sequence and the associated greedy set sequence.

PROPOSITION 2.6. If the Pareto front is described by a lower semi-continuous function f, then there exists a greedy sequence $\{v_n, n \ge 1\}$ associated to any valid reference point r.

Proof. If f is lower semi-continuous, then for any reference point r, the maximum of HV_r(.) exists, see [3, Theorem 1]. Therefore, there exists a vector verifying (2.5) and the problem of maximizing the maximum of a finite number of hypervolume functions defined in (3.4) admits a solution. Since the recurrence equation defining greedy sequences (2.6) is equivalent to (3.4) by Lemma 3.5, a lemma proven in the next section, we can build a sequence $\{v_n, n \ge 1\}$ verifying (2.5) and (2.6), namely a greedy sequence.

Additionally, since the hypervolume indicator associated to a valid reference point is strictly Pareto-compliant (see [14]), this sequence is composed of vectors of the Pareto front.

PROPOSITION 2.7. If the Pareto front is described by a lower semi-continuous function f, then any vector of a greedy sequence associated to a valid reference point r belongs to the Pareto front. Consequently, for such Pareto front and reference point and under Assumption 2.4 of perfect singleobjective optimization, all final incumbents Pareto front approximations \mathcal{I}_n of HV-ISOOMOO algorithms relative to r are subsets of the Pareto front.

247 Proof. Since for any valid reference point r, $HV_r(.)$ is strictly Pareto-compliant [14], 248 its maximum always belongs to the set of non-dominated feasible objective vectors, that 249 is the Pareto front. Thus, in particular, a vector v_1 verifying (2.5) belongs to the Pareto 250 front. Additionally, by Lemma 3.5, a lemma proven in the next section, every solution 251 of the update equation of greedy sequences (2.6) verifies (3.4). As a consequence, such 252 vectors are solution of at least one hypervolume maximization problem, and thus also 253 belong to the Pareto front.

Yet, in general, there exists more than one greedy sequence, and thus greedy set sequence. 254255For example, there are infinitely many greedy sequences associated to any affine Pareto 256front with a reference point dominating the nadir point. This statement relies on the fact that the unique maximizer of the hypervolume relative to a reference point r dominating 257the nadir point is the middle of the section of the Pareto front dominating r, see [2, 258Theorem 5]. As a consequence, the middle of the section of the Pareto front dominating r259is the only candidate for v_1 but v_2 can be either at 1/4 or at 3/4 of this section. Similarly, 260 261 v_3 has to be in the position where v_2 is not but v_4 can be at 1/8, 3/8, 5/8 or 7/8 of the section of the Pareto front dominating r. For any n, we can find an iteration m such 262that v_m can be placed at 2^n different points, whatever the m-1 first terms of the greedy 263sequence are. 264

2.3. Assumptions on the Pareto front and the objective functions. We 265present and discuss here the assumptions on the function f describing the Pareto front 266 under which we derive convergence results. We typically assume that the function f is 267268 bilipschitz, convex or simultaneously bilipschitz and with a Hölder continuous derivative, namely $\mathcal{C}^{1,\alpha}$. Under any of these three assumptions, f is continuous. For the sake of 269conciseness, we transfer the properties of f to the Pareto front. For example, we call 270convex Pareto front a Pareto front described by a convex function. We recall that a 271function q is Hölder continuous with exponent α when there exists H > 0 such that 272 $|g(x) - g(y)| \le H \times |x - y|^{\alpha}$ for all x, y [12]. We note $[g]_{\alpha}$ the minimum Hölder coefficient 273 of g with respect to the exponent α , that is $[g]_{\alpha} := \sup_{x \neq y} \frac{|g(x) - g(y)|}{|x - y|^{\alpha}}$. When needed, 274we detail the bilipschitz constants and say that a bilipschitz function f is (L_{\min}, L_{\max}) -275bilipschitz if for all $x, y \in [x_{\min}, x_{\max}]$, we have $L_{\min} \times |x-y| \le |f(x) - f(y)| \le L_{\max} \times |x-y|$ 276where $L_{\text{max}} \ge L_{\text{min}} > 0$. We also consider *affine Pareto fronts*, the simplest kind of Pareto 277front. As they form a line in the biobjective case, they are usually referred to as linear 278Pareto fronts. They provide good examples to illustrate a point and help to understand 279280the results we prove on the asymptotic convergence behavior.

We remind below sufficient conditions on the search-space and on the objective functions which guarantee that f is convex and bilipschitz.

PROPOSITION 2.8. Given a biobjective minimization problem as in (2.1) whose Pareto front is described by a function f. If F_1 and F_2 are respectively $(L_{\min,1}, L_{\max,1})$ -bilipschitz and $(L_{\min,2}, L_{\max,2})$ -bilipschitz, then f is $(\frac{L_{\min,2}}{L_{\max,1}}, \frac{L_{\max,2}}{L_{\min,1}})$ -bilipschitz.

286 PROPOSITION 2.9. Given a biobjective minimization problem as in (2.1) whose Pareto 287 front is described by a function f. If the search space Ω and the objective functions F_1

288 and F_2 are convex, then f is convex.

The proofs of both propositions can be found for instance in [15]. The conditions on F_1 , F_2 and Ω are sufficient but non-necessary conditions. Indeed, adding small discontinuity in the objective functions far from the Pareto set makes them non-convex and non-bilipschitz without modifying the Pareto front.

Representing F_1 values on the absciss and F_2 values on the ordinate instead of the 293converse is an arbitrary choice. When f is a bijection, had we chosen to represent the 294 F_2 values on the absciss instead of on the ordinate, we would have had another repre-295sentation of the Pareto front : $\{(y, f^{-1}(y)) : y \in [f(x_{\max}); f(x_{\min})]\}$. If so, the inverse 296 function f^{-1} would have played the role of f. It is interesting to notice that the choice 297of the objective function represented on the horizontal axis does not impact whether the 298function characterizing the Pareto front is bilipschitz or convex. Indeed, f being bilip-299 schitz is equivalent to both f and f^{-1} being lipschitz. Additionally, we can prove that 300 given that the function f is decreasing, f being convex is equivalent to its inverse f^{-1} 301 being convex. The proof of this property is straightforward. If f is convex, then for all $x, y \in \mathbb{R}$, $f\left(\frac{f^{-1}(x)+f^{-1}(y)}{2}\right)$ is smaller than $\frac{f(f^{-1}(x))+f(f^{-1}(y))}{2}$, that is $\frac{x+y}{2}$. Since f and 302 303 therefore f^{-1} are decreasing, by composing by f^{-1} each side of the inequality, we obtain 304 a characterization of the convexity of f^{-1} : for all $x, y \in \mathbb{R}$, $\frac{f^{-1}(x)+f^{-1}(y)}{2}$ is higher than 305 $f^{-1}(\frac{x+y}{2}).$ 306

307 3. Preliminary results. In this section, we present preliminary results which are 308 crucial for the analysis of the convergence of HV-ISOOMOO. While they are here exposed 309 as tools for convergence analysis, they are also interesting for their own sake.

3.1. Decomposition of the optimality gap using gap regions. We introduced \mathcal{D}_{S}^{r} in (2.3) as the region of the objective space dominating r and weakly dominated by S. Its Lebesgue measure is $HV_{r}(S)$. We now introduce *total gap regions*, whose Lebesgue 313 measure are optimality gaps.

314 DEFINITION 3.1. The total gap region of S with respect to a fixed valid reference 315 point r, \mathcal{G}_{S}^{r} , is defined as the region of the objective-space which dominates r and is 316 weakly dominated by PF_{f} but not by S, namely $\mathcal{D}_{PF_{f}}^{r} \setminus \mathcal{D}_{S}^{r}$.

When S is a subset of the Pareto front dominating the reference point r, the total gap region has a particular shape which can be visualized in the rightmost plot of Figure 1. The total gap region \mathcal{G}_{S}^{r} can be decomposed into the disjoint union of |S| + 1 sets of the form $\mathcal{D}_{S'}^{r'}$ that are formally defined below.

321 DEFINITION 3.2 (Gap regions, gaps and associated reference points). Let S =322 { $v_1, ..., v_n$ } be a set of n distinct vectors of the Pareto front dominating a valid refer-323 ence point r. Let σ be the permutation ordering the v_i by increasing F_1 values: $v_{\sigma(1),1} <$ 324 $v_{\sigma(2),1} < ... < v_{\sigma(n),1}$.

• For all $i \in [\![1, n+1]\!]$, the *i*-th gap region of the set S, $\mathcal{G}_{S,i}^r$, is the set $\mathcal{D}_{PF_f}^{r_i}$ with the associated reference points r_i being

327 (3.1)
$$r_1 = (v_{\sigma(1),1}, r_2), r_{n+1} = (r_1, v_{\sigma(n),2}) \text{ and} r_i = (v_{\sigma(i),1}, v_{\sigma(i-1),2}) \text{ for all } i \in [\![2,n]\!] .$$

325 326

• We refer to $\mathcal{G}_{S,1}^r$ and $\mathcal{G}_{S,n+1}^r$ as the left and the right extreme gap region of S, respectively. The left and the right extreme gap regions are empty when the left and the right extreme vectors of the Pareto front belong to S. Non-extreme gap regions are never empty. The total gap region is the disjoint union of the gap regions: $\mathcal{G}_{S_n}^r = \bigcup_{i=1}^{n+1} \mathcal{G}_{S_n,i}^r$. This decomposition of the total gap region, and thus of the optimality gap, is the cornerstone of the convergence analysis. The area of a gap region $\mathcal{G}_{S_n,i}^r$ is the hypervolume of the Pareto front with respect to the reference point r_i^n . As a consequence, we can write the optimality gap as the sum of n + 1 hypervolumes of the Pareto front with respect to different reference points.

338 LEMMA 3.3. At any iteration n, the optimality gap of a greedy set sequence with 339 respect to a valid reference point can be decomposed as the sum of n + 1 hypervolumes of 340 the Pareto front:

341 (3.2)
$$HV_r(PF_f) - HV_r(\mathcal{S}_n) = \sum_{i=1}^{n+1} HV_{r_i^n}(PF_f) .$$

Proof. The optimality gap at iteration n is the Lebesgue measure of the total gap region $\mathcal{G}_{\mathcal{S}_n}^r$, which is the disjoint union of the gap regions $\mathcal{G}_{\mathcal{S}_n,i}^r$ of \mathcal{S}_n . Since each gap region $\mathcal{G}_{\mathcal{S}_n,i}^r$ equals $\mathcal{D}_{\mathrm{PF}_f}^{r_i^n}$, the optimality gap is equal to $\sum_{i=1}^{n+1} \mathrm{HV}_{r_i^n}(\mathrm{PF}_f)$.

Additionally, we can express the hypervolume improvement of any vector to S_n as an hypervolume. It is immediate for vectors which do not dominate S_n . For other vectors, the reference point depends on the gap region to which the vector belongs.

LEMMA 3.4. Let $(S_n)_{n \in \mathbb{N}^*}$ be a greedy set sequence relative to a valid reference point r. At any iteration n, the hypervolume improvement to S_n of any u belonging to the *i*-th gap region of S_n , $\mathcal{G}_{S_n,i}^r$, satisfies

$$352 (3.3) HVI_r(u, \mathcal{S}_n) = HV_{r_i^n}(u)$$

Proof. The hypervolume improvement of any $u \in \mathcal{G}_{\mathcal{S}_n,i}^r$ is the Lebesgue-measure of the intersection between $\mathcal{G}_{\mathcal{S}_n,i}^r$ and \mathcal{D}_u^r . Therefore, it is equal to $\lambda(\mathcal{D}_u^{r_i^n})$, that is $\mathrm{HV}_{r_i^n}(u)$. We can now reformulate the recurrence relation defining a greedy sequence at iteration n + 1. Indeed, picking a vector maximizing the hypervolume improvement to \mathcal{S}_n is equivalent to pick a vector where the highest value of the maximum of the hypervolumes with respect to the r_i^n is reached.

LEMMA 3.5. At any iteration n, the recurrence formula satisfied by v_{n+1} , i.e. (2.6), can be reformulated as

$$\begin{array}{ll} 362 \\ 363 \end{array} (3.4) \qquad v_{n+1} \in \arg\max_{u \in PF_f} \max_{i \in [\![1,n+1]\!]} HV_{r_i^n}(u). \end{array}$$

364 Proof. The hypervolume improvement of any vector u to S_n is $\max_{i \in [\![1,n+1]\!]} \operatorname{HV}_{r_i^n}(u)$. 365 It is a consequence of Lemma 3.4 and of the fact that the hypervolume with respect to r_i^n 366 is null outside the *i*-th gap region of S_n . Additionally, v_{n+1} belongs to the Pareto front 367 by Proposition 2.7. Thus, (2.6) is equivalent to (3.4).

Similarly, we can express the decrease of the optimality gap at iteration n + 1, HV_r(S_{n+1}) – HV_r(S_n), as the maximum of n + 1 hypervolume maximization problems. 370 LEMMA 3.6. Let $(S_n)_{n \in \mathbb{N}^*}$ be a greedy set sequence relative to a valid reference point 371 r. The hypervolume improvement at iteration n + 1 equals

372 (3.5)
$$HV_r(\mathcal{S}_{n+1}) - HV_r(\mathcal{S}_n) = \max_{u \in PF_f} \max_{i \in [\![1,n+1]\!]} HV_{r_i^n}(u) .$$

Proof. The hypervolume improvement $\operatorname{HV}_r(\mathcal{S}_{n+1}) - \operatorname{HV}_r(\mathcal{S}_n)$ is the hypervolume improvement of v_{n+1} to \mathcal{S}_n . With the same arguments as in the proof of Lemma 3.5, we can prove that it equals $\max_{u \in \operatorname{PF}_f} \max_{i \in [1, n+1]} \operatorname{HV}_{r_i}^n(u)$.

377 3.2. Lower bound of the normalized maximum hypervolume for convex **Pareto fronts.** In this section and the next one, we provide bounds on the maximum hypervolume achievable by a single feasible vector normalized by the maximum hypervolume of a feasible set. We refer to this ratio as the *normalized maximum hypervolume* with respect to r:

$$\frac{\max_{u \in \mathrm{PF}_f} \mathrm{HV}_r(u)}{\mathrm{HV}_r(\mathrm{PF}_f)}$$

Bounds on the normalized maximum hypervolume are exploited in Section 4 to provide bounds on the speed of convergence of the greedy set sequence towards the Pareto front. The hypervolume relative to a reference point $r = (r_1, r_2)$ of a vector u = (x, f(x)) of the Pareto front is $HV_r(u) = (r_1 - x) \times (r_2 - f(x))$. From this simple formula, we derive in the next proposition necessary conditions for a vector of the Pareto front $u^* = (x^*, f(x^*))$ to be an hypervolume maximizer when f has at least left and right derivatives in x^* .

390 PROPOSITION 3.7. Let $x^* \in]x_{\min}, x_{\max}[$ such that $u^* := (x^*, f(x^*))$ maximizes the 391 hypervolume with respect to a valid reference point $r = (r_1, r_2)$. If the function f de-392 scribing the Pareto front admits left and right derivatives in x^* , respectively $f'_-(x^*)$ and 393 $f'_+(x^*)$, then

$$\begin{array}{c} 394 \\ 395 \end{array} (3.7) \qquad \qquad -f'_+(x^*) \le \frac{r_2 - f(x^*)}{r_1 - x^*} \le -f'_-(x^*) \ . \end{array}$$

³⁹⁶ Proof. We define the function $HV_{x,r}(.)$ as $x \mapsto HV_r((x, f(x)))$. If x^* maximizes ³⁹⁷ $HV_{x,r}(.)$, then the left and the right derivatives of $HV_{x,r}(.)$ are positive and negative, ³⁹⁸ respectively. By replacing the left and right derivatives of $HV_{x,r}(.)$ by their explicit ³⁹⁹ formulas and reorganizing the terms we obtain (3.7).

Equation (3.7) states that the slope of the diagonal of the rectangle $\mathcal{D}_{u^*}^r$ is between the absolute values of the slopes of the right and the left derivatives of f at x^* , see the middle plot of Figure 2. To the best of our knowledge, this geometric interpretation is new. It becomes simpler when f is differentiable. As soon as u^* is a non-extreme vector, the absolute value of the slope of the tangent of the front at u^* is equal to the slope of the diagonal of the rectangle $\mathcal{D}_{u^*}^r$, see the lefthand plot of Figure 2.

406 COROLLARY 3.8. Let $x^* \in]x_{\min}, x_{\max}[$ be such that $u^* := (x^*, f(x^*))$ maximizes the 407 hypervolume with respect to a valid reference point $r = (r_1, r_2)$. If the Pareto front is 408 described by a differentiable function f in x^* , then $f'(x^*)$ satisfies

409 (3.8)
410
$$-f'(x^*) = \frac{r_2 - f(x^*)}{r_1 - x^*}$$

10

412 A convex function may not be differentiable, but it always has left and right deriva-413 tives. It is also above its left and right tangent lines respectively on the left and on the right 414 of x^* . Therefore, (3.7) implies that the affine function $g: x \mapsto f(x^*) - \frac{r_2 - f(x^*)}{r_1 - x^*} \times (x - x^*)$ 415 is a minorant of f, and thus that the hypervolume of PF_f is smaller than the hypervol-416 ume of $\operatorname{PF}_g := \{g(x) : x \in [x_{\min}, x_{\max}]\}$. This upper bound on $\operatorname{HV}_r(\operatorname{PF}_f)$ involves the 417 lengths of the rectangle $\mathcal{D}_{u^*}^r$, whose area is $\operatorname{HV}_r(u^*)$. It is the key idea of the proof of the 418 following lower bound on the normalized maximum hypervolume.

419 PROPOSITION 3.9. If the Pareto front is described by a convex function f, then the 420 following lower bound on the normalized maximum hypervolume with respect to any valid 421 reference point r holds:

422 (3.9)
$$\frac{\max_{u \in PF_f} HV_r(u)}{HV_r(PF_f)} \ge \frac{1}{2}$$

where the inequality is an equality if and only if the Pareto front is affine and r dominates the nadir point.

426 Proof. The function f being convex, it has left and right derivatives and thus, so does 427 $\operatorname{HV}_{x,r}(.): x \mapsto \operatorname{HV}_r((x, f(x)))$. Thus, by Proposition 3.7, (3.7) holds. Since f is convex, 428 (3.7) implies that the affine function $g: x \mapsto f(x^*) - \frac{r_2 - f(x^*)}{r_1 - x^*} \times (x - x^*)$ is a minorant 429 of f. Therefore, $\operatorname{PF}_g := \{g(x): x \in [x_{\min}, x_{\max}]\}$ dominates PF_f , and thus has a higher 430 hypervolume.

431 We note $L_1 := r_1 - \tilde{x}_{\min,r}$ and $L_2 := r_2 - f(\tilde{x}_{\max,r})$ the lengths of the rectangle 432 $\mathcal{R} := [\tilde{x}_{\min,r}, r_1] \times [f(\tilde{x}_{\max,r}), r_2]$. We note $l_1 := r_1 - x^*$ and $l_2 := r_2 - f(x^*)$ the lengths 433 of the rectangle $\mathcal{D}_{u^*}^r$. The region of \mathcal{R} which dominates PF_g is a right-angled triangle. 434 Additionally, by definition, the slope of its hypotenuse is $\frac{l_2}{l_1}$, and thus the lengths of the 435 other sides are $L_1 - l_1 + (L_2 - l_2) \times \frac{l_1}{l_2}$ and $L_2 - l_2 + (L_1 - l_1) \times \frac{l_2}{l_1}$ (see the middle plot 436 of Figure 2). Therefore

437
$$\operatorname{HV}_{r}(\operatorname{PF}_{g}) = \lambda(\mathcal{R}) - \lambda(\{u \in \mathbb{R}^{2} : u \in \mathcal{R}, u \leq \operatorname{PF}_{g}\})$$

438
$$= L_{1}L_{2} - \frac{1}{2} \times (L_{1} - l_{1} + (L_{2} - l_{2}) \times \frac{l_{1}}{l_{2}}) \times (L_{2} - l_{2} + (L_{1} - l_{1}) \times \frac{l_{2}}{l_{1}})$$

439
440
$$= l_1 l_2 \times \left[-2 + 2 \times \frac{L_2}{l_2} - \frac{1}{2} \times \left(\frac{L_2}{l_2}\right)^2 + 2 \times \frac{L_1}{l_1} - \frac{1}{2} \times \left(\frac{L_1}{l_1}\right)^2 \right] .$$

For all x, we have $(x-2)^2 \ge 0$ and thus $2x - \frac{1}{2}x^2 \le 2$. Therefore, we can conclude 441 that $HV_r(PF_q)$, and thus $HV_r(PF_f)$ is lower than $2 \times l_1 l_2$, that is $2 \times HV_r(u^*)$. If either 442 $L_1/l_1 \neq 2$ or $L_2/l_2 \neq 2$, the inequality is strict. Thus, when the inequality is an equality, 443 the center of \mathcal{R} belongs to the Pareto front. Since f is convex, it requires from f to 444 be affine and from the reference point r to dominate the nadir point $(x_{\max}, f(x_{\min}))$. 445 Conversely, if the Pareto front is affine and the reference point r dominates the nadir 446 point, we know that the optimum is in the middle of the Pareto front and that we have 447 the equality (see [2, Theorem 5]). 448 Π

We just proved that one half is a tight lower bound on the normalized maximum hypervolume for convex Pareto fronts. However, except for the trivial upper bound 1, there is no upper bound valid for every convex Pareto front, even when r dominates the



Fig. 2: Left and middle: Two convex Pareto fronts and their respective hypervolume maximizers u^* , one differentiable (left) and one non-differentiable (middle). The slopes of the two dotted lines, namely PF_g and the diagonal of $\mathcal{D}_{u^*}^r$, are equal. Right: The Pareto front PF_{ϵ} and the hypervolume maximizer u^*_{ϵ} for $\epsilon = 1/4$ and r = (1, 1).

adir point. Here is a simple example which illustrates this. Let consider the convex Pareto front $PF_{\epsilon} := \{\max(1 - \frac{x}{\epsilon}, \epsilon - \epsilon \times x) : x \in [0, 1]\}$ represented in the righthand plot of Figure 2 and the reference point r = (1, 1). When $\epsilon \leq 1$, PF_{ϵ} is convex and (3.8) implies that $u_{\epsilon}^* = (\epsilon \times (1 - \epsilon), \epsilon \times (1 - \epsilon))$ is the unique hypervolume minimizer. Thus, the normalized maximum hypervolume of PF_{ϵ} for this reference point is equal to $\frac{(1-\epsilon+\epsilon^2)^2}{1-\epsilon \times (1-\epsilon)^2+(\epsilon-\epsilon^2)^2}$ and converges to 1 when ϵ goes to 0.

458 **3.3. Lower and upper bounds of the normalized maximum hypervolume**459 **for bilipschitz Pareto fronts.** In this section, we examine lower and upper bounds on
460 the normalized maximum hypervolume in the case of bilipschitz Pareto fronts.

461 We consider two affine fronts with the same left extreme vector as PF_f and slopes 462 $-L_{\min}$ and $-L_{\max}$, see Figure 3. We call them PF_{\min} and PF_{\max} , respectively. Formally:

463 (3.10)
$$PF_{\max} := \{ (x, f_{\max}(x)) : x \in [x_{\min}, x_{\max}] \}$$

464 (3.11)
$$\operatorname{PF}_{\min} := \{ (x, f_{\min}(x) : x \in [x_{\min}, x_{\max}] \}$$

466 with $f_{\min}(x) = f(x_{\min}) - (x - x_{\min}) \times L_{\min}$ and $f_{\max}(x) = f(x_{\min}) - (x - x_{\min}) \times L_{\min}$ 467 L_{\max} . For a (L_{\min}, L_{\max}) -bilipschitz function f, $f_{\min}(x) \leq f(x) \leq f_{\max}(x)$ for $x \in$ 468 $[x_{\min}, x_{\max}]$, and thus the Pareto front is dominated by PF_{\max} and dominates PF_{\min} . 469 These two affine fronts provide bounds on both the hypervolume of the Pareto front 470 and the highest hypervolume of a vector on the Pareto front. They are key to prove 471 the following lower bound on the normalized maximum hypervolume of a (L_{\min}, L_{\max}) -472 bilipschitz Pareto front.

473 PROPOSITION 3.10. If the Pareto front is described by a (L_{\min}, L_{\max}) -bilipschitz func-474 tion f, then the following bound on the normalized maximum hypervolume with respect to 475 any valid reference point r holds

476 (3.12)
$$\frac{\max_{u \in PF_f} HV_r(u)}{HV_r(PF_f)} \ge \frac{1}{2} \times \frac{L_{\min}}{L_{\max}}$$

478 Proof. The fronts PF_{max} and PF_{min} are defined respectively in (3.10) and (3.11). 479 We note $\Delta_1 := \tilde{x}_{max,r} - \tilde{x}_{min,r}, \ \Delta'_1 := r_1 - \tilde{x}_{max,r}, \ \Delta_2 := r_2 - f(\tilde{x}_{min,r})$ and V :=



Fig. 3: The Pareto front PF_f surrounded by PF_{max} and PF_{min} in the case where the reference point r does not dominate the nadir point (left) and in the case where it does (right).

 $\Delta_2 \times (r_1 - \tilde{x}_{\min,r})$, see the lefthand plot of Figure 3. Since the front PF_{max} dominates 480 the Pareto front, the hypervolume of PF_f is smaller than the hypervolume of PF_{max} , 481 $V + L_{\max} \times \Delta_1 \times \Delta'_1 + \frac{1}{2} \times L_{\max} \times \Delta_1^2$. Since each vector of PF_{\min} is dominated by a 482 vector of PF_f , the maximum hypervolume of a vector of PF_f is higher than the maximum 483 hypervolume of a vector of PF_{min} . The front PF_{min} being an affine and therefore convex 484 front, we know by Proposition 3.9 that the maximum hypervolume of a vector of PF_{min} is 485higher than half of $HV_r(PF_{min})$, which is equal to $\frac{1}{2} \times (V + L_{min} \times \Delta_1 \times \Delta'_1 + \frac{1}{2} \times L_{min} \times \Delta_1^2)$. 486To summarize, the maximum hypervolume of a vector of PF_f is higher than $\frac{1}{2} \times (V + L_{\min} \times$ 487 $\Delta_1 \times \Delta'_1 + \frac{1}{2} \times L_{\min} \times \Delta_1^2$). Combining the upper bound on the hypervolume of PF_f and the 488 lower bound on the maximum hypervolume of a vector of PF_{f} , the normalized maximum 489 hypervolume is higher than $\frac{\frac{1}{2} \times (V + L_{\min} \times \Delta_1 \times \Delta'_1 + \frac{1}{2} \times L_{\min} \times \Delta_1^2)}{V + L_{\max} \times \Delta_1 \times \Delta'_1 + \frac{1}{2} \times L_{\max} \times \Delta_1^2}$. This quantity is itself larger than $\frac{1}{2} \times \frac{L_{\min} \times \Delta_1 \times \Delta'_1 + \frac{1}{2} \times L_{\min} \times \Delta_1^2}{L_{\max} \times \Delta_1 \times \Delta'_1 + \frac{1}{2} \times L_{\max} \times \Delta_1^2}$. As $V \ge 0$ and $0 < \frac{L_{\max} \times \Delta_1 \times \Delta'_1 + \frac{1}{2} \times L_{\max} \times \Delta_1^2}{L_{\max} \times \Delta_1 \times \Delta'_1 + \frac{1}{2} \times L_{\max} \times \Delta_1^2} < 1$, we conclude that the normalized maximum hypervolume is higher than $\frac{1}{2} \times \frac{L_{\min}}{L_{\max}}$. 490 491 492 We cannot guarantee any upper bound strictly inferior to 1 on the normalized maximum 493

hypervolume without adding an assumption on the reference point. Indeed, for a given bounded Pareto front, it is easy to show that the normalized maximum hypervolume goes to 1 when $r \to \infty$. However, if f is (L_{\min}, L_{\max}) -bilipschitz and r dominates the nadir point, we can prove that the normalized maximum hypervolume is higher than $\frac{1}{2} \times \frac{L_{\max}}{L_{\min}}$. The proof relies on the fact that if the reference point dominates the nadir point, the vector of an affine front with the highest hypervolume is its middle (see [2, Theorem 5]), whose hypervolume is half of the hypervolume of the entire front.

501 PROPOSITION 3.11. If the Pareto front is described by a (L_{\min}, L_{\max}) -bilipschitz func-502 tion f and the reference point r is valid and dominates the nadir point, the following 503 upper-bound on the normalized maximum hypervolume with respect to r holds

$$\frac{\max_{u \in PF_f} HV_r(u)}{HV_r(PF_f)} \le \frac{1}{2} \times \frac{L_{\max}}{L_{\min}}$$

Proof. We use the same notations as in the proof of Proposition 3.10. Since r domi-506nates the nadir point, both Δ'_1 , Δ_2 and V equal 0, and thus the hypervolumes of PF_{max} 507and PF_{min} equal $\frac{1}{2} \times L_{max} \times \Delta_1^2$ and $\frac{1}{2} \times L_{min} \times \Delta_1^2$, respectively. The domination of 508 PF_{min} by PF_f implies that the hypervolume of the Pareto front is below $\frac{1}{2} \times L_{min} \times \Delta_1^2$. 509 Since PF_{max} is an affine front whose extremes dominate r, its middle is the unique hyper-510volume maximizer (see [1, Theorem 5]) with an hypervolume equal to $\frac{1}{4} \times L_{\max} \times \Delta_1$. The 511domination of PF_f by PF_{max} implies that the maximum hypervolume of a vector of PF_f is higher than $\frac{1}{4} \times L_{max} \times \Delta_1^2$. Gathering the lower bound on $HV_r(PF_f)$ and the upper 512513bound on the maximum hypervolume of a vector of PF_f , we retrieve (3.13). 514

515 This upper bound is only relevant for $L_{\text{max}}/L_{\text{min}} < 2$. The bound is the tightest for 516 $L_{\text{max}} = L_{\text{min}}$, where it achieves the value 1/2. In this paper, we use this upper bound for 517 $L_{\text{max}}/L_{\text{min}}$ close to 1 to analyze the asymptotic convergence behavior of HV-ISOOMOO.

4. Convergence of HV-ISOOMOO under perfect singleobjective optimization. We prove in this section various convergence results for HV-ISOOMOO algorithms under the assumption of perfect singleobjective optimization. We first prove that when the Pareto front is either convex or bilipschitz, these algorithms converge to the entire Pareto front with guaranteed convergence-speed. We transform the bounds on the normalized maximum hypervolume proven in Section 3 into lower bounds on the convergence speed. Second, we analyze the asymptotic convergence behavior when the Pareto front is bilipschitz with a Hölder continuous derivatives.

To analyze the decrease of the optimality gap with respect to n, we need to be able to 526track in which gap regions the vectors of the greedy sequence are inserted over multiple 527 iterations. Naturally, a gap region of \mathcal{S}_n persists in being a gap region in the following 528 iterations, as long as no greedy vector is added in this specific gap region. The greedy 529vector v_{n+1} is said to fill the gap region of S_n to which it belongs. At iteration n+1, 530this gap region disappears, replaced by two gap regions that we call its *children*. More 531532 generally, we say that a gap region is a *descendant* of another gap region when it is a proper subset of this gap region. 533

4.1. Convergence of HV-ISOOMOO with guaranteed convergence speed. We prove some upper bounds on the relation between the optimality gap at iteration 2n + 1 and at iteration n. These bounds translate into lower bounds on the speed of convergence of HV-ISOOMOO with perfect singleobjective optimization. The proof relies on inequalities of the form

539 (4.1)
$$\max_{u \in \mathrm{PF}_f} \mathrm{HV}_{r'}(u) \ge C \times \mathrm{HV}_{r'}(\mathrm{PF}_f)$$

stated in Propositions 3.9 and 3.10 and on the reformulation of optimality gaps, areas of gap regions and hypervolume improvement done in Subsection 3.1. A consequence of (4.1) being true for any reference point r' is that the optimality gap at iteration 2n + 1is at most (1 - C) times the optimality gap at iteration n.

We sketch the proof idea in the simple case where each of the v_k ($k \in [[n+1, 2n+1]]$) is inserted in a distinct gap region of S_n , see the lefthand plot of Figure 4. Inserting v_k in a gap region leads to an hypervolume improvement larger than C times the area of this gap region by (4.1). Thus, the hypervolume improvement from iteration n to 2n + 1 is larger than C times the area of the union of all gap regions of S_n , namely the optimality gap at iteration n. A detailed proof is presented after the theorem statement.



Fig. 4: Left: A Pareto front where each of the gap regions of S_3 is filled by one of the greedy vectors v_k for $k \in [\![4,7]\!]$. It is described by $f(x) = 1 - \sqrt{x}$ for $x \in [0,1]$. We represent the region $\mathcal{D}_{S_3}^r \square$, the regions corresponding to $\operatorname{HVI}_r(v_k, \mathcal{S}_{k-1})$ for $k \in [\![4,7]\!]$ \square and the gap regions of $\mathcal{S}_3 \square$. Right: The ordered greedy set along F_1 -values $w_{i,r}^n$ corresponding to the greedy set \mathcal{S}_3 . The Pareto front is described by $f(x) = \frac{e}{e-1} \times e^{-x} + 1 - \frac{e}{e-1}$ for $x \in [0,1]$.

551 PROPOSITION 4.1. Consider a biobjective optimization problem where the Pareto front 552 is described by a function f. Any greedy set sequence $(S_n)_{n \in \mathbb{N}^*}$ associated to a valid ref-553 erence point r satisfies for all n

554 (4.2)
$$\frac{HV_r(PF_f) - HV_r(\mathcal{S}_{2n+1})}{HV_r(PF_f) - HV_r(\mathcal{S}_n)} \le 1 - \frac{1}{2} \times \frac{L_{\min}}{L_{\max}} \text{ if } f \text{ is } (L_{\min}, L_{\max}) \text{-bilipschitz and}$$

555 (4.3)
$$\frac{HV_r(PF_f) - HV_r(\mathcal{S}_{2n+1})}{HV_r(PF_f) - HV_r(\mathcal{S}_n)} \le \frac{1}{2} \text{ if } f \text{ is convex.}$$

Proof. Fix $n \ge 1$. We note σ a permutation of [1, n+1] such that $n + \sigma(i)$ is the index of the first greedy vector v_k inserted in $\mathcal{G}_{S_n,i}^r$ when possible. With this choice of σ , 558 the *i*-th gap region of S_n is a gap region of $S_{n+\sigma(i)-1}$. As a consequence, the hypervolume 559improvement to $S_{n+\sigma(i)-1}$ of any vector u belonging to the *i*-th gap region of S_n is equal 560to $HV_{r_i^n}(u)$ by (3.3). The hypervolume improvement of the greedy vector $v_{n+\sigma(i)}$ to 561 $\mathcal{S}_{n+\sigma(i)-1}$ being maximal, it is in particular larger than the one of any vector of $\mathcal{G}^r_{\mathcal{S}_n,i}$, 562the *i*-th gap region of S_n , and thus than $\frac{1}{2} \times \frac{L_{\min}}{L_{\max}} \times \text{HV}_{r_i^n}(\text{PF}_f)$ by Proposition 3.10. In other words, the hypervolume improvement at any iteration $n + \sigma(i)$ is higher than 563 564 $\frac{1}{2} \times \frac{L_{\min}}{L_{\max}} \times HV_{r_i^n}(PF_f)$. By adding these inequations for all $i \in [[1, n+1]]$, we deduce 565that the hypervolume improvement from iteration n to 2n + 1 is larger than $\frac{1}{2} \times \frac{L_{\min}}{L_{\max}} \times$ 566 $\sum_{i=1}^{n+1} \mathrm{HV}_{r_i^n}(\mathrm{PF}_f)$. Since the sum of the $\mathrm{HV}_{r_i^n}(\mathrm{PF}_f)$ is the optimality gap at iteration 567 568n, we have (4.2). If f is convex instead of bilipschitz, we use Proposition 3.9 instead of Proposition 3.10 and obtain (4.3). П 569

570 Since the optimality gaps form a decreasing sequence, such lower bounds on the relation 571 between the optimality gaps at iteration 2n+1 and at iteration n imply that the optimality 572 gap associated to a greedy set sequence converges asymptotically to 0. Equivalently, HV-573 ISOOMOO algorithms converge to the entire Pareto front under Assumption 2.4, as stated 574 formally below. 575 THEOREM 4.2. Consider a biobjective optimization problem with a Pareto front de-576 scribed by a bilipschitz or convex function f.

The hypervolume of a greedy set sequence associated to a valid reference point r converges to the hypervolume of the entire Pareto front, i.e. $HV_r(\mathcal{S}_n) \xrightarrow[n \to \infty]{} HV_r(PF_f)$.

Equivalently, for such Pareto fronts and under Assumption 2.4 of perfect singleobjective optimization, HV-ISOOMOO algorithms associated to a valid reference point r converge to the Pareto front in the sense of Definition 2.3.

From the lower bounds on the relation between the optimality gaps at iteration 2n + 1and at iteration n, we deduce the following upper bounds on the normalized optimality gap at any iteration.

585 COROLLARY 4.3. Consider a biobjective optimization problem with a Pareto front 586 described by a (L_{\min}, L_{\max}) -bilipschitz function. A greedy set sequence $(S_n)_{n \in \mathbb{N}^*}$ associated 587 to a valid reference point r satisfies for all n

$$\frac{1}{588} \quad (4.4) \quad \frac{HV_r(PF_f) - HV_r(\mathcal{S}_n)}{HV_r(PF_f)} \le \left(1 - \frac{1}{2} \times \frac{L_{\min}}{L_{\max}}\right)^{\lfloor \log_2(n+1) \rfloor} \le (2n+2)^{\log_2(1-\frac{1}{2} \times \frac{L_{\min}}{L_{\max}})}$$

590 If the function f is convex, then any greedy set sequence associated to a valid reference 591 point r satisfies for all n

592 (4.5)
$$\frac{HV_r(PF_f) - HV_r(\mathcal{S}_n)}{HV_r(PF_f)} \le \left(\frac{1}{2}\right)^{\lfloor \log_2(n+1) \rfloor} \le \frac{1}{2n+2}$$

Consequently, under Assumption 2.4 of perfect singleobjective optimization, HV-ISOOMOO algorithms relative to a valid reference point r satisfy (4.4) if f is (L_{\min}, L_{\max}) -bilipschitz and (4.5) if f is convex where S_n is replaced by \mathcal{I}_n , the final incumbents Pareto front approximation at iteration n.

Proof. The k-th term of the sequence defined by $u_0 = 1$ and $u_{n+1} = 2 \times u_n + 1$ for all $n \ge 1$ is $2^k - 1$. Thus, (4.2) and (4.3) imply that when f is (L_{\min}, L_{\max}) -bilipschitz or convex, the normalized optimality gap at iteration $2^k - 1$ is inferior to $(1 - C)^k$ with C equal to $\frac{1}{2} \times \frac{L_{\min}}{L_{\max}}$ and $\frac{1}{2}$, respectively. Since the hypervolume of the greedy set increases with n, and thus the optimality gap decreases with n, we deduce the first inequalities in (4.4) and (4.5) via the change of variable $k = \lfloor \log_2(n+1) \rfloor$.

Additionally, for every n, $\lfloor \log_2(n+1) \rfloor$ is smaller than $\log_2(n+1)+1$, that is $\log_2(2n+605-2)$. For every C, $\log_2(2n+2)$ equals $\log_C(2n+2) \times \log_2(C)$, and thus $C^{\log_2(2n+2)}$ equals $(2n+2)^{\log_2(C)}$. Therefore, we can infer that $(2n+2)^{\log_2(C)}$ is an upper bound of the normalized optimality gap with $C := 1 - \frac{1}{2} \times \frac{L_{\min}}{L_{\max}}$ and $C := \frac{1}{2}$ when f is (L_{\min}, L_{\max}) -bilipschitz and convex, respectively.

We focus here on the relation between the optimality gap at iteration n and at iteration 2n + 1. We could similarly examine the relation between the optimality gap at iteration n and at any later iteration. For example, we could prove that if f is (L_{\min}, L_{\max}) bilipschitz, then for all n, for all $k \le n + 1$, $\frac{HV_r(PF_f) - HV_r(S_{n+k})}{HV_r(PF_f) - HV_r(S_n)}$ is lower than $1 - \frac{1}{2} \times \frac{L_{\min}}{2} \times \frac{k}{n+1}$.

613 $\frac{L_{\min}}{L_{\max}} \times \frac{k}{n+1}$. 614 $Sketch \ of \ proof.$ We consider the k gap regions of S_n with the highest areas. The 615 hypervolume improvement from iteration n to n+k is at least $\frac{1}{2} \times \frac{L_{\min}}{L_{\max}}$ times the area of 616 the union of these gap regions, which is at least $\frac{k}{n+1}$ times the optimality gap at iteration 617 n.

4.2. Asymptotical behavior of the convergence of $HV_r(\mathcal{S}_n)$ to $HV_r(PF_f)$. 618In this section, we analyze the asymptotic convergence behavior for a Pareto front de-619 scribed by a bilipschitz function with a Hölder continuous derivative. We prove that, in 620 this case, doubling the number of vectors in the greedy set divides the optimality gap by a 621 factor which converges asymptotically to two as stated in Theorem 4.10. This asymptotic 622 limit corresponds to the case of affine Pareto fronts with a reference point dominating 623 the nadir point. Indeed, for such Pareto fronts and reference points, the optimality gap 624 is always halved when the number of vectors in the greedy set goes from n to 2n + 1, see 625 626 Figure 5.

First, we study the properties of the part of the Pareto front corresponding to a specific gap region of S_n . For all n, let note σ_n the permutation of $[\![1, n]\!]$ which orders the vectors of S_n by increasing F_1 -values and the so-called *ordered greedy set along* F_1 -values:

630 (4.6)
$$w_{i,r}^n := v_{\sigma_n(i),1} \text{ for } i \in [\![1,n]\!]$$

$$\begin{array}{l} g_{31} \\ g_{32}^{31} \end{array} (4.7) \qquad \qquad w_{0,r}^n \coloneqq \tilde{x}_{\min,r} \text{ and } w_{n+1,r}^n \coloneqq \tilde{x}_{\max,r} \end{array} .$$

Naturally, we have $w_{0,r}^n \leq w_{1,r}^n \leq \ldots \leq w_{n+1,r}^n$, and the intervals $[w_{i-1,r}^n, w_{i,r}^n]$ for $i \in [1, n + 1]$ form a partition of $[\tilde{x}_{\min,r}, \tilde{x}_{\max,r}]$, see the righthand plot of Figure 4. The interval $[w_{i-1,r}^n, w_{i,r}^n]$ corresponds to the part of the Pareto front dominating r_i^n , the reference point associated to the *i*-th gap region of S_n . When the Pareto front is bilipschitz, the lengths of these intervals converge asymptotically to 0 as stated in the next lemma. It is a direct consequence of the convergence of $HV_r(S_n)$ to $HV_r(PF_f)$ stated in Theorem 4.2.

640 LEMMA 4.4. If the Pareto front is described by a bilipschitz function f and the greedy 641 set sequence is associated to a valid reference point r, then the ordered greedy set along 642 F_1 -values satisfy $\max_{i \in [1,n+1]} w_{i,r}^n - w_{i-1,r}^n \xrightarrow{n \to \infty} 0$ with the $w_{i,r}^n$ defined in (4.7).

643 Proof. Let L_{\min} and L_{\max} be constants such that f is (L_{\min}, L_{\max}) -bilipschitz. The 644 area of the *i*-th gap region of S_n is $\int_{w_{i-1,r}}^{w_{i,r}^n} (f(x) - f(w_{i,r}^n)) dx$. This is larger than 645 $\int_{w_{i-1,r}}^{w_{i,r}^n} L_{\min} \times (w_{i,r}^n - x) dx$, which equals $\frac{1}{2} \times L_{\min} \times (w_{i,r}^n - w_{i-1,r}^n)^2$. Since the area 646 of any gap region of S_n is inferior to the optimality gap at iteration n, this implies that 647 the difference $w_{i,r}^n - w_{i-1,r}^n$ is inferior to $\sqrt{2 \times (\text{HV}_r(\text{PF}_f) - \text{HV}_r(S_n))}$ for all n, for all 648 $i \in [1, n+1]$. Therefore, the convergence of $\text{HV}_r(S_n)$ to $\text{HV}_r(\text{PF}_f)$ stated in Theorem 4.2 649 implies that the maximum over i of $w_{i,r}^n - w_{i-1,r}^n$ converges to 0.

We prove in the next lemma that if the Pareto front is described by a bilipschitz function with a Hölder continuous derivative, then the restriction of this function associated to a gap region of S_n is bilipschitz for some constants whose ratio converges asymptotically to 1. Linear functions being the only functions to be (L_{\min}, L_{\max}) -bilipschitz with $L_{\min}/L_{\max} = 1$, it supports the interpretation that for such Pareto fronts, the convergence of the greedy set sequence is asymptotically similar as if they were affine Pareto fronts.

657 When the function describing the Pareto front is bilipschitz, its restriction to the part

of the Pareto front dominating r_i^n , that is $[w_{i-1,r}^n, w_{i,r}^n]$, is $(L_{\min}^{i,n}, L_{\max}^{i,n})$ -bilipschitz with

659 (4.8)
$$L_{\min}^{i,n} := \inf \left\{ \left| \frac{f(x) - f(y)}{x - y} \right|, x, y \in [w_{i-1,r}^n, w_{i,r}^n], x \neq y \right\} \text{ and}$$
$$L_{\max}^{i,n} := \sup \left\{ \left| \frac{f(x) - f(y)}{x - y} \right|, x, y \in [w_{i-1,r}^n, w_{i,r}^n], x \neq y \right\}.$$

660 At iteration n, the ratio between $L_{\max}^{i,n}$ and $L_{\min}^{i,n}$, the bilipschitz constants on the *i*-th 661 gap region of S_n , is by definition smaller than

662 (4.9)
$$q_n := \max\left\{\frac{L_{\max}^{i,n}}{L_{\min}^{i,n}}, i \in [\![1, n+1]\!] : [w_{i-1,r}^n, w_{i,r}^n] \neq \emptyset\right\}$$

The proof of the convergence of q_n to 1 relies on the fact that a derivable function can be approximated locally by an affine function. The quality of this approximation is guaranteed by the Hölder continuity of the derivative.

667 LEMMA 4.5. We consider a greedy set sequence $(S_n)_{n \in \mathbb{N}^*}$ relative to a valid reference 668 point r. If the Pareto front is described by a bilipschitz function with a Hölder continuous 669 derivative, then q_n defined in (4.9) converges asymptotically to 1.

Proof. We take α such that f' is Hölder continuous with exponent α , i.e f is $\mathcal{C}^{1,\alpha}$, 670 and $L_{\min}, L_{\max} > 0$ such that the function f describing the Pareto front is (L_{\min}, L_{\max}) -671 bilipschitz. We recall that f is decreasing, and thus for all x < y, we have $f(x) - f(y) \ge 0$. 672 Since f is $\mathcal{C}^{1,\alpha}$ and therefore \mathcal{C}^1 , the Taylor formula with Lagrange remainder states that 673 for all x < y, there exists $\xi \in [x, y]$ such that $f(y) = f(x) + (y - x) \times f'(\xi)$. Since f is $\mathcal{C}^{1,\alpha}$, 674 this implies that for all x < y, $|f(y) - f(x) - (y - x) \times f'(x)| \le (y - x)^{1+\alpha} \times [f']_{\mathcal{C}^{\alpha}}$. Thus, 675 $\frac{f(y)-f(x)}{x-y}$ is lower than $-f'(x) + [f']_{\mathcal{C}^{\alpha}} \times (y-x)^{\alpha}$. We now restrict ourselves to x and 676 y belonging to the non-empty interval $[w_{i-1,r}^n, w_{i,r}^n]$. Our goal is to find an upper bound 677 depending on i but not on either x or y. Since f is $\mathcal{C}^{1,\alpha}$, the difference between -f'(x)678and $-f'(w_{i-1,r}^n)$ is lower than $[f']_{\mathcal{C}^{\alpha}} \times (x - w_{i-1,1,r}^n)^{\alpha}$, and thus $[f']_{\mathcal{C}^{\alpha}} \times (w_{i,r}^n - w_{i-1,r}^n)^{\alpha}$. Additionally, the difference between x and y is lower than $w_{i,r}^n - w_{i-1,r}^n$. We conclude that 679 680 for $x, y \in [w_{i-1,r}^n, w_{i,r}^n], \frac{f(y) - f(x)}{x - y}$ is lower than $-f'(w_{i-1,1,r}^n) + 2[f']_{\mathcal{C}^{\alpha}} \times (w_{i,r}^n - w_{i-1,r}^n)^{\alpha}$, 681 and thus so is $L_{\max}^{i,n}$ defined in (4.8). 682

Following the same approach, we can also infer that $L_{\max}^{i,n}$ defined in (4.8) is greater than the symmetric quantity $-f'(w_{i-1,1,r}^n) - 2[f']_{\mathcal{C}^{\alpha}} \times (w_{i,r}^n - w_{i-1,r}^n)^{\alpha}$. The quantity $-f'(w_{i-1,1,r}^n)$ is greater than L_{\min} and $(w_{i,r}^n - w_{i-1,r}^n)^{\alpha}$ is smaller than $\max_{i \in [1,n+1]} (w_{i,r}^n - w_{i-1,r}^n)^{\alpha}$. $w_{i-1,r}^n)^{\alpha}$. As a consequence, q_n is lower than $\frac{L_{\min}+2[f']_{\mathcal{C}^{\alpha}} \times \max_{i \in [1,n+1]} (w_{i,r}^n - w_{i-1,r}^n)^{\alpha}}{L_{\min}-2[f']_{\mathcal{C}^{\alpha}} \times \max_{i \in [1,n+1]} (w_{i,r}^n - w_{i-1,r}^n)^{\alpha}}$. By Lemma 4.4, $\max_{i \in [1,n+1]} w_{i,r}^n - w_{i-1,r}^n$ converges to 0 and thus, this upper bound on q_n converges to 1. By definition, q_n is always higher than 1, and thus converges to 1.

A consequence of the previous lemma is that the bounds on the normalized hypervolume improvement of v_{n+1} to S_n that we can infer from Propositions 3.10 and 3.11 converge asymptotically to 1/2, see (4.10). Similarly, the bounds on the normalized area of the child of a gap region that we can infer from Lemma A.2 converge to 1/4, see (4.11). These asymptotic values correspond to the case of an affine Pareto front with a reference point dominating the nadir point, see Figure 5.



Fig. 5: The three greedy sets S_1 (left), S_2 (middle) and S_3 (right) and their gap regions for an affine Pareto front with a reference point r dominating the nadir point (1, 1). The area of the gap regions of S_1 are half of $HV_r(PF_f)$ (left). The area of the new gap regions of S_2 is a quarter of the area of their parents (middle). The optimality gap of S_3 (right) is half of the optimality gap of S_1 .

EEMMA 4.6. We consider a greedy set sequence $(S_n)_{n \in \mathbb{N}^*}$ relative to a valid reference point r. If the Pareto front is described by a bilipschitz function f with a Hölder continuous derivative, then for all $\epsilon > 0$, for n large enough, for every non-empty gap region $\mathcal{G}_{S_n,i}^r$ and every child $\mathcal{G}_{S_n,j}^r$ of $\mathcal{G}_{S_n,i}^r$, we have

699 (4.10)
$$\frac{1}{2} \times (1-\epsilon) \le \frac{\max_{u \in \mathcal{G}_{\mathcal{S}_{n,i}}^r} HVI_r(u, \mathcal{S}_n)}{\lambda(\mathcal{G}_{\mathcal{S}_{n,i}}^r)} \le \frac{1}{2} \times (1+\epsilon) \text{ and}$$

700 (4.11)
$$\frac{1}{4 \times (1+\epsilon)} \le \frac{\lambda(\mathcal{G}'_{\mathcal{S}_{m},j})}{\lambda(\mathcal{G}'_{\mathcal{S}_{n},i})} \le \frac{1}{4 \times (1-\epsilon)}$$

Proof. The set of the first coordinates of the vectors of the Pareto front which dominate r_i^n is the interval $[w_{i-1,r}^n, w_{i,r}^n]$. The restriction to $[w_{i-1,r}^n, w_{i,r}^n]$ of the function fdescribing the Pareto front is (L_{\min}, L_{\max}) -bilipschitz for some L_{\min} and L_{\max} such that $\frac{L_{\max}}{L_{\min}} = q_n$ with q_n defined in (4.9). Additionally, as stated in Proposition B.2, for n large enough, all the r_i^n corresponding to non-empty gap regions dominate the nadir point. It allows us to apply both Lemma A.2 and Proposition 3.11 to such gap regions. By Propositions 3.10 and 3.11, $\frac{\max_{u \in \mathrm{PF}_f} \mathrm{HV}_{r_i^n}(u)}{\mathrm{HV}_{r_i^n}(\mathrm{PF}_f)}$ is between $\frac{1}{2} \times \frac{1}{q_n}$ and $\frac{1}{2} \times q_n$.

Additionally, by Lemma A.2,
$$\frac{\lambda(\mathcal{G}_{\mathcal{S}_m,j}^r)}{\operatorname{HV}_{r_i^n}(\operatorname{PF}_f)}$$
 is between $\frac{1-\frac{1}{2}\times q_n}{1+q_n^2}$ and $\frac{1-\frac{1}{2}\times \frac{1}{q_n}}{1+\frac{1}{q_n^2}}$. The maxim

mum over the vectors u belonging to the Pareto front of $\operatorname{HV}_{r_i^n}(u)$ is equal to the maximum over u belonging to the *i*-th gap region of \mathcal{S}_n of $\operatorname{HVI}_r(u, \mathcal{S}_n)$. Indeed, $\operatorname{HV}_{r_i^n}(.)$ is null for vectors outside the *i*-th gap region of \mathcal{S}_n while it is nonnegative, equal to $\operatorname{HVI}_r(., \mathcal{S}_n)$, otherwise. Additionally, $\operatorname{HV}_{r_i^n}(\operatorname{PF}_f)$ equals $\lambda(\mathcal{G}_{\mathcal{S}_n,i}^r)$. The convergence of q_n to 1 stated in Lemma 4.5 imply that the bounds proven so far converge to a half and a quarter, respectively. Thus, we have (4.10) and (4.11) for n large enough.

The following lemma states that for n large enough, the area of two non-empty gap regions

relative to the same greedy set cannot be too different. More precisely, the area of any gap region of S_n cannot be more than $4 \times (1 + o(\epsilon))$ times greater than the area of another

⁷¹⁹ gap region of S_n . The proof relies on considering the parents of the gap regions.

T20 LEMMA 4.7. We consider a greedy set sequence $(S_n)_{n \in \mathbb{N}^*}$ relative to a valid reference point r. If the Pareto front is described by a bilipschitz function with a Hölder continuous derivative, then for all $\epsilon > 0$, for n large enough and for any non-empty gap regions of $S_n, \mathcal{G}^r_{S_n,i}$ and $\mathcal{G}^r_{S_n,j}$ with $i, j \in [1, n + 1]$, we have

724 (4.12)
$$\frac{\lambda(\mathcal{G}_{\mathcal{S}_n,i}^r)}{\lambda(\mathcal{G}_{\mathcal{S}_n,i}^r)} \le 4 \times \frac{(1+\epsilon)^2}{1-\epsilon}$$

Proof. Fix $\epsilon > 0$. By Lemma 4.6, there exists $N_1 \in \mathbb{N}^*$ such that for all n greater than N_1 , (4.10) and (4.11) are verified for any non-empty gap region of S_n and its children. Since $\max_{i \in [\![1,n+1]\!]} w_{i,r}^n - w_{i-1,r}^n$ converges to 0 by Lemma 4.4, every non-empty gap region is filled at some point. Take N_2 such that all the non-empty gap regions of S_{N_1} are filled at iteration N_2 . For all n greater than N_2 , (4.10) and (4.11) are true for any non-empty gap region of S_n and its children, but also for its parents.

732 Take $n \geq N_2$. We note $\mathcal{G}_1 := \mathcal{G}_{\mathcal{S}_n,i}^r$ and $\mathcal{G}_2 := \mathcal{G}_{\mathcal{S}_n,j}^r$ two distinct non-empty gap regions of \mathcal{S}_n , and \mathcal{P}_1 and \mathcal{P}_2 their respective parents. When two sets correspond to gap regions 733 734 relative to the same greedy set \mathcal{S}_m , we say that they cohabit at iteration m. Since only one vector is added to S_n at a time, the cohabitation of \mathcal{G}_1 and \mathcal{G}_2 implies that either \mathcal{G}_1 735and \mathcal{P}_2 or \mathcal{G}_2 and \mathcal{P}_1 cohabit at some earlier iteration. In the first case, there necessarily 736 exists $m \geq N_2$ such that \mathcal{P}_2 and \mathcal{G}_1 are gap regions relative to \mathcal{S}_m and v_{m+1} belongs 737 to \mathcal{P}_2 , otherwise, \mathcal{G}_1 and \mathcal{G}_2 would not cohabit. By (4.10), the maximum hypervolume 738 improvement to S_m of a vector of \mathcal{G}_1 and of a vector of \mathcal{P}_2 are at least $\frac{1}{2} \times (1-\epsilon) \times \lambda(\mathcal{G}_1)$ 739 and at most $\frac{1}{2} \times (1+\epsilon) \times \lambda(\mathcal{P}_2)$, respectively. Since a vector of \mathcal{P}_2 , v_{m+1} , maximizes the 740hypervolume improvement to \mathcal{S}_m , we have $\lambda(\mathcal{G}_1) \times \frac{1}{2} \times (1-\epsilon) \leq \lambda(\mathcal{P}_2) \times \frac{1}{2} \times (1+\epsilon)$. Since 741 $\lambda(\mathcal{P}_2)$ is lower than $4 \times (1 + \epsilon)$ times the area of its child $\lambda(\mathcal{G}_2)$ by (4.11), this inequality 742 implies (4.12). In the second case, \mathcal{P}_2 is filled before \mathcal{P}_1 . Thus, there exists $m \geq N_2$ such 743 that \mathcal{P}_1 and \mathcal{P}_2 cohabit at iteration m and v_{m+1} belongs to \mathcal{P}_2 . Since the area of \mathcal{P}_1 is 744higher than the one of its child \mathcal{G}_1 , the hypervolume improvement of v_{m+1} to \mathcal{S}_m is still 745higher than $\frac{1}{2} \times (1 - \epsilon) \times \lambda(\mathcal{G}_1)$. The rest of the argumentation remains valid. Π 746

747 We now have all the results needed to analyze the asymptotic impact of doubling the 748 number of points in the greedy set. To prove the following asymptotic upper bound, 749 we rely on similar arguments as for its nonasymptotic counterpart, Proposition 4.1. The 750 previous lemma guarantees that the impact of doubling the number of points in the greedy 751 set is asymptotically similar to the impact of passing from n points to 2n + 1.

752 PROPOSITION 4.8. Let $(S_n)_{n \in \mathbb{N}^*}$ be a greedy set sequence relative to valid reference 753 point r. If the Pareto front is described by a bilipschitz function f with a Hölder continuous 754 derivative, then for all $\epsilon > 0$, we have for n large enough

755 (4.13)
$$\frac{HV_r(PF_f) - HV_r(\mathcal{S}_{2n})}{HV_r(PF_f) - HV_r(\mathcal{S}_n)} \le \frac{1}{2} + o(\epsilon).$$

757 Proof. Fix $\epsilon > 0$. Fix n large enough to verify (4.10) and (4.12) for this particular ϵ . 758 Let σ be a permutation of $[\![1, n + 1]\!]$ such that the *i*-th gap region of S_n is filled by 759 $v_{n+\sigma(i)}$ when it is filled before iteration 2n + 1. With this choice of permutation, $\mathcal{G}_{S_n,i}^r$ 760 is always a gap region of $S_{n+\sigma(i)-1}$. Thus, $\text{HVI}_r(v_{n+\sigma(i)}, S_{n+\sigma(i)-1})$ is superior to the 761 maximum hypervolume improvement of a vector of $\mathcal{G}_{S_n,i}^r$ to $S_{n+\sigma(i)-1}$, which is superior 762 to $\frac{1}{2} \times (1-\epsilon) \times \lambda(\mathcal{G}_{S_n,i}^r)$ by (4.10). It is equivalent to say that the hypervolume improvement at iteration $n + \sigma(i)$ is higher than $\frac{1}{2} \times (1 - \epsilon) \times \lambda(\mathcal{G}_{\mathcal{S}_n,i}^r)$. Summing over $i \in [\![1, n + 1]\!]$, we obtain that the hypervolume improvement between iteration n and 2n + 1 is higher than the sum over i of $\frac{1}{2} \times (1 - \epsilon) \times \lambda(\mathcal{G}_{\mathcal{S}_n,i}^r)$, that is $\frac{1}{2} \times (1 - \epsilon)$ times the optimality gap at iteration n.

Now, we need to bound the hypervolume improvement at iteration 2n + 1, that is HVI_r (v_{2n+1}, S_{2n}) . It is lower than $\frac{1}{2} \times (1+\epsilon) \times \max_{i \in [\![1,2n+1]\!]} \lambda(\mathcal{G}_{\mathcal{S}_{2n},i}^{r})$ by (3.4) and (4.10). Since the area of a gap region is lower than the one of its parent, the maximum area of a gap region is lower at iteration 2n than at iteration n. The maximum area of one of the more than n-1 gap regions of \mathcal{S}_n is lower than $\frac{1}{n-1} \times \frac{4 \times (1+\epsilon)^2}{1-\epsilon}$ times the optimality gap at iteration n by (4.12). To summarize, the hypervolume improvement at iteration 2n+1 is lower than $\frac{2 \times (1+\epsilon)^3}{(n-1) \times (1-\epsilon)}$ times the optimality gap at iteration n.

773 2n+1 is lower than $\frac{2\times(1+\epsilon)^3}{(n-1)\times(1-\epsilon)}$ times the optimality gap at iteration n. 774 We conclude that the relation between the optimality gap at iteration 2n and at 775 iteration n is lower than $1-\frac{1}{2}\times(1-\epsilon)+\frac{1-\epsilon}{2\times(n-1)}$.

776 We broadly follow the same approach to obtain the following asymptotic lower bound on the impact of doubling the number of points in the greedy set. Lemmas 4.6 and 4.7 are 777 key to prove a lower bound on the hypervolume improvement at iteration k. They allow 778 to prove that filling a gap region of S_n more than once gives, up to a factor $1 + o(\epsilon)$, a 779 lower hypervolume improvement than filling a gap region which was not filled. Indeed, 780 the area of a descendant of a gap region of S_n is at most $\frac{1}{4} + o(\epsilon)$ times the area of its 781 parent by Lemma 4.6, which is itself at most $4 + o(\epsilon)$ times the area of any other gap 782 region of \mathcal{S}_n by Lemma 4.7. 783

784 PROPOSITION 4.9. Let $(S_n)_{n \in \mathbb{N}^*}$ be a greedy set sequence relative to a valid reference 785 point r. If the Pareto front is desribed by a bilipschitz function f with a Hölder continuous 786 derivative, then for all $\epsilon > 0$, we have for n large enough

787 (4.14)
$$\frac{HV_r(PF_f) - HV_r(\mathcal{S}_{2n})}{HV_r(PF_f) - HV_r(\mathcal{S}_n)} \ge \frac{1}{2} + o(\epsilon).$$

789 Proof. Fix $\epsilon > 0$. Fix n large enough to verify (4.10), (4.11) and (4.12) for this 790 particular ϵ . Let $\delta \in \{-1, 0, 1\}$ be such that S_n has $n + \delta$ non-empty gap regions. Let 791 $i_0 := 1$ when the left extreme gap region is empty and $i_0 := 0$ otherwise.

Let σ be a permutation of $[\![1,n+\delta]\!]$ such that the i-th non-empty gap region of 792 793 $\mathcal{S}_n, \mathcal{G}^r_{\mathcal{S}_n, i_0+i}$, is filled by the vector $v_{n+\sigma(i)}$ when it is filled before iteration $2n+\delta$. We distinguish two cases. In the first case, $v_{n+\sigma(i)}$ is the child of the *i*-th non-empty gap 794 region of \mathcal{S}_n , and consequently its hypervolume improvement to $\mathcal{S}_{n+\sigma(i)-1}$ is at most 795 $\frac{1}{2} \times (1+\epsilon) \times \lambda(\mathcal{G}^r_{\mathcal{S}_n, i_0+i})$ by (4.10). In the second case, $v_{n+\sigma(i)}$ belongs to $\mathcal{G}^r_{\mathcal{S}_n, i_0+j}$, the 796 \overline{j} -th non-empty gap region of \mathcal{S}_n , with $j \neq i$ and, by definition of σ , fills a descendant of 797 this gap region not $\mathcal{G}_{\mathcal{S}_n, i_0+j}^r$ itself. By (4.10), the hypervolume improvement of $v_{n+\sigma(i)}$ to 798 $S_{n+\sigma(i)-1}$ is still at most $\frac{1}{2} \times (1+\epsilon)$ times the area of the gap region it fills. By (4.11), 799the area of a descendant of $\mathcal{G}_{\mathcal{S}_n,i_0+j}^r$ is smaller than $\frac{1}{4\times(1-\epsilon)}$ times the area of its ancestor. 800 By (4.12), we also know that the area of the *i*-th non-empty gap region of S_n is at most 801 $4 \times \frac{(1+\epsilon)^2}{1-\epsilon}$ times the area of any other gap region of S_n , in particular its *i*-th non-empty 802 gap region. We conclude that the hypervolume improvement of $v_{n+\sigma(i)}$ to $\mathcal{S}_{n+\sigma(i)-1}$ is 803 lower than $\frac{1}{2} \times \frac{(1+\epsilon)^3}{(1-\epsilon)^2} \times \lambda(\mathcal{G}^r_{\mathcal{S}_n,i_0+i})$. To summarize, since $1+\epsilon$ is lower than $\frac{(1+\epsilon)^3}{(1-\epsilon)^2}$, the 804 hypervolume improvement at any iteration $n + \sigma(i)$ is lower than $\frac{1}{2} \times \frac{(1+\epsilon)^3}{(1-\epsilon)^2} \times \lambda(\mathcal{G}^r_{\mathcal{S}_n,i})$. 805

Summing over $i \in [\![1, n + \delta]\!]$, the hypervolume improvement from iteration n to $2n + \delta$ is lower than $\frac{1}{2} \times \frac{(1+\epsilon)^3}{(1-\epsilon)^2}$ times the sum over i of $\lambda(\mathcal{G}^r_{\mathcal{S}_n,i})$, that is the optimality gap at iteration n.

Now, it is left to prove an upper bound on $HV_r(S_{2n}) - HV_r(S_{2n+\delta})$. This quantity is the highest for $\delta = -1$, where it is simply the hypervolume improvement at iteration 2n. As in the previous proof, it is lower than $\frac{1+\epsilon}{2\times(n-1)}$ times the optimality gap at iteration n. Therefore, the relation between the optimality gap at iteration 2n and at iteration nis higher than $1 - \frac{1}{2} \times \frac{(1+\epsilon)^3}{(1-\epsilon)^2} - \frac{1+\epsilon}{2\times(n-1)}$.

814 We combine the lower and upper asymptotic bounds to obtain the following theorem.

THEOREM 4.10. Consider a biobjective optimization problem and a greedy set sequence $(S_n)_{n \in \mathbb{N}^*}$ relative to a valid reference point r. If the Pareto front is described by a bilipschitz function f with a Hölder continuous derivative, we have

Consequently, for such Pareto front and reference point and under Assumption 2.4 of perfect singleobjective optimization, HV-ISOOMOO algorithms relative to r satisfy (4.15) where S_n is replaced by \mathcal{I}_n , the final incumbents Pareto front approximation at iteration n.

5. Conclusion. We prove that HV-ISOOMOO algorithms coupled with a singleob-824 jective optimizer converge in O(1/n) on convex Pareto fronts and in $O(1/n^c)$ on bilipschitz 825 Pareto fronts with $c \leq 1$ depending on the bilipschitz constants where n is the number of 826 meta-iterations. Each meta-iteration corresponds to a singleobjective optimization run. 827 828 Both bounds are tight over the class of Pareto fronts and reference points considered. Indeed, they are reached for affine Pareto fronts and reference points dominating the nadir 829 point. On convex Pareto fronts, the highest achievable convergence-speed is reached 830 [15]. It shows that greedily adding points maximizing the hypervolume contribution as in 831 HV-ISOOMOO algorithms is an effective way to quickly increase the hypervolume. Ad-832 ditionally, we prove that for bilipschitz Pareto fronts with a Hölder continuous derivative, 833 834 asymptotically, doubling the number of meta-iterations halves the optimality gap. This asymptotic behavior resembles what we would observe with an affine Pareto front and a 835 reference point dominating the nadir point. Beware that this does not guarantee a rate of convergence in $\Theta(1/n)$. For example, the sequences $(\frac{\log(n)}{n})_{n \in \mathbb{N}^*}$ and $(\frac{1}{n \times \log(n)})_{n \in \mathbb{N}^*}$ both verify this property. The convergence rate on nonconvex Pareto fronts could theoretically 836 837 838 be slower than $\Theta(1/n)$, but not faster by [15]. 839

Appendix A. Normalized areas of the gap regions relative to an hypervolume maximizer. The goal of this section is to prove bounds on the normalized areas of the gap regions $\mathcal{G}_{\text{left}}^{u^*}$ and $\mathcal{G}_{\text{right}}^{u^*}$ relative to an hypervolume maximizer $u^* = (x^*, f(x^*))$ (see the lefthand plot of Figure 6) in the case of a bilipschitz Pareto front and of a reference point r dominating the nadir point. These bounds are stated in Lemma A.2.

We exploit the bounds on the normalized maximum hypervolume proven in Subsection 3.3 and the following lower and upper bounds on the relation between $\lambda(\mathcal{G}_{left}^{u^*})$ and $\lambda(\mathcal{G}_{right}^{u^*})$.



Fig. 6: Illustration of elements of the proofs of Proposition A.1 in the case $r_1 \leq x_{\max}$ (left and middle) and of Lemma B.1 (right). The Pareto front is described by $f(x) = \frac{e}{e-1} \times e^{-x} + 1 - \frac{e}{e-1}$ for $x \in [0, 1]$. Left: the gap regions $\mathcal{G}_{\text{left}}^{u^*}$ and $\mathcal{G}_{\text{right}}^{u^*}$ with a segment of slope $-L_{\min}$ passing through u^* . Middle: the hypervolume improvements $\text{HVI}_r(u^*, u_\epsilon)$ \blacksquare and $\text{HVI}_r(u_\epsilon, u^*) \blacksquare$. Right: the hypervolume improvement $\text{HVI}_r(u_{\min}, u) \blacksquare$ and its counterpart $\text{HVI}_r(u, u_{\min}) \blacksquare$ where u is a vector of the Pareto front which dominates r.

PROPOSITION A.1. We assume that the Pareto front is described by a (L_{\min}, L_{\max}) bilipschitz function f. Let u^* be a non-extreme vector of the Pareto front which maximizes the hypervolume with respect to a valid reference point r. If $r_1 \leq x_{\max}$, we have $\lambda(\mathcal{G}_{right}^{u^*}) \geq$ $\frac{L_{\min}^2}{L_{\max}^2} \times \lambda(\mathcal{G}_{left}^{u^*})$. If $r_2 \leq f(x_{\min})$, we have $\lambda(\mathcal{G}_{left}^{u^*}) \geq \frac{L_{\min}^2}{L_{\max}^2} \times \lambda(\mathcal{G}_{right}^{u^*})$.

Proof. We consider the case where $r_1 \leq x_{\text{max}}$. We note $L_1 := r_1 - x^*$ and $L_2 :=$ 852 $r_2 - f(x^*)$ the lengths of the sides of the rectangle $\mathcal{D}_{u^*}^r$. For all $x, y \in [x_{\min}, x_{\max}]$, we have 853 $|f(x) - f(y)| \ge L_{\min} \times |x - y|$. Additionally, since $r_1 \le x_{\max}$, the segment $[x^*, x^* + L_1]$ is 854 included in $[x_{\min}, x_{\max}]$. As a consequence, the section of the Pareto front on the right of 855 u^* dominates the segment between u^* and $u^* + L_1 \times (1, -L_{\min})$, see the lefthand plot of 856 Figure 6. Therefore, $\lambda(\mathcal{G}_{\text{right}}^{u^*})$ is larger than the area of the region of the objective space 857 dominated by this segment, not dominated by u^* and dominating r, that is $\frac{1}{2} \times L_{\min} \times L_1^2$. 858 For all $x, y \in [x_{\min}, x_{\max}]$, we also have $|f(x) - f(y)| \leq L_{\max} \times |x - y|$. Therefore, the part of the Pareto front on the left of u^* is dominated by the segment between u^* and $u^* + L_2 \times (-\frac{1}{L_{\min}}, 1)$, and $\lambda(\mathcal{G}_{\text{left}}^{u^*})$ is lower than $\frac{1}{2} \times \frac{1}{L_{\min}} \times L_2^2$. We have yet to prove a lower bound on $\frac{L_1}{L_2}$. The vector u^* being different from u_{\min} , for $\epsilon > 0$ small enough, the vector $u := (x^* - \epsilon)^* h_1(x^* - \epsilon)^*$ belows to the Dense front A859 860 861 862 vector $u_{\epsilon} := (x^* - \epsilon, f(x^* - \epsilon))$ belongs to the Pareto front. As we can see in the middle 863 plot of Figure 6, $HVI_r(u^*, u_{\epsilon})$ is lower than $L_1 \times L_{\max} \times \epsilon$ and $HVI_r(u_{\epsilon}, u^*)$ is higher than 864 $\epsilon \times (L_2 - \epsilon \times L_{\max})$. Additionally, u^* being an hypervolume maximizer, $HVI_r(u^*, u_{\epsilon})$ is 865 higher than $\text{HVI}_r(u_{\epsilon}, u^*)$, and thus $L_1 \times L_{\text{max}} \geq L_2 - \epsilon \times L_{\text{max}}$ for all $\epsilon > 0$. Taking 866 the limit of this inequality when $\epsilon \to 0$, we obtain that $L_1 \times L_{\max} \ge L_2$. Combining 867 the bounds on $\lambda(\mathcal{G}_{left}^{u^*})$ and $\lambda(\mathcal{G}_{right}^{u^*})$ with the lower-bound on $\frac{L_1}{L_2}$, we obtain the desired 868 lower bound on $\lambda(\mathcal{G}_{right}^{u^*})$. We can obtain the symmetric inequality when $r_2 \geq f(x_{\min})$ by 869 following the same approach. Π 870

In particular, when f is bilipschitz and r dominates the nadir point, both bounds hold. We are now able to prove the desired bounds on the normalized area of the gap regions $\mathcal{G}_{left}^{u^*}$ and $\mathcal{G}_{right}^{u^*}$. EEMMA A.2. Let u^* be a vector which maximizes the hypervolume with respect to a valid reference point r. If the Pareto front is described by a (L_{\min}, L_{\max}) -bilipschitz function f and the reference point r dominates the nadir point, both $\mathcal{G}_{left}^{u^*}$ and $\mathcal{G}_{right}^{u^*}$ are

877 between
$$(1 - \frac{1}{2} \times \frac{L_{\text{max}}}{L_{\text{min}}})/(1 + \frac{L_{\text{max}}}{L_{\text{min}}^2})$$
 and $(1 - \frac{1}{2} \times \frac{L_{\text{min}}}{L_{\text{max}}})/(1 + \frac{L_{\text{min}}}{L_{\text{max}}^2})$

Proof. Let note arbitrarily \mathcal{G}_1 and \mathcal{G}_2 the two gap regions of the set $S = \{u^*\}$. By Proposition A.1, $\lambda(\mathcal{G}_2)$ is between $\frac{L_{\min}^2}{L_{\max}^2} \times \lambda(\mathcal{G}_1)$ and $\frac{L_{\max}^2}{L_{\min}^2} \times \lambda(\mathcal{G}_1)$. Additionally, Propositions 3.10 and 3.11, the normalized maximum hypervolume $\max_{u \in \mathrm{PF}_f} \mathrm{HV}_r(u)$ over $\mathrm{HV}_r(\mathrm{PF}_f)$ is between $\frac{1}{2} \times \frac{L_{\min}}{L_{\max}}$ and $\frac{1}{2} \times \frac{L_{\min}}{L_{\min}}$. These bounds can be transformed into bounds on $\mathrm{HV}_r(\mathrm{PF}_f) - \max_{u \in \mathrm{PF}_f} \mathrm{HV}_r(u)$, that is $\lambda(\mathcal{G}_1) + \lambda(\mathcal{G}_2)$. As a consequence, $\lambda(\mathcal{G}_1)$ is between $(1 - \frac{1}{2} \times \frac{L_{\min}}{L_{\min}}) \times \mathrm{HV}_r(\mathrm{PF}_f) - \frac{L_{\max}^2}{L_{\min}^2} \times \lambda(\mathcal{G}_1)$ and $(1 - \frac{1}{2} \times \frac{L_{\min}}{L_{\max}}) \times \mathrm{HV}_r(\mathrm{PF}_f) - \frac{L_{\max}^2}{L_{\min}^2} \times \lambda(\mathcal{G}_1)$. Moving all the $\lambda(\mathcal{G}_1)$ terms on the same side and re-normalizing this side, we obtain the desired bounds for \mathcal{G}_1 , which can be chosen to be either $\mathcal{G}_{\mathrm{left}}^{u^*}$ or $\mathcal{G}_{\mathrm{right}}^{u^*}$.

Appendix B. The nadir point is dominated by all the r_i^n corresponding to non-empty gap regions for n large. We show in this section that for bilipschitz Pareto fronts, the nadir point is dominated by all the r_i^n corresponding to non-empty gap regions, for n large enough. This result is stated in Proposition B.2 and used in Subsection 4.2. It is equivalent to prove that the extreme vectors which dominate the reference point belong to the greedy set for n large enough.

First, we prove in the next proposition that if $r_1 > x_{\text{max}}$ (resp. $r_2 > f(x_{\min})$), then for r_2 (resp. r_1) close enough to $f(x_{\max})$ (resp. x_{\min}) the extreme vector u_{\max} (resp. u_{\min}) is the only hypervolume maximizer, see the righthand plot of Figure 6. There are similar statements in [8] for the set of μ points maximizing the hypervolume, but they only apply to $\mu \geq 2$.

EEMMA B.1. We assume that the Pareto front is described by a function f which is (L_{\min}, L_{\max}) -bilipschitz. For any valid reference point $r = (r_1, r_2)$ such that $r_1 > x_{\max}$ and $f(x_{\max}) < r_2 < f(x_{\max}) + L_{\min} \times (r_1 - x_{\max})$, the right extreme of the Pareto front u_{\max} is the only maximizer of $HV_r(.)$. Additionally, for any valid reference point $r = (r_1, r_2)$ such that $r_2 > f(x_{\min})$ and $x_{\min} < r_1 < x_{\min} + \frac{r_2 - f(x_{\min})}{L_{\max}}$, the vector $u_{\min} = (x_{\min}, f(x_{\min}))$ is the only maximizer of $HV_r(.)$.

Proof. This proof is illustrated in the righthand plot of Figure 6. Let r be a reference 903 point such that $r_2 > f(x_{\min})$ and $x_{\min} < r_1 < x_{\min} + \frac{r_2 - f(x_{\min})}{L_{\max}}$. Let $u = (x, f(x)) \neq u_{\min}$ be a vector of the Pareto front which dominates r. The hypervolume improvement of u_{\min} 904 905 to $\{u\}$ is $(r_2 - f(x_{\min})) \times (x - x_{\min})$. The hypervolume improvement of u to $\{u_{\min}\}$ is equal 906 to $(f(x_{\min}) - f(x)) \times (r_1 - x)$, which is smaller than $L_{\max} \times (x - x_{\min}) \times (r_1 - x_{\min})$ since u 907 dominates r and f is (L_{\min}, L_{\max}) -bilipschitz. Since we assume that $L_{\max} \times (r_1 - x_{\min}) < r_1 - r_{\min}$ 908 $r_2 - f(x_{\min})$, the upper bound on $HVI_r(u, u_{\min})$ is strictly lower than $HVI_r(u_{\min}, u)$. As 909 a consequence, the hypervolume of u_{\min} is strictly higher than the one of u. We conclude 910 that u_{\min} is the unique hypervolume maximizer. The symmetric result can be obtained 911 with the same approach. 912

913 It is left to prove that when $r_1 > x_{\max}$ (resp. $r_2 > f(x_{\min})$), the second coordinate of 914 r_{n+1}^n (resp. the first coordinate of r_0^n) indeed converge to $f(x_{\max})$ (resp. x_{\min}). It is a 915 straightforward consequence of Lemma 4.4. Therefore, we are able to conclude. 916 PROPOSITION B.2. We assume that the the Pareto front is described by a bilipschitz 917 function. Let $(S_n)_{n \in \mathbb{N}^*}$ be a greedy set sequence associated to a valid reference point r. 918 For n large enough, every reference point r_i^n corresponding to a non-empty gap region 919 $\mathcal{G}_{S_n,i}^r$ dominates the nadir point.

Proof. By Lemma 4.4, $w_{n,r}^n$ converges to x_{\max} , and thus the right extreme reference 920 point $r_{n+1}^n := (r_1, f(w_{n,r}^n))$ converges to $(r_1, f(x_{\max}))$ by continuity of f. Therefore, if r_1 921 is strictly higher than x_{\max} , then there exists N such that for all $n \geq N$, r_{n+1}^n verifies 922the assumptions on the reference point of Lemma B.1 which guarantee that $u_{\rm max}$ is the 923 unique maximizer of $HV_r(.)$ over the right extreme gap region $\mathcal{G}^r_{\mathcal{S}_n,n+1}$. Let assume that 924 u_{\max} does not belong to S_n . Then, $w_{N,r}^N \neq x_{\max}$, and since $w_{n,r}^n$ converges to x_{\max} , the left extreme gap region $\mathcal{G}_{S_n,i}^r$ is necessarily filled at some later iteration. When the right 925926 extreme gap region is filled, $u_{\rm max}$, the unique minimizer of $HV_r(.)$ over this gap region, 927 928 is added to the greedy set. To summarize, if $r_1 > x_{\text{max}}$, then for n large enough \mathcal{S}_n 929 contains $u_{\rm max}$, and thus the right extreme gap region is empty. We can prove with the 930 same approach that for $r_2 > f(x_{\min})$, S_n contains u_{\min} for n large enough.

At any iteration, the non-extreme reference points dominate the nadir point. Additionally, we proved that either $r_1 < x_{\text{max}}$ (resp. $r_2 < f(x_{\text{min}})$), and thus the left (resp. right) extreme reference point dominates the nadir point or for *n* large enough, the left (resp. right) extreme gap region is empty.

REFERENCES

936	[1] A. AUGER, J. BADER, AND D. BROCKHO	FF, Theoretically Investigating Optimal μ -Distributions
937	for the Hypervolume Indicator: First	Results for Three Objectives, in Parallel Problem Solving
938	from Nature, PPSN XI, D. Hutchison,	T. Kanade, J. Kittler, J. M. Kleinberg, F. Mattern, J. C.
939	Mitchell, M. Naor, O. Nierstrasz, C.	Pandu Rangan, B. Steffen, M. Sudan, D. Terzopoulos,
940	D. Tygar, M. Y. Vardi, G. Weikum, R	. Schaefer, C. Cotta, J. Kołodziej, and G. Rudolph, eds.,
941	vol. 6238, Springer Berlin Heidelberg,	Berlin, Heidelberg, 2010, pp. 586–596, https://doi.org/
942	$10.1007/978$ -3-642-15844-5_59.	
943	[2] A. Auger, J. Bader, D. Brockhoff, A	ND E. ZITZLER, Theory of the Hypervolume Indicator:
944	Optimal μ -Distributions and the Cho	ice of the Reference Point, in Proceedings of the Tenth
945	ACM SIGEVO Workshop on Foundat	cions of Genetic Algorithms, FOGA '09, New York, NY,
946	USA, Jan. 2009, Association for Com	puting Machinery, pp. 87–102, https://doi.org/10.1145/
947	1527125.1527138.	
948	[3] A. Auger, J. Bader, D. Brockhoff, A	ND E. ZITZLER, Hypervolume-based multiobjective opti-
949	mization: Theoretical foundations an	d practical implications, Theoretical Computer Science,
950	425 (2012), pp. 75–103, https://doi.or	g/10.1016/j.tcs.2011.03.012.
951	[4] N. BEUME, M. LAUMANNS, AND G. RUDOLF	PH, Convergence Rates of (1+1) Evolutionary Multiobjec-
952	tive Optimization Algorithms, in Para	llel Problem Solving from Nature, PPSN XI, R. Schaefer,
953	C. Cotta, J. Kołodziej, and G. Rudo	olph, eds., Lecture Notes in Computer Science, Berlin,
954	Heidelberg, 2010, Springer, pp. 597–6	06, https://doi.org/10.1007/978-3-642-15844-5_60.
955	[5] N. BEUME, B. NAUJOKS, AND M. EMMERI	CH, SMS-EMOA: Multiobjective selection based on dom-
956	inated hypervolume, European Journ	al of Operational Research, 181 (2007), pp. 1653–1669,
957	https://doi.org/10.1016/j.ejor.2006.08	.008.
958	[6] J. BIGEON, S. L. DIGABEL, AND L. SALOM	on, DMulti-MADS: Mesh adaptive direct multisearch for
959	$blackbox\ multiobjective\ optimization,$	p. 30.
960	[7] K. Bringmann and T. Friedrich, The m	aximum hypervolume set yields near-optimal approxima-
961	<i>tion</i> , in Proceedings of the 12th Annua	al Conference on Genetic and Evolutionary Computation
962	- GECCO '10, Portland, Oregon, US	5A, 2010, ACM Press, p. 511, https://doi.org/10.1145/

963 1830483.1830576.

935

 964 [8] D. BROCKHOFF, Optimal μ-Distributions for the Hypervolume Indicator for Problems with Lin-965 ear Bi-objective Fronts: Exact and Exhaustive Results, in Simulated Evolution and Learning,
 966 K. Deb, A. Bhattacharya, N. Chakraborti, P. Chakroborty, S. Das, J. Dutta, S. K. Gupta,

967		A. Jain, V. Aggarwal, J. Branke, S. J. Louis, and K. C. Tan, eds., vol. 6457, Springer Berlin
968		Heidelberg, Berlin, Heidelberg, 2010, pp. 24–34, https://doi.org/10.1007/978-3-642-17298-4_2.
969	[9]	A. L. CUSTÓDIO AND J. F. A. MADEIRA, MultiGLODS: Global and local multiobjective optimiza-
970		tion using direct search, J Glob Optim, 72 (2018), pp. 323–345, https://doi.org/10.1007/
971		s10898-018-0618-1.
972	[10]	A. L. CUSTÓDIO, J. F. A. MADEIRA, A. I. F. VAZ, AND L. N. VICENTE, Direct Multisearch for
973		Multiobjective Optimization, SIAM J. Optim., 21 (2011), pp. 1109–1140, https://doi.org/10.
974		1137/10079731X.
975	[11]	E. DE KLERK, F. GLINEUR, AND A. B. TAYLOR, On the worst-case complexity of the gradient method
976		with exact line search for smooth strongly convex functions, arXiv:1606.09365 [math], (2016),
977		https://arxiv.org/abs/1606.09365.
978	[12]	R. FIORENZA, Hölder and locally Hölder Continuous Functions, and Open Sets of Class C^k , $C^{k,\lambda}$,
979		Frontiers in Mathematics, Birkhäuser Basel, 2016, https://doi.org/10.1007/978-3-319-47940-8.
980	[13]	E. H. FUKUDA AND L. M. G. DRUMMOND, On the convergence of the projected gradient method
981		for vector optimization, Optimization, 60 (2011), pp. 1009–1021, https://doi.org/10.1080/
982		02331934.2010.522710.
983	[14]	J. KNOWLES AND D. CORNE, On metrics for comparing nondominated sets, in Proceedings of the
984		2002 Congress on Evolutionary Computation. CEC'02 (Cat. No.02TH8600), vol. 1, May 2002,
985		pp. 711–716 vol.1, https://doi.org/10.1109/CEC.2002.1007013.
986	[15]	E. MARESCAUX AND N. HANSEN, Hypervolume in biobjective optimization cannot converge faster
987		than $\Omega(1/p)$, in GECCO 2021 - The Genetic and Evolutionary Computation Conference, Lille
988		/ Virtual, France, July 2021.
989	[16]	G. L. NEMHAUSER, L. A. WOLSEY, AND M. L. FISHER, An analysis of approximations for max-
990		imizing submodular set functions—I, Mathematical Programming, 14 (1978), pp. 265–294,
991		https://doi.org/10.1007/BF01588971.
992	[17]	C. TOURÉ, N. HANSEN, A. AUGER, AND D. BROCKHOFF, Uncrowded Hypervolume Improvement:
993		COMO-CMA-ES and the Sofomore framework, arXiv:1904.08823 [cs, math], (2019), https:
994		//doi.org/10.1145/3321707.3321852, https://arxiv.org/abs/1904.08823.
995	[18]	D. A. V. VELDHUIZEN AND G. B. LAMONT, Evolutionary Computation and Convergence to a Pareto

967

- 996 Front, in Proceedings of the Third Annual Conference on Genetic Programming, San Francisco, 1998, Stanford University Bookstore, pp. 221–228. 997
- [19] E. ZITZLER, J. KNOWLES, AND L. THIELE, Quality Assessment of Pareto Set Approximations, in 998999 Multiobjective Optimization: Interactive and Evolutionary Approaches, J. Branke, K. Deb, 1000 K. Miettinen, and R. Słowiński, eds., Lecture Notes in Computer Science, Springer, Berlin, 1001 Heidelberg, 2008, pp. 373–404, https://doi.org/10.1007/978-3-540-88908-3_14.