



Multiobjective Hypervolume Based ISOOMOO Algorithms Converge with At Least Sublinear Speed to the Entire Pareto Front

Eugénie Marescaux, Anne Auger

► To cite this version:

Eugénie Marescaux, Anne Auger. Multiobjective Hypervolume Based ISOOMOO Algorithms Converge with At Least Sublinear Speed to the Entire Pareto Front. 2021. hal-03198414v2

HAL Id: hal-03198414

<https://hal.science/hal-03198414v2>

Preprint submitted on 16 Sep 2021 (v2), last revised 22 Apr 2022 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

MULTIOBJECTIVE HYPERVOLUME BASED ISOOMOO ALGORITHMS CONVERGE WITH AT LEAST SUBLINEAR SPEED TO THE ENTIRE PARETO FRONT

EUGÉNIE MARESCAUX AND ANNE AUGER

Abstract. In multiobjective optimization, one is interested in finding a good approximation of the Pareto set and the Pareto front, i.e the sets of best compromises in the decision and objective spaces, respectively. In this context, we introduce a new algorithm framework, Incremental SingleObjective Optimization for MultiObjective Optimization (ISOOMOO) for approximating the Pareto front with an increasing number of points. We focus on HV-ISOOMOO, its instantiation with the hypervolume indicator, a set-quality indicator which is widely used for algorithms design and performance assessment. HV-ISOOMOO algorithms approximate the Pareto front by greedily maximizing the hypervolume. We study the convergence to the entire Pareto front of HV-ISOOMOO coupled with perfect singleobjective optimization. The convergence is defined as the convergence of the hypervolume of the sets of all meta-iterations incumbents towards the hypervolume of the Pareto front. We prove tight lower bounds on the speed of convergence for convex and bilipschitz Pareto fronts in $O(1/n^c)$ with $c = 1$ and $c \leq 1$, respectively. The index n denotes the number of meta-iterations of HV-ISOOMOO. For convex Pareto fronts, the convergence is in $\Theta(1/n)$, namely the fastest convergence achievable by a biobjective optimization algorithm. These are the first results on the speed of convergence of multiobjective optimization algorithms towards the entire Pareto front. We also analyze theoretically the asymptotic convergence behavior.

Key words. multiobjective optimization, convergence, hypervolume, Pareto front

AMS subject classifications. 90C29, 90C30

1. Introduction. Real-world problems often involve the optimization of several conflicting objectives. The solution of such problems is the set of non-dominated *decision vectors* (vectors of the search space), the *Pareto set*. It is defined as the set of solutions that cannot be improved along one objective without degrading along another one. Its image in the objective space is the Pareto front. A decision maker is then often involved to choose, based on its preferences, a single best compromise. The shape of the Pareto front informs on the trade-off between objectives. Many algorithms such as evolutionary algorithms approximate the Pareto front with a number of points fixed in the beginning. But some algorithms, in particular stemming from direct search methods [1, 7, 10, 11] aim at approximating the Pareto set or Pareto front with as many well-distributed points as possible. Ideally, the quality of the Pareto front approximation increases with time without stagnating and such algorithms are referred to in the sequel as *anytime* algorithms.

The speed of convergence towards a critical decision vector or a vector of the Pareto front have been examined for many algorithms such as $(1+1)$ evolutionary multiobjective algorithms [5] or Newton's method [14]. Convergence speeds are typically similar to the ones obtained for singleobjective optimization. They both apply to a convergence towards a single point. Their analysis is sometimes reduced to the study of the convergence of a singleobjective optimization algorithm. The convergence of anytime algorithms towards the whole Pareto set or front is of a different kind because these are sets and not points. It has already been theoretically investigated for some algorithms [10] and more abstract frameworks [19], but analysis of the speed of convergence are missing. Additionally, empirical studies typically focus on determining which algorithm is faster and do not provide precise information on the speed of convergence such as order of convergence or

complexity. Yet, while largely overlooked, investigating the speed of convergence either theoretically or empirically is important. In this context, it has been proven that convergence of biobjective optimization algorithms towards the whole Pareto front is always sublinear in the number of function evaluations, at least when measuring convergence with the hypervolume indicator [16] or the multiplicative ϵ -indicator [8], and thus much smaller than typical speeds of convergence to a single point. The hypervolume and the multiplicative ϵ -indicator are set-quality indicators widely used in multiobjective optimization, both to guide algorithms and for performance assessment. The hypervolume is at the core of all known strictly Pareto-compliant indicators [20].

In this paper, we introduce a new anytime algorithm framework, Incremental SingleObjective Optimization for MultiObjective Optimization (ISOOMOO). We focus on its instantiation with the hypervolume indicator, namely HV-ISOOMOO algorithms. These algorithms try to greedily maximize the hypervolume by adding points with the largest hypervolume contribution. These points are obtained by running a singleobjective optimizer. A greedy idea has already been used in the selection part of some multiobjective optimization evolutionary algorithms such as SMS-EMOA to find a set of p points with a large hypervolume [6]. The hypervolume of such discrete greedy approximation is proven to be at least $(e-1)/e$ times the one of the p -optimal distribution [17]. To the best of our knowledge, we provide the first continuous equivalent of this result. We investigate the speed of convergence of HV-ISOOMOO towards the whole Pareto front in the ideal case of perfect singleobjective optimization, measuring the convergence with the hypervolume. For convex and bilipschitz Pareto fronts, we prove that the convergence is in $O(1/n^c)$ with $c = 1$ and $c \leq 1$, respectively, with n being the number of singleobjective optimization runs performed. For convex Pareto fronts, the convergence is exactly in $\Theta(1/n)$ as no biobjective algorithm can converge faster to the Pareto front [16]. Additionally, we prove that for simultaneously bilipschitz and smooth enough Pareto fronts doubling the number of points in the approximation divides the optimality gap by a factor which converges asymptotically to two. In the proof process, we obtain interesting intermediary results such as bounds on the normalized maximum hypervolume and a geometric interpretation of optimality conditions.

The paper is organized as follows. In Section 2, we lay the foundations of the problem. In Section 3, we prove preliminary results later used to investigate convergence. In Section 4, we derive lower bounds on the speed of convergence of HV-ISOOMOO coupled with perfect singleobjective optimization and an insight on its asymptotic convergence behavior.

Notations and conventions. For $a, b \in \mathbb{N}$, we note $\llbracket a; b \rrbracket$ the set $\{a, a+1, \dots, b-1, b\}$. For a vector $u \in \mathbb{R}^2$, we note u_1 and u_2 respectively its first and its second coordinate. If the vector notation already contains an index, we separate the two indices with a comma. For simplicity sake, we often replace the set $\{u\}$ by u in the notations. We say that a function $f : \mathbb{R} \rightarrow \mathbb{R}$ is decreasing (resp. strictly decreasing) when for all $x < y$, we have $f(x) \geq f(y)$ (resp. $f(x) > f(y)$). We only consider two-dimensional objective spaces and refer to the Lebesgue measure of a set as its area.

2. Background, algorithm framework and assumptions. We lay in this section the foundations of the problem we analyze. First, we recall some classic concepts of multiobjective optimization. Then, we introduce the ISOOMOO class of algorithms and its hypervolume based instantiation HV-ISOOMOO. We also formalize a mathematical abstraction of HV-ISOOMOO coupled with perfect singleobjective optimization, the

greedy set sequences. Finally, we examine our assumptions on the biobjective optimization problem.

2.1. Biobjective optimization problems, the Pareto front and the hypervolume indicator. We consider a biobjective minimization problem:

$$(2.1) \quad \min_{x \in \Omega \subset \mathbb{R}^d} F(x)$$

with $F : \Omega \subset \mathbb{R}^d \rightarrow \mathbb{R}^2 : x \mapsto (F_1(x), F_2(x))$. We define two dominance relations for vectors in the objective space. We say that u weakly dominates v denoted by $u \preceq v$ if $u_1 \leq v_1$ and $u_2 \leq v_2$ and that u dominates v denoted by $u \prec v$ if $u \preceq v$ and $u \neq v$. A vector of the objective space \mathbb{R}^2 is said *feasible* when it belongs to $F(\Omega)$. Solving the optimization problem consists in finding a good approximation of the *Pareto front*, the set of non-dominated feasible vectors, $\{F(X) : X \in \Omega, \forall Y \in \Omega, F(Y) \not\prec F(X)\}$. We restrict ourselves to Pareto fronts with an explicit representation:

$$(2.2) \quad \text{PF}_f = \{(x, f(x)) : x \in [x_{\min}, x_{\max}]\}$$

with $f : \mathbb{R} \mapsto \mathbb{R}$ decreasing. We denote by $u_{\min} := (x_{\min}, f(x_{\min}))$ and $u_{\max} := (x_{\max}, f(x_{\max}))$ the extreme vectors of the Pareto front. Likewise, we denote by $\tilde{u}_{\min, r} := (\tilde{x}_{\min, r}, f(\tilde{x}_{\min, r}))$ and $\tilde{u}_{\max, r} := (\tilde{x}_{\max, r}, f(\tilde{x}_{\max, r}))$ the extremes vectors of the part of the Pareto front dominating a reference point r , with $\tilde{x}_{\min, r} := \max(x_{\min}, f^{-1}(r_2))$ and $\tilde{x}_{\max, r} := \min(x_{\max}, r_1)$. The vector $(x_{\max}, f(x_{\min}))$ is called the *nadir point*. All these notations are illustrated in Figure 1.




Fig. 1: Illustration of notations. The extreme vectors u_{\min} and u_{\max} and the nadir point (leftmost) ; the extreme vectors relative to the reference point r $\tilde{u}_{\min, r}$ and $\tilde{u}_{\max, r}$ (left) ; three vectors u_1 , u_2 and u_3 and the regions weakly dominated by them and dominating r , $\mathcal{D}_{u_1}^r$ (red), $\mathcal{D}_{u_2}^r$ (blue) and $\mathcal{D}_{u_3}^r$ (dotted) (right) ; the greedy set $\mathcal{S}_3 = \{v_1, v_2, v_3\}$, its four gap regions (blue) and the associated reference points (rightmost).

The *hypervolume* with respect to a reference point r of a set S of objective vectors is the Lebesgue measure of the region of the objective space dominated by S and strictly dominating the reference point r . We denote it $\text{HV}_r(S)$. When no vector of the Pareto front dominates the reference point r , $\text{HV}_r(S) = 0$ for any set S of feasible points of the objective space. Since this particular case is not interesting, we only consider reference points dominated by at least one vector of the Pareto front from now on. We refer to such reference points as *valid*.

121 The region of the objective space dominated by S and dominating r (see the righthand
122 plot of Figure 1) is denoted by \mathcal{D}_S^r and formally defined as:

$$123 \quad (2.3) \quad \mathcal{D}_S^r = \{w \in \mathbb{R}^2 : \exists u \in S : u \preceq w \prec r\} .$$

125 The hypervolume of a set S relative to the reference point r equals $\lambda(\mathcal{D}_S^r)$ with $\lambda(\cdot)$ being
126 the Lebesgue measure. The set \mathcal{D}_S^r is the union of the \mathcal{D}_u^r for $u \in S$, \mathcal{D}_u^r being the
127 rectangle $[u_1, r_1] \times [u_2, r_2]$ when u dominates r and \emptyset otherwise, see the righthand plot of
128 Figure 1. Note that the \mathcal{D}_u^r are not disjoint.

129 We use the hypervolume to characterize the convergence of a set S of objective vectors
130 to the entire Pareto front. For a fixed valid reference point r , a set S is said to converge
131 to the Pareto front when the hypervolume difference $\text{HV}_r(\text{PF}_f) - \text{HV}_r(S)$ converges
132 to 0. We define the *optimality gap* of S with respect to a valid reference point r as
133 $\text{HV}_r(\text{PF}_f) - \text{HV}_r(S)$. Another quantity of interest is how much adding a vector to a set
134 affects its hypervolume. The *hypervolume improvement* with respect to r of the vector u
135 to the set S is $\text{HVI}_r(u, S) = \text{HV}_r(S \cup \{u\}) - \text{HV}_r(S)$. We also use the term hypervolume
136 improvement to refer to the hypervolume increase of an increasing¹ sequence of sets. More
137 precisely, the hypervolume improvement at iteration n of an increasing sequence $(\mathcal{S}_n)_{n \in \mathbb{N}^*}$
138 is $\text{HV}_r(\mathcal{S}_{n+1}) - \text{HV}_r(\mathcal{S}_n)$.

139 **2.2. The ISOOMOO framework, its HV-ISOOMOO instantiation and the**
140 **associated greedy set sequences.** The Incremental SingleObjective Optimization for
141 MultiObjective Optimization (ISOOMOO) framework builds incrementally an increasing
142 sequence $(\mathcal{I}_n)_{n \in \mathbb{N}^*}$ of sets of vectors of the objective space. The pseudocode of ISOOMOO
143 is given in Algorithm 2.1, where the current value of \mathcal{I}_n is denoted by \mathcal{I} . At each so-
144 called meta-iteration, a singleobjective maximization algorithm SOOPTIMIZER (line 2 in
145 Algorithm 2.1) is run on the criterion $X \in \Omega \subset \mathbb{R}^d \mapsto J(\mathcal{I}, X)$ and the resulting solution is
146 added to \mathcal{I} (line 3 in Algorithm 2.1). We use the term meta-iteration to separate between
147 the (meta-)iterations of ISOOMOO and the iterations of SOOPTIMIZER. Since the set
148 \mathcal{I} is composed of the final objective incumbents of previous runs of SOOPTIMIZER and
149 (ideally) provides an approximation of the Pareto front, we call it *final incumbents Pareto*
150 *front approximation*.

151 The singleobjective optimization procedure may vary between meta-iterations. More
152 precisely, the run of SOOPTIMIZER depends on data about precedent runs stored in
153 D (line 4 in Algorithm 2.1). This allows to alternate between various singleobjective
154 optimization algorithms with different features, but also to adapt their initialization. This
155 could be done by storing in D an iteration index and the final search space incumbents
156 of SOOPTIMIZER runs.

Algorithm 2.1 ISOOMOO Framework

```

1: while not stopping criterion do
2:    $Y, d \leftarrow \text{SOOPTIMIZER}(X \mapsto J(\mathcal{I}, X), D)$ 
3:    $\mathcal{I} \leftarrow \mathcal{I} \cup \{F(Y)\}$  # update of the approximation of the Pareto front
4:    $D \leftarrow D \cup \{d\}$  # update of the data collected
5: end while

```

¹A sequence of set $\{A_n, n \geq 0\}$ is increasing if the following inclusions $A_0 \subset A_1, \dots \subset A_n \subset \dots$ hold.

157 In this paper, we study HV-ISOOMOO, an instantiation of ISOOMOO for which the
 158 criterion $J(\mathcal{I}, \cdot)$ relates to the hypervolume improvement to \mathcal{I} . Formally, HV-ISOOMOO
 159 is a class of algorithms derived from ISOOMOO for which the maximization of the cri-
 160 terion J is *compliant* with the maximization of the hypervolume improvement as defined
 161 below.

162 *Assumption 2.1.* (Compliance to HVI_r maximization) The maximization of a crite-
 163 rion J as in ISOOMOO is *compliant* with the maximization of HVI_r if for any set \mathcal{I} of
 164 objective vectors, we have

$$165 \quad (2.4) \quad \operatorname{argmax}_{X \in \mathbb{R}^d} J(\mathcal{I}, X) = \operatorname{argmax}_{X \in \mathbb{R}^d} \text{HVI}_r(F(X), \mathcal{I}) .$$

166 We define an HV-ISOOMOO algorithm relative to the reference point r as an ISOOMOO
 167 algorithm as described in [Algorithm 2.1](#) where the criterion J satisfies [Assumption 2.1](#). At
 168 each meta-iteration n , an HV-ISOOMOO algorithm seeks a feasible vector maximizing
 169 the hypervolume improvement to the final incumbents Pareto front approximation \mathcal{I}_n .
 170 Ideally, when n goes to infinity, the non-dominated subset of $(\mathcal{I}_n)_{n \in \mathbb{N}^*}$ converges to the
 171 (entire) Pareto front which maximizes the hypervolume. In other words, HV-ISOOMOO
 172 algorithms try to approximate the Pareto front with a greedy approach.

173 *DEFINITION 2.2.* We define the convergence of an HV-ISOOMOO algorithm as the
 174 convergence of $\text{HV}_r(\mathcal{I}_n)$ towards $\text{HV}_r(\text{PF}_f)$.

175 The performance of a specific HV-ISOOMOO algorithm depends crucially on the
 176 choice of the criterion J . In this respect, $\text{HVI}_r(\mathcal{I}, F(\cdot))$ itself is not a good candidate for
 177 $J(\mathcal{I}, \cdot)$. Indeed, it is constant equal to zero in the region dominated by \mathcal{I} , which makes it
 178 difficult to optimize. A criterion whose maximization is compliant with the maximization
 179 of the hypervolume improvement and designed to be easier to optimize has already been
 180 introduced in [\[18\]](#) under the name *uncrowded hypervolume improvement* (UHVI). For
 181 $F(X)$ not dominated by \mathcal{I} , UHVI_r and HVI_r are equal. Otherwise, in the region where
 182 the hypervolume improvement is null, UHVI_r is negative and equals minus the distance
 183 to the empirical non-dominated front of the set \mathcal{I} relative to r . It is easy to see that
 184 UHVI_r satisfies [\(2.4\)](#).

185 The choice of SOOPTIMIZER also plays a key role in the performance of an HV-
 186 ISOOMOO algorithm. In this paper, we analyze the HV-ISOOMOO framework under
 187 the assumption of perfect singleobjective optimization formalized below.

188 *Assumption 2.3* (Perfect singleobjective optimization). At every meta-iteration n ,
 189 for any final incumbents Pareto front approximation \mathcal{I}_n , the run of SOOPTIMIZER (line 2
 190 in [Algorithm 2.1](#)) returns $Y \in \operatorname{argmax}_{X \in \Omega} J(\mathcal{I}_n, F(X))$.

191 The assumption of perfect singleobjective optimization is reminiscent to the assump-
 192 tion of perfect line search which is common in the analysis of gradient based methods [\[12\]](#).
 193 Under this assumption, all choices of criterions verifying [Assumption 2.1](#) are equivalent.
 194 The convergence of HV-ISOOMOO coupled with perfect singleobjective optimization is a
 195 necessary condition for the soundness of the approach. Furthermore, lower bounds on the
 196 speed of convergence of a real instantiation of HV-ISOOMOO could be obtained by com-
 197 bining lower bounds on the speed of convergence of HV-ISOOMOO under [Assumption 2.3](#)
 198 of perfect singleobjective optimization with the ones of singleobjective optimization algo-
 199 rithms.

200 We introduce below mathematical abstractions of the HV-ISOOMOO framework un-
 201 der [Assumption 2.3](#) of perfect singleobjective optimization, greedy sequences and greedy
 202 set sequences.

203 **DEFINITION 2.4** (Greedy sequence and greedy set sequence). *Given a valid reference*
 204 *point r , we define as greedy sequence relative to r , a sequence $(v_n)_{n \in \mathbb{N}^*}$ satisfying*

$$205 \quad (2.5) \quad v_1 \in \arg \max_{v \in F(\Omega)} HV_r(v) \text{ and}$$

$$206 \quad (2.6) \quad v_{n+1} \in \arg \max_{v \in F(\Omega)} HV_r(\{v_1, \dots, v_n, v\}) \text{ for all } n \geq 1 .$$

207
 208 *The greedy set sequence $(\mathcal{S}_n)_{n \in \mathbb{N}^*}$ associated to the greedy sequence $(v_n)_{n \in \mathbb{N}^*}$ is composed*
 209 *of the greedy sets $\mathcal{S}_n := \{v_k, k \leq n\}$.*

210 There is a bijection between greedy sequences and greedy set sequences. The n -th element
 211 of the greedy sequence $(v_n)_{n \in \mathbb{N}^*}$ associated to a greedy set sequence $(\mathcal{S}_n)_{n \in \mathbb{N}^*}$ is simply
 212 the unique element of $\mathcal{S}_n \setminus \mathcal{S}_{n-1}$ if $n > 1$ and of \mathcal{S}_1 if $n = 1$.

213 The recurrence relation of the greedy sequence (2.6) is equivalent to v_{n+1} belonging
 214 to $\arg \max_{v \in F(\Omega)} HVI_r(v, \mathcal{S}_n)$ for all $n \geq 1$. It is immediate to see that under [Assump-](#)
 215 [tion 2.3](#), the final incumbents generated by HV-ISOOMOO constitute a greedy sequence
 216 while the final incumbents Pareto front approximations form the associated greedy set
 217 sequence $(\mathcal{I}_n)_{n \in \mathbb{N}^*}$. The indices n of both sequences correspond to HV-ISOOMOO meta-
 218 iterations. In this paper, we derive convergence results for greedy set sequences, which
 219 transfer to HV-ISOOMOO under [Assumption 2.3](#).

220 As we will see in [Subsection 3.1](#), the problem of maximizing the hypervolume improve-
 221 ment to a fixed set can be rewritten as the maximum of a finite number of hypervolume
 222 maximization problems. Therefore, we can infer from [4, Theorem 1] that as soon as the
 223 Pareto front is lower semi-continuous, there exists a greedy sequence, and thus a greedy
 224 set sequence.

225 **PROPOSITION 2.5.** *If the Pareto front is described by a lower semi-continuous func-*
 226 *tion f , then there exists a greedy sequence $(v_n)_{n \in \mathbb{N}^*}$ relative to any valid reference point.*

227 *Proof.* If f is lower semi-continuous, then for any valid reference point r , the maxi-
 228 mum of $HV_r(\cdot)$ exists, see [4, Theorem 1]. Therefore, there exists a vector verifying (2.5)
 229 and the problem of maximizing the maximum of a finite number of hypervolume functions
 230 defined in (3.3) admits a solution. Since (2.6) and (3.3) are equivalent by [Lemma 3.5](#), we
 231 can build a sequence $(v_n)_{n \in \mathbb{N}^*}$ verifying (2.5) and (2.6), namely a greedy sequence. \square

232 Additionally, since the hypervolume indicator associated to a valid reference point is
 233 strictly Pareto-compliant (see [15]), this sequence is composed of vectors of the Pareto
 234 front.

235 **PROPOSITION 2.6.** *If the Pareto front is described by a lower semi-continuous func-*
 236 *tion f , then any vector of a greedy sequence relative to a valid reference point r belongs to*
 237 *the Pareto front. Consequently, for such Pareto front and reference point and under [As-](#)*
 238 [sumption 2.3](#), *all final incumbents Pareto front approximations \mathcal{I}_n of an HV-ISOOMOO*
 239 *algorithm relative to r are subsets of the Pareto front.*

240 *Proof.* Since for any valid reference point r , $HV_r(\cdot)$ is strictly Pareto-compliant [15],
 241 its maximum is non-dominated and belongs to the Pareto front. Thus, in particular, a
 242 vector v_1 verifying (2.5) belongs to the Pareto front. Additionally, by [Lemma 3.5](#), every

243 solution of (2.6) verifies (3.3) and is solution of at least one hypervolume maximization
 244 problem, and thus also belong to the Pareto front. \square

245 Yet, in general, there exists more than one greedy sequence, and thus greedy set sequence.
 246 For example, there are infinitely many greedy sequences associated to any affine Pareto
 247 front with a reference point dominating the nadir point. This statement relies on the fact
 248 that the unique maximizer of the hypervolume relative to a reference point r dominating
 249 the nadir point is the middle of the section of the Pareto front dominating r , see [3,
 250 Theorem 5]. As a consequence, the middle of the section of the Pareto front dominating r
 251 is the only candidate for v_1 but v_2 can be either at $1/4$ or at $3/4$ of this section. Similarly,
 252 v_3 has to be in the position where v_2 is not but v_4 can be at $1/8, 3/8, 5/8$ or $7/8$ of the
 253 section of the Pareto front dominating r . For any n , we can find an iteration m such
 254 that v_m can be placed at 2^n different points, whatever the $m - 1$ first terms of the greedy
 255 sequence are.

256 **2.3. Assumptions on the Pareto front and the objective functions.** We
 257 present and discuss here the assumptions on the function f describing the Pareto front
 258 under which we derive convergence results. We typically assume that the function f is
 259 bilipschitz, convex or simultaneously bilipschitz and with a Hölder continuous derivative.
 260 Under any of these three assumptions, f is continuous. For the sake of conciseness, we
 261 transfer the properties of f to the Pareto front. For example, we call *convex Pareto*
 262 *front* a Pareto front described by a convex function. We recall that a function f is
 263 Hölder continuous with exponent α , namely $\mathcal{C}^{1,\alpha}$, when there exists $H \geq 0$ such that
 264 $|f(x) - f(y)| \leq H \times |x - y|^\alpha$ for all x, y [13]. We note $[f]_\alpha$ the *minimum Hölder coefficient*
 265 with respect to the exponent α of a $\mathcal{C}^{1,\alpha}$ function f , that is $[f]_\alpha := \sup_{x \neq y} \frac{|f(x) - f(y)|}{|x - y|^\alpha}$.
 266 When needed, we detail the bilipschitz constants and say that a bilipschitz function f is
 267 (L_{\min}, L_{\max}) -bilipschitz if for all $x, y \in [x_{\min}, x_{\max}]$, we have $L_{\min} \times |x - y| \leq |f(x) - f(y)| \leq$
 268 $L_{\max} \times |x - y|$ where $L_{\max} \geq L_{\min} > 0$. We also talk of *affine Pareto fronts* when
 269 $f(x) = ax + b$ with $a < 0$ and $b \in \mathbb{R}$. As they form a line in the biobjective case, they are
 270 usually referred to as linear Pareto fronts. They provide good examples to illustrate a
 271 point and help to understand the results we prove on the asymptotic convergence behavior.
 272 We remind below sufficient conditions on the search space and on the objective func-
 273 tions which guarantee that f is convex and bilipschitz.

274 **PROPOSITION 2.7.** *Given a biobjective minimization problem as in (2.1) whose Pareto*
 275 *front is described by a function f . If F_1 and F_2 are respectively $(L_{\min,1}, L_{\max,1})$ -bilipschitz*
 276 *and $(L_{\min,2}, L_{\max,2})$ -bilipschitz, then f is $(\frac{L_{\min,2}}{L_{\max,1}}, \frac{L_{\max,2}}{L_{\min,1}})$ -bilipschitz. If the search space*
 277 *Ω and the objective functions F_1 and F_2 are convex, then f is convex.*

278 The proofs of this proposition can be found for instance in [16]. The conditions on F_1 ,
 279 F_2 and Ω are sufficient but non-necessary conditions. Indeed, adding small discontinuity in
 280 the objective functions far from the Pareto set makes them non-convex and non-bilipschitz
 281 without modifying the Pareto front.

282 Representing F_1 values on the absciss and F_2 values on the ordinate instead of the
 283 converse is an arbitrary choice. When f is a bijection, if we chose to represent the F_2
 284 values on the x -axis instead of on the y -axis, we would have another representation of
 285 the Pareto front : $\{(y, f^{-1}(y)) : y \in [f(x_{\max}); f(x_{\min})]\}$. If so, the inverse function f^{-1}
 286 would play the role of f . It is interesting to notice that the choice of the objective function
 287 represented on the horizontal axis does not impact whether the function characterizing

the Pareto front is bilipschitz or convex. Indeed, f being bilipschitz is equivalent to both f and f^{-1} being lipschitz. Additionally, we can prove that given that the function f is decreasing, f being convex is equivalent to its inverse f^{-1} being convex. The proof of this property is straightforward. If f is convex, then for all $x, y \in \mathbb{R}$, $f\left(\frac{f^{-1}(x)+f^{-1}(y)}{2}\right)$ is smaller than $\frac{f(f^{-1}(x))+f(f^{-1}(y))}{2}$, that is $\frac{x+y}{2}$. Since f and therefore f^{-1} are decreasing, by composing by f^{-1} each side of the inequality, we obtain a characterization of the convexity of f^{-1} : for all $x, y \in \mathbb{R}$, $\frac{f^{-1}(x)+f^{-1}(y)}{2}$ is larger than $f^{-1}\left(\frac{x+y}{2}\right)$.

3. Preliminary results. In this section, we present preliminary results which are crucial for the analysis of the convergence of HV-ISOOMOO. While we expose them as tools for convergence analysis, they are also interesting for their own sake.

3.1. Decomposition of the optimality gap using gap regions. The optimality gap is the Lebesgue measure of the *total gap region* introduced below.

DEFINITION 3.1. *The total gap region of S with respect to a fixed valid reference point r , \mathcal{G}_S^r , is defined as the region of the objective space which dominates r and is weakly dominated by PF_f but not by S , namely $\mathcal{D}_{PF_f}^r \setminus \mathcal{D}_S^r$.*

We introduced \mathcal{D}_S^r in (2.3). Its Lebesgue measure is $HV_r(S)$.

When S is a subset of the Pareto front dominating the reference point r , the total gap region \mathcal{G}_S^r has a particular shape which can be visualized in the rightmost plot of Figure 1. It can be decomposed into the disjoint union of $|S| + 1$ sets of the form $\mathcal{D}_{S'}^{r'}$ that are formally defined below.

DEFINITION 3.2 (Gap regions, gaps and associated reference points). *Let $S = \{v_1, \dots, v_n\}$ be a set of n distinct vectors of the Pareto front dominating a valid reference point r . Let σ be the permutation ordering the v_i by increasing F_1 values: $v_{\sigma(1),1} < v_{\sigma(2),1} < \dots < v_{\sigma(n),1}$.*

- *For all $i \in \llbracket 1, n+1 \rrbracket$, the i -th gap region of the set S , $\mathcal{G}_{S,i}^r$, is the set $\mathcal{D}_{PF_f}^{r_i}$ with the associated reference points r_i being $r_1 = (v_{\sigma(1),1}, r_2)$, $r_{n+1} = (r_1, v_{\sigma(n),2})$ and $r_i = (v_{\sigma(i),1}, v_{\sigma(i-1),2})$ for all $i \in \llbracket 2, n \rrbracket$.*
- *We refer to $\mathcal{G}_{S,1}^r$ and $\mathcal{G}_{S,n+1}^r$ as the left and the right extreme gap region of S , respectively.*

The left (resp. right) extreme gap region is empty when the left (resp. right) extreme vector of the Pareto front belongs to S . Non-extreme gap regions are never empty. The total gap region is the disjoint union of the gap regions: $\mathcal{G}_S^r = \dot{\cup}_{i=1}^{n+1} \mathcal{G}_{S,i}^r$. This decomposition of the total gap region, and thus of the optimality gap, is the cornerstone of the convergence analysis. Since the area of a gap region $\mathcal{G}_{S,i}^r$ is $HV_{r_i}^r(PF_f)$, we can write the optimality gap as the sum of $n+1$ hypervolumes of the Pareto front with respect to different reference points.

LEMMA 3.3. *At any iteration n , the optimality gap of a greedy set sequence with respect to a valid reference point can be decomposed as*

$$(3.1) \quad HV_r(PF_f) - HV_r(\mathcal{S}_n) = \sum_{i=1}^{n+1} HV_{r_i}^r(PF_f) .$$

328 *Proof.* The optimality gap at iteration n is the Lebesgue measure of the total gap
 329 region $\mathcal{G}_{\mathcal{S}_n}^r$, which is the disjoint union of the gap regions $\mathcal{G}_{\mathcal{S}_n,i}^r = \mathcal{D}_{\text{PF}_f}^{r_i^n}$. Therefore, the
 330 optimality gap equals $\sum_{i=1}^{n+1} \lambda(\mathcal{D}_{\text{PF}_f}^{r_i^n}) = \sum_{i=1}^{n+1} \text{HV}_{r_i^n}(\text{PF}_f)$. \square

331 Additionally, we can express the hypervolume improvement of any vector to \mathcal{S}_n as an
 332 hypervolume. It is a trivial assertion for vectors which do not dominate \mathcal{S}_n . For other
 333 vectors, the reference point depends on the gap region to which the vector belongs.

334 LEMMA 3.4. *Let $(\mathcal{S}_n)_{n \in \mathbb{N}^*}$ be a greedy set sequence relative to a valid reference point*
 335 *r . At any iteration n , for any u belonging to the i -th gap region of \mathcal{S}_n , $\mathcal{G}_{\mathcal{S}_n,i}^r$, we have*

$$336 \quad (3.2) \quad \text{HVI}_r(u, \mathcal{S}_n) = \text{HV}_{r_i^n}(u) \quad .$$

337 *Proof.* The hypervolume improvement of any $u \in \mathcal{G}_{\mathcal{S}_n,i}^r$ is the Lebesgue-measure of
 338 the intersection between $\mathcal{G}_{\mathcal{S}_n,i}^r$ and \mathcal{D}_u^r . Therefore, it is equal to $\lambda(\mathcal{D}_u^{r_i^n})$, that is $\text{HV}_{r_i^n}(u)$. \square
 339 We can now reformulate the recurrence relation defining a greedy sequence at iteration
 340 $n + 1$. Indeed, picking a vector maximizing the hypervolume improvement to \mathcal{S}_n is
 341 equivalent to pick a vector where the largest value of the maximum of the hypervolumes
 342 with respect to the r_i^n is reached.

343 LEMMA 3.5. *At any iteration n , the recurrence relation satisfied by v_{n+1} , i.e. (2.6),*
 344 *can be reformulated as*

$$346 \quad (3.3) \quad v_{n+1} \in \arg \max_{u \in \text{PF}_f} \max_{i \in [1, n+1]} \text{HV}_{r_i^n}(u) \quad .$$

347 *Proof.* The hypervolume improvement of any vector u to \mathcal{S}_n is $\max_{i \in [1, n+1]} \text{HV}_{r_i^n}(u)$.
 348 It is a consequence of Lemma 3.4 and of the fact that the hypervolume with respect to r_i^n
 349 is null outside the i -th gap region of \mathcal{S}_n . Additionally, v_{n+1} belongs to the Pareto front
 350 by Proposition 2.6. Thus, (2.6) is equivalent to (3.3). \square

351 Similarly, we can express the decrease of the optimality gap at iteration $n + 1$,
 352 $\text{HV}_r(\mathcal{S}_{n+1}) - \text{HV}_r(\mathcal{S}_n)$, as the maximum of $n + 1$ hypervolume maximization problems.

353 LEMMA 3.6. *Let $(\mathcal{S}_n)_{n \in \mathbb{N}^*}$ be a greedy set sequence relative to a valid reference point*
 354 *r . The hypervolume improvement at iteration $n + 1$ equals*

$$356 \quad (3.4) \quad \text{HV}_r(\mathcal{S}_{n+1}) - \text{HV}_r(\mathcal{S}_n) = \max_{u \in \text{PF}_f} \max_{i \in [1, n+1]} \text{HV}_{r_i^n}(u) \quad .$$

357 *Proof.* The hypervolume improvement $\text{HV}_r(\mathcal{S}_{n+1}) - \text{HV}_r(\mathcal{S}_n)$ is the hypervolume
 358 improvement of v_{n+1} to \mathcal{S}_n . With the same arguments as in the proof of Lemma 3.5, we
 359 can prove that it equals $\max_{u \in \text{PF}_f} \max_{i \in [1, n+1]} \text{HV}_{r_i^n}(u)$. \square

360 **3.2. Lower bound of the normalized maximum hypervolume for convex**
 361 **Pareto fronts.** In this section and the next one, we provide bounds on the maximum
 362 hypervolume achievable by a single feasible vector normalized by the maximum hypervol-
 363 ume of a feasible set: $\frac{\max_{u \in \text{PF}_f} \text{HV}_r(u)}{\text{HV}_r(\text{PF}_f)}$. We refer to this ratio as the *normalized maximum*
 364 *hypervolume* with respect to r . Bounds on the normalized maximum hypervolume are ex-
 365 ploited in Section 4 to provide bounds on the speed of convergence of greedy set sequences
 366 towards the Pareto front.

368 The hypervolume relative to a reference point r of a vector $u = (x, f(x))$ of the Pareto
 369 front is $\text{HV}_r(u) = (r_1 - x) \times (r_2 - f(x))$. From this simple formula, we derive in the next
 370 proposition necessary conditions for a vector of the Pareto front to be an hypervolume
 371 maximizer when f has at least left and right derivatives in x^* .

372 **PROPOSITION 3.7.** *Let $x^* \in]x_{\min}, x_{\max}[$ such that $u^* := (x^*, f(x^*))$ maximizes the*
 373 *hypervolume with respect to a valid reference point r . If the function f describing the*
 374 *Pareto front admits left and right derivatives in x^* , respectively $f'_-(x^*)$ and $f'_+(x^*)$, then*

$$375 \quad (3.5) \quad -f'_+(x^*) \leq \frac{r_2 - f(x^*)}{r_1 - x^*} \leq -f'_-(x^*) .$$

377 *Proof.* We define the function $\text{HV}_{x,r}(\cdot)$ as $x \mapsto \text{HV}_r((x, f(x)))$. If x^* maximizes
 378 $\text{HV}_{x,r}(\cdot)$, then the left and the right derivatives of $\text{HV}_{x,r}(\cdot)$ are positive and negative,
 379 respectively. By replacing the left and right derivatives of $\text{HV}_{x,r}(\cdot)$ by their explicit
 380 formulas and reorganizing the terms we obtain (3.5). \square

381 Equation (3.5) states that the slope of the diagonal of the rectangle $\mathcal{D}_{u^*}^r$ is between the
 382 absolute values of the slopes of the right and the left derivatives of f at x^* (see the middle
 383 plot of Figure 2). To the best of our knowledge, this geometric interpretation is new. It
 384 becomes simpler when f is differentiable. Then, the absolute value of the slope of the
 385 tangent of the front at a non-extreme vector u^* is equal to the slope of the diagonal of
 386 the rectangle $\mathcal{D}_{u^*}^r$ (see the lefthand plot of Figure 2).




Fig. 2: Left and middle: Two convex Pareto fronts and their respective hypervolume maximizers u^* , one differentiable (left) and one non-differentiable (middle). The slopes of the two dotted lines, namely PF_g and the diagonal of $\mathcal{D}_{u^*}^r$, are equal. Right: The Pareto front PF_ϵ and the hypervolume maximizer u_ϵ^* for $\epsilon = 1/3$ and $r = (1, 1)$.

387 **COROLLARY 3.8.** *Let $x^* \in]x_{\min}, x_{\max}[$ be such that $u^* := (x^*, f(x^*))$ maximizes the*
 388 *hypervolume with respect to a valid reference point r . If the Pareto front is described by*
 389 *a differentiable function f in x^* , then $f'(x^*)$ satisfies*

$$390 \quad (3.6) \quad -f'(x^*) = \frac{r_2 - f(x^*)}{r_1 - x^*} .$$

392 *Proof.* It is a direct consequence of Proposition 3.7 \square

393 A convex function may not be differentiable, but it always has left and right de-
 394 rivatives. It is also above its left and right tangent lines respectively on the left and

on the right of x^* . Therefore, [Proposition 3.7](#) implies that the affine function $g : x \mapsto f(x^*) - \frac{r_2 - f(x^*)}{r_1 - x^*} \times (x - x^*)$ is a minorant of f . It is the key idea of the proof of the following lower bound on the normalized maximum hypervolume.

PROPOSITION 3.9. *If the Pareto front is described by a convex function f , then the following lower bound on the normalized maximum hypervolume with respect to any valid reference point r holds:*

$$(3.7) \quad \frac{\max_{u \in PF_f} HV_r(u)}{HV_r(PF_f)} \geq \frac{1}{2}$$

where the inequality is an equality if and only if the Pareto front is affine and r dominates the nadir point.

Proof. As explained in the above paragraph, the convexity of f implies that the affine function $g : x \mapsto f(x^*) - \frac{r_2 - f(x^*)}{r_1 - x^*} \times (x - x^*)$ is a minorant of f . Therefore, $PF_g := \{g(x) : x \in [x_{\min}, x_{\max}]\}$ dominates PF_f , and thus has a higher hypervolume. We denote $L_1 := r_1 - \tilde{x}_{\min, r}$ and $L_2 := r_2 - f(\tilde{x}_{\max, r})$ the lengths of the rectangle $\mathcal{R} := [\tilde{x}_{\min, r}, r_1] \times [f(\tilde{x}_{\max, r}), r_2]$. We denote $l_1 := r_1 - x^*$ and $l_2 := r_2 - f(x^*)$ the lengths of the rectangle $\mathcal{D}_{u^*}^r$. The region of \mathcal{R} which dominates PF_g is a right-angled triangle. Additionally, by definition, the slope of its hypotenuse is l_2/l_1 , and thus the lengths of the other sides are $L_1 - l_1 + (L_2 - l_2) \times \frac{l_1}{l_2}$ and $L_2 - l_2 + (L_1 - l_1) \times \frac{l_2}{l_1}$ (see the middle plot of [Figure 2](#)). Therefore, we have

$$\begin{aligned} HV_r(PF_g) &= \lambda(\mathcal{R}) - \lambda(\{u \in \mathbb{R}^2 : u \in \mathcal{R}, u \preceq PF_g\}) \\ &= L_1 L_2 - \frac{1}{2} \times (L_1 - l_1 + (L_2 - l_2) \times \frac{l_1}{l_2}) \times (L_2 - l_2 + (L_1 - l_1) \times \frac{l_2}{l_1}) \\ &= l_1 l_2 \times \left[-2 + 2 \times \frac{L_2}{l_2} - \frac{1}{2} \times \left(\frac{L_2}{l_2} \right)^2 + 2 \times \frac{L_1}{l_1} - \frac{1}{2} \times \left(\frac{L_1}{l_1} \right)^2 \right]. \end{aligned}$$

For all x , we have $(x - 2)^2 \geq 0$ and thus $2x - \frac{1}{2}x^2 \leq 2$. Therefore, we can conclude that $HV_r(PF_g)$, and thus $HV_r(PF_f)$ is smaller than $2 \times l_1 l_2$, that is $2 \times HV_r(u^*)$. If either $L_1/l_1 \neq 2$ or $L_2/l_2 \neq 2$, the inequality is strict. Thus, when the inequality is an equality, the center of \mathcal{R} belongs to the Pareto front. Since f is convex, it happens only when f is affine and the reference point r dominates the nadir point. Conversely, if both conditions are met, we know that the optimum is in the middle of the Pareto front and that we have the equality (see [\[3, Theorem 5\]](#)). \square

We just proved that one half is a tight lower bound on the normalized maximum hypervolume for convex Pareto fronts. However, except for the trivial upper bound 1, there is no upper bound valid for every convex Pareto front, even when r dominates the nadir point. Here is a simple example which illustrates this. Let consider the convex Pareto front $PF_\epsilon := \{\max(1 - \frac{x}{\epsilon}, \epsilon - \epsilon \times x) : x \in [0, 1]\}$ represented in the righthand plot of [Figure 2](#) and the reference point $r = (1, 1)$. When $\epsilon \leq 1$, PF_ϵ is convex and [\(3.6\)](#) implies that $u_\epsilon^* = (\epsilon \times (1 - \epsilon), \epsilon \times (1 - \epsilon))$ is the unique hypervolume minimizer. Thus, the normalized maximum hypervolume of PF_ϵ for this reference point is equal to $\frac{(1 - \epsilon + \epsilon^2)^2}{1 - \epsilon \times (1 - \epsilon)^2 + (\epsilon - \epsilon^2)^2}$ and converges to 1 when ϵ goes to 0.

3.3. Lower and upper bounds of the normalized maximum hypervolume for bilipschitz Pareto fronts. In this section, we examine lower and upper bounds on the normalized maximum hypervolume in the case of bilipschitz Pareto fronts.

437 We consider two affine fronts with the same left extreme vector as PF_f and slopes
 438 $-L_{\min}$ and $-L_{\max}$, see the lefthand plot of Figure 3. We call them PF_{\min} and PF_{\max} ,
 439 respectively. Formally:

$$440 \quad (3.8) \quad \text{PF}_{\max} := \{(x, f_{\max}(x)) : x \in [x_{\min}, x_{\max}]\} \text{ and}$$

$$441 \quad (3.9) \quad \text{PF}_{\min} := \{(x, f_{\min}(x)) : x \in [x_{\min}, x_{\max}]\}$$

443 with $f_{\min}(x) = f(x_{\min}) - (x - x_{\min}) \times L_{\min}$ and $f_{\max}(x) = f(x_{\min}) - (x - x_{\min}) \times$
 444 L_{\max} . For a (L_{\min}, L_{\max}) -bilipschitz function f , $f_{\min}(x) \leq f(x) \leq f_{\max}(x)$ for all $x \in$
 445 $[x_{\min}, x_{\max}]$, and thus the Pareto front is dominated by PF_{\max} and dominates PF_{\min} .
 These two affine fronts provide bounds on both the hypervolume of the Pareto front




Fig. 3: Left : The Pareto front PF_f surrounded by PF_{\max} (below) and PF_{\min} (above).
 Right : An illustration that $\text{HV}_r(u_{\min}) - \text{HV}_r(\text{PF}_f)$ (■) becomes negligible compared to
 $\text{HV}_r(u_{\min})$ (■) for $r_1 = 1$ and $r_2 \rightarrow \infty$.

446 and the largest hypervolume of a vector on the Pareto front. They are key to prove
 447 the following lower bound on the normalized maximum hypervolume of a (L_{\min}, L_{\max}) -
 448 bilipschitz Pareto front.

450 PROPOSITION 3.10. *If the Pareto front is described by a (L_{\min}, L_{\max}) -bilipschitz func-*
 451 *tion f , then for any valid reference point r , we have*

$$452 \quad (3.10) \quad \frac{\max_{u \in \text{PF}_f} \text{HV}_r(u)}{\text{HV}_r(\text{PF}_f)} \geq \frac{1}{2} \times \frac{L_{\min}}{L_{\max}} .$$

454 *Proof.* The fronts PF_{\max} and PF_{\min} are defined respectively in (3.8) and (3.9). We
 455 note $\Delta_1 := \tilde{x}_{\max, r} - \tilde{x}_{\min, r}$, $\Delta'_1 := r_1 - \tilde{x}_{\max, r}$, $\Delta_2 := r_2 - f(\tilde{x}_{\min, r})$ and $V := \Delta_2 \times (r_1 -$
 456 $\tilde{x}_{\min, r})$, see the lefthand plot of Figure 3. Since the front PF_{\max} dominates the Pareto
 457 front, the hypervolume of PF_f is smaller than the hypervolume of PF_{\max} , $V + L_{\max} \times$
 458 $\Delta_1 \times \Delta'_1 + \frac{1}{2} \times L_{\max} \times \Delta_1^2$. Additionally, since each vector of PF_{\min} is dominated by a
 459 vector of PF_f , the maximum hypervolume of a vector of PF_f is larger than the maximum
 460 hypervolume of a vector of PF_{\min} . The front PF_{\min} being an affine and therefore convex
 461 front, we know by Proposition 3.9 that the maximum hypervolume of a vector of PF_{\min} is
 462 larger than half of $\text{HV}_r(\text{PF}_{\min})$, which is equal to $\frac{1}{2} \times (V + L_{\min} \times \Delta_1 \times \Delta'_1 + \frac{1}{2} \times L_{\min} \times \Delta_1^2)$.
 463 To summarize, the maximum hypervolume of a vector of PF_f is larger than $\frac{1}{2} \times (V + L_{\min} \times$
 464 $\Delta_1 \times \Delta'_1 + \frac{1}{2} \times L_{\min} \times \Delta_1^2)$. Combining the upper bound on the hypervolume of PF_f and the
 465 lower bound on the maximum hypervolume of a vector of PF_f , the normalized maximum

hypervolume is larger than $\frac{\frac{1}{2} \times (V + L_{\min} \times \Delta_1 \times \Delta'_1 + \frac{1}{2} \times L_{\min} \times \Delta_1^2)}{V + L_{\max} \times \Delta_1 \times \Delta'_1 + \frac{1}{2} \times L_{\max} \times \Delta_1^2}$. This quantity is itself larger than $\frac{1}{2} \times \frac{L_{\min} \times \Delta_1 \times \Delta'_1 + \frac{1}{2} \times L_{\min} \times \Delta_1^2}{L_{\max} \times \Delta_1 \times \Delta'_1 + \frac{1}{2} \times L_{\max} \times \Delta_1^2}$. As $V \geq 0$ and $0 < \frac{L_{\min} \times \Delta_1 \times \Delta'_1 + \frac{1}{2} \times L_{\min} \times \Delta_1^2}{L_{\max} \times \Delta_1 \times \Delta'_1 + \frac{1}{2} \times L_{\max} \times \Delta_1^2} < 1$, we conclude that the normalized maximum hypervolume is larger than $\frac{1}{2} \times \frac{L_{\min}}{L_{\max}}$. \square

We cannot guarantee any upper bound strictly inferior to 1 on the normalized maximum hypervolume without adding an assumption on the reference point. Indeed, for a given bounded Pareto front, it is easy to show that the normalized maximum hypervolume goes to 1 for $r_1 = x_{\max}$ and $r_2 \rightarrow \infty$ (see the righthand plot of Figure 3). However, if f is (L_{\min}, L_{\max}) -bilipschitz and r dominates the nadir point, we can prove that the normalized maximum hypervolume is larger than $\frac{1}{2} \times \frac{L_{\max}}{L_{\min}}$. The proof relies on the fact that if the reference point r dominates the nadir point, the vector of an affine front with the largest hypervolume relative to r is its middle (see [3, Theorem 5]), whose hypervolume is half of the hypervolume of the entire Pareto front.

PROPOSITION 3.11. *If the Pareto front is described by a (L_{\min}, L_{\max}) -bilipschitz function f and the reference point r is valid and dominates the nadir point, the following upper-bound on the normalized maximum hypervolume with respect to r holds*

$$(3.11) \quad \frac{\max_{u \in PF_f} HV_r(u)}{HV_r(PF_f)} \leq \frac{1}{2} \times \frac{L_{\max}}{L_{\min}}.$$

Proof. We use the same notations as in the proof of Proposition 3.10. Since r dominates the nadir point, both Δ'_1 , Δ_2 and V equal 0, and thus the hypervolumes of PF_{\max} and PF_{\min} equal $\frac{1}{2} \times L_{\max} \times \Delta_1^2$ and $\frac{1}{2} \times L_{\min} \times \Delta_1^2$, respectively. The domination of PF_{\min} by PF_f implies that the hypervolume of the Pareto front is below $\frac{1}{2} \times L_{\min} \times \Delta_1^2$. Since PF_{\max} is an affine front whose extremes dominate r , its middle is the unique hypervolume maximizer (see [2, Theorem 5]) with an hypervolume equal to $\frac{1}{4} \times L_{\max} \times \Delta_1$. The domination of PF_f by PF_{\max} implies that the maximum hypervolume of a vector of PF_f is larger than $\frac{1}{4} \times L_{\max} \times \Delta_1^2$. Gathering the lower bound on $HV_r(PF_f)$ and the upper bound on the maximum hypervolume of a vector of PF_f , we retrieve (3.11). \square

This upper bound is only relevant for $L_{\max}/L_{\min} < 2$ and is the tightest for $L_{\max} = L_{\min}$, where it achieves the value 1/2. In this paper, we use this upper bound for L_{\max}/L_{\min} close to 1 to analyze the asymptotic convergence behavior of HV-ISOOMOO.

4. Convergence of HV-ISOOMOO coupled with perfect singleobjective optimization. We prove in this section various convergence results for HV-ISOOMOO algorithms coupled with perfect singleobjective optimization. We first prove that when the Pareto front is either convex or bilipschitz, these algorithms converge to the entire Pareto front. We transform the bounds on the normalized maximum hypervolume proven in Section 3 into lower bounds on the speed of convergence. Then, we analyze the asymptotic convergence behavior when the Pareto front is bilipschitz with a Hölder continuous derivative.

To analyze the decrease of the optimality gap with n , we track in which gap regions the vectors of the greedy sequence are inserted over multiple iterations. Naturally, a gap region of \mathcal{S}_n persists in being a gap region as long as no greedy vector is added in this specific gap region. The greedy vector v_{n+1} is said to *fill* the gap region of \mathcal{S}_n to which it belongs. At iteration $n + 1$, this gap region disappears, replaced by two gap regions that we call its *children*. More generally, we say that a gap region is a *descendant* of another gap region when it is a proper subset of this gap region.

510 **4.1. Convergence of HV-ISOOMOO with guaranteed speed of conver-**
511 **gence.** We prove some upper bounds on the relation between the optimality gap at
512 iteration $2n + 1$ and at iteration n . These bounds translate into lower bounds on the
513 speed of convergence of HV-ISOOMOO under [Assumption 2.3](#) of perfect singleobjective
514 optimization. The proof relies on inequalities of the form

$$515 \quad (4.1) \quad \max_{u \in \text{PF}_f} \text{HV}_{r'}(u) \geq C \times \text{HV}_{r'}(\text{PF}_f)$$

517 stated in [Propositions 3.9](#) and [3.10](#) and equations regarding optimality gaps, areas of gap
518 regions and hypervolume improvement presented in [Subsection 3.1](#). A consequence of
519 (4.1) being true for any reference point r' is that the optimality gap at iteration $2n + 1$
520 is at most $(1 - C)$ times the optimality gap at iteration n .

521 We sketch the proof idea in the simple case where each of the v_k ($k \in \llbracket n + 1, 2n + 1 \rrbracket$)
522 is inserted in a distinct gap region of \mathcal{S}_n , see the lefthand plot of [Figure 4](#). Inserting v_k in
523 a gap region leads to an hypervolume improvement larger than C times the area of this
524 gap region by (4.1). Thus, the hypervolume improvement from iteration n to $2n + 1$ is
525 larger than C times the area of the union of all gap regions of \mathcal{S}_n , namely the optimality
gap at iteration n . A detailed proof is presented after the theorem statement.




Fig. 4: Left: A Pareto front where each of the gap regions of \mathcal{S}_3 is filled by one of the greedy vectors v_k for $k \in \llbracket 4, 7 \rrbracket$. It is described by $f(x) = 1 - \sqrt{x}$ for $x \in [0, 1]$. We represent the region $\mathcal{D}_{\mathcal{S}_3}^r$ (yellow), the gap regions of \mathcal{S}_3 (blue) and the regions corresponding to $\text{HVI}_r(v_k, \mathcal{S}_{k-1})$ for $k \in \llbracket 4, 7 \rrbracket$ (orange). Right: The ordered greedy set F_1 -values $w_{i,r}^n$ corresponding to the greedy set \mathcal{S}_3 . The Pareto front is described by $f(x) = \frac{e}{e-1} \times e^{-x} + 1 - \frac{e}{e-1}$ for $x \in [0, 1]$.

526
527 **PROPOSITION 4.1.** *Consider a biobjective optimization problem with a Pareto front*
528 *described by a function f . Any greedy set sequence $(\mathcal{S}_n)_{n \in \mathbb{N}^*}$ relative to a valid reference*
529 *point r satisfies for all n*

$$530 \quad (4.2) \quad \frac{\text{HV}_r(\text{PF}_f) - \text{HV}_r(\mathcal{S}_{2n+1})}{\text{HV}_r(\text{PF}_f) - \text{HV}_r(\mathcal{S}_n)} \leq 1 - \frac{1}{2} \times \frac{L_{\min}}{L_{\max}} \text{ if } f \text{ is } (L_{\min}, L_{\max})\text{-bilipschitz and}$$

$$531 \quad (4.3) \quad \frac{\text{HV}_r(\text{PF}_f) - \text{HV}_r(\mathcal{S}_{2n+1})}{\text{HV}_r(\text{PF}_f) - \text{HV}_r(\mathcal{S}_n)} \leq \frac{1}{2} \text{ if } f \text{ is convex.}$$

533 *Proof.* Fix $n \geq 1$. We note σ a permutation of $\llbracket 1, n + 1 \rrbracket$ such that $n + \sigma(i)$ is the
534 index of the first greedy vector v_k inserted in $\mathcal{G}_{\mathcal{S}_n, i}^r$ when possible. With this choice of σ ,

the i -th gap region of \mathcal{S}_n is a gap region of $\mathcal{S}_{n+\sigma(i)-1}$. As a consequence, the hypervolume improvement to $\mathcal{S}_{n+\sigma(i)-1}$ of any vector u belonging to the i -th gap region of \mathcal{S}_n , $\mathcal{G}_{\mathcal{S}_n,i}^r$, is equal to $\text{HV}_{r_i^n}(u)$ by Lemma 3.5. The hypervolume improvement of the greedy vector $v_{n+\sigma(i)}$ to $\mathcal{S}_{n+\sigma(i)-1}$ being maximal, it is in particular larger than the one of any vector of $\mathcal{G}_{\mathcal{S}_n,i}^r$ and thus than $\frac{1}{2} \times \frac{L_{\min}}{L_{\max}} \times \text{HV}_{r_i^n}(\text{PF}_f)$ by Proposition 3.10. In other words, the hypervolume improvement at any iteration $n + \sigma(i)$ is larger than $\frac{1}{2} \times \frac{L_{\min}}{L_{\max}} \times \text{HV}_{r_i^n}(\text{PF}_f)$. By adding these inequations for all $i \in \llbracket 1, n + 1 \rrbracket$, we deduce that the hypervolume improvement from iteration n to $2n + 1$ is larger than $\frac{1}{2} \times \frac{L_{\min}}{L_{\max}} \times \sum_{i=1}^{n+1} \text{HV}_{r_i^n}(\text{PF}_f)$. Since the sum of the $\text{HV}_{r_i^n}(\text{PF}_f)$ is the optimality gap at iteration n , we have (4.2). If f is convex instead of bilipschitz, we use Proposition 3.9 instead of Proposition 3.10 and obtain (4.3). \square

Since the optimality gaps form a decreasing sequence, such lower bounds on the relation between the optimality gaps at iteration $2n+1$ and at iteration n imply that the optimality gap associated to a greedy set sequence converges asymptotically to 0. Equivalently, HV-ISOOMOO algorithms coupled with perfect singleobjective optimization converge to the entire Pareto front as stated formally below.

THEOREM 4.2. *Consider a biobjective optimization problem with a Pareto front described by a bilipschitz or convex function f .*

The hypervolume of a greedy set sequence relative to a valid reference point r converges to the hypervolume of the entire Pareto front, i.e. $\text{HV}_r(\mathcal{S}_n) \xrightarrow{n \rightarrow \infty} \text{HV}_r(\text{PF}_f)$.

Equivalently, for such Pareto fronts and under Assumption 2.3 of perfect singleobjective optimization, HV-ISOOMOO algorithms relative to a valid reference point r converge to the Pareto front in the sense of Definition 2.2.

From the lower bounds on the relation between the optimality gaps at iteration $2n + 1$ and at iteration n , we deduce the following upper bounds on the normalized optimality gap at any iteration.

COROLLARY 4.3. *Consider a biobjective optimization problem with a Pareto front described by a (L_{\min}, L_{\max}) -bilipschitz function. A greedy set sequence $(\mathcal{S}_n)_{n \in \mathbb{N}^*}$ relative to a valid reference point r satisfies for all n*

$$(4.4) \quad \frac{\text{HV}_r(\text{PF}_f) - \text{HV}_r(\mathcal{S}_n)}{\text{HV}_r(\text{PF}_f)} \leq \left(1 - \frac{1}{2} \times \frac{L_{\min}}{L_{\max}}\right)^{\lfloor \log_2(n+1) \rfloor} \leq (2n+2)^{\log_2(1 - \frac{1}{2} \times \frac{L_{\min}}{L_{\max}})}.$$

If the function f is convex, then any greedy set sequence relative to a valid reference point r satisfies for all n

$$(4.5) \quad \frac{\text{HV}_r(\text{PF}_f) - \text{HV}_r(\mathcal{S}_n)}{\text{HV}_r(\text{PF}_f)} \leq \left(\frac{1}{2}\right)^{\lfloor \log_2(n+1) \rfloor} \leq \frac{1}{2n+2}.$$

Hence, for such reference points and under Assumption 2.3 of perfect singleobjective optimization, HV-ISOOMOO algorithms relative to r satisfy (4.4) if f is (L_{\min}, L_{\max}) -bilipschitz and (4.5) if f is convex where \mathcal{S}_n is replaced by \mathcal{I}_n , the final incumbents Pareto front approximation at iteration n .

Proof. The k -th term of the sequence defined by $u_0 = 1$ and $u_{n+1} = 2 \times u_n + 1$ for all $n \geq 1$ is $2^k - 1$. Thus, (4.2) and (4.3) imply that when f is (L_{\min}, L_{\max}) -bilipschitz or convex, the normalized optimality gap at iteration $2^k - 1$ is inferior to $(1 - C)^k$ with C

equal to $\frac{1}{2} \times \frac{L_{\min}}{L_{\max}}$ and $\frac{1}{2}$, respectively. Since the hypervolume of the greedy set increases with n , and thus the optimality gap decreases with n , we deduce the first inequalities in (4.4) and (4.5) via the change of variable $k = \lfloor \log_2(n+1) \rfloor$.

Additionally, for every n , $\lfloor \log_2(n+1) \rfloor$ is smaller than $\log_2(n+1)+1$, that is $\log_2(2n+2)$. For every C , $\log_2(2n+2)$ equals $\log_C(2n+2) \times \log_2(C)$, and thus $C^{\log_2(2n+2)}$ equals $(2n+2)^{\log_2(C)}$. Therefore, we can infer that $(2n+2)^{\log_2(C)}$ is an upper bound of the normalized optimality gap with $C = 1 - \frac{1}{2} \times \frac{L_{\min}}{L_{\max}}$ and $C = \frac{1}{2}$ when f is (L_{\min}, L_{\max}) -bilipschitz and convex, respectively. \square

We focus here on the relation between the optimality gap at iteration n and at iteration $2n+1$. We could similarly examine the relation between the optimality gap at iteration n and at any later iteration. For example, we could prove that if f is (L_{\min}, L_{\max}) -bilipschitz, then for all n , for all $k \leq n+1$, $\frac{\text{HV}_r(\text{PF}_f) - \text{HV}_r(\mathcal{S}_{n+k})}{\text{HV}_r(\text{PF}_f) - \text{HV}_r(\mathcal{S}_n)}$ is smaller than $1 - \frac{1}{2} \times \frac{L_{\min}}{L_{\max}} \times \frac{k}{n+1}$.

Consider the k gap regions of \mathcal{S}_n with the largest areas. The hypervolume improvement from iteration n to $n+k$ is at least $\frac{1}{2} \times \frac{L_{\min}}{L_{\max}}$ times the area of the union of these gap regions, which is at least $\frac{k}{n+1}$ times the optimality gap at iteration n .

4.2. Asymptotical behavior of the convergence of $\text{HV}_r(\mathcal{S}_n)$ to $\text{HV}_r(\text{PF}_f)$.

In this section, we analyze the asymptotic convergence behavior for a Pareto front described by a bilipschitz function with a Hölder continuous derivative. We prove that, in this case, doubling the number of vectors in the greedy set divides the optimality gap by a factor which converges asymptotically to two as stated in Theorem 4.10. This asymptotic limit corresponds to the case of affine Pareto fronts with a reference point dominating the nadir point. For such Pareto fronts and reference points, the optimality gap is always halved when the number of vectors in the greedy set goes from n to $2n+1$, see Figure 5.

First, we study the properties of the part of the Pareto front corresponding to a specific gap region of \mathcal{S}_n . For all n , let note σ_n the permutation of $\llbracket 1, n \rrbracket$ which orders the vectors of \mathcal{S}_n by increasing F_1 -values and the so-called *ordered greedy set F_1 -values*:

$$(4.6) \quad w_{i,r}^n := v_{\sigma_n(i),1} \text{ for } i \in \llbracket 1, n \rrbracket, w_{0,r}^n := \tilde{x}_{\min,r} \text{ and } w_{n+1,r}^n := \tilde{x}_{\max,r}.$$

Naturally, we have $w_{0,r}^n \leq w_{1,r}^n \leq \dots \leq w_{n+1,r}^n$, and the intervals $[w_{i-1,r}^n, w_{i,r}^n[$ for $i \in \llbracket 1, n+1 \rrbracket$ form a partition of $[\tilde{x}_{\min,r}, \tilde{x}_{\max,r}[$, see the righthand plot of Figure 4. The interval $[w_{i-1,r}^n, w_{i,r}^n[$ corresponds to the part of the Pareto front belonging to the i -th gap region of \mathcal{S}_n . When the Pareto front is bilipschitz, the lengths of these intervals converge asymptotically to 0 as stated in the next lemma. It is a direct consequence of the convergence of $\text{HV}_r(\mathcal{S}_n)$ to $\text{HV}_r(\text{PF}_f)$ stated in Theorem 4.2.

LEMMA 4.4. *If the Pareto front is described by a bilipschitz function f and the greedy set sequence is associated to a valid reference point r , then the ordered greedy set F_1 -values satisfy $\max_{i \in \llbracket 1, n+1 \rrbracket} w_{i,r}^n - w_{i-1,r}^n \xrightarrow{n \rightarrow \infty} 0$ with the $w_{i,r}^n$ defined in (4.6).*

Proof. Let L_{\min} and L_{\max} be constants such that f is (L_{\min}, L_{\max}) -bilipschitz. The area of the i -th gap region of \mathcal{S}_n is $\int_{w_{i-1,r}^n}^{w_{i,r}^n} (f(x) - f(w_{i-1,r}^n))dx$. This is larger than $\int_{w_{i-1,r}^n}^{w_{i,r}^n} L_{\min} \times (w_{i,r}^n - x)dx$, which equals $\frac{1}{2} \times L_{\min} \times (w_{i,r}^n - w_{i-1,r}^n)^2$. Since the area of any gap region of \mathcal{S}_n is inferior to the optimality gap at iteration n , this implies that the difference $w_{i,r}^n - w_{i-1,r}^n$ is inferior to $\sqrt{2 \times (\text{HV}_r(\text{PF}_f) - \text{HV}_r(\mathcal{S}_n))}$ for all n , for all

620 $i \in \llbracket 1, n+1 \rrbracket$. Therefore, the convergence of $\text{HV}_r(\mathcal{S}_n)$ to $\text{HV}_r(\text{PF}_f)$ stated in [Theorem 4.2](#)
 621 implies that the maximum over i of $w_{i,r}^n - w_{i-1,r}^n$ converges to 0. \square

622 We prove in the next lemma that if the Pareto front is described by a bilipschitz
 623 function f with a Hölder continuous derivative, then the the part of the Pareto front
 624 belonging to a specific gap region of \mathcal{S}_n is bilipschitz for some constants whose ratio con-
 625 verges asymptotically to 1. Affine functions being the only functions to be (L_{\min}, L_{\max}) -
 626 bilipschitz with $L_{\min}/L_{\max} = 1$, it supports the interpretation that the convergence of
 627 a greedy set sequence for such Pareto fronts and for affine Pareto fronts share some
 628 asymptotic similarities.

629 When f is bilipschitz, its restriction to the part of the Pareto front dominating r_i^n ,
 630 that is $[w_{i-1,r}^n, w_{i,r}^n]$, is $(L_{\min}^{i,n}, L_{\max}^{i,n})$ -bilipschitz with

$$\begin{aligned} L_{\min}^{i,n} &:= \inf \left\{ \left| \frac{f(x) - f(y)}{x - y} \right|, x, y \in [w_{i-1,r}^n, w_{i,r}^n], x \neq y \right\} \text{ and} \\ L_{\max}^{i,n} &:= \sup \left\{ \left| \frac{f(x) - f(y)}{x - y} \right|, x, y \in [w_{i-1,r}^n, w_{i,r}^n], x \neq y \right\}. \end{aligned} \quad (4.7)$$

632 At iteration n , the ratio between $L_{\max}^{i,n}$ and $L_{\min}^{i,n}$, the bilipschitz constants on the i -th gap
 633 region of \mathcal{S}_n , is by definition smaller than

$$q_n := \max \left\{ \frac{L_{\max}^{i,n}}{L_{\min}^{i,n}}, i \in \llbracket 1, n+1 \rrbracket : [w_{i-1,r}^n, w_{i,r}^n] \neq \emptyset \right\}. \quad (4.8)$$

636 The proof of the convergence of q_n to 1 relies on the fact that a derivable function can
 637 be approximated locally by an affine function. The quality of this approximation is
 638 guaranteed by the Hölder continuity of the derivative.

639 **LEMMA 4.5.** *We consider a greedy set sequence $(\mathcal{S}_n)_{n \in \mathbb{N}^*}$ relative to a valid reference*
 640 *point r . If the Pareto front is described by a bilipschitz function with a Hölder continuous*
 641 *derivative, then q_n defined in (4.8) converges asymptotically to 1.*

642 *Proof.* We take α such that f' is Hölder continuous with exponent α , i.e f is $\mathcal{C}^{1,\alpha}$,
 643 and $L_{\min}, L_{\max} > 0$ such that the function f describing the Pareto front is (L_{\min}, L_{\max}) -
 644 bilipschitz. We recall that f is decreasing, and thus for all $x < y$, we have $f(x) - f(y) \geq 0$.
 645 Since f is $\mathcal{C}^{1,\alpha}$ and therefore \mathcal{C}^1 , the Taylor formula with Lagrange remainder states that
 646 for all $x < y$, there exists $\xi \in [x, y]$ such that $f(y) = f(x) + (y - x) \times f'(\xi)$. Since f is $\mathcal{C}^{1,\alpha}$,
 647 this implies that for all $x < y$, $|f(y) - f(x) - (y - x) \times f'(x)| \leq (y - x)^{1+\alpha} \times [f']_{\mathcal{C}^\alpha}$. Thus,
 648 $\frac{f(y) - f(x)}{x - y}$ is smaller than $-f'(x) + [f']_{\mathcal{C}^\alpha} \times (y - x)^\alpha$. We now restrict ourselves to x and
 649 y belonging to the non-empty interval $[w_{i-1,r}^n, w_{i,r}^n]$. Our goal is to find an upper bound
 650 depending on i but not on either x or y . Since f is $\mathcal{C}^{1,\alpha}$, the difference between $-f'(x)$
 651 and $-f'(w_{i-1,r}^n)$ is smaller than $[f']_{\mathcal{C}^\alpha} \times (x - w_{i-1,r}^n)^\alpha$, and thus $[f']_{\mathcal{C}^\alpha} \times (w_{i,r}^n - w_{i-1,r}^n)^\alpha$
 652 . Additionally, the difference between x and y is smaller than $w_{i,r}^n - w_{i-1,r}^n$. We conclude
 653 that for $x, y \in [w_{i-1,r}^n, w_{i,r}^n]$, $\frac{f(y) - f(x)}{x - y}$ is smaller than $-f'(w_{i-1,r}^n) + 2[f']_{\mathcal{C}^\alpha} \times (w_{i,r}^n - w_{i-1,r}^n)^\alpha$,
 654 and thus so is $L_{\max}^{i,n}$ defined in (4.7).

655 Following the same approach, we can also infer that $L_{\max}^{i,n}$ defined in (4.7) is greater
 656 than the symmetric quantity $-f'(w_{i-1,r}^n) - 2[f']_{\mathcal{C}^\alpha} \times (w_{i,r}^n - w_{i-1,r}^n)^\alpha$. The quantity
 657 $-f'(w_{i-1,r}^n)$ is greater than L_{\min} and $(w_{i,r}^n - w_{i-1,r}^n)^\alpha$ is smaller than $\max_{i \in \llbracket 1, n+1 \rrbracket} (w_{i,r}^n -$

658 $w_{i-1,r}^n)^\alpha$. As a consequence, q_n is smaller than $\frac{L_{\min}+2[f']_{C^\alpha} \times \max_{i \in \llbracket 1, n+1 \rrbracket} (w_{i,r}^n - w_{i-1,r}^n)^\alpha}{L_{\min}-2[f']_{C^\alpha} \times \max_{i \in \llbracket 1, n+1 \rrbracket} (w_{i,r}^n - w_{i-1,r}^n)^\alpha}$. By
 659 Lemma 4.4, $\max_{i \in \llbracket 1, n+1 \rrbracket} w_{i,r}^n - w_{i-1,r}^n$ converges to 0 and thus, this upper bound on q_n
 660 converges to 1. Since q_n is always larger than 1, it converges to 1. \square

661 A consequence of the previous lemma is that the bounds on the hypervolume improvement
 662 of v_{n+1} to \mathcal{S}_n normalized by the area of the gap region filled by v_{n+1} that we can infer
 663 from Propositions 3.10 and 3.11 converge asymptotically to 1/2, see (4.9). Similarly,
 664 the bounds on the normalized area of the child of a gap region that we can infer from
 665 Lemma A.2 converge to 1/4, see (4.10). These asymptotic values correspond to the case
 666 of an affine Pareto front with a reference point dominating the nadir point, see Figure 5.




Fig. 5: The three greedy sets \mathcal{S}_1 (left), \mathcal{S}_2 (middle) and \mathcal{S}_3 (right) and their gap regions for an affine Pareto front with a reference point r dominating the nadir point. The area of any of the gap regions of \mathcal{S}_1 are half of $\text{HV}_r(\text{PF}_f)$ (left). The area of any of the new gap regions of \mathcal{S}_2 is a quarter of the area of their parents (middle). The optimality gap of \mathcal{S}_3 (right) is half of the optimality gap of \mathcal{S}_1 .

667

668 LEMMA 4.6. We consider a greedy set sequence $(\mathcal{S}_n)_{n \in \mathbb{N}^*}$ relative to a valid reference
 669 point r . If the Pareto front is described by a bilipschitz function f with a Hölder continuous
 670 derivative, then for all $\epsilon > 0$, for n large enough, for every non-empty gap region $\mathcal{G}_{\mathcal{S}_n,i}^r$
 671 and every child $\mathcal{G}_{\mathcal{S}_n,j}^r$ of $\mathcal{G}_{\mathcal{S}_n,i}^r$, we have

$$672 \quad (4.9) \quad \frac{1}{2} \times (1 - \epsilon) \leq \frac{\max_{u \in \mathcal{G}_{\mathcal{S}_n,i}^r} \text{HVI}_r(u, \mathcal{S}_n)}{\lambda(\mathcal{G}_{\mathcal{S}_n,i}^r)} \leq \frac{1}{2} \times (1 + \epsilon) \text{ and}$$

$$673 \quad (4.10) \quad \frac{1}{4 \times (1 + \epsilon)} \leq \frac{\lambda(\mathcal{G}_{\mathcal{S}_n,j}^r)}{\lambda(\mathcal{G}_{\mathcal{S}_n,i}^r)} \leq \frac{1}{4 \times (1 - \epsilon)} .$$

674

675 *Proof.* The interval $[w_{i-1,r}^n, w_{i,r}^n]$ is the set of the first coordinates of the vectors of
 676 the Pareto front which dominate r_i^n . The restriction to $[w_{i-1,r}^n, w_{i,r}^n]$ of f is (L_{\min}, L_{\max}) -
 677 bilipschitz for some L_{\min} and L_{\max} such that $\frac{L_{\max}}{L_{\min}} = q_n$ with q_n defined in (4.8). Ad-
 678 ditionally, as stated in Proposition B.2, for n large enough, all the r_i^n corresponding to
 679 non-empty gap regions dominate the nadir point. As a consequence, the conditions to
 680 apply Lemma A.2 and Proposition 3.11 are met by non-extremes gap regions.

681 By Propositions 3.10 and 3.11, $\frac{\max_{u \in \text{PF}_f} \text{HVI}_{r_i^n}(u)}{\text{HV}_{r_i^n}(\text{PF}_f)}$ is between $\frac{1}{2} \times \frac{1}{q_n}$ and $\frac{1}{2} \times q_n$. Ad-
 682 ditionally, by Lemma A.2, $\frac{\lambda(\mathcal{G}_{\mathcal{S}_n,j}^r)}{\text{HV}_{r_i^n}(\text{PF}_f)}$ is between $\frac{1 - \frac{1}{2} \times q_n}{1 + q_n^2}$ and $\frac{1 - \frac{1}{2} \times \frac{1}{q_n}}{1 + \frac{1}{q_n^2}}$. The maximum

over the vectors u belonging to the Pareto front of $\text{HV}_{r_i^n}(u)$ is equal to the maximum over u belonging to the i -th gap region of \mathcal{S}_n of $\text{HVI}_r(u, \mathcal{S}_n)$. Indeed, $\text{HV}_{r_i^n}(\cdot)$ is null for vectors outside the i -th gap region of \mathcal{S}_n while it is nonnegative, equal to $\text{HVI}_r(\cdot, \mathcal{S}_n)$, otherwise. Additionally, $\text{HV}_{r_i^n}(\text{PF}_f)$ equals $\lambda(\mathcal{G}_{\mathcal{S}_n, i}^r)$. The convergence of q_n to 1 stated in Lemma 4.5 imply that the bounds proven so far converge to a half and a quarter, respectively. Thus, we have (4.9) and (4.10) for n large enough. \square

The following lemma states that for n large enough, the area of two non-empty gap regions relative to the same greedy set cannot be too different. More precisely, the area of any gap region of \mathcal{S}_n cannot be more than $4 \times (1 + o(\epsilon))$ times greater than the area of another gap region of \mathcal{S}_n . The proof relies on considering the parents of the gap regions.

LEMMA 4.7. *We consider a greedy set sequence $(\mathcal{S}_n)_{n \in \mathbb{N}^*}$ relative to a valid reference point r . If the Pareto front is described by a bilipschitz function with a Hölder continuous derivative, then for all $\epsilon > 0$, for n large enough and for any non-empty gap regions of \mathcal{S}_n , $\mathcal{G}_{\mathcal{S}_n, i}^r$ and $\mathcal{G}_{\mathcal{S}_n, j}^r$ with $i, j \in \llbracket 1, n+1 \rrbracket$, we have*

$$(4.11) \quad \frac{\lambda(\mathcal{G}_{\mathcal{S}_n, i}^r)}{\lambda(\mathcal{G}_{\mathcal{S}_n, j}^r)} \leq 4 \times \frac{(1 + \epsilon)^2}{1 - \epsilon}.$$

Proof. Fix $\epsilon > 0$. By Lemma 4.6, there exists $N_1 \in \mathbb{N}^*$ such that for all n greater than N_1 , (4.9) and (4.10) are verified for any non-empty gap region of \mathcal{S}_n and its children. Since $\max_{i \in \llbracket 1, n+1 \rrbracket} w_{i, r}^n - w_{i-1, r}^n$ converges to 0 by Lemma 4.4, every non-empty gap region is filled at some point. Take N_2 such that all the non-empty gap regions of \mathcal{S}_{N_1} are filled at iteration N_2 . For all n greater than N_2 , (4.9) and (4.10) are true for any non-empty gap region of \mathcal{S}_n and its children, but also for its parents.

Take $n \geq N_2$. We note $\mathcal{G}_1 := \mathcal{G}_{\mathcal{S}_n, i}^r$ and $\mathcal{G}_2 := \mathcal{G}_{\mathcal{S}_n, j}^r$ two distinct non-empty gap regions of \mathcal{S}_n , and \mathcal{P}_1 and \mathcal{P}_2 their respective parents. When two sets correspond to gap regions relative to the same greedy set \mathcal{S}_m , we say that they cohabit at iteration m . Since only one vector is added to \mathcal{S}_n at a time, the cohabitation of \mathcal{G}_1 and \mathcal{G}_2 implies that either \mathcal{G}_1 and \mathcal{P}_2 or \mathcal{G}_2 and \mathcal{P}_1 cohabit at some earlier iteration. In the first case, there necessarily exists $m \geq N_2$ such that \mathcal{P}_2 and \mathcal{G}_1 are gap regions relative to \mathcal{S}_m and v_{m+1} belongs to \mathcal{P}_2 , otherwise, \mathcal{G}_1 and \mathcal{G}_2 would not cohabit. By (4.9), the maximum hypervolume improvement to \mathcal{S}_m of a vector of \mathcal{G}_1 and of a vector of \mathcal{P}_2 are at least $\frac{1}{2} \times (1 - \epsilon) \times \lambda(\mathcal{G}_1)$ and at most $\frac{1}{2} \times (1 + \epsilon) \times \lambda(\mathcal{P}_2)$, respectively. Since a vector of \mathcal{P}_2 , v_{m+1} , maximizes the hypervolume improvement to \mathcal{S}_m , we have $\lambda(\mathcal{G}_1) \times \frac{1}{2} \times (1 - \epsilon) \leq \lambda(\mathcal{P}_2) \times \frac{1}{2} \times (1 + \epsilon)$. Since $\lambda(\mathcal{P}_2)$ is smaller than $4 \times (1 + \epsilon)$ times the area of its child $\lambda(\mathcal{G}_2)$ by (4.10), this inequality implies (4.11). In the second case, \mathcal{P}_2 is filled before \mathcal{P}_1 . Thus, there exists $m \geq N_2$ such that \mathcal{P}_1 and \mathcal{P}_2 cohabit at iteration m and v_{m+1} belongs to \mathcal{P}_2 . Since the area of \mathcal{P}_1 is larger than the one of its child \mathcal{G}_1 , the hypervolume improvement of v_{m+1} to \mathcal{S}_m is still larger than $\frac{1}{2} \times (1 - \epsilon) \times \lambda(\mathcal{G}_1)$. The rest of the argumentation remains valid. \square

We now have all the results needed to analyze the asymptotic impact of doubling the number of points in the greedy set. To prove the following asymptotic upper bound, we rely on similar arguments as for its nonasymptotic counterpart, Proposition 4.1. The previous lemma guarantees that the impact of doubling the number of points in the greedy set is asymptotically similar to the impact of passing from n points to $2n + 1$.

PROPOSITION 4.8. *Let $(\mathcal{S}_n)_{n \in \mathbb{N}^*}$ be a greedy set sequence relative to valid reference point r . If the Pareto front is described by a bilipschitz function f with a Hölder continuous*

727 derivative, then for all $\epsilon > 0$, we have for n large enough

$$728 \quad (4.12) \quad \frac{HV_r(PF_f) - HV_r(\mathcal{S}_{2n})}{HV_r(PF_f) - HV_r(\mathcal{S}_n)} \leq \frac{1}{2} + o(\epsilon) .$$

730 *Proof.* Fix $\epsilon > 0$. Fix n large enough to verify (4.9) and (4.11) for this particular ϵ .

731 Let σ be a permutation of $\llbracket 1, n+1 \rrbracket$ such that the i -th gap region of \mathcal{S}_n is filled by
 732 $v_{n+\sigma(i)}$ when it is filled before iteration $2n+1$. With this choice of permutation, $\mathcal{G}_{\mathcal{S}_n, i}^r$
 733 is always a gap region of $\mathcal{S}_{n+\sigma(i)-1}$. Thus, $HVI_r(v_{n+\sigma(i)}, \mathcal{S}_{n+\sigma(i)-1})$ is superior to the
 734 maximum hypervolume improvement of a vector of $\mathcal{G}_{\mathcal{S}_n, i}^r$ to $\mathcal{S}_{n+\sigma(i)-1}$, which is superior
 735 to $\frac{1}{2} \times (1-\epsilon) \times \lambda(\mathcal{G}_{\mathcal{S}_n, i}^r)$ by (4.9). It is equivalent to say that the hypervolume improvement
 736 at iteration $n+\sigma(i)$ is larger than $\frac{1}{2} \times (1-\epsilon) \times \lambda(\mathcal{G}_{\mathcal{S}_n, i}^r)$. Summing over $i \in \llbracket 1, n+1 \rrbracket$, we
 737 obtain that the hypervolume improvement between iteration n and $2n+1$ is larger than
 738 the sum over i of $\frac{1}{2} \times (1-\epsilon) \times \lambda(\mathcal{G}_{\mathcal{S}_n, i}^r)$, that is $\frac{1}{2} \times (1-\epsilon)$ times the optimality gap at
 739 iteration n .

740 Now, we need to bound the hypervolume improvement at iteration $2n+1$, that is
 741 $HVI_r(v_{2n+1}, \mathcal{S}_{2n})$. It is smaller than $\frac{1}{2} \times (1+\epsilon) \times \max_{i \in \llbracket 1, 2n+1 \rrbracket} \lambda(\mathcal{G}_{\mathcal{S}_{2n}, i}^r)$ by (3.3) and
 742 (4.9). Since the area of a gap region is smaller than the one of its parent, the maximum
 743 area of a gap region is lower at iteration $2n$ than at iteration n . The maximum area of
 744 one of the more than $n-1$ gap regions of \mathcal{S}_n is itself smaller than $\frac{1}{n-1} \times \frac{4 \times (1+\epsilon)^2}{1-\epsilon}$ times
 745 the optimality gap at iteration n by (4.11).

746 We conclude that the relation between the optimality gap at iteration $2n$ and at
 747 iteration n is smaller than $1 - \frac{1}{2} \times (1-\epsilon) + \frac{1-\epsilon}{2 \times (n-1)}$. \square

748 We roughly follow the same approach to obtain the following asymptotic lower bound on
 749 the impact of doubling the number of points in the greedy set. Lemmas 4.6 and 4.7 are
 750 key to prove an upper bound on the hypervolume improvement at iteration k . They allow
 751 to prove that filling a gap region of \mathcal{S}_n more than once gives, up to a factor $1+o(\epsilon)$, a
 752 lower hypervolume improvement than filling a gap region which was not filled. Indeed,
 753 the area of a descendant of a gap region of \mathcal{S}_n is at most $\frac{1}{4} + o(\epsilon)$ times the area of its
 754 parent by Lemma 4.6, which is itself at most $4 + o(\epsilon)$ times the area of any other gap
 755 region of \mathcal{S}_n by Lemma 4.7.

756 PROPOSITION 4.9. Let $(\mathcal{S}_n)_{n \in \mathbb{N}^*}$ be a greedy set sequence relative to a valid reference
 757 point r . If the Pareto front is described by a bilipschitz function f with a Hölder continuous
 758 derivative, then for all $\epsilon > 0$, we have for n large enough

$$759 \quad (4.13) \quad \frac{HV_r(PF_f) - HV_r(\mathcal{S}_{2n})}{HV_r(PF_f) - HV_r(\mathcal{S}_n)} \geq \frac{1}{2} + o(\epsilon) .$$

761 *Proof.* Fix $\epsilon > 0$. Fix n large enough to verify (4.9), (4.10) and (4.11) for this
 762 particular ϵ . Let $\delta \in \{-1, 0, 1\}$ be such that \mathcal{S}_n has $n+\delta$ non-empty gap regions. Let
 763 $i_0 := 1$ when the left extreme gap region is empty and $i_0 := 0$ otherwise.

764 Let σ be a permutation of $\llbracket 1, n+\delta \rrbracket$ such that the i -th non-empty gap region of
 765 \mathcal{S}_n , $\mathcal{G}_{\mathcal{S}_n, i_0+i}^r$, is filled by the vector $v_{n+\sigma(i)}$ when it is filled before iteration $2n+\delta$. We
 766 distinguish two cases. In the first case, $v_{n+\sigma(i)}$ is the child of the i -th non-empty gap
 767 region of \mathcal{S}_n , and consequently its hypervolume improvement to $\mathcal{S}_{n+\sigma(i)-1}$ is at most
 768 $\frac{1}{2} \times (1+\epsilon) \times \lambda(\mathcal{G}_{\mathcal{S}_n, i_0+i}^r)$ by (4.9). In the second case, $v_{n+\sigma(i)}$ belongs to $\mathcal{G}_{\mathcal{S}_n, i_0+j}^r$, the
 769 j -th non-empty gap region of \mathcal{S}_n , with $j \neq i$ and, by definition of σ , fills a descendant of

770 this gap region not $\mathcal{G}_{\mathcal{S}_n, i_0+j}^r$ itself. By (4.9), the hypervolume improvement of $v_{n+\sigma(i)}$ to
 771 $\mathcal{S}_{n+\sigma(i)-1}$ is still at most $\frac{1}{2} \times (1+\epsilon)$ times the area of the gap region it fills. By (4.10),
 772 the area of a descendant of $\mathcal{G}_{\mathcal{S}_n, i_0+j}^r$ is smaller than $\frac{1}{4 \times (1-\epsilon)}$ times the area of its ancestor.
 773 By (4.11), we also know that the area of the i -th non-empty gap region of \mathcal{S}_n is at most
 774 $4 \times \frac{(1+\epsilon)^2}{1-\epsilon}$ times the area of any other gap region of \mathcal{S}_n , in particular its i -th non-empty
 775 gap region. We conclude that the hypervolume improvement of $v_{n+\sigma(i)}$ to $\mathcal{S}_{n+\sigma(i)-1}$ is
 776 smaller than $\frac{1}{2} \times \frac{(1+\epsilon)^3}{(1-\epsilon)^2} \times \lambda(\mathcal{G}_{\mathcal{S}_n, i_0+i}^r)$. To summarize, since $1+\epsilon$ is smaller than $\frac{(1+\epsilon)^3}{(1-\epsilon)^2}$, the
 777 hypervolume improvement at any iteration $n+\sigma(i)$ is smaller than $\frac{1}{2} \times \frac{(1+\epsilon)^3}{(1-\epsilon)^2} \times \lambda(\mathcal{G}_{\mathcal{S}_n, i}^r)$.
 778 Summing over $i \in \llbracket 1, n+\delta \rrbracket$, the hypervolume improvement from iteration n to $2n+\delta$
 779 is smaller than $\frac{1}{2} \times \frac{(1+\epsilon)^3}{(1-\epsilon)^2}$ times the sum over i of $\lambda(\mathcal{G}_{\mathcal{S}_n, i}^r)$, that is the optimality gap at
 780 iteration n .

781 Now, it is left to prove an upper bound on $\text{HV}_r(\mathcal{S}_{2n}) - \text{HV}_r(\mathcal{S}_{2n+\delta})$. This quantity
 782 is maximal for $\delta = -1$, where it is simply the hypervolume improvement at iteration $2n$.
 783 As in the previous proof, it is smaller than $\frac{1+\epsilon}{2 \times (n-1)}$ times the optimality gap at iteration
 784 n . Therefore, the relation between the optimality gap at iteration $2n$ and at iteration n
 785 is larger than $1 - \frac{1}{2} \times \frac{(1+\epsilon)^3}{(1-\epsilon)^2} - \frac{1+\epsilon}{2 \times (n-1)}$. \square

786 We combine the lower and upper asymptotic bounds to obtain the following theorem.

787 **THEOREM 4.10.** *Consider a biobjective optimization problem and a greedy set se-*
 788 *quence $(\mathcal{S}_n)_{n \in \mathbb{N}^*}$ relative to a valid reference point r . If the Pareto front is described by a*
 789 *bilipschitz function f with a Hölder continuous derivative, we have*

$$790 \quad (4.14) \quad \frac{\text{HV}_r(\text{PF}_f) - \text{HV}_r(\mathcal{S}_{2n})}{\text{HV}_r(\text{PF}_f) - \text{HV}_r(\mathcal{S}_n)} \xrightarrow{n \rightarrow \infty} \frac{1}{2} .$$

792 *Consequently, for such Pareto front and reference point and under [Assumption 2.3](#) of*
 793 *perfect singleobjective optimization, HV-ISOOMOO algorithms relative to r satisfy (4.14)*
 794 *where \mathcal{S}_n is replaced by \mathcal{I}_n , the final incumbents Pareto front approximation at iteration*
 795 *n .*

796 **5. Conclusion.** We prove that HV-ISOOMOO algorithms coupled with a singleob-
 797 jective optimizer converge in $O(1/n)$ on convex Pareto fronts and in $O(1/n^c)$ on bilipschitz
 798 Pareto fronts with $c \leq 1$ depending on the bilipschitz constants where n is the number
 799 of meta-iterations. Each meta-iteration corresponds to a singleobjective optimization
 800 run. Both bounds are tight over the class of Pareto fronts and reference points consid-
 801 ered. They are reached for affine Pareto fronts and reference points dominating the nadir
 802 point. On convex Pareto fronts, the convergence is exactly in $\Theta(1/p)$, the fastest con-
 803 vergence achievable by biobjective optimization algorithms [16]. It shows that greedily
 804 adding points maximizing the hypervolume contribution as in HV-ISOOMOO algorithms
 805 is an effective way to quickly increase the hypervolume. Additionally, we prove that for
 806 bilipschitz Pareto fronts with a Hölder continuous derivative, doubling the number of
 807 meta-iterations divides the optimality gap by a factor which converges asymptotically to
 808 two. This asymptotic behavior resembles what we would observe with an affine Pareto
 809 front and a reference point dominating the nadir point. Yet, it does not guarantee con-
 810 vergence in $\Theta(1/n)$. Both $\left(\frac{\log(n)}{n}\right)_{n \in \mathbb{N}^*}$ and $\left(\frac{1}{n \times \log(n)}\right)_{n \in \mathbb{N}^*}$ are examples of sequences
 811 verifying this property without converging in $\Theta(1/n)$. The convergence on nonconvex

Pareto fronts could theoretically be slower than in $\Theta(1/n)$, but not faster [16].

Appendix A. Normalized areas of the gap regions relative to an hypervolume maximizer. The goal of this section is to prove bounds on the normalized areas of the gap regions $\mathcal{G}_{\text{left}}^{u^*}$ and $\mathcal{G}_{\text{right}}^{u^*}$ relative to an hypervolume maximizer u^* (see the lefthand plot of Figure 6) in the case of a bilipschitz Pareto front and of a reference point r dominating the nadir point. These bounds are stated in Lemma A.2. The proof relies on the bounds on the normalized maximum hypervolume proven in Subsection 3.3 and the following lower and upper bounds on the relation between $\lambda(\mathcal{G}_{\text{left}}^{u^*})$ and $\lambda(\mathcal{G}_{\text{right}}^{u^*})$.




Fig. 6: Illustration of elements of the proofs of Proposition A.1 in the case $r_1 \leq x_{\max}$ (left and middle) and of Lemma B.1 (right). The Pareto front is described by $f(x) = \frac{e}{e-1} \times e^{-x} + 1 - \frac{e}{e-1}$ for $x \in [0, 1]$. Left: the gap regions $\mathcal{G}_{\text{left}}^{u^*}$ and $\mathcal{G}_{\text{right}}^{u^*}$ with a segment of slope $-L_{\min}$ passing through u^* . Middle: the hypervolume improvements $\text{HVI}_r(u^*, u_\epsilon)$ (green) and $\text{HVI}_r(u_\epsilon, u^*)$ (purple). Right: the hypervolume improvement $\text{HVI}_r(u_{\min}, u)$ (green) and its counterpart $\text{HVI}_r(u, u_{\min})$ (purple) where u is a vector of the Pareto front which dominates r .

PROPOSITION A.1. We assume that the Pareto front is described by a (L_{\min}, L_{\max}) -bilipschitz function f . Let u^* be a non-extreme vector of the Pareto front which maximizes the hypervolume with respect to a valid reference point r . If $r_1 \leq x_{\max}$, we have $\lambda(\mathcal{G}_{\text{right}}^{u^*}) \geq \frac{L_{\min}^2}{L_{\max}^2} \times \lambda(\mathcal{G}_{\text{left}}^{u^*})$. If $r_2 \leq f(x_{\min})$, we have $\lambda(\mathcal{G}_{\text{left}}^{u^*}) \geq \frac{L_{\min}^2}{L_{\max}^2} \times \lambda(\mathcal{G}_{\text{right}}^{u^*})$.

Proof. We consider the case where $r_1 \leq x_{\max}$. Let x^* be the first coordinate of u^* . We denote $L_1 := r_1 - x^*$ and $L_2 := r_2 - f(x^*)$ the lengths of the sides of the rectangle $\mathcal{D}_{u^*}^r$. For all $x, y \in [x_{\min}, x_{\max}]$, we have $|f(x) - f(y)| \geq L_{\min} \times |x - y|$. Additionally, since $r_1 \leq x_{\max}$, the segment $[x^*, x^* + L_1]$ is included in $[x_{\min}, x_{\max}]$. As a consequence, the section of the Pareto front on the right of u^* dominates the segment between u^* and $u^* + L_1 \times (1, -L_{\min})$, see the lefthand plot of Figure 6. Therefore, $\lambda(\mathcal{G}_{\text{right}}^{u^*})$ is larger than the area of the region of the objective space dominated by this segment, not dominated by u^* and dominating r , that is $\frac{1}{2} \times L_{\min} \times L_1^2$. For all $x, y \in [x_{\min}, x_{\max}]$, we also have $|f(x) - f(y)| \leq L_{\max} \times |x - y|$. Therefore, the part of the Pareto front on the left of u^* is dominated by the segment between u^* and $u^* + L_2 \times (-\frac{1}{L_{\min}}, 1)$, and $\lambda(\mathcal{G}_{\text{left}}^{u^*})$ is smaller than $\frac{1}{2} \times \frac{1}{L_{\min}} \times L_2^2$. We have yet to prove a lower bound on $\frac{L_1}{L_2}$. The vector u^* being different from u_{\min} , for $\epsilon > 0$ small enough, the vector $u_\epsilon := (x^* - \epsilon, f(x^* - \epsilon))$ belongs to the Pareto front. As we can see in the middle plot of Figure 6, $\text{HVI}_r(u^*, u_\epsilon)$ is smaller

than $L_1 \times L_{\max} \times \epsilon$ and $\text{HVI}_r(u_\epsilon, u^*)$ is larger than $\epsilon \times (L_2 - \epsilon \times L_{\max})$. Additionally, u^* being an hypervolume maximizer, $\text{HVI}_r(u^*, u_\epsilon)$ is larger than $\text{HVI}_r(u_\epsilon, u^*)$, and thus $L_1 \times L_{\max} \geq L_2 - \epsilon \times L_{\max}$ for all $\epsilon > 0$. Taking the limit of this inequality when $\epsilon \rightarrow 0$, we obtain that $L_1 \times L_{\max} \geq L_2$. Combining the bounds on $\lambda(\mathcal{G}_{\text{left}}^{u^*})$ and $\lambda(\mathcal{G}_{\text{right}}^{u^*})$ with the lower-bound on $\frac{L_1}{L_2}$, we obtain the desired lower bound on $\lambda(\mathcal{G}_{\text{right}}^{u^*})$. We can obtain the symmetric inequality when $r_2 \geq f(x_{\min})$ by following the same approach. \square

In particular, when f is bilipschitz and r dominates the nadir point, both bounds hold. We now prove the desired bounds on the normalized area of the gap regions $\mathcal{G}_{\text{left}}^{u^*}$ and $\mathcal{G}_{\text{right}}^{u^*}$.

LEMMA A.2. *Let u^* be a vector which maximizes the hypervolume with respect to a valid reference point r . If the Pareto front is described by a (L_{\min}, L_{\max}) -bilipschitz function f and the reference point r dominates the nadir point, both $\lambda(\mathcal{G}_{\text{left}}^{u^*})$ and $\lambda(\mathcal{G}_{\text{right}}^{u^*})$ are between $(1 - \frac{1}{2} \times \frac{L_{\max}}{L_{\min}}) / (1 + \frac{L_{\max}^2}{L_{\min}^2})$ and $(1 - \frac{1}{2} \times \frac{L_{\min}}{L_{\max}}) / (1 + \frac{L_{\min}^2}{L_{\max}^2})$.*

Proof. Let note arbitrarily \mathcal{G}_1 and \mathcal{G}_2 the two gap regions of the set $S = \{u^*\}$. By Proposition A.1, $\lambda(\mathcal{G}_2)$ is between $\frac{L_{\min}^2}{L_{\max}^2} \times \lambda(\mathcal{G}_1)$ and $\frac{L_{\max}^2}{L_{\min}^2} \times \lambda(\mathcal{G}_1)$. Additionally, by Propositions 3.10 and 3.11, the normalized maximum hypervolume $\max_{u \in \text{PF}_f} \text{HV}_r(u)$ over $\text{HV}_r(\text{PF}_f)$ is between $\frac{1}{2} \times \frac{L_{\min}}{L_{\max}}$ and $\frac{1}{2} \times \frac{L_{\max}}{L_{\min}}$. These bounds can be transformed into bounds on $\text{HV}_r(\text{PF}_f) - \max_{u \in \text{PF}_f} \text{HV}_r(u)$, that is $\lambda(\mathcal{G}_1) + \lambda(\mathcal{G}_2)$. As a consequence, $\lambda(\mathcal{G}_1)$ is between $(1 - \frac{1}{2} \times \frac{L_{\max}}{L_{\min}}) \times \text{HV}_r(\text{PF}_f) - \frac{L_{\max}^2}{L_{\min}^2} \times \lambda(\mathcal{G}_1)$ and $(1 - \frac{1}{2} \times \frac{L_{\min}}{L_{\max}}) \times \text{HV}_r(\text{PF}_f) - \frac{L_{\min}^2}{L_{\max}^2} \times \lambda(\mathcal{G}_1)$. Moving all the $\lambda(\mathcal{G}_1)$ terms on the same side and re-normalizing this side, we obtain the desired bounds for \mathcal{G}_1 , which can be chosen to be either $\mathcal{G}_{\text{left}}^{u^*}$ or $\mathcal{G}_{\text{right}}^{u^*}$. \square

Appendix B. The nadir point is dominated by all the r_i^n corresponding to non-empty gap regions for n large. We show in this section that for bilipschitz Pareto fronts, the nadir point is dominated by all the r_i^n corresponding to non-empty gap regions, for n large enough. This result is stated in Proposition B.2 and used in Subsection 4.2. It is equivalent to prove that the extreme vectors which dominate the reference point belong to the greedy set for n large enough.

First, we prove in the next proposition that if $r_1 > x_{\max}$ (resp. $r_2 > f(x_{\min})$), then for r_2 (resp. r_1) close enough to $f(x_{\max})$ (resp. x_{\min}) the extreme vector u_{\max} (resp. u_{\min}) is the only hypervolume maximizer, see the righthand plot of Figure 6. There are similar statements in [9] for the set of μ points maximizing the hypervolume, but they only apply to $\mu \geq 2$.

LEMMA B.1. *We assume that the Pareto front is described by a function f which is (L_{\min}, L_{\max}) -bilipschitz and that the reference point r is valid. If $r_1 > x_{\max}$ and $f(x_{\max}) < r_2 < f(x_{\max}) + L_{\min} \times (r_1 - x_{\max})$, the right extreme of the Pareto front u_{\max} is the only maximizer of $\text{HV}_r(\cdot)$. Additionally, if $r_2 > f(x_{\min})$ and $x_{\min} < r_1 < x_{\min} + \frac{r_2 - f(x_{\min})}{L_{\max}}$, the vector $u_{\min} = (x_{\min}, f(x_{\min}))$ is the only maximizer of $\text{HV}_r(\cdot)$.*

Proof. This proof is illustrated in the righthand plot of Figure 6. Let r be a reference point such that $r_2 > f(x_{\min})$ and $x_{\min} < r_1 < x_{\min} + \frac{r_2 - f(x_{\min})}{L_{\max}}$. Let $u = (x, f(x)) \neq u_{\min}$ be a vector of the Pareto front which dominates r . The hypervolume improvement of u_{\min} to $\{u\}$ is $(r_2 - f(x_{\min})) \times (x - x_{\min})$. The hypervolume improvement of u to $\{u_{\min}\}$ is $(f(x_{\min}) - f(x)) \times (r_1 - x)$, which is smaller than $L_{\max} \times (x - x_{\min}) \times (r_1 - x_{\min})$ since u

dominates r and f is (L_{\min}, L_{\max}) -bilipschitz. Since we assume that $L_{\max} \times (r_1 - x_{\min}) < r_2 - f(x_{\min})$, the upper bound on $\text{HVI}_r(u, u_{\min})$ is strictly smaller than $\text{HVI}_r(u_{\min}, u)$. Therefore, the hypervolume of u_{\min} is strictly larger than the one of u . We conclude that u_{\min} is the unique hypervolume maximizer. The symmetric result can be obtained with the same approach. \square

It is left to prove that when $r_1 > x_{\max}$ (resp. $r_2 > f(x_{\min})$), the second coordinate of r_{n+1}^n (resp. the first coordinate of r_0^n) indeed converge to $f(x_{\max})$ (resp. x_{\min}). It is a straightforward consequence of Lemma 4.4. Therefore, we are able to conclude.

PROPOSITION B.2. *We assume that the Pareto front is described by a bilipschitz function f . Let $(\mathcal{S}_n)_{n \in \mathbb{N}^*}$ be a greedy set sequence relative to a valid reference point r . For n large enough, every reference point r_i^n corresponding to a non-empty gap region $\mathcal{G}_{\mathcal{S}_n, i}^r$ dominates the nadir point.*

Proof. By Lemma 4.4, $w_{n,r}^n$ converges to x_{\max} , and thus the right extreme reference point $r_{n+1}^n := (r_1, f(w_{n,r}^n))$ converges to $(r_1, f(x_{\max}))$ by continuity of f . Therefore, if r_1 is strictly larger than x_{\max} , then there exists N such that for all $n \geq N$, r_{n+1}^n verifies the assumptions on the reference point of Lemma B.1 which guarantee that u_{\max} is the unique maximizer of $\text{HV}_r(\cdot)$ over the right extreme gap region $\mathcal{G}_{\mathcal{S}_n, n+1}^r$. Let assume that u_{\max} does not belong to \mathcal{S}_n . Then, $w_{N,r}^N \neq x_{\max}$, and since $w_{n,r}^n$ converges to x_{\max} , the left extreme gap region $\mathcal{G}_{\mathcal{S}_n, i}^r$ is necessarily filled at some later iteration. When the right extreme gap region is filled, u_{\max} , the unique minimizer of $\text{HV}_r(\cdot)$ over this gap region, is added to the greedy set. To summarize, if $r_1 > x_{\max}$, then for n large enough \mathcal{S}_n contains u_{\max} , and thus the right extreme gap region is empty. We can prove with the same approach that for $r_2 > f(x_{\min})$, \mathcal{S}_n contains u_{\min} for n large enough.

At any iteration, the non-extreme reference points dominate the nadir point. Additionally, we proved that either $r_1 < x_{\max}$ (resp. $r_2 < f(x_{\min})$), and thus the left (resp. right) extreme reference point dominates the nadir point or for n large enough, the left (resp. right) extreme gap region is empty. \square

Acknowledgments. We would like to thank Dima Brockhoff and Nikolaus Hansen for several helpful discussions, notably about the importance of decomposing the optimality gap and for the conjecture of (4.14).

REFERENCES

- [1] C. AUDET, G. SAVARD, AND W. ZGHAL, *Multiobjective Optimization Through a Series of Single-Objective Formulations*, SIAM Journal on Optimization, 19 (2008), pp. 188–210, <https://doi.org/10.1137/060677513>.
- [2] A. AUGER, J. BADER, AND D. BROCKHOFF, *Theoretically investigating optimal μ -distributions for the hypervolume indicator: first results for three objectives*, in Proceedings of the 11th international conference on Parallel problem solving from nature: Part I, PPSN'10, Berlin, Heidelberg, Sept. 2010, Springer-Verlag, pp. 586–596.
- [3] A. AUGER, J. BADER, D. BROCKHOFF, AND E. ZITZLER, *Theory of the Hypervolume Indicator: Optimal μ -Distributions and the Choice of the Reference Point*, in Proceedings of the Tenth ACM SIGEVO Workshop on Foundations of Genetic Algorithms, FOGA '09, New York, NY, USA, Jan. 2009, Association for Computing Machinery, pp. 87–102, <https://doi.org/10.1145/1527125.1527138>.
- [4] A. AUGER, J. BADER, D. BROCKHOFF, AND E. ZITZLER, *Hypervolume-based multiobjective optimization: Theoretical foundations and practical implications*, Theoretical Computer Science, 425 (2012), pp. 75–103, <https://doi.org/10.1016/j.tcs.2011.03.012>.

- 925 [5] N. BEUME, M. LAUMANN, AND G. RUDOLPH, *Convergence Rates of (1+1) Evolutionary Multiobjec-*
926 *tive Optimization Algorithms*, in Parallel Problem Solving from Nature, PPSN XI, R. Schaefer,
927 C. Cotta, J. Kołodziej, and G. Rudolph, eds., Lecture Notes in Computer Science, Berlin,
928 Heidelberg, 2010, Springer, pp. 597–606, https://doi.org/10.1007/978-3-642-15844-5_60.
- 929 [6] N. BEUME, B. NAUJOKS, AND M. EMMERICH, *SMS-EMOA: Multiobjective selection based on dom-*
930 *inated hypervolume*, European Journal of Operational Research, 181 (2007), pp. 1653–1669,
931 <https://doi.org/10.1016/j.ejor.2006.08.008>.
- 932 [7] J. BIGEON, S. LE DIGABEL, AND L. SALOMON, *DMulti-MADS: Mesh adaptive direct multisearch*
933 *for bound-constrained blackbox multiobjective optimization*, Computational Optimization and
934 Applications, 79 (2021), pp. 301–338, <https://doi.org/10.1007/s10589-021-00272-9>.
- 935 [8] K. BRINGMANN AND T. FRIEDRICH, *The maximum hypervolume set yields near-optimal approx-*
936 *imation*, in Proceedings of the 12th Annual Conference on Genetic and Evolutionary Com-
937 *putation - GECCO '10*, Portland, Oregon, USA, 2010, ACM Press, pp. 511 – 518, <https://doi.org/10.1145/1830483.1830576>.
- 938 [9] D. BROCKHOFF, *Optimal μ -distributions for the hypervolume indicator for problems with linear bi-*
939 *objective fronts: exact and exhaustive results*, in Proceedings of the 8th international conference
940 *on Simulated evolution and learning, SEAL'10*, Berlin, Heidelberg, Dec. 2010, Springer-Verlag,
941 pp. 24–34.
- 942 [10] A. L. CUSTÓDIO AND J. F. A. MADEIRA, *MultiGLODS: Global and local multiobjective optimization*
943 *using direct search*, Journal of Global Optimization, 72 (2018), pp. 323–345, <https://doi.org/10.1007/s10898-018-0618-1>.
- 944 [11] A. L. CUSTÓDIO, J. F. A. MADEIRA, A. I. F. VAZ, AND L. N. VICENTE, *Direct Multisearch for*
945 *Multiobjective Optimization*, SIAM Journal on Optimization, 21 (2011), pp. 1109–1140, <https://doi.org/10.1137/10079731X>.
- 946 [12] E. DE KLERK, F. GLINEUR, AND A. B. TAYLOR, *On the worst-case complexity of the gradient method*
947 *with exact line search for smooth strongly convex functions*, Optimization Letters, 11 (2017),
948 pp. 1185–1199, <https://doi.org/10.1007/s11590-016-1087-4>.
- 949 [13] R. FIORENZA, *Hölder and locally Hölder Continuous Functions, and Open Sets of Class C^k , $C^{k,\lambda}$* ,
950 *Frontiers in Mathematics*, Birkhäuser Basel, 2016, <https://doi.org/10.1007/978-3-319-47940-8>.
- 951 [14] E. H. FUKUDA AND L. M. G. DRUMMOND, *On the convergence of the projected gradient method*
952 *for vector optimization*, Optimization, 60 (2011), pp. 1009–1021, <https://doi.org/10.1080/02331934.2010.522710>.
- 953 [15] J. KNOWLES AND D. CORNE, *On metrics for comparing nondominated sets*, in Proceedings of the
954 *2002 Congress on Evolutionary Computation. CEC'02*, vol. 1, May 2002, pp. 711–716 vol.1,
955 <https://doi.org/10.1109/CEC.2002.1007013>.
- 956 [16] E. MARESCAUX AND N. HANSEN, *Hypervolume in biobjective optimization cannot converge faster*
957 *than $\Omega(1/p)$* , in Proceedings of the Genetic and Evolutionary Computation Conference, GECCO
958 '21, New York, NY, USA, June 2021, Association for Computing Machinery, pp. 430–438,
959 <https://doi.org/10.1145/3449639.3459371>.
- 960 [17] G. L. NEMHAUSER, L. A. WOLSEY, AND M. L. FISHER, *An analysis of approximations for max-*
961 *imizing submodular set functions—I*, Mathematical Programming, 14 (1978), pp. 265–294,
962 <https://doi.org/10.1007/BF01588971>.
- 963 [18] C. TOURÉ, N. HANSEN, A. AUGER, AND D. BROCKHOFF, *Uncrowded hypervolume improvement:*
964 *COMO-CMA-ES and the sofomore framework*, in Proceedings of the Genetic and Evolution-
965 *ary Computation Conference, GECCO '19*, New York, NY, USA, July 2019, Association for
966 *Computing Machinery*, pp. 638–646, <https://doi.org/10.1145/3321707.3321852>.
- 967 [19] D. A. V. VELDHUIZEN AND G. B. LAMONT, *Evolutionary Computation and Convergence to a Pareto*
968 *Front*, in Proceedings of the Third Annual Conference on Genetic Programming, San Francisco,
969 1998, Stanford University Bookstore, pp. 221–228.
- 970 [20] E. ZITZLER, J. KNOWLES, AND L. THIELE, *Quality Assessment of Pareto Set Approximations*, in
971 *Multiobjective Optimization: Interactive and Evolutionary Approaches*, J. Branke, K. Deb,
972 K. Miettinen, and R. Słowiński, eds., Lecture Notes in Computer Science, Springer, Berlin,
973 Heidelberg, 2008, pp. 373–404, https://doi.org/10.1007/978-3-540-88908-3_14.