



**HAL**  
open science

## Designing a blueprint for coral reef survival

Joan Kleypas, Denis Allemand, Ken Anthony, Andrew Baker, Michael Beck, Lynne Zeitlin Hale, Nathalie Hilmi, Ove Hoegh-Guldberg, Terry Hughes, Les Kaufman, et al.

► **To cite this version:**

Joan Kleypas, Denis Allemand, Ken Anthony, Andrew Baker, Michael Beck, et al.. Designing a blueprint for coral reef survival. *Biological Conservation*, 2021, 257, pp.109107. 10.1016/j.biocon.2021.109107 . hal-03201225

**HAL Id: hal-03201225**

**<https://hal.science/hal-03201225>**

Submitted on 3 May 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Designing a blueprint for coral reef survival

The Pew Marine Fellows Program and The Ocean Solutions Initiative

- Joan Kleypas:** National Center for Atmospheric Research, Boulder, Colorado, USA  
**Denis Allemand:** Centre Scientifique de Monaco, Monaco, Principality of Monaco  
**Ken Anthony:** Australian Institute of Marine Science, Townsville, Queensland, Australia  
**Andrew C. Baker:** Department of Marine Biology and Ecology, Rosenstiel School of Marine and Atmospheric Science, University of Miami, Miami, Florida, USA  
**Michael Beck:** University of California, Santa Cruz, California, USA  
**Lynne Zeitlin Hale:** The Nature Conservancy, Narragansett, Rhode Island, USA  
**Nathalie Hilmi:** Centre Scientifique de Monaco, Monaco, Principality of Monaco  
**Ove Hoegh-Guldberg:** Global Change Institute, University of Queensland, St Lucia, Queensland, Australia  
**Terry Hughes:** ARC Centre of Excellence for Coral Reef Studies, James Cook University, Townsville, Queensland, Australia  
**Les Kaufman:** Boston University Marine Program, Boston, Massachusetts, USA  
**Hajime Kayanne:** Department of Earth and Planetary Science, University of Tokyo, Tokyo, Japan  
**Alexandre K. Magnan:** IDDRI (Sciences Po), CNRS and University of la Rochelle, France  
**Elizabeth Mcleod:** The Nature Conservancy, Austin, Texas, USA  
**Peter Mumby:** School of Biological Sciences, University of Queensland, St. Lucia, Queensland, Australia  
**Stephen Palumbi:** Department of Biology, Stanford University, Hopkins Marine Station, Pacific Grove, California, USA  
**Robert H. Richmond:** University of Hawaii at Manoa, Kewalo Marine Laboratory, Honolulu, Hawaii, USA  
**Baruch Rinkevich:** Israel Oceanographic & Limnological Research, National Institute of Oceanography, Tel Shikmona, Haifa, Israel  
**Robert S. Steneck:** University of Maine, School of Marine Sciences, Darling Marine Center, Walpole, Maine, USA  
**Christian R. Voolstra:** Department of Biology, University of Konstanz, Konstanz, Germany  
**David Wachenfeld:** Great Barrier Reef Marine Park Authority, Townsville, Queensland, Australia  
**Jean-Pierre Gattuso:** CNRS, Sorbonne Université and IDDRI (Sciences Po), France

## Abstract

Maintaining coral reef ecosystems is a social imperative, because so many people depend on coral reefs for food production, shoreline protection, and livelihoods. The survival of reefs this century, however, is threatened by the mounting effects of climate change. Climate mitigation is the foremost and essential action to prevent coral reef ecosystem collapse. Without it, reefs will become extremely diminished within the next 20-30 years. Even with strong climate mitigation, however, existing conservation measures such as marine protected areas and fisheries management are no longer sufficient to sustain the ecosystem and many additional and innovative actions to increase reef resilience must also be taken. In this paper we assess the suite of protections and actions in terms of their potential be effective according to a set of criteria that include effectiveness, readiness, co-benefits and disbenefits. Even with the best scientific innovation, saving coral reefs will require a well-funded, well-designed, and rapidly executed strategy with political and social commitments at the level of other grand challenges.

## Keywords

Coral reefs, climate change, conservation strategy, ecosystem restoration

## 1. Introduction

Over two recent meetings, the authors of this Perspective assessed (a) the need for an ambitious but realistic plan to sustain coral reefs through the 21<sup>st</sup> century; and (b) the need to objectively evaluate interventions that should be included in the plan. The urgency of taking action is clear based on many previous projections of coral decline. Here we present additional projections to illustrate that a strategy to save coral reefs requires both a reduction in ocean warming and an increase coral resiliency. For the latter we provide a summary of an assessment of various interventions through expert opinion. We then outline the principles of a coordinated strategy to sustain reefs, with the goal of stimulating the design of a blueprint with input from many more stakeholders.

## 2. Background

Coral reefs are succumbing rapidly to rising ocean temperatures, and the recent and rapid degradation of reefs worldwide is well documented (Bindoff et al., 2019). Passive recovery (i.e., natural recovery without human intervention) is proving increasingly inadequate (Ortiz et al., 2018) as coral bleaching and mortality events become more frequent and severe, adding to the impacts of local anthropogenic stressors such as overfishing and pollution (Hughes et al., 2018). As stressors increase and coral cover declines, so do reproductive and recruitment success, preventing many reefs from recovering (Richmond et al., 2018). Major losses of reef corals and reef structure are expected if global warming exceeds 1.5°C above the pre-industrial average temperature (Hoegh-Guldberg et al., 2018), and the most effective action to reduce the decline of coral reefs remains rapid and effective mitigation of greenhouse gas emissions (Hoegh-Guldberg et al., 2019). Even if the Paris Agreement is fully implemented, however, warming over the coming two-three decades will increasingly threaten the ability of reefs to recover.

To illustrate this trajectory, a new modelling analysis was performed by combining a coral bleaching model (van Hooidonk et al., 2013) with sea surface temperature projections for two emission scenarios from the NCAR Community Earth System Model Version 1.2 (Fig. 1A; see Supplementary 2 for details): high (RCP 8.5; RCP=Representative Carbon Pathway) and low emissions (RCP 2.6). We also allowed corals and coral communities to adapt, both naturally and through human actions, at a rate consistent with the change in the average annual maximum temperature of the previous 50 years (see Logan et al., 2014), which for this analysis infers a variable rate of adaptation with a maximum of 0.2°C per decade (Fig. 1B). While some species may adapt more quickly, and others more slowly, we assume this rate includes the overall ability of reef communities to persist due to community shifts, natural adaptation, and resilience gained from human interventions (discussed below).

Regardless of the shape of the trajectory, reefs will experience a rapid increase in exposure to temperature anomalies predictive of severe bleaching (more than two severe bleaching events per decade), which will be unsustainable for most coral species (Fig. 1B).

The main advantage of the low-emissions trajectory is that it buys time for corals to adapt to rising temperature through both natural processes and human interventions (Bindoff et al., 2019). Relative to the high-emissions trajectory, adaptation could greatly reduce the percentage of reefs experiencing high-frequency bleaching, despite exposure to increasingly elevated temperature, through 2040–2050 (Fig. 1B, see also Logan et al., 2014). However, the difference between the high and low emissions trajectories becomes more apparent after 2050. Under the high-emissions trajectory, temperature continues to increase, and adaptation and interventions can delay but not prevent the eventual high-frequency bleaching of all reefs. Under the low-emissions trajectory, the temperature increase slows, allowing some corals to adapt (Fig. 1B). These results are similar to adaptation models that incorporate phenotype shifts (Logan et al., 2014; Walsworth et al., 2019) or genomic-based adaptation to warming temperatures (Bay et al., 2017); i.e., adaptation delays the loss of coral populations but even these fail under all but the low emission climate scenarios (Fig. 1B, C).

The window for opportunity to act on both mitigation and adaptation is thus between now and 2050 (Anthony et al., 2017; Hardisty et al., 2020). How fast we act within this window will determine what reefs, species, and ecosystem functions and services can be sustained. Without action, coral reefs as we know them today – and their associated services to humankind (food production, shoreline protection, tourism, economic resources, cultural values) – could therefore be one of the first major ecosystems in this century to collapse under the weight of climate change.

### **3. Evaluating existing and new interventions to improve the outlook for reefs**

#### **3.1 Actions to sustain coral reefs**

In addition to mitigation of greenhouse gas emissions, we consider a suite of actions (Fig. 2) that fall within two categories: (1) solar radiation modification (SRM), and (2) enhancing biological, ecological, and socio-economic adaptation. SRM consists of untested strategies to reflect sunlight back to space, such as stratospheric aerosol injection that acts at the global scale, marine cloud brightening at the regional scale, surface albedo enhancement, and shading of the ocean surface that work at the local scale. We do not condone SRM because it does not reduce ocean acidification, and presents significant uncertainties and risks (Gattuso et al., 2018). SRM and other physical interventions to cool reef waters, such as pumping deep seawater into reef areas, were not included in the modeling predictions in Fig. 1B, but we list them here because they are considered by some proponents as local, high-cost actions to reduce heat stress in tourist zones or other high value areas.

Biological, ecological, and socio-economic actions are necessary partners to mitigation of emissions. These aim to enhance the natural resilience and stabilization of corals, reefs and societies (Gattuso et al., 2018). Many of the interventions are existing, proven practices that reduce reef stressors, such as management of watersheds, coastal zones, and fisheries, often as part of the establishment of marine protected areas, and often

reinforced with socio-economic incentives and regulatory measures to protect coastal ecosystems, and maintain ecosystem services (Hilmi et al., 2019). An additional suite of interventions is being developed to support biological and ecological adaptation of reef organisms, including ecological engineering, assisted adaptation, and managed relocation (Rinkevich, 2020). While most of the existing practices aim to indirectly increase coral reef resilience by improving environmental conditions, most of the newer interventions aim to directly increase resilience of corals and other reef building organisms to environmental change.

The US National Academy of Sciences reviewed the science and risks/benefits of interventions to enhance the persistence and resilience of coral reefs (National Academies of Sciences Engineering and Medicine, 2018, 2019). Many of these interventions build on the practice of active reef restoration (Rinkevich, 2014), which entails sexual and asexual propagation of corals for outplanting into the natural environment. Active reef restoration will most likely succeed in the long term if it also incorporates (a) innovations such as assisted fertilization and recruitment (de la Cruz et al., 2015; Nakamura et al., 2011); (b) actions to speed natural selection and adaptation through assisted gene flow and managed relocation (Aitken and Whitlock, 2013; Van Oppen et al., 2017) with possible incorporation of assisted adaptation and evolution (Van Oppen et al., 2017); and (c) a focus on preserving and enhancing species diversity and genetic variation within populations, and managed species assembly to maximize the best adapted species consortia (National Academies of Sciences Engineering and Medicine, 2019). A plan for a World Coral Conservatory, i.e., *ex situ* coral conservation in aquaria, has also been proposed (Zoccola et al., 2020).

Deployment of most of these interventions will rely on active restoration in regions where pollution, overfishing, and other human-driven reef stressors are well managed. Active restoration still focuses on too few species and is limited to small spatial scales: even large efforts struggle to restore a hectare per year. The effectiveness of active restoration thus depends on strategies that harness coral population connectivity to accelerate the spread of heat-tolerant corals, i.e., focusing restoration on reefs where larval dispersal and recruitment can spark year-to-year spread over larger areas (Hock et al., 2017; Walsworth et al., 2019). In some regions, structural restoration will be needed to enhance ecological recovery of shorelines (Beck et al., 2018) and fish production (Rogers et al., 2018). All of these innovations, which are broadly reviewed in the context of management, policymaking, and fund-raising (Hein et al., 2020), will require rigorous field testing prior to large-scale deployment.

### **3.2 Assessment of actions**

Although coral reef scientists and managers agree that an ambitious plan is needed to fortify coral reefs against climate change and other stressors, there is considerable debate on the elements that should be included in such a plan. The participants of both teams of this paper (n=21) were asked to provide their expert opinion for each of the actions in Fig. 2 based on

six criteria and with the next 20-30 years as a timeframe: (1) *Effectiveness* in moderating drivers of ocean changes and/or impacts on coral reefs; (2) *Readiness*: the current state of technological development; (3) *Co-benefits*, or additional benefits to other ecosystems; (4) *Disbenefits*, or adverse consequences; (5) *Acceptability* of the action culturally, societally, politically; and (6) *Scale* (see Supplementary 3). A 0-5 scoring system (unknown, very low to very high) was used except for Scale, which was scored from 1-3 (local, regional, global) (Fig. 3A, B). An overall score was determined as the sum of the first five criteria, while scale was addressed separately (Fig. 3C).

Addressing the causes of climate change ranked the highest overall. Among the suite of local to regional actions to support biological, ecological, and societal adaptation, those that have been in place for longest (e.g., improved governance, coastal zone management, fisheries management) were scored as effective, ready, with mostly high co-benefits and few disbenefits, but had mixed scores in terms of acceptability, which reflect conflicts associated with increased regulations. Removal of predators and invasive species was broadly considered to be less effective as a stand-alone action, but perhaps necessary to guarantee success of other actions (e.g., reef restoration; assisted fertilization and recruitment). The four remaining biological and ecological interventions (restoration, managed relocation, assisted adaptation, and assisted fertilization) are untested, or still confined to local deployment. There was relatively low agreement amongst the responders on their effectiveness, readiness and co-benefits, although most actions scored highly in terms of lacking disbenefits. Among these, assisted adaptation and evolution received the lowest overall scores, and reef restoration the highest. The issue of scalability (across both space and across a range of species) was cited as a major reason for the low effectiveness scores, except for assisted fertilization and recruitment, which has the potential to operate over larger scales.

The geoengineering approaches received the lowest overall scores, reflecting a high degree of uncertainty and difficulty with evaluating them. Notably, co-benefits were considered to be low but disbenefits high, particularly with the regional to global scale actions of cloud brightening, aerosol-based SRM and space-based SRM.

The overall score (Fig. 3C) is an insufficient assessment of each action because one cannot assume that the criteria of evaluation are equally important. However, it does provide a coarse insight into expert opinions. Among the two global-scale actions, only reducing carbon emissions scored highly (aggregated score of 24.5 out of a maximum score of 25) while the other (aerosol-based SRM) scored very low (< 7 out of 25). Most of the regional scale actions are already well-established within “coastal zone management.” Among the local actions, a few would rank more highly if scalability could be increased, such as through networks of local actions (restoration, assisted reproduction, assisted gene flow) to increase resilience within a reproductively connected population or metapopulation.

This expert assessment can serve as an example framework for constructing a blueprint for sustaining coral reefs. The evaluation highlighted the consensus that mitigation

of greenhouse gas emissions is not only essential to reef survival, but is also the most wide-reaching, effective, achievable, beneficial, and acceptable action. It also highlighted that reduced emissions must be paired with continued local to regional actions to protect the coral reef environment and to increase coral reef resilience. Many of these actions need improvement, and some need further development and testing with the acknowledgment that some may never be implemented if risks are deemed to be too high.

#### **4. A blueprint for reef survival**

At the outset of designing a blueprint to sustain coral reefs into the future, three key constraints must be acknowledged. First, fully protecting and restoring the global coral reef estate of the 1890s or 1990s is impossible, even if global warming is kept to 1.5°C. Areas for protection and restoration must be prioritised and one must accept that future reefs will be very different than reefs of the past. Second, the goals of reef protection and restoration are already shifting from what is desirable (e.g., species protection, maintaining species diversity, and returning the ecosystem to its former state) to what is possible (e.g., maintaining sustainable reef function, supporting fisheries, culture and tourism, and enhancing reef-building to protect shorelines). Third, current approaches to reef conservation must be rapidly converted to an organized strategy, underwritten by strong governance and investment, to sustain coral reefs and the services they render to humankind.

##### **4.1 An organized strategy**

We stress the need for an organized strategy that coordinates the science, policy, governance, and investment to achieve the common goal to sustain coral reefs as viable socio-ecological systems into the future. Overall, the *scientific commitment* to the blueprint should build on and improve previous actions to reduce local to regional stresses on coral reefs but will also need to encourage and test new innovations, as has been required for all previous world challenges. Even the best scientific innovation, however, requires a level of political, social, and economic commitment similar to when society came together to confront other huge challenges. A few prominent examples include the US Apollo program, efforts to cure cancer and AIDS, and eliminating famine, each of which have achieved different levels of success. The two necessary actions to save coral reefs, mitigating climate change and increasing coral resilience, require different models.

*The challenge of mitigating climate change* is often compared to the challenge of the Apollo program to put a human on the moon (King et al., 2015). The Apollo program was largely an engineering challenge, received tremendous financial backing (about 4% of the US federal budget in each of the years 1964-1967), and employed thousands of scientists and engineers. Solving climate change through innovations in renewable energy and carbon capture is similarly considered a technological challenge, although failures in the political and economic systems have greatly impeded progress (van Renssen, 2018).



*The challenge of saving an ecosystem*, however, requires more than technology and must be deployable at hundreds to thousands of locations in developed and developing countries. An effort to save the reef ecosystem is actually more akin to global efforts to cure disease. Using cancer as an example, research led by the U.S. National Cancer Institute has led to a substantial decline in mortality due to progress not only in treatment, but also in social programs that promote positive human behaviors such as healthy eating and reduced smoking (Kort et al. 2009). Some of the biggest hurdles to finding cures for cancer include the disease's inherent biological complexity, the need for translational research (applying biological findings and clinical trials to improve health outcomes), and the drug approval process. Other hurdles include delivering these benefits to communities throughout the world, often at medical technology centers. A strategy to save coral reefs should build on these lessons, for example: new technologies that can scale up reef restoration and adaptation, policies and strategies to accelerate widespread deployment while minimizing risks; and incentivizing local communities and global society to adopt behaviors that improve the environmental conditions on reefs. The latter requires efforts to vastly improve public literacy on ocean issues (Steel et al., 2005) with an approach tailored to different nationalities, cultures and demographic groups (Buckley et al., 2017).

The urgency of the coral reef crisis also calls for much stronger coordination across disciplines. Traditional coral reef science is mostly carried out by biological researchers who are rarely organized within a larger plan to help sustain reefs. The basis for this coordination would ideally take advantage of multiple existing organizations and networks (e.g., International Coral Reef Society, Reef Resilience Network, International Coral Reef Initiative, the Coral Restoration Consortium), and numerous government organizations, NGOs, and private institutions that are already partnering to achieve common objectives. Such partnerships must be strengthened with a clearly defined mission of top priorities over the next 20–30 years, and a flexible plan for coordinating the multiple measures of adaptation (Fig. 2).

#### **4.2 Strong Backing by Governments and Investment**

Large-scale missions are more likely to succeed when they have well-defined goals and outcomes, and “when government plays a deliberate role in directing innovation and growth to achieve a desired objective” (Mazzucato, 2016). Policies to sustain coral reefs should be based on achieving a series of prioritized objectives rather than a set of poorly defined goals (e.g., “save coral reefs”). This may be the most difficult challenge, because priorities (ecosystem function, biodiversity, fisheries support, tourism, reduction in shoreline flooding and erosion) will vary from region to region and across coral population boundaries, which are inherently transnational, or even trans-governance within national jurisdictions (e.g., Great Barrier Reef; village or atoll-level governance across wide stretches of the Indo-Pacific).

Previous multi-national agreements to protect marine resources provide examples and lessons in how this challenge can be achieved (e.g., the Micronesia Challenge, Coral Triangle Initiative, Western Indian Ocean Challenge, Caribbean Challenge). One can also build on several regional plans under development, such as Australia's Reef Restoration and Adaptation Plan (RRAP) for the Great Barrier Reef (Hardisty et al., 2020). The Australian government has begun to identify potentially scalable actions and develop an economic case for further research, followed by an AU\$100m investment to build a decision-support system that allows potential restoration options to be integrated with other management actions. In the U.S.A., Congress is considering modest funding (US\$36m) to strengthen coral reef conservation and restoration, and increase collaboration among stakeholders (S2429, Restoring Resilient Reefs Act). The newly developed Global Fund for Coral Reefs, a partnership of private philanthropy, finance institutions, and UN agencies, seeks to invest US\$500m through blended finance in coral reef conservation over the next decade. These efforts are a good start, but the cost of sustaining enough of the coral reef ecosystem to prevent collapse will require billions in US dollars. While this value seems high, by comparison hundreds of billions of US dollars will be required to protect U.S. coastal infrastructure from rising sea level alone over the next 20 years (LeRoy et al., 2019).

## **5. Conclusions**

The scientific vision for a 20 to 30-year blueprint should prioritize reducing greenhouse gas emissions and enhancing the capacity of reefs to recover from climate impacts, adapt to stress, and respond rapidly to their changing environment. Some strategies, such as supported breeding and assisted gene flow, can be deployed now. As demonstrated for biomedical research (Chalmers et al., 2014), new technologies will require translational research to speed up their testing, risk assessment, approvals, and deployment in the field. One must hasten natural selection of a wide array of species that are better adapted to the new biophysical regime, and then find ways to sustain those adaptations throughout entire populations, which can range over hundreds, even thousands of kilometers. Dissemination of adapted corals and other innovations could be modeled after agricultural and forest extension services, which provide guidance to cultivators toward improving their productivity, including in the context of climate change, and in the value of species and genetic diversity in the face of an uncertain future. Finally, strong governance that includes a supportive regulatory framework for implementing and enforcing policies must be addressed concurrently with the scientific approaches and public communication and engagement; otherwise, critical time will be lost along with the potential for success.

We present this Perspective as a broad view of the global challenge to save coral reefs, and to help define the elements needed to design a comprehensive plan. What we propose is possible with a clear 20 to 30-year plan that attracts the necessary public interest, investment, and policies on par with major efforts in space research and human health. It requires a 'Grand Bargain' with world leaders: if they reduce global emissions to keep

warming below 1.5°C, reef scientists will work in partnership with management agencies, conservation NGOs, and communities to save a sufficient number of reefs to sustain key services to coastal communities and allow their global recovery once emissions are effectively mitigated. The economic value of coral reefs in US waters alone is estimated at US\$3.6 billion annually (Brander and Van Beukering, 2013), and globally the value is at least an order of magnitude higher. With support equal to the scale of this problem, coral reefs can be protected for future generations. Developing a blueprint to save coral reefs may be the first major attempt to save an ecosystem, but it will also provide a roadmap of action for other ecosystems at risk (Duarte et al., 2020).

## **Acknowledgements**

This article is dedicated to the memory of our colleague and friend Ruth Gates. We thank M. Khamla for help with artwork and Ruben van Hooidonk for providing the reef locations. This work is sponsored by the Pew Marine Fellows program and "The Oceans Solutions Initiative" supported by the Prince Albert II of Monaco Foundation, the Ocean Acidification International Coordination Centre of the International Atomic Energy Agency, the Veolia Foundation, the French Facility for Global Environment, and the STORISK project (ANR-15CE03-0003). Modeled data presented in Fig. 1A,B are based upon work supported by the National Center for Atmospheric Research, which is a major facility sponsored by the National Science Foundation under Cooperative Agreement No. 1852977. No authors have competing interests.

## **Supporting Information**

**Supplementary 1.** List of contributors and their affiliations

**Supplementary 2.** Methods used to calculate modeled projections of coral bleaching

**Supplementary 3.** Survey methods and results of expert opinions of innovations to increase the resilience of coral reefs based on: *Effectiveness, Readiness, Co-benefits, Disbenefits, Acceptability, and Scale.*

## **References**


Aitken, S.N., Whitlock, M.C., 2013. Assisted gene flow to facilitate local adaptation to climate change. *Annual Review of Ecology, Evolution, and Systematics* 44, 367-388. DOI: 10.1146/annurev-ecolsys-110512-135747

- Anthony, K., Bay, L.K., Costanza, R., Firm, J., Gunn, J., Harrison, P., Heyward, A., Lundgren, P., Mead, D., Moore, T., Mumby, P.J., van Oppen, M.J.H., Robertson, J., Runge, M.C., Suggett, D.J., Schaffelke, B., Wachenfeld, D., Walshe, T., 2017. New interventions are needed to save coral reefs. *Nature Ecology & Evolution* 1, 1420-1422. DOI: 10.1038/s41559-017-0313-5
- Bay, R.A., Rose, N.H., Logan, C.A., Palumbi, S.R., 2017. Genomic models predict successful coral adaptation if future ocean warming rates are reduced. *Science Advances* 3, 10.1126/sciadv.1701413. DOI: 10.1086/691233
- Beck, M.W., Losada, I.J., Menéndez, P., Reguero, B.G., Díaz-Simal, P., Fernández, F., 2018. The global flood protection savings provided by coral reefs. *Nature Communications* 9, doi: 10.1038/s41467-41018-04568-z. DOI: 10.1038/s41467-018-04568-z
- Bindoff, N.L., Cheung, W.W.L., Kairo, J.G., Arístegui, J., Guinder, V.A., Hallberg, R., Hilmi, N., Jiao, N., Karim, M.S., Levin, L., O'Donoghue, S., Purca Cuicapusa, S.R., Rinkevich, B., Suga, T., Tagliabue, T., Williamson, P., 2019. Changing ocean, marine ecosystems, and dependent communities, In IPCC Special Report on the Ocean and Cryosphere in a Changing Climate. eds H.-O. Pörtner, D.C. Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E. Poloczanska, K. Mintenbeck, A. Alegría, M. Nicolai, A. Okem, J. Petzold, B. Rama, N.M. Weyer.
- Brander, L.M., Van Beukering, P., 2013. The total economic value of U.S. coral reefs: A review of the literature, In NOAA Coral Reef Conservation Program. p. 32. NOAA, Silver Spring, MD.
- Buckley, P.J., Pinnegar, J.K., Painting, S.J., Terry, G., Chilvers, J., Lorenzoni, I., Gelcich, S., Duarte, C.M., 2017. Ten thousand voices on marine climate change in Europe: Different perceptions among demographic groups and nationalities. *Frontiers in Marine Science* 4. DOI: 10.3389/fmars.2017.00206
- Chalmers, I., Bracken, M.B., Djulbegovic, B., Garattini, S., Grant, J., Gulmezoglu, A.M., Howells, D.W., Ioannidis, J.P.A., Oliver, S., 2014. How to increase value and reduce waste when research priorities are set. *Lancet* 383, 156-165. DOI: 10.1016/S0140-6736(13)62229-1
- dela Cruz, D.W., Rinkevich, B., Gomez, E.D., Yap, H.T., 2015. Assessing an abridged nursery phase for slow growing corals used in coral restoration. *Ecological Engineering* 84, 408-415. DOI: 10.1016/j.ecoleng.2015.09.042
- Duarte, C.M., Agusti, S., Barbier, E., Britten, G.L., Castilla, J.C., Gattuso, J.-P., Fulweiler, R.W., Hughes, T.P., Knowlton, N., Lovelock, C.E., Lotze, H.K., Predragovic, M., Poloczanska, E., Roberts, C., Worm, B., 2020. Rebuilding marine life. *Nature* 580, 39-51. DOI: 10.1038/s41586-020-2146-7
- Gattuso, J.-P., Magnan, A.K., Bopp, L., Cheung, W.W.L., Duarte, C.M., Hinkel, J., McLeod, E., Micheli, F., Oschlies, A., Williamson, P., Billé, R., Chalastani, V.I., Gates, R.D., Irsson, J.-O., Middelburg, J.J., Pörtner, H.-O., Rau, G.H., 2018. Ocean solutions to

- address climate change and its effects on marine ecosystems. *Frontiers in Marine Science* 5, art. 337. DOI: 10.3389/fmars.2018.00337
- Hardisty, P., Roth, C.H., Silvey, P.J., Mead, D., Anthony, K.R.N., 2020. Reef Restoration and Adaptation Program – Investment Case. A report provided to the Australian Government from the Reef Restoration and Adaptation Program p. 100, <http://www.gbrrestoration.org/reports>.
- Hein, M.Y., McLeod, I.M., Shaver, E.C., Vardi, T., Pioch, S., Boström-Einarsson, L., Ahmed, M., Grimsditch, G., 2020. Coral Reef Restoration as a strategy to improve ecosystem services – A guide to coral restoration methods, ed. United Nations Environment Program, Nairobi, Kenya, 60 p.
- Hilmi, N., Osborn, D., Acar, S., Bambridge, T., Chlous, F., Cinar, M., Djoundourian, S., Haraldsson, G., Lam, V.W.Y., Maliki, S., Mantuano, A.D.M., Marshall, N., Marshall, P., Pascal, N., Recuero-Virto, L., Rehdanz, K., Safa, A., 2019. Socio-economic tools to mitigate the impacts of ocean acidification on economies and communities reliant on coral reefs - a framework for prioritization. *Regional Studies in Marine Science* 28, 10.1016/j.rsma.2019.100559.
- Hock, K., Wolff, N.H., Ortiz, J.C., Condie, S.A., Anthony, K.R.N., Blackwell, P.G., Mumby, P.J., 2017. Connectivity and systemic resilience of the Great Barrier Reef. *Plos Biology* 15, doi: 10.1371/journal.pbio.2003355.
- Hoegh-Guldberg, O., Jacob, D., Taylor, M., Bindi, M., Brown, S., Camilloni, I., Diedhiou, A., Djalante, R., Ebi, K., Engelbrecht, F., Guiot, J., Hijioka, Y., Mehrotra, S., Payne, A., Seneviratne, S.I., Thomas, A., Warren, R., Zhou, G., 2018. Impacts of 1.5°C global warming on natural and human systems, In *Global warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty*. eds V. Masson-Delmotte, P. Zhai, H.O. Pörtner, D. Roberts, J. Skea, P.R. Shukla, A. Pirani, W. Moufouma-Okia, C. Péan, R. Pidcock, S. Connors, J.B.R. Matthews, Y. Chen, X. Zhou, M.I. Gomis, E. Lonnoy, T. Maycock, M. Tignor, T. Waterfield.
- Hoegh-Guldberg, O., Jacob, D., Taylor, M., Guillén Bolaños, T., Bindi, M., Brown, S., Camilloni, I.A., Diedhiou, A., Djalante, R., Ebi, K., Engelbrecht, F., Guiot, J., Hijioka, Y., Mehrotra, S., Hope, C.W., Payne, A.J., Pörtner, H.O., Seneviratne, S.I., Thomas, A., Warren, R., Zhou, G., 2019. The human imperative of stabilizing global climate change at 1.5°C. *Science* 365, eaaw6974. DOI: 10.1126/science.aaw6974
- Hughes, T.P., Anderson, K.D., Connolly, S.R., Heron, S.F., Kerry, J.T., Lough, J.M., Baird, A.H., Baum, J.K., Berumen, M.L., Bridge, T.C., Claar, D.C., Eakin, C.M., Gilmour, J.P., Graham, N.A.J., Harrison, H., Hobbs, J.P.A., Hoey, A.S., Hoogenboom, M., Lowe, R.J., McCulloch, M.T., Pandolfi, J.M., Pratchett, M., Schoepf, V., Torda, G.,


- Wilson, S.K., 2018. Spatial and temporal patterns of mass bleaching of corals in the Anthropocene. *Science* 359, 80-83. DOI: 10.1126/science.aan8048
- King, D., Browne, J., Layard, R., O'Donnell, G., M., R., Stern, N., Turner, A., 2015. A Global Apollo Programme to combat climate change, p. 38. Center for Economic Performance. [http://cep.lse.ac.uk/pubs/download/special/Global\\_Apollo\\_Programme\\_Report.pdf](http://cep.lse.ac.uk/pubs/download/special/Global_Apollo_Programme_Report.pdf)
- Kort, E.J., Paneth, N., Woude, G.F.V., 2009. The decline in US cancer mortality in people born since 1925. *Cancer Research* 69, 6500-6505. DOI: 10.1158/0008-5472.Can-09-0357
- LeRoy, S., Wiles, W., Chinowsky, P., Helman, J., 2019. High tide tax. The cost to protect coastal communities from rising seas, ed. The Center for Climate Integrity, p. 25, [https://www.climatecosts2040.org/files/ClimateCosts2040\\_Report.pdf](https://www.climatecosts2040.org/files/ClimateCosts2040_Report.pdf).
- Logan, C.A., Dunne, J.P., Eakin, C.M., Donner, S.D., 2014. Incorporating adaptive responses into future projections of coral bleaching. *Global Change Biology* 20, 125-139. DOI: 10.1111/gcb.12390
- Mazzucato, M., 2016. Innovation, the State, and Patient Capital, In *Rethinking Capitalism*. eds M. Jacobs, M. Mazzucato, pp. 98-118. Wiley-Blackwell, United Kingdom.
- Nakamura, R., Ando, W., Yamamoto, H., Kitano, M., Sato, A., Nakamura, M., Kayanne, H., Omori, M., 2011. Corals mass-cultured from eggs and transplanted as juveniles to their native, remote coral reef. *Marine Ecology Progress Series* 436, 161-168. DOI: 10.3354/meps09257
- National Academies of Sciences Engineering and Medicine, 2018. A Research Review of Interventions to Increase the Persistence and Resilience of Coral Reefs, In *The National Academies Press.*, p. 258, Washington, DC. <https://doi.org/10.17226/25279>
- National Academies of Sciences Engineering and Medicine, 2019. A Decision Framework for Interventions to Increase the Persistence and Resilience of Coral Reefs, p. 200, Washington, DC. <https://doi.org/10.17226/25424>
- Ortiz, J.C., Wolff, N.H., Anthony, K.R.N., Devlin, M., Lewis, S., Mumby, P.J., 2018. Impaired recovery of the Great Barrier Reef under cumulative stress. *Science Advances* 4, 10.1126/sciadv.aar6127.
- Richmond, R.H., Tisthammer, K.H., Spies, N.P., 2018. The effects of anthropogenic stressors on reproduction and recruitment of corals and reef organisms. *Frontiers in Marine Science* 5, 10.3389/fmars.2018.00226.
- Rinkevich, B., 2014. Rebuilding coral reefs: does active reef restoration lead to sustainable reefs? *Current Opinion in Environmental Sustainability* 7, 28-36. DOI: 10.1016/j.cosust.2013.11.018
- Rinkevich, B., 2020. Ecological engineering approaches in coral reef restoration. *Ices Journal of Marine Science*, doi: 10.1093/icesjms/fsaa1022. DOI: 10.1093/icesjms/fsaa022

- Rogers, A., Blanchard, J.L., Mumby, P.J., 2018. Fisheries productivity under progressive coral reef degradation. *Journal of Applied Ecology* 55, 1041-1049. DOI: 10.1111/1365-2664.13051
- Steel, B.S., Smith, C., Opsommer, L., Curiel, S., Warner-Steel, R., 2005. Public ocean literacy in the United States. *Ocean and Coastal Management* 48, 97-114. DOI: 10.1016/j.ocecoaman.2005.01.002
- van Hooidek, R., Maynard, J.A., Planes, S., 2013. Temporary refugia for coral reefs in a warming world. *Nature Climate Change* 3, 508-511. DOI: 10.1038/nclimate1829
- Van Oppen, M.J.H., Gates, R.D., Blackall, L.L., Cantin, N., Chakravarti, L.J., Chan, W.Y., Cormick, C., Crean, A., Damjanovic, K., Epstein, H., Harrison, P.L., Jones, T.A., Miller, M., Pears, R.J., Peplow, L.M., Raftos, D.A., Schaffelke, B., Stewart, K., Torda, G., Wachenfeld, D., Weeks, A.R., Putnam, H.M., 2017. Shifting paradigms in restoration of the world's coral reefs. *Global Change Biology* 23, 3437-3448. DOI: 10.1111/gcb.13647
- van Renssen, S., 2018. The inconvenient truth of failed climate policies. *Nature Climate Change* 8, 355-358. DOI: 10.1038/s41558-018-0155-4
- Walsworth, T.E., Schindler, D.E., Colton, M.A., Webster, M.S., Palumbi, S.R., Mumby, P.J., Essington, T.E., Pinsky, M.L., 2019. Management for network diversity speeds evolutionary adaptation to climate change. *Nature Climate Change* 9, 632–636. DOI: 10.1038/s41558-019-0518-5
- Zoccola, D., Ounais, N., Barthelemy, D., Calcagno, R., Gaill, F., Henard, S., Hoegh-Guldberg, O., Janse, M., Jaubert, J., Putnam, H., Salvat, B., Voolstra, C.R., Allemand, D., 2020. The World Coral Conservatory (WCC): A Noah's ark for corals to support survival of reef ecosystems. *Plos Biology* 18, e3000823. DOI: 10.1371/journal.pbio.3000823




**Figure 1.** Modeled projections of average sea surface temperature and bleaching: (A) temperatures for high (RCP8.5) and low (RCP2.6) emissions scenarios, and (B) the percentage of 1x1 degree reef cells experiencing high-frequency bleaching ( $\geq 2$  severe events per decade) for four combinations of mitigation and adaptation; lines are smoothed with 10-year moving averages. For comparison, (C) is the change in modeled coral population size under a multi-locus genetic adaptation model at different future emission levels (Bay et al., 2017), beginning in year 2010. The dashed lines in (B) showing bleaching frequency with adaptation are mirrored by the lines in (C) showing percentage of original coral population size.





**Figure 2.** Potential actions to sustain coral reefs and the benefits they provide, ranging from mitigation (addressing the cause of climate change), to a variety of biological, ecological, and social interventions and adaptations, and to solar radiation manipulation. An evaluation of these measures is illustrated in Fig. 3.



**Figure 3.** Results of expert ranking of potential actions shown in Figure 2 on (A) effectiveness, readiness, co-benefits, disbenefits (lack of), and acceptability, as well as on (B) scalability. Sum of scores in A are shown in (C).

## **Supplementary 1: List of contributors and their affiliations**

- This material is withheld during the double-blind review -

## Supplementary 2: Methods used to calculate modeled projections of coral bleaching

### Methods

Projections of temperature projections for Representative Carbon Pathways (RCP) 8.5 and 2.6 were based on sea surface temperature (SST) from the Community Earth System Model version 1.3 (CESM1.3) a fully coupled global climate model with atmosphere, land, sea ice, and ocean components, that provides state-of-the-art computer simulations of the Earth's past, present, and future climate states (Hurrell et al., 2013). This version uses the community atmospheric model version 5 (CAM5) with a 0.25° horizontal resolution the Parallel Ocean Program version 2 (POP2) with a nominal horizontal resolution of 1° (complete model details are described in Meehl et al. (2019)). Two simulations were analyzed for years 1950-2100, with years 1950-2005 constituting the historical climate simulation, and then branching at year 2005 into two 21<sup>st</sup> Century simulations. The first is forced by RCP2.6, which simulates a median globally averaged warming of 1.6°C in 2100 (likely range: 0.9–2.3°C) relative to 1850–1900, close to the target of the Paris Agreement to limit average global warming to 1.5°C. The second is forced by RCP8.5, which simulates a worst-case in average global warming of 4.3°C (likely range: 3.2–5.4°C) relative to 1850–1900 (IPCC, 2014). At the time of this paper, each scenario included a single ensemble member, and although these results are in line with similar model results, they should be considered as illustrative of the magnitude and timing of temperature projections for coral reefs.

Monthly SST fields were regridded to 1x1° spatial resolution, and the 1850-2000 SST time-series were extracted for each of 1-deg cells that included coral reef (using the same 1x1° data set of reef locations processed by (van Hooidonk et al., 2013)). The SST fields were bias corrected against the HadISST version 1 observed SSTs (Rayner et al., 2003) according to a procedure first described in (Dunne et al., 2013) and adapted by (Logan et al., 2014).

For each scenario, SSTs were averaged across all reef sites (Fig. 2A) over the 1950-2100 time period. For each reef location, coral bleaching was calculated for each year based on a determination of the Degree Heating Month (DHM), following the method of van Hooidonk, et al., (2013) where DHM was calculated as the accumulation of heat stress above the climatological maximum. A DHM value  $\geq 2$  indicated severe bleaching. Two different SST climatologies were used to calculate DHM. To determine bleaching with “no adaptation” a DHM climatology for each grid cell was calculated as the average of the warmest months for each year for the period 1980-1999. To determine bleaching “with adaptation” a rolling DHM climatology was calculated for each grid cell based on the average of the monthly maxima over some “adaptation period” determined as a previous number of years ( $n$ ). We examined adaptation periods ( $n$ ) of 10, 20, 30, 40, and 50 years and found that the adaptation rates for the 50-year period remained less than 0.2°C per decade;

this is in line with maximum acclimation and adaptation rates in corals, although it is likely that adaptation rates vary widely amongst the range of coral species on any single reef.

The number of severe bleaching events was determined for both the RCP2.6 and RCP8.5 scenarios, for both the “no adaptation” and “50-year adaptation” climatologies. The number of reefs experiencing two or more severe bleaching events in any one 10-y (van Hooijdonk et al., 2013) for each of the four conditions, as illustrated in Figure 2B.

## S2 References

- Dunne, J.P., Stouffer, R.J., John, J.G., 2013. Reductions in labour capacity from heat stress under climate warming. *Nature Climate Change* 3, 563-566. DOI: 10.1038/nclimate1827
- Hurrell, J.W., Holland, M.M., Gent, P.R., Ghan, S., Kay, J.E., Kushner, P.J., Lamarque, J.F., Large, W.G., Lawrence, D., Lindsay, K., Lipscomb, W.H., Long, M.C., Mahowald, N., Marsh, D.R., Neale, R.B., Rasch, P., Vavrus, S., Vertenstein, M., Bader, D., Collins, W.D., Hack, J.J., Kiehl, J., Marshall, S., 2013. The Community Earth System Model: A framework for collaborative research. *Bulletin of the American Meteorological Society* 94, 1339-1360. DOI: 10.1175/BAMS-D-12-00121.1
- IPCC, 2014. *Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Core Writing Team, R. K. Pachauri and L. A. Meyer (eds.)], 151 pp, IPCC, Geneva, Switzerland.
- Logan, C.A., Dunne, J.P., Eakin, C.M., Donner, S.D., 2014. Incorporating adaptive responses into future projections of coral bleaching. *Global Change Biology* 20, 125-139. DOI: 10.1111/gcb.12390
- Meehl, G.A., Yang, D., Arblaster, J.M., Bates, S.C., Rosenbloom, N., Neale, R., Bacmeister, J., Lauritzen, P.H., Bryan, F., Small, J., Truesdale, J., Hannay, C., Shields, C., Strand, W.G., Dennis, J., Danabasoglu, G., 2019. Effects of Model Resolution, Physics, and Coupling on Southern Hemisphere Storm Tracks in CESM1.3. *Geophysical Research Letters* 46: 12408-12416. DOI: 10.1029/2019GL084057
- Rayner, N.A., Parker, D.E., Horton, E.B., Folland, C.K., Alexander, L.V., Rowell, D.P., Kent, E.C., Kaplan, A., 2003. Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century. *Journal of Geophysical Research: Atmospheres* 108, 4407-. DOI: 10.1029/2002JD002670
- van Hooijdonk, R., Maynard, J.A., Planes, S., 2013. Temporary refugia for coral reefs in a warming world. *Nature Climate Change* 3, 508-511. DOI: 10.1038/nclimate1829

## **Supplementary 3: Survey methods and results of expert opinions of innovations to increase the resilience of coral reefs based on: *Effectiveness, Readiness, Co-benefits, Disbenefits, Acceptability, and Scale.***

### **Methods**

A blind assessment was performed in order to eliminate influence of others' authors scoring and reduce observer bias. The 21 authors were sent an empty spreadsheet with the following information on the criteria and scales.

#### **1. Effectiveness**

Potential to moderate the drivers of ocean changes (warming, acidification, sea level rise) and/or reducing the impacts of these changes on coral reefs, over the next 20-30 years.

Scoring scale: 0 to 5

0: no or negative effectiveness

1: very low effectiveness

2: low effectiveness

3: moderate effectiveness

4: high effectiveness

5: very high effectiveness

#### **2. Readiness**

This criterion describes the act current stage of technological development of the measure.

Scoring scale: 0 to 5

0: unknown

1: concept stage

2: only laboratory experiments

3: first real-world experiments

4: significant deployment already, with needs for improvements

5: fully ready

#### **3. Co-benefits**

This criterion describes the additional benefits of the measure, that is the benefits not related to those from the direct reduction of climate drivers or their impacts over the next 20-30 years. Co-benefits could include the improvement of the status of the ecosystems and/or the generation of ecosystem services

Scoring scale: 0 to 5

- 0: unknown level of co-benefits
- 1: very low level of co-benefits
- 2: low level of co-benefits
- 3: moderate level of co-benefits
- 4: high level of co-benefits
- 5: very high level of co-benefits

#### **4. Disbenefits**

This criterion describes the demonstrated or potential adverse consequences or negative impacts of the measure on ecosystems and their services over the next 20-30 years.

Scoring scale: 1 to 5.:

- 0: unknown level of disbenefits
- 1: very low level of disbenefits
- 2: low level of disbenefits
- 3: moderate level of disbenefits
- 4: high level of disbenefits
- 5: very high level of disbenefits

#### **5. Acceptability**

This criterion describes the expected degree of acceptance for each method culturally, societally, and politically over the next 20-30 years.

Scoring scale: 1 to 5.:

- 0: unknown level of acceptance
- 1: very low level of acceptance
- 2: low level of acceptance
- 3: moderate level of acceptance
- 4: high level of acceptance
- 5: very high level of acceptance

#### **6. Scale**

This criterion describes the operating scale of the measure over the next 20-30 years.

Scoring scale: 0 to 3

- 0: unknown
- 1: local
- 2: regional

3: global

## 7. Global score

A global score was calculated adding the scores of each criterion for each measure. Importantly, the scale for the Disbenefits has been reversed prior to the calculation in order to ensure that high scores are always more desirable than lower ones. Scores of ‘0’ were ignored in the calculation of the medians. The maximum value is 25.

## Summary of Results

**Table S3.1.** Median values of scores for each criterion and measure. Values of ‘0’ are excluded.

Measure	Effectiveness	Readiness	Cobenefits	Disbenefits	Acceptability	Scale	"Score" *
Addressing causes of climate change	5.0	4.0	5.0	1.0	4.0	3.0	23.0
Improved governance and regulation	4.0	4.0	5.0	1.0	4.0	2.0	22.0
Marine protected areas	3.0	5.0	4.0	1.0	3.0	2.0	20.0
Pollution reduction	4.0	4.0	4.0	1.0	4.0	1.5	21.0
Coastal zone management	3.0	4.0	4.0	1.0	3.0	2.0	19.0
Fisheries management	3.3	4.0	4.0	1.0	3.0	2.0	19.3
Removal of predators/invasive species	2.0	4.0	3.0	1.0	4.0	1.0	18.0
Reef community & structural restoration	3.0	3.5	4.0	1.0	4.0	1.0	19.5
Managed relocation & assisted gene flow	3.0	3.0	3.0	2.0	3.0	1.0	16.0
Assisted adaptation and evolution	3.0	2.0	3.0	2.0	2.0	1.0	14.0
Assisted fertilization and recruitment	3.0	3.0	3.0	2.0	3.0	1.0	16.0
Surface albedo enhancement	2.0	2.0	1.0	4.0	2.0	1.0	9.0
Cloud brightening	3.0	2.0	2.0	4.0	1.3	2.0	10.3
Aerosol-based SRM	2.0	1.0	2.0	4.0	1.0	2.5	8.0
Space-based SRM	2.5	1.0	2.0	4.5	1.0	3.0	8.0
Artificial upwelling	2.0	2.0	1.0	3.5	2.0	1.0	9.5

\*Score was calculated as the sum of Effectiveness, Readiness, Cobenefits, LACK of Disbenefits (i.e., the inverse of Disbenefits), and Acceptability.

**Table S3.2.** Number of scores available for each criterion and measure. Values of ‘0’ are excluded.



<b>Measure</b>	<b>Effectiveness</b>	<b>Readiness</b>	<b>Cobenefits</b>	<b>Disbenefits</b>	<b>Acceptability</b>	<b>Scale</b>
Addressing causes of climate change	20	18	18	19	19	18
Improved governance and regulation	21	20	21	21	21	21
Marine protected areas	21	21	21	21	21	21
Pollution reduction	21	21	21	21	21	21
Coastal zone management	21	21	21	21	21	21
Fisheries management	20	20	20	20	20	20
Removal of predators/invasive species	19	20	20	19	20	20
Reef community & structural restoration	20	20	20	19	20	20
Managed relocation & assisted gene flow	19	19	20	19	20	20
Assisted adaptation and evolution	19	19	20	19	20	20
Assisted fertilization and recruitment	20	20	20	20	20	20
Surface albedo enhancement	19	17	18	18	19	18
Cloud brightening	19	17	18	18	19	18
Aerosol-based SRM	19	17	18	18	19	17
Space-based SRM	19	17	18	18	19	17
Artificial upwelling	19	17	18	18	19	18