



HAL
open science

On the quenched functional CLT in random sceneries

Guy Cohen, Jean-Pierre Conze

► **To cite this version:**

Guy Cohen, Jean-Pierre Conze. On the quenched functional CLT in random sceneries. *Studia Mathematica*, 2023, 269 (3), pp.261-303. 10.4064/sm210421-13-10 . hal-03205934v2

HAL Id: hal-03205934

<https://hal.science/hal-03205934v2>

Submitted on 15 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON THE QUENCHED FUNCTIONAL CLT IN RANDOM SCENERIES

GUY COHEN AND JEAN-PIERRE CONZE

ABSTRACT. We prove a quenched functional central limit theorem (quenched FCLT) for the sums of a random field (r.f.) along a \mathbb{Z}^d -random walk in different frameworks: probabilistic (when the r.f. is i.i.d. or a moving average of i.i.d. random variables) and algebraic (when the r.f. is generated by commuting automorphisms of a torus or by commuting hyperbolic flows on homogeneous spaces).

CONTENTS

Introduction	2
1. Summation along a r.w. and variance	3
1.1. Random walks and sums along random walks	3
1.2. Variance for quenched processes	4
2. Random walk in random scenery	13
3. Cumulants and CLT	17
3.1. Moments and cumulants	17
3.2. A sufficient condition for the CLT	19
4. Moving averages of i.i.d. random variables	21
5. Tightness and 4th-moment	23
6. Automorphisms of a torus	26
7. Exponential mixing of all orders	29
7.1. FCLT and exponential mixing of all orders	29
7.2. Translations on homogeneous spaces	32
References	34

Date: September 29, 2022.

2010 Mathematics Subject Classification. Primary: 60F05, 28D05, 22D40, 60G50; Secondary: 47B15, 37A25, 37A30.

Key words and phrases. quenched functional central limit theorem, \mathbb{Z}^d -action, random walk in random scenery, self-intersections of a r.w., toral automorphisms, exponential mixing, flows on homogeneous spaces, S -unit, cumulant.

Introduction

Let G be a group acting on a probability space (E, \mathcal{B}, μ) by a measure preserving action $(g, x) \in G \times E \rightarrow T^g x \in E$. A random walk (Z_n) defined on a probability space (Ω, \mathbb{P}) with values in the group G induces a random walk on E . For $f \in L^2(E, \mathcal{B}, \mu)$, we can consider the sums along the random walk: $\sum_{k=0}^{n-1} f(T^{Z_k} x)$.

This general framework leads in practice to different situations and methods in the proof of a central limit theorem (CLT) and a functional central limit theorem (FCLT) along the paths of the random walk. In particular the proof of the tightness for the FCLT requires specific tools which it seems interesting to present in examples.

A first situation is that of a random walk in random sceneries (cf. [18], [5]). For $d \geq 1$, let $X = (X_\ell)_{\ell \in \mathbb{Z}^d}$ be a strictly \mathbb{Z}^d -stationary real random field (r.f.). One can assume that the r.v.s X_ℓ are defined on a probability space (E, \mathcal{B}, μ) on which commuting measure preserving maps T_1, \dots, T_d act in such a way¹ that $X_\ell = T^\ell X_0$.

Conversely, given commuting measure preserving invertible maps T_1, \dots, T_d and a measurable f on a probability space (E, \mathcal{B}, μ) , $(T^\ell f)_{\ell \in \mathbb{Z}^d}$ is a strictly \mathbb{Z}^d -stationary random field. If (Z_n) is a random walk in the group \mathbb{Z}^d , then the sums along Z_n read $\sum_{k=0}^{n-1} X_{Z_k} = \sum_{k=0}^{n-1} T^{Z_k} X_0$, or $\sum_{k=0}^{n-1} T^{Z_k} f$. When (X_ℓ) is a d -dimensional random field of i.i.d. random variables, we obtain the classical random walk in random sceneries.

Another kind of examples in the algebraic case can be obtained as follows. Suppose that $G = \text{SL}(\rho, \mathbb{Z})$ and that (E, \mathcal{B}, μ) is the torus \mathbb{T}^ρ , $\rho \geq 2$, endowed with the Borel σ -algebra and the Lebesgue measure. The map $x \rightarrow Ax$, where A is a matrix in $\text{SL}(\rho, \mathbb{Z})$, defines an automorphism of \mathbb{T}^ρ which preserves μ . When a spectral gap property is available for the transition operator associated to the random walk on G , the previous sums for f in a convenient class of observables satisfy a CLT (cf. [1], [12]) for \mathbb{P} -a.e. $\omega \in \Omega$.

In the commutative case there is no spectral gap and we use a method based on cumulants. A first example considered here is the action of commuting matrices in $\text{SL}(\rho, \mathbb{Z})$ acting on \mathbb{T}^ρ , for which we prove for \mathbb{P} -a.e. ω a functional CLT, extending previous results in [8]. A second algebraic example comes from commuting flows on homogeneous spaces. Based on the exponential mixing of all orders proved in [3], a CLT has been shown in [4] for ergodic sums on Følner sets when the observables are smooth. Likewise we prove here a CLT and its functional version for the sums along a random walk.

The result, a functional CLT for the models described above, is presented for a general aperiodic random walk in dimension $d > 1$ with a moment of order 2, but the detailed proofs are given in the case of a centered 2-dimensional r.w. The proofs can be adapted easily to the case of transient random walks. We say also some words in the i.i.d. case, when the usual random walk is replaced by a plane Lorentz process generated by a periodic billiard with dispersive obstacles (cf. [24], [25]).

Beyond the CLT, tightness is a main step in the proof of a FCLT. To show it we use the method based on the maximal inequality for associated r.v.s due to Newman and Wright [23]

¹Underlined letters represent elements of \mathbb{Z}^d or \mathbb{T}^d . We write $\underline{\ell}$ for (ℓ_1, \dots, ℓ_d) and $T^{\underline{\ell}}$ for $T_1^{\ell_1} \dots T_d^{\ell_d}$. The euclidean norm of $\underline{\ell} \in \mathbb{Z}^d$ is denoted by $|\underline{\ell}|$ or $\|\underline{\ell}\|$.

or, in the algebraic case, the method based on norm estimates for the maximum of partial sums (cf. Billingsley [2], Móricz [22]). A difficulty which occurs is that the estimates available for the random walk involve constants depending on the trajectory.

The content of the paper is the following. Section 1 contains results on the variance of sums along a random walk. The independent case is presented in Section 2. Some facts on cumulants are recalled in Section 3, then applied to moving averages in Section 4 and to the algebraic models in Sections 6 and 7. For the tightness in the latter cases, we use the method of maximum of partial sums in an adapted version presented in Section 5.

This paper extends a previous preprint [10]. We have added the FCLT along a random walk for flows on homogeneous spaces, using the recent results in [3] and [4] on the multiple mixing and on the CLT for group actions which are exponentially mixing of all orders. We have also added some remarks about the non nullity of the variance, in particular the observation that there is no degeneracy for the sums along a transient random walk.

Acknowledgements. This research started during visits of the first author to the IRMAR at the University of Rennes 1 and of the second author to the Center for Advanced Studies in Mathematics at Ben Gurion University. The authors are grateful to their hosts for their support. They thank B. Bekka, Y. Guivarc'h and M. Lin for helpful discussions.

1. Summation along a r.w. and variance

1.1. Random walks and sums along random walks.

First we recall some definitions and results about the random walks on \mathbb{Z}^d (see [27], details on the results recalled here can also be found in [8]).

Let $(\zeta_i)_{i \geq 0}$ be a sequence of i.i.d. random vectors on a probability space (Ω, \mathbb{P}) with values in \mathbb{Z}^d and common probability distribution ν . The associated *random walk* (r.w.) $Z = (Z_n)$ in \mathbb{Z}^d starting from $\underline{0}$ is defined by $Z_0 := \underline{0}$, $Z_n := \zeta_0 + \dots + \zeta_{n-1}$, $n \geq 1$.

The r.v.s ζ_i can be viewed as the coordinate maps on (Ω, \mathbb{P}) obtained as $(\mathbb{Z}^d)^{\mathbb{Z}}$ equipped with the product measure $\nu^{\otimes \mathbb{Z}}$ and with the shift θ acting on the coordinates. We have $\zeta_i = \zeta_0 \circ \theta^i$ and the cocycle relation $Z_{n+n'} = Z_n + Z_{n'} \circ \theta^n$, $\forall n, n' \geq 0$.

Let denote by $\mathcal{S} := \{\underline{\ell} \in \mathbb{Z}^d : \mathbb{P}(\zeta_0 = \underline{\ell}) > 0\}$ the support of ν and by L the sub-lattice of \mathbb{Z}^d generated by \mathcal{S} . Without loss of generality, we can assume that Z is reduced which means that L is cofinite in \mathbb{Z}^d . Therefore the vector space generated by L is \mathbb{R}^d and d is the ‘genuine’ dimension of the random walk Z .

For simplicity, we will assume that $L = \mathbb{Z}^d$ (then the random walk Z is said to be *aperiodic*). Observe that one can replace a reduced r.w. Z by an aperiodic one, again without loss of generality.

Let D be the sub-lattice of \mathbb{Z}^d generated by $\{\underline{\ell} - \underline{\ell}', \underline{\ell}, \underline{\ell}' \in \mathcal{S}\}$. We denote by Γ_1 the annihilator in \mathbb{T}^d of D , that is the closed subgroup of \mathbb{T}^d defined by $\{\underline{t} \in \mathbb{T}^d : e^{2\pi i \langle \underline{r}, \underline{t} \rangle} = 1, \forall \underline{r} \in D\}$ and by $d\gamma_1$ the Haar probability measure of the group Γ_1 . The r.w. is said to be *strictly (or strongly) aperiodic*, if $D = \mathbb{Z}^d$. For example, the r.w. on \mathbb{Z} defined by $\mathbb{P}(\zeta_0 = \pm 1) = \frac{1}{2}$ is aperiodic, but not strictly aperiodic.

Sums along random walks

Given a strictly \mathbb{Z}^d -stationary random field $X = (X_{\underline{\ell}})_{\underline{\ell} \in \mathbb{Z}^d}$, where the real random variables $X_{\underline{\ell}}$ are defined on a probability space (E, \mathcal{B}, μ) , the process of ‘ergodic sums’ along the random walk (Z_n) is

$$(1) \quad S_n^{\omega, X}(x) = S_n^{\omega}(x) := \sum_{k=0}^{n-1} X_{Z_k(\omega)}(x), \quad n \geq 1, \quad \omega \in \Omega.$$

If the random field is represented as $X_{\underline{\ell}} = T^{\underline{\ell}} f$, where T_1, \dots, T_d are commuting measure preserving maps and $f \in L^2(E, \mathcal{B}, \mu)$, the sums read:

$$(2) \quad S_n^{\omega} f = \sum_{k=0}^{n-1} T^{Z_k(\omega)} f = \sum_{\underline{\ell} \in \mathbb{Z}^d} w_n(\omega, \underline{\ell}) T^{\underline{\ell}} f,$$

where $w_n(\omega, \underline{\ell})$ (denoted also by $w_n^{\omega}(\underline{\ell})$) is the local time of the random walk at time $n \geq 1$:

$$(3) \quad w_n(\omega, \underline{\ell}) = \#\{k < n : Z_k(\omega) = \underline{\ell}\} = \sum_{k=0}^{n-1} 1_{Z_k(\omega) = \underline{\ell}}.$$

Summing along the random walk amounts to take the ergodic sums for the skew product $(\omega, x) \rightarrow T_{\zeta_0}(\omega, x) = (\theta\omega, T^{\zeta_0(\omega)}x)$ on $\Omega \times E$. Putting $F_f(\omega, x) = F(\omega, x) = f(x)$ for an observable f on E , we get that the ergodic sums of F for T_{ζ_0} read:

$$(4) \quad S_n F(\omega, x) = \sum_{k=0}^{n-1} F(T_{\zeta_0}^k(\omega, x)) = \sum_{k=0}^{n-1} f(T^{Z_k(\omega)}x) = (S_n^{\omega} f)(x).$$

A limit theorem in distribution for the sums $S_n^{\omega} f$ (with respect to the measure μ on E) obtained for \mathbb{P} -a.e. ω is sometimes called *quenched*. We will use this terminology². If the random variables $S_n F(\omega, x)$ are viewed as defined on $\Omega \times E$ endowed with the probability $\mathbb{P} \times \mu$, a limit theorem under $\mathbb{P} \times \mu$ for these sums is called *annealed*.

1.2. Variance for quenched processes.

Let f be a function in $L^2(E, \mathcal{B}, \mu)$ with real values. Everywhere we assume (or prove) the absolute summability of the series of decorrelations

$$(5) \quad \sum_{\underline{\ell} \in \mathbb{Z}^d} \left| \int_X T^{\underline{\ell}} f f \, d\mu \right| < \infty,$$

which implies existence and continuity of the spectral density, the even function given by

$$(6) \quad \varphi_f(\underline{t}) = \sum_{\underline{\ell} \in \mathbb{Z}^d} \langle T^{\underline{\ell}} f f \rangle e^{2\pi i \langle \underline{\ell}, \underline{t} \rangle}.$$

The computation of the variance $\int_E \left| \sum_{k=0}^{n-1} T^{Z_k(\omega)} f \right|^2 d\mu$, is related to the number of self-intersections of the random walk at time $n \geq 1$:

$$(7) \quad V_n(\omega) := \#\{0 \leq u, v < n : Z_u(\omega) = Z_v(\omega)\} = \sum_{\underline{\ell} \in \mathbb{Z}^d} w_n(\omega, \underline{\ell})^2 = \int_{\mathbb{T}^d} \left| \sum_{\underline{\ell} \in \mathbb{Z}^d} w_n(\underline{\ell}) e^{2\pi i \langle \underline{\ell}, \underline{t} \rangle} \right|^2 d\underline{t}.$$

²We follow here the terminology of [1] used in several papers. The term ‘quenched’ is also used in the random scenery when a limit theorem is shown for the distribution with respect to ω , conditionally to the scenery X .

Let us consider the kernels (which are even functions)

$$(8) \quad K(w_n^\omega)(\underline{t}) = \left| \sum_{k=0}^{n-1} e^{2\pi i \langle Z_k(\omega), \underline{t} \rangle} \right|^2 = \left| \sum_{\underline{\ell} \in \mathbb{Z}^d} w_n(\omega, \underline{\ell}) e^{2\pi i \langle \underline{\ell}, \underline{t} \rangle} \right|^2, \quad \tilde{K}(w_n^\omega)(\underline{t}) = V_n(\omega)^{-1} K(w_n^\omega)(\underline{t}).$$

We say that the summation along the r.w. Z is ξ -regular, where ξ is a probability measure on \mathbb{T}^d , if (for \mathbb{P} -a.e. ω) the normalised kernel $(\tilde{K}(w_n^\omega))_{n \geq 1}$ converges weakly to ξ when n tends to infinity, i.e., $\lim_{n \rightarrow \infty} \int_{\mathbb{T}^d} \tilde{K}(w_n^\omega) \varphi \, d\underline{t} = \xi(\varphi)$ for every continuous function φ on \mathbb{T}^d .

This property is equivalent to (for \mathbb{P} -a.e. ω):

$$(9) \quad \lim_{n \rightarrow \infty} \int_{\mathbb{T}^d} \tilde{K}(w_n^\omega)(\underline{t}) e^{-2\pi i \langle \underline{p}, \underline{t} \rangle} \, d\underline{t} = \lim_{n \rightarrow \infty} \int_{\mathbb{T}^d} \tilde{K}(w_n^\omega)(\underline{t}) \cos(2\pi \langle \underline{p}, \underline{t} \rangle) \, d\underline{t} = \hat{\xi}(\underline{p}), \quad \forall \underline{p} \in \mathbb{Z}^d.$$

Another equivalent formulation is

$$(10) \quad \lim_{n \rightarrow \infty} \frac{V_n(\omega, \underline{p})}{V_n(\omega)} = \hat{\xi}(\underline{p}), \quad \forall \underline{p} \in \mathbb{Z}^d, \text{ for a.e. } \omega,$$

with $V_n(\omega, \underline{p}) := \#\{0 \leq u, v < n : Z_u(\omega) - Z_v(\omega) = \underline{p}\}, \underline{p} \in \mathbb{Z}^d.$

Using the spectral theorem, if f satisfies (5), ξ -regularity implies that the (asymptotic) normalised variance is, for a.e. ω ,

$$(11) \quad \sigma^2(f) := \lim_n \frac{\left\| \sum_{\underline{\ell} \in \mathbb{Z}^d} w_n(\underline{\ell}, \omega) T^{\underline{\ell}} f \right\|_2^2}{\sum_{\underline{\ell} \in \mathbb{Z}^d} |w_n(\underline{\ell}, \omega)|^2} = \lim_n \int_{\mathbb{T}^d} \tilde{K}(w_n^\omega)(\underline{t}) \varphi_f(\underline{t}) \, d\underline{t} = \xi(\varphi_f).$$

It can be shown that the summation along any random walk in \mathbb{Z}^d is ξ -regular for some measure ξ (cf. [8]).

After some reminders on recurrence/transience, we summarize below the results on the asymptotic variance.

1.2.1. Recurrence/transience.

(cf. [6], [7], [27])

Recall that a r.w. $Z = (Z_n)$ is recurrent if $\sum_{n=1}^{\infty} \mathbb{P}(Z_n = \underline{0}) = +\infty$ and otherwise transient.

Recurrence occurs if and only if $\mathbb{P}(Z_n = \underline{0} \text{ infinitely often}) = 1$, and transience if and only if $\mathbb{P}(Z_n = \underline{0} \text{ infinitely often}) = 0$.

The random walk is said to have a moment of order 2 if $\sum_{\underline{\ell} \in \mathbb{Z}^d} \mathbb{P}(\zeta_0 = \underline{\ell}) \|\underline{\ell}\|^2 < \infty$.

Assume that Z is reduced in dimension d with a moment of order 2 (for $d = 1$, a moment of order 1 suffices). Then for $d=1, 2$, Z is recurrent if and only if it is centered. For $d \geq 3$, it is always transient.

We denote by $\Psi(\underline{t}) = \mathbb{E}[e^{2\pi i \langle \zeta_0, \underline{t} \rangle}]$, $\underline{t} \in \mathbb{T}^d$, the characteristic function of the r.w.

Observe that $\Psi(\underline{t}) \neq 1$ for $\underline{t} \neq \underline{0}$ in \mathbb{T}^d , since Z is aperiodic. We put

$$(12) \quad \Phi(\underline{t}) = \frac{1 - |\Psi(\underline{t})|^2}{|1 - \Psi(\underline{t})|^2} \text{ for } \underline{t} \neq \underline{0}, = 0 \text{ for } \underline{t} = \underline{0}.$$

Remark 1.1. Φ is nonnegative and $\Phi(\underline{t}) = 0$ only on Γ_1 . Hence, using again aperiodicity, it is positive for a.e. \underline{t} , except when the r.w. is ‘deterministic’ (i.e., if $\mathbb{P}(\zeta_0 = \underline{\ell}) = 1$ for some $\underline{\ell} \in \mathbb{Z}^d$, so that $|\Psi(\underline{t})| \equiv 1$ in this case).

A r.w. of genuine dimension d which is aperiodic is transient or recurrent depending on whether $\Re(\frac{1}{1-\Psi})$ is integrable or not on the d -dimensional unit cube ([27]).

Transient case

In the transient case, one can show:

Theorem 1.2. ([27]) *Let $Z = (Z_n)$ be a transient aperiodic random walk in \mathbb{Z}^d .*

a) *The function Φ is integrable on \mathbb{T}^d and, with a nonnegative constant K , we have*

$$I(\underline{\ell}) := 1_{\underline{\ell}=0} + \sum_{k=1}^{\infty} [\mathbb{P}(Z_k = \underline{\ell}) + \mathbb{P}(Z_k = -\underline{\ell})] = \int_{\mathbb{T}^d} \cos(2\pi\langle \underline{\ell}, \underline{t} \rangle) \Phi(\underline{t}) d\underline{t} + K, \forall \underline{\ell} \in \mathbb{Z}^d.$$

b₁) *Suppose $d = 1$. If $m_1(Z) = +\infty$, then $K = 0$. If $m_1(Z) < \infty$, then Z is non centered (because it is transient) and $K = |\sum_{\ell \in \mathbb{Z}} \mathbb{P}(\zeta_0 = \ell) \ell|^{-1}$.*

b₂) *If $d > 1$, then $K = 0$.*

c) *Denoting by $d\xi_0(\underline{t})$ the measure $\Phi(\underline{t})d\underline{t} + K\delta_0(\underline{t})$, we have, for a.e. ω ,*

$$(13) \quad \begin{aligned} & \int \frac{1}{n} K_n^\omega(\underline{t}) \cos(2\pi\langle \underline{\ell}, \underline{t} \rangle) d\underline{t} = 1_{\underline{\ell}=0} + \frac{1}{n} \sum_{k=1}^{n-1} \sum_{j=0}^{n-k-1} [1_{Z_k(\theta^j \omega) = \underline{\ell}} + 1_{Z_k(\theta^j \omega) = -\underline{\ell}}] \\ & \xrightarrow{n \rightarrow \infty} I(\underline{\ell}) = \int \cos(2\pi\langle \underline{\ell}, \underline{t} \rangle) d\xi_0(\underline{t}). \end{aligned}$$

It follows that the summation along a transient r.w. behaves for the normalisation like the iteration of a single transformation, is ξ_0 -regular (up to a constant factor) and that

$$(14) \quad \lim_n \frac{1}{n} \left\| \sum_{k=0}^{n-1} T^{Z_k(\omega)} f \right\|_2^2 = \int \Phi(\underline{t}) \varphi_f(\underline{t}) d\underline{t} + K \varphi_f(\underline{0}).$$

From (13), (14) and the expression of φ_f , $\varphi_f(\underline{t}) = \sum_{\underline{\ell} \in \mathbb{Z}^d} \langle T^{\underline{\ell}} f, f \rangle \cos(2\pi\langle \underline{\ell}, \underline{t} \rangle)$, we deduce:

$$\begin{aligned} & \int \Phi(\underline{t}) \varphi_f(\underline{t}) d\underline{t} + K \varphi_f(\underline{0}) = \int \varphi_f(\underline{t}) d\xi_0(\underline{t}) = \sum_{\underline{\ell} \in \mathbb{Z}^d} \langle T^{\underline{\ell}} f, f \rangle \int \cos(2\pi\langle \underline{\ell}, \underline{t} \rangle) d\xi_0(\underline{t}) = \\ & \sum_{\underline{\ell} \in \mathbb{Z}^d} I(\underline{\ell}) \langle T^{\underline{\ell}} f, f \rangle = \|f\|_2^2 + 2 \sum_{k \geq 1} \left(\sum_{\underline{\ell} \in \mathbb{Z}^d} \mathbb{P}(Z_k = \underline{\ell}) \langle T^{\underline{\ell}} f, f \rangle \right). \end{aligned}$$

Remark 1.3. (about the variance in the non deterministic transient case) Let f be in $L^2(E, \mu)$ with real values and satisfying (5). Suppose that f is not a.e. equal to 0.

a) By Remark 1.1, Φ is positive a.e. Therefore $\int \Phi \varphi_f d\underline{t} = 0$ implies $\varphi_f(\underline{t}) = 0$ a.e. which is not possible since $\int \varphi_f d\underline{t} = \|f\|_2$.

Therefore the (quenched) asymptotic variance, which is given by (14), is > 0 .

b) Let $F_f(\omega, x) = F(\omega, x) = f(x)$. For the map T_{ζ_0} acting on the product space $\Omega \times E$ endowed with the product measure $\mathbb{P} \times \mu$, we have

$$\int T_{\zeta_0}^n F F d\mu d\mathbb{P} = \sum_{\underline{\ell}} (\mathbb{E} 1_{Z_n = \underline{\ell}}) \langle T^{\underline{\ell}} f f \rangle = \sum_{\underline{\ell} \in \mathbb{Z}^d} \mathbb{P}(Z_n = \underline{\ell}) \langle T^{\underline{\ell}} f, f \rangle, \quad n \geq 0.$$

In the transient case it holds, for every $\underline{\ell}$, $\sum_{k \geq 1} \mathbb{P}(Z_k = \underline{\ell}) \leq \sum_{k \geq 1} \mathbb{P}(Z_k = \underline{0}) < +\infty$. Therefore the density of the spectral measure for F_f and the map T_{ζ_0} is

$$\|f\|_2^2 + 2 \sum_{k \geq 1} \left(\sum_{\underline{\ell} \in \mathbb{Z}^d} \mathbb{P}(Z_k = \underline{\ell}) \langle T^{\underline{\ell}} f, f \rangle \right) \cos 2\pi k t.$$

The asymptotic variance $\lim_n \frac{1}{n} \left\| \sum_{k=0}^{n-1} T_{\zeta_0}^k F_f \right\|_2^2$ for the annealed model is the same as for the quenched model and is equal to $\|f\|_2^2 + 2 \sum_{k \geq 1} \left(\sum_{\underline{\ell} \in \mathbb{Z}^d} \mathbb{P}(Z_k = \underline{\ell}) \langle T^{\underline{\ell}} f, f \rangle \right)$.

It follows that the function F_f on $\Omega \times E$ (which depends only on the second coordinate), with f as above and non a.e. null, is never a coboundary in $L^2(\mathbb{P} \times \mu)$ for T_{ζ_0} , because the asymptotic variance is non null. Observe also that F_f is not even a measurable coboundary, at least when the CLT holds, which is the case of the situations that we are going to consider here. This follows from the fact that, for a single measure preserving transformation, if an observable is a coboundary in the space of measurable functions, then the limiting distribution of the ergodic sums after normalisation by any sequence tending to infinity is the Dirac mass at 0, which is excluded here.

Recurrent case

Let us consider now the case $d = 2$ and a centered random walk Z with a moment of order 2. By the local limit theorem (LLT), Z is recurrent.

A non standard normalization occurs in the CLT for sums along Z_n as recalled below. There are C_0, C finite positive constants³ such that (cf. [5] Lemma 2.6, [21] Proposition 1.4 for (15) and (16), [8] Theorem 4.13 for (17)):

$$(15) \quad \mathbb{E}(V_n) \sim C_0 n \log n, \quad \text{Var}(V_n) \leq C n^2,$$

$$(16) \quad \varphi_n(\omega) := \frac{V_n(\omega)}{C_0 n \log n} \rightarrow 1, \quad \text{for a.e. } \omega,$$

$$(17) \quad \varphi_n(\omega, \underline{p}) := \frac{V_n(\omega, \underline{p})}{C_0 n \log n} \rightarrow 1, \quad \forall \underline{p} \in \mathbb{Z}^d, \quad \text{for a.e. } \omega.$$

Therefore the summation along the r.w. Z is δ_0 -regular: the normalised kernel satisfies $\lim_n \int \tilde{K}(w_n^\omega)(\underline{t}) e^{-2\pi i \langle \underline{p}, \underline{t} \rangle} d\underline{t} = 1, \quad \forall \underline{p} \in \mathbb{Z}^d$ and the asymptotic variance is

$$(18) \quad \sigma^2(f) = \lim_n (C_0 n \log n)^{-1} \left\| \sum_{k=0}^n T^{Z_k(\omega)} f \right\|_2^2 = \sum_{\underline{k} \in \mathbb{Z}^d} \langle T^{\underline{k}} f f \rangle = \varphi_f(\underline{0}).$$

³If the r.w. is strongly aperiodic, $C_0 = (\pi \sqrt{\det \Sigma})^{-1}$, where Σ is the covariance matrix of the increments of the r.w.

The results presented below are valid for the cases covered above, excludes only the one-dimensional recurrent case. We stress that, in the recurrent 2-dimensional case, the variance can be degenerate, while this does not occur in the transient case unless $f = 0$.

1.2.2. Number of self-intersections of a 2-dimensional centered r.w.

In this subsection, we study more precisely the case $d = 2$ and a centered random walk (Z_n) with a moment of order 2.

If I, J are intervals, the quantity $V(\omega, I, J, \underline{p}) :=$

$$(19) \int \left(\sum_{u \in I} e^{2\pi i \langle Z_u(\omega), \underline{t} \rangle} \right) \left(\sum_{v \in J} e^{-2\pi i \langle Z_v(\omega), \underline{t} \rangle} \right) e^{-2\pi i \langle \underline{p}, \underline{t} \rangle} d\underline{t} = \#\{(u, v) \in I \times J : Z_u(\omega) - Z_v(\omega) = \underline{p}\}$$

is non negative and increases when I or J increases for the inclusion order.

We write simply $V(\omega, I, \underline{p})$ if $I = J$, $V(\omega, I)$ for $V(\omega, I, \underline{0})$, $V_n(\omega)$ and $V_n(\omega, \underline{p})$ as above for $V(\omega, [0, n[)$ and $V(\omega, [0, n[, \underline{p})$.

Observe that $V(\omega, J) = \sum_{\underline{\ell} \in \mathbb{Z}^2} w(\omega, J, \underline{\ell})^2$, where $w(\omega, J, \underline{\ell}) = \sum_{i \in J} 1_{Z_i(\omega) = \underline{\ell}}$. Notice also that $V(\omega, [b, b+k[) = V(\theta^b \omega, [0, k[) = V_k(\theta^b \omega)$, for $b \geq 0, k \geq 1$.

Let A, B be in $[0, 1]$. We have:⁴

$$\begin{aligned} V(\omega, [nA, nB], \underline{p}) &= \int \left(\sum_{u \in [nA, nB]} e^{2\pi i \langle Z_u(\omega), \underline{t} \rangle} \right) \left(\sum_{v \in [nA, nB]} e^{-2\pi i \langle Z_v(\omega), \underline{t} \rangle} \right) e^{-2\pi i \langle \underline{p}, \underline{t} \rangle} d\underline{t} \\ &= \#\{u, v \in [0, n(B-A)] : \sum_{i=0}^{u-1} \zeta_0(\theta^{i+nA} \omega) - \sum_{i=0}^{v-1} \zeta_0(\theta^{i+nA} \omega) = \underline{p}\} = V(\theta^{nA} \omega, [0, n(B-A)], \underline{p}). \end{aligned}$$

By (16) and (17) there a set Ω_0 of full probability such that

$$(20) \quad V_n(\omega) \leq K(\omega) n \log n, \quad \forall n \geq 2, \quad \text{where the function } K \geq 0 \text{ is finite on } \Omega_0,$$

$$(21) \quad \text{for any fixed } A \in]0, 1[, \quad V(\omega, [1, nA], \underline{p}) \sim C_0 n A \log n, \quad \text{for } \omega \in \Omega_0.$$

By [5, Lemma 2.5] we have

$$(22) \quad \sup_{\underline{\ell} \in \mathbb{Z}^2} w_n(\omega, \underline{\ell}) = o(n^\varepsilon), \quad \text{for a.e. } \omega, \quad \text{for every } \varepsilon > 0.$$

For a simple r.w. on \mathbb{Z}^2 , Erdős and Taylor [15] have shown: $\limsup_n \sup_{\underline{\ell} \in \mathbb{Z}^2} \frac{w_n(\omega, \underline{\ell})}{(\log n)^2} \leq \frac{1}{\pi}$.

The result has been extended by Dembo, Peres, Rosen and Zeitouni [14]. They proved for an aperiodic centered random walk on \mathbb{Z}^2 with moments of all orders:

$$\limsup_n \sup_{\underline{\ell} \in \mathbb{Z}^2} \frac{w_n(\omega, \underline{\ell})}{(\log n)^2} = \frac{1}{\pi}.$$

Remark 1.4. In Section 7 we will use the upper bound

$$(23) \quad \limsup_n \sup_{\underline{\ell} \in \mathbb{Z}^2} \frac{w_n(\omega, \underline{\ell})}{(\log n)^2} \leq \frac{1}{\pi}.$$

⁴For simplicity, in the formulas above and below, we write nA, nB instead of $[nA]$ or $[nA] + 1, [nB]$, θ^t instead of $\theta^{\lfloor t \rfloor}$. The equalities are satisfied up to the addition of quantities which are bounded independently from A, B, n .

As shown in the proof in [14], it suffices for the validity of (23) to suppose that the 2-dimensional r.w. is aperiodic. Moreover, the proof of the upper bound is based on the local limit theorem which uses only the existence of the moments of order 2.

We will need also to bound, for $\underline{\ell}_1, \underline{\ell}_2, \underline{\ell}_3 \in \mathbb{Z}^d$, $W_n(\omega, \underline{\ell}_1, \underline{\ell}_2, \underline{\ell}_3) :=$

$$(24) \quad \#\{1 \leq i_0, i_1, i_2, i_3 < n : Z_{i_1}(\omega) - Z_{i_0}(\omega) = \underline{\ell}_1, Z_{i_2}(\omega) - Z_{i_1}(\omega) = \underline{\ell}_2, Z_{i_3}(\omega) - Z_{i_2}(\omega) = \underline{\ell}_3\}.$$

Lemma 1.5. *There exists a positive integrable function C_3 such that*

$$(25) \quad W_n(\omega, \underline{\ell}_1, \underline{\ell}_2, \underline{\ell}_3) \leq C_3(\omega) n (\log n)^5, \quad \forall n \geq 1.$$

Proof. It suffices to bound the sum with strict inequality between indices

$$W'_n(\omega) = \sum_{1 \leq i_0 < i_1 < i_2 < i_3 \leq n} \mathbf{1}_{Z_{i_1} - Z_{i_0} = \underline{\ell}_1} \cdot \mathbf{1}_{Z_{i_2} - Z_{i_1} = \underline{\ell}_2 - \underline{\ell}_1} \cdot \mathbf{1}_{Z_{i_3} - Z_{i_2} = \underline{\ell}_3 - \underline{\ell}_2}.$$

Using independence and the local limit theorem for the random walk, we find the bound

$$\int W'_n(\omega) d\mathbb{P}(\omega) \leq C_1 \sum_{i_0, i_1, i_2, i_3 \in [1, n]} (i_1 i_2 i_3)^{-1} \leq C_2 n (\log n)^3.$$

Therefore $\sum_{p=1}^{\infty} \int 2^{-p} (\log(2^p))^{-5} W'_{2^p} d\mathbb{P} < \infty$. The function $C(\omega) := \sum_{p=1}^{\infty} 2^{-p} (\log(2^p))^{-5} W'_{2^p}$ is integrable and we have: $W'_{2^p}(\omega) \leq C(\omega) 2^p (\log(2^p))^5, \forall p \geq 1$.

Let p_n be such that: $2^{p_n-1} \leq n < 2^{p_n}$. Since W'_n is increasing with n , we obtain:

$$W'_n(\omega) \leq W'_{2^{p_n}}(\omega) \leq C(\omega) 2^{p_n} (\log(2^{p_n}))^5 \leq C(\omega) 2n (\log(2n))^5 \leq C'(\omega) n (\log n)^5. \quad \square$$

Variance for the finite dimensional distributions

The following lemma will be applied to the successive return times of a point ω into a set under the iteration of the shift θ .

Lemma 1.6. *Let $(y(j), j \geq 1)$ be a sequence with values in $\{0, 1\}$ such that $\lim_n \frac{1}{n} \sum_{j=1}^n y(j) = a > 0$. If (k_r) is the sequence of successive times such that $y(k_r) = 1$, then, for every $\delta > 0$, there is $n(\delta)$ such that, for $n \geq n(\delta)$, $k_{r+1} - k_r \leq \delta n$, for all $r \in [1, n]$.*

Proof. Since $r = \sum_{j=1}^{k_r} y(j)$, we have: $k_r/r = k_r / \sum_{j=1}^{k_r} y(j) \rightarrow a^{-1}$. Hence, for every $\delta > 0$, there is $n_1(\delta)$ such that $0 < k_{r+1} - k_r \leq \delta r$, for $r \geq n_1(\delta)$. Therefore, if $n \geq n_1(\delta)$, then $0 < k_{r+1} - k_r \leq \delta r \leq \delta n$, for $r \in [n_1(\delta), n]$.

If $n(\delta) \geq n_1(\delta)$ is such that $k_{r+1} - k_r \leq \delta n(\delta)$ for $r \leq n_1(\delta)$, we get the result of the lemma. \square

Lemma 1.7. *Let Λ be a measurable set in Ω of positive measure. Let $k_r = k_r(\omega)$ be the successive times such that $\theta^{k_r} \omega \in \Lambda$. For ω in a set Ω_Λ of full measure, for every positive small enough δ , there is $n(\delta)$ such that for $n \geq n(\delta)$*

- 1) $k_{r+1} - k_r \leq \delta n$, for all $r \in [1, n]$; moreover, $k_n \sim cn$, with $c = \mathbb{P}(\Lambda)^{-1}$, when $n \rightarrow \infty$;
- 2) there are integers $v < 2/\delta$ and $0 = \rho_1^{(n)} < \rho_2^{(n)} < \dots < \rho_v^{(n)} \leq n < \rho_{v+1}^{(n)}$, such that $\theta^{\rho_i^{(n)}} \omega \in \Lambda$ and $\frac{1}{2}\delta n \leq \rho_{i+1}^{(n)} - \rho_i^{(n)} \leq \frac{3}{2}\delta n$, for $i = 1, \dots, v$.

Proof. Since θ is ergodic on (Ω, \mathbb{P}) , Birkhoff ergodic theorem implies $\lim_n \frac{1}{n} \sum_0^{n-1} 1_\Lambda(\theta^k \omega) = \mathbb{P}(\Lambda) > 0$, for a.e. ω and $k_n/n \rightarrow \mathbb{P}(\Lambda)^{-1}$. Hence Lemma 1.6 implies 1). For 2), we select in the sequence (k_r) an increasing sequence of visit times to the set Λ satisfying the prescribed conditions by eliminating successive times which are at a distance $< \frac{1}{2}\delta n$. \square

Asymptotic orthogonality of the cross terms

Proposition 1.8. *For $0 < A < B < C < D < 1$, $\underline{p} \in \mathbb{Z}$, it holds a.e.*

$$(26) \quad \int \left[\left(\sum_{v=nA}^{nB} e^{2\pi i \langle Z_v(\omega), \underline{u} \rangle} \right) \left(\sum_{w=nC}^{nD} e^{-2\pi i \langle Z_w(\omega), \underline{u} \rangle} \right) e^{-2\pi i \langle \underline{p}, \underline{u} \rangle} \right] d\underline{u} = \varepsilon_n(\omega) n \log n, \quad \text{with } \varepsilon_n(\omega) \rightarrow 0.$$

The above integral is the non negative self-intersection quantity: $V(\omega, [nA, nB], [nC, nD], \underline{p})$. By (19), $V(\omega, I, J, \underline{p})$ increases when I or J increases. Hence, it suffices to show (26) for the intervals $[1, nA], [nA, n]$, for $0 < A < 1$. The proof below is based on (17) and (21).

Lemma 1.9. *There is a set $\hat{\Omega} \subset \Omega$ such that $\mathbb{P}(\hat{\Omega}) = 1$ and for all $\omega \in \hat{\Omega}$, the following holds:*

$$(27) \quad \lim_n \varphi_{nB}(\theta^{nA} \omega, \underline{p}) = \lim_n \frac{V(\omega, [nA, n], \underline{p})}{C_0 n B \log n} = 1, \quad \text{for } A \in]0, 1[, B = 1 - A;$$

$$(28) \quad V(\omega, [1, nA], [nA, n], \underline{p}) + V(\omega, [nA, n], [1, nA], \underline{p}) = \varepsilon_n(\omega) n \log n, \quad \text{with } \varepsilon_n(\omega) \rightarrow 0.$$

Proof. 1) *The set $\hat{\Omega}$.* For every $L \geq 1$ and $\delta > 0$, let $\Lambda(L, \delta) := \{\omega : \varphi_n(\omega, \underline{p}) - 1 \in [-\delta, \delta], \forall n \geq L\}$. We have $\lim_{L \uparrow \infty} \mathbb{P}(\Lambda(L, \delta)) = 1$. There is $L(\delta)$ such that $\mathbb{P}(\Lambda(L(\delta), \delta)) \geq \frac{1}{2}$.

Let (δ_j) be a sequence tending to 0. We apply Lemma 1.7 to $\Lambda(L(\delta_j), \delta_j)$ for each j . By taking the intersection of the sets of full measure $\Omega_{\Lambda(L(\delta_j), \delta_j)}$, we get a set Ω_1 of full measure. The set $\hat{\Omega}$ is the intersection of Ω_1 with the set Ω_0 (of full measure) for which the law of large numbers holds for $(V_n(\omega))$. Let $\omega \in \hat{\Omega}$.

2) *Proof of (27).* We have $V(\omega, [nA, n], \underline{p}) = V(\theta^{nA} \omega, [0, n(1-A)], \underline{p})$ and

$$(29) \quad \begin{aligned} & V(\omega, [1, n], \underline{p}) - V(\omega, [1, nA], \underline{p}) - V(\omega, [nA, n], \underline{p}) \\ &= V(\omega, [1, nA], [nA, n], \underline{p}) + V(\omega, [nA, n], [1, nA], \underline{p}) \geq 0. \end{aligned}$$

Claim: for an absolute constant C_1 depending on A and \underline{p} , for every δ , for n big enough,

$$(30) \quad \varphi_{nB}(\theta^{nA} \omega, \underline{p}) = \frac{V(\omega, [nA, n], \underline{p})}{C_0 n(1-A) \log n} \in [1 - C_1 \delta, 1 + C_1 \delta].$$

Let us show the claim. We put $B = 1 - A$ and take $\delta \in]0, B[$.

Upper bound: The law of large numbers for $V_n(\omega, \underline{p})$ implies, with $|\varepsilon_n|, |\varepsilon'_n| \leq \delta$ for n big enough,

$$C_0^{-1} V(\omega, [1, n], \underline{p}) = (1 + \varepsilon_n) n \log n, \quad C_0^{-1} V(\omega, [1, nA], \underline{p}) = (1 + \varepsilon'_n) nA \log n.$$

This implies by (29)

$$\frac{V(\omega, [nA, n], \underline{p})}{C_0 n B \log n} \leq \frac{(1 + \varepsilon_n) n \log n - (1 + \varepsilon'_n) nA \log n}{nB \log n} \leq 1 + \frac{|\varepsilon_n|}{B} + \frac{|\varepsilon'_n| A}{B} \leq 1 + \frac{1 + A}{B} \delta.$$

Lower bound: We apply Lemma 1.7 to $\Lambda(L(\delta), \delta)$. Let n_A, n'_A be two consecutive visit times $\leq n$ such that $n_A \leq nA < n'_A$. For n big enough, we have $0 < n'_A - n_A \leq \delta n$ and

$$n_A = nA(1 - \rho_n), \quad n'_A = nA(1 + \rho'_n), \quad \text{with } 0 \leq A\rho_n, A\rho'_n \leq \delta.$$

Since $\omega \in \hat{\Omega}$, we have for n big enough, with $|\delta'_n| \leq \delta$,

$$C_0^{-1} V(\omega, [n'_A, n], \underline{p}) \geq (1 - \delta'_n)(n - n'_A) \log(n - n'_A) = (1 - \delta'_n)(nB - nA\rho'_n) \log(nB - nA\rho'_n).$$

It follows, for δ (hence ρ'_n) small:

$$\begin{aligned} \frac{V(\omega, [n'_A, n], \underline{p})}{C_0(1 - \delta'_n)nB \log(nB)} &\geq \frac{(nB - nA\rho'_n) \log(nB - nA\rho'_n)}{nB \log(nB)} = \frac{(B - A\rho'_n) [\log(nB) + \log(1 - \frac{A}{B}\rho'_n)]}{B \log(nB)} \\ &\geq (1 - \frac{A}{B}\rho'_n) - 2(1 - \frac{A}{B}\rho'_n) \frac{\frac{A}{B}\rho'_n}{\log(nB)} \geq 1 - \frac{A}{B}\rho'_n - 2 \frac{\frac{A}{B}\rho'_n}{\log(nB)} \geq 1 - B^{-1}\delta(1 + \frac{2}{\log(nB)}). \end{aligned}$$

As $V(\omega, J, \underline{p})$ increases when the set J increases, we have by the choice of n_A and n'_A :

$$V(\omega, [n'_A, n], \underline{p}) \leq V(\omega, [nA, n], \underline{p}).$$

Therefore, for n such that $\log(nB) \geq 2$, we have

$$\frac{V(\omega, [nA, n], \underline{p})}{C_0 nB \log(nB)} \geq (1 - \delta) (1 - \frac{2}{B}\delta) \geq 1 - \delta(1 + \frac{2}{B}).$$

This shows the lower bound. Altogether with the upper bound, this proves the claim (30).

3) *Proof of (28).* Let $\delta > 0$. According to (29) and (30), for n big enough, we have with $|\varepsilon''_n| \leq C_1\delta$:

$$\begin{aligned} V(\omega, [1, nA], [nA, n], \underline{p}) + V(\omega, [nA, n], [1, nA], \underline{p}) &= V(\omega, [1, n], \underline{p}) - V(\omega, [1, nA], \underline{p}) - V(\omega, [nA, n], \underline{p}) \\ &= C_0[(1 + \varepsilon_n)n \log n - (1 + \varepsilon'_n)nA \log n - (1 + \varepsilon''_n)n(1 - A) \log n] \leq (2 + C_1) C_0 \delta n \log n. \quad \square \end{aligned}$$

Let a_1, \dots, a_s be real numbers and $0 = t_0 < t_1 < \dots < t_{s-1} < t_s = 1$ a subdivision of $[0, 1]$. For the asymptotic variance of $\sum_{j=0}^s a_j \sum_{nt_{j-1} \leq k < nt_j} T^{Z_k(\omega)} f$, which is used later, we need the following lemma.

Notation: With the convention of footnote 4, we will write $\sum_{k=nt_{j-1}}^{nt_j}$ instead of $\sum_{nt_{j-1} \leq k < nt_j}$.

Recall that f has a continuous spectral density φ_f .

Lemma 1.10. *For a.e. ω and for every partition (t_j) , we have*

$$(31) \quad (C_0 n \log n)^{-1} \left\| \sum_{j=1}^s a_j \sum_{k=nt_{j-1}}^{nt_j} T^{Z_k(\omega)} f \right\|_2^2 \rightarrow \varphi_f(\underline{0}) \sum_{j=1}^s a_j^2 (t_j - t_{j-1}).$$

Proof. 1) Recall that proving (31) amounts to prove

$$(C_0 n \log n)^{-1} \int \left| \sum_{j=1}^s a_j \sum_{k=nt_{j-1}}^{nt_j} e^{2\pi i \langle Z_k(\omega), \underline{u} \rangle} \right|^2 \varphi_f(\underline{u}) d\underline{u} \rightarrow \varphi_f(\underline{0}) \sum_{j=1}^s a_j^2 (t_j - t_{j-1}).$$

1) First suppose that φ_f is a trigonometric polynomial ρ , which allows to use (26) for a finite set of characters $e^{-2\pi i \langle \underline{p}, \underline{u} \rangle}$. Using (18) for the asymptotic variance starting from 0, we have

$(C_0 n \log n)^{-1} \left\| \sum_{k=0}^{\lfloor tn \rfloor} T^{Z_k(\omega)} f \right\|_2^2 \rightarrow t \rho(\underline{0})$, for $t \in]0, 1[$. By Lemma 1.9,

$$(C_0 n \log n)^{-1} \left\| \sum_{k=\lfloor sn \rfloor}^{\lfloor tn \rfloor} T^{Z_k(\omega)} f \right\|_2^2 \rightarrow (t-s) \rho(\underline{0}), \text{ for } 0 < s < t < 1.$$

Expanding the square and using that the cross terms are asymptotically negligible, we have

$$\begin{aligned} & (C_0 n \log n)^{-1} \int \left| \sum_{j=1}^s a_j \sum_{k=nt_{j-1}}^{nt_j} e^{2\pi i \langle Z_k(\omega), \underline{u} \rangle} \right|^2 \rho(\underline{u}) d\underline{u} \\ & \sim (C_0 n \log n)^{-1} \left(\sum_{j=1}^s a_j^2 \int \left| \sum_{k=nt_{j-1}}^{nt_j} e^{2\pi i \langle Z_k(\omega), \underline{u} \rangle} \right|^2 \rho(\underline{u}) d\underline{u} \right) \rightarrow \rho(\underline{0}) \sum_{j=1}^s a_j^2 (t_j - t_{j-1}). \end{aligned}$$

This shows (31) for trigonometric polynomials.

2) For a general continuous spectral density φ_f , for $\varepsilon > 0$, let ρ be a trigonometric polynomial, such that $\|\varphi_f - \rho\|_\infty < \varepsilon$. Remark that

$$\int \left| \sum_{j=1}^s a_j \sum_{k=nt_{j-1}}^{nt_j} e^{2\pi i \langle Z_k(\omega), \underline{u} \rangle} \right|^2 d\underline{u} \leq \sum_{j,j'=1}^s a_j a_{j'} V(\omega, [nt_{j-1}, nt_j], [nt_{j'-1}, nt_{j'}], \underline{0}) \leq \left(\sum_{j=1}^s |a_j| \right)^2 V_n(\omega).$$

Therefore we have:

$$\begin{aligned} & \left| (C_0 n \log n)^{-1} \int \left| \sum_{j=1}^s a_j \sum_{k=nt_{j-1}}^{nt_j} e^{2\pi i \langle Z_k(\omega), \underline{u} \rangle} \right|^2 \varphi_f(\underline{u}) d\underline{u} - \varphi_f(\underline{0}) \sum_{j=1}^s a_j^2 (t_j - t_{j-1}) \right| \\ & \leq \left| (C_0 n \log n)^{-1} \int \left| \sum_{j=1}^s a_j \sum_{k=nt_{j-1}}^{nt_j} e^{2\pi i \langle Z_k(\omega), \underline{u} \rangle} \right|^2 \rho(\underline{u}) d\underline{u} - \rho(\underline{0}) \sum_{j=1}^s a_j^2 (t_j - t_{j-1}) \right| \\ & + \varepsilon \left[(C_0 n \log n)^{-1} \int \left| \sum_{j=1}^s a_j \sum_{k=nt_{j-1}}^{nt_j} e^{2\pi i \langle Z_k(\omega), \underline{u} \rangle} \right|^2 d\underline{u} + \sum_{j=1}^s a_j^2 (t_j - t_{j-1}) \right]. \end{aligned}$$

By the remark, the above quantity inside [] is less than $(\sum_{j=1}^s |a_j|)^2 (C_0 n \log n)^{-1} V_n(\omega) + \sum_{j=1}^s a_j^2 (t_j - t_{j-1})$, which is bounded uniformly with respect to n by (20). Therefore we can conclude for a general continuous spectral density by step 1). \square

Remarks 1.11. 1) In Lemma 1.7, the dynamical system $(\Omega, \theta, \mathbb{P})$ can be replaced by any ergodic dynamical system.

2) If the spectral density is constant (i.e., when the X_k 's are pairwise orthogonal), (26) and (31) are a consequence of the law of large numbers for the number of self-intersections, that is $\frac{V_n(\omega)}{C_0 n \log n} \rightarrow 1$. The law of large numbers for $V_n(\omega, \underline{p})$, $\underline{p} \neq \underline{0}$, is not needed.

3) A result analogous to Proposition 1.8 is valid if the r.w. Z is transient: for $0 < A < B < C < D < 1$, $\underline{p} \in \mathbb{Z}$,

$$(32) \quad \int \left[\left(\sum_{v=nA}^{nB} e^{2\pi i \langle Z_v(\omega), \underline{u} \rangle} \right) \left(\sum_{w=nC}^{nD} e^{-2\pi i \langle Z_w(\omega), \underline{u} \rangle} \right) e^{-2\pi i \langle \underline{p}, \underline{u} \rangle} \right] d\underline{u} = \varepsilon_n(\omega) n, \text{ with } \varepsilon_n(\omega) \rightarrow 0.$$

4) The quenched FCLT shown in the different examples below is valid for a set of ω 's of \mathbb{P} -measure 1 given by the results of this Section 1. This set does not depend on the \mathbb{Z}^d -dynamical systems considered in the further sections. The joint distribution on $\Omega \times E$ is used only when the annealed model is mentioned when Bolthausen's result is recalled in Section 2.

1.2.3. Formulation of the quenched FCLT.

(For the reminders below, see Theorem 7.5, p. 84 in [2], as well as Theorems 15.1 and 15.5 in the first edition of this book.)

Let $(Y_n(t), t \in [0, 1])$ be a process on (E, μ) with values in the space $C[0, 1]$ of real valued continuous functions on $[0, 1]$ or in the space $D[0, 1]$ of right continuous real valued functions on $[0, 1]$ with left limits, endowed with the uniform norm.

Let $(W(t), t \in [0, 1])$ be the Wiener process on $[0, 1]$. To show a functional limit theorem (FCLT) for $(Y_n(t), t \in [0, 1])$, i.e., weak convergence to the Wiener process, it suffices to prove the two following properties (" \implies " denotes the convergence in distribution):

1) *Convergence of the finite dimensional distributions:*

$$\forall 0 = t_0 < t_1 < \dots < t_r = 1, (Y_n(t_1), \dots, Y_n(t_r)) \xrightarrow[n \rightarrow \infty]{} (W_{t_1}, \dots, W_{t_r}),$$

a property which follows (by the Cramér-Wold device) from

$$(33) \quad \sum_{j=1}^r a_j (Y_n(t_j) - Y_n(t_{j-1})) \implies \mathcal{N}(0, \sum_{j=1}^r a_j^2 (t_j - t_{j-1})), \forall (a_j)_{1 \leq j \leq r} \in \mathbb{R}.$$

2) *Tightness of the process:* The condition of tightness reads:

$$(34) \quad \forall \varepsilon > 0, \lim_{\delta \rightarrow 0^+} \limsup_n \mu(x \in E : \sup_{|t'-t| \leq \delta} |Y_n(x, t') - Y_n(x, t)| \geq \varepsilon) = 0.$$

Let (Z_n) be a random walk on (Ω, \mathbb{P}) with values in \mathbb{Z}^d , $d \geq 1$, and let $X = (X_\ell(x))_{\ell \in \mathbb{Z}^d} = (T^\ell f(x))_{\ell \in \mathbb{Z}^d}$ be a d -dimensional random field defined on a probability space (E, \mathcal{B}, μ) . We take sums along Z_n , introducing so another alea ω (which will be fixed in the quenched setting).

A quenched FCLT is satisfied by the sums along Z_n if, for \mathbb{P} -a.e. ω , the functional central limit theorem holds for the process $(S_{[nt]}^{\omega, X}(x))$ (cf. Notation (1)) after normalisation.

For $d = 2$, it means that, for \mathbb{P} -a.e. ω , the functional central limit theorem holds for the normalised process

$$(35) \quad (Y_n(\omega, x, t))_{t \in [0, 1]} := \left(\frac{S_{[nt]}^{\omega, X}(x)}{\sqrt{C_0 n \log n}} \right)_{t \in [0, 1]}.$$

2. Random walk in random scenery

We consider in this section $d = 2$ and the random walk in random scenery $S_n^{\omega, X}(x)$, that is the process (Y_n) defined by (35) when X is a 2-dimensional random field of i.i.d. real variables with $\mathbb{E}(X_0^2) = 1$ and mean 0 on a probability space (E, \mathcal{B}, μ) .

As the random field is i.i.d centered with $\mathbb{E}(X_0^2) = 1$, the corresponding spectral density is constant and identically = 1.

It was shown by E. Bolthausen [5] that this process satisfies an annealed FCLT: with respect to the probability $\mathbb{P} \times \mu$, the law of Y_n converges weakly to the Wiener measure.

We show a quenched FCLT for the r.f. X (and when X is a r.f. of moving averages of i.i.d. random variables in Section 4). As for the annealed FCLT in [5] and for Theorem 2.2 in [13] for a 1-dimensional stable r.w., the proof of Theorem 2.2 below is based on the maximal inequality shown by Newman and Wright [23] for associated r.v.s.

Definition 2.1. (cf. [16]) Recall that real random variables X_1, \dots, X_n are *associated* if, for every $n \geq 1$, for all non-decreasing (in each coordinate) functions $f, g : \mathbb{R}^n \mapsto \mathbb{R}$, we have $\text{Cov}(f(X_1, \dots, X_n), g(X_1, \dots, X_n)) \geq 0$ (if the covariance exists).

Non-decreasing functions of a family of associated random variables are associated [16]. Independent variables are associated. A family made out of a family of associated (in particular independent) variables with possible repetition is associated.

It follows that, if $(X_{\underline{\ell}}, \underline{\ell} \in \mathbb{Z}^2)$, are associated r.v.s, in particular independent, then the r.v.s $(X_{Z_k(\omega)}, k \geq 0)$, are associated for every $\omega \in \Omega$.

Theorem 2.2. *If $\mathbb{E}(X_0^2) = 1$, for \mathbb{P} -a.e. ω , the process $(Y_n(\omega, x, t))_{t \in [0,1]} = \left(\frac{S_{[nt]}^{\omega, X}(x)}{\sqrt{n \log n}} \right)_{t \in [0,1]}$ satisfies a FCLT with asymptotic variance $\sigma^2 = (\pi \sqrt{\det \Sigma})^{-1}$.*

Proof. 1) For the convergence of the finite dimensional distributions, the proof, relying on Cramér-Wold's theorem and Lindeberg's CLT, is as in [5]. Another proof, based on truncation and cumulants, can be given, like for the more general case of moving averages in Section 4.

2) *Tightness of the process (Y_n)*

The following is shown in [23, p. 673]:

Let U_1, U_2, \dots be centered associated random variables with finite second order moment. Put $S_k = \sum_{j=1}^k U_j$, for $k \geq 1$. Then, for every $\lambda > 0$ and $n \geq 1$, we have

$$(36) \quad \mu(\max_{1 \leq k \leq n} |S_k| \geq \lambda \|S_n\|_2) \leq 2\mu(|S_n| \geq (\lambda - \sqrt{2}) \|S_n\|_2).$$

Inequality (36) can be applied for every fixed ω to $U_j = X_{Z_j(\omega)}$ and to the sums $S_J = \sum_{j=b}^{b+k} X_{Z_j(\omega)}$ for any interval $J = [b, b+k] \subset [0, n]$. We also note that $\mathbb{E}(S_J^2) = \|X_0\|_2^2 V(\omega, J)$.

a) First, let us assume that $\mathbb{E}(X_0^4) < \infty$. With $K(\omega)$ given by (20), we have

$$(37) \quad \begin{aligned} & \left\| \sum_{i \in J} X_{Z_i(\omega)} \right\|_{4, \mu}^4 = 3\mathbb{E}(X_0^2)^2 \sum_{\underline{\ell}_1 \neq \underline{\ell}_2} w(\omega, J, \underline{\ell}_1)^2 w(\omega, J, \underline{\ell}_2)^2 + \mathbb{E}(X_0^4) \sum_{\underline{\ell}} w(\omega, J, \underline{\ell})^4 \\ & \leq 4\mathbb{E}(X_0^4) V(\omega, J)^2 \leq 4\mathbb{E}(X_0^4) (K(\theta^b \omega))^2 (k \log k)^2. \end{aligned}$$

Let C_1 be a constant > 0 such that $\mathbb{P}\{\omega : K(\omega) \leq C_1\} > 0$. Using Lemma 1.7, for $\delta \in]0, 1[$ and n big enough, there are times $0 = \rho_1 < \rho_2 < \dots < \rho_v \leq n < \rho_{v+1}$, with $v < 2/\delta$, such that $K(\theta^{\rho_i} \omega) \leq C_1$ and $\frac{1}{2}\delta n \leq \rho_{i+1} - \rho_i \leq \frac{3}{2}\delta n$, for $i = 1, \dots, v$.

Let $t_i = \frac{\rho_i}{n}$, $\lambda = \frac{\varepsilon}{\sqrt{\delta}}$, $J_i = [\rho_{i-1}, \dots, \rho_i]$, $m_i = \frac{2}{3}(\rho_{i+1} - \rho_i) \leq \delta n$. By (20) and (37), with $C = (2C_1)^{\frac{1}{2}}$, we have:

$$(38) \quad \left\| \sum_{j=\rho_{i-1}}^{\rho_i} X_{Z_j(\omega)} \right\|_2 \leq C \|X_0\|_2 (n \delta \log(n\delta))^{\frac{1}{2}},$$

$$(39) \quad \left\| \sum_{j=\rho_{i-1}}^{\rho_i} X_{Z_j(\omega)} \right\|_4 \leq C \|X_0\|_4 (n \delta \log(n\delta))^{\frac{1}{2}}, \forall i.$$

We can assume that $\delta \leq \frac{1}{8} \frac{\varepsilon^2}{C^2 \|X_0\|_2^2}$. This implies the inequality $\lambda_i - \sqrt{2} \geq \frac{1}{2} \lambda_i$ used below.

Using (36), we get (with $\sigma^{(i)} := \left\| \sum_{j=\rho_{i-1}}^{\rho_i} X_{Z_j(\omega)} \right\|_2$ and $\lambda_i := \varepsilon \sqrt{n \log n} / \sigma^{(i)}$), by Chebyshev's inequality (for a moment of order 4) and (39):

$$(40) \quad \begin{aligned} & \mu\left(\sup_{t_{i-1} \leq s \leq t_i} \left| \sum_{j=[t_{i-1}n]}^{[sn]} X_{Z_j(\omega)} \right| \geq \varepsilon \sqrt{n \log n}\right) = \mu\left(\max_{\rho_{i-1} \leq k \leq \rho_i} \left| \sum_{j=\rho_{i-1}}^k X_{Z_j(\omega)} \right| \geq \lambda_i \sigma^{(i)}\right) \\ & \leq 2\mu\left(\left| \sum_{j=\rho_{i-1}}^{\rho_i} X_{Z_j(\omega)} \right| \geq (\lambda_i - \sqrt{2}) \sigma^{(i)}\right) \leq 2\mu\left(\left| \sum_{j=\rho_{i-1}}^{\rho_i} X_{Z_j(\omega)} \right| \geq \frac{1}{2} \lambda_i \sigma^{(i)}\right) \\ & \leq 2\mu\left(\left| \sum_{j=\rho_{i-1}}^{\rho_i} X_{Z_j(\omega)} \right| \geq \frac{1}{2} \varepsilon \sqrt{n \log n}\right) \leq 2C^4 \|X_0\|_4^4 \frac{(n \delta \log(n\delta))^2}{\frac{1}{16} \varepsilon^4 (n \log n)^2} \leq 32C^4 \|X_0\|_4^4 \frac{\delta^2}{\varepsilon^4}. \end{aligned}$$

Observe now that (cf. [2, Theorem 7.4, p. 83])

$$\mu\left(\sup_{|t'-t| \leq \delta} |Y_n(t) - Y_n(s)| \geq 3\varepsilon\right) \leq \sum_{i=1}^v \mu\left(\sup_{t_{i-1} \leq s \leq t_i} \left| \sum_{j=[t_{i-1}n]}^{[sn]} X_{Z_j(\omega)} \right| \geq \varepsilon \sqrt{C_0 n \log n}\right).$$

Hence, by (40) we get

$$\mu\left(\sup_{|t'-t| \leq \delta} |Y_n(t) - Y_n(s)| \geq 3\varepsilon\right) \leq 32 \frac{C^4}{C_0^2} \|X_0\|_4^4 \frac{2 \delta^2}{\delta \varepsilon^4} = 64 \frac{C^4}{C_0^2} \|X_0\|_4^4 \frac{\delta}{\varepsilon^4}.$$

b) Now we use a truncation. For $L > 0$, let

$$\begin{aligned} \hat{X}_{\underline{k}}^L &:= X_{\underline{k}} \mathbf{1}_{\{|X_{\underline{k}}| \leq L\}} - \mathbb{E}(X_{\underline{k}} \mathbf{1}_{\{|X_{\underline{k}}| \leq L\}}), \quad \tilde{X}_{\underline{k}}^L := X_{\underline{k}} - \hat{X}_{\underline{k}}^L = X_{\underline{k}} \mathbf{1}_{\{|X_{\underline{k}}| > L\}} - \mathbb{E}(X_{\underline{k}} \mathbf{1}_{\{|X_{\underline{k}}| > L\}}), \\ \hat{Y}_n^L(t) &= \frac{1}{\sqrt{C_0 n \log n}} \sum_{j=0}^{[tn]} \hat{X}_{Z_j(\omega)}^L \quad \text{and} \quad \tilde{Y}_n^L(t) := Y_n(t) - \hat{Y}_n^L(t) = \frac{1}{\sqrt{C_0 n \log n}} \sum_{j=0}^{[tn]} \tilde{X}_{Z_j(\omega)}^L. \end{aligned}$$

Since we have still sums of associated random variables, all what we have done above (including (36)) holds for both sums, except that for the unbounded part of the truncation we only have a moment of order 2. We use Chebyshev's inequality (for a moment of order 2) and (38) to control the unbounded truncated part:

$$\mu\left(\left| \sum_{j=\rho_{i-1}}^{\rho_i} \tilde{X}_{Z_j(\omega)}^L \right| \geq \frac{1}{2} \varepsilon \sqrt{n \log n}\right) \leq C^2 \|\tilde{X}_0^L\|_2^2 \frac{n \delta \log(n\delta)}{\frac{1}{4} \varepsilon^2 n \log n} \leq 4C^2 \|\tilde{X}_0^L\|_2^2 \frac{\delta}{\varepsilon^2}.$$

Hence, for n large, the sum over i of the above measures at left is comparable for some constant C' to

$$C' \sum_{i=1}^v \|\tilde{X}_0^L\|_2^2 \leq \frac{C'}{\delta} \frac{\delta \|\tilde{X}_0^L\|_2^2}{\varepsilon^2} = C' \frac{\|\tilde{X}_0^L\|_2^2}{\varepsilon^2}.$$

Applying the inequality $\mu(|f + g| \geq \varepsilon) \leq \mu(|f| \geq \frac{\varepsilon}{2}) + \mu(|g| \geq \frac{\varepsilon}{2})$ to $Y_n(t) = \hat{Y}_n^L(t) + \tilde{Y}_n^L(t)$, we obtain the bound:

$$\mu\left(\sup_{|t'-t| \leq \delta} |Y_n(t') - Y_n(t)| \geq 3\varepsilon\right) \leq 2^{10} \frac{C^4}{C_0^2} \frac{L^4 \delta}{\varepsilon^4} + 4C' \frac{\|\tilde{X}_0^L\|_2^2}{\varepsilon^2}.$$

We need, for fixed $\varepsilon > 0$, $\lim_{\delta \rightarrow 0^+} \limsup_n \mu(\sup_{|t'-t| \leq \delta} |Y_n(t') - Y_n(t)| \geq 3\varepsilon) = 0$.

Let $\eta > 0$. First we take L such that $4C' \frac{\|\tilde{X}_0^L\|_2^2}{\varepsilon^2} < \frac{1}{2}\eta$, then δ such that $2^{10} \frac{C^4}{C_0^2} \frac{L^4 \delta}{\varepsilon^4} \leq \frac{1}{2}\eta$. \square

A model based on the Lorentz process

We sketch briefly how to obtain a version of a FCLT when the random walk is replaced by the movement of a particle in a dispersing periodic billiard. We refer to [24] and [25] for more details on this model.

Let be given a “billiard table” in the plane, union of \mathbb{Z}^2 -periodically distributed obstacles with pairwise disjoint closures. We consider a point particle moving in the complementary Q of the billiard table in \mathbb{R}^2 with unit speed and elastic reflection off the obstacles. By sampling the flow at the successive times of impact with the obstacles, we obtain a Poincaré’s section of the billiard flow, the billiard transformation.

We assume that the obstacles are strictly convex with pairwise disjoint closures and boundaries of class C^{r+1} with curvature > 0 (Sinai’s billiard or Lorentz’s process). Moreover we make the hypothesis of finite horizon (the time between two subsequent reflections is uniformly bounded).

Suppose that to each obstacle is associated a real random variable with zero expectation, positive and finite variance, independent of the motion of the particle and that the family of these r.v.s is i.i.d.

Like in an infinite “pinball” with random gain, at each collision with an obstacle, the particle wins the amount given by the random variable associated with the obstacle which is met. Let W_n be the total amount won by the particle after n reflections. An annealed FCLT for W_n has been shown by F. Pène ([24]): there exists $\beta_0 > 0$ such that $\frac{W_{[nt]}}{\beta_0 n \log n}$ converges weakly to the standard Wiener process.

In order to prove a quenched version, we use [24, Proposition 7], in place of (22) for the r.w., and [25, Corollary 4] (the main and most difficult step), which gives for the self-intersections of the billiard transformation a law of large numbers replacing (16). Then, by 1) in Remarks 1.11 and by the preceding method for the r.w. in random sceneries, we obtain the quenched version of the FCLT for this model.

3. Cumulants and CLT

For the models of random fields in Sections 4, 6, 7, we need to recall some tools, in particular the method of cumulants which can be used to prove a CLT for dynamical systems satisfying a mixing property of all orders.

In 1960, Leonov ([19], [20]) applied it to a single algebraic endomorphism of a compact abelian group. In [8], [9], it was applied to multidimensional actions given by algebraic endomorphisms in the connected case and in some non connected cases. Recently the method of cumulants has been used in [4] to prove a CLT for multiple mixing actions with exponential rate. Using the tightness criterium given in Section 5, we will obtain a functional version in these examples.

3.1. Moments and cumulants.

For $r \geq 1$, let X_1, \dots, X_r be r real centered bounded random variables. We denote by J_r the set $\{1, \dots, r\}$. For any subset $I = \{i_1, \dots, i_p\} \subset J_r$, we put $m(I) := \mathbb{E}(X_{i_1} \dots X_{i_p})$. The *cumulant* of order r is

$$(41) \quad \mathcal{C}(X_1, \dots, X_r) = \sum_{Q=\{I_1, \dots, I_p\} \in \mathcal{Q}} (-1)^{p-1} (p-1)! m(I_1) \dots m(I_p).$$

Putting $s(I) := \mathcal{C}(X_{i_1}, \dots, X_{i_p})$ for $I = \{i_1, \dots, i_p\}$, we have

$$(42) \quad \mathbb{E}(X_1 \dots X_r) = m(J_r) = \sum_{Q=\{I_1, \dots, I_p\} \in \mathcal{Q}} s(I_1) \dots s(I_p),$$

where in both formulas, \mathcal{Q} is the set of partitions $Q = \{I_1, I_2, \dots, I_p\}$ of J_r into $p \leq r$ nonempty subsets, with p varying from 1 to r .

For a single random variable Y , the cumulant of order r is defined by $\mathcal{C}^{(r)}(Y) := \mathcal{C}((Y, \dots, Y)_r)$, where $(Y, \dots, Y)_r$ is the vector with r components equal to Y . If Y is centered, we have $\|Y\|_2^2 = \mathcal{C}^{(2)}(Y)$ and

$$(43) \quad \mathbb{E}(Y^4) = 3\mathbb{E}(Y^2)^2 + \mathcal{C}^{(4)}(Y).$$

In the next sections, we are going to consider random fields obtained by a measure preserving action $(T^h, h \in \mathcal{H})$ of a group \mathcal{H} (with a left-invariant metric d on \mathcal{H}) on a probability space (E, \mathcal{B}, μ) . If f be a measurable bounded centered function on (E, μ) , the composed function $f \circ T^h$ is denoted by $T^h f$. For $\underline{H} = (h_1, \dots, h_r)$ in \mathcal{H}^r , we can apply the definition of moments and cumulants to $(T^{h_1} f, \dots, T^{h_r} f)$.

Notation 3.1. For the purpose of Section 7, we introduce some notations as in [4]. Let \underline{H} be in \mathcal{H}^r and I, J non empty subsets of J_r . We set

$$d^r(\underline{H}) := \max_{i,j} d(h_i, h_j), \quad d^I(\underline{H}) := \max_{i,j \in I} d(h_i, h_j), \quad d_{I,J}(\underline{H}) := \min_{i \in I, j \in J} d(h_i, h_j)$$

Let Q be a partition of J_r , with $|Q| \geq 2$. We set

$$d^Q(\underline{H}) := \max_{I \in Q} d^I(\underline{H}), \quad d_Q(\underline{H}) := \min_{I \neq J, I, J \in Q} d_{I,J}(\underline{H}),$$

and, for $0 \leq \alpha < \beta$,

$$\Delta(\beta) := \{\underline{H} \in \mathcal{H}^r : d^r(\underline{H}) \leq \beta\}, \quad \Delta_Q(\alpha, \beta) := \{\underline{H} \in \mathcal{H}^r : d^Q(\underline{H}) \leq \alpha \text{ and } d_Q(\underline{H}) > \beta\}.$$

The elements of a configuration $\underline{H} = (h_1, h_2, \dots, h_r)$ in $\Delta(\beta)$ can be viewed as ‘clustered’, since $(h_1, h_2, \dots, h_r) \in \Delta(\beta)$ implies $(h_2, \dots, h_r) \subset B_d(h_1, \beta)^{r-1}$. The configurations in $\Delta_Q(\alpha, \beta)$ for some partition Q and $0 < \alpha < \beta$ are made of ‘well-separated’ clusters.

Moment of order 4

The moment of order 4 plays a special role in the proof of tightness.

Let $J = [b, b+k]$ be an interval. We will bound the moment of order 4 of $Y = \sum_{j \in J} T^{Z_j(\omega)} f$, by using (43) and by bounding the cumulant of order 4: $\mathcal{C}^{(4)}(\sum_{j \in J} T^{Z_j(\omega)} f)$. We have:

$$|\mathcal{C}^{(4)}(\sum_{j \in J} T^{Z_j(\omega)} f)| \leq H(\omega, b, k) := \sum_{s, t, v, w \in J} |\mathcal{C}(T^{Z_s(\omega)} f, T^{Z_t(\omega)} f, T^{Z_v(\omega)} f, T^{Z_w(\omega)} f)|.$$

Observe that H is super-additive in the sense of the definition given later in Section 5.

Partitions of $\{1, 2, 3, 4\}$

In Formula (41) of cumulants, the contribution of a partition such that one of its atoms is a singleton is 0, and so does not appear. The partitions of $\{1, 2, 3, 4\}$ without atoms reduced to a singleton are $\{\{1, 2, 3, 4\}\}$, $\{\{1, 2\}, \{3, 4\}\}$, $\{\{1, 3\}, \{2, 4\}\}$, $\{\{1, 4\}, \{2, 3\}\}$.

The cumulant of order 4 of $(T^{h_i} f, T^{h_k} f, T^{h_\ell} f, T^{h_r} f)$ reads:

$$(44) \quad \mathcal{C}(T^{h_i} f, T^{h_k} f, T^{h_\ell} f, T^{h_r} f) = \mathbb{E}(T^{h_i} f T^{h_k} f T^{h_\ell} f T^{h_r} f) \\ - [\mathbb{E}(T^{h_i} f T^{h_k} f) \mathbb{E}(T^{h_\ell} f T^{h_r} f) + \mathbb{E}(T^{h_i} f T^{h_\ell} f) \mathbb{E}(T^{h_k} f T^{h_r} f) + \mathbb{E}(T^{h_i} f T^{h_r} f) \mathbb{E}(T^{h_k} f T^{h_\ell} f)].$$

Well separated configurations

The following proposition is a key step in the proof of the CLT shown in [4].

Proposition 3.2. (Proposition 6.2 in [4]) For every sequence $(\beta_j)_{j=0, \dots, r}$ such that

$$\beta_0 = 0 < \beta_1 < 3\beta_1 < \beta_2 < \dots < \beta_{r-1} < 3\beta_{r-1} < \beta_r,$$

we have

$$(45) \quad \mathcal{H}^r = \Delta(\beta_r) \cup \left(\bigcup_{j=0}^{r-1} \bigcup_{|Q| \geq 2} \Delta_Q(3\beta_j, \beta_{j+1}) \right).$$

Below, as an illustration, we give a proof for $r = 4$.

Proof. ($r = 4$) We represent an element of \mathcal{H} by \bullet and, given two elements $a, b \in \mathcal{H}$, draw $\bullet\bullet$ or $\bullet \bullet$ depending on whether they are close or far from each other.

Given a set $\underline{H} = \{t, u, v, w\}$ with 4 elements in \mathcal{H} , we will show that (up to a permutation) \underline{H} belongs to one of the configuration types $S_0 = \bullet\bullet\bullet\bullet$, $S_1 = \bullet\bullet\bullet \bullet$, $S_2 = \bullet\bullet \bullet\bullet$, $S_3 = \bullet\bullet \bullet \bullet$, or $S_4 = \bullet \bullet \bullet \bullet$.

The configurations of type S_0 are the ‘clustered’ configurations. This is quantified by saying that these configurations are in $\Delta(\beta)$ for some $\beta > 0$. The configurations of type S_4 are the configurations with pairwise distant elements. They correspond to $\Delta_Q(0, \beta)$ for some β .

We show that \underline{H} is either

of type S_0 , with $d(x, y) < \beta_4$, for every x, y in \underline{H} ,

or of type S_4 , with $d(x, y) \geq \beta_1$, for every $x \neq y$ in \underline{H} ,
or of one of the types S_1, S_2, S_3 .

If \underline{H} is not of type S_0 or S_4 , there are distinct elements, let call them t, u, w , such that $d(t, u) < \beta_1$ and $d(t, w) > \beta_4$, which implies

$$(46) \quad d(t, u) < \beta_1, d(t, w) > \beta_4 \geq \beta_3, d(u, w) > \beta_4 - \beta_1 \geq \beta_3.$$

One of the following cases occurs:

1) (type $S_1 = \bullet \bullet \bullet \bullet$) $d(v, t) < 2\beta_2$, which implies:

$$d(v, u) < 2\beta_2 + \beta_1 < 3\beta_2, d(v, w) > \beta_4 - 2\beta_2 > 3\beta_3 - 2\beta_2 > \beta_3.$$

In this case, for the partition $Q = \{\{t, u, v\}, \{w\}\}$, we get $d^Q(\underline{H}) < 3\beta_2$ and $d_Q(\underline{H}) > \beta_3$.

2) $d(v, t) > 2\beta_2$, which implies: $d(v, u) > 2\beta_2 - \beta_1 > \beta_2$,

2a) (type $S_2 = \bullet \bullet \bullet \bullet$) $d(v, w) < \beta_2$, which implies:

$$d(v, t) > d(t, w) - d(v, w) > \beta_4 - \beta_2 > 3\beta_3 - \beta_2 > \beta_3,$$

$$d(v, t) > d(t, w) - d(v, w) > \beta_4 - \beta_1 - \beta_2 > 3\beta_3 - \beta_1 - \beta_2 > \beta_3.$$

For $Q = \{\{t, u\}, \{v, w\}\}$, we get $d^Q(\underline{H}) < 3\beta_2$ and $d_Q(\underline{H}) > \beta_3$.

2b) (type $S_3 = \bullet \bullet \bullet \bullet$) $d(v, w) > \beta_2$:

For $Q = \{\{t, u\}, \{v\}, \{w\}\}$, we get $d^Q(\underline{H}) < 3\beta_1$ and $d_Q(\underline{H}) > \beta_2$. \square

3.2. A sufficient condition for the CLT.

Let us recall a criterium in terms of cumulants for the CLT (cf. [20, Th. 7], [8, Th. 6.2]). It is convenient to formulate the criterium by using a summation sequence, $w = (w_n)_{n \geq 1}$, i.e., for each n a function $w_n : \underline{\ell} \in \mathbb{Z}^d \rightarrow w_n(\underline{\ell}) \in \mathbb{R}$, with $0 < \sum_{\underline{\ell} \in \mathbb{Z}^d} |w_n(\underline{\ell})| < +\infty$.

The associated normalized non-negative kernel is $\tilde{K}(w_n)(\underline{t}) = \frac{|\sum_{\underline{\ell} \in \mathbb{Z}^d} w_n(\underline{\ell}) e^{2\pi i \langle \underline{\ell}, \underline{t} \rangle}|^2}{\sum_{\underline{\ell} \in \mathbb{Z}^d} |w_n(\underline{\ell})|^2}$, $\underline{t} \in \mathbb{T}^d$.

As for the summation along a random walk which is a special case, we say that the summation is ξ -regular for a probability measure ξ on \mathbb{T}^d , if the normalised kernel $(\tilde{K}(w_n)_{n \geq 1})$ converges weakly to ξ , i.e.,

$$(47) \quad \lim_{n \rightarrow \infty} \int_{\mathbb{T}^d} \tilde{K}(w_n) \varphi d\underline{t} = \xi(\varphi), \text{ for every continuous function } \varphi \text{ on } \mathbb{T}^d.$$

This implies that for f , under Condition (5), the asymptotic variance for the normalised sums is

$$\sigma_w^2(f) := \lim_n \frac{\|\sum_{\underline{\ell} \in \mathbb{Z}^d} w_n(\underline{\ell}) T^{\underline{\ell}} f\|_2^2}{\sum_{\underline{\ell} \in \mathbb{Z}^d} |w_n(\underline{\ell})|^2} = \xi(\varphi_f).$$

Theorem 3.3. *If $(w_n)_{n \geq 1}$ is a summation sequence on \mathbb{Z}^d such that (47) holds for a probability measure ξ on \mathbb{T}^d , the condition*

$$(48) \quad \sum_{(\underline{\ell}_1, \dots, \underline{\ell}_r) \in (\mathbb{Z}^d)^r} w_n(\underline{\ell}_1) \dots w_n(\underline{\ell}_r) \mathcal{C}(T^{\underline{\ell}_1} f, \dots, T^{\underline{\ell}_r} f) = o\left(\left(\sum_{\underline{\ell} \in \mathbb{Z}^d} w_n^2(\underline{\ell})\right)^{\frac{r}{2}}\right), \quad \forall r \geq 3,$$

$$(49) \quad \text{implies} \quad \left(\sum_{\underline{\ell} \in \mathbb{Z}^d} w_n^2(\underline{\ell})\right)^{-\frac{1}{2}} \sum_{\underline{\ell} \in \mathbb{Z}^d} w_n(\underline{\ell}) f(T^{\underline{\ell}} \cdot) \xrightarrow[n \rightarrow \infty]{} \mathcal{N}(0, \xi(\varphi_f)).$$

The following result (cf. [8, Lemma 6.6]) shows that mixing of all orders implies the asymptotic nullity of the cumulants.

Proposition 3.4. *Let $(T^\ell, \ell \in \mathbb{Z}^d)$ be a \mathbb{Z}^d -measure preserving action on a probability space (E, μ) . If it is mixing of order $r \geq 2$, then, for any $f \in L_0^\infty(X)$,*

$$(50) \quad \lim_{\max_{i \neq j} \|\ell_i - \ell_j\| \rightarrow \infty} \mathcal{C}(T^{\ell_1} f, \dots, T^{\ell_r} f) = 0.$$

Remark that (50) does not give the quantitative estimate (48). Nevertheless, in Section 6, (50) will be sufficient for an action by automorphisms of a connected compact abelian group (in particular of a torus) which is mixing, when f is a trigonometric polynomial. For general exponentially mixing actions, a quantitative formulation is needed as in [4], using Proposition 3.2.

Array of sequences and finite dimensional distributions

For $s \geq 1$ and $j = 1, \dots, s$, let $(w_{n,j}, n \geq 1)$ be s summation sequences, satisfying (47) with respectively $\xi = \xi_j$, where the ξ_j 's are probability measures on \mathbb{T}^d .

Using Theorem 3.3, we are going to deduce from the following two conditions the asymptotic normality (after normalization) of the vectorial process $(\sum_{\ell \in \mathbb{Z}^d} w_{n,1}(\ell) T^\ell f, \dots, \sum_{\ell \in \mathbb{Z}^d} w_{n,s}(\ell) T^\ell f)$:

- asymptotic orthogonality:

$$(51) \quad \int_{\mathbb{T}^d} \left(\sum_{\ell \in \mathbb{Z}^d} w_{n,j}(\ell) e^{2\pi i \langle \ell, t \rangle} \right) \left(\sum_{\ell \in \mathbb{Z}^d} w_{n,j'}(\ell) e^{-2\pi i \langle \ell, t \rangle} \right) e^{-2\pi i \langle p, t \rangle} dt \\ = o\left(\sum_{\ell \in \mathbb{Z}^d} |w_{n,j}(\ell)|^2 + \sum_{\ell \in \mathbb{Z}^d} |w_{n,j'}(\ell)|^2 \right), \forall j \neq j', \forall p \in \mathbb{Z}^d.$$

- convergence to 0 of the normalized cumulants of order ≥ 3 :

$$(52) \quad \sum_{(\ell_1, \dots, \ell_r) \in (\mathbb{Z}^d)^r} w_{n,i_1}(\ell_1) \dots w_{n,i_r}(\ell_r) \mathcal{C}(T^{\ell_1} f, \dots, T^{\ell_r} f) \\ = o\left(\sum_{\ell \in \mathbb{Z}^d} \left[\sum_{j=1}^s |w_{n,j}(\ell)|^2 \right]^{r/2} \right), \forall (i_1, \dots, i_r) \in \{1, \dots, s\}^r, \forall r \geq 3.$$

Proposition 3.5. *Under Conditions (51) and (52), the vectorial process*

$$\left(\frac{\sum_{\ell \in \mathbb{Z}^d} w_{n,1}(\ell) T^\ell f}{\left(\sum_{\ell \in \mathbb{Z}^d} |w_{n,1}(\ell)|^2 \right)^{\frac{1}{2}}}, \dots, \frac{\sum_{\ell \in \mathbb{Z}^d} w_{n,s}(\ell) T^\ell f}{\left(\sum_{\ell \in \mathbb{Z}^d} |w_{n,s}(\ell)|^2 \right)^{\frac{1}{2}}} \right)_{n \geq 1}$$

is asymptotically distributed as $\mathcal{N}(0, J_s)$, where J_s is the s -dimensional diagonal matrix with diagonal $(\xi_j(\varphi_f), j = 1, \dots, s)$.

Proof. The hypothesis (51) implies, for s non zero real parameters a_1, \dots, a_s :

$$(53) \quad \left(\sum_j a_j^2 \sum_{\ell \in \mathbb{Z}^d} |w_{n,j}(\ell)|^2 \right)^{-1} \sum_{\ell \in \mathbb{Z}^d} \sum_j a_j w_{n,j}(\ell) e^{2\pi i \langle \ell, t \rangle} \xrightarrow[n \rightarrow \infty]{\text{weakly}} \left(\sum_j a_j^2 \right)^{-1} \sum_j a_j^2 \xi_j.$$

Putting $w_n^{a_1, \dots, a_s}(\underline{\ell}) = a_1 w_{n,1}(\underline{\ell}) + \dots + a_s w_{n,s}(\underline{\ell})$, by the Cramér-Wold theorem, to conclude it suffices to show

$$(54) \quad \frac{\sum_{\underline{\ell} \in \mathbb{Z}^d} w_n^{a_1, \dots, a_s}(\underline{\ell}) T^{\underline{\ell}} f}{(a_1^2 \sum_{\underline{\ell} \in \mathbb{Z}^d} |w_{n,1}(\underline{\ell})|^2 + \dots + a_s^2 \sum_{\underline{\ell} \in \mathbb{Z}^d} |w_{n,s}(\underline{\ell})|^2)^{\frac{1}{2}}} \implies \mathcal{N}(0, \sum_{j=1}^s a_j^2 \xi_j(\varphi_f) / \sum_{j=1}^s a_j^2).$$

By (52), the sum $\sum_{i_1, \dots, i_r \in \{1, \dots, s\}^r} \sum_{(\underline{\ell}_1, \dots, \underline{\ell}_r) \in (\mathbb{Z}^d)^r} w_{n,i_1}(\underline{\ell}_1) \dots w_{n,i_r}(\underline{\ell}_r) \mathcal{C}(T^{\underline{\ell}_1} f, \dots, T^{\underline{\ell}_r} f)$ satisfies (48) and the result follows from Theorem 3.3. \square

We will use also the following lemma. Let $(w_n)_{n \geq 1}$ be a summation sequence on \mathbb{Z}^d such that (47) holds for a probability measure ξ on \mathbb{T}^d . For $f \in L^2(\mu)$, we put $\sigma_n(f) := \|\sum_{\underline{\ell}} w_n(\underline{\ell}) T^{\underline{\ell}} f\|_2$. We can suppose $\xi(\varphi_f) > 0$, since otherwise the limiting distribution is δ_0 .

Lemma 3.6. *Let $f, f_k, k \geq 1$ be in $L^2(\mu)$ and satisfying (5) such that $\|\varphi_{f-f_k}\|_\infty \rightarrow 0$. Then*

$$\frac{\sum_{\underline{\ell} \in \mathbb{Z}^d} w_n(\underline{\ell}) T^{\underline{\ell}} f_k}{\sigma_n(f_k)} \xrightarrow[n \rightarrow \infty]{} \mathcal{N}(0, 1), \forall k \geq 1, \text{ implies } \frac{\sum_{\underline{\ell} \in \mathbb{Z}^d} w_n(\underline{\ell}) T^{\underline{\ell}} f}{\sigma_n(f)} \xrightarrow[n \rightarrow \infty]{} \mathcal{N}(0, 1).$$

Proof. Let us consider the processes defined respectively by

$$U_n^k := \left(\sum_{\underline{\ell} \in \mathbb{Z}^d} w_n^2(\underline{\ell}) \right)^{-\frac{1}{2}} \sum_{\underline{\ell} \in \mathbb{Z}^d} w_n(\underline{\ell}) T^{\underline{\ell}} f_k, \quad U_n := \left(\sum_{\underline{\ell} \in \mathbb{Z}^d} w_n^2(\underline{\ell}) \right)^{-\frac{1}{2}} \sum_{\underline{\ell} \in \mathbb{Z}^d} w_n(\underline{\ell}) T^{\underline{\ell}} f.$$

By (47) we have:

$$\left(\sum_{\underline{\ell} \in \mathbb{Z}^d} w_n^2(\underline{\ell}) \right)^{-1} \left\| \sum_{\underline{\ell} \in \mathbb{Z}^d} w_n(\underline{\ell}) T^{\underline{\ell}} f \right\|_2^2 = \int_{\mathbb{T}^d} \tilde{K}(w_n) \varphi_f dt \xrightarrow[n \rightarrow \infty]{} \xi(\varphi_f).$$

Observe that, if ξ is a probability measure on \mathbb{T}^d , $f \rightarrow (\xi(\varphi_f))^{\frac{1}{2}}$ satisfies the triangular inequality; hence $\xi(\varphi_{f_k}) \geq \xi(\varphi_f) - \xi(\varphi_{f-f_k})$. Since $\xi(\varphi_{f-f_k}) \leq \|\varphi_{f-f_k}\|_\infty \rightarrow 0$, this implies $\xi(\varphi_{f_k}) \neq 0$ for k big enough.

The hypotheses imply “ $U_n^k \xrightarrow[n \rightarrow \infty]{} \mathcal{N}(0, \xi(\varphi_{f_k}))$ ” for every k . Moreover, since

$$\lim_n \int |U_n^k - U_n|^2 d\mu = \lim_n \int_{\mathbb{T}^d} \tilde{K}(w_n) \varphi_{f-f_k} dt = \xi(\varphi_{f-f_k}) \leq \|\varphi_{f-f_k}\|_\infty,$$

we have $\limsup_n \mu[|U_n^k - U_n| > \delta] \leq \delta^{-2} \limsup_n \int |U_n^k - U_n|^2 d\mu \xrightarrow[k \rightarrow \infty]{} 0$, for every $\delta > 0$.

Therefore, using [2, Theorem 3.2], the conclusion “ $U_n \xrightarrow[n \rightarrow \infty]{} \mathcal{N}(0, \xi(\varphi_f))$ ” follows. \square

4. Moving averages of i.i.d. random variables

Let $(X_\underline{\ell})_{\underline{\ell} \in \mathbb{Z}^2}$ be a r.f. of centered i.i.d. real random variables on a probability space (E, \mathcal{B}, μ) such that $\|X_0\|_2 = 1$. Let $(a_q)_{q \in \mathbb{Z}^2}$ be an array of real numbers such that $\sum_{q \in \mathbb{Z}^2} |a_q| < \infty$ and let $\Xi = (\Xi_\underline{\ell})_{\underline{\ell} \in \mathbb{Z}^2}$ be the random field defined by $\Xi_\underline{\ell}(x) = \sum_{q \in \mathbb{Z}^2} a_q X_{\underline{\ell}-q}(x)$.

The correlation is $\widehat{\varphi}_\Xi(\underline{\ell}) = \left\langle \sum_{q \in \mathbb{Z}^2} a_q X_{\underline{\ell}-q}, \sum_{q' \in \mathbb{Z}^2} a_{q'} X_{-q'} \right\rangle = \sum_{q \in \mathbb{Z}^2} a_q a_{q-\underline{\ell}}$. We have

$$\sum_{\underline{\ell}} |\widehat{\varphi}_\Xi(\underline{\ell})| \leq \sum_{\underline{\ell}} \sum_{q \in \mathbb{Z}^2} |a_q| |a_{q-\underline{\ell}}| = \left(\sum_{q \in \mathbb{Z}^2} |a_q| \right)^2 < +\infty.$$

The continuous spectral density of the process is $\varphi_{\Xi}(\underline{t}) = |\sum_{\underline{q}} a_{\underline{q}} e^{2\pi i \langle \underline{q}, \underline{t} \rangle}|^2$. The asymptotic variance for the summation along the r.w. (with the normalisation by $C_0 n \log n$) is $(\sum_{\underline{q} \in \mathbb{Z}^2} a_{\underline{q}})^2$ that we suppose $\neq 0$.

Using the method of associated r.v.s we obtain a quenched FCLT for $S_{[nt]}^{\omega, \Xi}$ (cf. Notation (1)):

Theorem 4.1. *Let $(\Xi_{\underline{\ell}}, \underline{\ell} \in \mathbb{Z}^2)$ be the random field defined by moving averages as above. Under the assumption $\sum_{\underline{q} \in \mathbb{Z}^2} |a_{\underline{q}}| < \infty$, the process $(\frac{S_{[nt]}^{\omega, \Xi}(x)}{\sqrt{n \log n}})_{t \geq 0}$ satisfies a quenched FCLT with asymptotic variance $\sigma^2 = |\sum_{\underline{q} \in \mathbb{Z}^2} a_{\underline{q}}|^2 (\pi \sqrt{\det \bar{\Sigma}})^{-1}$.*

Proof. 1) *Convergence of the finite dimensional distributions*

a) First we assume that the random variables $X_{\underline{\ell}}, \underline{\ell} \in \mathbb{Z}^2$, are bounded. Moreover let us consider first a finite sum $F = \sum_{\underline{s} \in S} a_{\underline{s}} X_{\underline{s}}$, where S is a finite subset of \mathbb{Z}^2 . The case of the series, $\Xi_0 = \sum_{\underline{s} \in \mathbb{Z}^2} a_{\underline{s}} X_{\underline{s}}$, will follow by an approximation argument.

As we have seen, for $Y_n(\omega, x, t) = \frac{S_{[nt]}^{\omega, X}(x)}{\sqrt{C_0 n \log n}}$, we have to show:

$$\forall 0 = t_0 < t_1 < \dots < t_r = 1, (Y_n(t_1), \dots, Y_n(t_r)) \xrightarrow[n \rightarrow \infty]{} (W_{t_1}, \dots, W_{t_r}).$$

For it, we use Proposition 3.5. Condition (51) follows from Lemma 1.10. Let us check (52).

There is M such that the cumulant $\mathcal{C}(T^{\underline{\ell}_1} F, \dots, T^{\underline{\ell}_r} F) = 0$, if $\max_{i,j} \|\underline{\ell}_i - \underline{\ell}_j\| > M$, because if M is big enough, there is a random variable $T^{\underline{\ell}_i} F$ which is independent from σ -algebra generated by the others in the collection $T^{\underline{\ell}_1} F, \dots, T^{\underline{\ell}_r} F$ (by finiteness of S).

Let $w_n^u(\omega, \underline{\ell}) := \sum_{j=nt_u}^{nt_{u+1}} 1_{Z_j = \underline{\ell}}$, for $u = 0, \dots, r-1$. Then we have $w_n^u(\omega, \underline{\ell}) \leq w_n(\omega, \underline{\ell})$ and, since

$$\sup_{\underline{\ell}_1, \dots, \underline{\ell}_r} |\mathcal{C}(T^{\underline{\ell}_1} F, T^{\underline{\ell}_2} F, \dots, T^{\underline{\ell}_r} F)| < \infty:$$

$$\begin{aligned} & \left| \sum_{\max_{i,j} \|\underline{\ell}_i - \underline{\ell}_j\| \leq M} \mathcal{C}(T^{\underline{\ell}_1} F, T^{\underline{\ell}_2} F, \dots, T^{\underline{\ell}_r} F) w_n^{i_1}(\omega, \underline{\ell}_1) w_n^{i_2}(\omega, \underline{\ell}_2) \dots w_n^{i_r}(\omega, \underline{\ell}_r) \right. \\ & \leq \sum_{\underline{\ell}} \sum_{\|\underline{j}_2\|, \dots, \|\underline{j}_r\| \leq M, \underline{j}_1 = 0} |\mathcal{C}(T^{\underline{\ell}} F, T^{\underline{\ell} + \underline{j}_2} F, \dots, T^{\underline{\ell} + \underline{j}_r} F)| \prod_{k=1}^r w_n^{i_k}(\omega, \underline{\ell} + \underline{j}_k) \\ & \leq C \sum_{\underline{\ell}} \sum_{\|\underline{j}_2\|, \dots, \|\underline{j}_r\| \leq M, \underline{j}_1 = 0} \prod_{k=1}^r w_n^{i_k}(\omega, \underline{\ell} + \underline{j}_k) \leq C \sum_{\underline{\ell}} \sum_{\|\underline{j}_2\|, \dots, \|\underline{j}_r\| \leq M, \underline{j}_1 = 0} \prod_{k=1}^r w_n(\omega, \underline{\ell} + \underline{j}_k). \end{aligned}$$

The right hand side is less than a finite sum of sums of the form $\sum_{\underline{\ell} \in \mathbb{Z}^d} \prod_{k=1}^r w_n(\omega, \underline{\ell} + \underline{j}_k)$ with $\{\underline{j}_1, \dots, \underline{j}_r\} \in \mathbb{Z}^2$.

By (22), for every $\varepsilon > 0$, there is $C_{\varepsilon}(\omega)$ a.e. finite such that $\sup_{\underline{\ell}} w_n(\omega, \underline{\ell}) \leq C_{\varepsilon}(\omega) n^{\varepsilon}$. For $r \geq 3$, take $\varepsilon < \frac{r-2}{2(r-1)}$. We have then $\sum_{\underline{\ell} \in \mathbb{Z}^d} \prod_{k=1}^r w_n(\omega, \underline{\ell} + \underline{j}_k) \leq C_{\varepsilon}(\omega)^{r-1} n^{\varepsilon(r-1)} n = o(n^{r/2})$ and (52) is satisfied.

Using Lemma 3.6, the result can be extended to a general sum $\sum_{\underline{s} \in S} a_{\underline{s}} X_{\underline{s}}$, such that $\sum_{\underline{s} \in S} |a_{\underline{s}}| < \infty$.

b) Now if we assume only the condition $\|X_0\|_2 < \infty$, we apply a truncation argument to the independent r.v.s $X_{\underline{\ell}}$ and use again Lemma 3.6.

2) *Tightness* Let $a_{\underline{q}}^+ = \max(a_{\underline{q}}, 0)$, $a_{\underline{q}}^- = \max(-a_{\underline{q}}, 0)$. Observe that the random variables $\sum_{\underline{q} \in \mathbb{Z}^2} a_{\underline{q}}^+ X_{\underline{\ell}-\underline{q}}(x) = \sum_{\underline{q} \in \mathbb{Z}^2} a_{\underline{q}+\underline{\ell}}^+ X_{-\underline{q}}(x)$, for $\underline{\ell} \in \mathbb{Z}^2$, are associated (as non decreasing functions of associated r.v.s), as well as $\sum_{\underline{q} \in \mathbb{Z}^2} a_{\underline{q}}^- X_{\underline{\ell}-\underline{q}}(x)$, for $\underline{\ell} \in \mathbb{Z}^2$.

Therefore tightness can be proved separately for both processes. Now, again we use a truncation of the independent r.v.s $X_{\underline{\ell}}$. For $L > 0$, we put

$$\begin{aligned}\hat{X}_{\underline{k}}^L &:= X_{\underline{k}} \mathbf{1}_{\{|X_{\underline{k}}| \leq L\}} - \mathbb{E}(X_{\underline{k}} \mathbf{1}_{\{|X_{\underline{k}}| \leq L\}}), \\ \tilde{X}_{\underline{k}}^L &:= X_{\underline{k}} - \hat{X}_{\underline{k}}^L = X_{\underline{k}} \mathbf{1}_{\{|X_{\underline{k}}| > L\}} - \mathbb{E}(X_{\underline{k}} \mathbf{1}_{\{|X_{\underline{k}}| > L\}}).\end{aligned}$$

The r.v.s $(\sum_{\underline{q} \in \mathbb{Z}^2} a_{\underline{q}+\underline{\ell}}^+ \hat{X}_{-\underline{q}}(x), \underline{\ell} \in \mathbb{Z}^2)$ are still associated, as well as $(\sum_{\underline{q} \in \mathbb{Z}^2} a_{\underline{q}+\underline{\ell}}^+ \tilde{X}_{-\underline{q}}(x), \underline{\ell} \in \mathbb{Z}^2)$.

Moreover, we control the norms in $L^2(\mu)$ and $L^4(\mu)$, since by the triangular inequality, it holds:

$$\begin{aligned}\left\| \sum_{j=A}^B \sum_{\underline{q} \in \mathbb{Z}^2} a_{\underline{q}}^+ \hat{X}_{Z_j(\omega)-\underline{q}} \right\|_{\mu,4} &\leq \left(\sum_{\underline{q} \in \mathbb{Z}^2} a_{\underline{q}}^+ \right) \left\| \sum_{j=A}^B \hat{X}_{Z_j(\omega)} \right\|_{\mu,4}, \\ \left\| \sum_{j=A}^B \sum_{\underline{q} \in \mathbb{Z}^2} a_{\underline{q}}^+ \tilde{X}_{Z_j(\omega)-\underline{q}} \right\|_{\mu,2} &\leq \left(\sum_{\underline{q} \in \mathbb{Z}^2} a_{\underline{q}}^+ \right) \left\| \sum_{j=A}^B \tilde{X}_{Z_j(\omega)} \right\|_{\mu,2}.\end{aligned}$$

Now the proof is like the proof of tightness in Theorem 2.2. □

5. Tightness and 4th-moment

In this section, we show a criterium of tightness based on the 4th-moment.

Let $n \geq 1$. We say that a nonnegative function $G_0 = (G_0(b, k))$, defined for b, k such that $0 \leq b \leq b+k \leq n$, is super-additive if $G_0(b, 0) = 0$ and

$$(55) \quad G_0(b, k) + G_0(b+k, \ell) \leq G_0(b, k+\ell), \quad \forall b \geq 0, \forall k, \ell \geq 1 \text{ such that } b+k+\ell \leq n.$$

Let (W_k) be a sequence of real or complex random variables on a probability space (E, μ) . We set

$$S_{b,k} = \sum_{r=b+1}^{b+k} W_r, \quad M_{b,n} = \max_{1 \leq k \leq n} |S_{b,k}|.$$

The following result is adapted from [22]:

Theorem 5.1. (*F. Móricz*) *Let $n \geq 1$. Suppose that there exists G_0 a super-additive function such that*

$$(56) \quad \mathbb{E}_{\mu}(|S_{b,k}|^4) \leq G_0^2(b, k), \quad \forall b, k \text{ such that } 0 \leq b \leq b+k \leq n.$$

Then, with the constant $C_{max} = (1 - 2^{-\frac{1}{4}})^{-4}$,

$$(57) \quad \mathbb{E}_{\mu}(|M_{b,n}|^4) \leq C_{max} G_0^2(b, n), \quad \forall b \leq n.$$

Let $X = (X_\ell)_{\ell \in \mathbb{Z}^2}$ be a strictly stationary real random field on a probability space (E, μ) , where the X_ℓ 's have zero mean and finite second moment. Setting $S_J^\omega(x) = \sum_{i \in J} X_{Z_i(\omega)}(x)$ if J is an interval, we deduce from (57) a criterium for tightness adapted to the sums along a random walk.

Proposition 5.2. *Let $G(\omega, \cdot, \cdot)$, $H(\omega, \cdot, \cdot)$ be super-additive functions such that for a parameter γ and $K_1(\omega), K_2(\omega)$ a.e. finite functions on (Ω, \mathbb{P}) ,*

$$(58) \quad G(\omega, b, k) \leq K_1(\theta^b \omega) k \log k, \quad H(\omega, b, k) \leq K_2(\theta^b \omega) k (\log k)^\gamma, \quad G(\omega, b, k) \geq k.$$

Suppose that the r.v.s X_ℓ are bounded and satisfy

$$(59) \quad \mathbb{E}_\mu(|S_J^{\omega, X}|^4) \leq G(\omega, b, k)^2 + n^{\frac{1}{2}} (\log n)^{-(\gamma+1)} H(\omega, b, k), \quad \forall J = [b, b+k] \subset [1, n], \text{ for a.e. } \omega.$$

Then, for every $\varepsilon > 0$, $Y_n(\omega, x, t) = \frac{1}{\sqrt{n \log n}} \sum_{j=1}^{\lfloor nt \rfloor} X_{Z_j(\omega)}$ satisfies

$$(60) \quad \lim_{\delta \rightarrow 0^+} \limsup_n \mu(x \in E : \sup_{|t'-t| \leq \delta} |Y_n(\omega, x, t') - Y_n(\omega, x, t)| \geq \varepsilon) = 0.$$

Proof. 1) Let $c \geq 0$, $\Delta_n = \lceil n^{\frac{1}{2}} (\log n)^{-2} \rceil + 1$, $\nu = \nu_n \geq \Delta_n$, $L_n = \lfloor \frac{\nu_n}{\Delta_n} \rfloor$, $\nu' = \nu'_n = \lfloor \frac{\nu_n}{\Delta_n} \rfloor \Delta_n + \Delta_n - 1$. The integer ν_n will be chosen of order δn . Since Δ_n is an integer ≥ 1 , we have $\nu \leq \nu'$. We can write, with the convention that $\sum_{r=0}^{-1} = 0$:

$$\begin{aligned} \max_{0 \leq k \leq \nu} \left| \sum_{j=0}^k X_{Z_{j+c}(\omega)} \right| &\leq \max_{0 \leq k \leq \nu'} \left| \sum_{j=0}^k X_{Z_{j+c}(\omega)} \right| = \max_{0 \leq u \leq \lfloor \frac{\nu'}{\Delta_n} \rfloor, 1 \leq k \leq \Delta_n} \left| \sum_{r=0}^{u-1} \sum_{j=r\Delta_n}^{(r+1)\Delta_n-1} X_{Z_{j+c}(\omega)} + \sum_{j=u\Delta_n}^{u\Delta_n+k-1} X_{Z_{j+c}(\omega)} \right| \\ &\leq \max_{0 < u \leq L_n, 1 \leq k \leq \Delta_n} \left| \sum_{r=0}^{u-1} \sum_{j=r\Delta_n}^{(r+1)\Delta_n-1} X_{Z_{j+c}(\omega)} \right| + \max_{0 \leq u \leq L_n, 1 \leq k \leq \Delta_n} \left| \sum_{j=u\Delta_n}^{u\Delta_n+k-1} X_{Z_{j+c}(\omega)} \right| \\ &= \max_{0 < u \leq L_n} \left| \sum_{j=0}^{u\Delta_n} X_{Z_{j+c}(\omega)} \right| + \max_{0 \leq u \leq L_n, 1 \leq k \leq \Delta_n} \left| \sum_{j=u\Delta_n}^{u\Delta_n+k-1} X_{Z_{j+c}(\omega)} \right|. \end{aligned}$$

Let \hat{A}_n and \tilde{A}_n be respectively defined as the first and the second term above. This previous inequality implies

$$(61) \quad \mu\left(\max_{0 \leq k \leq \nu} \left| \sum_{j=0}^k X_{Z_{j+c}(\omega)} \right| \geq \varepsilon \sqrt{n \log n}\right) \leq \mu(\hat{A}_n \geq \frac{1}{2} \varepsilon \sqrt{n \log n}) + \mu(\tilde{A}_n \geq \frac{1}{2} \varepsilon \sqrt{n \log n}).$$

For \tilde{A}_n , since the X_ℓ 's are bounded (uniformly in ℓ by stationarity), by the choice of Δ_n there is $N_1(\varepsilon, \delta)$ such that $\mu(\tilde{A}_n \geq \frac{1}{2} \varepsilon \sqrt{n \log n}) = 0$, for $n \geq N_1(\varepsilon, \delta)$.

For \hat{A}_n we will apply Theorem 5.1 to $W_r = \sum_{j=r\Delta_n}^{(r+1)\Delta_n-1} X_{Z_{j+c}(\omega)}$, with

$$(62) \quad G_0(b, k) := G(\omega, c + b\Delta_n, k\Delta_n) + (\log n)^{-\gamma+1} H(\omega, c + b\Delta_n, k\Delta_n),$$

which is super-additive as G and H .

Since $G(\omega, c + b\Delta_n, k\Delta_n) \geq k\Delta_n \geq \Delta_n = \lceil n^{\frac{1}{2}} (\log n)^{-2} \rceil$, we have for $k \geq 1$:

$$G^2(\omega, c + b\Delta_n, k\Delta_n) + n^{\frac{1}{2}} (\log n)^{-\gamma-1} H(\omega, c + b\Delta_n, k\Delta_n) \leq G_0^2(b, k).$$

Therefore, by (59),

$$\mathbb{E}_\mu(|\sum_{r=b+1}^{b+k} W_r|^4) = \mathbb{E}_\mu(|\sum_{j=(b+1)\Delta_n}^{(b+k+1)\Delta_n-1} X_{Z_{j+c}(\omega)}|^4) \leq G_0(b, k)^2, \forall b \geq 0, \forall k \geq 1,$$

which implies by (57) of Theorem 5.1:

$$\mathbb{E}_\mu(\max_{1 \leq k \leq p} |\sum_{j=(b+1)\Delta_n}^{(b+k+1)\Delta_n-1} X_{Z_{j+c}(\omega)}|^4) \leq C_{\max} G_0(b, p)^2, \forall b \geq 0, \forall p \geq 1.$$

Putting $K(\omega) := \max(K_1(\omega), K_2(\omega))$ and using (58), we get the bound

$$\begin{aligned} & \|\max_{u=1}^{L_n} |\sum_{j=0}^{u\Delta_n} X_{Z_{j+c}(\omega)}|\|_4^4 \leq C_{\max} [G(\omega, c, L_n\Delta_n) + (\log n)^{-\gamma+1} H(\omega, c, L_n\Delta_n)]^2 \\ (63) \quad & \leq C_{\max} K(\theta^c\omega)^2 [L_n\Delta_n \log(L_n\Delta_n) + (\log n)^{-\gamma+1} (L_n\Delta_n)(\log(L_n\Delta_n))^\gamma]^2. \end{aligned}$$

2) For $M > 0$ big enough, the set $\Omega_M := \{\omega : K(\omega) \leq M\}$ has a probability $\mathbb{P}(\Omega_M) \geq \frac{1}{2}$. We apply Lemma 1.7 to Ω_M . Given $\delta > 0$, there is $N_2(\delta)$ such that for $n \geq N_2(\delta)$, we can find a sequence $0 = \rho_{1,n} < \rho_{2,n} < \dots < \rho_{v,n} \leq n < \rho_{v+1,n}$ of visit times of $\theta^k\omega$ in Ω_M under the iteration of the shift θ , such that $\frac{1}{2}\delta n \leq \rho_{i+1,n} - \rho_{i,n} \leq \frac{3}{2}\delta n$ and $v < 2/\delta$. By construction, $K(\theta^{\rho_{i,n}}\omega) \leq M, \forall i$.

With $c = \rho_{i,n}$, $\nu_n = \nu_{i,n} = \rho_{i+1,n} - \rho_{i,n} \leq \frac{3}{2}\delta n$, $L_{i,n} = [\frac{\nu_{i,n}}{\Delta_n}]$ (so that $L_{i,n}\Delta_n \leq \frac{3}{2}\delta n$), we deduce from the upper bound (63) (for n big enough and using $0 \leq \log(\delta n) \leq \log n$, if $n \geq \delta^{-1}$):

$$\begin{aligned} & \|\max_{u=1}^{L_{i,n}} |\sum_{j=0}^{u\Delta_n} X_{Z_{\rho_{i,n}+j}(\omega)}|\|_4^4 \leq C_{\max} M [\nu_{i,n} \log \nu_{i,n} + (\log n)^{-\gamma+1} \nu_{i,n} (\log \nu_{i,n})^\gamma]^2 \\ & \leq C_{\max} M [\frac{3}{2}\delta n \log(\delta n) + (\log n)^{-\gamma+1} \frac{3}{2}\delta n (\log(\delta n))^\gamma]^2 \leq C_{\max} M [3\delta n \log n]^2. \end{aligned}$$

This implies that there is a constant C such that, for $n \geq N_2(\delta)$, for $i = 1, \dots, v$:

$$\mu(\max_{0 \leq u \leq L_{i,n}} |\sum_{j=0}^{u\Delta_n-1} X_{Z_{j+\rho_i}(\omega)}| \geq \frac{1}{2}\varepsilon \sqrt{n \log n}) \leq \frac{C_{\max} M (3\delta n \log n)^2}{(\frac{1}{2}\varepsilon \sqrt{n \log n})^4} \leq C \varepsilon^{-4} \delta^2.$$

This allows to bound the term $\mu(\hat{A}_n \geq \frac{1}{2}\varepsilon \sqrt{n \log n})$ in (61): Putting $t_i = \rho_{i,n}/n$, we obtain, for $n \geq \max(N_1(\varepsilon, \delta), N_2(\delta))$,

$$\mu(\sup_{|t'-t| \leq \delta} |Y_n(t') - Y_n(t)| \geq 3\varepsilon) \leq \sum_{i=1}^v \mu(\sup_{t_{i-1} \leq s \leq t_i} |Y_n(s) - Y_n(t_{i-1})| \geq \varepsilon) \leq 2C \varepsilon^{-4} \delta^2 v \leq 2C \frac{\delta}{\varepsilon^4}. \quad \square$$

Remark 5.3. Let be given for each s in a set of indices S a process $X^s = (X_{s,\underline{\ell}})_{\underline{\ell} \in \mathbb{Z}^2}$ satisfying the hypotheses of the proposition, with the same uniform bound and the same G, H, γ . Then, if $X_{\underline{\ell}} = \sum_s a_s X_{s,\underline{\ell}}^s$ with $\sum_s |a_s| \leq 1$, the r.f. $X = (X_{\underline{\ell}})$ satisfies the conditions of the proposition and therefore the conclusion (60). This follows from Minkowski inequality:

$$\begin{aligned} \|S_J^{\omega, X}\|_4^4 & \leq (\sum_s |a_s| \|S_J^{\omega, X^s}\|_4)^4 \leq (\sum_s |a_s| [G(\omega, b, k)^2 + n^{\frac{1}{2}} (\log n)^{-(\gamma+1)} H(\omega, b, k)]^{\frac{1}{4}})^4 \\ & = (\sum_s |a_s|)^4 [G(\omega, b, k)^2 + n^{\frac{1}{2}} (\log n)^{-(\gamma+1)} H(\omega, b, k)]. \end{aligned}$$

6. Automorphisms of a torus

We consider now a random field generated by the action of commuting automorphisms on a torus. Let us first present the model. We will give the details of the proof for $d = 2$.

Actions by endomorphisms on a compact abelian group: Let G be a compact abelian group with Haar measure μ . The group of characters of G is denoted by \hat{G} and the set of non trivial characters by \hat{G}^* . The Fourier coefficients of a function f in $L^1(G, \mu)$ (denoted also $\hat{f}(\underline{k})$ when G is a torus) are $c_f(\chi) := \int_G \bar{\chi} f d\mu$, $\chi \in \hat{G}$.

Every surjective endomorphism of G defines a measure preserving transformation on (G, μ) and a dual injective endomorphism on \hat{G} .

Let (T_1, \dots, T_d) be a finite family of d commuting surjective endomorphisms of G and $T^\ell = T_1^{\ell_1} \dots T_d^{\ell_d}$, for $\underline{\ell} = (\ell_1, \dots, \ell_d) \in \mathbb{N}^d$. We obtain a \mathbb{N}^d -action $(T^\ell, \underline{\ell} \in \mathbb{Z}^d)$ on G , which is totally ergodic if and only if the dual action is free.

Let $AC_0(G)$ denote the space of real functions on G with *absolutely convergent Fourier series* and $\mu(f) = 0$, endowed with the norm: $\|f\|_c := \sum_{\chi \in \hat{G}} |c_f(\chi)| < +\infty$.

Recall that the action on G is mixing of all orders when it is totally ergodic and G is connected.

Proposition 6.1. *If f is in $AC_0(G)$, the spectral density φ_f is continuous on \mathbb{T}^d and $\|\varphi_f\|_\infty \leq \|f\|_c^2$. For every $\varepsilon > 0$ there is a trigonometric polynomial P such that $\|\varphi_{f-P}\|_\infty \leq \varepsilon$.*

Proof. Since by total ergodicity the characters $T^\ell \chi$ for $\underline{\ell} \in \mathbb{Z}^d$ are pairwise distinct, we have

$$\sum_{\underline{\ell} \in \mathbb{Z}^d} |\langle T^\ell f, f \rangle| \leq \sum_{\underline{\ell} \in \mathbb{Z}^d} \sum_{\chi \in \hat{G}^*} |c_f(T^\ell \chi)| |c_f(\chi)| \leq \sum_{\chi \in \hat{G}^*} \left(\sum_{\underline{\ell} \in \mathbb{Z}^d} |c_f(T^\ell \chi)| \right) |c_f(\chi)| \leq \left(\sum_{\chi \in \hat{G}^*} |c_f(\chi)| \right)^2.$$

Therefore, if f is in $AC_0(G)$, then $\sum_{\underline{\ell} \in \mathbb{Z}^d} |\langle T^\ell f, f \rangle| < \infty$, the spectral density is continuous and $\|\varphi_f\|_\infty \leq \varepsilon$. By this inequality, we can take for P the restriction of the Fourier series of f to a finite set \mathcal{E} in \hat{G} , where \mathcal{E} is such that $\|\varphi_{f-P}\|_\infty \leq \left(\sum_{\chi \in \hat{G} \setminus \mathcal{E}} |c_f(\chi)| \right)^2 \leq \varepsilon$. \square

For compact abelian groups which are connected (cf. [8]) or which belong to a special family of non connected groups (cf. [9]), a CLT has been shown for summation either over sets or along a random walk. Our aim is to extend this latter result to a functional CLT at least in the case of automorphisms of a torus.

Matrices and automorphisms of a torus:

Now we will restrict to the special case of $G = \mathbb{T}^\rho$, $\rho > 1$.

Every A in the semigroup $\mathcal{M}^*(\rho, \mathbb{Z})$ of non singular $\rho \times \rho$ matrices with coefficients in \mathbb{Z} defines a surjective endomorphism of \mathbb{T}^ρ and a measure preserving transformation on (\mathbb{T}^ρ, μ) . It defines also a dual endomorphism of the group of characters $\widehat{\mathbb{T}^\rho}$ identified with \mathbb{Z}^ρ (this is the action by the transposed matrix, but since we compose commuting matrices, for simplicity we do not write the transposition). The linear operator on \mathbb{C}^ρ defined by A is denoted by \tilde{A} .

When A is in the group $GL(\rho, \mathbb{Z})$ of matrices with coefficients in \mathbb{Z} and determinant ± 1 , it defines an automorphism of \mathbb{T}^ρ . Recall that the action of $A \in \mathcal{M}^*(\rho, \mathbb{Z})$ on (\mathbb{T}^ρ, μ) is ergodic if and only if A has no eigenvalue root of unity.

Here we present the proof for the case of automorphisms and for $d = 2$ (in the recurrent case for the random walk). Let (A_1, A_2) be two commuting matrices in $GL(\rho, \mathbb{Z})$ and $A^\ell = A_1^{\ell_1} A_2^{\ell_2}$, for $\ell = (\ell_1, \ell_2) \in \mathbb{Z}^2$. It defines a \mathbb{Z}^2 -action $(A^\ell, \ell \in \mathbb{Z}^2)$ on (\mathbb{T}^ρ, μ) , which is totally ergodic if and only if A^ℓ has no eigenvalue root of unity for $\ell \neq \underline{0}$.

Explicit totally ergodic \mathbb{Z}^2 -actions can be computed (cf. [8]) like the example below (see the book of H. Cohen on computational algebraic number theory [11]):

$$A_1 = \begin{pmatrix} -3 & -3 & 1 \\ 10 & 9 & -3 \\ -30 & -26 & 9 \end{pmatrix}, \quad A_2 = \begin{pmatrix} 11 & 1 & -1 \\ -10 & -1 & 1 \\ 10 & 2 & -1 \end{pmatrix}.$$

We will need an algebraic result based on the following theorem on S-unit equations ([26]):

Theorem 6.2. ([17, Th. 1.1]) *Let K be an algebraically closed field of characteristic 0 and for $r \geq 2$, let Γ_r be a subgroup of the multiplicative group $(K^*)^r$ of finite rank. For any $(a_1, \dots, a_r) \in (K^*)^r$, the number of solutions $x = (x_1, \dots, x_r) \in \Gamma_r$ of the equation $a_1 x_1 + \dots + a_r x_r = 1$ such that no proper subsum of $a_1 x_1 + \dots + a_r x_r$ vanishes, is finite.*

Corollary 6.3. *Suppose that the \mathbb{Z}^2 -action $(A^\ell, \ell \in \mathbb{Z}^2)$ is totally ergodic. The set F of triples $(\ell_1, \ell_2, \ell_3) \in (\mathbb{Z}^2)^3$ for which there is $\gamma \in \mathbb{Z}^2 \setminus \{\underline{0}\}$ such that*

$$(64) \quad A^{\ell_1} \gamma - A^{\ell_2} \gamma + A^{\ell_3} \gamma - \gamma = 0,$$

without vanishing proper sub-sum, is finite.

Proof. There exists a decomposition of $E = \mathbb{C}^\rho$ into vectorial subspaces $\mathbb{C}^\rho = \bigoplus_k E_k$ which are simultaneously invariant by \tilde{A}_i , $i = 1, 2$, and such that there is a basis B_k in which \tilde{A}_i restricted to E_k is represented in a triangular form with an eigenvalue of \tilde{A}_i on the diagonal.

This follows from the fact that the commuting matrices A_i have a common non trivial space W of eigenvectors, and then by an induction on the dimension of the vector space, applying the induction hypothesis to the action of the quotient map of \tilde{A}_i on E/W .

For $\gamma \in \mathbb{Z}^\rho \setminus \{0\}$, there is k_0 such that the component γ_0 of γ in E_{k_0} is $\neq 0$. Let δ_0 be the dimension of E_{k_0} . In the basis $B_{k_0} = \{e_{k_0,1}, \dots, e_{k_0,\delta_0}\}$ of E_{k_0} , we denote the coordinates of γ_0 by $(\gamma_0^1, \dots, \gamma_0^{\delta_0})$. There is $\delta'_0 \in \{1, \dots, \delta_0\}$ such that $\gamma_0^i = 0$, $\forall i < \delta'_0$, and $\underline{v}_0 := \gamma_0^{\delta'_0} \neq 0$.

Due to the triangular form, for $j = 1, 2$, we have $A_j^\ell \gamma_0 = \alpha_{k_0,j}^\ell \gamma_0^{\delta'_0} e_{k_0,\delta'_0} + \zeta(j, \ell)$, $\forall \ell \in \mathbb{Z}$, where $\alpha_{k_0,j}$ is an eigenvalue of A_j and where $\zeta(j, \ell)$ belongs to the subspace generated by $\{e_{k_0,\delta'_0+1}, \dots, e_{k_0,\delta_0}\}$.

Using the notation $\underline{\alpha}_u^\ell = \alpha_{u,1}^{\ell_1} \alpha_{u,2}^{\ell_2}$, if $\alpha_{u,1}$ (resp. $\alpha_{u,2}$) is an eigenvalue of A_1 (resp. A_2), if (64) holds, then $(\underline{\alpha}_{k_0}^{\ell_1} - \underline{\alpha}_{k_0}^{\ell_2} + \underline{\alpha}_{k_0}^{\ell_3}) \underline{v}_0 = \underline{v}_0$. This equation is still without vanishing proper sub-sum, because of the assumption of total ergodicity. By Theorem 6.2 applied to the multiplicative group (of finite rank) generated by $\alpha_{k_0,j}$, $j = 1, 2$, the number of solutions of the previous equation is finite. Hence the result, since k_0 takes a finite number of values. \square

Random walks and quenched CLT

Our aim is to replace the r.f. of i.i.d. variables $(X_\ell, \ell \in \mathbb{Z}^2)$ discussed in Section 2 by the random field generated by an observable f on a torus \mathbb{T}^ρ under the action of commuting automorphisms.

More precisely, we consider $\underline{\ell} \mapsto A^{\underline{\ell}}$ a totally ergodic \mathbb{Z}^2 -action by algebraic automorphisms of \mathbb{T}^ρ , $\rho > 1$, defined by commuting $\rho \times \rho$ matrices A_1, A_2 with integer entries, determinant ± 1 such that the eigenvalues of $A^{\underline{\ell}} = A_1^{\ell_1} A_2^{\ell_2}$ are $\neq 1$, if $\underline{\ell} = (\ell_1, \ell_2) \neq (0, 0)$.

The composition with a function f defined on \mathbb{T}^ρ is denoted by $A^{\underline{\ell}}f$ as well as $T^{\underline{\ell}}f$. We consider the random field $(X_{\underline{\ell}} = A^{\underline{\ell}}f, \underline{\ell} \in \mathbb{Z}^2)$, with $f \in AC_0(\mathbb{T}^\rho)$.

A sufficient condition for f with 0 integral to be in $AC_0(\mathbb{T}^\rho)$ is $|\hat{f}(\underline{k})| = O(\|\underline{k}\|^{-\beta})$, with $\beta > \rho$.

For a.e. ω the following asymptotic variance exists:

$$\lim_n \frac{1}{n \log n} \left\| \sum_{k=1}^n A^{Z_k(\omega)} f \right\|_2^2 = C_0 \sum_{\underline{k} \in \mathbb{Z}^d} \langle T^{\underline{k}} f, f \rangle,$$

where the constant C_0 is defined in Subsection 1.2.

The following quenched FCLT extends for the torus the CLT proved in [8]. Remark that the CLT is proved therein for a general compact abelian group. The extension to a functional version of the CLT holds in this general case when f is a trigonometric polynomial.

Theorem 6.4. *Let (Z_n) be a 2-dimensional reduced centered random walk with a finite moment of order 2 and let f be a real function in $AC_0(\mathbb{T}^\rho)$ with spectral density φ_f and $\varphi_f(\underline{0}) \neq 0$. Denoting by $S_n^\omega(f) := \sum_{k=1}^n A^{Z_k(\omega)} f$ the sums along the r.w., the process $\left(\frac{1}{\sqrt{n \log n}} S_{[nt]}^\omega(f) \right)_{t \in [0,1]}$ satisfies the FCLT for a.e. ω .*

Proof. 1) *Convergence of the finite dimensional distributions*

1a) First suppose that f is a trigonometric polynomial: $f = \sum_{\underline{k} \in \Lambda} c_{\underline{k}}(f) \chi_{\underline{k}}$, where $(\chi_{\underline{k}}, \underline{k} \in \Lambda)$ is a finite set of characters on \mathbb{T}^ρ and χ_0 the trivial character.

We use Proposition 3.5: (51) follows from (26) and Lemma 1.10. For (52), we have to show

$$(65) \quad \mathcal{C}^{(r)}(\sum_{\underline{\ell} \in \mathbb{Z}^2} w_n(\omega, \underline{\ell}) T^{\underline{\ell}} f) = o((n \log n)^{r/2}), \forall r \geq 3, \text{ for a.e. } \omega.$$

We apply Theorem 3.3. Let us check (48). For r fixed, the function $(\underline{n}_1, \dots, \underline{n}_r) \rightarrow m_f(\underline{n}_1, \dots, \underline{n}_r) := \int_X T^{\underline{n}_1} f \cdots T^{\underline{n}_r} f d\mu$ takes a finite number of values, since m_f is a sum with coefficients 0 or 1 of the products $c_{k_1} \cdots c_{k_r}$ with k_j in a finite set. The cumulants of a given order take also a finite number of values according to (42).

Therefore, since mixing of all orders implies $\lim_{\max_{i,j} \|\underline{\ell}_i - \underline{\ell}_j\| \rightarrow \infty} \mathcal{C}(T^{\underline{\ell}_1} f, \dots, T^{\underline{\ell}_r} f) = 0$ by Proposition 3.4, there is M_r such that $\mathcal{C}(T^{\underline{\ell}_1} f, \dots, T^{\underline{\ell}_r} f) = 0$ if $\max_{i,j} \|\underline{\ell}_i - \underline{\ell}_j\| > M_r$. The end of the proof is then like in Theorem 4.1.

1b) For $f \in AC_0(\mathbb{T}^\rho)$, using Proposition 6.1 and Lemma 3.6, the convergence follows by approximation of f by a sequence of trigonometric polynomials f_L in such a way that $\lim_L \|\varphi_f - \varphi_{f_L}\| = 0$.

2) *Moment of order 4 and tightness*

We use Proposition 5.2. Taking into account Remark 5.3, it suffices for the tightness to take for f a character and show that the bounds are independent of the character.

Let χ_v be a character on the torus \mathbb{T}^ρ , $\chi_v : x \rightarrow \exp(2\pi i \langle v, x \rangle)$, where $v \in \mathbb{Z}^\rho \setminus \{0\}$.

For an interval $J = [b, b + k] \subset [1, n]$, we have:

$$\|\sum_{i=b}^{b+k} A^{Z_i} \chi_v\|_4^4 = \#\{(i_1, i_2, i_3, i_4) \in J^4 : (A^{Z_{i_1}} - A^{Z_{i_2}} + A^{Z_{i_3}} - A^{Z_{i_4}}) v = 0\}.$$

This number is less than $2G^2(\omega, b, k) + H(\omega, b, k)$, with

$$\begin{aligned} G(\omega, b, k) &:= \#\{(i_1, i_2) \in J^2 : (A^{Z_{i_1}} - A^{Z_{i_2}}) \underline{v} = 0\}, \\ H(\omega, b, k) &:= \#\{(i_1, i_2, i_3, i_4) \in J^4 : (A^{Z_{i_1}} - A^{Z_{i_2}} + A^{Z_{i_3}} - A^{Z_{i_4}}) \underline{v} = 0\}, \end{aligned}$$

where above in H we count the number of solutions without vanishing proper sub-sums.

By assumption of total ergodicity, if $(A^{Z_{i_1}} - A^{Z_{i_2}}) \underline{v} = 0$, then $Z_{i_1} = Z_{i_2}$, so that G is the number of self-intersections of the r.w. on $[b, b + k]$: $G(\omega, b, k) = V(\omega, [b, b + k]) = V_k(\theta^b \omega)$. A bound of these quantities is given by (20).

For H , by Corollary 6.3, there is a finite set F (independent of \underline{v}) such that

$$H(\omega, b, k) \leq \#\{(i_1, i_2, i_3, i_4) \in J^4 : (Z_{i_1}(\omega) - Z_{i_4}(\omega), Z_{i_2}(\omega) - Z_{i_4}(\omega), Z_{i_3}(\omega) - Z_{i_4}(\omega)) \in F\}.$$

Therefore, with the notation (24), $H(\omega, b, k) \leq \sum_{(\underline{\ell}_1, \underline{\ell}_2, \underline{\ell}_3) \in F} W_k(\theta^b \omega, \underline{\ell}_1, \underline{\ell}_2, \underline{\ell}_3)$.

By (25) in Lemma 1.5, there exists a positive integrable function C_3 such that $W_n(\omega, \underline{\ell}_1, \underline{\ell}_2, \underline{\ell}_3) \leq C_3(\omega) n (\log n)^5$, $\forall n \geq 1$, which implies $H(\omega, b, k) \leq (\text{Card } F) C_3(\theta^b \omega) k (\log k)^5$. Remark that the bounds do not depend on the character, but only on the matrices A_1, A_2 .

Since G and H are super-additive (Condition (55)), the tightness property follows now from Proposition 5.2 with $\gamma = 5$. \square

Remarks 6.5. 1) An analogous result is valid for any transient random walk in dimension $d \geq 1$, with the standard normalisation by \sqrt{n} . In this case, if the observable is non null a.e., the asymptotic variance for the sums along the r.w. is different from 0 (cf. Subsection 1.2).

2) For automorphisms of a torus \mathbb{T}^ρ , in the recurrent 2-dimensional model studied above, if f satisfies the regularity condition $|c_f(\underline{k})| = O(\|\underline{k}\|^{-\beta})$, with $\beta > \rho$, then the asymptotic variance is given by $\varphi_f(0)$ and it is null, if and only if f is a mixed coboundary: there are continuous functions u_1, u_2 such that $f = (I - A_1)u_1 + (I - A_2)u_2$ (cf. [8]).

7. Exponential mixing of all orders

7.1. FCLT and exponential mixing of all orders.

Our last example is given by commuting translations on homogeneous spaces. It relies on recent results on the exponential mixing of all orders for flows on homogeneous spaces shown in [3] and their application to the CLT in [4]. Closely following the latter reference, we recall first the notion of exponential mixing of all orders for a group action by measure preserving transformations.

Exponential mixing of all orders

Let \mathcal{H} be a group with a left invariant distance d . Let $h \rightarrow T^h$ be a homomorphism of \mathcal{H} in the group of measure preserving invertible transformations of a probability space (E, \mathcal{B}, μ) . We denote by \mathcal{A} a sub-algebra in $L^\infty(E, \mu)$ which is \mathcal{H} -invariant. Let $\mathcal{N} = (N_s)$ be a family of semi-norms on \mathcal{A} , indexed by positive integers s .

Hypothesis 7.1. *The following conditions are assumed to hold for all $s > 1$ and all $f, g \in \mathcal{A}$ with constant factors depending only on s (all denoted by C_s):*

- 1) $N_s(f) \leq C_s N_{s+1}(f)$; 2) $\|f\|_{L^\infty} \leq C_s N_s(f)$; 3) $N_s(fg) \leq C_s N_{s+1}(f) N_{s+1}(g)$;
- 4) *there exists $\zeta_s > 0$ such that $N_s(T^h f) \leq C_s e^{\zeta_s d(h,e)} N_s(f)$, $\forall h \in \mathcal{H}$.*

Let $r \geq 2$ be an integer. For $\underline{H} = (h_1, \dots, h_r) \in \mathcal{H}^r$, we set $d_r(\underline{H}) := \min_{i \neq j} d(h_i, h_j)$.

Definition 7.2. *We say that the \mathcal{H} -action on (E, μ) is exponentially mixing of order r , with respect to d and $(\mathcal{A}, \mathcal{N})$, if there exist constants $C_{r,s}, \delta_r > 0$ and an integer $s_r > 0$ such that for all $s \geq s_r$ and $f_1, \dots, f_r \in \mathcal{A}$,*

$$(66) \quad \left| \mu\left(\prod_{i=1}^r T^{h_i} f_i\right) - \prod_{i=1}^r \mu(f_i) \right| \leq C_{r,s} e^{-\delta_r d_r(\underline{H})} \prod_{i=1}^r N_s(f_i),$$

for all $\underline{H} = (h_1, \dots, h_r) \in \mathcal{H}^r$. The constant C depends only on r and s .

We may assume that ζ_s is increasing with s , δ_r decreasing with r and $\delta_r < r\zeta_s$, $\forall r, s$.

In what follows, we will consider, for $d \geq 2$, a measure preserving \mathbb{Z}^d -action on a probability space (E, \mathcal{B}, μ) generated by d commuting invertible maps T_1, \dots, T_d . Therefore the group \mathcal{H} in Definition 7.2 is going to be the group \mathbb{Z}^d still denoted also by \mathcal{H} .

If (Z_n) is a random walk on \mathbb{Z}^d , then we get a random walk $(T^{Z_n(\omega)})$ on the group of measure preserving invertible transformations on (E, \mathcal{B}, μ) .

With a distance on \mathbb{Z}^d associated to a norm equivalent to the Euclidean norm, the volume of a big ball is of order the number of integral points in the ball. It is important to relate this distance to the distance d of Definition 7.2. We will assume that the action satisfies:

Hypothesis 7.3. *For an \mathcal{H} -invariant sub-algebra \mathcal{A} in $L^\infty(E, \mu)$ and a family $\mathcal{N} = (N_s)$ of semi-norms on \mathcal{A} , we assume that the \mathbb{Z}^d -action on (E, \mathcal{B}, μ) is exponentially mixing of order r for every $r \geq 2$ in the sense of Definition 7.2 with a distance d equivalent to the Euclidean distance.*

With this assumption, for simplicity of notation we can assume that the distance d in Definition 7.2 applied to \mathbb{Z}^d is the Euclidean distance on \mathbb{Z}^d .

Spectral density and cumulants

Let f be a centered function in \mathcal{A} . Its spectral density is $\varphi_f = \sum_{\underline{\ell} \in \mathbb{Z}^d} a_{\underline{\ell}} e^{2\pi i \langle \underline{\ell}, \underline{t} \rangle}$, with $a_{\underline{\ell}} = \langle T^{\underline{\ell}} f, f \rangle$. The absolute summability $\sum_{\underline{\ell}} |a_{\underline{\ell}}| < \infty$ is a consequence of (66) for $r = 2$.

Moreover, there exists s_2 and a constant C such that $\|\varphi_f\|_\infty \leq C N_{s_2}(f)^2$.

Let $J = [b, b+k] \subset [1, n]$. With the notation of 1.2.2, we have

$$\int \left| \sum_{j \in J} T^{Z_j(\omega)} f \right|^2 d\mu = \int_{\mathbb{T}^d} \left| \sum_{j \in J} e^{2\pi i \langle Z_j(\omega), \underline{t} \rangle} \right|^2 \varphi_f(\underline{t}) d\underline{t} \leq \sum_{\underline{\ell} \in \mathbb{Z}^d} |a_{\underline{\ell}}| V(\omega, J, \underline{\ell}) =: G(\omega, b, k).$$

For a fixed ω , the bound $G(\omega, b, k)$ is super-additive. By (43), we have

$$(67) \quad \mathbb{E}_\mu \left(\left| \sum_{j \in J} T^{Z_j(\omega)} f \right|^4 \right) = 3 \left(\int \left(\sum_{j \in J} T^{Z_j(\omega)} f \right)^2 d\mu \right)^2 + C^{(4)} \left(\sum_{j \in J} T^{Z_j(\omega)} f \right).$$

The next proposition shows that the cumulants $\mathcal{C}(T^{h_1} f, \dots, T^{h_r} f)$ are small for all well-separated r -tuples $\underline{H} = (h_1, \dots, h_r)$. The constants δ_r and ζ_r are those of Definition 7.2.

Proposition 7.4. *(Proposition 6.1 in [4]) For $r \geq 2$, let Q be a partition of $\{1, 2, \dots, r\}$ with $|Q| \geq 2$ and let s be an integer $> s_r + r$. Then, if $\underline{H} \in \Delta_Q(\alpha, \beta)$ with $0 < \alpha < \beta$, for f centered in \mathcal{A} we have*

$$(68) \quad |\mathcal{C}(T^{h_1} f, \dots, T^{h_r} f)| \leq C e^{-(\delta_r \beta - r \alpha \zeta_s)} N_s(f)^r.$$

where the constant C depends only on r and s .

Proof. ($r = 4$) Let us give the proof for $r = 4$ and in the case of the configurations \underline{H} of type $S_3 = \bullet \bullet \bullet \bullet$, that is (up to a permutation): $\{\{h_i, h_k\}, \{h_\ell\}, \{h_r\}\}$ with, for some j , $d(h_i, h_k) < 3\beta_j$ and $d(h_\ell, \{h_i, h_k\}) > \beta_{j+1}$, $d(h_r, \{h_i, h_k\}) > \beta_{j+1}$, $d(h_\ell, h_r) > \beta_{j+1}$.

We may write the formula for the cumulants in the following way:

$$\begin{aligned} \mathcal{C}(T^{h_i} f, T^{h_k} f, T^{h_\ell} f, T^{h_r} f) &= (A) - (B) - (C) - (D), \text{ with} \\ (A) &= \mathbb{E}(T^{h_i} f T^{h_k} f T^{h_\ell} f T^{h_r} f), \quad (B) = \mathbb{E}(T^{h_i} f T^{h_k} f) \mathbb{E}(T^{h_\ell} f T^{h_r} f), \\ (C) &= \mathbb{E}(T^{h_i} f T^{h_\ell} f) \mathbb{E}(T^{h_k} f T^{h_r} f), \quad (D) = \mathbb{E}(T^{h_i} f T^{h_r} f) \mathbb{E}(T^{h_k} f T^{h_\ell} f). \end{aligned}$$

We use the exponential mixing of order 2 for $(B), (C), (D)$ and of order 3 for (A) . More precisely, we have $(A) = \mathbb{E}((f T^{h_k - h_i} f) T^{h_\ell - h_i} f T^{h_r - h_i} f)$ and

$$N_s(f T^{h_k - h_i} f) \leq C N_{s+1}(f) N_{s+1}(T^{h_k - h_i} f) \leq C e^{3\zeta_s \beta_j} N_{s+1}(f) N_{s+1}(f).$$

Therefore, by the 3-mixing applied to $f T^{h_k - h_i} f, T^{h_\ell - h_i} f, T^{h_r - h_i} f$, we have

$$(A) \leq C e^{3\zeta_s \beta_j} e^{-\delta_r \beta_{j+1}} N_{s+1}(f)^2 N_s(f)^2. \quad \square$$

Let us now fix $\gamma > 0$. We define β_j by $\beta_0 = 0$ and $\beta_{j+1} = 3r\beta_j\zeta_s/\delta_r + \gamma$, for $j > 0$.

As $\delta_r < r\zeta_s$, we have $\beta_{j+1} > 3\beta_j$. Moreover $\beta_r = \gamma \sum_{j=0}^{r-1} (3r\zeta_s/\delta_r)^j =: \gamma c_{r,s}$.

Let $\nu = \sum_{\underline{\ell}} \nu(\underline{\ell}) \delta_{\underline{\ell}}$ be a positive finite measure on $\mathcal{H} = \mathbb{Z}^d$ with mass $\|\nu\|$. Let $B(h, R)$ be the euclidean ball of center h and radius R . The following bound on cumulants is given in [4] (cf. Proposition 7.4 above and (45)). The first term at right in (69) comes from the clustered configurations and the second term from the well separated configurations. The constants s_r and δ_r are introduced in Definition 7.2.

Proposition 7.5. *(Proposition 5.2 in [4]) For every $r \geq 3$ and $s \geq s_r + r$, there exist $C_{r,s}$ and $c_{r,s} > 0$ (not depending on ν) such that, for all $\gamma > 0$ and $f \in \mathcal{A}$,*

$$(69) \quad |\mathcal{C}^{(r)}(\nu * f)| \leq C_{r,s} \left(\int_{\mathcal{H}} \nu(B(h, c_{r,s}\gamma))^{r-1} d\nu(h) + e^{-\delta_r \gamma} \|\nu\|^r \right) N_s(f)^r.$$

Under the assumption of exponential mixing of all orders, it is shown in [4] that the CLT holds for $\nu_n * f$, when $f \in \mathcal{A}$ and (ν_n) a sequence of measures satisfying a certain condition.

In our framework, ν is the measure $\nu_n^\omega = \sum_{j=0}^{n-1} \delta_{Z_j(\omega)}$, where $(Z_j)_{j \geq 0}$ is a r.w. on \mathbb{Z}^2 . Its mass is $\|\nu_n^\omega\| = n$. The convolution “ $\nu * f$ ” in (69) for ν_n^ω means “ $\sum_{j=0}^{n-1} f(T^{Z_j(\omega)} \cdot)$ ”.

Let us assume that the r.w. Z is a centered random walk on \mathbb{Z}^2 with moments of order 2. By (23) (cf. Remark 1.4 in 1.2.2), it holds with a constant $K(\omega)$ finite for a.e. ω , uniformly in

$h \in \mathcal{H}$:

$$\nu_n^\omega(B(h, c\gamma)) = \#\{j \leq n : Z_j(\omega) \in B(h, c\gamma)\} \leq \sup_{\underline{\ell}} w_n(\omega, \underline{\ell}) \text{Card}(B(h, c\gamma)) \leq K(\omega)(c\gamma)^2 \log^2 n.$$

Due to Hypothesis 7.3 and by (69), this implies that there is a constant C_1 (depending only on r and s) such that

$$|\mathcal{C}^{(r)}(\nu_n^\omega * f)| \leq C_1 (K(\omega)^{r-1} (\log n)^{2(r-1)} (c\gamma)^{2(r-1)} n + e^{-\delta_r \gamma} n^r) N_s(f)^r.$$

Taking $\gamma = \frac{r-1}{\delta_r} \log n$, the first term above gives the bound $(c \frac{r-1}{\delta_r})^{2(r-1)} (\log n)^{4(r-1)} n$ and the second term is n . It follows, for a constant C_2 depending on r and s :

$$|\mathcal{C}^{(r)}(\nu_n^\omega * f)| \leq C_2 N_s(f)^r K(\omega)^{r-1} (\log n)^{4(r-1)} n.$$

Likewise for an interval $J = [b, b+k]$ and $\nu_J^\omega = \sum_{j \in J} \delta_{Z_j(\omega)}$, we get

$$|\mathcal{C}^{(r)}(\nu_J^\omega * f)| \leq C_2 N_s(f)^r K(\theta^b \omega)^{r-1} (\log k)^{4(r-1)} k.$$

So we get the same type of upper bound for the cumulants as for the automorphisms of the torus. Therefore we have convergence of the finite dimensional distributions and tightness.

For a \mathbb{Z}^2 -action satisfying the exponential mixing condition on an algebra of functions as presented at the beginning of this section, we can state now a functional version of a CLT result for the summation along a random walk.

Theorem 7.6. *Let (Z_n) be a 2-dimensional aperiodic centered random walk with moments of order 2. Let f be a real centered function in \mathcal{A} with spectral density φ_f such that $\varphi_f(\mathbf{0}) \neq 0$. Under Hypothesis 7.3, denoting by $S_n^\omega(f) := \sum_{k=1}^n T^{Z_k(\omega)} f$ the sums along the r.w., the process $\left(\frac{1}{\sqrt{n \log n}} S_{[nt]}^\omega(f) \right)_{t \in [0,1]}$ satisfies the FCLT for a.e. ω .*

Remarks 7.7. 1) As suggested in a referee's remark, the above result can be extended to a space of functions larger than \mathcal{A} . For $r = 4$, by Proposition 7.5, there is s_4 such that (69) holds for $g \in \mathcal{A}$. Let \mathcal{A}_1 be the space obtained as closure of \mathcal{A} for the norm N_{s_4} .

Let us sketch a proof of the extension of Theorem 7.6 to functions $f \in \mathcal{A}_1$. By 7.2 we still have fast decorrelation for $f \in \mathcal{A}_1$. We can use Lemma 3.6 for $f \in \mathcal{A}_1$ and (g_n) a sequence in \mathcal{A} such that $\|f - g_n\|_{N_{s_4}} \rightarrow 0$ (hence also $\|\varphi_{f-g_n}\|_\infty \rightarrow 0$ by (66) and hypothesis 7.1). Therefore f in \mathcal{A}_1 satisfies the convergence of the finite dimensional distributions, as the functions g_n in \mathcal{A} . Moreover, by (69) the estimate for the cumulant of order 4, which is satisfied by functions in \mathcal{A} , still holds for \mathcal{A}_1 and this, as above, implies tightness.

2) The result is formulated for \mathbb{Z}^2 , but an analogous result with a normalisation in \sqrt{n} can be proved for a transient random walk in dimension ≥ 3 with the same method. To do it, we use the bound (67) for the 4th moment. In the transient case, the first term behaves like n^2 because of (14). The second term behaves as $n(\log n)^{3(r-1)}$ by using Prop. 7.3 and Theorem 13 in Erdos-Taylor [15] (which can be extended to a r.w. as for the upper bound in [14], according to Remark 1.4). Observe also that Proposition 5.2 can be adapted to the normalisation corresponding to the transient case.

7.2. Translations on homogeneous spaces.

The following example is an action which is exponentially mixing of all orders on an algebra \mathcal{A} of functions according to [3].

We take the group $G = \mathrm{SL}(n, \mathbb{R})$ and a lattice Γ in G , i.e., a discrete subgroup such that G/Γ has a finite volume for the measure μ induced by the Haar measure of G , for example $\mathrm{SL}(n, \mathbb{Z})$. The space E is the quotient G/Γ . The action on E will be given by left multiplication $g\Gamma \rightarrow hg\Gamma$ where h is in the diagonal subgroup D of G .

The algebra \mathcal{A} in the example is the algebra of C^∞ -functions with compact support on G/Γ , and \mathcal{N} is a family of Sobolev norms as in [3].

Let us explicit the distance in the example.

Left invariant pseudo-metric on $G = \mathrm{SL}(n, \mathbb{R})$

Recall that there is a distance on D , induced by a canonical left invariant distance on G . The left invariant distance d_G defined on G is comparable on D to the ‘pseudo-metric’ $\delta(\cdot, \cdot)$ defined as follows:

For $A \in G$, let $\|A\| = \max_{x \neq 0} \frac{\|Ax\|}{\|x\|}$ be the norm of the $n \times n$ matrix A as operator on \mathbb{R}^n endowed with the euclidian norm. Since the determinant is 1, A has an eigenvalue of modulus ≥ 1 , which implies $\|A\| \geq 1$. For $A, B \in G$, we put $\delta(A, B) = \log(\|A^{-1}B\|)$.

Clearly, $\delta(A, B) \geq 0$, the triangular inequality is satisfied by sub-multiplicativity of the operator norm, and δ is left invariant on G . If $\|A\| = 1$, the iterates of A are bounded, so all eigenvalues must have a modulus ≤ 1 . As the determinant is 1, the modulus of the eigenvalues is 1. Now considering the Jordan form of A over \mathbb{C} , it must be diagonal. Finally we conclude that $\{A : \|A\| = 1\}$ is the orthogonal group in G .

We take two elements $A_1 = \exp(U_1)$, $A_2 = \exp(U_2)$, where U_1, U_2 , in the sub-algebra \mathfrak{D} corresponding to D in the Lie algebra \mathfrak{G} of G , are such that U_1, U_2 generate a 2-dimensional vectorial space of \mathfrak{D} . The group $(A_1^{\ell_1} A_2^{\ell_2}, (\ell_1, \ell_2) \in \mathbb{Z}^2)$ yields a totally ergodic action on G/Γ .

For instance in $\mathrm{SL}(3, \mathbb{R})$, we can take with $a_1, a_2 > 1$: $A_1 = \begin{pmatrix} a_1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & a_1^{-1} \end{pmatrix}$, $A_2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & a_2 & 0 \\ 0 & 0 & a_2^{-1} \end{pmatrix}$.

The distance $d_G(A_1^{\ell_1} A_2^{\ell_2}, Id)$ is equivalent to $\|\ell\| = (\ell_1^2 + \ell_2^2)^{\frac{1}{2}}$. The measure $\nu_n(B(Id, R))$ is the counting measure with some weight applied to the ball, therefore up to the weight it is the number of elements of the form $h_1^{\ell_1} h_2^{\ell_2}$ in the ball, and finally the (weighted) number of integers $\ell = (\ell_1, \ell_2)$ of norm $\leq R$.

By what precedes, Hypothesis 7.3 is satisfied and Theorem 7.6 yields a functional CLT in the class of centered compactly supported C^∞ -functions for the action of a 2-dimensional random walk on the diagonal subgroup on G/Γ .

A result analogous to Theorem 7.6 holds for the sums along a transient random walk: the only change is the estimate of the number of self-intersection (normalisation by \sqrt{n}).

In Theorem 7.6, in particular in the example provided by homogeneous spaces, as in the CLT in [4], the statement says nothing about the non-nullity of the variance, for a 2-dimensional recurrent r.w. The question of degeneracy of the asymptotic variance (for a recurrent r.w.) is the same as for the sums over squares: it depends of the nullity of $\varphi(\underline{0})$. This contrasts with the action of commuting automorphisms of a torus, for which (as recalled in Remark 6.5.2) there is a description of the degenerate case in terms of mixed coboundaries.

In the case of a transient random walk on \mathbb{Z}^d , as noticed in Subsection 1.2, if the observable is non null a.e., the asymptotic variance for the sums along a transient r.w. is different from 0.

REFERENCES

- [1] A. Ayer, C. Liverani and M. Stenlund: Quenched CLT for random toral automorphism, *Discrete Contin. Dyn. Syst.* 24 (2009), no. 2, p. 331-348.
- [2] P. Billingsley: *Convergence of probability measures*, 2d edition. John Wiley & Sons, Inc., NY, 1999.
- [3] M. Björklund, M. Einsiedler, A. Gorodnik: Quantitative multiple mixing. *J. Eur. Math. Soc. (JEMS)* 22 (2020), no. 5, 1475-1529.
- [4] M. Björklund, A. Gorodnik: Central limit theorems for group actions which are exponentially mixing of all orders. *J. Anal. Math.* 141 (2020), no. 2, 457-482.
- [5] E. Bolthausen: A central limit theorem for two-dimensional random walks in random sceneries, *Ann. Probab.* 17, no. 1, 108-115 (1989).
- [6] K. L. Chung, W. H. J. Fuchs: On the distribution of values of sums of random variables. *Mem. Amer. Math. Soc.* 6 (1951), 12 pp.
- [7] K. L. Chung, D. Ornstein: On the recurrence of sums of random variables. *Bull. Amer. Math. Soc.* 68 (1962), 30-32.
- [8] G. Cohen, J.-P. Conze: CLT for random walks of commuting endomorphisms on compact abelian groups, *J. Theoret. Probab.* 30 (2017), no. 1, 143-195.
- [9] G. Cohen, J.-P. Conze: Almost mixing of all orders and CLT for some \mathbb{Z}^d -actions on subgroups of $\mathbb{F}_p^{\mathbb{Z}^d}$ (2016), <https://arxiv.org/abs/1609.06484>
- [10] G. Cohen, J.-P. Conze: On the quenched functional CLT in $2d$ -random sceneries, examples (2019), <https://arxiv.org/abs/arXiv:1908.03777>
- [11] H. Cohen: *A course in computational algebraic number theory*. Graduate Texts in Mathematics, 138. Springer-Verlag, Berlin (1993). doi: 10.1007/978-3-662-02945-9
- [12] J.-P. Conze, S. Le Borgne: Quenched central limit theorem for random walks with a spectral gap, *CRAS*, vol. 349, no 13-14, juillet 2011, 801-805.
- [13] G. Deligiannidis, S.A. Utev: Computation of the asymptotics of the variance of the number of self-intersections of stable random walks using the Wiener-Darboux theory. (Russian) *Sibirsk. Mat. Zh.* 52 (2011), no. 4, 809-822; translation in *Sib. Math. J.* 52 (2011), no. 4, 639-650.
- [14] A. Dembo, Y. Peres, J. Rosen, O. Zeitouni: Thick points for planar Brownian motion and the Erdős-Taylor conjecture on random walk. *Acta Math.* 186 (2001), no. 2, 239-270.
- [15] P. Erdős and S. J. Taylor: Some problems concerning the structure of random walk paths. *Acta Math. Acad. Sci. Hungar.* 11 (1960), 137-162.
- [16] J.D. Esary, F. Proschan and D.W. Walkup: Association of Random Variables with Applications, *Ann. Math. Stat.* 38 (1967) no. 5, 1466-1474.
- [17] J.-H. Evertse, H.P. Schlickewei, W.M. Schmidt: Linear equations in variables which lie in a multiplicative group, *Ann. of Math.* 155, no. 3, 807-836 (2002). doi: 10.2307/3062133
- [18] H. Kesten, F. Spitzer: A Limit Theorem Related to a New Class of Self Similar Processes. *Z. Wahrsch. Verw. Gebiete.* 50, 5-25 (1979). doi: 10.1007/BF00535672
- [19] V.P. Leonov: The use of the characteristic functional and semi-invariants in the ergodic theory of stationary processes. *Dokl. Akad. Nauk SSSR* 133, 523-526; transl. as *Soviet Math. Dokl.* 1, 878-881 (1960).
- [20] V.P. Leonov: On the central limit theorem for ergodic endomorphisms of compact commutative groups, *Dokl. Akad. Nauk SSSR* 135, 258-261 (1960).
- [21] T.M. Lewis: A law of the iterated logarithm for random walk in random scenery with deterministic normalizers, *J. Theoret. Probab.* 6, no. 2, 209-230 (1993). doi: 10.1007/BF01047572
- [22] F. Móricz: Moment inequalities and the strong laws of large numbers, *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* 35 (1976), no. 4, 299-314.

- [23] C.M. Newman, A.L. Wright: An invariance principle for certain dependent sequences, *Ann. Probab.* 9 (1981), no. 4, 671-675.
- [24] F. Pène: Planar Lorentz process in a random scenery, *Ann. IHP, Probab. Stat.*, 45 (2009), 818-839.
- [25] F. Pène: Self-intersections of trajectories of the Lorentz process, *Discrete Contin. Dyn. Syst.* 34 (2014), 11, 4781-4806.
- [26] H.P. Schlickewei: S-unit equations over number fields, *Invent. Math.* 102 (1990), 95-107.
- [27] F. Spitzer: Principles of random walk. The University Series in Higher Mathematics D. Van Nostrand Co., Inc., Princeton, N.J.-Toronto-London (1964). doi: 10.1007/978-1-4757-4229-9

GUY COHEN,
DEPT. OF ELECTRICAL ENGINEERING,
BEN-GURION UNIVERSITY, ISRAEL

Email address: guycohen@bgu.ac.il

JEAN-PIERRE CONZE,
IRMAR, CNRS UMR 6625,
UNIVERSITY OF RENNES I, CAMPUS DE BEAULIEU, 35042 RENNES CEDEX, FRANCE

Email address: conze@univ-rennes1.fr