



HAL
open science

Weighted- L^2 polynomial approximation in C

Séverine Biard, John Erik Fornæss, Jujie Wu

► **To cite this version:**

Séverine Biard, John Erik Fornæss, Jujie Wu. Weighted- L^2 polynomial approximation in C . Transactions of the American Mathematical Society, 2020, 373 (2), pp.919-938. 10.1090/tran/7935 . hal-03207780

HAL Id: hal-03207780

<https://hal.science/hal-03207780v1>

Submitted on 31 Oct 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

WEIGHTED- L^2 POLYNOMIAL APPROXIMATION IN \mathbb{C}

SÉVERINE BIARD, JOHN ERIK FORNÆSS, AND JUJIE WU

ABSTRACT. We study the density of polynomials in $H^2(\Omega, e^{-\varphi})$, the space of square integrable holomorphic functions in a bounded domain Ω in \mathbb{C} , where φ is a subharmonic function. In particular, we prove that the density holds in Carathéodory domains for any subharmonic function φ in a neighborhood of $\bar{\Omega}$. In non-Carathéodory domains, we prove that the density depends on the weight function, giving examples.

CONTENTS

| | |
|--------------------------|----|
| 1. Introduction | 1 |
| 2. Proof of Theorem 1.3 | 4 |
| 3. Proof of Theorem 1.6 | 8 |
| 4. Proof of Theorem 1.7 | 14 |
| 5. Example 1.10 | 18 |
| 6. Proof of Theorem 1.11 | 20 |
| References | 22 |

1. INTRODUCTION

Let Ω be a domain in \mathbb{C} . We denote by $L^2(\Omega, e^{-\varphi})$ the space of measurable functions f such that

$$\|f\|_{\Omega, \varphi}^2 := \int_{\Omega} |f|^2 e^{-\varphi} d\lambda < +\infty,$$

Date: Received by the editors June 19, 2018, and, in revised form, Feb 27, 2019.

2000 Mathematics Subject Classification. Primary 30D15, 30E10; Secondary 32A10, 32E30, 32W05.

Key words and phrases. Carathéodory domain, weighted L^2 -estimates, polynomial approximation, moon-shaped domain .

Jujie Wu is the corresponding author.

The first author was supported in part by Rannis-grant 152572-051. The second author was supported by the Norwegian Research Council grant 240569. The third author was supported by the Norwegian Research Council grant 240569 and NSFC grant 11601120.

where φ is a measurable function on Ω , and $d\lambda$ is the Lebesgue measure. Let $H^2(\Omega, e^{-\varphi})$ (resp. $H^2(\overline{\Omega}, e^{-\varphi})$) be the space of holomorphic functions on a domain Ω (resp. holomorphic functions on a neighborhood of $\overline{\Omega}$), which are in $L^2(\Omega, e^{-\varphi})$, that is

$$H^2(\Omega, e^{-\varphi}) := \mathcal{O}(\Omega) \cap L^2(\Omega, e^{-\varphi}).$$

Recall that a Carathéodory domain Ω is a simply-connected bounded planar domain whose boundary $\partial\Omega$ is also the boundary of an unbounded domain. An unbounded domain Ω is said to be Carathéodory if its image under the map $z \mapsto (z - z^0)^{-1}$ (z^0 being a fixed point in $\mathbb{C} \setminus \overline{\Omega}$) is Carathéodory. Every Jordan domain is a Carathéodory domain. The domains, for example, of a snake winding infinitely often around the outside of a circle and approaching this circle (“outer snake”) are Carathéodory, but not a snake winding infinitely often inside a circle and approaching it from the inside (“inner snake”). For more relevant references about Carathéodory domains, please see [11] on page 17.

Questions of completeness for polynomials were first studied by T. Carleman [4] in 1923 who proved that if Ω is a Jordan domain and $\varphi \equiv 0$, then every L^2 holomorphic function on Ω can be approximated by polynomials in $L^2(\Omega, 1)$, and this result was later extended by Farrell [8] and Markushevitch [14] independently to Carathéodory domains. For more general non-Carathéodory domains it is well known that this property need not hold. In [12], Hedberg proved that if Ω is a Carathéodory domain, the weight $e^{-\varphi}$ is continuous and satisfies some conditions then polynomials are dense in $H^2(\Omega, e^{-\varphi})$. For non-Carathéodory domains, the weighted approximation is usually considered when the weight $e^{-\varphi}$ is essentially bounded and satisfies additional conditions (see [3]). Based on Hörmander’s L^2 -estimates for the $\bar{\partial}$ -operator, Taylor [17] proved that polynomials are dense in $H^2(\mathbb{C}^n, e^{-\varphi})$ when φ is convex, which allows the weight to have singularity and can be seen as a major breakthrough for weighted L^2 approximation (see also [19]). Sibony [16] generalized Taylor’s result and obtained that if φ is plurisubharmonic (psh) on \mathbb{C}^n and complex homogeneous of order $\rho > 0$, i.e. $\varphi(uz) = |u|^\rho \varphi(z)$ for $u \in \mathbb{C}$, $z \in \mathbb{C}^n$ then polynomials are dense in $H^2(\mathbb{C}^n, e^{-\varphi})$ (see also [9], section 8). It is well known that each convex function is psh, but the converse is not true. Thus it is natural to ask

Question 1.1. Can we generalize Taylor’s result to any psh function or can we find some non-convex psh function φ on $\Omega \subset \mathbb{C}^n$ satisfying the property that $H^2(\Omega, e^{-\varphi})$ contains all the polynomials but polynomials are not dense in it?

Our first result is

Proposition 1.2. *Let $\Omega = \bigcap_{\nu=0}^N G_\nu$ be a bounded domain in \mathbb{C} where G_0 is a bounded Carathéodory domain and G_ν , $1 \leq \nu \leq N$, are unbounded*

Carathéodory domains. If φ is a subharmonic function on $\overline{\Omega}$, i.e. in a neighborhood V of $\overline{\Omega}$, then $H^2(\overline{\Omega}, e^{-\varphi})$ is dense in $H^2(\Omega, e^{-\varphi})$.

Our proof depends heavily on the Donnelly-Fefferman L^2 -estimate for the $\bar{\partial}$ -operator. In contrast with known results on weighted L^2 approximation of holomorphic functions, we allow singularities of the weight function, which makes the result useful. An application of Proposition 1.2 is the following

Theorem 1.3. *Let Ω be a bounded Carathéodory domain and φ a subharmonic function on $\overline{\Omega}$. Then polynomials are dense in $H^2(\Omega, e^{-\varphi})$.*

Especially we will have the following

Corollary 1.4. *Let Ω be a bounded Jordan domain and let φ be as in Theorem 1.3. Then polynomials are dense in $H^2(\Omega, e^{-\varphi})$.*

Remark 1.5. Under the assumptions of Corollary 1.4, let $f \in H^2(\Omega, e^{-\varphi})$. Then f can be approximated by polynomials in $H^2(\Omega, e^{-\varphi})$ such that the Taylor series of the polynomials around a given point $p \in \Omega$ agrees with the one for f to any given order.

It's not the case that polynomials are dense for general psh weight functions so that the corresponding Hilbert spaces contain the polynomials. An example is provided by the following

Theorem 1.6. *Let $\varphi(z) = |\Im m(z)| + |z|^p$, with $0 < p < 1$. Then the holomorphic polynomials are in $H^2(\mathbb{C}, e^{-\varphi})$, but they are not dense in $H^2(\mathbb{C}, e^{-\varphi})$.*

A general moon-shaped domain is a bounded domain whose boundary consists of two Jordan curves having exactly one point in common. This point is called the multiple boundary point. The moon-shaped domain is an example of a non-Carathéodory Runge domain in \mathbb{C} . Keldych [13] was the first to study the L^2 polynomials approximation property of moon-shaped domains without weight. Here we generalize his result with singular weight as in the following.

Theorem 1.7. *Let Ω be a moon-shaped domain with the origin inside the inner Jordan curve of the boundary $\partial\Omega$. Let φ be a subharmonic function on $\overline{\Omega}$. Then polynomials are dense in $H^2(\Omega, e^{-\varphi})$ if and only if the function $\frac{1}{\sqrt{z}}$ can be approximated arbitrarily well by polynomials in $L^2(\Omega, e^{-\varphi})$.*

We give two concrete examples of moon-shaped domains where density holds. The first example is based on Keldych [13] and is an application of Theorem 1.7.

Definition 1.8. Let ϕ be a subharmonic function on \mathbb{C} . Let μ denote the Laplacian of $\frac{1}{2\pi}\phi$. We say that ϕ satisfies condition (A) if the mass of μ on the closed unit disc is strictly less than 2.

Example 1.9. There exists a moon-shaped domain Ω with the unit circle being the outer Jordan curve, such that for any subharmonic function φ on \mathbb{C} satisfying condition (A), polynomials are in $H^2(\Omega, e^{-\varphi})$ and dense in it.

Example 1.10. There exist a moon-shaped domain Ω bounded by two circles and a subharmonic function φ on Ω so that polynomials are in $H^2(\Omega, e^{-\varphi})$ and are dense in it.

Theorem 1.11. *Let Ω be a moon-shaped domain bounded by two circles and let φ be a subharmonic function on Ω which is uniformly bounded above. Then the set of polynomials which is in $H^2(\Omega, e^{-\varphi})$ is never a dense subset of $H^2(\Omega, e^{-\varphi})$.*

This paper is set-up as follows. In Section 2, we prove Proposition 1.2 and Theorem 1.3, exploiting the property of Carathéodory domains in order to be able to exhaust them from outside by Jordan domains that are conformally equivalent to the unit disc. Then, we apply Donnelly-Fefferman's L^2 -estimates on each of those to obtain the weighted L^2 approximation. In Section 3, we give a counterexample on \mathbb{C} where we exhibit a subharmonic function φ for which the polynomials are in $H^2(\mathbb{C}, e^{-\varphi})$ but they are not dense in it (Theorem 1.6). In Section 4, we prove Theorem 1.7 and give an example of a moon-shaped domain where the density is proved by approximating $\frac{1}{\sqrt{z}}$ (Example 1.9). In Section 5, we present Example 1.10 and we finally prove, in Section 6, Theorem 1.11.

2. PROOF OF THEOREM 1.3

We observe that it suffices to prove Proposition 1.2 and Theorem 1.3 when the subharmonic function φ is globally defined. To see this, let ϕ be a subharmonic function defined on a bounded open set V containing $\bar{\Omega}$ and choose an open set $U, \bar{\Omega} \subset U \subset\subset V$. Then $\mu := \Delta(\phi)|_U$ is a positive measure with bounded mass on \mathbb{C} . Hence there is a globally defined subharmonic function φ such that $\Delta(\varphi) = \mu$. But then $\phi = \varphi + h$ on U for some harmonic function h . Since h is uniformly bounded on Ω , it follows that the Hilbert spaces $H^2(\Omega, e^{-\phi})$ and $H^2(\Omega, e^{-\varphi})$ are the same and the norms are equivalent.

We use, in the next Lemma, the following classical result from one complex variable:

Theorem 2.1 (cf. [18], p. 382). *Let $\{\Omega_n\}_{n=1}^\infty$ be a sequence of uniformly bounded simply connected domains in \mathbb{C} and Ω a bounded simply connected domain, all containing the origin, so that the Hausdorff distance between $\partial\Omega_n$ and $\partial\Omega$ tends to zero as $n \rightarrow \infty$. If we map \mathbb{D} conformally onto Ω_n by $w = f_n(z)$, $f_n(0) = 0$, $f'_n(0) > 0$, then f_n converges locally uniformly to $f \in \mathcal{O}(\mathbb{D})$ such that $w = f(z)$ maps \mathbb{D} conformally onto Ω .*

For a planar domain Ω , let $SH^-(\Omega)$ denote the set of negative subharmonic functions on Ω . Our key observation to prove Proposition 1.2 is the following:

Lemma 2.2. *Let Ω be a bounded Carathéodory domain. Then there exists a sequence of bounded simply-connected domains $\Omega_n \supset \bar{\Omega}$, a sequence of positive numbers $\varepsilon_n \rightarrow 0$ ($n \rightarrow \infty$), and a sequence of continuous functions $\rho_n \in SH^-(\Omega_n)$ such that*

- (1) $\Omega_{n,-\varepsilon_n} := \{z \in \Omega_n : \rho_n(z) < -\varepsilon_n\} \subset \Omega$,
- (2) the volume of $\Omega_n \setminus \Omega_{n,-2\varepsilon_n}$ tends to 0 as $n \rightarrow \infty$.

Before proving the Lemma we recall the following L^2 -estimates for the $\bar{\partial}$ -operator which will be used here.

Proposition 2.3 (Donnelly-Fefferman, [7]). *Let $\Omega \subset \mathbb{C}^n$ be a pseudoconvex domain and $\varphi \in psh(\Omega)$. Suppose that ψ is a C^2 strictly psh function which satisfies*

$$(2.1) \quad i\bar{\partial}\bar{\partial}\psi \geq i\partial\psi \wedge \bar{\partial}\psi.$$

Then for each $\bar{\partial}$ -closed $(0,1)$ -form v , there exists a solution u to $\bar{\partial}u = v$ satisfying

$$(2.2) \quad \int_{\Omega} |u|^2 e^{-\varphi} d\lambda \leq C \int_{\Omega} |v|_{i\bar{\partial}\bar{\partial}\psi}^2 e^{-\varphi} d\lambda,$$

where $C > 0$ is an absolute constant, provided that the right hand side of (2.2) is finite.

The norm $\|\alpha\|_{i\bar{\partial}\bar{\partial}\psi}$ for a $(0,1)$ -form α is the smallest function H that satisfies

$$i\bar{\alpha} \wedge \alpha \leq H(i\bar{\partial}\bar{\partial}\psi).$$

In particular, on \mathbb{C} , we have

$$(2.3) \quad |\alpha|_{i\bar{\partial}\bar{\partial}\psi} = \left(\frac{\partial^2 \psi}{\partial z \partial \bar{z}} \right)^{-1} |\alpha|^2.$$

For more details, see for example [6, 2].

Proof of Lemma 2.2. Since Ω is a Carathéodory domain, there exists a sequence $\{\Omega_n\}$ of bounded simply-connected domains such that $\bar{\Omega} \subset \Omega_n$ and $\bar{\Omega}_{n+1} \subset \Omega_n$ and the Hausdorff distance between $\partial\Omega_n$ and $\partial\Omega$ tends to zero as $n \rightarrow \infty$ (see for example [11], p.17). Without loss of generality, we may assume that $0 \in \Omega$. By virtue of Riemann's mapping theorem, there are conformal mappings

$$w = f_n(z), \quad f_n(0) = 0, \quad f'_n(0) > 0$$

which maps Ω_n onto \mathbb{D} , and

$$w = f(z), \quad f(0) = 0, \quad f'(0) > 0,$$

which maps Ω onto \mathbb{D} . Set

$$\rho(z) = |f(z)| - 1, \quad \rho_n(z) = |f_n(z)| - 1.$$

Clearly, ρ (resp. ρ_n) is a negative continuous subharmonic function on Ω (resp. Ω_n). Let

$$\varepsilon_n := \max \{1 - |f_n(z)| : z \in \overline{\Omega_n} \setminus \Omega\}.$$

By Theorem 2.1, the sequence of Riemann mappings $f_n^{-1} : \mathbb{D} \rightarrow \Omega_n$ converges u.c.c. to the Riemann mapping $f^{-1} : \mathbb{D} \rightarrow \Omega$. Suppose $z \in \Omega_n$ and $\rho_n(z) < -\varepsilon_n$. Then $|f_n(z)| - 1 < -\varepsilon_n$ and hence $1 - |f_n(z)| > \varepsilon_n$. This implies that $z \in \Omega$ which proves (1). Next we prove that $\varepsilon_n \rightarrow 0$. Let $0 < a < b < c < 1$. By the open mapping theorem, there exists $n_0 \in \mathbb{N}$ so that

$$f^{-1}(|z| < a) \subset f_n^{-1}(|z| < b) \subset f^{-1}(|z| < c), \quad \forall n \geq n_0.$$

Hence $f_n^{-1}(|z| < b) \subset \Omega$ so $f_n(\Omega) \supset (|z| < b)$. Therefore if $z \in \Omega_n \setminus \Omega$, then $|f_n(z)| \geq b$, so $1 - |f_n(z)| \leq 1 - b$. This shows that $\varepsilon_n \leq 1 - b$ if $n \geq n_0$. It follows that $\varepsilon_n \rightarrow 0$ as b tends to 1.

Finally we show (2).

Let $\delta > 0$. Then if $0 < a < 1$ is chosen large enough, then the area $|f^{-1}(|z| > a)| < \delta$. Choose $0 < a < b < c < 1$ as above with $1 - b < (1 - a)/2$. Then for all large enough n , $2\varepsilon_n \leq 2(1 - b) < 1 - a$. Suppose that $z \in \Omega \setminus \Omega_{n, -2\varepsilon_n}$. Then $\rho_n(z) \geq -2\varepsilon_n$ so $|f_n(z)| \geq 1 - 2\varepsilon_n > a$ which implies that (for large n) $|f(z)| > a$. Hence $|\Omega \setminus \Omega_{n, -2\varepsilon_n}| < \delta$. □

Now we can prove Proposition 1.2.

Proof of Proposition 1.2. In view of Lemma 2.2, there exist for each $0 \leq \nu \leq N$ a sequence of Jordan domains $G_n^\nu \supset \overline{G}_\nu$, a sequence of positive numbers $\varepsilon_n^\nu \rightarrow 0$ ($n \rightarrow \infty$), and a sequence of continuous functions $\rho_n^\nu \in SH^-(G_n^\nu)$ such that

- (i) $G_{n, -\varepsilon_n^\nu}^\nu := \{z \in G_n^\nu : \rho_n^\nu(z) < -\varepsilon_n^\nu\} \subset G_\nu$,
- (ii) $\text{vol}(G_\nu \setminus G_{n, -2\varepsilon_n^\nu}^\nu) \rightarrow 0$ as $n \rightarrow \infty$.

(In (ii) we can use the spherical metric near ∞ .) Set

$$\rho_n(z) = \max_{0 \leq \nu \leq N} \{\rho_n^\nu(z)\}, \quad \varepsilon_n = \max_{0 \leq \nu \leq N} \{\varepsilon_n^\nu\}, \quad \Omega_n = \bigcap_{\nu=0}^N G_n^\nu.$$

It is easy to verify that $\overline{\Omega} \subset \Omega_n$, $\rho_n \in SH^-(\Omega_n) \cap C(\Omega_n)$ and

- (iii) $\Omega_{n, -\varepsilon_n} := \{z \in \Omega_n : \rho_n(z) < -\varepsilon_n\} \subset \Omega$,
- (iv) $\text{vol}(\Omega \setminus \Omega_{n, -2\varepsilon_n}) \rightarrow 0$ as $n \rightarrow \infty$.

We continue with the proof in a similar way as in the proof in [5]. Choose a family of negative C^∞ subharmonic functions $\{\rho_{n,s}\}$ on Ω_n such that

$\rho_{n,s} \downarrow \rho_n$ uniformly on $\overline{\Omega}_{n+1}$ as $s \downarrow 0$. Put $\psi_n^s = -\log(-\rho_{n,s})$. Clearly, we have

$$(2.4) \quad i\partial\bar{\partial}\psi_n^s \geq i\partial\psi_n^s \wedge \bar{\partial}\psi_n^s.$$

Now choose a cut-off function $\chi : \mathbb{R} \rightarrow [0, 1]$ such that $\chi|_{(-\infty, -\log 3/2)} \equiv 1$ and $\chi|_{[0, \infty)} \equiv 0$. Set $\eta_n^s = \chi(\psi_n^s + \log \varepsilon_n)$ on Ω_n . Then we have

$$\text{supp } \eta_n^s \subset \{z \in \Omega_n \mid \rho_{n,s}(z) < -\varepsilon_n\} \subset \Omega_{n, -\varepsilon_n} \subset \Omega$$

and

$$|\bar{\partial}\eta_n^s|_{i\partial\bar{\partial}\psi_n^s}^2 \leq \sup |\chi'|^2$$

in view of (2.4). Here $|\cdot|_{i\partial\bar{\partial}\psi_n^s}$ stands for the point-wise norm with respect to the metric $i\partial\bar{\partial}\psi_n^s$, like in (2.3). For each $f \in H^2(\Omega, e^{-\varphi})$, we define

$$v_n^s := f\bar{\partial}\eta_n^s.$$

Clearly, v_n^s is a well-defined C^∞ , $\bar{\partial}$ -closed (0,1) form on Ω_n satisfying

$$(2.5) \quad \begin{aligned} \int_{\Omega_{n+1}} |f|^2 |\bar{\partial}\eta_n^s|_{i\partial\bar{\partial}\psi_n^s}^2 e^{-\varphi} d\lambda &\leq \sup |\chi'|^2 \int_{\{z \in \Omega_{n+1} : -\frac{3}{2}\varepsilon_n \leq \rho_{n,s}(z) < -\varepsilon_n\}} |f|^2 e^{-\varphi} d\lambda \\ &\leq \sup |\chi'|^2 \int_{\Omega \setminus \Omega_{n, -2\varepsilon_n}} |f|^2 e^{-\varphi} d\lambda \end{aligned}$$

provided s sufficiently small. By Proposition 2.3, there exists a solution u_n^s to the equation $\bar{\partial}u_n^s = v_n^s$ on Ω_{n+1} verifying, by using (2.5),

$$\int_{\Omega_{n+1}} |u_n^s|^2 e^{-\varphi} d\lambda \leq C \sup |\chi'|^2 \int_{\Omega \setminus \Omega_{n, -2\varepsilon_n}} |f|^2 e^{-\varphi} d\lambda.$$

Let $f_n^s = f\eta_n^s - u_n^s$. Then, $f_n^s \in \mathcal{O}(\Omega_{n+1})$ and

$$\int_{\Omega} |f_n^s - f|^2 e^{-\varphi} d\lambda \leq C \int_{\Omega \setminus \Omega_{n, -2\varepsilon_n}} |f|^2 e^{-\varphi} d\lambda \rightarrow 0 \text{ as } n \rightarrow \infty$$

□

Proof of Theorem 1.3. Let φ be a subharmonic function on $\overline{\Omega}$. If for each $x \in \overline{\Omega}$, the Lelong number $\nu(\varphi)(x)$ of φ satisfies $\nu(\varphi)(x) < 2$ then we have $1 \in H^2(\Omega, e^{-\varphi})$. So there exists a constant $M > 0$ such that $\int_{\Omega} e^{-\varphi} d\lambda < M$.

Let $f \in H^2(\Omega, e^{-\varphi})$. According to the proof of Proposition 1.2, for each $\varepsilon > 0$ there exists $F \in H^2(\Omega_n, e^{-\varphi})$ satisfying

$$\int_{\Omega} |F(z) - f(z)|^2 e^{-\varphi(z)} d\lambda < \frac{\varepsilon}{4}.$$

where Ω_n is a simply connected domain containing $\overline{\Omega}$. We apply Runge's theorem to F and see that for $\delta = \sqrt{\frac{\varepsilon}{4M}}$ there exists a polynomial P such that

$$|F(z) - P(z)| < \delta, \quad z \in \overline{\Omega}.$$

Consequently we have

$$\int_{\Omega} |F(z) - P(z)|^2 e^{-\varphi} d\lambda < \delta^2 \cdot M = \frac{\varepsilon}{4}.$$

Thus we have

$$\begin{aligned} \int_{\Omega} |f(z) - P(z)|^2 e^{-\varphi(z)} d\lambda &\leq 2 \int_{\Omega} |f(z) - F(z)|^2 e^{-\varphi(z)} d\lambda \\ &\quad + 2 \int_{\Omega} |F(z) - P(z)|^2 e^{-\varphi(z)} d\lambda \\ &< \varepsilon \end{aligned}$$

If there exist finitely many points x_0, x_1, \dots, x_N with $\nu(\varphi)(x_j) \geq 2$ for each $0 \leq j \leq N$, then we may choose some polynomial Q so that $Q(x_j) = 0$ and $\varphi = \psi + \log|Q|^2$ with $\nu(\psi) < 2$ at each point x_j . Then for each $f \in H^2(\Omega, e^{-\varphi})$ we have

$$\int_{\Omega} \left| \frac{f}{Q} \right|^2 e^{-\psi} d\lambda < \infty.$$

Since ψ is bounded above, $\int_{\Omega} \left| \frac{f}{Q} \right|^2 d\lambda < \infty$. Hence $\frac{f}{Q}$ is holomorphic on Ω . Based on the above discussion, for each $\varepsilon > 0$ we can find some polynomial P satisfying

$$\int_{\Omega} \left| \frac{f}{Q} - P \right|^2 e^{-\psi} d\lambda < \varepsilon.$$

That is

$$\int_{\Omega} |f - PQ|^2 e^{-\varphi} d\lambda < \varepsilon.$$

□

3. PROOF OF THEOREM 1.6

In this section, we denote $x = \Re e(z)$ and $y = \Im m(z)$.

Before proving Theorem 1.6 by contradiction, we need a couple of Lemmas:

Lemma 3.1. $\cos \frac{z}{2} \in H^2(\mathbb{C}, e^{-\varphi})$.

Proof. Since $\cos \frac{z}{2} = \frac{e^{i\frac{z}{2}} + e^{-i\frac{z}{2}}}{2}$, we have

$$\begin{aligned}
\int_{\mathbb{C}} \left| \cos \frac{z}{2} \right|^2 e^{-\varphi} d\lambda &= \int_{\mathbb{C}} \left| \frac{e^{i\frac{z}{2}} + e^{-i\frac{z}{2}}}{2} \right|^2 e^{-\varphi} d\lambda \\
&= \frac{1}{4} \int_{\mathbb{C}} \left(\left| e^{i\frac{z}{2}} \right|^2 + \left| e^{-i\frac{z}{2}} \right|^2 + e^{i\frac{z}{2}} \overline{e^{-i\frac{z}{2}}} + e^{-i\frac{z}{2}} \overline{e^{i\frac{z}{2}}} \right) e^{-\varphi} d\lambda \\
&\leq \frac{1}{4} \int_{\mathbb{C}} (e^{-y} + e^y + e^{ix} + e^{-ix}) e^{-\varphi} d\lambda \\
(3.1) \quad &= \frac{1}{4} \int_{\mathbb{C}} (e^{-y} + e^y) e^{-|y|-|z|^p} d\lambda + \frac{1}{2} \int_{\mathbb{C}} \cos x e^{-|y|-|z|^p} d\lambda.
\end{aligned}$$

Remark that $(e^{-y} + e^y)e^{-|y|} = 1 + e^{-2|y|} \leq 2$ so we obtain for the first term of the right-hand side of (3.1)

$$\begin{aligned}
\int_{\mathbb{C}} (e^{-y} + e^y) e^{-|y|-|z|^p} d\lambda &= \int_{\mathbb{C}} (e^{-2|y|} + 1) e^{-|z|^p} d\lambda \\
&\leq 2 \int_{\mathbb{C}} e^{-|z|^p} d\lambda \\
&\leq 4\pi \int_0^{+\infty} e^{-r^p} r dr < \infty,
\end{aligned}$$

and for the second term of (3.1),

$$\int_{\mathbb{C}} |\cos x| e^{-|y|-|z|^p} d\lambda \leq \int_{\mathbb{C}} e^{-|y|-|z|^p} d\lambda \leq 2\pi \int_0^{+\infty} r e^{-r^p} dr < \infty.$$

□

We will need the following integral representation too:

Lemma 3.2 (see Chapter 7 of [1]). *Let $u(t) : \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function satisfying*

$$\int_{\mathbb{R}} \frac{u(t)}{1 + |t|^2} dt < \infty.$$

Then

$$U(x + iy) = \frac{1}{\pi} \int_{-\infty}^{+\infty} u(t) \frac{y}{(t-x)^2 + y^2} dt$$

is a harmonic extension of u to the upper half plane.

In particular, if $u(t) = |t|^p$, $0 < p < 1$, then

$$(3.2) \quad U(x + iy) = \frac{1}{\pi} \int_{-\infty}^{+\infty} \frac{y|t|^p}{(x-t)^2 + y^2} dt, \quad y > 0,$$

is a harmonic extension of $|t|^p$ to the upper half plane.

Proposition 3.3. *Let U be as in (3.2). Then there exists constant $C_p > \frac{1}{4}$ so that on the upper half plane $y > 0$,*

$$\frac{1}{4}|z|^p < U(x + iy) < C_p|z|^p.$$

Proof. The right hand side inequality is direct:

$$\begin{aligned}
(3.3) \quad U(x + iy) &= \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{y|t|^p}{(t-x)^2 + y^2} dt \\
&= \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{y|s+x|^p}{s^2 + y^2} ds \\
&= \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{y|y\tau + x|^p}{(y\tau)^2 + y^2} y d\tau \\
&= \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{|y\tau + x|^p}{\tau^2 + 1} d\tau \\
&\leq \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{y^p |\tau|^p}{\tau^2 + 1} d\tau + \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{|x|^p}{\tau^2 + 1} d\tau \\
&\leq \frac{2}{\pi} |y|^p \cdot \pi \frac{\sin(\frac{p}{2}\pi)}{\sin(p\pi)} + |x|^p \\
&= \frac{1}{\cos(\frac{p}{2}\pi)} |y|^p + |x|^p \\
&\leq \frac{2}{\cos(\frac{p}{2}\pi)} |z|^p \\
(3.4) \quad &:= C_p |z|^p.
\end{aligned}$$

To prove the left-hand side inequality, we prove the following inequalities:

(i) $U(x + iy) \geq \frac{1}{2}|x|^p$ and (ii) $U(x + iy) \geq \frac{1}{2}|y|^p$.

We prove (i) as follows: from (3.3) τ inherits the sign of x , then we only have to study $x \geq 0$. We get

$$\begin{aligned}
U(x + iy) &= \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{|y\tau + x|^p}{\tau^2 + 1} d\tau \\
&\geq \frac{1}{\pi} \int_0^{\infty} \frac{|y\tau + x|^p}{\tau^2 + 1} d\tau \\
&\geq \frac{1}{\pi} \int_0^{\infty} \frac{|x|^p}{\tau^2 + 1} d\tau \\
&\geq \frac{1}{2}|x|^p.
\end{aligned}$$

For (ii), we also start from (3.3) and by a similar argument, we may assume that $x \geq 0$. Then,

$$\begin{aligned}
U(x + iy) &= \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{|y\tau + x|^p}{\tau^2 + 1} d\tau \\
&\geq \frac{1}{\pi} \int_0^{\infty} \frac{|y\tau + x|^p}{\tau^2 + 1} d\tau \\
&\geq \frac{1}{\pi} \int_0^{\infty} \frac{|y|^p \tau^p}{\tau^2 + 1} d\tau \\
&= \frac{|y|^p}{\pi} \int_0^{\infty} \frac{\tau^p}{\tau^2 + 1} d\tau \\
&= \frac{1}{2} \frac{|y|^p}{\cos(\frac{p}{2}\pi)} \\
&\geq \frac{1}{2} |y|^p.
\end{aligned}$$

Finally, combining (i) and (ii) and by concavity for $0 < p < 1$, we get

$$U(x + iy) \geq \frac{1}{4}(|x|^p + |y|^p) \geq \frac{1}{4}(|x| + |y|)^p \geq \frac{1}{4}|z|^p.$$

□

Proof of Theorem 1.6. First, remark that the holomorphic polynomials are in $L^2(\mathbb{C}, e^{-\varphi})$ thanks to the exponential rate $e^{-|z|^p}$.

We prove Theorem 1.6 by contradiction. Assume now that holomorphic polynomials are dense in $H^2(\mathbb{C}, e^{-\varphi})$. Since $\cos \frac{z}{2} \in H^2(\mathbb{C}, e^{-\varphi})$ by Lemma 3.1, for all $\varepsilon > 0$, there exist a sequence of polynomials (P_n) and $N \in \mathbb{N}$ such that for all $n \geq N$,

$$(3.5) \quad \int_{\mathbb{C}} \left| P_n(z) - \cos \frac{z}{2} \right|^2 e^{-\varphi} d\lambda < \varepsilon.$$

Note that for n sufficiently large,

$$\begin{aligned}
\|P_n(z)\|_{\mathbb{C}, \varphi} &= \left\| P_n(z) - \cos \frac{z}{2} + \cos \frac{z}{2} \right\|_{\mathbb{C}, \varphi} \\
&\leq \left\| P_n(z) - \cos \frac{z}{2} \right\|_{\mathbb{C}, \varphi} + \left\| \cos \frac{z}{2} \right\|_{\mathbb{C}, \varphi} \\
(3.6) \quad &\leq 1 + \left\| \cos \frac{z}{2} \right\|_{\mathbb{C}, \varphi}.
\end{aligned}$$

We deduce from (3.6) that there exists $M > 1$ such that $\|P_n(z)\|_{\mathbb{C}, \varphi} \leq M$ for all n .

Since P_n is analytic, we have

$$P_n(z) = \frac{1}{\pi} \int_{|\zeta| \leq 1} P_n(z + \zeta) d\lambda_{\zeta}.$$

So

$$\begin{aligned}
|P_n(z)| &\leq \frac{1}{\pi} \int_{|\zeta| \leq 1} |P_n(z + \zeta)| e^{\frac{\varphi(z+\zeta)}{2}} e^{-\frac{\varphi(z+\zeta)}{2}} d\lambda_\zeta \\
(3.7) \quad &\leq \frac{1}{\pi} \sup_{|\zeta| \leq 1} e^{\frac{\varphi(z+\zeta)}{2}} \int_{|\zeta| \leq 1} |P_n(z + \zeta)| e^{-\frac{\varphi(z+\zeta)}{2}} d\lambda_\zeta.
\end{aligned}$$

By Cauchy-Schwarz's inequality, we get from (3.7) and for all n ,

$$\begin{aligned}
|P_n(z)| &\leq \frac{1}{\pi} \sup_{|\zeta| \leq 1} e^{\frac{\varphi(z+\zeta)}{2}} \left(\int_{|\zeta| \leq 1} |P_n(z + \zeta)|^2 e^{-\varphi(z+\zeta)} d\lambda_\zeta \right)^{\frac{1}{2}} \left(\int_{|\zeta| \leq 1} d\lambda_\zeta \right)^{\frac{1}{2}} \\
(3.8) \quad &\leq \frac{1}{\sqrt{\pi}} \|P_n(z)\|_{\mathbb{C}, \varphi} \sup_{|\zeta| \leq 1} e^{\frac{\varphi(z+\zeta)}{2}} \leq \frac{M}{\sqrt{\pi}} \sup_{|\zeta| \leq 1} e^{\frac{\varphi(z+\zeta)}{2}}.
\end{aligned}$$

Note that for $0 < p < 1$,

$$\begin{aligned}
\sup_{|\zeta| \leq 1} \varphi(z + \zeta) &= \sup_{|\zeta| \leq 1} (|\Im m(z + \zeta)| + |z + \zeta|^p) \\
&\leq |\Im m(z)| + 1 + (|z| + 1)^p \\
&\leq |\Im m(z)| + 1 + 1 + |z|^p \\
&= |y| + 2 + |z|^p.
\end{aligned}$$

Plugging it in (3.8), we get for all n

$$|P_n(x + iy)| \leq \frac{M}{\sqrt{\pi}} e^{\frac{1}{2}(2+|y|+|x+iy|^p)}.$$

Then it follows that on the real axis ($y = 0$),

$$|P_n(x)| \leq \frac{M}{\sqrt{\pi}} e^{\frac{1}{2}(2+|x|^p)} \leq \frac{M}{\sqrt{\pi}} e^{\frac{1}{2}|x|^p},$$

so

$$(3.9) \quad \log |P_n(x)| \leq \log M + 1 - \log \sqrt{\pi} + \frac{1}{2}|x|^p.$$

By Lemma 3.2 applied to $\tilde{u}(x) = \log M + 1 - \log \sqrt{\pi} + \frac{1}{2}|x|^p$,

$$\tilde{U}(x + iy) = \frac{1}{\pi} \int_{-\infty}^{+\infty} \frac{y(\log M + 1 - \log \sqrt{\pi} + \frac{1}{2}|t|^p)}{(x-t)^2 + y^2} dt, \quad y > 0,$$

is a harmonic extension of \tilde{u} to the upper half plane. And from Proposition 3.3, we obtain

$$\log M + 1 - \log \sqrt{\pi} + \frac{1}{8}|z|^p \leq \tilde{U}(z) \leq \log M + 1 - \log \sqrt{\pi} + C_p |z|^p.$$

From the fact that for any positive constant C , $C \log |z|$ is much smaller than $|z|^p$ for large enough $|z|$, we deduce that for large enough $|z|$, $\log |P_n|(z) \leq \tilde{U}(z)$. Because \tilde{U} is harmonic and $\log |P_n|$ is subharmonic, we obtain

$$(3.10) \quad \log |P_n(x + iy)| \leq \tilde{U}(x + iy) \leq C_1 + C_p |z|^p \quad \text{on } \{z \in \mathbb{C} \mid y > 0\},$$

where $C_1 = \log M + 1 - \log \sqrt{\pi}$. From (3.10), we get

$$\log |P_n(z)| \leq C_1 + C_p |z|^p,$$

so

$$|P_n(z)| \leq e^{C_1} e^{C_p |z|^p}.$$

But remark that

$$\lim_{|z| \rightarrow +\infty} \frac{e^{|z|/4} - 1}{e^{C_1} e^{C_p |z|^p}} = +\infty$$

so there exists a positive constant $Y > 1$ such that for all $|z| > Y$,

$$(3.11) \quad e^{|z|/4} - 1 > 4e^{C_1} e^{C_p |z|^p}.$$

Note also that if $x^2 < 3y^2$ then $4y^2 = y^2 + 3y^2 > x^2 + y^2 = |z|^2$ so if in addition $y > 0$, we get $y > \frac{1}{2}|z|$. Hence, on $\{z \in \mathbb{C} \mid x^2 < 3y^2, y > 0\}$, we have

$$(3.12) \quad \begin{aligned} \left| \cos \frac{z}{2} \right| &= \left| \frac{1}{2} e^{ix/2 - y/2} + \frac{1}{2} e^{ix/2 + y/2} \right| \\ &\geq \left| \frac{1}{2} e^{ix/2 + y/2} \right| - \left| \frac{1}{2} e^{ix/2 - y/2} \right| \\ &= \frac{1}{2} (e^{y/2} - e^{-y/2}) \\ &\geq \frac{1}{2} (e^{|z|/4} - 1). \end{aligned}$$

Combining (3.12) with (3.11), we obtain on $W := \{z \in \mathbb{C} \mid x^2 < 3y^2, y > 0, |z| > Y\} = \{(r, \theta) \mid r > Y, \frac{\pi}{6} < \theta < \frac{5\pi}{6}\}$,

$$(3.13) \quad \left| \cos \frac{z}{2} - P_n(z) \right| \geq \left| \cos \frac{z}{2} \right| - |P_n(z)| \geq e^{C_1} e^{C_p |z|^p} \geq e^{C_1} e^{C_p Y^p}.$$

Hence, from (3.13), we have on W ,

$$\begin{aligned} \int_{\mathbb{C}} \left| \cos \frac{z}{2} - P_n(z) \right|^2 e^{-\phi(z)} d\lambda &\geq \int_W \left| \cos \frac{z}{2} - P_n(z) \right|^2 e^{-\varphi(z)} d\lambda \\ &\geq e^{2C_1} e^{2C_p Y^p} \int_W e^{-\varphi(z)} d\lambda \\ &= e^{2C_1 + 2C_p Y^p} \int_{\pi/6}^{5\pi/6} d\theta \int_Y^{+\infty} r e^{-|r \sin \theta| - |r|^p} dr \\ &\geq \frac{2\pi}{3} e^{2C_1 + 2C_p Y^p} \int_Y^{+\infty} r e^{-|r| - |r|^p} dr \\ &\geq \frac{2\pi}{3} e^{2C_1 + 2C_p Y^p} \cdot \frac{1}{2} e^{-2Y} \\ &= \frac{\pi}{3} e^{2C_1 + 2C_p Y^p - 2Y}. \end{aligned}$$

This is a contradiction with the formula (3.5) when n is large enough. \square

4. PROOF OF THEOREM 1.7

Recall the following classical fact:

Lemma 4.1. *Let Ω be an open set in \mathbb{C} , h a holomorphic on Ω and φ a subharmonic function on an open set $V \supset h(\Omega)$. Then $\varphi \circ h(z) = \varphi(h(z))$ is also subharmonic on Ω .*

Proof of Theorem 1.7. Since $\frac{1}{\sqrt{z}} \in H^2(\Omega, e^{-\varphi})$, the condition is obviously necessary.

We then need to prove the sufficiency. The mapping $w = \sqrt{z}$ transforms Ω into a Jordan domain Ω' and for each $f \in H^2(\Omega, e^{-\varphi})$,

$$\int_{\Omega} |f(z)|^2 e^{-\varphi(z)} d\lambda_z = 4 \int_{\Omega'} |f(w^2)w|^2 e^{-\varphi(w^2)} d\lambda_w < \infty.$$

Put $h(w) = w^2$, we know that $wf(w^2) \in H^2(\Omega', e^{-\varphi \circ h(w)})$, where $\varphi \circ h(w) = \varphi(h(w))$ is subharmonic on the closure of the bounded Jordan domain Ω' . In particular, $\varphi \circ h(w) = \varphi(h(w))$ is subharmonic in a neighborhood of $\overline{\Omega'}$. According to Corollary 1.4, for each $\varepsilon > 0$ there exists a polynomial $P(w)$ such that

$$4 \int_{\Omega'} |wf(w^2) - P(w)|^2 e^{-\varphi \circ h(w)} d\lambda_w < \varepsilon.$$

Therefore

$$(4.1) \quad \int_{\Omega} |\sqrt{z}f(z) - P(\sqrt{z})|^2 e^{-\varphi(z)} \cdot \frac{1}{|z|} d\lambda_z = \int_{\Omega} \left| f(z) - \frac{P(\sqrt{z})}{\sqrt{z}} \right|^2 e^{-\varphi(z)} d\lambda_z < \varepsilon.$$

Separate the polynomial P into even and odd parts:

$$P(\sqrt{z}) = P_1(z) + \sqrt{z}P_2(z),$$

then the formula (4.1) implies that

$$\int_{\Omega} \left| f(z) - P_2(z) - \frac{P_1(z)}{\sqrt{z}} \right|^2 e^{-\varphi(z)} d\lambda_z < \varepsilon.$$

In order to find some polynomial $Q(z)$ such that

$$\int_{\Omega} |f(z) - Q(z)|^2 e^{-\varphi(z)} d\lambda_z < 2\varepsilon.$$

It is sufficient to know that

$$\int_{\Omega} \left| \frac{1}{\sqrt{z}} - R(z) \right|^2 e^{-\varphi(z)} d\lambda < \varepsilon$$

for some polynomial $R(z)$ but this holds by assumption. \square

We give now an application of Theorem 1.7: Example 1.9. First we need the following lemmas.

Lemma 4.2 (Riesz Decomposition Theorem, see for example Theorem 3.7.9 in [15]). *Let u be a subharmonic function on a domain D in \mathbb{C} , with $u \not\equiv -\infty$. Then, given a relatively compact open subset U of D , we can decompose u as*

$$u = \int_{\zeta \in U} \log |z - \zeta| d\mu(\zeta) + h$$

on U , where $\mu = \frac{1}{2\pi} \Delta u|_U$ and h is harmonic on U .

Let D, R be positive numbers so that $D = \pi R^2$. Fix $\alpha, 0 < \alpha < 2$.

Lemma 4.3. *Let $z_0 \in \mathbb{C}$ and A be a measurable set in \mathbb{C} with bounded area D . Then $\int_A \frac{1}{|z - z_0|^\alpha} d\lambda \leq \frac{R^{2-\alpha}}{2-\alpha}$. Moreover this estimate is sharp.*

Proof. It suffices to consider the case when $z_0 = 0$. The largest integral occurs when the area is a disc centered at 0. The radius then is given by $\pi R^2 = D$. We get $\int_{|z| \leq R} \frac{1}{|z|^\alpha} d\lambda = \int_0^R t^{1-\alpha} dt = \frac{R^{2-\alpha}}{2-\alpha}$. \square

Using the convexity of the exponential function we apply Lemma 2.1 in [10] to obtain:

Lemma 4.4. *Suppose $0 < \alpha_i, \sum_i \alpha_i = \alpha < 2$. Then*

$$\int_A \prod_i \frac{1}{|z - z_i|^{\alpha_i}} d\lambda \leq \frac{R^{2-\alpha}}{2-\alpha}$$

as in the above Lemma.

Corollary 4.5. *Let μ be any nonnegative measure with total mass $\alpha, 0 < \alpha < 2$. A be a measurable set in \mathbb{C} with bounded area D . Then if $\varphi(z) = \int \log |z - \zeta| d\mu(\zeta)$, we have that $\int_A e^{-\varphi} d\lambda \leq \frac{R^{2-\alpha}}{2-\alpha}$.*

Proof. Define $\psi_n(z, \zeta) = \max\{\log |z - \zeta|, -n\}$ and

$$\varphi_n(z) = \int \psi_n(z, \zeta) d\mu(\zeta).$$

It suffices to show that

$$\int_A e^{-\varphi_n(z)} d\lambda(z) \leq \frac{R^{2-\alpha}}{2-\alpha} + \frac{1}{n} \quad \forall n \in \mathbb{N}^*.$$

We fix n . Let $\varepsilon > 0$. We apply Lemma 2.4 [10]: there exists a finite positive measure $\beta := \sum_{i=1}^N \alpha_i \delta_{z_i}$ with $\sum_{i=1}^N \alpha_i = \alpha$ such that

$$\int \psi_n(z, \zeta) d\beta(\zeta) \leq \int \psi_n(z, \zeta) d\mu(\zeta) + \varepsilon.$$

By Lemma 4.4 we know that

$$\int_A e^{-\sum_i \alpha_i \log |z-z_i|} d\lambda \leq \frac{R^{2-\alpha}}{2-\alpha}.$$

So

$$\int_A e^{-\int \log |z-\zeta| d\beta(\zeta)} d\lambda(z) \leq \frac{R^{2-\alpha}}{2-\alpha}.$$

Hence $\int_A e^{-\int \psi_n(z,\zeta) d\beta(\zeta)} d\lambda \leq \frac{R^{2-\alpha}}{2-\alpha}$. Finally, by choosing ε small enough we get

$$\int_A e^{-\varphi_n} d\lambda \leq \frac{R^{2-\alpha}}{2-\alpha} + \frac{1}{n}.$$

□

Define for $n \in \mathbb{N}^*$

$$S_n = \left\{ \varphi \text{ is subharmonic on } \mathbb{C} : \begin{array}{l} \text{the mass of } \mu \left(|z| < 1 + \frac{1}{n} \right) < 2 - \frac{1}{n}, \\ \varphi(z) = \psi_n + h_n, \quad \psi_n = \int_{|\zeta| < 1 + \frac{1}{n}} \log |z - \zeta| d\mu \text{ and } |h_n| < n \text{ on } |z| < 1 \end{array} \right\}.$$

Lemma 4.6. *Suppose φ is subharmonic on \mathbb{C} satisfying condition (A), then for all large enough n , $\varphi \in S_n$.*

Proof. Pick m so that the mass of $\mu = \frac{1}{2\pi} \Delta \varphi$ is strictly less than 2 on the disc $\Delta \left(1 + \frac{1}{m} \right)$. Increasing m , we may assume

$$\mu \left(|z| < 1 + \frac{1}{m} \right) < 2 - \frac{1}{m}.$$

This remains true for all large m , let φ be subharmonic on \mathbb{C} satisfying condition (A), write $\varphi = \psi_m + h_m$ and set

$$K = \sup_{|z| \leq 1} |h_m(z)|.$$

For $n > m$, we may also write $\varphi = \psi_n + h_n$. We have that

$$|\psi_n - \psi_m| \leq 2 \log n \quad \text{on } |z| < 1.$$

Hence

$$|h_n| \leq K + 2 \log n \quad \text{on } |z| < 1.$$

We may choose n so large that

$$K + 2 \log n \leq n.$$

□

Now we begin to construct Example 1.9. By Lemma 4.6, we may assume $\varphi \in S_1$. Then, $\varphi = \varphi_1 + h_1$ where $\varphi_1 = \int_{|\zeta| < 2} \log |z - \zeta| d\mu$ and $\mu(|z| < 2) < 1$. Let D_1 be the domain composed of points satisfying the inequalities

$$|z| < 1, \quad \left| z - \frac{1}{2^2} \right| > \frac{3}{4}, \quad \frac{\pi}{2^2} < \arg z < 2\pi - \frac{\pi}{2^2}.$$

Then D_1 is a bounded simply connected domain so that $\mathbb{C} \setminus \overline{D_1}$ is connected. By Corollary 4.5 we know that $\int_{D_1} e^{-\varphi_1} d\lambda$ is uniformly bounded. Since $|h_1| < 1$, we also get that $\int_{D_1} e^{-\varphi} d\lambda$ is uniformly bounded for all $\varphi \in S_1$. By the Runge theorem there exists a polynomial $P_{n_1}(z)$ with degree n_1 such that

$$\int_{D_1} \left| \frac{1}{\sqrt{z}} - P_{n_1}(z) \right|^2 e^{-\varphi(z)} d\lambda < \frac{1}{2^2}$$

for all $\varphi \in S_1$. Choose $0 < \alpha_1 < \frac{1}{2^2}$ is sufficiently small so that for the above polynomial $P_{n_1}(z)$ satisfying

$$\begin{aligned} \int_{\Delta_1} \left| \frac{1}{\sqrt{z}} - P_{n_1}(z) \right|^2 e^{-\varphi(z)} d\lambda &\leq \sup_{\Delta_1} \left| \frac{1}{\sqrt{z}} - P_{n_1}(z) \right|^2 \int_{\Delta_1} e^{-\varphi_1 - h_1} d\lambda \\ &\leq \sup_{\Delta_1} \left| \frac{1}{\sqrt{z}} - P_{n_1}(z) \right|^2 \frac{R_1^{2-\alpha_1}}{2-\alpha_1} \cdot e \\ &< \frac{1}{2^2}, \end{aligned}$$

where Δ_1 , of area πR_1^2 being the points satisfying the inequalities

$$(4.2) \quad |z| < 1, \quad |z - \alpha_1| > 1 - \alpha_1, \quad |\arg z| \leq \frac{\pi}{2^2}.$$

Let \widetilde{D}_2 be the domain composed of the points verifying the inequalities

$$|z| < 1, \quad |z - \alpha_1| > 1 - \alpha_1, \quad \frac{\pi}{2^3} < \arg z < 2\pi - \frac{\pi}{2^3}.$$

Set $D_2 = D_1 \cup \widetilde{D}_2$, this is a Jordan domain. We now consider any $\varphi \in S_2$. Then there exists a polynomial $P_{n_2}(z)$ with degree n_2 such that

$$\int_{D_2} \left| \frac{1}{\sqrt{z}} - P_{n_2}(z) \right|^2 e^{-\varphi(z)} d\lambda < \frac{1}{2^3}.$$

Choose $0 < \alpha_2 < \alpha_1$ so that for the above polynomial $P_{n_2}(z)$ satisfying

$$\begin{aligned} \int_{\Delta_2} \left| \frac{1}{\sqrt{z}} - P_{n_2}(z) \right|^2 e^{-\varphi(z)} d\lambda &\leq \sup_{\Delta_2} \left| \frac{1}{\sqrt{z}} - P_{n_2}(z) \right|^2 \int_{\Delta_2} e^{-\varphi} d\lambda \\ &\leq \sup_{\Delta_2} \left| \frac{1}{\sqrt{z}} - P_{n_2}(z) \right|^2 \frac{R_2^{2-\alpha_2}}{2-\alpha_2} \cdot e^2 \\ &< \frac{1}{2^3}, \end{aligned}$$

where Δ_2 , of area πR_2^2 being the points satisfying the inequalities

$$|z| < 1, \quad |z - \alpha_2| > 1 - \alpha_2, \quad |\arg z| \leq \frac{\pi}{2^3}.$$

Let $\widetilde{D}_3 := \{|z| < 1, |z - \alpha_2| > 1 - \alpha_2, \frac{\pi}{2^4} < \arg z < 2\pi - \frac{\pi}{2^4}\}$. Set $D_3 = D_2 \cup \widetilde{D}_3$. Proceed as above, we can find a sequence of bounded simply connected domains $D_1, D_2, \dots, D_n, \dots$. Let D be the limit domain of D_n . Then the domain D is bounded by the circle $|z| = 1$ and a simple Jordan curve Γ tangent to $|z| = 1$ in $z = 1$. Thus D is a bounded very thin moon-shaped domain and we know that the limit domain D is contained in the limit domain $D_k \cup \Delta_k$, from which it follows that for all $\varphi \in S_k$,

$$\int_D \left| \frac{1}{\sqrt{z}} - P_{n_k}(z) \right|^2 e^{-\varphi_1(z)} d\lambda < \frac{1}{2^{k+1}} + \frac{1}{2^{k+1}} = \frac{1}{2^k}.$$

Then polynomials are dense in $H^2(D, e^{-\varphi})$ for any φ satisfying condition (A).

5. EXAMPLE 1.10

First, we prove the following Lemma, already known for Bergman spaces ($\varphi \equiv 0$).

Lemma 5.1. *Let $\Omega \subset \subset \mathbb{C}$ and φ a subharmonic function on Ω . Assume that $1 \in H^2(\Omega, e^{-\varphi})$ and $p \in \Omega$. For each n , let $f_n \in H^2(\Omega, e^{-\varphi})$ be a function $f_n = a_n(z-p)^n + O((z-p)^{n+1})$, $a_n > 0$ maximal with $\|f_n\|_{\Omega, \varphi} = 1$. Then $\{f_n\}_{n=0}^\infty$ is an orthonormal basis for $H^2(\Omega, e^{-\varphi})$.*

Proof. We show that $f_n \perp \{g \in H^2(\Omega, e^{-\varphi}); g = O((z-p)^{n+1})\}$.

By contradiction, suppose that there exists such $g \in H^2(\Omega, e^{-\varphi})$ that is not orthogonal to f_n . Then for complex-valued ε small enough,

$$(5.1) \quad \langle f_n + \varepsilon g, f_n + \varepsilon g \rangle = 1 - 2\Re(\varepsilon \langle f_n, g \rangle) + |\varepsilon|^2 \langle g, g \rangle,$$

$$(5.2) \quad = 1 - 2|\varepsilon| |\langle f_n, g \rangle| + |\varepsilon g|^2,$$

$$(5.3) \quad = t < 1.$$

Hence, $\widetilde{f}_n = \frac{f_n + \varepsilon g}{\sqrt{t}} = \frac{a_n}{\sqrt{t}}(z-p)^n + O_t((z-p)^{n+1})$ contradicts the maximality of the coefficients.

Suppose now that $g \perp \{f_n\}$ such that $g = b_n(z-p)^n + O((z-p)^{n+1})$ with $b_n = a_n$. We get $g - f_n = O((z-p)^{n+1})$ that is orthogonal to f_n i.e

$$\langle g - f_n, f_n \rangle = \langle g, f_n \rangle - \langle f_n, f_n \rangle = 0.$$

but $\langle g, f_n \rangle = 0 \neq 1 = \langle f_n, f_n \rangle$. □

Let M be a moon-shaped domain with multiple boundary point Q , and p be an interior point of M . For each $n \geq 2$, denote $M_n = M \setminus B(Q, 1/n)$ where $B(Q, 1/n)$ is the disc of center Q and radius $1/n$. Let φ be a non-negative subharmonic function on \mathbb{C} and $f \in H^2(M_n, e^{-\varphi})$. By Corollary 1.4, for every $\varepsilon > 0$, there exists a polynomial P such that $\|f - P\|_{M_n, \varphi} < \varepsilon$.

Lemma 5.2. *Under the previous assumptions, there exists a subharmonic function $\tilde{\varphi}$ on \mathbb{C} such that $\tilde{\varphi} = \varphi$ on M_n and*

$$\|P\|_{B(Q, 1/n) \cap M, \tilde{\varphi}} < \varepsilon.$$

Proof. Because $\overline{M_n}$ is polynomially convex, there exists a subharmonic function γ on \mathbb{C} such that $\gamma = 0$ on M_n and $\gamma > 0$ outside: To find γ , note that for every point q outside $\overline{M_n}$ there exists a polynomial P_q such that $|P_q(q)| > 1$ and $|P_q| < 1$ on $\overline{M_n}$. We choose a convex function $\chi(x)$ which vanishes when $x \leq 1$ and is strictly positive when $x > 1$. Then $\chi \circ |P_q|^2$ is subharmonic and vanishes on M_n while it is strictly positive in a neighborhood of q . Then one can define $\gamma = \sum_m \varepsilon_m \chi \circ |P_{q_m}|^2$ for suitable choices. By choosing $\tilde{\varphi} = \varphi + L\gamma$ for L large enough, we get

$$\|P\|_{B(Q, 1/n) \cap M, \tilde{\varphi}} \leq \|P\|_{B(Q, 1/n) \cap M, L\gamma}.$$

By taking the limit as L tends to $+\infty$, the second term of the previous estimate tends to 0. \square

By Lemma 5.2, we can construct inductively an increasing sequence of non-negative subharmonic functions φ_n on M and by Lemma 5.1, we can find $f_0^n, \dots, f_n^n \in H^2(M_n, e^{-\varphi_n})$ such that $f_j^n = a_j^n (z - p)^j + O((z - p)^{j+1})$ with $a_j^n > 0$ maximal and $\|f_j^n\|_{M_n, \varphi_n} = 1$. Then, there exist polynomials P_j^n and φ_{n+1} large enough by Lemma 5.2 such that

$$(5.4) \quad \|f_j^n - P_j^n\|_{M_n, \varphi_n} < \frac{1}{n}$$

and

$$(5.5) \quad \|P_j^n\|_{B(Q, 1/n) \cap M, \varphi_{n+1}} < \frac{1}{n}$$

Moreover, by Remark 1.5, we can choose

$$P_j^n(z) = a_j^n (z - p)^j + O((z - p)^{j+1}).$$

Let $\varphi := \lim_{n \rightarrow +\infty} \varphi_n$. Hence $\|f_j^n\|_{M_n, \varphi} = 1$ and (5.4) and (5.5) hold with respect to φ .

Lemma 5.3. *Under the previous assumptions, the polynomials P_j^n , $j = 0, \dots, n$ built inductively as previously verify the following property*

$$\|P_j^n\|_{M, \varphi} \leq 1 + \frac{2}{n}.$$

Proof. Combining the properties of f_j^n seen in Lemma 5.1 and Lemma 5.2

$$\begin{aligned} \|P_j^n\|_{M,\varphi} &\leq \|P_j^n\|_{M_n,\varphi} + \|P_j^n\|_{B(Q,1/n)\cap M,\varphi}, \\ &\leq \|P_j^n\|_{M_n,\varphi_n} + \|P_j^n\|_{B(Q,1/n)\cap M,\varphi_{n+1}}, \\ &\leq \|P_j^n - f_j^n\|_{M_n,\varphi_n} + \|f_j^n\|_{M_n,\varphi_n} + \frac{1}{n}, \\ &\leq 1 + \frac{2}{n}. \end{aligned}$$

□

We are now able to give the details of the construction of Example 1.10.

By Lemma 5.3, P_j^n converges weakly to P_j in $H^2(M, e^{-\varphi})$ such that

$$\|P_j\|_{M,\varphi} \leq 1$$

and

$$P_j = \left(\lim_{n \rightarrow +\infty} a_j^n \right) (z-p)^j + O((z-p)^{j+1}).$$

In particular, the limit $\lim_{n \rightarrow +\infty} a_j^n$ gives optimal coefficients for P_j . It follows that $\|P_j\|_{M,\varphi} = 1$ and P_j is an orthonormal basis for $H^2(M, e^{-\varphi})$ by Lemma 5.1.

Let $f \in H^2(M, e^{-\varphi})$ and $\varepsilon > 0$. We can express f as $f = \sum_{j=0}^{\infty} A_j P_j$.

For N large enough,

$$\left\| \sum_{j=0}^N A_j P_j - f \right\|_{M,\varphi} < \frac{\varepsilon}{2}.$$

For n large enough, we get finally

$$\left\| \sum_{j=0}^N A_j P_j^n - f \right\|_{M,\varphi} < \varepsilon.$$

6. PROOF OF THEOREM 1.11

Proof of Theorem 1.11. The proof relies on [11], Chapter I, section 3. We reason by contradiction.

Let Γ be a circle as in Figure 1. Then there exists a positive constant C such that $d_{\partial\Omega}(z) \geq C|z-1|^2$ for all $z \in \Gamma$. Choose a function $f \in H^2(\Omega, e^{-\varphi})$ which does not extend holomorphically to the inside of the small circle. (One can first choose any nontrivial holomorphic function in the $H^2(\Omega, e^{-\varphi})$. Pick a suitable point p inside the inner circle and observe that $\frac{f}{z-p}$ is still in the H^2 space.)

Assume that there exists a sequence of polynomials P_n so that $\|f - P_n\|_{\Omega,\varphi}$ tends to 0 as n tends to $+\infty$. We obtain a contradiction by showing that f extends analytically in the interior of Γ so in the hole (see Figure 1), which




FIGURE 1. A general moon-shaped domain including the curve Γ passing through the multiple boundary point Q .

is impossible.

For each $w \in \Gamma \setminus \{Q\}$, by the mean value property for subharmonic functions, we get for any $n, m \in \mathbb{N}$,

$$(6.1) \quad \begin{aligned} |P_n(w) - P_m(w)|^2 &\leq \frac{1}{\pi d_{\partial\Omega}^2(w)} \int_{B(w, d_{\partial\Omega}(w))} |P_n(z) - P_m(z)|^2 d\lambda, \\ &\leq \frac{\tilde{C}}{\pi d_{\partial\Omega}^2(w)} \int_{B(w, d_{\partial\Omega}(w))} |P_n(z) - P_m(z)|^2 e^{-\varphi(z)} d\lambda, \end{aligned}$$

$$(6.2) \quad \leq \frac{\tilde{C}}{\pi d_{\partial\Omega}^2(w)} \|P_n(z) - P_m(z)\|_{\Omega, \varphi}^2,$$

where we have used in (6.1) the fact that φ is bounded above in Ω with \tilde{C} a positive constant. Because $d_{\partial\Omega}(z) \geq C|z - 1|^2$ by assumption, we obtain from (6.2) and for each $w \in \Gamma \setminus \{Q\}$,

$$(6.3) \quad \begin{aligned} |(w - 1)^2(P_n(w) - P_m(w))| &\leq \frac{d_{\partial\Omega}(w)}{C} \frac{\sqrt{\tilde{C}}}{\sqrt{\pi} d_{\partial\Omega}(w)} \|P_n(z) - P_m(z)\|_{\Omega, \varphi}, \\ &\leq C' \|P_n(z) - P_m(z)\|_{\Omega, \varphi} \end{aligned}$$

where $C' > 0$. The inequality (6.3) holds for each point $w \in \Gamma$ so in the interior of Γ . Hence, the sequence $((z - 1)^2 P_n)$ converges uniformly in the interior of Γ to $(z - 1)^2 f$.

□

Acknowledgements The authors would like to thank Zhonghua Wang for useful comments and the referees for their valuable suggestions.

REFERENCES

- [1] S. Axler, P. Bourdon and R. Wade, Harmonic function theory. Springer Science & Business Media, 2013.
- [2] Z. Błocki, Cauchy-Riemann meet Monge-Ampère. *Bull. Math. Sci* **4** (2014), 433–480.
- [3] J. Brennan, Approximation in the mean of polynomials on non-Caratheodory domains, *Ark. Mat* **15**, no.1 (1977), 117–168.
- [4] T. Carleman, Über die Approximation analytischer Funktionen durch lineare Aggregate von vorgegebenen Potenzen[M]. Almqvist and Wiksell, 1923.
- [5] B.Y. Chen and J.H. Zhang, On Bergman completeness and Bergman Stability, *Math. Ann.*, **318**, 2000, 517-526.
- [6] J.P. Demailly, Estimations L^2 pour l'opérateur $\bar{\partial}$ d'un fibré vectoriel holomorphe semi-positif au dessus d'une variété kählérienne complète, *Annales Sc. de l'E.N.S*, **15**, no.3 (1982), 457–511.
- [7] H. Donnelly and C. Fefferman, L^2 -cohomology and index theorem for the Bergman metric, *Ann. of Math.* **118** (1983), 593-618.
- [8] O. Farrell, On approximation to an analytic function by polynomials, Bull. Amer. Math. Soc. **40**, no. 12 (1934), 908–914.
- [9] J. E. Fornæss, F. Forstneric and E.F. Wold, Holomorphic approximation: the legacy of Weierstrass, Runge, Oka-Weil, and Mergelyan, arXiv:1802.03924v2.
- [10] J.E. Fornæss and J. Wu, Weighted approximation in \mathbb{C} , *J. Math. Z.* (2019) <https://doi.org/10.1007/s00209-019-02321-w>.
- [11] D. Gaier, Lectures on Complex Approximation, Boston. Basel. Stuttgart. 1980.
- [12] L.I. Hedberg, Weighted mean square approximation in plane regions and generators of an algebra of analytic functions, *Arkiv för Matematik* band, **5**, no. 36, (1965), 541–552.
- [13] M. Keldych, "Sur l'approximation en moyenne quadratique des fonctions analytiques." *Rec. Math. [Mat. Sb.]* **5(47)**, no. 2 (1939): 391-401.
- [14] A.I. Markushevitch, Conformal mapping of regions with variable boundaries for the approximation of analytic functions by polynomials (russian), PhD. Thesis, (1934).
- [15] T. Ransford, Potential theory in the complex plane, Vol. 28, Cambridge University press, 1995.
- [16] N. Sibony, Approximation polynomiale pondérée dans un domaine d'holomorphic de \mathbb{C}^n , *Annales de l'Institut Fourier*, tome 26, 1976, 71-79.
- [17] B.A. Taylor, On weighted polynomial approximation of entire functions, *Pacific. J. Math.*, **36**, 1971, 523-539.
- [18] M. Tsuji, Potential Theory in Modern Function Theory, Maruzen Co., LTD. Tokyo, 1959.
- [19] D. Wohlgeleinter, Weighted L^2 approximation of entire functions, *Transactions Of The American Mathematical Society.*, **202**, 1975, 211-219.

SCIENCE INSTITUTE, UNIVERSITY OF ICELAND, DUNHAGI 3, IS-107 REYKJAVIK, ICELAND

Email address: `biard@hi.is`

DEPARTMENT OF MATHEMATICAL SCIENCES, NTNU, SENTRALBYGG 2, ALFRED GETZ VEI 1, 7034 TRONDHEIM, NORWAY

Email address: `john.fornass@ntnu.no`

SCHOOL OF MATHEMATICS AND STATISTICS, HENAN UNIVERSITY, KAIFENG 475001, HENAN PROVINCE, PEOPLE'S REPUBLIC OF CHINA -AND- DEPARTMENT OF MATHEMATICAL SCIENCES, NTNU, SENTRALBYGG 2, ALFRED GETZ VEI 1, 7034 TRONDHEIM, NORWAY

Email address: `jujie.wu@ntnu.no`

Email address: `99jujiwu@tongji.edu.cn`