



HAL
open science

Multi-Component Reaction for the Preparation of 1,5-Disubstituted 1,2,3-Triazoles by In-Situ Generation of Azides and Nickel-Catalyzed Azide-Alkyne Cycloaddition

Virgyl Camberlein, Nicolas Kraupner, Nour Bou Karroum, Emmanuelle Lipka, Rebecca Deprez-Poulain, Benoit Deprez, Damien Bosc

► **To cite this version:**

Virgyl Camberlein, Nicolas Kraupner, Nour Bou Karroum, Emmanuelle Lipka, Rebecca Deprez-Poulain, et al.. Multi-Component Reaction for the Preparation of 1,5-Disubstituted 1,2,3-Triazoles by In-Situ Generation of Azides and Nickel-Catalyzed Azide-Alkyne Cycloaddition. *Tetrahedron Letters*, In press, 10.1016/j.tetlet.2021.153131 . hal-03207833

HAL Id: hal-03207833

<https://hal.science/hal-03207833>

Submitted on 26 Apr 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Multi-Component Reaction for the Preparation of 1,5-Disubstituted 1,2,3-Triazoles by In-Situ Generation of Azides and Nickel-Catalyzed Azide-Alkyne Cycloaddition.

Virgyl Camberlein,^{a,1} Nicolas Kraupner,^{a,1} Nour Bou Karroum,^a Emmanuelle Lipka,^{b,c}
Rebecca Deprez-Poulain,^{a,d} Benoit Deprez,^{a,d} and Damien Bosc*^a

^aUniv. Lille, Inserm, Institut Pasteur de Lille, U1177—Drugs and Molecules for Living Systems, Lille, France

^bUniv. Lille, Inserm, RID-AGE U1167, F-59000, Lille, France

^cUFR Pharmacie, Laboratoire de Chimie Analytique, BP 83, F-59006, Lille, France

^dInstitut Universitaire de France, F- 75231, Paris, France.

^eEuropean Genomic Institute for Diabetes, EGID, University of Lille, F-59000, France

¹These authors contributed equally to this work.

Abstract


An efficient one-pot procedure combining bromide conversion into azide followed by NiAAC for the preparation of 1,5-disubstituted 1,2,3-triazoles has been developed. This procedure prevents the use of isolated azides, which are insufficiently commercially available and could be potentially unstable and difficult to handle. Moreover, this one-pot method tolerates a broad range of functional moieties including ester, carbamate or alcohol. Diverse 1,5-disubstituted 1,2,3-triazoles can be obtained from functionalized aryl and alkyl alkynes and bromides with modest to excellent yields and regioselectivities. This procedure will enable the synthesis of libraries of functionalizable 1,5-disubstituted 1,2,3-triazoles particularly helpful for diverse applications such as medicinal chemistry and chemical biology purposes.

click chemistry ; molecular diversity ; multicomponent reactions ; regioselectivity ; synthetic methods

Introduction

Spearhead of the heterocycles generated by click chemistry, 1,2,3-triazoles are broadly found in several domains such as materials science,^[1] organometallic science^[2] but also chemical biology^[3] and medicinal chemistry^[4]. This considerable importance is due to the favorable properties linked to this heteroaromatic ring but also thanks to the easiness of access of the 1,4-disubstituted 1,2,3-triazoles by copper-catalyzed azide-alkyne cycloaddition (CuAAC).^[5] This methodology is indeed powerful with excellent regioselectivity, high yield and is easy to implement. Noteworthy, the 1,5-disubstituted 1,2,3-triazoles are less represented. As there is a high interest for 1,5-disubstituted regioisomers in drug discovery as disulfide bond surrogate^[6] or cis-peptide bond mimetic^[7] or enzyme modulator^[8], it is valuable to facilitate the access of this scaffold. Several teams have worked on this purpose and to date there are few methods to synthesize 1,5-disubstituted 1,2,3-triazoles, which are for most of them with very limited scope. The main methodologies use azide-alkyne cycloadditions either mediated by metal acetylide or metal catalysts.^[9] The alkynes can also be replaced by nitroolefins with the use of metallic additives to afford 1,5-disubstituted 1,2,3-triazoles substituted by aryl groups.^[10] Moreover, in a lesser extent, other functional moieties have also been used for playing the role of dipolarophile in 1,5-disubstituted 1,2,3-triazole synthesis.^[11] An alternative method based on a cascade Michael addition was also described for the construction of 1,5-disubstituted 1,2,3-triazoles.^[12] Regrettably, most of these methodologies have scope constraints. Besides, most of them require the handling of isolated organic azides, which are not often commercially available and are known to be unstable, especially those of low molecular weight.^[13]

In order to bypass these drawbacks, only few methods, most of them being multicomponent, have been developed.^[14] These methodologies have the advantage to be safer by preventing the manipulation of organic azides. However, they suffer from a very limited substrate scope, often requesting high temperatures, strict inert atmosphere conditions or an inconvenient sequential pattern, all of which have confined their use. In order to provide an extensive procedure that circumvent all these drawbacks and that afford functionalized 1,5-disubstituted 1,2,3-triazoles suitable for diverse applications, we investigated the feasibility to implement an efficient and safe one-pot, two-step method. This procedure would allow the in situ bromides conversion by nucleophilic substitution with sodium azide into azides, which react consecutively with alkynes by NiAAC (Scheme 1).




Scheme 1. Strategies for the synthesis of 1,5-disubstituted 1,2,3-triazoles.

Results and discussion

First we performed the screening of different reaction conditions on the model substrates 4-tert-butylbenzyl bromide **1a** and phenylacetylene **2A** (Table 1). Reaction conditions close to the recently published NiAAC ones^[9g] were applied with the combination of bromide **1a** (1.0 equiv), sodium azide (1.0 equiv), alkyne **2A** (1.2 equiv) in presence of nickelocene (Cp₂Ni) precatalyst (20 mol%), bidentate Xantphos ligand (20 mol%) and Cs₂CO₃ (1.0 equiv) at room temperature for 2 h in water. Under these conditions, the bromide **1a** was converted into azide but the cycloaddition was not observed (entry 1). To facilitate the triazole formation, several conditions were screened. First, dimethylformamide, often used as a solvent for azide formation from bromide and shown to be efficient for the NiAAC, was evaluated (entry 2). Pleasantly, this solvent allowed the triazole formation with good yield (71 %) contrary to other solvents such as dichloromethane, ethanol and tetrahydrofuran (entries 2-5). In order to improve the reaction conversion into triazole, the reaction vessel was slightly heated at 50 °C under microwave irradiation for 1 h (entry 6). This heating condition was beneficial for the regioselectivity (from 88:12 to 93:7) without compromising the yield (entries 2 vs 6). Higher temperature (80 °C or 100 °C) resulted to a lower yield and a poorer regioselectivity (entries 7 and 8), probably due to the catalyst degradation. Increasing reaction time to 4 h improved yield (86 %, entry 9) but slightly decreased the regioselectivity (from 93:7 to 92:8). In the attempt to improve the yield of the 1,5-disubstituted 1,2,3-triazole **3aA**, the bromide **1a** and the sodium azide were defined as excess reagents (1.2 equiv and 1.5 equiv respectively) (entries 10). Appealingly, after 4 h under microwave irradiation at 50 °C, the calculated yield of triazole **3aA** was excellent (89 %) with no secondary products and

excellent regioselectivity (92:8). To investigate the mechanism of this multicomponent method, 4-tert-butylbenzyl bromide **1a** was submitted to sodium azide at room temperature for 2 h in DMF with or without the Cp₂Ni/Xantphos catalytic system. For both conditions, 4-tert-butylbenzyl azide was formed in quantitative yield. Thus, the nickelocene precatalyst in association with its Xantphos ligand has no influence on the SN₂ event and only has positive effect for the azide-alkyne cycloaddition and its regioisomerism.

Table 1. Condition screening^a.



Entry	Solvent	Temp (°C)	Time (h)	<i>c</i> Yield (%) ^c	Ratio 3aA/4aA ^d
1	H ₂ O	rt	2	— ^e	— ^e
2	DMF	rt	2	71	88:12
3	EtOH	rt	2	— ^e	— ^e
4	DCM	rt	2	— ^e	— ^e
5	THF	rt	2	— ^e	— ^e
6	DMF	50 °C (MW)	1	73	93:7
7	DMF	80 °C (MW)	1	49	64:36
8	DMF	100 °C (MW)	1	— ^e	— ^e
9	DMF	50 °C (MW)	4	86	92:8
10 ^b	DMF	50 °C (MW)	4	89	92:8

^aReaction conditions: Run on a 0.5 mmol scale of alkyne **2A** (1.2 equiv), bromide **1a** (1.0 equiv), NaN₃ (1.0 equiv), solvent (2 mL), Cp₂Ni (20 mol%), Xantphos (20 mol%), Cs₂CO₃ (1.0 equiv). ^bRun on a 0.5 mmol scale of alkyne **2A** (1.0 equiv), bromide **1a** (1.2 equiv), NaN₃ (1.5 equiv), solvent (2 mL), Cp₂Ni (20 mol%), Xantphos (20 mol%), Cs₂CO₃ (1.0 equiv). ^cCalculated yield of **3aA** determined by LCMS analysis from the crude product. ^dDetermined by LCMS analysis from the crude product. ^eNo triazole formation. Cp, cyclopentadienyl; *c*Yield, calculated yield; DCM, dichloromethane; DMF, dimethylformamide; MW, microwave irradiation; temp, temperature; rt, room temperature.

After determining the optimized conditions for this one-pot procedure, we explored the scope with various alkynes and bromides. The integrity of the 1,5-disubstituted 1,2,3-triazole template was validated by NMR NOESY and HSQC experiments to corroborate the distinctive ¹³C chemical shift of the triazole CH at ~133 ppm.^[14]

The evaluation of substrate scope with various aromatic (**2A–2E**) and aliphatic alkynes (**2F–2H**) in reaction with bromide **1a** is summarized in Table 2. Except for alkynes **2D** and **2E**, the attempts to separate and isolate the two regioisomers **3** and **4** were not fruitful either by classical flash column chromatography, or by C-18 reversed-phase flash column chromatography with the use of classical or pentafluorophenyl columns. However, because of the high regioselectivity rates associated with this methodology (from 66:34 to >99:1), flash chromatography was still performed to afford **3** in mixture with **4** in order to evaluate the efficiency of this one-pot multicomponent procedure. Modest to excellent calculated yields ranging from 27% to 84% were obtained. Besides, in order to isolate and characterize properly the 1,5-regioisomers **3**, the reaction crude mixtures were also purified by semi-preparative supercritical fluid chromatography (SFC) and comparable isolated yields (24% to 72%) were obtained.

The regioselectivity of the reactions was determined by supercritical fluid chromatography (SFC) analysis of the crude reaction mixtures obtained after the work-up of the reactions carried out with alkynes **2A–2E** and **2H**, or according to the ¹H-NMR spectra for reactions carried out with alkynes **2F** and **2G**, whose separation of the two regioisomers by SFC was not optimal. Modest to excellent regioselectivities were obtained. Considering the regioselectivity obtained with the *p*-nitrophenylacetylene **2B** (ratio 66:34) compared to the phenylacetylene **2A** (ratio 89:11) and then the *p*-fluorophenylacetylene **2C** (ratio 91:9) and the *p*-methoxyphenylacetylene **2D** (ratio >99:1), a positive mesomeric effect on the phenyl ring seems to induce a positive contribution to the regioselectivity without having an effect on the yield. Noteworthy, heteroaryl and alkyl chains are accepted for this procedure (**2E–2H**). Interestingly, the scope was widened by hydroxyl and carbamate moieties, which were compatible with this process.

Table 2. Substrate scope with various alkynes^a.

1a	2A-2H	3A-3A'H	4A-4A'H
	3A	72 (64)	89:11
	3B	24 (37)	66:34
	3C	34 (84)	91:9
	3D	76 ^b	>99:1
	3E	73 ^b	92:8
	3F	36 (27)	86:14 ^c
	3G	56 (70)	89:11 ^c
	3H	34 (34)	67:33

^aReaction conditions: Run on a 0.3–0.6 mmol scale of alkyne **2** with **1a** (1.2 equiv), NaN₃ (1.5 equiv), Cp₂Ni (20 mol%), Xantphos (20 mol%), Cs₂CO₃ (1.0 equiv), DMF (2.0 mL), 50 °C (MW), 4 h. Isolated yields of **3** obtained after SFC purification; **3/4** ratio determined from the crude product by SFC analysis

unless noted otherwise; in brackets: calculated yields of **3** obtained after flash column chromatography purification (in mixture with **4**); ^bIsolated yield of **3** obtained after flash column chromatography purification; ^cDetermined by ¹H-NMR. Boc, tert-butoxycarbonyl; Cp, cyclopentadienyl; t-Bu, tert-butyl.

The substrate scope was also examined with various bromides (Table 3). Aromatic benzyl bromide possessing either electron-donating (**1a** and **1b**) or electron-withdrawing (**1c–1g**) groups are tolerated and afforded the 1,5-regioisomers (**3aA–3gA**) with modest to high yields both after flash chromatography purification (calculated yield: 64% – 79%) and after semi-preparative SFC purification (isolated yield: 31% – 84%). The ortho isomer **3gA** was obtained with the lowest yield in the series, probably due to steric effect. Besides, regioisomerism of the phenyl substitution (**1e–1g**) has only minor impact on the yield and does not influence the regioselectivity. Strikingly, high to excellent regioselectivities from 85:15 to 97:3 ratios were also obtained. Interestingly, the scope of this one-pot procedure could be efficiently enlarged with non-saturated bromide (**1i–1l**). Noteworthy, functional groups such as carbamate-protected amine and ester are well tolerated with these reaction conditions (**1k–1l**) providing the related 1,5-disubstituted 1,2,3-triazoles with high to excellent yields and regioselectivities.

Table 3. Substrate scope with various alkynes^a.

R-Br		2A		3aA-3lA		4aA-4lA	
	1a		2A		3aA		4aA
Reaction conditions: NaN_3 , Cp_2Ni (20 mol%), Xantphos (20 mol%), Cs_2CO_3 (1.0 equiv), DMF, 50 °C (MW), 4 h.							
	1b		2A		3bA		4bA
	1c		2A		3cA		4cA
	1d		2A		3dA		4dA
	1e		2A		3eA		4eA
	1f		2A		3fA		4fA
	1g		2A		3gA		4gA
	1h		2A		3hA		4hA
	1i		2A		3iA		4iA
	1j		2A		3jA		4jA
	1k		2A		3kA		4kA

^aReaction conditions: Run on a 0.49 mmol scale of alkyne **2A** with **1** (1.2 equiv), NaN_3 (1.5 equiv), Cp_2Ni (20 mol%), Xantphos (20 mol%), Cs_2CO_3 (1.0 equiv), DMF (2.0 mL), 50 °C (MW), 4 h.

Isolated yields of **3** obtained after SFC purification; **3/4** ratio determined from the crude product by SFC analysis unless noted otherwise; in brackets: calculated yields of **3** obtained after flash column chromatography purification (in mixture with **4**); ^bIsolated yield of **3** obtained after flash column chromatography. Boc, tert-butoxycarbonyl; Me, methyl; Ph, phenyl; t-Bu, tert-butyl.

Briefly, we succeeded in providing a procedure to efficiently access diverse 1,5-disubstituted 1,2,3-triazoles in a fast and simple manner. Remarkably, one of the advantages of this one-pot reaction is the in situ generation of azides from bromides, which are commercially available with a large diversity. Thus, this one-pot method prevents the sequential and time-consuming synthesis and isolation of azides, which could be unstable and are rarely commercially available. It makes this procedure very convenient, especially when sensitive low-molecular weight azides are required. A diverse range of functional moieties including N-protected amines, hydroxyls and esters are tolerated by these reaction conditions. This method will enable the synthesis of libraries of functionalizable 1,5-regioisomers especially useful for drug discovery and chemical biology purposes.

Acknowledgements

We are grateful to the institutions that support our laboratory: Université de Lille, Institut Pasteur de Lille, INSERM. The authors thank Nicolas Lefebvre for technical assistance. The authors thank the members of the Laboratoire d'Application de Résonance Magnétique Nucléaire (LARMN), Lille, France for NMR acquisitions and for their technical assistance. The postdoctoral fellowship of N.B.K. is funded by I-SITE-ULNE (contract CSB0120_ I-SITE Molecular Design For Health). V.C. is a recipient of a doctoral fellowship from Université de Lille and Universität des Saarlandes. N.K. is a recipient of a doctoral fellowship from Université de Lille.

References

- [1] a) P. Nezhad-Mokhtari, M. Ghorbani., L. Roshangar, J. Soleimani Rad, *Eur. Polym. J.* **2019**, *117*, 64–76; b) M. Arslan, G. Acik, M. A. Tasdelen, *Polym. Chem.* **2019**, *10*, 3806–3821.
- [2] a) P.-Z. Li, -J. Wang, Y. Zhao, *Coord. Chem. Rev.* **2019**, *380*, 484–518; b) D. Huang, P. Zhao, D. Astruc, *Coord. Chem. Rev.* **2014**, *272*, 145–165.
- [3] a) Kenry, B. Liu, *Trends Chem.* **2019**, *1*, 763–778; b) E. Kim, H. Koo, *Chem. Sci.* **2019**, *10*, 7835–7851.
- [4] a) A. Rani, G. Singh, A. Singh, U. Maqbool, G. Kaur, J. Singh, *RSC Adv.* **2020**, *10*, 5610–5635; b) E. Bonandi, M. S. Christodoulou, G. Fumagalli, D. Perdicchia, G. Rastelli, D. Passarella, *Drug Discov. Today* **2017**, *22*, 1572–1581; c) D. Bosc, V. Camberlein, R. Gealageas, O. Castillo-Aguilera, B. Deprez, R. Deprez-Poulain, *J. Med. Chem.* **2020**, *63*, 3817–3833; d) R. Deprez-Poulain, N. Hennuyer, D. Bosc, W. G. Liang, E. Enée, X. Marechal, J. Charton, J. Totobenazara, G. Berte, J. Jahklal, T. Verdelet, J. Dumont, S. Dassonneville, E. Woittrain, M. Gauriot, C. Paquet, I. Duplan, P. Hermant, F.-X. Cantrelle, E. Sevin, M. Culot, V. Landry, A. Herledan, C. Piveteau,

- G. Lippens, F. Leroux, W.-J Tang, P. van Endert, B. Staels, B. Deprez, *Nat. Commun.* **2015**, *6*, 8250.
- [5] a) V. V. Rostovtsev, L. G. Green, V. V. Fokin, K. B. Sharpless, *Angew. Chem. Int. Ed Engl.* **2002**, *41*, 2596–2599; b) C. W. Tornøe, C. Christensen, M. Meldal, *J. Org. Chem.* **2002**, *67*, 3057–3064.
- [6] A. M. White, S. J. Veer, G. Wu, P. J. Harvey, K. Yap, G. J. King, J. E. Swedberg, C. K. Wang, R. H. P. Law, T. Durek, D. J. Craik, *Angew. Chem. Int. Ed.* **2020**, *59*, 11273–11277.
- [7] O. Kracker, J. Góra, J. Krzciuk-Gula, A. Marion, B. Neumann, H.-G. Stammer, A. Nieß, I. Antes, R. Latajka, N. Sewald, *Chem. - Eur. J.* **2018**, *24*, 953–961.
- [8] [a] D. P. Becker, M. R. Lutz, S. Flieger, A. Colorina, J. Wozny, N. S. Hosmane, *ChemMedChem*, **2020**, *15*, 1897–1908; [b] S. C. C. Lucas, S. J. Atkinson, P. Bamborough, H. Barnett, C.-W. Chung, L. Gordon, D. J. Mitchell, A. Phillipou, R. K. Prinjha, R. J. Sheppard, N. C. O. Tomkinson, R. J. Watson, E. H. Demont, *J. Med. Chem.* **2020**, *63*, 5212–5241.
- [9] a) G. Himbert, D. Frank, M. Regit, *Chem. Ber.* **1976**, *109*, 370–394; b) A. Krasinski, V. V. Fokin, K. B. Sharpless, *Org. Lett.* **2004**, *6*, 1237–1240; c) C. D. Smith, M. F. Greaney, *Org. Lett.* **2013**, *15*, 4826–4829; d) B. C. Boren, S. Narayan, L. K. Rasmussen, L. Zhang, H. Zhao, Z. Lin, G. Jia, V. V. Fokin, *J. Am. Chem. Soc.* **2008**, *130*, 8923–8930; e) L. Zhang, X. Chen, P. Xue, H. H. Y. Sun, I. D. Williams, K. B. Sharpless, V. V. Fokin, V. V.; G. Jia, *J. Am. Chem. Soc.* **2005**, *127*, 15998–15999; f) L. Hong, W. Lin, F. Zhang, R. Liu, X. Zhou, *Chem. Commun.* **2013**, *49*, 5589; g) W. G. Kim, M. E. Kang, J. B. Lee, M. H. Jeon, S. Lee, J. Lee, B. Choi, P. M. S. D. Cal, S. Kang, J.-M. Kee, G. J. L. Bernardes, J.-U. Rohde, W. Choe, S. Y. Hong, *J. Am. Chem. Soc.* **2017**, *139*, 12121–12124.
- [10] a) L. Maiuolo, B. Russo, V. Algieri, M. Nardi, M. L. Di Gioia, M. A. Tallarida, A. De Nino, *Tetrahedron Lett.* **2019**, *60*, 672–674; b) Y.-C. Wang, Y.-Y. Xie, H.-E. Qu, H.-S. Wang, Y.-M. Pan, F.-P. Huang, *J. Org. Chem.* **2014**, *79*, 4463–4469; c) H. M. Nanjundaswamy, H. Abrahamse, *Synth. Commun.* **2015**, *45*, 967–974; d) A. De Nino, P. Merino, V. Algieri, M. Nardi, M. Di gioia, B. Russo, M. Tallarida, L. Maiuolo, *Catalysts* **2018**, *8*, 364.
- [11] a) Ahsanullah; P. Schmieder, R. Kühne, J. Rademann, *Angew. Chem. Int. Ed.* **2009**, *48*, 5042–5045; b) X. Zhang, K.P. Rakesh, H.-L. Qin, *Chem. Commun.* **2019**, *55*, 2845–2848; c) L. Wu, Y. Chen, J. Luo, Q. Sun, M. Peng, Q. Lin, *Tetrahedron Lett.* **2014**, *55*, 3847–3850; d) A. Banday, V. Hruby, *Synlett* **2014**, *25*, 1859–1862; e) N. Kumar, M. Y. Ansari, R. Kant, A. Kumar, *Chem. Commun.* **2018**, *54*, 2627–2630.
- [12] G. Cheng, X. Zeng, J. Shen, X. Wang, X. Cui, *Angew. Chem. Int. Ed.* **2013**, *52*, 13265–13268.
- [13] S. Bräse, C. Gil, K. Knepper, V. Zimmermann, *Angew. Chem. Int. Ed.* **2005**, *44*, 5188–5240.

- [14] a) H.-W Bai, Z.-J. Cai, S.-Y. Wang, S.-J. Ji, *Org. Lett.* **2015**, *17*, 2898–2901; b) J.-P. Wan, S. Cao, Y. Liu, *J. Org. Chem.* **2015**, *80*, 9028–9033; c) S. Cao, Y. Liu, C. Hu, C. Wen, J. Wan, *ChemCatChem* **2018**, *10*, 5007–5011; d) W. Huang, C. Zhu, M. Li, Y. Yu, W. Wu, Z. Tu, H. Jiang, *Adv. Synth. Catal.* **2018**, *360*, 3117–3123; e) S. Dey, T. A. Pathak, *RSC Adv.* **2014**, *4*, 9275; f) J. R. Johansson, P. Lincoln, B. Nordén, N. Kann, *J. Org. Chem.* **2011**, *76*, 2355–2359.
- [15] X. Creary, A. Anderson, C. Brophy, F. Crowell, Z. Funk, *J. Org. Chem.* **2012**, *77*, 8756–8761.