



**HAL**  
open science

## Gaia EDR3 Proper Motions of Milky Way Dwarfs. I. 3D Motions and Orbits

Hefan Li, Francois Hammer, Carine Babusiaux, Marcel S. Pawlowski, Yanbin Yang, Frederic Arenou, Cuihua Du, Jianling Wang

► **To cite this version:**

Hefan Li, Francois Hammer, Carine Babusiaux, Marcel S. Pawlowski, Yanbin Yang, et al.. Gaia EDR3 Proper Motions of Milky Way Dwarfs. I. 3D Motions and Orbits. *The Astrophysical Journal*, 2021, 916 (1), pp.8. 10.3847/1538-4357/ac0436 . hal-03210349

**HAL Id: hal-03210349**

**<https://hal.science/hal-03210349>**

Submitted on 9 Feb 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Gaia EDR3 proper motions of Milky Way dwarfs I: 3D Motions and Orbits**HEFAN LI,<sup>1</sup> FRANCOIS HAMMER,<sup>2</sup> CARINE BABUSIAUX,<sup>3,2</sup> MARCEL S. PAWLOWSKI,<sup>4</sup> YANBIN YANG,<sup>2</sup>  
FREDERIC ARENOU,<sup>2</sup> CUIHUA DU,<sup>1</sup> AND JIANLING WANG<sup>5</sup><sup>1</sup>*School of Physical Sciences, University of Chinese Academy of Sciences, Beijing 100049, P. R. China*<sup>2</sup>*GEPI, Observatoire de Paris, Université PSL, CNRS, Place Jules Janssen 92195, Meudon, France*<sup>3</sup>*Université de Grenoble-Alpes, CNRS, IPAG, F-38000 Grenoble, France*<sup>4</sup>*Leibniz-Institut fuer Astrophysik Potsdam (AIP), An der Sternwarte 16, D-14482 Potsdam Germany*<sup>5</sup>*CAS Key Laboratory of Optical Astronomy, National Astronomical Observatories, Beijing 100101, China*

## ABSTRACT

Based on Gaia Early Data Release 3 (EDR3), we estimate the proper motions for 46 dwarf galaxies of the Milky Way. The uncertainties in proper motions, determined by combining both statistical and systematic errors, are smaller by a factor 2.5, when compared with Gaia Data Release 2. We have derived orbits in four Milky Way potential models that are consistent with the MW rotation curve, with total mass ranging from  $2.8 \times 10^{11} M_{\odot}$  to  $15 \times 10^{11} M_{\odot}$ . Although the type of orbit (ellipse or hyperbola) are very dependent on the potential model, the pericenter values are firmly determined, largely independent of the adopted MW mass model. By analyzing the orbital phases, we found that the dwarf galaxies are highly concentrated close to their pericenter, rather than to their apocenter as expected from Kepler's law. This may challenge the fact that most dwarf galaxies are Milky Way satellites, or alternatively indicates an unexpected large number of undiscovered dwarf galaxies lying very close to their apocenters. Between half and two thirds of the satellites have orbital poles that indicate them to orbit along the Vast Polar Structure (VPOS), with the vast majority of these co-orbiting in a common direction also shared by the Magellanic Clouds, which is indicative of a real structure of dwarf galaxies.

*Keywords:* galaxies: dwarf — galaxies: kinematics and dynamics

## 1. INTRODUCTION

Gaia DR2 (Helmi et al. 2018) revolutionized the knowledge of Milky Way (MW) dwarf galaxies by revealing their proper motions (PMs) (Fritz et al. 2018). Because they lie in our neighborhood, MW dwarf galaxies are unique galaxies for which we can firmly establish 6D phase diagrams (3D locations and 3D velocities). Gaia EDR3 (Gaia Collaboration et al. 2016, 2020) provides a significant step forward by improving PM precision by a factor  $\sim 2$ , including for very faint dwarf galaxies.

In the meantime between Gaia DR2 and EDR3 epochs, several new dwarf galaxies have been discovered, or have been spectroscopically observed to derive their radial velocities, which is the essential complement to Gaia PMs for establishing their 3D motions (see references in Table 1). Combined with Gaia EDR3 this allows the determination of accurate orbits for several tens of dwarf galaxies much less massive than the classical dwarf spheroidal galaxies<sup>1</sup>, a yet unprecedented number. In principle, one may now investigate their past and future history.

However, besides observed (instantaneous) quantities (distances and velocities), evaluating integrated orbital quantities (pericenter, apocenter, orbit shape and eccentricity) requires the knowledge of the potential, i.e., namely that of the main Galaxy, the MW. It has been shown (Fritz et al. 2018; Hammer et al. 2020) that the determination of dwarf galaxy orbital quantities is limited by Gaia DR2 PM uncertainties, but perhaps even more by the systematics due to the inaccurate knowledge of the MW mass profile. Reducing PM errors by at least a factor  $\sim 2$  with EDR3, leads to a problem for which the limited knowledge of the MW mass-profile dominates the uncertainties in the derived orbital properties. Another possible limitation is the impact of the second massive body in the MW halo, the Large Magellanic Cloud (LMC), which can, if it has a total mass exceeding  $10^{11} M_{\odot}$ , also affect stellar streams such as the Orphan stream (Erkal et al. 2019).

In other words, the MW potential uncertainties hamper our knowledge of dwarf galaxy orbits. The circular velocity curve of the MW has been accurately provided by Eilers et al. (2019) and Mróz et al. (2019). This result can be associated to other possible probes of the MW potential, for example dwarf galaxy (Cautun et al. 2020) or globular cluster motions Wang & Hammer (2021), leading to quite discrepant values for the total MW mass. This is not unexpected because these methods depend on whether or not these additional probes share a similar equilibrium than rotating disk stars in the MW potential. Jiao et al. (2021) showed that MW mass profiles derived from the rotation curve are also affected by the choice of the dark-matter mass profile. In a generalized study, they determine the range of total MW mass that can be consistent with the MW rotation curve fit. They found that the MW mass can be as small as  $2.8 \times 10^{11} M_{\odot}$  (see also de Salas et al. 2019 and Karukes et al. 2020) or as large as  $15 \times 10^{11} M_{\odot}$ . This somewhat constraints the available range of MW potential, but it also emphasizes that it is still required to consider a wide range of total MW masses when performing orbital analyses.

Dwarf galaxies present a large variety of properties in mass, radius and velocity dispersion (see an analysis in Hammer et al. 2019). Few of them (Hercules, Tucana III) show sign of tidal disruption, which for Tucana III could be related to its passage at low pericenter (Li et al. 2018). More enigmatic is the fact that most dwarf galaxies appear to lie and to orbit within a gigantic disk almost perpendicular to the MW disk, the Vast Polar Structure (Pawlowski & Kroupa 2014). Dwarf galaxies appear also to be excessively close to their pericenters (Fritz et al. 2018; Simon 2018). Their internal properties also show unusual properties. For example, their gravity at half-light radius declines with Galactic distance (Hammer et al. 2019). DSphs have had their own complex star-formation history, which is revealed when they possess two distinct main stellar populations (Sculptor and Ursa Minor, Tolstoy et al. 2004; Pace et al. 2020), or more than two (Carina and Fornax, Savino et al. 2015; de Boer et al. 2012). However, Draco appears to possess only a single stellar population (Aparicio et al. 2001).

<sup>1</sup> Following Tolstoy et al. (2009) MW dwarfs include dwarf spheroidals (dSphs) that are the 9 classical dSphs and Canes Venatici I, and ultra-faint dwarfs (UFDs), which are less luminous, with  $L_V < 10^5 M_{\odot}$ , or  $M_V > -7.5$ .

This paper aims to provide updated and new proper motion measurements for MW dwarf galaxies, and determines their resulting integrated orbital parameters in the widest range of MW masses that remain consistent with the MW rotation curve. The goal is to avoid possible systematics or biases that could affect our understanding of dwarf galaxies. Section 2 describes the dataset, the determination of observed properties deduced from Gaia EDR3 and radial velocity measurements. In Section 3, we have calculated integrated orbit properties for four different MW potentials derived from the [Jiao et al. \(2021\)](#) analysis. Results are given and discussed in Section 4, including the dwarf galaxy 3D phase-diagram, the VPOS, and the fraction of dwarf galaxies lying near their pericenters.

## 2. DATA

### 2.1. *Sample of dwarf galaxies*

The sample has been selected from the literature ([Fritz et al. 2018](#); [Simon 2019](#); [McConnachie & Venn 2020](#)), with the supplementary request to have at least four spectroscopically measured stars detected by Gaia EDR3 (the full reference list is given in Table 1). To this we have added five dwarf galaxies having three measured stars or less to investigate the behavior of PM accuracy at low signal (name in *italic* in Table 1). We have also added Eridanus II, even though its very large distance precludes accurate velocity measurements. Following [Fritz et al. \(2018\)](#) we have kept Crater I and Draco II in the dwarf galaxy list, since their precise nature (globular cluster or dwarf galaxy) is still under discussion. The total list includes 46 galaxies.

### 2.2. *Proper Motions, uncertainties, and comparison with other studies*

We have first selected dwarf galaxy member stars from literature catalogues based, among others, on their radial velocities (see references in Table 1), for which we have calculated the median. We then have further selected only stars with Gaia parallaxes and proper motions consistent with the median value of the dwarf galaxy at  $5\sigma$ . Finally, stars with a dubious astrometry have been also removed if the Gaia renormalized unit weight error (ruwe) is larger than 1.4 ([Lindgren et al. 2020](#)).

To compute the dwarf galaxy proper motions and their errors, we have adopted the method described in [Vasiliev \(2019\)](#) who used both the statistical covariance matrix (as derived from the formal errors and correlations provided in the Gaia catalogue) and the systematic one derived from the spatial correlations. The proper motion covariance function has been constructed using the formulae of [Vasiliev \(2019\)](#) adapted to EDR3 and using the values of [Lindgren et al. \(2020\)](#):

$$V_{\mu}(\theta) = 292 \exp(-\theta/12^{\circ}) + 258 \exp(-\theta/0.25^{\circ}) \mu\text{as}^2 \text{yr}^{-2} \quad (1)$$

[McConnachie & Venn \(2020\)](#) promptly published PM estimates just after the Gaia EDR3. Appendix A compares their values to those in Table 1, which show an excellent agreement, except for error bars that are systematically smaller in [McConnachie & Venn \(2020\)](#) by large factors, almost always larger than 2 but that can reach 11 (for Sculptor), and even 20 for Fornax. We interpret this as a difference in the treatment of systematics (not accounted by [McConnachie & Venn \(2020\)](#)), which is still an important limitation for determining PMs of dwarf galaxies, and especially to evaluate their orbital motions. Figure A1 indicates the name of the few dwarf galaxies for which both studies show disagreement, which further points out the uncertain determination of PMs for 5 galaxies with 3 stars or less: Aquarius II, Columba I, Horologium II, Pisces II, and Reticulum III.

Appendix A also compares EDR3 results to [Fritz et al. \(2018\)](#) DR2 values. EDR3 and DR2 PM values are often consistent within the pretty conservative DR2 and EDR3 error bars (see Figure A2)

. However, for some dwarf galaxies there is some changes of the PM from DR2 to EDR3. Specifically, changes exceeding the DR2  $1\sigma$  error bars from [Fritz et al. \(2018\)](#) are affecting RA PMs of Aquarius II, Crater II, Draco, Grus I, Hercules, Leo IV, Pisces II, Segue I, Segue II, Tucana III, UMa I, and UMi, and for DEC PMs, Eridanus II, Hydrus, and Triangulum II. Top panels of Figure A2 also show that on average, typical errors (including systematics) have decreased by a factor of about 2.5 from DR2 to EDR3. Note however that the number of dwarf galaxies with different measurements from DR2 to EDR3 is consistent with expectations: there are 37 satellites in common between the two data sets, and one expects on average 32% of data points to disagree by  $1\sigma$  or more, which corresponds to about 12 dwarf galaxies in this case. This number well consistent with the numbers of measurements differing by  $1\sigma$  or more.

**Table 1.** Origin data of dwarf galaxies.

name	$dm^a$	$N_{\text{star}}$	$\varpi$	$\mu_{\alpha^*}$	$\mu_{\delta}$	$\rho_{\mu_{\alpha^*}}^{\mu_{\delta}}$	$rv$	Ref.
(1)	(2)	(3)	mas	(mas yr <sup>-1</sup> )	(mas yr <sup>-1</sup> )		(km s <sup>-1</sup> )	(9)
AntII	$20.6 \pm 0.11$	157	$-0.017 \pm 0.018$	$-0.101 \pm 0.02$	$0.113 \pm 0.02$	0.044	$294.0 \pm 0.1$	1
<i>AquII</i>	$20.17 \pm 0.07$	2	$-0.845 \pm 0.564$	$0.647 \pm 0.588$	$-0.298 \pm 0.548$	0.267	$-74.4 \pm 1.0$	2
BooI	$19.1 \pm 0.07$	37	$-0.04 \pm 0.049$	$-0.307 \pm 0.052$	$-1.157 \pm 0.043$	-0.107	$100.3 \pm 0.1$	3,4
BooII	$18.12 \pm 0.05$	4	$0.215 \pm 0.148$	$-2.273 \pm 0.151$	$-0.361 \pm 0.115$	-0.164	$-114.2 \pm 1.4$	5
CVenI	$21.62 \pm 0.06$	53	$-0.019 \pm 0.059$	$-0.084 \pm 0.052$	$-0.127 \pm 0.037$	0.149	$26.4 \pm 0.2$	6,3
CVenII	$21.02 \pm 0.05$	15	$0.15 \pm 0.124$	$-0.138 \pm 0.111$	$-0.32 \pm 0.082$	0.333	$-129.5 \pm 0.7$	7
CarI	$20.13 \pm 0.1$	882	$-0.004 \pm 0.018$	$0.533 \pm 0.021$	$0.12 \pm 0.021$	-0.009	$221.8 \pm 0.1$	8,9
CarII	$17.79 \pm 0.04$	18	$0.05 \pm 0.036$	$1.887 \pm 0.043$	$0.164 \pm 0.043$	0.024	$480.3 \pm 0.3$	10
CarIII	$17.22 \pm 0.05$	4	$0.041 \pm 0.054$	$3.082 \pm 0.068$	$1.394 \pm 0.072$	0.021	$283.2 \pm 0.6$	10
<i>ColI</i>	$21.31 \pm 0.12$	3	$0.065 \pm 0.133$	$0.189 \pm 0.119$	$-0.556 \pm 0.134$	-0.069	$152.3 \pm 3.7$	11
CberI	$18.12 \pm 0.08$	17	$0.03 \pm 0.056$	$0.374 \pm 0.059$	$-1.699 \pm 0.056$	-0.293	$97.0 \pm 0.6$	7
CraI	$20.82 \pm 0.03$	6	$0.017 \pm 0.151$	$0.056 \pm 0.127$	$-0.122 \pm 0.111$	-0.292	$149.3 \pm 0.7$	12,13
CraII	$20.35 \pm 0.02$	59	$0.004 \pm 0.029$	$-0.072 \pm 0.032$	$-0.123 \pm 0.025$	-0.024	$87.9 \pm 0.1$	14
DraI	$19.57 \pm 0.16$	536	$-0.021 \pm 0.018$	$0.039 \pm 0.02$	$-0.181 \pm 0.02$	0.021	$-292.1 \pm 0.0$	15,16,17
DraII	$16.66 \pm 0.04$	7	$-0.024 \pm 0.105$	$1.011 \pm 0.115$	$0.956 \pm 0.126$	-0.082	$-347.1 \pm 1.3$	18
EriII	$22.82 \pm 0.1$	13	$-0.256 \pm 0.127$	$0.095 \pm 0.133$	$-0.175 \pm 0.164$	-0.213	$74.6 \pm 0.3$	19
FnxI	$20.72 \pm 0.05$	2463	$-0.01 \pm 0.016$	$0.384 \pm 0.018$	$-0.364 \pm 0.019$	-0.014	$55.3 \pm 0.0$	8,20
GruI	$20.4 \pm 0.21$	7	$-0.068 \pm 0.078$	$0.071 \pm 0.056$	$-0.27 \pm 0.078$	0.071	$-139.6 \pm 0.3$	21
GruII	$18.62 \pm 0.21$	45	$-0.032 \pm 0.04$	$0.389 \pm 0.034$	$-1.526 \pm 0.036$	0.159	$-109.9 \pm 0.3$	22
HerI	$20.6 \pm 0.1$	20	$0.11 \pm 0.07$	$-0.051 \pm 0.059$	$-0.332 \pm 0.051$	0.431	$44.7 \pm 0.4$	23
HorI	$19.7 \pm 0.3$	5	$-0.058 \pm 0.045$	$0.865 \pm 0.047$	$-0.601 \pm 0.048$	0.029	$116.5 \pm 0.1$	24
<i>HorII</i>	$19.46 \pm 0.21$	1	$0.469 \pm 0.331$	$0.915 \pm 0.33$	$-0.913 \pm 0.439$	-0.113	$158.3 \pm 11.3$	11
HyaII	$20.89 \pm 0.11$	6	$0.48 \pm 0.246$	$-0.576 \pm 0.28$	$-0.101 \pm 0.212$	0.054	$303.8 \pm 1.0$	12
HyI	$17.2 \pm 0.04$	32	$-0.004 \pm 0.026$	$3.775 \pm 0.027$	$-1.513 \pm 0.027$	0.004	$80.3 \pm 0.2$	25

**Table 1** continued on next page

**Table 1** (*continued*)

name	$dm^a$	$N_{\text{star}}$	$\varpi$	$\mu_{\alpha^*}$	$\mu_{\delta}$	$\rho_{\mu_{\alpha^*}}^{\mu_{\delta}}$	$rv$	Ref.
			mas	(mas yr <sup>-1</sup> )	(mas yr <sup>-1</sup> )		(km s <sup>-1</sup> )	
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
LeoI	22.02 ± 0.13	368	-0.065 ± 0.032	-0.066 ± 0.029	-0.107 ± 0.026	-0.169	283.0 ± 0.1	26,27
LeoII	21.84 ± 0.13	221	0.031 ± 0.037	-0.125 ± 0.039	-0.122 ± 0.036	-0.158	79.6 ± 0.1	28,29
LeoIV	20.94 ± 0.07	8	0.118 ± 0.178	0.007 ± 0.174	-0.261 ± 0.134	-0.168	131.2 ± 0.5	7,30
LeoV	21.14 ± 0.05	7	0.067 ± 0.173	0.118 ± 0.212	-0.387 ± 0.151	-0.137	174.0 ± 0.4	31,30
PhxII	19.63 ± 0.1	5	0.042 ± 0.078	0.501 ± 0.062	-1.199 ± 0.076	-0.389	32.2 ± 1.5	11
<i>PisII</i>	21.31 ± 0.18	3	0.135 ± 0.223	0.675 ± 0.299	-0.631 ± 0.212	-0.016	-227.5 ± 1.1	12
RetII	17.5 ± 0.1	28	0.025 ± 0.029	2.391 ± 0.029	-1.379 ± 0.032	-0.068	63.0 ± 0.3	32
<i>RetIII</i>	19.82 ± 0.31	3	-0.129 ± 0.194	0.519 ± 0.222	-0.173 ± 0.247	0.116	274.4 ± 4.5	11
SgrII	19.23 ± 0.07	7	0.015 ± 0.072	-0.71 ± 0.077	-0.905 ± 0.051	-0.047	-175.3 ± 0.4	33
ScII	19.67 ± 0.13	1405	0.011 ± 0.018	0.096 ± 0.019	-0.159 ± 0.019	-0.020	111.5 ± 0.0	8,34
SegI	16.81 ± 0.19	22	-0.023 ± 0.057	-2.074 ± 0.052	-3.411 ± 0.043	-0.284	208.9 ± 0.7	35
SegII	17.84 ± 0.18	14	-0.075 ± 0.057	1.425 ± 0.061	-0.313 ± 0.052	0.207	-41.0 ± 0.6	36
SxtI	19.89 ± 0.07	511	-0.007 ± 0.019	-0.403 ± 0.021	0.029 ± 0.021	-0.090	224.9 ± 0.1	8,37
TriII	17.27 ± 0.12	7	-0.008 ± 0.076	0.602 ± 0.081	0.085 ± 0.093	0.229	-381.3 ± 0.9	38
TucII	18.82 ± 0.3	15	-0.011 ± 0.037	0.935 ± 0.031	-1.243 ± 0.036	-0.168	-127.8 ± 0.2	21
TucIII	16.99 ± 0.17	44	0.014 ± 0.021	-0.111 ± 0.023	-1.629 ± 0.023	-0.098	-104.3 ± 0.2	39,40
TucIV	18.41 ± 0.18	39	0.038 ± 0.039	0.618 ± 0.036	-1.696 ± 0.038	-0.111	13.7 ± 0.3	22
TucV	18.7 ± 0.36	6	-0.134 ± 0.094	-0.27 ± 0.083	-1.253 ± 0.11	-0.097	-36.7 ± 0.7	22
UMaI	19.94 ± 0.13	10	0.111 ± 0.07	-0.387 ± 0.058	-0.641 ± 0.068	0.004	-58.3 ± 0.4	3
UMaII	17.7 ± 0.12	5	-0.237 ± 0.141	1.701 ± 0.12	-1.845 ± 0.128	0.061	-115.6 ± 0.9	3
UMiI	19.4 ± 0.11	782	-0.019 ± 0.018	-0.114 ± 0.02	0.069 ± 0.02	-0.009	-245.1 ± 0.1	17,41
Will	18.27 ± 0.49	7	0.084 ± 0.123	0.295 ± 0.086	-1.074 ± 0.131	-0.098	-20.7 ± 0.6	3

NOTE—Column 1 lists the abbreviated dwarf galaxy name, with names in italics represents dwarf galaxies for which the number of representative stars is small; Column 2-6 gives the distance modulus, number of member stars, parallax, and proper motions in both dimension; Column 7 is the correlation coefficient between  $\mu_{\alpha^*}$  and  $\mu_{\delta}$ ; Column 8-9 is the heliocentric radial velocity and the references of member stars.

**References**—(1) Torrealba et al. (2019); (2) Torrealba et al. (2016); (3) Martin et al. (2007); (4) Koposov et al. (2011); (5) Koch et al. (2009); (6) Ural et al. (2010); (7) Simon & Geha (2007); (8) Walker et al. (2009b); (9) Muñoz et al. (2006); (10) Li et al. (2018); (11) Fritz et al. (2019); (12) Kirby et al. (2015); (13) Voggel et al. (2016); (14) Caldwell et al. (2017); (15) Kleyna et al. (2002); (16) Walker et al. (2015); (17) Armandroff et al. (1995); (18) Martin et al. (2016); (19) Li et al. (2017); (20) Battaglia et al. (2006); (21) Walker et al. (2016); (22) Simon et al. (2020); (23) Adén et al. (2009); (24) Koposov et al. (2015); (25) Koposov et al. (2018); (26) Mateo et al. (2008); (27) Sohn et al. (2007); (28) Spencer et al. (2017); (29) Koch et al. (2007); (30) Jenkins et al. (2020); (31) Walker et al. (2009a); (32) Kacharov et al. (2017); (33) Simon et al. (2015); (34) Hill et al. (2019); (35) Simon et al. (2011); (36) Kirby et al. (2013); (37) Battaglia et al. (2011); (38) Longeard et al. (2020); (39) Kirby et al. (2017); (40) Simon et al. (2017); (41) Li et al. (2018); (42) Pace et al. (2020).

<sup>a</sup>The reference of distance modulus is the same as Fritz et al. (2018) and updated with reference of Simon (2019), except the following galaxy: Antlia II (Torrealba et al. 2019).



### 2.3. Galactocentric coordinates and velocities

Heliocentric distance ( $d$ ) can be derived from the distance modulus ( $dm$ ):

$$d = 10^{dm/5-2} \text{ kpc.} \quad (2)$$

We then transform it and the Galactic coordinates ( $l, b$ ) for dwarf galaxies into a Galactocentric Cartesian coordinate system ( $x, y, z$ ):

$$\begin{aligned} x &= R_{\odot} - d \cos(b) \cos(l) \\ y &= -d \cos(b) \sin(l) \\ z &= d \sin(b) + z_{\odot}, \end{aligned} \quad (3)$$

where  $R_{\odot} = 8.122$  kpc is the distance from the Sun to Galactic center (Gravity Collaboration et al. 2018) and  $z_{\odot} = 25$  pc is the solar offset from the Galactic midplane (Jurić et al. 2008). We use a right-handed Galactocentric frame, the sign of  $x$  and  $z$  are positive in the directions of the Sun, and the North Galactic Pole (NGP), respectively. The Galactic space-velocity components ( $U, V, W$ ) can be calculated from proper motions, radial velocities and distances (Johnson & Soderblom 1987).  $U, V$  and  $W$  point toward the Galactic center, Galactic rotation, and the North Galactic Pole (NGP), respectively. We assume a solar motion  $(U_{\odot}, V_{\odot}, W_{\odot}) = (10., 11., 7.)$  km s<sup>-1</sup> (Bland-Hawthorn & Gerhard 2016) and a circular velocity at the location of the Sun  $V_{\text{LSR}} = 229$  km s<sup>-1</sup> (Eilers et al. 2019).

We then transform the Cartesian coordinate system ( $x, y, z, v_x, v_y, v_z$ ) to a spherical coordinate system ( $r_{\text{gc}}, \theta, \phi, v_r, v_{\theta}, v_{\phi}$ ), where  $(v_x, v_y, v_z)$  are related to the  $(U, V, W)$  by,  $v_x = -U, v_y = -V$ , and  $v_z = W$ . We generate 2,000 realizations for each galaxy using a Monte Carlo (MC) method and calculate their 3D position  $(r, \theta, \phi)$  and 3D velocity  $(v_r, v_{\theta}, v_{\phi})$  in the spherical coordinate system. The result is listed in Table 2.

The Galactocentric tangential velocity is given by:


$$v_{\text{tan}} = \sqrt{v_{\theta}^2 + v_{\phi}^2}. \quad (4)$$

The value of  $v_{\text{tan,obs}}$  is a biased estimator of the true tangential velocity  $v_{\text{tan}}$  (van der Marel & Guhathakurta 2008), especially when the relative error is close to or greater than 1 (Fritz et al. 2018). In order to correct this bias and derive accurate estimates, van der Marel & Guhathakurta (2008) use a Bayesian approach to infer tangential velocity. The posterior PDF is:

$$P(v_{\text{tan}}|v_{\text{tan,obs}}) \propto P(v_{\text{tan,obs}}|v_{\text{tan}})P(v_{\text{tan}}). \quad (5)$$

We assume uniform priors on  $v_{\text{tan}}$ . The likelihood distribution is obtained by sampling the observational data using a MC method. We characterize the posterior via rejection sampling and obtain 2,000 samples for each galaxy.

In the Bayesian approach, the effect of proper motions and other observed data is included in the likelihood function. For the 46 dwarf galaxies this does not affect the value of  $v_{\text{tan,obs}}$ , but only the error bars for the few objects with very large uncertainties. We adopt the Bayesian calculations in Table 2 for error bars of tangential velocity  $v_{\text{tan}}$  and of total velocity  $v_{3\text{D}}$ . Figure 1 shows the Galactocentric total velocity  $v_{3\text{D}}$  as a function of Galactocentric distance  $r_{\text{gc}}$  for all galaxies.



**Figure 1.** 3D phase diagram for dwarf galaxies, which are represented by open and full triangles for visible luminosity smaller or larger than  $10^5 L_{\odot}$ , respectively. The triangle orientation towards  $r_{gc} = 0$  or in the opposite direction indicates whether dwarf galaxies are on an approaching ( $v_r < 0$ ) or on a receding ( $v_r > 0$ ) orbits, respectively. Green, cyan, magenta, and blue lines indicate the escape velocity of Einasto high-mass (PE<sub>HM</sub>), PNFW, intermediate mass (PE<sub>IM</sub>), and low mass (PE<sub>LM</sub>), respectively. Star-points indicates Sagittarius (red) and the LMC (green) positions.

**Table 2.** Kinematic properties of dwarf galaxies in Galactocentric spherical coordinate system.

name	$r_{GC}$	$\theta$	$\phi$	$v_r$	$v_{\theta}$	$v_{\phi}$	$v_{tan}$	$v_{3D}$
	(kpc)	(deg)	(deg)	(km s <sup>-1</sup> )	(km s <sup>-1</sup> )	(km s <sup>-1</sup> )	(km s <sup>-1</sup> )	(km s <sup>-1</sup> )
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
AntII	132.8 <sup>+6.7</sup> <sub>-6.3</sub>	78.8 <sup>+0.0</sup> <sub>-0.0</sub>	81.3 <sup>+0.2</sup> <sub>-0.2</sub>	66.4 <sup>+0.8</sup> <sub>-0.8</sub>	-61.2 <sup>+12.7</sup> <sub>-12.5</sub>	-102.6 <sup>+12.8</sup> <sub>-13.3</sub>	119.7 <sup>+12.9</sup> <sub>-13.1</sub>	136.9 <sup>+11.7</sup> <sub>-11.7</sub>
AquII	105.4 <sup>+3.2</sup> <sub>-3.3</sub>	144.9 <sup>+0.1</sup> <sub>-0.0</sub>	241.4 <sup>+0.2</sup> <sub>-0.2</sub>	59.3 <sup>+24.0</sup> <sub>-23.5</sub>	182.4 <sup>+241.9</sup> <sub>-257.2</sub>	191.3 <sup>+323.5</sup> <sub>-326.1</sub>	300.3 <sup>+288.5</sup> <sub>-205.7</sub>	305.2 <sup>+288.6</sup> <sub>-201.0</sub>
BooI	63.7 <sup>+2.0</sup> <sub>-1.9</sub>	13.5 <sup>+0.2</sup> <sub>-0.2</sub>	177.0 <sup>+0.0</sup> <sub>-0.0</sub>	87.4 <sup>+1.9</sup> <sub>-1.9</sub>	178.4 <sup>+16.1</sup> <sub>-16.4</sub>	-92.6 <sup>+17.0</sup> <sub>-17.1</sub>	201.7 <sup>+16.7</sup> <sub>-17.6</sub>	219.7 <sup>+15.0</sup> <sub>-15.2</sub>
BooII	39.8 <sup>+1.0</sup> <sub>-1.0</sub>	10.3 <sup>+0.3</sup> <sub>-0.3</sub>	166.6 <sup>+0.4</sup> <sub>-0.4</sub>	-54.2 <sup>+5.7</sup> <sub>-5.8</sub>	-283.9 <sup>+27.1</sup> <sub>-29.1</sub>	-169.4 <sup>+26.1</sup> <sub>-26.4</sub>	331.7 <sup>+31.4</sup> <sub>-32.0</sub>	336.1 <sup>+30.5</sup> <sub>-30.4</sub>

Table 2 continued on next page



Table 2 (continued)

name	$r_{GC}$	$\theta$	$\phi$	$v_r$	$v_\theta$	$v_\phi$	$v_{tan}$	$v_{3D}$
	(kpc)	(deg)	(deg)	(km s <sup>-1</sup> )	(km s <sup>-1</sup> )	(km s <sup>-1</sup> )	(km s <sup>-1</sup> )	(km s <sup>-1</sup> )
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
CVenI	210.7 <sup>+6.3</sup> <sub>-6.1</sub>	9.8 <sup>+0.0</sup> <sub>-0.0</sub>	266.9 <sup>+0.4</sup> <sub>-0.4</sub>	73.8 <sup>+1.6</sup> <sub>-1.6</sub>	85.7 <sup>+46.6</sup> <sub>-45.5</sub>	-18.2 <sup>+40.5</sup> <sub>-40.9</sub>	91.2 <sup>+49.2</sup> <sub>-47.5</sub>	117.6 <sup>+40.7</sup> <sub>-31.2</sub>
CVenII	160.7 <sup>+4.4</sup> <sub>-4.0</sub>	8.9 <sup>+0.0</sup> <sub>-0.1</sub>	311.1 <sup>+0.4</sup> <sub>-0.4</sub>	-97.8 <sup>+3.4</sup> <sub>-3.6</sub>	-51.4 <sup>+57.9</sup> <sub>-56.1</sub>	-26.1 <sup>+88.7</sup> <sub>-82.6</sub>	71.3 <sup>+64.5</sup> <sub>-46.7</sub>	121.0 <sup>+48.1</sup> <sub>-22.3</sub>
CarI	107.6 <sup>+4.9</sup> <sub>-4.9</sub>	111.9 <sup>+0.0</sup> <sub>-0.0</sub>	75.5 <sup>+0.2</sup> <sub>-0.2</sub>	0.8 <sup>+0.8</sup> <sub>-0.7</sub>	-191.6 <sup>+17.5</sup> <sub>-16.4</sub>	-8.6 <sup>+9.8</sup> <sub>-10.2</sub>	192.2 <sup>+17.4</sup> <sub>-15.9</sub>	192.2 <sup>+17.4</sup> <sub>-15.9</sub>
CarII	37.1 <sup>+0.6</sup> <sub>-0.6</sub>	106.7 <sup>+0.0</sup> <sub>-0.0</sub>	76.8 <sup>+0.2</sup> <sub>-0.2</sub>	217.6 <sup>+1.6</sup> <sub>-1.7</sub>	-231.4 <sup>+9.7</sup> <sub>-9.5</sub>	183.5 <sup>+7.4</sup> <sub>-6.7</sub>	295.4 <sup>+8.8</sup> <sub>-8.6</sub>	366.8 <sup>+6.8</sup> <sub>-6.5</sub>
CarIII	29.0 <sup>+0.6</sup> <sub>-0.6</sub>	106.1 <sup>+0.0</sup> <sub>-0.0</sub>	73.0 <sup>+0.3</sup> <sub>-0.3</sub>	46.6 <sup>+2.6</sup> <sub>-2.7</sub>	-379.9 <sup>+13.8</sup> <sub>-13.4</sub>	57.1 <sup>+9.1</sup> <sub>-8.4</sub>	384.0 <sup>+13.5</sup> <sub>-12.8</sub>	386.9 <sup>+13.3</sup> <sub>-12.7</sub>
ColI	187.1 <sup>+10.5</sup> <sub>-9.3</sub>	118.1 <sup>+0.0</sup> <sub>-0.0</sub>	49.3 <sup>+0.1</sup> <sub>-0.1</sub>	-34.4 <sup>+5.6</sup> <sub>-5.9</sub>	84.0 <sup>+97.8</sup> <sub>-105.5</sub>	360.6 <sup>+124.1</sup> <sub>-122.4</sub>	372.9 <sup>+122.7</sup> <sub>-113.3</sub>	374.4 <sup>+122.8</sup> <sub>-113.4</sub>
CberI	43.3 <sup>+1.5</sup> <sub>-1.6</sub>	14.9 <sup>+0.4</sup> <sub>-0.3</sub>	21.8 <sup>+0.6</sup> <sub>-0.6</sub>	32.5 <sup>+2.6</sup> <sub>-2.6</sub>	-230.6 <sup>+12.3</sup> <sub>-12.8</sub>	111.9 <sup>+18.7</sup> <sub>-17.0</sub>	256.9 <sup>+16.0</sup> <sub>-15.2</sub>	258.9 <sup>+15.7</sup> <sub>-14.7</sub>
CraI	145.9 <sup>+1.9</sup> <sub>-2.1</sub>	40.5 <sup>+0.0</sup> <sub>-0.0</sub>	88.8 <sup>+0.1</sup> <sub>-0.1</sub>	-5.8 <sup>+5.4</sup> <sub>-5.2</sub>	-134.5 <sup>+69.7</sup> <sub>-66.2</sub>	88.0 <sup>+92.0</sup> <sub>-95.5</sub>	163.8 <sup>+79.1</sup> <sub>-70.4</sub>	163.9 <sup>+79.1</sup> <sub>-70.5</sub>
CraII	116.5 <sup>+1.1</sup> <sub>-1.1</sub>	47.5 <sup>+0.0</sup> <sub>-0.0</sub>	97.6 <sup>+0.0</sup> <sub>-0.0</sub>	-81.7 <sup>+1.3</sup> <sub>-1.1</sub>	-85.7 <sup>+15.0</sup> <sub>-14.6</sub>	37.5 <sup>+15.7</sup> <sub>-18.4</sub>	93.6 <sup>+15.8</sup> <sub>-16.5</sub>	124.2 <sup>+12.6</sup> <sub>-12.1</sub>
DraI	81.8 <sup>+6.1</sup> <sub>-5.7</sub>	55.3 <sup>+0.0</sup> <sub>-0.0</sub>	273.3 <sup>+0.5</sup> <sub>-0.5</sub>	-97.4 <sup>+0.8</sup> <sub>-0.8</sub>	136.9 <sup>+7.9</sup> <sub>-7.4</sub>	-66.2 <sup>+9.5</sup> <sub>-9.7</sub>	153.1 <sup>+8.3</sup> <sub>-8.6</sub>	181.5 <sup>+7.2</sup> <sub>-7.5</sub>
DraII	23.8 <sup>+0.4</sup> <sub>-0.4</sub>	52.0 <sup>+0.1</sup> <sub>-0.1</sub>	303.7 <sup>+0.4</sup> <sub>-0.4</sub>	-153.9 <sup>+4.5</sup> <sub>-4.7</sub>	299.5 <sup>+11.7</sup> <sub>-11.7</sub>	-43.5 <sup>+11.7</sup> <sub>-12.2</sub>	302.0 <sup>+11.7</sup> <sub>-11.3</sub>	338.9 <sup>+10.2</sup> <sub>-9.6</sub>
EriII	367.9 <sup>+17.4</sup> <sub>-16.2</sub>	141.3 <sup>+0.0</sup> <sub>-0.0</sub>	67.9 <sup>+0.1</sup> <sub>-0.1</sub>	-76.7 <sup>+5.7</sup> <sub>-5.4</sub>	-21.2 <sup>+225.2</sup> <sub>-249.0</sub>	193.2 <sup>+280.3</sup> <sub>-264.6</sub>	267.8 <sup>+269.7</sup> <sub>-184.1</sub>	278.8 <sup>+265.2</sup> <sub>-167.7</sub>
FnxI	141.0 <sup>+3.1</sup> <sub>-3.1</sub>	153.9 <sup>+0.0</sup> <sub>-0.0</sub>	50.9 <sup>+0.1</sup> <sub>-0.1</sub>	-36.8 <sup>+0.7</sup> <sub>-0.7</sub>	-100.0 <sup>+14.6</sup> <sub>-12.9</sub>	78.5 <sup>+12.8</sup> <sub>-13.2</sub>	126.9 <sup>+14.7</sup> <sub>-14.2</sub>	132.2 <sup>+14.1</sup> <sub>-13.7</sub>
GruI	116.0 <sup>+11.9</sup> <sub>-10.7</sub>	151.4 <sup>+0.3</sup> <sub>-0.3</sub>	155.6 <sup>+0.3</sup> <sub>-0.3</sub>	-187.9 <sup>+2.1</sup> <sub>-2.1</sub>	21.2 <sup>+37.0</sup> <sub>-42.1</sub>	73.0 <sup>+41.0</sup> <sub>-41.6</sub>	76.9 <sup>+41.6</sup> <sub>-40.2</sub>	203.0 <sup>+19.1</sup> <sub>-11.7</sub>
GruII	48.6 <sup>+5.4</sup> <sub>-4.6</sub>	149.4 <sup>+0.8</sup> <sub>-0.7</sub>	168.2 <sup>+0.3</sup> <sub>-0.4</sub>	-125.9 <sup>+1.2</sup> <sub>-1.2</sub>	17.5 <sup>+9.4</sup> <sub>-9.3</sub>	-162.9 <sup>+36.8</sup> <sub>-40.0</sub>	166.1 <sup>+40.7</sup> <sub>-35.1</sub>	208.3 <sup>+33.8</sup> <sub>-26.5</sub>
HerI	126.5 <sup>+6.4</sup> <sub>-6.2</sub>	51.2 <sup>+0.1</sup> <sub>-0.1</sub>	211.0 <sup>+0.1</sup> <sub>-0.1</sub>	143.8 <sup>+1.2</sup> <sub>-1.3</sub>	130.9 <sup>+30.4</sup> <sub>-31.6</sub>	-4.2 <sup>+38.7</sup> <sub>-38.9</sub>	131.7 <sup>+31.1</sup> <sub>-28.4</sub>	195.0 <sup>+21.8</sup> <sub>-17.6</sub>
HorI	86.6 <sup>+13.6</sup> <sub>-10.9</sub>	144.5 <sup>+0.1</sup> <sub>-0.1</sub>	82.1 <sup>+1.3</sup> <sub>-1.3</sub>	-27.5 <sup>+1.8</sup> <sub>-1.9</sub>	-238.2 <sup>+57.2</sup> <sub>-69.2</sub>	24.2 <sup>+21.4</sup> <sub>-18.2</sub>	249.9 <sup>+65.7</sup> <sub>-57.7</sub>	251.3 <sup>+65.5</sup> <sub>-57.0</sub>
HorII	79.2 <sup>+7.6</sup> <sub>-7.0</sub>	143.1 <sup>+0.1</sup> <sub>-0.1</sub>	72.7 <sup>+0.8</sup> <sub>-0.9</sub>	2.1 <sup>+19.0</sup> <sub>-19.1</sub>	-236.6 <sup>+134.1</sup> <sub>-148.5</sub>	163.9 <sup>+157.1</sup> <sub>-157.9</sub>	310.7 <sup>+167.9</sup> <sub>-153.5</sub>	310.8 <sup>+168.1</sup> <sub>-152.9</sub>
HyaII	147.8 <sup>+7.8</sup> <sub>-7.1</sub>	58.9 <sup>+0.0</sup> <sub>-0.0</sub>	112.3 <sup>+0.2</sup> <sub>-0.2</sub>	138.7 <sup>+9.6</sup> <sub>-9.6</sub>	11.6 <sup>+164.4</sup> <sub>-154.0</sub>	-286.1 <sup>+195.3</sup> <sub>-194.2</sub>	299.2 <sup>+190.6</sup> <sub>-178.0</sub>	329.7 <sup>+181.1</sup> <sub>-151.7</sub>
HyII	25.7 <sup>+0.5</sup> <sub>-0.5</sub>	129.9 <sup>+0.0</sup> <sub>-0.0</sub>	96.0 <sup>+0.4</sup> <sub>-0.4</sub>	-47.6 <sup>+1.1</sup> <sub>-1.0</sub>	-327.9 <sup>+8.6</sup> <sub>-8.8</sub>	-176.1 <sup>+5.8</sup> <sub>-7.1</sub>	372.8 <sup>+9.8</sup> <sub>-9.9</sub>	375.8 <sup>+9.7</sup> <sub>-9.7</sub>
LeoI	257.9 <sup>+17.0</sup> <sub>-14.6</sub>	41.8 <sup>+0.1</sup> <sub>-0.1</sub>	44.0 <sup>+0.1</sup> <sub>-0.1</sub>	172.4 <sup>+1.0</sup> <sub>-1.0</sub>	-11.8 <sup>+32.9</sup> <sub>-32.2</sub>	-79.2 <sup>+36.2</sup> <sub>-31.8</sub>	80.5 <sup>+31.8</sup> <sub>-32.7</sub>	190.2 <sup>+15.9</sup> <sub>-11.9</sub>
LeoII	235.6 <sup>+13.9</sup> <sub>-14.1</sub>	24.2 <sup>+0.1</sup> <sub>-0.1</sub>	37.1 <sup>+0.2</sup> <sub>-0.2</sub>	25.7 <sup>+1.4</sup> <sub>-1.5</sub>	23.7 <sup>+43.1</sup> <sub>-39.8</sub>	-91.1 <sup>+44.0</sup> <sub>-42.2</sub>	94.5 <sup>+43.2</sup> <sub>-43.1</sub>	97.9 <sup>+42.5</sup> <sub>-41.2</sub>
LeoIV	154.4 <sup>+5.1</sup> <sub>-4.8</sub>	33.8 <sup>+0.0</sup> <sub>-0.0</sub>	80.0 <sup>+0.2</sup> <sub>-0.2</sub>	-0.6 <sup>+6.8</sup> <sub>-6.5</sub>	-55.0 <sup>+102.9</sup> <sub>-102.5</sub>	94.8 <sup>+128.6</sup> <sub>-131.0</sub>	131.2 <sup>+128.6</sup> <sub>-90.4</sub>	131.2 <sup>+128.6</sup> <sub>-90.3</sub>
LeoV	169.8 <sup>+3.9</sup> <sub>-3.8</sub>	31.9 <sup>+0.0</sup> <sub>-0.0</sub>	76.7 <sup>+0.1</sup> <sub>-0.1</sub>	44.1 <sup>+7.6</sup> <sub>-8.0</sub>	-27.9 <sup>+131.7</sup> <sub>-127.3</sub>	233.2 <sup>+158.7</sup> <sub>-154.5</sub>	261.5 <sup>+156.7</sup> <sub>-154.0</sub>	265.1 <sup>+154.6</sup> <sub>-145.9</sub>
PhxII	81.4 <sup>+4.3</sup> <sub>-3.7</sub>	154.3 <sup>+0.2</sup> <sub>-0.2</sub>	136.9 <sup>+0.4</sup> <sub>-0.4</sub>	-37.6 <sup>+2.5</sup> <sub>-2.5</sub>	-202.2 <sup>+30.3</sup> <sub>-29.3</sub>	-210.0 <sup>+36.0</sup> <sub>-36.8</sub>	291.7 <sup>+42.8</sup> <sub>-41.0</sub>	294.2 <sup>+42.5</sup> <sub>-40.7</sub>
PisII	181.9 <sup>+16.5</sup> <sub>-15.0</sub>	137.4 <sup>+0.0</sup> <sub>-0.0</sub>	262.9 <sup>+0.3</sup> <sub>-0.3</sub>	-60.0 <sup>+10.6</sup> <sub>-10.6</sub>	606.5 <sup>+226.6</sup> <sub>-210.0</sub>	188.9 <sup>+241.0</sup> <sub>-224.9</sub>	641.0 <sup>+257.6</sup> <sub>-234.3</sub>	644.0 <sup>+256.2</sup> <sub>-231.8</sub>
RetII	32.9 <sup>+1.4</sup> <sub>-1.4</sub>	137.0 <sup>+0.2</sup> <sub>-0.2</sub>	65.1 <sup>+0.9</sup> <sub>-0.9</sub>	-98.7 <sup>+1.1</sup> <sub>-1.2</sub>	-224.2 <sup>+18.4</sup> <sub>-19.7</sub>	68.2 <sup>+7.1</sup> <sub>-6.8</sub>	235.9 <sup>+20.4</sup> <sub>-19.0</sub>	255.7 <sup>+18.9</sup> <sub>-17.4</sub>
RetIII	91.6 <sup>+13.2</sup> <sub>-11.7</sub>	135.7 <sup>+0.0</sup> <sub>-0.0</sub>	86.6 <sup>+0.9</sup> <sub>-1.1</sub>	102.7 <sup>+10.7</sup> <sub>-10.9</sub>	-69.6 <sup>+99.4</sup> <sub>-103.9</sub>	2.4 <sup>+113.7</sup> <sub>-105.7</sub>	99.0 <sup>+104.4</sup> <sub>-69.4</sub>	143.3 <sup>+84.2</sup> <sub>-35.5</sub>
SgrII	63.1 <sup>+2.3</sup> <sub>-2.3</sub>	115.9 <sup>+0.1</sup> <sub>-0.1</sub>	200.6 <sup>+0.1</sup> <sub>-0.1</sub>	-113.4 <sup>+1.7</sup> <sub>-1.6</sub>	-143.5 <sup>+24.0</sup> <sub>-25.6</sub>	-138.1 <sup>+22.1</sup> <sub>-21.7</sub>	200.8 <sup>+27.2</sup> <sub>-27.2</sub>	230.7 <sup>+24.7</sup> <sub>-24.1</sub>
SclII	86.0 <sup>+5.2</sup> <sub>-4.7</sub>	172.7 <sup>+0.1</sup> <sub>-0.1</sub>	62.7 <sup>+2.2</sup> <sub>-2.2</sub>	75.9 <sup>+0.7</sup> <sub>-0.7</sub>	154.2 <sup>+8.2</sup> <sub>-8.4</sub>	-54.3 <sup>+11.4</sup> <sub>-10.9</sub>	162.7 <sup>+9.1</sup> <sub>-9.2</sub>	179.6 <sup>+8.3</sup> <sub>-8.2</sub>
SegI	27.9 <sup>+2.0</sup> <sub>-1.8</sub>	50.4 <sup>+0.8</sup> <sub>-0.8</sub>	26.3 <sup>+0.8</sup> <sub>-0.8</sub>	130.4 <sup>+1.6</sup> <sub>-1.7</sub>	182.9 <sup>+30.5</sup> <sub>-28.6</sub>	133.2 <sup>+24.7</sup> <sub>-24.0</sub>	229.0 <sup>+38.6</sup> <sub>-37.6</sub>	263.5 <sup>+34.4</sup> <sub>-31.8</sub>
SegII	42.8 <sup>+3.0</sup> <sub>-2.9</sub>	122.1 <sup>+0.4</sup> <sub>-0.4</sub>	336.0 <sup>+0.4</sup> <sub>-0.4</sub>	60.4 <sup>+2.1</sup> <sub>-2.1</sub>	-128.1 <sup>+11.8</sup> <sub>-12.0</sub>	23.8 <sup>+24.5</sup> <sub>-21.1</sub>	132.2 <sup>+17.8</sup> <sub>-13.1</sub>	145.5 <sup>+17.0</sup> <sub>-12.7</sub>
SxtI	98.1 <sup>+3.0</sup> <sub>-2.9</sub>	49.3 <sup>+0.1</sup> <sub>-0.0</sub>	57.9 <sup>+0.2</sup> <sub>-0.2</sub>	83.3 <sup>+0.8</sup> <sub>-0.8</sub>	-8.8 <sup>+10.4</sup> <sub>-10.4</sub>	-219.7 <sup>+9.9</sup> <sub>-11.0</sub>	220.7 <sup>+11.3</sup> <sub>-10.4</sub>	235.9 <sup>+10.8</sup> <sub>-10.0</sub>
TriII	34.7 <sup>+1.6</sup> <sub>-1.5</sub>	109.3 <sup>+0.2</sup> <sub>-0.2</sub>	329.9 <sup>+0.4</sup> <sub>-0.4</sub>	-259.4 <sup>+2.2</sup> <sub>-2.3</sub>	-117.1 <sup>+12.2</sup> <sub>-12.8</sub>	-89.0 <sup>+10.3</sup> <sub>-10.0</sub>	147.4 <sup>+12.1</sup> <sub>-12.0</sub>	298.5 <sup>+5.9</sup> <sub>-5.9</sub>
TucII	54.0 <sup>+8.8</sup> <sub>-7.4</sub>	147.9 <sup>+0.8</sup> <sub>-0.7</sub>	139.4 <sup>+1.4</sup> <sub>-1.6</sub>	-182.9 <sup>+1.1</sup> <sub>-1.0</sub>	31.9 <sup>+13.8</sup> <sub>-16.1</sub>	-235.6 <sup>+51.3</sup> <sub>-61.0</sub>	247.6 <sup>+63.7</sup> <sub>-47.0</sub>	307.5 <sup>+53.2</sup> <sub>-35.6</sub>
TucIII	23.0 <sup>+2.0</sup> <sub>-1.7</sub>	154.4 <sup>+0.2</sup> <sub>-0.3</sub>	100.2 <sup>+3.6</sup> <sub>-3.7</sub>	-229.2 <sup>+1.0</sup> <sub>-1.0</sub>	25.7 <sup>+17.8</sup> <sub>-21.1</sub>	57.2 <sup>+9.9</sup> <sub>-11.5</sub>	64.3 <sup>+7.7</sup> <sub>-3.8</sub>	238.2 <sup>+2.8</sup> <sub>-1.8</sub>

Table 2 continued on next page

**Table 2** (*continued*)

name	$r_{\text{GC}}$	$\theta$	$\phi$	$v_r$	$v_\theta$	$v_\phi$	$v_{\text{tan}}$	$v_{3\text{D}}$
	(kpc)	(deg)	(deg)	(km s <sup>-1</sup> )	(km s <sup>-1</sup> )	(km s <sup>-1</sup> )	(km s <sup>-1</sup> )	(km s <sup>-1</sup> )
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
TucIV	45.6 <sup>+4.1</sup> <sub>-4.0</sub>	150.2 <sup>+0.4</sup> <sub>-0.3</sub>	118.2 <sup>+1.4</sup> <sub>-1.7</sub>	-93.5 <sup>+1.4</sup> <sub>-1.5</sub>	-194.3 <sup>+31.4</sup> <sub>-32.5</sub>	-58.4 <sup>+19.4</sup> <sub>-21.4</sub>	207.2 <sup>+35.3</sup> <sub>-33.7</sub>	227.3 <sup>+32.4</sup> <sub>-30.0</sub>
TucV	52.2 <sup>+9.0</sup> <sub>-7.9</sub>	146.5 <sup>+0.7</sup> <sub>-0.6</sub>	125.1 <sup>+1.8</sup> <sub>-2.3</sub>	-166.8 <sup>+3.3</sup> <sub>-3.3</sub>	-175.9 <sup>+60.4</sup> <sub>-67.5</sub>	105.6 <sup>+22.6</sup> <sub>-25.7</sub>	221.0 <sup>+62.5</sup> <sub>-53.3</sub>	276.9 <sup>+51.7</sup> <sub>-40.5</sub>
UMaI	102.0 <sup>+5.9</sup> <sub>-5.5</sub>	39.1 <sup>+0.2</sup> <sub>-0.2</sub>	342.0 <sup>+0.1</sup> <sub>-0.1</sub>	-2.2 <sup>+1.8</sup> <sub>-1.9</sub>	61.9 <sup>+28.3</sup> <sub>-26.7</sub>	114.7 <sup>+37.9</sup> <sub>-36.0</sub>	133.3 <sup>+34.9</sup> <sub>-32.1</sub>	133.3 <sup>+34.9</sup> <sub>-32.1</sub>
UMaII	40.8 <sup>+2.0</sup> <sub>-1.9</sub>	58.9 <sup>+0.3</sup> <sub>-0.3</sub>	338.6 <sup>+0.3</sup> <sub>-0.3</sub>	-59.7 <sup>+3.1</sup> <sub>-3.2</sub>	-280.5 <sup>+26.5</sup> <sub>-28.2</sub>	30.7 <sup>+26.3</sup> <sub>-25.4</sub>	285.9 <sup>+29.3</sup> <sub>-26.7</sub>	292.1 <sup>+28.9</sup> <sub>-26.3</sub>
UMiI	77.8 <sup>+4.2</sup> <sub>-3.9</sub>	46.5 <sup>+0.1</sup> <sub>-0.1</sub>	293.0 <sup>+0.4</sup> <sub>-0.4</sub>	-76.9 <sup>+0.8</sup> <sub>-0.8</sub>	143.9 <sup>+7.3</sup> <sub>-6.7</sub>	-31.9 <sup>+8.1</sup> <sub>-8.0</sub>	147.6 <sup>+7.1</sup> <sub>-7.4</sub>	166.4 <sup>+6.4</sup> <sub>-6.5</sub>
WiII	49.3 <sup>+11.6</sup> <sub>-8.8</sub>	40.7 <sup>+1.7</sup> <sub>-1.4</sub>	343.9 <sup>+0.9</sup> <sub>-0.9</sub>	8.1 <sup>+2.9</sup> <sub>-2.8</sub>	-124.7 <sup>+36.9</sup> <sub>-49.7</sub>	-67.8 <sup>+49.3</sup> <sub>-39.3</sub>	159.2 <sup>+53.7</sup> <sub>-29.3</sub>	159.4 <sup>+53.7</sup> <sub>-29.2</sub>

NOTE—Column 1 lists the abbreviated dwarf galaxy name; Column 2-4 is the Galactocentric distance, angle with respect to the North Galactic Pole and azimuthal angle; Column 5-7 gives the velocities in three dimensions; Column 8 provides the Galactocentric tangential velocity and Column 9 lists the total velocity in the Galactic rest frame.

### 3. INTEGRATED ORBITAL PARAMETERS

#### 3.1. *Four flavors of the Milky Way potential*

Based on APOGEE, WISE, 2MASS and Gaia data, [Eilers et al. \(2019\)](#) derived the rotation curve of the Milky Way at Galactocentric distances between  $5 \leq R \leq 25$  kpc. Combined with the baryonic components of [Pouliasis et al. \(2017, their Model I\)](#), they estimate the parameters of a NFW ([Navarro et al. 1997](#)) dark matter halo by fitting the rotation curve. Model I of [Pouliasis et al. \(2017\)](#) includes three baryonic components: a Plummer ([Binney & Tremaine 1987](#)) bulge, and two Miyamoto-Nagai ([Miyamoto & Nagai 1975](#)) for the thin and thick disk. We choose this model (*PNFW*, MW total mass:  $8.1 \times 10^{11} M_\odot$ ) that is similar to that of [Bovy \(2015, see a comparison of the two mass profiles in Figure 1 of Hammer et al. \(2020\)\)](#).

[Jiao et al. \(2021\)](#) determined the range of Milky Way mass that can reproduce the rotation curve of [Eilers et al. \(2019\)](#). They found that an Einasto profile ([Retana-Montenegro et al. 2012](#)) for the dark matter is more appropriate to define this range, which includes MW total masses from  $2.8 \times 10^{11} M_\odot$  to  $15 \times 10^{11} M_\odot$ . We adopt these two models ( $\text{PE}_{\text{LM}}$  and  $\text{PE}_{\text{HM}}$ , also associated to Model I of [Pouliasis et al. 2017](#)).  $\text{PE}_{\text{LM}}$  provides the best fit of the MW rotation curve, while  $\text{PE}_{\text{HM}}$  is among the most massive MW mass available for such a fitting. Both *PNFW* and  $\text{PE}_{\text{HM}}$  are comparable to the 'low' and 'high' MW mass models used by [Fritz et al. \(2018\)](#), respectively. For a better sampling of possible MW masses that fit the rotation curve, we also consider an intermediate mass model ( $\text{PE}_{\text{IM}}$ ) from [Wang & Hammer \(2021\)](#), obtained after combining constraints from the MW rotation curve and globular cluster orbital motions. Properties of the dark matter halo of the four models are listed in Table 3, all of them being associated to Model I of [Pouliasis et al. \(2017\)](#) for baryons, with a total baryonic mass of  $0.89 \times 10^{11} M_\odot$ .

#### 3.2. *Orbit integration*

We have used `galpy` ([Bovy 2015](#)) to investigate the orbital properties of the dwarf galaxies by adopting a Milky Way potential model. We use the sample derived from the Bayesian method, because the orbital properties are related to total energy, which is also affected by the bias in  $v_{\text{tan}}$ . For example, for the *PNFW* model, we have integrated the orbit from  $-10$  to  $10$  Gyr for each dwarf

**Table 3.** Properties of the four Milky Way dark-matter mass models.

Parameters	PE <sub>HM</sub>	PNFW	PE <sub>IM</sub>	PE <sub>LM</sub>
$M_{\text{DM}}(10^{11}M_{\odot})$	14.1	7.2	4.2	1.9
$M_{\text{tot}}(10^{11}M_{\odot})$	15	8.1	5.1	2.8
$r_{200}(\text{kpc})$	236	189	164	135

galaxy. Thus, we can derive the pericenter,  $r_{\text{peri}}$ , the closest approach of an orbit to the the Milky Way center (i.e., the Galaxy center, GC) for each orbit. However, due to long orbital periods or hyperbolic orbits, it is not always possible to derive the apocenter,  $r_{\text{apo}}$ , the farthest extent of an orbit from the GC. In case the chosen integration time is too short to fully cover one orbit, we at least may ensure that each dwarf galaxy can reach 300 kpc from the Milky Way center so that the Milky Way can be considered as a point source.

By assuming conservation of energy and angular momentum,  $r_{\text{apo}}$  can be derived from the following equation<sup>2</sup>:

$$\begin{aligned}
 E &= \Phi(\vec{r}_{\text{apo}}) + \frac{v_{\text{apo}}^2}{2} \\
 &= -\frac{GM}{r_{\text{apo}}} + \frac{L_{\text{apo}}^2}{2r_{\text{apo}}^2} \\
 &= -\frac{GM}{r_{\text{apo}}} + \frac{L_{10\text{Gyr}}^2}{2r_{\text{apo}}^2},
 \end{aligned} \tag{6}$$

where  $E$  is the total energy,  $M$  is the total mass of the MW, and  $L_{10\text{Gyr}}$  is the dwarf galaxy's angular momentum at 10 Gyr.

We define the eccentricity of an elliptical orbit by:

$$e = \frac{r_{\text{apo}} - r_{\text{peri}}}{r_{\text{apo}} + r_{\text{peri}}}. \tag{7}$$

Hyperbolic orbits in extended mass profile require another definition of the eccentricity. This is why hyperbolic orbits are characterized in Tables 4 and 5 by  $e > 1$  for purely hyperbolic orbits (hence without values for apocenter) and  $e \geq 1$  for orbits being hyperbolic on average but having few MC solutions consistent with elliptical orbits (and then with a quoted value for apocenter).

We notice that pericenters are well determined and do not depend much on the adopted MW mass profile (compare Tables 4 and 5), while the opposite is true for eccentricity and apocenter values. Such a property will be further investigated in a future paper (Hammer et al. 2021). We also notice that the more precise Gaia-EDR3 values have led to only two dwarf galaxies (Triangulum II and Tucana III) having pericenters lower than 20 kpc. This contrasts with Fritz et al. (2018) who were finding between seven to eight such low-pericenter dwarf galaxies, and this may affect some analyses based on the tidal disruption scenario caused by, e.g., the MW disk.

<sup>2</sup>  $r_{\text{apo}}$  is the larger one of the two roots.

**Table 4.** Orbital properties of dwarf galaxies for Model PE<sub>HM</sub> and Model PNF<sub>W</sub>.

name	Model PE <sub>HM</sub>					Model PNF <sub>W</sub>				
	$r_{\text{peri}}$ (kpc)	$r_{\text{apo}}^{a,c}$ (kpc)	$e$	$P_{\text{unb}}$	$P^{b,c}$	$r_{\text{peri}}$ (kpc)	$r_{\text{apo}}^{a,c}$ (kpc)	$e$	$P_{\text{unb}}$	$P^{b,c}$
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)
AntII	56 <sup>+12</sup> <sub>-10</sub>	149 <sup>+8</sup> <sub>-7</sub>	0.45 <sup>+0.06</sup> <sub>-0.06</sub>	0.0%	0.63 <sup>+0.03</sup> <sub>-0.04</sub>	69 <sup>+15</sup> <sub>-13</sub>	168 <sup>+19</sup> <sub>-12</sub>	0.41 <sup>+0.06</sup> <sub>-0.04</sub>	0.0%	0.48 <sup>+0.06</sup> <sub>-0.08</sub>
AquII	101 <sup>+4</sup> <sub>-71</sub>	116 <sup>+190</sup> <sub>-8</sub>	0.95 <sup>+6.15</sup> <sub>-0.61</sub>	47.3%	0.07 <sup>+0.72</sup> <sub>-0.02</sub>	102 <sup>+4</sup> <sub>-66</sub>	116 <sup>+163</sup> <sub>-7</sub>	≥ 1	58.9%	0.07 <sup>+0.65</sup> <sub>-0.02</sub>
BooI	49 <sup>+5</sup> <sub>-6</sub>	97 <sup>+15</sup> <sub>-10</sub>	0.33 <sup>+0.02</sup> <sub>-0.01</sub>	0.0%	0.28 <sup>+0.08</sup> <sub>-0.07</sub>	52 <sup>+4</sup> <sub>-5</sub>	135 <sup>+39</sup> <sub>-25</sub>	0.44 <sup>+0.07</sup> <sub>-0.04</sub>	0.0%	0.15 <sup>+0.08</sup> <sub>-0.06</sub>
BooII	39 <sup>+1</sup> <sub>-1</sub>	181 <sup>+104</sup> <sub>-56</sub>	0.65 <sup>+0.11</sup> <sub>-0.11</sub>	0.45%	0.02 <sup>+0.02</sup> <sub>-0.01</sub>	39 <sup>+1</sup> <sub>-1</sub>	387 <sup>+738</sup> <sub>-198</sub>	0.92 <sup>+0.84</sup> <sub>-0.12</sub>	36.95%	0.02 <sup>+0.0</sup> <sub>-0.0</sub>
CVenI	62 <sup>+59</sup> <sub>-39</sub>	242 <sup>+26</sup> <sub>-8</sub>	0.6 <sup>+0.23</sup> <sub>-0.2</sub>	0.25%	0.6 <sup>+0.04</sup> <sub>-0.15</sub>	81 <sup>+73</sup> <sub>-55</sub>	290 <sup>+220</sup> <sub>-31</sub>	0.66 <sup>+0.25</sup> <sub>-0.12</sub>	6.5%	0.48 <sup>+0.03</sup> <sub>-0.06</sub>
CVenII	31 <sup>+48</sup> <sub>-23</sub>	193 <sup>+27</sup> <sub>-7</sub>	0.73 <sup>+0.19</sup> <sub>-0.23</sub>	0.3%	0.55 <sup>+0.03</sup> <sub>-0.13</sub>	37 <sup>+59</sup> <sub>-28</sub>	222 <sup>+103</sup> <sub>-16</sub>	0.77 <sup>+0.18</sup> <sub>-0.15</sub>	4.55%	0.4 <sup>+0.06</sup> <sub>-0.08</sub>
CarI	108 <sup>+5</sup> <sub>-17</sub>	117 <sup>+36</sup> <sub>-13</sub>	0.08 <sup>+0.09</sup> <sub>-0.05</sub>	0.0%	0.02 <sup>+0.96</sup> <sub>-0.02</sub>	108 <sup>+5</sup> <sub>-5</sub>	201 <sup>+154</sup> <sub>-61</sub>	0.3 <sup>+0.22</sup> <sub>-0.15</sub>	0.7%	0.0 <sup>+0.01</sup> <sub>-0.0</sub>
CarII	27 <sup>+1</sup> <sub>-1</sub>	252 <sup>+34</sup> <sub>-26</sub>	0.8 <sup>+0.02</sup> <sub>-0.01</sub>	0.0%	0.04 <sup>+0.01</sup> <sub>-0.01</sub>	28 <sup>+1</sup> <sub>-1</sub>	—	> 1	94.05%	—
CarIII	29 <sup>+1</sup> <sub>-0</sub>	227 <sup>+64</sup> <sub>-42</sub>	0.78 <sup>+0.04</sup> <sub>-0.04</sub>	0.0%	0.01 <sup>+0.0</sup> <sub>-0.0</sub>	29 <sup>+1</sup> <sub>-0</sub>	1196 <sup>+2881</sup> <sub>-651</sub>	≥ 1	65.65%	0.01 <sup>+0.0</sup> <sub>-0.0</sub>
ColII	186 <sup>+8</sup> <sub>-8</sub>	404 <sup>+1734</sup> <sub>-208</sub>	≥ 1	83.6%	0.08 <sup>+0.03</sup> <sub>-0.01</sub>	186 <sup>+8</sup> <sub>-8</sub>	—	> 1	94.1%	—
CberI	42 <sup>+1</sup> <sub>-1</sub>	83 <sup>+18</sup> <sub>-13</sub>	0.32 <sup>+0.07</sup> <sub>-0.06</sub>	0.0%	0.06 <sup>+0.04</sup> <sub>-0.02</sub>	43 <sup>+1</sup> <sub>-1</sub>	119 <sup>+41</sup> <sub>-26</sub>	0.47 <sup>+0.1</sup> <sub>-0.09</sub>	0.0%	0.03 <sup>+0.02</sup> <sub>-0.01</sub>
CraI	119 <sup>+27</sup> <sub>-75</sub>	147 <sup>+144</sup> <sub>-2</sub>	0.35 <sup>+0.41</sup> <sub>-0.24</sub>	5.9%	0.91 <sup>+0.08</sup> <sub>-0.89</sub>	144 <sup>+2</sup> <sub>-89</sub>	149 <sup>+338</sup> <sub>-3</sub>	0.53 <sup>+1.04</sup> <sub>-0.38</sub>	24.35%	0.04 <sup>+0.94</sup> <sub>-0.02</sub>
CraII	33 <sup>+8</sup> <sub>-8</sub>	133 <sup>+2</sup> <sub>-2</sub>	0.6 <sup>+0.08</sup> <sub>-0.07</sub>	0.0%	0.61 <sup>+0.02</sup> <sub>-0.02</sub>	38 <sup>+11</sup> <sub>-10</sub>	145 <sup>+7</sup> <sub>-4</sub>	0.58 <sup>+0.08</sup> <sub>-0.07</sub>	0.0%	0.51 <sup>+0.03</sup> <sub>-0.04</sub>
DraI	45 <sup>+6</sup> <sub>-5</sub>	108 <sup>+10</sup> <sub>-7</sub>	0.41 <sup>+0.02</sup> <sub>-0.02</sub>	0.0%	0.45 <sup>+0.03</sup> <sub>-0.03</sub>	50 <sup>+6</sup> <sub>-5</sub>	133 <sup>+19</sup> <sub>-13</sub>	0.46 <sup>+0.01</sup> <sub>-0.01</sub>	0.0%	0.3 <sup>+0.04</sup> <sub>-0.05</sub>
DraII	20 <sup>+1</sup> <sub>-1</sub>	98 <sup>+12</sup> <sub>-10</sub>	0.66 <sup>+0.03</sup> <sub>-0.02</sub>	0.0%	0.06 <sup>+0.01</sup> <sub>-0.01</sub>	20 <sup>+1</sup> <sub>-0</sub>	146 <sup>+32</sup> <sub>-22</sub>	0.76 <sup>+0.04</sup> <sub>-0.03</sub>	0.0%	0.03 <sup>+0.01</sup> <sub>-0.01</sub>
EriII	345 <sup>+23</sup> <sub>-237</sub>	485 <sup>+822</sup> <sub>-50</sub>	≥ 1	67.85%	1.0 <sup>+0.0</sup> <sub>-0.0</sub>	348 <sup>+21</sup> <sub>-187</sub>	617 <sup>+745</sup> <sub>-102</sub>	≥ 1	78.3%	1.0 <sup>+0.0</sup> <sub>-0.0</sub>
FnxI	69 <sup>+16</sup> <sub>-13</sub>	147 <sup>+4</sup> <sub>-3</sub>	0.36 <sup>+0.08</sup> <sub>-0.08</sub>	0.0%	0.77 <sup>+0.03</sup> <sub>-0.04</sub>	92 <sup>+21</sup> <sub>-19</sub>	157 <sup>+16</sup> <sub>-7</sub>	0.26 <sup>+0.09</sup> <sub>-0.05</sub>	0.0%	0.61 <sup>+0.09</sup> <sub>-0.16</sub>
GruI	22 <sup>+15</sup> <sub>-13</sub>	224 <sup>+28</sup> <sub>-20</sub>	0.82 <sup>+0.1</sup> <sub>-0.08</sub>	0.0%	0.25 <sup>+0.02</sup> <sub>-0.04</sub>	24 <sup>+17</sup> <sub>-14</sub>	430 <sup>+281</sup> <sub>-91</sub>	0.91 <sup>+0.05</sup> <sub>-0.03</sub>	3.35%	0.23 <sup>+0.04</sup> <sub>-0.05</sub>
GruII	27 <sup>+11</sup> <sub>-9</sub>	74 <sup>+28</sup> <sub>-13</sub>	0.48 <sup>+0.07</sup> <sub>-0.03</sub>	0.05%	0.35 <sup>+0.08</sup> <sub>-0.13</sub>	28 <sup>+12</sup> <sub>-10</sub>	87 <sup>+57</sup> <sub>-21</sub>	0.54 <sup>+0.07</sup> <sub>-0.02</sub>	0.75%	0.26 <sup>+0.11</sup> <sub>-0.14</sub>
HerI	52 <sup>+17</sup> <sub>-15</sub>	212 <sup>+37</sup> <sub>-20</sub>	0.61 <sup>+0.07</sup> <sub>-0.04</sub>	0.0%	0.31 <sup>+0.05</sup> <sub>-0.07</sub>	59 <sup>+18</sup> <sub>-17</sub>	390 <sup>+378</sup> <sub>-107</sub>	0.76 <sup>+0.11</sup> <sub>-0.03</sub>	6.95%	0.25 <sup>+0.03</sup> <sub>-0.05</sub>
HorI	87 <sup>+13</sup> <sub>-17</sub>	156 <sup>+258</sup> <sub>-72</sub>	0.37 <sup>+0.5</sup> <sub>-0.22</sub>	12.2%	0.05 <sup>+0.37</sup> <sub>-0.02</sub>	87 <sup>+13</sup> <sub>-13</sub>	194 <sup>+517</sup> <sub>-104</sub>	0.79 <sup>+1.61</sup> <sub>-0.53</sub>	41.3%	0.03 <sup>+0.1</sup> <sub>-0.0</sub>
HorII	78 <sup>+7</sup> <sub>-26</sub>	127 <sup>+305</sup> <sub>-51</sub>	0.74 <sup>+2.99</sup> <sub>-0.5</sub>	39.45%	0.01 <sup>+0.88</sup> <sub>-0.01</sub>	79 <sup>+6</sup> <sub>-13</sub>	107 <sup>+384</sup> <sub>-31</sub>	≥ 1	57.3%	0.01 <sup>+0.74</sup> <sub>-0.01</sub>
HyaII	128 <sup>+14</sup> <sub>-71</sub>	254 <sup>+429</sup> <sub>-58</sub>	≥ 1	59.35%	0.24 <sup>+0.12</sup> <sub>-0.07</sub>	130 <sup>+12</sup> <sub>-63</sub>	291 <sup>+627</sup> <sub>-63</sub>	≥ 1	75.4%	0.24 <sup>+0.08</sup> <sub>-0.07</sub>
HyiI	25 <sup>+0</sup> <sub>-0</sub>	167 <sup>+29</sup> <sub>-23</sub>	0.74 <sup>+0.03</sup> <sub>-0.03</sub>	0.0%	0.01 <sup>+0.0</sup> <sub>-0.0</sub>	25 <sup>+0</sup> <sub>-0</sub>	518 <sup>+678</sup> <sub>-201</sub>	0.91 <sup>+0.06</sup> <sub>-0.06</sub>	9.6%	0.01 <sup>+0.0</sup> <sub>-0.0</sub>
LeoI	56 <sup>+30</sup> <sub>-27</sub>	888 <sup>+578</sup> <sub>-185</sub>	0.9 <sup>+0.04</sup> <sub>-0.02</sub>	3.75%	0.81 <sup>+0.05</sup> <sub>-0.05</sub>	67 <sup>+35</sup> <sub>-33</sub>	—	> 1	100.0%	—
LeoII	79 <sup>+72</sup> <sub>-47</sub>	242 <sup>+14</sup> <sub>-11</sub>	0.51 <sup>+0.26</sup> <sub>-0.25</sub>	0.05%	0.85 <sup>+0.03</sup> <sub>-0.11</sub>	114 <sup>+104</sup> <sub>-75</sub>	254 <sup>+146</sup> <sub>-16</sub>	0.47 <sup>+0.33</sup> <sub>-0.22</sub>	4.0%	0.72 <sup>+0.1</sup> <sub>-0.34</sub>
LeoIV	84 <sup>+72</sup> <sub>-69</sub>	157 <sup>+116</sup> <sub>-5</sub>	0.59 <sup>+0.37</sup> <sub>-0.41</sub>	11.75%	0.97 <sup>+0.02</sup> <sub>-0.95</sub>	123 <sup>+33</sup> <sub>-107</sub>	156 <sup>+154</sup> <sub>-5</sub>	0.74 <sup>+1.32</sup> <sub>-0.49</sub>	27.2%	0.94 <sup>+0.05</sup> <sub>-0.92</sub>
LeoV	164 <sup>+6</sup> <sub>-101</sub>	197 <sup>+460</sup> <sub>-18</sub>	0.97 <sup>+3.16</sup> <sub>-0.64</sub>	48.15%	0.13 <sup>+0.58</sup> <sub>-0.07</sub>	165 <sup>+5</sup> <sub>-81</sub>	203 <sup>+462</sup> <sub>-18</sub>	≥ 1	65.1%	0.13 <sup>+0.44</sup> <sub>-0.07</sub>
PhxII	80 <sup>+4</sup> <sub>-4</sub>	276 <sup>+379</sup> <sub>-117</sub>	0.58 <sup>+0.33</sup> <sub>-0.22</sub>	10.0%	0.03 <sup>+0.03</sup> <sub>-0.0</sub>	80 <sup>+4</sup> <sub>-4</sub>	491 <sup>+1283</sup> <sub>-279</sub>	≥ 1	66.9%	0.04 <sup>+0.0</sup> <sub>-0.0</sub>
PisII	181 <sup>+12</sup> <sub>-12</sub>	—	> 1	95.85%	—	181 <sup>+12</sup> <sub>-12</sub>	—	> 1	97.9%	—
RetII	28 <sup>+2</sup> <sub>-3</sub>	63 <sup>+15</sup> <sub>-10</sub>	0.39 <sup>+0.05</sup> <sub>-0.03</sub>	0.0%	0.18 <sup>+0.07</sup> <sub>-0.05</sub>	28 <sup>+2</sup> <sub>-3</sub>	77 <sup>+28</sup> <sub>-17</sub>	0.47 <sup>+0.08</sup> <sub>-0.06</sub>	0.0%	0.13 <sup>+0.06</sup> <sub>-0.05</sub>
RetIII	27 <sup>+46</sup> <sub>-22</sub>	114 <sup>+50</sup> <sub>-15</sub>	0.67 <sup>+0.25</sup> <sub>-0.22</sub>	3.05%	0.53 <sup>+0.05</sup> <sub>-0.32</sub>	30 <sup>+47</sup> <sub>-25</sub>	124 <sup>+74</sup> <sub>-18</sub>	0.74 <sup>+0.23</sup> <sub>-0.21</sub>	11.2%	0.43 <sup>+0.07</sup> <sub>-0.28</sub>

Table 4 continued on next page

**Table 4** (*continued*)

name	Model PE <sub>HM</sub>					Model P <sub>NFW</sub>				
	$r_{\text{peri}}$ (kpc)	$r_{\text{apo}}^{a,c}$ (kpc)	$e$	$P_{\text{unb}}$	$P^{b,c}$	$r_{\text{peri}}$ (kpc)	$r_{\text{apo}}^{a,c}$ (kpc)	$e$	$P_{\text{unb}}$	$P^{b,c}$
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)
SgrII	45 <sup>+6</sup> <sub>-7</sub>	109 <sup>+28</sup> <sub>-17</sub>	0.42 <sup>+0.04</sup> <sub>-0.01</sub>	0.0%	0.25 <sup>+0.1</sup> <sub>-0.09</sub>	47 <sup>+5</sup> <sub>-7</sub>	158 <sup>+91</sup> <sub>-41</sub>	0.54 <sup>+0.12</sup> <sub>-0.05</sub>	0.4%	0.13 <sup>+0.1</sup> <sub>-0.06</sub>
ScII	54 <sup>+3</sup> <sub>-3</sub>	107 <sup>+4</sup> <sub>-4</sub>	0.32 <sup>+0.03</sup> <sub>-0.02</sub>	0.0%	0.48 <sup>+0.04</sup> <sub>-0.04</sub>	62 <sup>+3</sup> <sub>-3</sub>	133 <sup>+9</sup> <sub>-7</sub>	0.36 <sup>+0.01</sup> <sub>-0.01</sub>	0.0%	0.3 <sup>+0.05</sup> <sub>-0.05</sub>
SegI	21 <sup>+4</sup> <sub>-5</sub>	58 <sup>+26</sup> <sub>-15</sub>	0.47 <sup>+0.07</sup> <sub>-0.02</sub>	0.05%	0.19 <sup>+0.11</sup> <sub>-0.08</sub>	21 <sup>+4</sup> <sub>-5</sub>	68 <sup>+46</sup> <sub>-21</sub>	0.53 <sup>+0.12</sup> <sub>-0.04</sub>	0.5%	0.14 <sup>+0.11</sup> <sub>-0.08</sub>
SegII	19 <sup>+6</sup> <sub>-4</sub>	47 <sup>+4</sup> <sub>-3</sub>	0.42 <sup>+0.06</sup> <sub>-0.08</sub>	0.0%	0.69 <sup>+0.03</sup> <sub>-0.06</sub>	20 <sup>+6</sup> <sub>-4</sub>	48 <sup>+5</sup> <sub>-4</sub>	0.4 <sup>+0.06</sup> <sub>-0.08</sub>	0.0%	0.65 <sup>+0.04</sup> <sub>-0.08</sub>
SxtI	84 <sup>+4</sup> <sub>-4</sub>	188 <sup>+28</sup> <sub>-21</sub>	0.39 <sup>+0.04</sup> <sub>-0.03</sub>	0.0%	0.17 <sup>+0.04</sup> <sub>-0.04</sub>	87 <sup>+3</sup> <sub>-3</sub>	592 <sup>+713</sup> <sub>-217</sub>	0.75 <sup>+0.14</sup> <sub>-0.11</sub>	4.3%	0.12 <sup>+0.01</sup> <sub>-0.01</sub>
TriII	12 <sup>+1</sup> <sub>-1</sub>	110 <sup>+7</sup> <sub>-6</sub>	0.8 <sup>+0.01</sup> <sub>-0.01</sub>	0.0%	0.13 <sup>+0.01</sup> <sub>-0.01</sub>	12 <sup>+1</sup> <sub>-1</sub>	161 <sup>+18</sup> <sub>-14</sub>	0.86 <sup>+0.0</sup> <sub>-0.0</sub>	0.0%	0.07 <sup>+0.01</sup> <sub>-0.01</sub>
TucII	39 <sup>+13</sup> <sub>-11</sub>	191 <sup>+243</sup> <sub>-73</sub>	0.68 <sup>+0.24</sup> <sub>-0.06</sub>	11.7%	0.1 <sup>+0.06</sup> <sub>-0.03</sub>	40 <sup>+13</sup> <sub>-11</sub>	287 <sup>+630</sup> <sub>-138</sub>	0.96 <sup>+1.48</sup> <sub>-0.22</sub>	45.75%	0.09 <sup>+0.01</sup> <sub>-0.03</sub>
TucIII	3 <sup>+1</sup> <sub>-0</sub>	47 <sup>+6</sup> <sub>-4</sub>	0.87 <sup>+0.01</sup> <sub>-0.02</sub>	0.0%	0.24 <sup>+0.0</sup> <sub>-0.01</sub>	3 <sup>+1</sup> <sub>-0</sub>	51 <sup>+8</sup> <sub>-4</sub>	0.88 <sup>+0.01</sup> <sub>-0.01</sub>	0.0%	0.21 <sup>+0.01</sup> <sub>-0.01</sub>
TucIV	36 <sup>+8</sup> <sub>-9</sub>	72 <sup>+31</sup> <sub>-16</sub>	0.36 <sup>+0.07</sup> <sub>-0.02</sub>	0.0%	0.27 <sup>+0.15</sup> <sub>-0.13</sub>	37 <sup>+7</sup> <sub>-9</sub>	90 <sup>+66</sup> <sub>-28</sub>	0.43 <sup>+0.14</sup> <sub>-0.05</sub>	0.8%	0.18 <sup>+0.16</sup> <sub>-0.11</sub>
TucV	37 <sup>+14</sup> <sub>-14</sub>	138 <sup>+149</sup> <sub>-52</sub>	0.61 <sup>+0.14</sup> <sub>-0.03</sub>	5.6%	0.14 <sup>+0.11</sup> <sub>-0.06</sub>	38 <sup>+14</sup> <sub>-14</sub>	182 <sup>+385</sup> <sub>-81</sub>	0.77 <sup>+1.04</sup> <sub>-0.11</sub>	26.9%	0.1 <sup>+0.07</sup> <sub>-0.03</sub>
UMaI	53 <sup>+33</sup> <sub>-20</sub>	102 <sup>+5</sup> <sub>-5</sub>	0.32 <sup>+0.19</sup> <sub>-0.19</sub>	0.0%	0.98 <sup>+0.01</sup> <sub>-0.01</sub>	68 <sup>+37</sup> <sub>-30</sub>	103 <sup>+24</sup> <sub>-5</sub>	0.25 <sup>+0.22</sup> <sub>-0.17</sub>	0.15%	0.97 <sup>+0.01</sup> <sub>-0.96</sub>
UMaII	39 <sup>+2</sup> <sub>-2</sub>	110 <sup>+52</sup> <sub>-29</sub>	0.48 <sup>+0.12</sup> <sub>-0.11</sub>	0.05%	0.06 <sup>+0.04</sup> <sub>-0.03</sub>	39 <sup>+2</sup> <sub>-2</sub>	178 <sup>+206</sup> <sub>-68</sub>	0.65 <sup>+0.2</sup> <sub>-0.15</sub>	5.35%	0.03 <sup>+0.03</sup> <sub>-0.01</sub>
UMiI	42 <sup>+3</sup> <sub>-3</sub>	92 <sup>+4</sup> <sub>-3</sub>	0.37 <sup>+0.03</sup> <sub>-0.02</sub>	0.0%	0.55 <sup>+0.02</sup> <sub>-0.03</sub>	48 <sup>+4</sup> <sub>-4</sub>	105 <sup>+6</sup> <sub>-5</sub>	0.37 <sup>+0.02</sup> <sub>-0.01</sub>	0.0%	0.41 <sup>+0.04</sup> <sub>-0.04</sub>
WiII	35 <sup>+31</sup> <sub>-14</sub>	54 <sup>+30</sup> <sub>-9</sub>	0.25 <sup>+0.14</sup> <sub>-0.16</sub>	0.55%	0.94 <sup>+0.02</sup> <sub>-0.87</sub>	40 <sup>+27</sup> <sub>-17</sub>	53 <sup>+55</sup> <sub>-9</sub>	0.25 <sup>+0.17</sup> <sub>-0.16</sub>	2.95%	0.91 <sup>+0.04</sup> <sub>-0.89</sub>

NOTE—Column 1 lists the abbreviated dwarf galaxy name; Column 2 and 7 gives the pericenter of the orbit for the two potential models; Column 3 and 8 gives the apocenter of the orbit for the two potential models; Column 4 and 9 is the eccentricity of the orbit for the two potential models; in Column 5 and 9, we provide the probability of the galaxy being unbound for the two potential models. Column 5 and 11 gives the orbital phase.

<sup>a</sup>The apocenter is only for samples with elliptical orbit.

<sup>b</sup>The orbital phase probability defined in Eq. 8 to characterize the chance for a dwarf galaxy to be that close to its pericenter.

<sup>c</sup>When the value of total energy minus  $1\sigma$  is larger than  $0 \text{ km}^2 \text{ s}^{-2}$  ( $P_{\text{unb}} > 84.13\%$ ), we only quote “–” in the table.

**Table 5.** Orbital properties of dwarf galaxies for Model PE<sub>IM</sub> and Model PE<sub>LM</sub>.

name	Model PE <sub>IM</sub>					Model PE <sub>LM</sub>				
	$r_{\text{peri}}$ (kpc)	$r_{\text{apo}}^{a,c}$ (kpc)	$e$	$P_{\text{unb}}$	$P^{b,c}$	$r_{\text{peri}}$ (kpc)	$r_{\text{apo}}^{a,c}$ (kpc)	$e$	$P_{\text{unb}}$	$P^{b,c}$
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)
AntII	83 <sup>+16</sup> <sub>-17</sub>	237 <sup>+117</sup> <sub>-47</sub>	0.49 <sup>+0.07</sup> <sub>-0.02</sub>	0.25%	0.25 <sup>+0.1</sup> <sub>-0.05</sub>	102 <sup>+11</sup> <sub>-13</sub>	1078 <sup>+2506</sup> <sub>-584</sub>	$\geq 1$	56.8%	0.26 <sup>+0.02</sup> <sub>-0.02</sub>
AquII	102 <sup>+4</sup> <sub>-62</sub>	117 <sup>+185</sup> <sub>-7</sub>	$\geq 1$	66.9%	0.07 <sup>+0.57</sup> <sub>-0.02</sub>	102 <sup>+3</sup> <sub>-43</sub>	124 <sup>+167</sup> <sub>-10</sub>	$\geq 1$	75.35%	0.07 <sup>+0.31</sup> <sub>-0.02</sub>
BooI	53 <sup>+4</sup> <sub>-5</sub>	244 <sup>+280</sup> <sub>-88</sub>	0.65 <sup>+0.18</sup> <sub>-0.12</sub>	3.25%	0.07 <sup>+0.05</sup> <sub>-0.01</sub>	55 <sup>+3</sup> <sub>-4</sub>	–	$> 1$	95.5%	–
BooII	39 <sup>+1</sup> <sub>-1</sub>	–	$> 1$	86.35%	–	39 <sup>+1</sup> <sub>-1</sub>	–	$> 1$	99.95%	–

**Table 5** continued on next page

Table 5 (continued)

name	Model PE <sub>IM</sub>					Model PE <sub>LM</sub>				
	$r_{\text{peri}}$ (kpc)	$r_{\text{apo}}^{a,c}$ (kpc)	$e$	$P_{\text{unb}}$	$P^{b,c}$	$r_{\text{peri}}$ (kpc)	$r_{\text{apo}}^{a,c}$ (kpc)	$e$	$P_{\text{unb}}$	$P^{b,c}$
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)
CVenI	104 <sup>+67</sup> <sub>-75</sub>	392 <sup>+678</sup> <sub>-87</sub>	0.83 <sup>+0.55</sup> <sub>-0.14</sub>	25.95%	0.51 <sup>+0.05</sup> <sub>-0.11</sub>	136 <sup>+44</sup> <sub>-96</sub>	679 <sup>+1315</sup> <sub>-232</sub>	≥ 1	61.3%	0.55 <sup>+0.05</sup> <sub>-0.1</sub>
CVenII	42 <sup>+66</sup> <sub>-32</sub>	288 <sup>+269</sup> <sub>-40</sub>	0.87 <sup>+0.25</sup> <sub>-0.11</sub>	17.75%	0.31 <sup>+0.07</sup> <sub>-0.04</sub>	56 <sup>+63</sup> <sub>-46</sub>	546 <sup>+945</sup> <sub>-119</sub>	1.0 <sup>+1.37</sup> <sub>-0.06</sub>	48.8%	0.39 <sup>+0.04</sup> <sub>-0.06</sub>
CarI	108 <sup>+5</sup> <sub>-5</sub>	589 <sup>+1348</sup> <sub>-324</sub>	0.87 <sup>+0.49</sup> <sub>-0.35</sub>	38.05%	0.0 <sup>+0.0</sup> <sub>-0.0</sub>	108 <sup>+5</sup> <sub>-5</sub>	–	> 1	99.15%	–
CarII	28 <sup>+1</sup> <sub>-1</sub>	–	> 1	100.0%	–	28 <sup>+1</sup> <sub>-1</sub>	–	> 1	100.0%	–
CarIII	29 <sup>+1</sup> <sub>-0</sub>	–	> 1	100.0%	–	29 <sup>+1</sup> <sub>-0</sub>	–	> 1	100.0%	–
ColI	186 <sup>+8</sup> <sub>-7</sub>	–	> 1	97.65%	–	187 <sup>+8</sup> <sub>-7</sub>	–	> 1	99.4%	–
CberI	43 <sup>+1</sup> <sub>-1</sub>	201 <sup>+227</sup> <sub>-72</sub>	0.66 <sup>+0.18</sup> <sub>-0.14</sub>	3.6%	0.01 <sup>+0.02</sup> <sub>-0.0</sub>	43 <sup>+1</sup> <sub>-1</sub>	–	> 1	94.9%	–
CraI	145 <sup>+2</sup> <sub>-74</sub>	150 <sup>+557</sup> <sub>-5</sub>	0.9 <sup>+2.12</sup> <sub>-0.66</sub>	45.65%	0.03 <sup>+0.94</sup> <sub>-0.01</sub>	145 <sup>+2</sup> <sub>-3</sub>	167 <sup>+751</sup> <sub>-22</sub>	≥ 1	68.0%	0.03 <sup>+0.12</sup> <sub>-0.01</sub>
CraII	43 <sup>+13</sup> <sub>-12</sub>	170 <sup>+24</sup> <sub>-12</sub>	0.6 <sup>+0.07</sup> <sub>-0.04</sub>	0.0%	0.36 <sup>+0.05</sup> <sub>-0.08</sub>	58 <sup>+15</sup> <sub>-17</sub>	396 <sup>+448</sup> <sub>-137</sub>	0.76 <sup>+0.13</sup> <sub>-0.03</sub>	7.55%	0.23 <sup>+0.02</sup> <sub>-0.05</sub>
DraI	54 <sup>+7</sup> <sub>-6</sub>	198 <sup>+79</sup> <sub>-38</sub>	0.57 <sup>+0.07</sup> <sub>-0.03</sub>	0.0%	0.15 <sup>+0.05</sup> <sub>-0.04</sub>	61 <sup>+6</sup> <sub>-5</sub>	–	> 1	86.7%	–
DraII	20 <sup>+1</sup> <sub>-0</sub>	304 <sup>+308</sup> <sub>-102</sub>	0.88 <sup>+0.07</sup> <sub>-0.05</sub>	3.95%	0.02 <sup>+0.0</sup> <sub>-0.0</sub>	20 <sup>+0</sup> <sub>-0</sub>	–	> 1	100.0%	–
EriII	349 <sup>+20</sup> <sub>-138</sub>	–	> 1	84.95%	–	349 <sup>+20</sup> <sub>-104</sub>	–	> 1	93.55%	–
FnxI	116 <sup>+13</sup> <sub>-22</sub>	218 <sup>+148</sup> <sub>-48</sub>	0.33 <sup>+0.16</sup> <sub>-0.04</sub>	0.35%	0.24 <sup>+0.23</sup> <sub>-0.09</sub>	130 <sup>+5</sup> <sub>-7</sub>	904 <sup>+2232</sup> <sub>-500</sub>	≥ 1	55.95%	0.18 <sup>+0.02</sup> <sub>-0.02</sub>
GruI	25 <sup>+18</sup> <sub>-15</sub>	–	> 1	88.15%	–	28 <sup>+21</sup> <sub>-18</sub>	–	> 1	100.0%	–
GruII	29 <sup>+12</sup> <sub>-10</sub>	102 <sup>+118</sup> <sub>-31</sub>	0.59 <sup>+0.17</sup> <sub>-0.03</sub>	6.05%	0.19 <sup>+0.13</sup> <sub>-0.11</sub>	31 <sup>+11</sup> <sub>-11</sub>	165 <sup>+352</sup> <sub>-76</sub>	0.88 <sup>+0.74</sup> <sub>-0.19</sub>	40.1%	0.09 <sup>+0.06</sup> <sub>-0.03</sub>
HerI	64 <sup>+17</sup> <sub>-19</sub>	1336 <sup>+3149</sup> <sub>-677</sub>	≥ 1	71.2%	0.3 <sup>+0.02</sup> <sub>-0.02</sub>	73 <sup>+15</sup> <sub>-19</sub>	–	> 1	100.0%	–
HorI	88 <sup>+13</sup> <sub>-12</sub>	212 <sup>+588</sup> <sub>-126</sub>	≥ 1	67.7%	0.04 <sup>+0.01</sup> <sub>-0.0</sub>	88 <sup>+13</sup> <sub>-11</sub>	–	> 1	90.15%	–
HorII	79 <sup>+6</sup> <sub>-10</sub>	87 <sup>+238</sup> <sub>-12</sub>	≥ 1	70.3%	0.01 <sup>+0.21</sup> <sub>-0.01</sub>	79 <sup>+6</sup> <sub>-8</sub>	84 <sup>+304</sup> <sub>-9</sub>	≥ 1	80.85%	0.01 <sup>+0.02</sup> <sub>-0.01</sub>
HyaII	131 <sup>+12</sup> <sub>-56</sub>	–	> 1	85.45%	–	132 <sup>+11</sup> <sub>-45</sub>	–	> 1	96.85%	–
HyII	25 <sup>+0</sup> <sub>-0</sub>	–	> 1	98.85%	–	25 <sup>+0</sup> <sub>-0</sub>	–	> 1	100.0%	–
LeoI	77 <sup>+37</sup> <sub>-39</sub>	–	> 1	100.0%	–	89 <sup>+36</sup> <sub>-43</sub>	–	> 1	100.0%	–
LeoII	182 <sup>+46</sup> <sub>-134</sub>	274 <sup>+498</sup> <sub>-30</sub>	0.61 <sup>+0.59</sup> <sub>-0.29</sub>	19.0%	0.51 <sup>+0.24</sup> <sub>-0.2</sub>	215 <sup>+18</sup> <sub>-134</sub>	326 <sup>+870</sup> <sub>-69</sub>	0.95 <sup>+1.9</sup> <sub>-0.51</sub>	47.05%	0.42 <sup>+0.2</sup> <sub>-0.16</sub>
LeoIV	150 <sup>+7</sup> <sub>-132</sub>	156 <sup>+110</sup> <sub>-5</sub>	0.9 <sup>+2.92</sup> <sub>-0.57</sub>	40.1%	0.03 <sup>+0.95</sup> <sub>-0.01</sub>	152 <sup>+5</sup> <sub>-131</sub>	157 <sup>+219</sup> <sub>-5</sub>	≥ 1	52.35%	0.02 <sup>+0.94</sup> <sub>-0.01</sub>
LeoV	166 <sup>+5</sup> <sub>-56</sub>	218 <sup>+429</sup> <sub>-23</sub>	≥ 1	75.15%	0.12 <sup>+0.2</sup> <sub>-0.06</sub>	166 <sup>+4</sup> <sub>-28</sub>	–	> 1	84.4%	–
PhxII	80 <sup>+4</sup> <sub>-4</sub>	–	> 1	94.35%	–	81 <sup>+4</sup> <sub>-4</sub>	–	> 1	99.95%	–
PisII	181 <sup>+12</sup> <sub>-12</sub>	–	> 1	98.75%	–	181 <sup>+12</sup> <sub>-12</sub>	–	> 1	99.3%	–
RetII	28 <sup>+2</sup> <sub>-3</sub>	93 <sup>+57</sup> <sub>-26</sub>	0.54 <sup>+0.13</sup> <sub>-0.09</sub>	0.35%	0.09 <sup>+0.06</sup> <sub>-0.05</sub>	29 <sup>+2</sup> <sub>-2</sub>	235 <sup>+488</sup> <sub>-122</sub>	0.92 <sup>+0.43</sup> <sub>-0.23</sub>	39.95%	0.03 <sup>+0.01</sup> <sub>-0.01</sub>
RetIII	33 <sup>+47</sup> <sub>-27</sub>	136 <sup>+109</sup> <sub>-23</sub>	0.84 <sup>+0.59</sup> <sub>-0.19</sub>	20.85%	0.3 <sup>+0.12</sup> <sub>-0.16</sub>	40 <sup>+43</sup> <sub>-34</sub>	183 <sup>+269</sup> <sub>-42</sub>	0.96 <sup>+2.0</sup> <sub>-0.14</sub>	39.65%	0.19 <sup>+0.06</sup> <sub>-0.05</sub>
SgrII	49 <sup>+5</sup> <sub>-6</sub>	287 <sup>+560</sup> <sub>-131</sub>	0.77 <sup>+0.42</sup> <sub>-0.17</sub>	20.3%	0.09 <sup>+0.04</sup> <sub>-0.02</sub>	51 <sup>+4</sup> <sub>-5</sub>	–	> 1	92.95%	–
ScII	66 <sup>+3</sup> <sub>-3</sub>	209 <sup>+44</sup> <sub>-29</sub>	0.52 <sup>+0.05</sup> <sub>-0.04</sub>	0.0%	0.13 <sup>+0.04</sup> <sub>-0.04</sub>	73 <sup>+3</sup> <sub>-3</sub>	–	> 1	98.25%	–
SegI	21 <sup>+4</sup> <sub>-5</sub>	76 <sup>+82</sup> <sub>-27</sub>	0.57 <sup>+0.19</sup> <sub>-0.07</sub>	3.8%	0.11 <sup>+0.12</sup> <sub>-0.08</sub>	22 <sup>+4</sup> <sub>-5</sub>	106 <sup>+273</sup> <sub>-50</sub>	0.82 <sup>+0.64</sup> <sub>-0.23</sub>	34.1%	0.04 <sup>+0.1</sup> <sub>-0.01</sub>
SegII	21 <sup>+7</sup> <sub>-4</sub>	49 <sup>+6</sup> <sub>-4</sub>	0.4 <sup>+0.06</sup> <sub>-0.07</sub>	0.0%	0.62 <sup>+0.05</sup> <sub>-0.1</sub>	23 <sup>+8</sup> <sub>-5</sub>	53 <sup>+14</sup> <sub>-6</sub>	0.4 <sup>+0.06</sup> <sub>-0.03</sub>	0.0%	0.5 <sup>+0.09</sup> <sub>-0.18</sub>
SxtI	88 <sup>+3</sup> <sub>-3</sub>	–	> 1	99.75%	–	90 <sup>+3</sup> <sub>-3</sub>	–	> 1	100.0%	–
TriII	12 <sup>+1</sup> <sub>-1</sub>	363 <sup>+189</sup> <sub>-87</sub>	0.93 <sup>+0.02</sup> <sub>-0.01</sub>	0.55%	0.04 <sup>+0.01</sup> <sub>-0.01</sub>	12 <sup>+1</sup> <sub>-1</sub>	–	> 1	100.0%	–
TucII	41 <sup>+13</sup> <sub>-11</sub>	350 <sup>+1109</sup> <sub>-169</sub>	≥ 1	79.8%	0.1 <sup>+0.01</sup> <sub>-0.02</sub>	42 <sup>+13</sup> <sub>-11</sub>	–	> 1	99.45%	–

Table 5 continued on next page



**Table 5** (*continued*)

name	Model PE <sub>IM</sub>					Model PE <sub>LM</sub>				
	$r_{\text{peri}}$ (kpc)	$r_{\text{apo}}^{a,c}$ (kpc)	$e$	$P_{\text{unb}}$	$P^{b,c}$	$r_{\text{peri}}$ (kpc)	$r_{\text{apo}}^{a,c}$ (kpc)	$e$	$P_{\text{unb}}$	$P^{b,c}$
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)
TucIII	$3_{-0}^{+1}$	$53_{-5}^{+9}$	$0.89_{-0.01}^{+0.01}$	0.0%	$0.19_{-0.02}^{+0.01}$	$3_{-0}^{+1}$	$68_{-9}^{+22}$	$0.91_{-0.0}^{+0.0}$	0.0%	$0.12_{-0.03}^{+0.02}$
TucIV	$37_{-9}^{+7}$	$109_{-42}^{+157}$	$0.52_{-0.1}^{+0.29}$	7.95%	$0.11_{-0.07}^{+0.17}$	$39_{-8}^{+6}$	$176_{-96}^{+395}$	$\geq 1$	52.35%	$0.06_{-0.01}^{+0.04}$
TucV	$38_{-14}^{+14}$	$207_{-101}^{+472}$	$\geq 1$	55.75%	$0.1_{-0.03}^{+0.01}$	$40_{-14}^{+13}$	—	$> 1$	89.4%	—
UMaI	$90_{-46}^{+17}$	$104_{-6}^{+104}$	$0.27_{-0.19}^{+0.29}$	4.25%	$0.92_{-0.92}^{+0.05}$	$102_{-28}^{+5}$	$161_{-63}^{+507}$	$0.56_{-0.41}^{+0.96}$	30.05%	$0.01_{-0.01}^{+0.9}$
UMaII	$40_{-2}^{+2}$	$301_{-158}^{+673}$	$0.93_{-0.27}^{+0.72}$	41.55%	$0.02_{-0.0}^{+0.0}$	$40_{-2}^{+2}$	—	$> 1$	96.95%	—
UMiI	$52_{-4}^{+4}$	$129_{-11}^{+14}$	$0.43_{-0.01}^{+0.01}$	0.0%	$0.27_{-0.05}^{+0.05}$	$60_{-3}^{+3}$	$702_{-282}^{+1059}$	$0.86_{-0.09}^{+0.11}$	10.35%	$0.12_{-0.01}^{+0.01}$
Will	$44_{-21}^{+23}$	$53_{-9}^{+70}$	$0.27_{-0.16}^{+0.3}$	8.2%	$0.86_{-0.85}^{+0.08}$	$52_{-25}^{+15}$	$53_{-10}^{+132}$	$0.33_{-0.22}^{+1.27}$	23.9%	$0.07_{-0.06}^{+0.84}$

NOTE—Similar to Table 4, but using two lighter MW potential models.

<sup>a</sup>The apocenter is only for samples with elliptical orbit.

<sup>b</sup>The orbital phase probability defined in Eq. 8 to characterize the chance for a dwarf galaxy to be that close to its pericenter.

<sup>c</sup>When the value of total energy minus  $1\sigma$  is larger than  $0 \text{ km}^2 \text{ s}^{-2}$  ( $P_{\text{unb}} > 84.13\%$ ), we only quote “—” in the table.

## 4. RESULTS AND DISCUSSION

### 4.1. Phase diagram and the bound nature of dwarf galaxies

Figure 1 shows the escape velocity curves of the four MW potential models superposed to the dwarf galaxy phase diagram. It demonstrates that the dwarf galaxies can be either almost all bound to the MW (models Model PE<sub>HM</sub> & PNFW), or half of them (model PE<sub>IM</sub>) or even most of them (model PE<sub>LM</sub>) could be unbound. This underlines how the nature of dwarf galaxies depends on our knowledge of the MW potential (Hammer et al. 2021). This calls for caution when interpreting the dwarf galaxy orbital properties. In discussing the dwarf galaxy orbits, one thus needs to always account for their dependency to the adopted MW mass profile, and also consider the possible impact of the LMC.

The position of the LMC is also given in Figure 1. Kallivayalil et al. (2013) discussed whether the LMC is bound or not to the MW, and concluded that for most MW mass models, it is likely at its first passage. In fact, their result depends on the LMC mass, for which they assumed a sufficiently high value ( $> 10^{11} M_{\odot}$ ) to keep the SMC bound to the LMC for more than 2 Gyr. The goal was to perform the modeling of the Magellanic Stream (Besla et al. 2012), assumed to be a tidal tail induced 1 Gyr ago, during an interaction between the Clouds before they entered the MW halo.

However, the Magellanic stream properties are apparently better reproduced by a “ram-pressure + collision” model (Hammer et al. 2015), which only requires the unquestionable collision between the two Clouds 250 to 300 Myr ago. This type of model recovers (Wang et al. 2019) the HI and stellar properties of the Magellanic Bridge, and additionally the unusual 30 kpc elongated shape of the SMC on the line of sight, discovered after examining the young variable star distribution (Ripepi et al. 2017). Wang et al. (2019) argued that it is the recent collision between the Clouds that dominates their past orbital history, and that a light LMC ( $\leq 2 \times 10^{10} M_{\odot}$ ) is required to let large amounts

of ionized gas ( $> 10^9 M_{\odot}$ , see Fox et al. 2014) be stripped from both LMC and SMC due to the ram-pressure exerted by the MW halo gas.

In Figure 1 there are several dwarf galaxies that share a kinetic energy similar to that of the LMC, namely Bootes II, Carina II, Carina III, Hydrus & Tucana II. Carina II, Carina III & Hydrus may have their orbits significantly affected by the LMC, or they could be bound to it (Patel et al. 2020), an issue that also depends on the total LMC mass. Therefore, and despite the accuracy of the Gaia EDR3 3D velocities (see error bars in Figure 1), the uncertainties in the mass and potential of the MW are too large for a robust conclusion on the fraction of dwarf galaxies that are bound to the MW.

In contrast to this, we confirm that the pericenter determination is very robust, and almost independent of the mass model (Simon 2018). Half of the dwarf galaxies have the same pericenters for all the different MW mass models, within less than 10% and well within their error bars. Only 3 dwarf galaxies (Antlia II, Crater II, and Fornax) show a pericenter almost 2 times smaller when using the high MW mass model ( $PE_{\text{HM}}$ ) compared to result from the low MW mass model ( $PE_{\text{LM}}$ ).

#### 4.2. Gaia EDR3 confirms the prominence of the VPOS

Figure 2 presents the orbital poles of all 46 dwarf galaxies, projected in an Hammer-Aitoff diagram. The angular momenta are calculated as the cross-product of the Galactocentric position vector  $\vec{r}_{\text{gc}}$ , and the 3D velocity in the MW frame,  $\vec{v}_{3\text{D}}$ , for each of the 2000 Monte-Carlo realizations. Figure 2 shows that the Gaia EDR3 accuracy is not sufficient to efficiently constraint the poles of dwarf galaxies beyond  $r_{\text{gc}} > 200$  kpc.

Whether a dwarf galaxy is part of the VPOS depends on its 3D position (it has to lie close to the VPOS plane), and orbital pole (which should be close to the VPOS normal vector). Since the latter contains information on both the position and velocity, we here focus on it to judge possible VPOS membership. Figure 2 thus already allows a simple visual confirmation of a preferred alignment of orbital poles with the VPOS. It indicates in pink the areas corresponding to the VPOS (from Figure 3 of Fritz et al. 2018), containing 10% of the area on the sphere around the adopted VPOS normal vector pointing towards Galactic coordinates  $(l, b) = (169.3^{\circ}, -2.8^{\circ})$ .

For a more quantitative analysis, we follow the method outlined in Fritz et al. (2018). Table 6 lists for all 46 dwarf galaxies in our sample, the best-possible alignment  $\theta_{\text{VPOS-3}}^{\text{predicted}}$  of their orbital pole with the VPOS normal vector that is defined by their spatial position (following the geometric method presented in Pawlowski & Kroupa 2013; Pawlowski et al. 2015), and the actually observed angular separation  $\theta_{\text{VPOS-3}}^{\text{measured}}$  between the median orbital pole and the VPOS normal. The latter is positive if the object co-orbits in the same sense as the majority of VPOS members (including the LMC and SMC), and negative if the object counter-orbits. We furthermore count the fraction  $f_{\text{inVPOS}}$  of Monte-Carlo realizations which result in orbital poles that are aligned with the VPOS normal vector to within an angle  $\theta_{\text{inVPOS}}$ . For consistency and direct comparability we adopt the same values as in Fritz et al. (2018):  $\theta_{\text{inVPOS}} = 36.9^{\circ}$  corresponding to 10% of the area of the sky, and a VPOS normal vector pointing to Galactic coordinates  $(l, b) = (169.3^{\circ}, -2.8^{\circ})$ . To further assess the degree to which a misalignment might simply be due to proper motion uncertainties, we also generate 2000 mock orbital pole realizations by placing the object’s intrinsic orbital pole in its predicted (best-aligned with the VPOS) position, and then vary its direction by drawing from the orbital pole uncertainty of the observed data. This way we determine the probability  $p_{\text{outsideVPOS}}$  of these mock-realizations to be found outside of  $\theta_{\text{inVPOS}}$  despite their intrinsic close alignment (i.e. the false negative rate), and



**Table 6** (*continued*)

Name	$\theta_{\text{VPOS-3}}^{\text{predicted}}$	$\theta_{\text{VPOS-3}}^{\text{measured}}$	$f_{\text{inVPOS}}$	$p_{\text{outsideVPOS}}$	$p_{>\text{obs}}$	volume-complete member
	( $^{\circ}$ )	( $^{\circ}$ )				
(1)	(2)	(3)	(4)	(5)	(6)	(7)
CarI	4.7	4.7	1.000	0.000	0.790	<i>Near – S, Dist – S</i>
CarII	3.5	40.5	0.005	0.000	0.000	<i>Near – S, Dist – S</i>
CarIII	7.1	11.7	1.000	0.000	0.000	<i>Near – S</i>
ColI	27.6	–89.6	0.005	0.128	0.000	
CberI	9.6	82.9	0.000	0.000	0.000	<i>Near – S, Dist – S</i>
CraI	9.6	28.9	0.555	0.253	0.366	<i>Dist – S</i>
CraII	15.2	20.0	0.991	0.002	0.187	<i>Near – S, Dist – S</i>
DraI	10.4	18.2	1.000	0.000	0.000	<i>Near – S, Dist – S</i>
DraII	29.9	39.3	0.036	0.000	0.000	
EriII	9.3	76.1	0.305	0.458	0.106	
FnxI	14.6	19.8	1.000	0.000	0.017	<i>Dist – S</i>
GruI	24.9	–25.0	0.675	0.323	0.933	<i>Near – S</i>
GruII	27.6	28.7	0.998	0.001	0.085	<i>Near – S</i>
HerI	37.7	49.5	0.000	1.000	0.050	<i>Near – S, Dist – S</i>
HorI	1.0	10.0	1.000	0.000	0.050	<i>Near – S, Dist – S</i>
HorII	6.2	31.8	0.551	0.201	0.267	
HyaII	28.9	–73.9	0.104	0.443	0.066	<i>Dist – S</i>
HyiI	10.5	18.7	1.000	0.000	0.000	<i>Near – S, Dist – S</i>
LeoI	20.4	54.4	0.196	0.219	0.065	
LeoII	13.4	65.3	0.129	0.189	0.035	<i>Dist – S</i>
LeoIV	2.7	59.9	0.300	0.398	0.189	<i>Dist – S</i>
LeoV	1.1	87.1	0.143	0.269	0.009	<i>Dist – S</i>
PhxII	19.5	20.9	1.000	0.000	0.060	<i>Near – S</i>
PisII	4.6	17.2	0.853	0.081	0.383	<i>Dist – S</i>
RetII	11.9	14.3	1.000	0.000	0.000	<i>Near – S, Dist – S</i>
RetIII	3.0	10.2	0.443	0.549	0.866	<i>Near – S</i>
SciI	47.5	–85.1	0.000	1.000	0.000	<i>Near – S, Dist – S</i>
SegI	5.0	–5.8	1.000	0.000	0.251	
SegII	36.1	–36.2	0.710	0.001	0.077	
SxtI	58.7	–81.9	0.000	1.000	0.000	<i>Near – S, Dist – S</i>
SgrII	14.7	71.8	0.000	0.000	0.000	<i>Near – S, Dist – S</i>
TriII	64.5	–64.9	0.000	1.000	0.053	
TucII	25.3	46.6	0.000	0.001	0.000	<i>Near – S</i>
TucIII	6.1	–46.0	0.280	0.017	0.002	
TucIV	15.4	24.7	1.000	0.000	0.000	<i>Near – S</i>
TucV	20.6	74.3	0.000	0.015	0.000	

**Table 6** *continued on next page*

**Table 6** (*continued*)

Name	$\theta_{\text{VPOS-3}}^{\text{predicted}}$	$\theta_{\text{VPOS-3}}^{\text{measured}}$	$f_{\text{inVPOS}}$	$p_{\text{outsideVPOS}}$	$p_{>\text{obs}}$	volume-complete member
	( $^{\circ}$ )	( $^{\circ}$ )				
(1)	(2)	(3)	(4)	(5)	(6)	(7)
UMaI	36.0	-50.3	0.018	0.537	0.012	<i>Near - S</i>
UMaII	55.4	-76.0	0.000	1.000	0.000	<i>Near - S</i>
UMiI	21.7	25.7	1.000	0.000	0.000	<i>Near - S, Dist - S</i>
Will	39.1	73.1	0.000	1.000	0.000	<i>Near - S</i>

NOTE— Column 1: name of the object, Col. 2: angle between VPOS normal and predicted orbital pole (best-possible alignment), Col. 3: angle between VPOS normal and median measured orbital pole, Col. 4: fraction of realizations whose orbital pole falls into the 10% circles around the VPOS normal vector, Col. 5: probability that an intrinsically perfectly aligned orbital pole is found outside of the 10% circles given the proper motion measurement uncertainties, Col. 6: probability to find an orbital pole at least as far inclined from the VPOS as the median measured orbital pole if the intrinsic alignment is as close as possible, and Col. 7 membership to the nearby volume-complete sub-sample (quoted *Near - S*) up to the 90.5 kpc slice from [Drlica-Wagner et al. 2020](#), or the distant (quoted *Dist - S*) volume complete sub-sample up to the 181 kpc slice.

Of the 46 dwarf galaxies in our sample, six cannot orbit within the VPOS because their positions place them well outside of the structure ( $\theta_{\text{VPOS-3}}^{\text{predicted}} > \theta_{\text{inVPOS}}$ ). These are Hercules I, Segue II, Sagittarius II, Triangulum II, Ursa Major II, and Willman 1. Of the remaining 40 objects for which an alignment is feasible, 20 have median orbital poles that align to better than  $\theta_{\text{inVPOS}}$  with the VPOS normal vector<sup>3</sup>. Most of these are well constrained, with 13 having almost all their Monte Carlo realizations within this angle from the VPOS plane ( $f_{\text{inVPOS}} \geq 0.99$ ). Only one object, RetIII with  $f_{\text{inVPOS}} = 0.44$ , is relatively poorly constrained while the remaining six have  $0.85 > f_{\text{inVPOS}} > 0.55$ . Only three (Grus I, Sculptor, Segue I) of these 20 are counter-orbiting with respect to the bulk orbital sense of the VPOS members. Together with the LMC and SMC, which also orbit along the VPOS, the counter-orbiting fraction is thus  $f_{\text{counter}} = \frac{3}{22} = 0.14$ . This is intriguingly close to the counter-orbiting fraction considering only the 11 bright, classical MW satellites ([Pawlowski & Kroupa 2020](#)), of which 8 have orbital poles aligned with the VPOS with only Sculptor orbiting on the opposite sense than the others ( $f_{\text{counter}} = \frac{1}{8} = 0.13$ ).

The remaining 20 dwarf galaxies have median orbital poles that do not align with the VPOS. Of these, 11 can confidently be ruled out as VPOS members (by our adopted criterion) because their chance of aligning is smaller than  $f_{\text{inVPOS}} < 0.05$ . Note, however, that the orbital poles of Crater II and Draco II are well constrained but only marginally outside of our adopted maximum accepted alignment angle of  $\theta_{\text{inVPOS}}$ , and a small change in the adopted direction of the VPOS normal vector would place both inside of the 10% region.

The remaining nine dwarf galaxies have orbital poles that are only poorly constrained, mainly due to their large distance and thus large relative proper motion error<sup>4</sup>. However, these nine are all well consistent with being aligned with the VPOS within their uncertainties, with  $0.1 < f_{\text{inVPOS}} < 0.4$ .

<sup>3</sup> The dwarf galaxies with median orbital poles aligned with the VPOS are: Canes Venaciti I, Carina I, Carina III, Crater I, Crater II, Draco I, Fornax, Grus I, Grus II, Horologium I, Horologium II, Hydrus, Phoenix II, Pisces II, Reticulum II, Reticulum III, Sculptor, Segue I, Tucana IV, and Ursa Minor.

<sup>4</sup> These dwarf galaxies with orbital poles that are poorly constrained but consistent with VPOS alignment are: Aquarius II, Canes Venaciti I, Eridanus II, Leo I, Leo II, Leo IV, Leo V, Tucana III

This is further demonstrated by their high  $p_{\text{outsideVPOS}}$  values of 19 to 60% (except Tucana III with only  $p_{\text{outsideVPOS}} = 0.02$ ), indicating that even if they were intrinsically well aligned, the substantial proper motion uncertainties would likely place their derived median orbital pole outside of the region around the VPOS normal vector. Six of the nine objects most-likely co-orbit.

The similarity of our study with that of Fritz et al. (2018) in sample and data analysis also allows us to judge how the improved data quality of Gaia eDR3 over DR2 affects the VPOS signal, without the danger of being affected by biases due to differing methodologies. The two studies have 37 objects in common, of which 32 can possibly orbit along the VPOS. Of these 32, we find that 16 have orbital poles that are most-likely aligned with the VPOS normal ( $f_{\text{inVPOS}} > 0.5$ ), and 10 of these are definitively aligned ( $f_{\text{inVPOS}} > 0.98$ ). In Fritz et al. (2018), these numbers were 14 and 7, respectively. Thus, the improved data has resulted in a substantial increase in the number of orbitally aligned dwarf galaxies, as to be expected if an underlying correlation is obscured by initially larger measurement errors. The same trend has previously been found for the classical MW satellites, whose orbital poles clustered subsequently tighter around the VPOS normal direction as their proper motion errors have decreased (Pawlowski & Kroupa 2020). Furthermore, it is noteworthy that  $f_{\text{counter}}$  dropped considerably compared to Fritz et al. (2018), who found  $f_{\text{counter}} = 0.32$  among their 17 likely VPOS members and the Magallanic Clouds. This change is also consistent with the VPOS being an intrinsic, correlated structure exhibiting a preferred orbital direction that is becoming more apparent as proper motion measurements improve.

In summary, of the 40 Milky Way dwarf galaxies that have spatial positions consistent with being members of the VPOS, at least 20 to 29 objects have orbital poles aligned with the VPOS normal vector. Thus, from 50 to 73% of potentially aligned dwarf galaxies (or 43 to 63 % of all) indeed lie and orbit in the VPOS, which confirms the prominence of the VPOS for the MW dwarf galaxy spatial and orbital distributions<sup>5</sup>. Such a large fraction is indicative of a real structure of dwarf galaxies, which could not be associated to the expectations for a cosmological infall of primordial dwarfs, including if they had been accreted along cold streams (Pawlowski & Kroupa 2014, and references therein).

#### 4.3. *Locations of dwarf galaxies are excessively near their orbital pericenter*

Based on Gaia DR2, Fritz et al. (2018) and Simon (2018) noticed that dwarf galaxy locations are excessively concentrated near their pericenters, which is at odd for satellites that are expected to lie mostly near their apocenters. Fritz et al. (2018) argued that this effect may be caused by the existence of non-detected ultra faint dwarfs, mostly those lying beyond 100 kpc. The problem is also less pronounced when considering high MW mass (Fritz et al. 2018; Hammer et al. 2020, see their Figure 5 and Figure 1, respectively). If persistent, the dwarf galaxy excess near pericenter may challenge their commonly assumed nature as long-lived MW satellites since such a realization would be associated to a very small probabilities especially for moderate MW mass (Hammer et al. 2020,  $P \sim 2 \times 10^{-7}$  for MW mass  $\leq 10^{12} M_{\odot}$ ).

Here we try to reevaluate these statistics using Gaia EDR3 data and by accounting for possible biases linked to the observability of dwarf galaxies at larger distances. Indeed, one may consider that

<sup>5</sup> This fraction can be considered a lower limit: For Leo II, one of the most distant dwarf galaxies in our sample that has a poorly constrained orbital pole, more accurate Hubble Space Telescope proper motion (Piatek et al. 2016) place its orbital pole firmly in the VPOS (Pawlowski & Kroupa 2020). Furthermore, discussed above Crater II and Draco II are just marginally outside of the adopted VPOS alignment criterion, and both the LMC and SMC are known to co-orbit along the VPOS (though Sagittarius does not).



we are only able to see the closest dwarf galaxies, which then could be those close to their pericenters. Following [Hammer et al. \(2020\)](#) we first define the orbital phase probability:

$$P = \frac{t_{\text{peri}}}{t_{\text{peri-apo}}}, \quad (8)$$


to estimate the chance for a dwarf galaxy to be close to its pericenter. In Equation 8,  $t_{\text{peri}}$  is the orbital time for reaching or leaving the pericenter from the dwarf galaxy's current position (whichever is shorter), and  $t_{\text{peri-apo}}$  is the time to complete half an orbit. We have used the calculated dwarf orbits from the publicly available code *galpy* ([Bovy 2015](#)) for deriving the time spent to reach the pericenter, as well as that from apocenter to pericenter. We further limit the volume and distance to 300 kpc because beyond this, most dwarf galaxies would have been hard to be detected ([Simon 2019](#)). It leads to change  $t_{\text{peri-apo}}$  in Equation 8 by the time taken from pericenter to  $\min(\text{apocenter}, 300 \text{ kpc})$ :

$$P = \frac{t_{\text{peri}}}{t_{\text{peri-min(apo,300kpc)}}}, \quad (9)$$

In the following, we examine the properties of complete subsamples using the probabilities based on Equation 9. Non-parametric Kolmogorov-Smirnov tests presented in Figure 3 are made by comparing the cumulative probabilities calculated from Equation 9 to those expected in the null hypothesis, which represents a random distribution of time between  $t_{\text{peri}}$  and  $t_{\text{peri-min(apo,300kpc)}}$ .

The proximity of dwarf galaxies to their pericenter could be caused by biases, i.e., that we preferentially detect nearby dwarf galaxies more likely to be close to their pericenters, and this at the cost of missing a population of faint dwarfs that would lie at larger distance, e.g., beyond 100 kpc ([Fritz et al. 2018](#)). To test the possibility of missing non-detected ultra faint dwarfs, we have built 'volume-complete samples' of dwarf galaxies, i.e., samples including only the dwarfs that could be detected within a given volume, independently of their actual distances. We have made use of the detectability study of [Drlica-Wagner et al. \(2020\)](#) and have applied their Equation 2 for estimating the  $(M_V, r_{\text{half}})$  range to which dwarf galaxies can be observed. It leads to volume-complete samples of 26 (23) dwarf galaxies that can be observed up to the whole volume defined by the distance slices of 90.5 (181) kpc, respectively (see [Drlica-Wagner et al. 2020's](#) Table 5 and Fig. 6). Membership to the two sub-samples are tabulated in the last column of Table 6, and V-absolute magnitude and  $r_{\text{half}}$  values used to define the dwarf galaxy detectability have been taken from [Simon \(2019\)](#).

Figure 3 presents the the cumulative probability of having an excess of dwarf galaxy locations near the pericenter, assuming they are satellites of the MW modeled with *PNFW* (green curve, see Table 3) and with the *PE<sub>HM</sub>* (red curve, highest mass). As expected (see also [Fritz et al. 2018](#)), the higher the MW mass, the more likely is the satellite hypothesis for dwarf galaxies as it is illustrated in Figure 1. Gaia EDR3 provides sufficiently accurate orbits and pericenter to gather samples of 26 and 23 dwarf galaxies that are essentially complete in the sense that all included dwarf galaxies can be seen in the whole considered small ( $\sim 100$  kpc) and large ( $\sim 200$  kpc) volume, respectively. Furthermore, this allows us to test the hypothesis of missing ultra-faint dwarfs that would lie at apocenter and then that have supposedly not been detected yet ([Fritz et al. 2018](#)).



**Figure 3.** Cumulative distribution of time to reach the pericenter divided by the time taken from pericenter to min(apocenter, 300 kpc). The black solid line shows the null hypothesis, i.e., dwarf galaxies have location randomly distributed from pericenter to min(apocenter, 300 kpc). Green and red lines represent the  $PNFW$  model ( $M_{tot} = 8.1 \times 10^{11} M_{\odot}$ , see also Eilers et al. 2019, and  $PE_{HM}$  the most massive mass able to fit the MW rotation curve ( $M_{tot} = 15 \times 10^{11} M_{\odot}$ , see Jiao et al. 2021), respectively.  $D_{max}$  values and associated probabilities of the Kolmogorov Smirnov tests are given in the Figures, top and bottom panels representing the 90.5 (dubbed as small volume) and the 181 kpc (dubbed as large volume) distance slices of Drlica-Wagner et al. 2020, respectively.

Let us consider the small volume-complete sample (top panel of Figure 3) of 26 dwarf galaxies, which leads to very small probabilities. Examining the apocenters of these dwarf galaxies and considering the  $PNFW$  MW mass model, we find that only 34% of them are within the small volume, i.e., letting open the possibility that there are undetected missing dwarf galaxies beyond  $\sim 100$  kpc. However, for the most massive MW mass model ( $PE_{HM}$ ), most (58%) apocenters lie within the small volume, and it becomes non plausible that the small probabilities ( $P = 9.3 \times 10^{-4}$ ) can be due to missing ultra faint dwarfs. Now let consider the large volume-complete sample of 23 dwarf galaxies, which is undoubtedly more affected by the difficulty in detecting distant faint dwarfs. Assuming the  $PNFW$  MW mass model, 66% of apocenters are within that volume, which means that there could be only few missing dwarfs lying further away near their apocenter to explain the small probability

given in the bottom panel of Figure 3. At first glance, the probability for the most massive MW mass model ( $PE_{\text{HM}}$ ) could be interpreted as a significant alleviation of the proximity-to-pericenter problem. However, in such a case, 92% of the apocenters are within the large volume, letting no possibility for missing dwarfs at larger distances.

In summary, we have examined the the proximity-to-pericenter problem that may affect the satellite nature of most dwarf galaxies. For this, we have only considered the two most massive MW mass models of this paper, because they are the ones for which most dwarf galaxies are bound. Having defined volume-complete samples following the detectability procedure of [Drlica-Wagner et al. \(2020\)](#), we find that by increasing the MW mass, and by assuming a putative population of distant faint dwarfs may alleviate part of the problem, but certainly not completely. For example, assuming the largest MW mass available to fit the MW rotation curve, the probability that MW dwarf galaxies behave as satellites is well below 1 in the two considered complete volumes of the above analysis.

## 5. CONCLUSION

We have determined the proper motions of 46 dwarf galaxies using Gaia EDR3, with a robust evaluation of errors accounting both for effects of their statistics and of Gaia systematics. The gain compared to former DR2 analysis by [Fritz et al. \(2018\)](#) is twofold. First, the accuracy for objects in common between the two studies has improved by a factor  $\sim 2.5$ , which corresponds to the actual reduction of total errors for the analysis of their tangential and 3D velocities. Second, it allows to increase to 40 the number of dwarf galaxies for which analysis of their motions can be robustly made, i.e., more than twice what was done with Gaia DR2.

We have then derived the 3D phase-space diagram of dwarf galaxies that illustrates the sequence delineated by most dwarf galaxies. Gaia EDR3 errors on 3D velocities are systematically smaller than differences between expectations from various possible MW mass models. It implies that almost all dwarf galaxies are gravitationally bound if the MW total mass is larger than  $8 \times 10^{11} M_{\odot}$ , while they would be increasingly unbound for MW mass values going down to  $5.1 \times 10^{11} M_{\odot}$  and  $2.8 \times 10^{11} M_{\odot}$ .

In this paper, we have incorporated calculations of integrated orbital parameters after considering the whole range of four MW mass models that can be consistent with the MW rotation curves ([Eilers et al. 2019](#); [Mróz et al. 2019](#)). It provides a useful library of orbits with different schemes for the MW mass, some of them could be either rosette or hyperbolic. While apocenters and eccentricities are very dependent on the adopted MW potential, we show that pericenter values are very robustly determined, whatever the MW mass model, and down to an accuracy of 10% for a majority of dwarf galaxies.

We also confirm that many of the dwarf galaxies lie near their pericenters, which cannot simply be due to either selection effects or to an underestimate of the MW mass. This later result appears to be problematic if most dwarf galaxies are MW satellites. Finally we identify the strong prominence of the VPOS in the distribution of orbital poles, that includes a majority of dwarf galaxy locations and orbits, confirming it as an important feature of the outer Galactic halo. Such a large fraction of VPOS members and their strong kinematic correlation is indicative of a real structure of dwarf galaxies, which could be in conflict with expectations for a cosmological infall of primordial dwarfs.

## ACKNOWLEDGMENTS

This work has made use of data from the European Space Agency (ESA) mission *Gaia* (<https://www.cosmos.esa.int/gaia>), processed by the *Gaia* Data Processing and Analysis Consortium (DPAC, <https://www.cosmos.esa.int/web/gaia/dpac/consortium>).

Funding for the DPAC has been provided by national institutions, in particular the institutions participating in the *Gaia* Multilateral Agreement. This work has been supported by the National Natural Foundation of China (NSFC No. 11973042 and No. 11973052). It was also supported by the Fundamental Research Funds for the Central Universities and the China Scholarship Council (CSC). We also thank for its support the International Research Program Tianguan, which is an agreement between the CNRS, NAOC and the Yunnan University. MSP thanks the Klaus Tschira Stiftung gGmbH and German Scholars Organization e.V. for support via a Klaus Tschira Boost Fund.

## APPENDIX


### A. COMPARISON OF PMS WITH OTHER PAPERS.

McConnachie & Venn (2020) published PM based on Gaia EDR3 for a large number of dwarf galaxies. Figure A1 compares values of Table 1 to theirs. Although values are often consistent within error bars found in this paper, those tabulated by McConnachie & Venn (2020) appear to be extremely small. This is because they do not account for Gaia systematics as we have made in using Vasiliev (2019, see also Section 2.2 of this paper), though mentioning their impact in the text. We also notice a good agreement within our uncertainties between our values for 9 dwarfs that have been studied by Vitral (2021), though our uncertainties are larger and there is a small,  $< 2\sigma$  discrepancy for Bootes I.

Figure A2 compares PM values and their errors from EDR3 to those from Gaia DR2 from Fritz et al. (2018).

## REFERENCES

- Adén, D., Feltzing, S., Koch, A., et al. 2009, A&A, 506, 1147, doi: [10.1051/0004-6361/200912718](https://doi.org/10.1051/0004-6361/200912718)
- Aparicio, A., Carrera, R., & Martínez-Delgado, D. 2001, AJ, 122, 2524, doi: [10.1086/323535](https://doi.org/10.1086/323535)
- Armandroff, T. E., Olszewski, E. W., & Pryor, C. 1995, AJ, 110, 2131, doi: [10.1086/117675](https://doi.org/10.1086/117675)
- Battaglia, G., Tolstoy, E., Helmi, A., et al. 2011, MNRAS, 411, 1013, doi: [10.1111/j.1365-2966.2010.17745.x](https://doi.org/10.1111/j.1365-2966.2010.17745.x)
- . 2006, A&A, 459, 423, doi: [10.1051/0004-6361:20065720](https://doi.org/10.1051/0004-6361:20065720)
- Besla, G., Kallivayalil, N., Hernquist, L., et al. 2012, MNRAS, 421, 2109, doi: [10.1111/j.1365-2966.2012.20466.x](https://doi.org/10.1111/j.1365-2966.2012.20466.x)
- Binney, J., & Tremaine, S. 1987, Princeton, NJ, Princeton University Press, 1987, 747 p.
- Bland-Hawthorn, J., & Gerhard, O. 2016, ARA&A, 54, 529, doi: [10.1146/annurev-astro-081915-023441](https://doi.org/10.1146/annurev-astro-081915-023441)
- Bovy, J. 2015, ApJS, 216, 29, doi: [10.1088/0067-0049/216/2/29](https://doi.org/10.1088/0067-0049/216/2/29)
- Caldwell, N., Walker, M. G., Mateo, M., et al. 2017, ApJ, 839, 20, doi: [10.3847/1538-4357/aa688e](https://doi.org/10.3847/1538-4357/aa688e)
- Cautun, M., Benítez-Llambay, A., Deason, A. J., et al. 2020, MNRAS, 494, 4291, doi: [10.1093/mnras/staa1017](https://doi.org/10.1093/mnras/staa1017)
- de Boer, T. J. L., Tolstoy, E., Hill, V., et al. 2012, A&A, 544, A73, doi: [10.1051/0004-6361/201219547](https://doi.org/10.1051/0004-6361/201219547)
- de Salas, P. F., Malhan, K., Freese, K., Hattori, K., & Valluri, M. 2019, JCAP, 10, 037, doi: [10.1088/1475-7516/2019/10/037](https://doi.org/10.1088/1475-7516/2019/10/037)



**Figure A1.** Comparison of proper motions (left: RA, right: DEC) of Table 1 with those calculated by McConnachie & Venn (2020). Few dwarf galaxies names are indicated, pointing out to dwarf galaxies for which the two measurements are discrepant at  $> 1 \sigma$  level. The top panels indicate the ratio of error from this paper to that tabulated by McConnachie & Venn (2020) both extracted from Gaia EDR3. Notice that the error on PM RA of Fornax is so small ( $1 \mu\text{s yr}^{-1}$ ) in McConnachie & Venn (2020) that it leads to a ratio in excess of 20 (not shown in the Figure).

Drlica-Wagner, A., Bechtol, K., Mau, S., et al. 2020, *ApJ*, 893, 47,

doi: [10.3847/1538-4357/ab7eb9](https://doi.org/10.3847/1538-4357/ab7eb9)

Eilers, A.-C., Hogg, D. W., Rix, H.-W., & Ness, M. K. 2019, *ApJ*, 871, 120,

doi: [10.3847/1538-4357/aaf648](https://doi.org/10.3847/1538-4357/aaf648)

Erkal, D., Belokurov, V., Laporte, C. F. P., et al. 2019, *MNRAS*, 487, 2685,

doi: [10.1093/mnras/stz1371](https://doi.org/10.1093/mnras/stz1371)

Fox, A. J., Wakker, B. P., Barger, K. A., et al. 2014, *ApJ*, 787, 147,

doi: [10.1088/0004-637X/787/2/147](https://doi.org/10.1088/0004-637X/787/2/147)

Fritz, T. K., Battaglia, G., Pawlowski, M. S., et al. 2018, *A&A*, 619, A103,

doi: [10.1051/0004-6361/201833343](https://doi.org/10.1051/0004-6361/201833343)

Fritz, T. K., Carrera, R., Battaglia, G., & Taibi, S. 2019, *A&A*, 623, A129,

doi: [10.1051/0004-6361/201833458](https://doi.org/10.1051/0004-6361/201833458)

Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2020, arXiv e-prints, arXiv:2012.01533.

<https://arxiv.org/abs/2012.01533>

Gaia Collaboration, Prusti, T., de Bruijne, J. H. J., et al. 2016, *A&A*, 595, A1,

doi: [10.1051/0004-6361/201629272](https://doi.org/10.1051/0004-6361/201629272)

Gravity Collaboration, Abuter, R., Amorim, A., et al. 2018, *A&A*, 615, L15,

doi: [10.1051/0004-6361/201833718](https://doi.org/10.1051/0004-6361/201833718)

Hammer, F., Wang, J., Pawlowski, M. S., et al. 2021, submitted to *ApJ*

Hammer, F., Yang, Y., Arenou, F., et al. 2020, *The Astrophysical Journal*, 892, 3,

doi: [10.3847/1538-4357/ab77be](https://doi.org/10.3847/1538-4357/ab77be)


Hammer, F., Yang, Y., Wang, J., et al. 2019, *ApJ*, 883, 171, doi: [10.3847/1538-4357/ab36b6](https://doi.org/10.3847/1538-4357/ab36b6)

Hammer, F., Yang, Y. B., Flores, H., Puech, M., & Fouquet, S. 2015, *ApJ*, 813, 110,

doi: [10.1088/0004-637X/813/2/110](https://doi.org/10.1088/0004-637X/813/2/110)

Helmi, A., van Leeuwen, F., McMillan, P. J., et al. 2018, *A&A*, 616, A12,

doi: [10.1051/0004-6361/201832698](https://doi.org/10.1051/0004-6361/201832698)



**Figure A2.** Comparison of proper motions (left: RA, right: DEC) of Table 1 with those calculated by Fritz et al. (2018) with Gaia DR2. The top panels indicate the ratio of error from Fritz et al. (2018) Gaia DR2 to that of this paper, Gaia EDR3.

- Hill, V., Skúladóttir, Á., Tolstoy, E., et al. 2019, *A&A*, 626, A15, doi: [10.1051/0004-6361/201833950](https://doi.org/10.1051/0004-6361/201833950)
- Jenkins, S., Li, T. S., Pace, A. B., et al. 2020, arXiv e-prints, arXiv:2101.00013, <https://arxiv.org/abs/2101.00013>
- Jiao, Y., Hammer, F., Wang, J., & Yang, Y. 2021, arXiv e-prints, accepted in *Astronomy & Astrophysics*, arXiv:2107.00014
- Johnson, D. R. H., & Soderblom, D. R. 1987, *AJ*, 93, 864, doi: [10.1086/114370](https://doi.org/10.1086/114370)
- Jurić, M., Ivezić, Ž., Brooks, A., et al. 2008, *ApJ*, 673, 864, doi: [10.1086/523619](https://doi.org/10.1086/523619)
- Kacharov, N., Battaglia, G., Rejkuba, M., et al. 2017, *MNRAS*, 466, 2006, doi: [10.1093/mnras/stw3188](https://doi.org/10.1093/mnras/stw3188)
- Kallivayalil, N., van der Marel, R. P., Besla, G., Anderson, J., & Alcock, C. 2013, *ApJ*, 764, 161, doi: [10.1088/0004-637X/764/2/161](https://doi.org/10.1088/0004-637X/764/2/161)
- Karukes, E. V., Benito, M., Iocco, F., Trotta, R., & Geringer-Sameth, A. 2020, *JCAP*, 05, 033, doi: [10.1088/1475-7516/2020/05/033](https://doi.org/10.1088/1475-7516/2020/05/033)
- Kirby, E. N., Boylan-Kolchin, M., Cohen, J. G., et al. 2013, *ApJ*, 770, 16, doi: [10.1088/0004-637X/770/1/16](https://doi.org/10.1088/0004-637X/770/1/16)
- Kirby, E. N., Cohen, J. G., Simon, J. D., et al. 2017, *ApJ*, 838, 83, doi: [10.3847/1538-4357/aa6570](https://doi.org/10.3847/1538-4357/aa6570)
- Kirby, E. N., Simon, J. D., & Cohen, J. G. 2015, *ApJ*, 810, 56, doi: [10.1088/0004-637X/810/1/56](https://doi.org/10.1088/0004-637X/810/1/56)
- Kleyna, J., Wilkinson, M. I., Evans, N. W., Gilmore, G., & Frayn, C. 2002, *MNRAS*, 330, 792, doi: [10.1046/j.1365-8711.2002.05155.x](https://doi.org/10.1046/j.1365-8711.2002.05155.x)
- Koch, A., Kleyna, J. T., Wilkinson, M. I., et al. 2007, *AJ*, 134, 566, doi: [10.1086/519380](https://doi.org/10.1086/519380)
- Koch, A., Wilkinson, M. I., Kleyna, J. T., et al. 2009, *ApJ*, 690, 453, doi: [10.1088/0004-637X/690/1/453](https://doi.org/10.1088/0004-637X/690/1/453)
- Koposov, S. E., Gilmore, G., Walker, M. G., et al. 2011, *ApJ*, 736, 146, doi: [10.1088/0004-637X/736/2/146](https://doi.org/10.1088/0004-637X/736/2/146)
- Koposov, S. E., Casey, A. R., Belokurov, V., et al. 2015, *ApJ*, 811, 62, doi: [10.1088/0004-637X/811/1/62](https://doi.org/10.1088/0004-637X/811/1/62)



- Koposov, S. E., Walker, M. G., Belokurov, V., et al. 2018, *MNRAS*, 479, 5343, doi: [10.1093/mnras/sty1772](https://doi.org/10.1093/mnras/sty1772)
- Li, T. S., Simon, J. D., Drlica-Wagner, A., et al. 2017, *ApJ*, 838, 8, doi: [10.3847/1538-4357/aa6113](https://doi.org/10.3847/1538-4357/aa6113)
- Li, T. S., Simon, J. D., Kuehn, K., et al. 2018, *ApJ*, 866, 22, doi: [10.3847/1538-4357/aadf91](https://doi.org/10.3847/1538-4357/aadf91)
- Lindegren, L., Klioner, S. A., Hernández, J., et al. 2020, arXiv e-prints, arXiv:2012.03380
- Longeard, N., Martin, N., Starkenburg, E., et al. 2020, *MNRAS*, 491, 356, doi: [10.1093/mnras/stz2854](https://doi.org/10.1093/mnras/stz2854)
- Martin, N. F., Ibata, R. A., Chapman, S. C., Irwin, M., & Lewis, G. F. 2007, *MNRAS*, 380, 281, doi: [10.1111/j.1365-2966.2007.12055.x](https://doi.org/10.1111/j.1365-2966.2007.12055.x)
- Martin, N. F., Geha, M., Ibata, R. A., et al. 2016, *MNRAS*, 458, L59, doi: [10.1093/mnrasl/slw013](https://doi.org/10.1093/mnrasl/slw013)
- Mateo, M., Olszewski, E. W., & Walker, M. G. 2008, *ApJ*, 675, 201, doi: [10.1086/522326](https://doi.org/10.1086/522326)
- McConnachie, A. W., & Venn, K. A. 2020, arXiv e-prints, arXiv:2012.03904
- Miyamoto, M., & Nagai, R. 1975, *Publications of the Astronomical Society of Japan*, 27, 533
- Mróz, P., Udalski, A., Skowron, D. M., et al. 2019, *ApJL*, 870, L10, doi: [10.3847/2041-8213/aaf73f](https://doi.org/10.3847/2041-8213/aaf73f)
- Muñoz, R. R., Majewski, S. R., Zaggia, S., et al. 2006, *ApJ*, 649, 201, doi: [10.1086/505620](https://doi.org/10.1086/505620)
- Navarro, J. F., Frenk, C. S., & White, S. D. M. 1997, *ApJ*, 490, 493, doi: [10.1086/304888](https://doi.org/10.1086/304888)
- Pace, A. B., Kaplinghat, M., Kirby, E., et al. 2020, *MNRAS*, 495, 3022, doi: [10.1093/mnras/staa1419](https://doi.org/10.1093/mnras/staa1419)
- Patel, E., Kallivayalil, N., Garavito-Camargo, N., et al. 2020, *ApJ*, 893, 121, doi: [10.3847/1538-4357/ab7b75](https://doi.org/10.3847/1538-4357/ab7b75)
- Pawlowski, M. S., & Kroupa, P. 2013, *MNRAS*, 435, 2116, doi: [10.1093/mnras/stt1429](https://doi.org/10.1093/mnras/stt1429)
- Pawlowski, M. S., & Kroupa, P. 2014, *ApJ*, 790, 74, doi: [10.1088/0004-637X/790/1/74](https://doi.org/10.1088/0004-637X/790/1/74)
- Pawlowski, M. S., & Kroupa, P. 2020, *MNRAS*, 491, 3042, doi: [10.1093/mnras/stz3163](https://doi.org/10.1093/mnras/stz3163)
- Pawlowski, M. S., McGaugh, S. S., & Jerjen, H. 2015, *MNRAS*, 453, 1047, doi: [10.1093/mnras/stv1588](https://doi.org/10.1093/mnras/stv1588)
- Piatek, S., Pryor, C., & Olszewski, E. W. 2016, *AJ*, 152, 166, doi: [10.3847/0004-6256/152/6/166](https://doi.org/10.3847/0004-6256/152/6/166)
- Pouliasis, E., Matteo, P. D., & Haywood, M. 2017, *A&A*, 598, A66, doi: [10.1051/0004-6361/201527346](https://doi.org/10.1051/0004-6361/201527346)
- Retana-Montenegro, E., van Hese, E., Gentile, G., Baes, M., & Frutos-Alfaro, F. 2012, *A&A*, 540, A70, doi: [10.1051/0004-6361/201118543](https://doi.org/10.1051/0004-6361/201118543)
- Ripepi, V., Cioni, M.-R. L., Moretti, M. I., et al. 2017, *MNRAS*, 472, 808, doi: [10.1093/mnras/stx2096](https://doi.org/10.1093/mnras/stx2096)
- Savino, A., Salaris, M., & Tolstoy, E. 2015, *A&A*, 583, A126, doi: [10.1051/0004-6361/201527072](https://doi.org/10.1051/0004-6361/201527072)
- Simon, J. D. 2018, *ApJ*, 863, 89, doi: [10.3847/1538-4357/aacdfb](https://doi.org/10.3847/1538-4357/aacdfb)
- Simon, J. D. 2019, *Annual Review of Astronomy and Astrophysics*, 57, 375, doi: [10.1146/annurev-astro-091918-104453](https://doi.org/10.1146/annurev-astro-091918-104453)
- Simon, J. D., & Geha, M. 2007, *ApJ*, 670, 313, doi: [10.1086/521816](https://doi.org/10.1086/521816)
- Simon, J. D., Geha, M., Minor, Q. E., et al. 2011, *ApJ*, 733, 46, doi: [10.1088/0004-637X/733/1/46](https://doi.org/10.1088/0004-637X/733/1/46)
- Simon, J. D., Drlica-Wagner, A., Li, T. S., et al. 2015, *ApJ*, 808, 95, doi: [10.1088/0004-637X/808/1/95](https://doi.org/10.1088/0004-637X/808/1/95)
- Simon, J. D., Li, T. S., Drlica-Wagner, A., et al. 2017, *ApJ*, 838, 11, doi: [10.3847/1538-4357/aa5be7](https://doi.org/10.3847/1538-4357/aa5be7)
- Simon, J. D., Li, T. S., Erkal, D., et al. 2020, *ApJ*, 892, 137, doi: [10.3847/1538-4357/ab7ccb](https://doi.org/10.3847/1538-4357/ab7ccb)
- Sohn, S. T., Majewski, S. R., Muñoz, R. R., et al. 2007, *ApJ*, 663, 960, doi: [10.1086/518302](https://doi.org/10.1086/518302)
- Spencer, M. E., Mateo, M., Walker, M. G., & Olszewski, E. W. 2017, *ApJ*, 836, 202, doi: [10.3847/1538-4357/836/2/202](https://doi.org/10.3847/1538-4357/836/2/202)
- Tolstoy, E., Hill, V., & Tosi, M. 2009, *ARA&A*, 47, 371, doi: [10.1146/annurev-astro-082708-101650](https://doi.org/10.1146/annurev-astro-082708-101650)
- Tolstoy, E., Irwin, M. J., Helmi, A., et al. 2004, *ApJL*, 617, L119, doi: [10.1086/427388](https://doi.org/10.1086/427388)
- Torrealba, G., Koposov, S. E., Belokurov, V., et al. 2016, *MNRAS*, 463, 712, doi: [10.1093/mnras/stw2051](https://doi.org/10.1093/mnras/stw2051)
- Torrealba, G., Belokurov, V., Koposov, S. E., et al. 2019, *MNRAS*, 488, 2743, doi: [10.1093/mnras/stz1624](https://doi.org/10.1093/mnras/stz1624)
- Torrealba, G., Belokurov, V., Koposov, S. E., et al. 2019, *MNRAS*, 488, 2743, doi: [10.1093/mnras/stz1624](https://doi.org/10.1093/mnras/stz1624)
- Ural, U., Wilkinson, M. I., Koch, A., et al. 2010, *MNRAS*, 402, 1357, doi: [10.1111/j.1365-2966.2009.15975.x](https://doi.org/10.1111/j.1365-2966.2009.15975.x)
- van der Marel, R. P., & Guhathakurta, P. 2008, *ApJ*, 678, 187, doi: [10.1086/533430](https://doi.org/10.1086/533430)

- Vasiliev, E. 2019, MNRAS, 489, 623,  
doi: [10.1093/mnras/stz2100](https://doi.org/10.1093/mnras/stz2100)
- Vitral, E. 2021, arXiv e-prints, arXiv:2102.04841.  
<https://arxiv.org/abs/2102.04841>
- Voggel, K., Hilker, M., Baumgardt, H., et al. 2016,  
MNRAS, 460, 3384,  
doi: [10.1093/mnras/stw1132](https://doi.org/10.1093/mnras/stw1132)
- Walker, M. G., Belokurov, V., Evans, N. W., et al.  
2009a, ApJL, 694, L144,  
doi: [10.1088/0004-637X/694/2/L144](https://doi.org/10.1088/0004-637X/694/2/L144)
- Walker, M. G., Mateo, M., & Olszewski, E. W.  
2009b, AJ, 137, 3100,  
doi: [10.1088/0004-6256/137/2/3100](https://doi.org/10.1088/0004-6256/137/2/3100)
- Walker, M. G., Olszewski, E. W., & Mateo, M.  
2015, MNRAS, 448, 2717,  
doi: [10.1093/mnras/stv099](https://doi.org/10.1093/mnras/stv099)
- Walker, M. G., Mateo, M., Olszewski, E. W.,  
et al. 2016, ApJ, 819, 53,  
doi: [10.3847/0004-637X/819/1/53](https://doi.org/10.3847/0004-637X/819/1/53)
- Wang, J., & Hammer, F. 2021, to be submitted
- Wang, J., Hammer, F., Yang, Y., et al. 2019,  
MNRAS, 486, 5907, doi: [10.1093/mnras/stz1274](https://doi.org/10.1093/mnras/stz1274)