



HAL
open science

Exploring the contribution of the mitochondrial disulfide relay system to Parkinson's disease: the PINK1/CHCHD4 interplay

Giuseppe Arena, Nazanine Modjtahedi, Rejko Kruger

► To cite this version:

Giuseppe Arena, Nazanine Modjtahedi, Rejko Kruger. Exploring the contribution of the mitochondrial disulfide relay system to Parkinson's disease: the PINK1/CHCHD4 interplay. *Neural Regeneration Research*, 2021, 16 (11), pp.2222. 10.4103/1673-5374.310679 . hal-03211905

HAL Id: hal-03211905

<https://hal.science/hal-03211905>

Submitted on 10 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Neural Regeneration Research

Exploring the contribution of the mitochondrial disulfide relay system to Parkinson's disease: the PINK1/CHCHD4 interplay --Manuscript Draft--

Manuscript Number:	
Full Title:	Exploring the contribution of the mitochondrial disulfide relay system to Parkinson's disease: the PINK1/CHCHD4 interplay
Article Type:	Invited Paper (Only solicited by the editor)
Section/Category:	Neurodegenerative Disease and Neural Regeneration
Corresponding Author:	Giuseppe Arena University of Luxembourg: Universite du Luxembourg Esch-sur-Alzette, Esch-sur-Alzette LUXEMBOURG
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	University of Luxembourg: Universite du Luxembourg
Corresponding Author's Secondary Institution:	
First Author:	Giuseppe Arena
First Author Secondary Information:	
Order of Authors:	Giuseppe Arena Nazanine Modjtahedi Rejko Kruger
Order of Authors Secondary Information:	
Suggested Reviewers:	Nobutaka Hattori Juntendo University School of Medicine, Tokyo, Japan nhattori@juntendo.ac.jp Matthew J Farrer University of Florida McKnight Brain Institute m.farrer@ufl.edu

Exploring the contribution of the mitochondrial disulfide relay system to Parkinson's disease: the PINK1/CHCHD4 interplay

Parkinson's disease (PD) is a common movement disorder of the elderly caused by the degeneration of dopaminergic neurons in the *substantia nigra pars compacta* of the brain.

Both environmental and genetic factors pointed out mitochondrial dysfunction as a major cause of neurodegeneration in PD. Pioneering studies using mitochondrial toxins revealed their ability to trigger dopaminergic cell death and irreversible parkinsonism in different animal models (1).

Typical features of mitochondrial dysfunction have been also observed in the human brain of idiopathic PD cases, showing alterations of respiratory chain complex I and IV activity, accumulation of mtDNA deletions and increased oxidative stress (2). Moreover, a number of genes found mutated in familial PD forms encode for proteins involved in the maintenance of mitochondrial homeostasis and quality control. Among these, the PINK1 gene encodes a mitochondrial serine/threonine kinase implicated in key neuroprotective functions, including mitophagy, regulation of mitochondrial transport, control of the mitochondria/endoplasmic reticulum crosstalk and calcium homeostasis (3).

Like the majority of nuclear-encoded mitochondrial proteins containing an N-terminal mitochondrial localization signal (MLS), PINK1 is actively transported into mitochondria through the TOM/TIM complex (translocases of the outer and inner mitochondrial membranes, respectively), driven by the electrical potential ($\Delta\psi$) across the inner mitochondrial membrane (IMM). Once the N-terminal domain of PINK1 reaches the mitochondrial matrix, the full-length 63kDa protein is first cleaved by the MPP α/β protease and then by the IMM-resident PGAM5-associated rhomboid-like (PARL) protease, generating an unstable 52kDa product that is retro-translocated to the cytosol and finally degraded by the proteasome via the N-end rule pathway.

Other proteases embedded in the IMM, such as YME1L and AFG3L2, have been reported to

1
2
3
4 cooperate with PARL in order to fulfill the processing of PINK1 in healthy mitochondria.
5
6 Conversely, PINK1 cleavage is impaired upon dissipation of the mitochondrial membrane
7 potential ($\Delta\psi_m$) by oxidative phosphorylation uncoupling agents, which lead to the accumulation
8
9 of full-length PINK1 on the outer mitochondrial membrane (OMM). This induces the translocation
10
11 of the ubiquitin-ligase Parkin from the cytosol to the surface of depolarized organelles, thus
12
13 activating a complex signaling cascade that (i) excludes damaged mitochondria from the network
14
15 by preventing their movement and (ii) ensures their selective elimination through the autophagy
16
17 pathway (4).
18
19
20

21
22 It is worth noting that not only the collapse of $\Delta\psi_m$, but also generation of mitochondrial reactive
23
24 oxygen species (ROS) or even the accumulation of misfolded proteins in the mitochondrial matrix
25
26 have been previously described as potential inducers of PINK1-dependent mitophagy. Further
27
28 expanding the plethora of stress conditions and signals triggering this pathway, Gao and
29
30 colleagues recently found that the redox-regulated CHCHD4/GFER mitochondrial import pathway
31
32 was required for PINK1 stabilization and mitophagy execution (5). This evolutionary conserved
33
34 disulfide relay system, mammalian homologue of the yeast Mia40/Erv1 pathway, operates in the
35
36 mitochondrial intermembrane space (IMS) to regulate the import and/or proper folding of a set of
37
38 substrates carrying specific cysteine motifs; the CHCHD4 protein is the core component of this
39
40 machinery that catalyzes the oxidation of the cysteine residues to disulfide bridges. Notably, many
41
42 of these substrates participate in potentially relevant pro-survival and neuroprotective activities,
43
44 including biogenesis and assembly of the respiratory chain complexes, regulation of mitochondrial
45
46 ultrastructure and dynamics, general mitochondrial import control, calcium storage, lipid
47
48 homeostasis and mitochondrial protein translation (6). In this light, impairment of the CHCHD4-
49
50 dependent pathway would have profound effects on mitochondrial function and neuronal
51
52 metabolism, thus representing a candidate mechanism contributing to neurodegeneration in PD.
53
54 In accordance with this hypothesis, heterozygous mutations in the CHCHD4/Mia40 substrates
55
56
57
58 CHCHD2/MIX17A and CHCHD10/MIX17B have been detected in distinct families affected by
59
60
61
62
63
64
65

1
2
3
4 late-onset autosomal dominant PD as well as in sporadic PD patients (6). Both CHCHD2 and
5
6 CHCHD10 mutations are suggested to disrupt the integrity of respiratory complexes by impinging
7
8 on the mitochondrial contact site and cristae organization system (MICOS). Of note, CHCHD2
9
10 and CHCHD10 are binding partners and recent studies suggest that mitochondrial dysfunction
11
12 caused by PD-associated mutations of CHCHD2 could implicate the perturbation of the physical
13
14 and functional interaction with CHCHD10 (6). Deficiency in the unique *Drosophila* orthologue of
15
16 CHCHD2/CHCHD10 results in loss of dopaminergic neurons and motor dysfunction, a phenotype
17
18 likely due to alteration of cristae ultrastructure, which in turn leads to impaired mitochondrial
19
20 respiration and increased oxidative stress. In line with this, mitochondrial phenotypes observed in
21
22 mutant flies are rescued by overexpressing the wild-type human CHCHD2, but not its PD-
23
24 associated mutant variants (7).
25
26
27

28
29 CHCHD2 and CHCHD10 are prototypes of proteins that carry simultaneously an N-terminal MLS
30
31 and a coiled-coil-helix-coiled-coil-helix (CHCH) domain, characterized by four cysteine residues
32
33 that form two disulfide bonds. Interestingly, a recent study of CHCHD10 indicated that the CHCH
34
35 domain of the protein was required for its CHCHD4-mediated import to the IMS, rather than its N-
36
37 terminal positively charged MLS (7). A similar mechanism could be also responsible for PINK1
38
39 stabilization on the OMM of depolarized mitochondria during mitophagy, as demonstrated by
40
41 inhibition of PINK1 accumulation upon CHCHD4 or GFER knockdown, or after treatment with
42
43 chemical inhibitors of the CHCHD4/GFER disulfide relay system (5).
44
45
46

47
48 Structural analysis of the PINK1 protein as well as sub-mitochondrial localization studies of PINK1
49
50 deletion mutants allowed to identify a putative OMM localization signal (OMS) between the amino
51
52 acids 74 and 93 of PINK1, which deletion impairs PINK1 retention on the surface of damaged
53
54 mitochondria and the subsequent mitophagy cascade (4). It is worth noting that a cysteine (amino
55
56 acid 92) is present in this region, whereas two other cysteine residues (amino acids 96 and 125)
57
58 are located in the just adjacent transmembrane domain (TMD) of PINK1. In this light, one could
59
60 speculate that, upon mitochondrial damage, activation of the CHCHD4/GFER disulfide relay
61
62
63
64
65

1
2
3
4 system may induce the formation of disulfide bonds in the region encompassing the OMS and the
5
6 TMD of PINK1. This could in turn inhibit the mitochondrial import and full proteolytic processing
7
8 of PINK1 and thus promote its accumulation on the OMM required for mitophagy induction.
9
10 According to this model, two PD-associated mutations involving cysteine residues located within
11
12 this region have been described, namely the C92F and C125G substitutions. Importantly, both
13
14 display reduced stabilization of PINK1 and, limited to the C125G mutant, impaired Parkin
15
16 recruitment to mitochondria in response to CCCP treatment (8).
17
18

19
20 Another PINK1 domain that could interact with CHCHD4 is the amphipathic helix (amino acids
21
22 166-172) containing hydrophobic and aromatic residues in the vicinity of a potential docking
23
24 cysteine 166. This segment of PINK1 could represent a putative internal IMS-targeting sequence,
25
26 which is proposed to allow the incoming protein substrate to recognize and interact with the
27
28 CHCHD4 oxidase (6). Indeed, deletion of this region (PINK1 Δ 166-171) strongly decrease the
29
30 PINK1 binding to CHCHD4 and, most importantly, two PD-related mutations in PINK1 (i.e. A168P
31
32 and V170G) also display a reduced interaction with CHCHD4 and impaired accumulation upon
33
34 CCCP treatment (5).
35
36

37
38 If, on the one hand, the CHCHD4/GFER mitochondrial import machinery seems to be important
39
40 to assist PINK1 accumulation on the OMM in the early phase of mitophagy, on the other hand
41
42 PINK1 could also regulate the activity of CHCHD4/GFER, thus providing a positive feedback loop
43
44 that sustains the import of other CHCHD4 substrates. In fact, the interaction between endogenous
45
46 PINK1 and CHCHD4 was detected not only in cells treated with CCCP but also under basal
47
48 conditions, suggesting a potential interplay between these two proteins that extends beyond the
49
50 mere regulation of PINK1 (5). By controlling the import and/or proper folding of specific subsets
51
52 of proteins, impairment of CHCHD4/Mia40 activity is known to (i) alter mitochondrial morphology
53
54 and dynamics, (ii) decrease the formation of electron transport chain complexes, (iii) induce
55
56 mitochondrial calcium dyshomeostasis and (iv) disrupt phospholipid metabolism, thus impinging
57
58 on the integrity of the IMM and cristae ultrastructure (6). Notably, many of these alterations are
59
60
61
62
63
64
65

1
2
3
4 often observed in PINK1 deficient cells as well. At least six CHCHD4 substrates (NDUFA8,
5
6 NDUFAF8, NDUFB7, NDUFB10, NDUFS5 and NDUFS8) are directly involved in the biogenesis
7
8 and proper assembly of mitochondrial respiratory complex I, which activity is impaired in PD
9
10 patients carrying distinct PINK1 loss-of-function mutations and, more in general, in the brain of
11
12 sporadic PD patients (3,6). Decreased complex IV activity is also observed in PINK1-KO flies and
13
14 in PINK1 null dopaminergic neurons, an effect previously linked to down-regulation of specific
15
16 mitochondrial chaperons involved in the assembly of complex IV subunits (9). Interestingly, the
17
18 CHCHD4/Mia40 substrates CMC1, CMC2, COX6B1 (otherwise called COXG), COA4, COA5,
19
20 COA6, COX17 and the copper chaperone CHCHD7 (else known as COX23) also participate in
21
22 the biogenesis and assembly of mitochondrial complex IV (6). Another CHCHD4/Mia40 substrate,
23
24 namely TRIAP1/Mdm35, could indirectly influence the activity of electron transport chain by
25
26 regulating the phospholipid composition (i.e. cardiolipin) of the IMM, which is essential for the
27
28 accurate assembly of respiratory complexes and supercomplexes (6,10). To this regard,
29
30 cardiolipin (CL) levels are significantly reduced in mitochondrial fractions from PINK1-KO mouse
31
32 embryonic fibroblasts (MEFs) compared to control MEFs, whereas CL supplementation rescues
33
34 the impairment of complex I activity observed in PINK1 deficient flies (3). As a structural
35
36 component of the IMM, CL also play a role in other key functions that were found altered in PINK1
37
38 mutant cells, including mitochondrial dynamics and apoptotic cell death.
39
40
41
42
43

44 Even if it appears quite clear that PINK1 and CHCHD4 regulate common downstream processes
45
46 maintaining mitochondrial homeostasis and function, further efforts are needed to provide a more
47
48 comprehensive characterization of the molecular mechanisms regulating the PINK1/CHCHD4
49
50 interplay as well as its physiological relevance in the neuronal context. To this regard, useful hints
51
52 may come from the discovery of Apoptosis-Inducing Factor (AIF) as a major CHCHD4 binding
53
54 partner in the IMS (6). *Aifm1* depletion in the mouse forebrain results in a neurodegenerative
55
56 phenotype characterized by altered cortical development and decreased neurons survival, an
57
58 effect likely due to the accumulation of dysfunctional mitochondria, which appeared fragmented,
59
60
61
62
63
64
65

1
2
3
4 with aberrant cristae ultrastructure and reduced OXPHOS capacity. A viral insertion in the *Aifm1*
5
6 gene, which causes an 80% reduction in the expression of AIF, underlies the progressive
7
8 neuronal degeneration observed in the *Harlequin (Hq)* mutant mice, one of the most reliable
9
10 models of mitochondrial complex I deficiency. Moreover, *Hq* mice are highly susceptible to the
11
12 MPTP toxin previously associated to PD, as demonstrated by the strong nigrostriatal
13
14 dopaminergic loss even in presence of subtoxic doses (6). Based on the current knowledge, AIF
15
16 acts upstream of CHCHD4 to regulate the co-translational import of CHCHD4 itself and, in turn,
17
18 the activity of the whole CHCHD4/GFER disulfide relay system. Thus, mitochondrial dysfunctions
19
20 observed in AIF deficient models, including impaired complex I activity, are significantly restored
21
22 upon CHCHD4 over-expression, whereas the beneficial effects of ectopically expressed AIF are
23
24 abolished after CHCHD4 downregulation (6).
25
26
27

28
29 The exact molecular mechanism through which AIF may control the CHCHD4-dependent
30
31 mitochondrial import machinery has not been fully elucidated yet. Previous findings revealed that,
32
33 upon binding to the reduced pyridine nucleotide cofactors NADH and NADPH, AIF undergoes a
34
35 conformational modification, which appears to enhance its interaction with CHCHD4 (6).
36
37 Interestingly, PD-related mutations in PINK1 and GBA genes decreased NAD levels, leading to
38
39 mitochondrial dysfunction and neurodegeneration. In contrast, boosting the intracellular pool of
40
41 pyridine nucleotide cofactors upon supplementation with the NAD precursors Nicotinamide (NAM)
42
43 or Nicotinamide Riboside (NR) rescued mitochondrial defects and neuronal loss in *Drosophila*
44
45 and human induced pluripotent stem cells (iPSC)-derived models of PD (11). If this translates in
46
47 increased AIF/CHCHD4 binding and enhanced CHCHD4 activity remains to be demonstrated, as
48
49 well as the potential contribution of this pathway in the neuroprotective effects of NAD precursors.
50
51 Another mechanism underlying regulation of the CHCHD4/GFER disulfide relay system could
52
53 involve potential post-translational modifications of either AIF or CHCHD4, which may control their
54
55 protein-protein interaction and modulate the activity of AIF/CHCHD4 complex. If PINK1 could play
56
57 a role in that sense, for instance by means of its kinase activity, still needs further investigation,
58
59
60
61
62
63
64
65

1
2
3
4 but distinct observations make this possibility worthy of being explored. First, PINK1 and CHCHD4
5
6 bind to each other not only in dysfunctional mitochondria, destined to mitophagy, but also in
7
8 healthy polarized organelles, thus extending the functional consequences of their interaction
9
10 behind the simple control of PINK1 accumulation on the OMM by CHCHD4 (5). This also
11
12 highlights the importance, often underestimated, of the mitochondrial intermembrane space
13
14 (IMS), where PINK1 localizes during its mitochondrial import, just before its processing and retro-
15
16 translocation to the cytosol. Moreover, even if some in vitro experiments suggested that the C-
17
18 terminal domain of PINK1 could remain in the cytosol during the import process, previous findings
19
20 reported that PINK1 interacts with and phosphorylates a number of IMS proteins, such as
21
22 HtrA2/Omi and TRAP1 (12).
23
24

25
26 Future research aiming to better characterize the PINK1 activity in the IMS of healthy
27
28 mitochondria and its relationship with the AIF/CHCHD4 machinery is warranted, and will answer
29
30 the question whether stimulating the disulfide relay system could represent an effective strategy
31
32 to rescue mitochondrial dysfunction in PD.
33
34
35
36
37
38
39
40

41 REFERENCES

- 42
43 1) Poewe W, Seppi K, Tanner CM, et al. Parkinson disease. *Nat Rev Dis Prim* (2017) **3**:1-
44 21.
- 45 2) Bender A, Krishnan KJ, Morris CM, et al. High levels of mitochondrial DNA deletions in
46 substantia nigra neurons in aging and Parkinson disease. *Nat Genet* (2006) **38**:515-7.
- 47 3) Brunelli F, Valente EM, and Arena G. Mechanisms of neurodegeneration in Parkinson's
48 disease: keep neurons in the PINK1. *Mech Ageing Dev* (2020), **189**:111277.
- 49 4) Sekine S and Youle RJ. PINK1 import regulation; a fine system to convey mitochondrial
50 stress to the cytosol. *BMC Biology* (2018) **16**:2.
- 51 5) Gao F, Zhang Y, Hou X, et al. Dependence of PINK1 accumulation on mitochondrial redox
52 system. *Aging Cell* (2020) **19**:e13211.
- 53 6) Reinhardt C, Arena G, Nedara K, et al. AIF meets the CHCHD4/Mia40-dependent
54 mitochondrial import pathway. *Biochim Biophys Acta Mol Basis Dis* (2020) **1866**:165746.
- 55 7) Imai Y, Meng H, Shiba-Fukushima K, et al. Twin CHCH Proteins, CHCHD2, and
56 CHCHD10: Key Molecules of Parkinson's Disease, Amyotrophic Lateral Sclerosis, and
57 Frontotemporal Dementia, *Int J Mol Sci* (2019) **20**: 908.
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

8) Sekine S, Wang C, Sideris DP, et al. Reciprocal Roles of Tom7 and OMA1 during Mitochondrial Import and Activation of PINK1. *Mol Cell* (2019) **73**:1028–1043.

9) Kim KH, Song S, Yoon SH, et al. Rescue of PINK1 Protein Null-specific Mitochondrial Complex IV Deficits by Ginsenoside Re Activation of Nitric Oxide Signaling. *J Biol Chem* (2012) **287**:44109-44120.

10) Maguire JJ, Tyurina YY, Mohammadyani D, et al. Known unknowns of cardiolipin signaling: The best is yet to come. *Biochim Biophys Acta Mol Cell Biol Lipids* (2017) **1862**:8-24.

11) Schondorf DC, Ivanyuk D, Baden P, et al. The NAD⁺ Precursor Nicotinamide Riboside Rescues Mitochondrial Defects and Neuronal Loss in iPSC and Fly Models of Parkinson's Disease. *Cell Reports* (2018) **23**:2976–2988.

12) Arena G & Valente EM. PINK1 in the limelight: multiple functions of an eclectic protein in human health and disease. *J Pathol* (2017) **241**: 251–263.