



On the logarithmic coarse structures of Lie groups and hyperbolic spaces

Gabriel Pallier

► To cite this version:

Gabriel Pallier. On the logarithmic coarse structures of Lie groups and hyperbolic spaces. *Geometriae Dedicata*, 2022, 216 (5), pp.56. 10.1007/s10711-022-00714-6 . hal-03214831

HAL Id: hal-03214831

<https://hal.science/hal-03214831>

Submitted on 3 May 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON THE LOGARITHMIC COARSE STRUCTURES OF LIE GROUPS AND HYPERBOLIC SPACES

GABRIEL PALLIER

ABSTRACT. We characterize the Lie groups with finitely many connected components that are $O(u)$ -bilipschitz equivalent (almost quasiisometric in the sense that the sublinear function u replaces the additive bounds of quasiisometry) to the real hyperbolic space, or to the complex hyperbolic plane. The characterizations are expressed in terms of deformations of Lie algebras and in terms of pinching of sectional curvature of left-invariant Riemannian metrics in the real case. We also compare sublinear bilipschitz equivalence and coarse equivalence, and prove that every coarse equivalence between the logarithmic coarse structures of geodesic spaces is a $O(\log)$ -bilipschitz equivalence. The Lie groups characterized are exactly those whose logarithmic coarse structure is equivalent to that of a real hyperbolic space or the complex hyperbolic plane.

1. INTRODUCTION

1.1. Background. Let X and Y be metric spaces. A map $\phi : X \rightarrow Y$ is a quasi-isometry if there exists $\lambda \geq 1$ and $c \geq 0$ such that $\lambda^{-1}d(x, x') - c \leq d(\phi(x), \phi(x')) \leq \lambda d(x, x') + c$ and for every y in Y , $d(y, \phi(X)) \leq c$. Let a locally compact, compactly generated group G act continuously co-compactly properly by isometries on a locally compact geodesic space X ; we call X a geometric model of G . Every such G has a geometric model (e.g. Cayley graphs if it is finitely generated, Riemannian metrics if it is connected Lie), and two geometric models of a given G will always be equivariantly quasiisometric. Thus one can speak of quasiisometries between compactly generated locally compact groups.

Quasiisometries arose from the interpretation by Margulis of the work of Mostow on the rigidity of locally symmetric spaces [Mar70]. Specifically, Margulis conjectured that a quasiisometry of a higher rank symmetric space X should lie at bounded distance from an isometry, implying Mostow rigidity for the co-compact lattices in X , but also the fact that any finitely G quasiisometric to X must surject with finite kernel onto such a uniform lattice. This was first proved by Kleiner and Leeb using asymptotic cones, a tool formerly introduced by Gromov, in the form recast by van den Dries and Wilkie [KL97]. The interplay of quasiisometries and asymptotic cones can actually be expressed in the following way: between geodesic metric spaces, a map is a quasiisometry if and only if it goes through any asymptotic cone (with possibly moving observation centers); see §2.3 for a precise

Date: May 2, 2021.

2010 *Mathematics Subject Classification.* Primary 20F69, 20F67; Secondary 53C24, 53C30, 57T10, 22E25.

Author supported by the European Research Council, ERC Starting Grant 713998 GeoMeG ‘Geometry of Metric Groups’.

statement. Kleiner and Leeb's theorem is part of a more general principle which, in contrast with Mostow rigidity, makes sense (and is stated below) for locally compact compactly generated groups.

Theorem (Many authors, see [Cor18, Theorem 19.25] and the references there). *Let G be a compactly generated locally compact group and let X be a Riemannian symmetric space of non-compact type. The following are equivalent:*

- (1) G is quasiisometric to X .
- (2) X is a Riemannian geometric model for G .

Moreover, if G is a Lie group isomorphic to a closed subgroup of upper triangular real matrices (call such groups completely solvable), then the former are equivalent to:

- (3) G is isomorphic to a maximal completely solvable¹ subgroup of $\text{Isom}(X)$.

The case G finitely generated and $X = \mathbb{H}_{\mathbb{R}}^n$, $n \geq 3$ is up to formulation due to Tukia [Tuk86] and was among the early results motivating the first formulation of quasiisometric rigidity by Gromov [Gro84]. Gromov almost simultaneously proposed a vast programme of classifying finitely generated groups and isometrically homogeneous spaces up to quasiisometry [Gro83]. For nonsemisimple connected or nonarchimedean Lie groups and their lattices, this is far from being achieved today.

Between geodesic metric spaces, quasiisometries are exactly the coarse equivalences, that is, they respect the bounded coarse structure described as the family of entourages

$$\mathcal{E}^{O(1)} = \left\{ E \subseteq X \times X : \exists D > 0, \sup_{(x,x') \in E} d_X(x, x') \leq D \right\}.$$

A broad interpretation of Gromov's programme is the following: classify the coarse structures generated by compactly generated groups, and characterize those that are generated by particular geometric models, especially the Riemannian symmetric or homogeneous spaces. Recently, certain extensions of Gromov's questions have been addressed where coarse surjectivity is relaxed. These are the study of the rigidity of quasiisometric embeddings (see [FN15] and [FW18] for symmetric spaces) and of the (non)-existence of coarse embeddings (see [HMT20b] for connected Lie groups).

1.2. Main results. In this paper, we are interested in maps more general than quasiisometries. In contrast with quasiisometries, these can still be characterized as going through asymptotic cones, though not through asymptotic cones for any sequence of basepoints (we elaborate on [Cor11] for this; see §2.3 for a precise statement). The coarse surjectivity assumption is not exactly relaxed, but adapted accordingly.

For the needs of the next definition, say that a function $u : [0, +\infty) \rightarrow (0, +\infty)$ is admissible if $\limsup_{r \rightarrow +\infty} u(r)/r = 0$ (that is, u is sublinear) and for every $A \geq 1$ there exists $B < +\infty$ such that for all sequences (r_n, s_n) with $1/A \leq \inf s_n/r_n \leq$

¹Beware that the maximal solvable subgroups of $\text{Isom}(X)$ (which is a real Lie group) are not always completely solvable; they only have a co-compactly embedded such subgroup.

$\sup s_n/r_n \leq A$, $\sup u(s_n)/u(r_n) \leq B$. Examples of admissible function include $u(r) = r^\alpha \log^\beta(r)$ for $r \geq 2$ (and $u(r) = 1$ otherwise) when $\alpha \in (-\infty, 1)$ and $\beta \in (-\infty, +\infty)$.

Definition (After² [Cor19]). Let u be an admissible function. A map $\phi : (X, o_X) \rightarrow (Y, o_Y)$ between pointed metric spaces realizes a (large-scale) $O(u)$ -bilipschitz equivalence if there are $\kappa \geq 1$ and $c \geq 0$ such that for every r large enough

$$(1.1) \quad -cu(r) + \frac{d_X(x, x')}{\kappa} \leq d_Y(\phi(x), \phi(x')) \leq \kappa d_X(x, x') + cu(r)$$

$$(1.2) \quad d_Y(y, \phi(x)) \leq cu(r),$$

for all $x, x' \in B(o_X, r)$ and $y \in B(o_Y, r)$.

We also call $o(r)$ -bilipschitz equivalence, or sublinear bilipschitz equivalence (abbreviated SBE in some places), a ϕ such that (1.1) and (1.2) hold with some unspecified strictly sublinear function in lieu of cu .

Quasiisometries correspond to $u \equiv 1$. Of particular importance in this paper is $u = \log$. Given an admissible function u , we consider the coarse structure on metric spaces with the following entourages:

$$\mathcal{E}^{O(u)} = \left\{ E \subseteq X \times X : \sup_{(x, x') \in E, (x, x') \rightarrow \infty} \limsup \frac{d_X(x, x')}{u(|x|)} < +\infty \right\}.$$

These are quantitative refinements of the coarse structure introduced in [DS07]. $O(u)$ -bilipschitz equivalences are always $\mathcal{E}^{O(u)}$ -coarse equivalences. We prove that the converse holds between geodesic spaces when $u = \log$:

Theorem A. *Assume that X and Y are geodesic. Then $\phi : X \rightarrow Y$ is $O(\log)$ -bilipschitz if and only if it is a coarse equivalence of $\mathcal{E}^{O(\log)}$.*

This is a variant of the well-known fact that coarse equivalences between geodesic spaces are quasiisometries, however the proof is significantly more involved.

Keeping quasiisometric rigidity and classification in mind, it is natural to ask:

Question 1.1 (Rigidity). Let u be as above, $u \geq 1$. Which compactly generated locally compact groups G are $O(u)$ -bilipschitz equivalent to a given symmetric space X ?

Question 1.2 (Classification). Given u as above, $u \geq 1$, classify isometrically homogeneous spaces up to $O(u)$ -bilipschitz equivalence.

The following theorem was stated in the introduction of the author's thesis. While essentially following from the combination of [DS07], [HP13] and the coarse interpretation of $o(r)$ -bilipschitz equivalences, it was not extracted at first sight from the literature, so we provide a proof here (relying on the above cited works). Recall for the statement that all the maximal compact subgroups of a connected Lie groups are conjugated [Bor95].

²This is [Cor19, Definition 2.1] with a mild difference in the definition of the class of admissible functions that we make in order to include functions with limit 0 at ∞ (see §2 for why).

Theorem B (After [DS07] and [HP13]). *Let G and H be connected Lie groups. If there exists a $o(r)$ -bilipschitz equivalence $\phi : G \rightarrow H$, then*

$$(1.3) \quad \text{geodim}(G) = \text{geodim}(H),$$

where $\text{geomdim}(G)$ denotes $\dim G/K$ if K is any maximal compact subgroup of G . Especially, if G and H are solvable and simply connected, then $\dim G = \dim H$.

The theorem actually holds for every $o(r)$ -coarse equivalences ϕ , see §2.5. If G and H are nilpotent, then geodim is the covering dimension of their asymptotic cones and Theorem B also follows from [Pan83].

Next, building on [Cor11], [CT11] and [Pal20] (which was already concerned with Question 1.2) we formulate below a partial answer to Question 1.1 for connected Lie groups G and real hyperbolic space X . While this is not made apparent in the statement, all the groups obtained are either of Heintze or rank-one type, in the typology of [CT11] and [CCMT15].

Theorem C. *Let G be a Lie group with finitely many connected components and $n \geq 2$ an integer. The following are equivalent:*

- (C.1) *G is $O(u)$ -bilipschitz equivalent to $\mathbb{H}_{\mathbf{R}}^n$, for some sublinear admissible u .*
- (C.2) *G is $O(\log)$ -bilipschitz equivalent to $\mathbb{H}_{\mathbf{R}}^n$.*
- (C.3) *For every $\varepsilon > 0$, G has a Riemannian model X with $-1 \leq K \leq -1 + \varepsilon$.*

Moreover, if G is completely solvable with Lie algebra $\mathfrak{g}$, the former conditions are equivalent to:

- (C.4) *$\mathfrak{g}$ degenerates to the (isomorphism class of a) maximal completely solvable subalgebra $\mathfrak{g}_{\infty}$ of $\mathfrak{o}(n, 1)$.*
- (C.5) *The Lie algebra $\mathfrak{g}$ decomposes as $[\mathfrak{g}, \mathfrak{g}] \oplus \mathbf{R}A$, where $[\mathfrak{g}, \mathfrak{g}]$ is abelian and ad_A is unipotent on $[\mathfrak{g}, \mathfrak{g}]$.*

Here saying that $\mathfrak{g}$ degenerates to $\mathfrak{g}_{\infty}$ means that the Zariski closure of the orbit of $\mathfrak{g}$ in the variety of Lie algebra laws contains $\mathfrak{g}_{\infty}$, which occurs especially if there is a continuous $(\varphi_t)_{t \in [0, +\infty)}$ in $\text{GL}(\mathfrak{g})$ and a linear isomorphism $\psi : \mathfrak{g} \rightarrow \mathfrak{g}_{\infty}$ such that for every $X, Y \in \mathfrak{g}$,

$$\lim_{t \rightarrow +\infty} \varphi_t^{-1}[\varphi_t X, \varphi_t Y]_{\mathfrak{g}} = \psi^{-1}[\psi X, \psi Y]_{\mathfrak{g}_{\infty}}.$$

Theorem C combines known results. That (C.1) implies (C.3) rests on [CT11] and [Pal20], the equivalence of the last two conditions (C.4) and (C.5) is [Lau03, Theorem 6.2] with minor enhancement, the implication from (C.3) to (C.5) uses [Pan89a], while the fact that (C.5) implies (C.2) is a consequence of [Cor11]. When $n = 2$, Theorem C reduces to a weak form of [Cor19, Corollary 1.10(2)]. The statement is simpler for homogeneous metrics have constant curvature, hold with the mere assumption that G be compactly generated locally compact, and the techniques are specific, relying essentially on [Gab92], [CJ94].

For general connected Lie groups, the process of going from $\mathfrak{g}$ to a less complicated $\mathfrak{g}_{\infty}$ so that the simply connected G and G_{∞} remain $O(u)$ -bilipschitz equivalent has an alternative description given in [Cor11] (recalled here in Theorem 5.1) which does not require degenerations. Our formulation using degeneration is half-successful in this generality. While it also applies well when $\mathfrak{g}$ is nilpotent (in

this case it is due to Pansu [Pan83]), we do not know whether $\mathfrak{g}_\infty$ is a degeneration of $\mathfrak{g}$ in general. This will be discussed in §5.1.

The appearance of the sectional curvature pinching in characterization (C.3) might appeal to some comments. The sphere theorem of Berger and Klingenberg implies that on a positively curved Riemannian manifold of positive curvature, a pinching sufficiently close to 1 determines the homotopy type of the (finite) universal cover. Namely, the latter must be a sphere. As demonstrated by Gromov and Thurston, there is no counterpart for this in negative curvature as one constructs sequences of closed manifolds supporting negatively curved metrics, arbitrarily pinched close to 1, albeit with vanishing first cohomology, hence not homotopy equivalent to any locally symmetric space of constant negative curvature [GT87].

This is not even repaired if one replaces homotopy equivalence with quasiisometry, as one constructs isometrically homogeneous manifolds with pinching $> -1/4$ or even arbitrarily close to -1 (characterized in [EH96], see §3.1), that are not quasiisometric to $\mathbb{H}_{\mathbf{R}}^n$ [Xie14]. Theorem C implies the following as far as Lie groups are concerned.

Corollary D (of Theorem C). *If a connected Lie group G has Riemannian models with pinching arbitrarily close to -1 , then its sublinear Higson corona $\nu_L G$ is homeomorphic to that of a real hyperbolic space.*

(We recall the definition of the sublinear Higson corona in §2.5.)

Finally, we also characterize the Lie groups $O(u)$ -bilipschitz equivalent to $\mathbb{H}_{\mathbf{C}}^2$. Following [Cor15], say that the locally compact G and H are commable if there exists a finite sequence of homomorphisms with compact kernels and co-compact images (both directions allowed) between G and H .

Theorem E. *Let G be a Lie group. with finitely many connected components. The following are equivalent:*

- (E.1) G is $O(u)$ -bilipschitz equivalent to $\mathbb{H}_{\mathbf{C}}^2$
- (E.2) G is $O(\log)$ -bilipschitz equivalent to $\mathbb{H}_{\mathbf{C}}^2$
- (E.3) G is commable either to the semisimple $\mathrm{SU}(2,1)$ or to the solvable $S' = H_3 \rtimes \mathbf{R}$, where H_3 is the 3-dimensional Heisenberg group and $t \in \mathbf{R}$ acts by

$$t \cdot \exp(x, y, z) = \exp(e^t x + t e^t y, e^t y, e^{2t} z)$$

in a basis of infinitesimal generators X, Y, Z such that $[X, Y] = Z$.

Moreover, if G is completely solvable, the former conditions are equivalent to:

- (E.4) $\mathfrak{g}$ degenerates to the maximal completely solvable subalgebra of $\mathfrak{u}(2,1)$

where $\mathfrak{g}$ denotes the Lie algebra of G .

The restriction that G be a connected Lie group makes Theorems C and E very special compared to the QI rigidity recalled above, and we benefit from some constraints of the structure theory of Lie groups. Unlike Theorem C, Theorem E requires some additional technical work, done in §4.

1.3. Other spaces. We know little even about Question 1.2 for higher rank symmetric spaces and other settings, even when quasiisometric rigidity is known to

hold. In the end of this paper, we summarize the current situation for symmetric space of higher rank and Fuchsian buildings; especially we explain why their classification is still open at the time of writing.

1.4. Organization of the paper. §2 is a general discussion on the theoretical status of SBE (especially, as compared to QI). It is not concerned with Lie groups and can be read independently. §2.1 provides some preliminaries for §2. §3 and §4 establish the characterizations of Lie groups $O(u)$ -bilipschitz equivalent to real, resp. complex hyperbolic space, and follow a similar scheme, so we advise to read §3 first. Most of the technical input in this paper serve the proofs of Theorems A and E and is concentrated in §2.4 and §4.1 respectively. SBE appears to be quite a new notion and some of the contents of this paper are rather expository in nature, including especially §2.5 on Theorem B, §3.1 and §3.2 preparing the proof of Theorem C, and §5.1 on general connected Lie groups. §5.2 and §5.3 gather a collection of independent remarks. Finally, a certain amount of actual Lie algebra cohomology computations (for trivial and adjoint modules) are required in particular in Lemma 4.5 and Example 5.8; we summarize these in Appendix A.

Convention, notation. When $G, H, \dots$ are simply connected Lie groups, then $\mathfrak{g}, \mathfrak{h}, \dots$ denote their Lie algebra. We often consider semi-direct products of the form $N \rtimes \mathbf{R}$ or $\mathfrak{n} \rtimes \mathbf{R}$; we then write $N \rtimes_{\alpha} \mathbf{R}$ or $\mathfrak{n} \rtimes_{\alpha} \mathbf{R}$ meaning that the Lie algebra representation $\rho : \mathbf{R} \rightarrow \text{Der}(\mathfrak{n})$ (and not the Lie group representation) is determined by $1 \mapsto \alpha$. If V is a module and n a nonnegative integer, we denote by $\Lambda^n V$ its n -fold exterior product and by $\Lambda^n V^*$ the n -fold exterior product of its dual. If $\mathfrak{g}$ is a Lie (sub)algebra, $\text{Vect}(\mathfrak{g})$ will denote its underlying vector (sub)space. (This is useful to avoid confusions because we may sometimes consider several Lie brackets on a given space.)

CONTENTS

1. Introduction	1
2. Coarse geometry and Theorems A and B	6
3. Real hyperbolic spaces and Theorem C	20
4. Proof of Theorem E	29
5. Some remarks on spaces other than $\mathbb{H}_{\mathbf{R}}^n$ and $\mathbb{H}_{\mathbf{C}}^n$	32
Appendix A. Methods used for the cohomology computations	40
References	45

2. COARSE GEOMETRY AND THEOREMS A AND B

This section motivates sublinear bilipschitz equivalence (defined in §1.2) by comparing it to the more standard notions of quasiisometry and coarse equivalence. This comparison will be made through the relations that sublinear bilipschitz equivalence enjoys with asymptotic cones and certain coarse structures. The relation to asymptotic cones is the reason why they were introduced by Cornuier

in the first place, in [Cor08] and then more explicitly³ in [Cor11], [Cor19] (See §2.3.1 for precisely why). In the end of this section, we show that the geometric dimension of connected Lie groups is a SBE invariant.

2.1. Preliminaries.

Definition 2.1 (Coarse equivalence and quasiisometry). Let X and Y be two metric spaces and let $\phi : X \rightarrow Y$. ϕ is a (uniform) coarse embedding if there exists two proper functions ρ_- and $\rho_+ : [0, +\infty) \rightarrow [0, +\infty)$ such that for every $x, x' \in X$

$$(2.1) \quad \rho_-(d_X(x, x')) \leq d_Y(\phi(x), \phi(x')) \leq \rho_+(d_X(x, x')).$$

ϕ is a coarse equivalence if moreover, there exists a coarse embedding $\psi : Y \rightarrow X$ and a constant $R \geq 0$ such that for all $x \in X$, $d_X(\psi \circ \phi(x), x) \leq R$ and for all $y \in Y$, $d_Y(\phi \circ \psi(y), y) \leq R$; we call g a coarse inverse. ϕ is a (κ, c) -quasiisometric embedding if ρ_- and ρ_+ can be taken affine in (2.1), namely $\rho_{\pm}(r) = \kappa^{\pm 1}r \pm c$. If in addition ϕ a coarse equivalence, ϕ is called a quasiisometry and any coarse inverse g is also a quasiisometry; equivalently a quasiisometry is a quasiisometric embedding ϕ such that $\sup_{y \in Y} d_Y(y, \phi(X)) < +\infty$. We may define a quasiisometry only on a net, that is, a closed subspace $X^{(0)} \subseteq X$ such that $\sup_{x \in X} d(x, X^{(0)}) < +\infty$.

Proposition 2.2 (See e.g. [CdlH16, 3.B.9]). *Let X and Y be two metric spaces and let $\phi : X \rightarrow Y$ be a coarse equivalence. If X and Y are geodesic, then any coarse equivalence $X \rightarrow Y$ is a quasiisometry.*

Proposition 2.3. *Let G be a compactly generated locally compact group. Then*

- (1) *If G acts continuously, properly cocompactly on the locally compact geodesic spaces X and Y , then there exists a quasiisometry $\phi : X \rightarrow Y$ such that $\sup_{(g,x) \in G \times X} d_Y(\phi(g.x), g.\phi(x)) < +\infty$.*
- (2) *There exists X locally compact geodesic metric space and an isometric proper co-compact continuous action of G on X .*

(1) is a consequence of [CdlH16, Theorem 4.C.5]. For (2), see [CCMT15, Proposition 2.1]. In this paper we call X and Y as in the previous proposition geometric models for G .

2.2. Admissible sublinear functions.

Definition 2.4. Call $u : [0, +\infty) \rightarrow (0, +\infty)$ admissible if $\limsup_{r \rightarrow +\infty} u(r)/r = 0$ and for every $A \geq 1$ there exists $B < +\infty$ (only depending on A) such that for all sequences (r_n, s_n) with $r_n \rightarrow +\infty$ and $1/A < \inf s_n/r_n \leq \sup s_n/r_n < A$, one has

$$(2.2) \quad 1/B \leq \liminf \frac{u(s_n)}{u(r_n)} \leq \limsup \frac{u(s_n)}{u(r_n)} \leq B.$$

Lemma 2.5. *Let $u : [0, +\infty) \rightarrow (0, +\infty)$ be a sublinear function. If u is nondecreasing and $\limsup u(2r)/u(r) < +\infty$, resp. if u is nonincreasing and $\liminf u(2r)/u(r)$, then u is admissible.*

³We should warn the reader about terminology: they were called “cone bilipschitz” in [Cor11] and “asymptotically bilipschitz” in [DK18].

Proof. Let us consider only the case where u is nondecreasing, the proof going the same way. Let $A > 1$ and (r_n, s_n) be such that $r_n \rightarrow +\infty$ and $\{s_n/r_n\} \in [1/A, A]$. Set $\beta = \limsup u(2r)/u(r)$. Since $u(s_n)/u(r_n) \leq 1$ when $s_n \leq r_n$, one has

$$\limsup \frac{u(s_n)}{u(r_n)} = \sup \left(1, \limsup_{n: s_n \geq r_n} \frac{u(s_n)}{u(r_n)} \right) \leq \beta^{\lceil \log_2 A \rceil}$$

This is the inequality on the right in (2.2) with $B = \beta^{\lceil \log_2 A \rceil}$. The left inequality is obtained by reversing r_n and s_n . $\square$

The usefulness of Lemma 2.5 may not be obvious. Let us give two motivations. The first is that it ensures that the functions u considered in [Cor19, Definition 2.4] are admissible in our sense. The second is that, while Definition 2.4 allows a unified treatment for sublinear functions u with $u(r) \rightarrow +\infty$ or $u(r) \rightarrow 0$ and is sufficient for our purposes in §2.3 and §2.4, it appears that it is often easier to argue and prove the main statement of this section with monotonic functions u .

The above notion of admissible function resembles the much-studied class of (not necessarily sublinear) regularly varying function in real analysis, but we found no implication between the two without further assumptions.

2.3. Going through cones. Let (σ_n) be a sequence of positive real numbers. For $x_n, x'_n \in X^{\mathbf{N}}$, denote $x_n \sim_{\sigma_n} x'_n$ if $\sup d(x_n, x'_n)/\sigma_n < +\infty$ and $x_n \approx_{\sigma_n} x'_n$ if $\limsup d(x_n, x'_n)/\sigma_n = 0$. $\text{Precone}(X, x_n, \sigma_n)$ denotes the $\sim_{\sigma_n}$ equivalence class of X . Further, given a nonprincipal ultrafilter $\omega \in \beta\mathbf{N} \setminus \mathbf{N}$, $\text{Cone}_{\omega}(X, x_n, \sigma_n)$ is⁴ the largest quotient of $\text{Precone}(X, x_n, \sigma_n)$ whose points are separated by distance $d_{\omega}((x'_n), (x''_n)) = \lim_{n \rightarrow \omega} d(x'_n, x''_n)/\sigma_n$. When $\sigma_n \rightarrow +\infty$ and x_n is bounded (or $|x_n| \ll \sigma_n$), the cone does not depend on x_n so we write it $\text{Cone}_{\omega}^{\bullet}(X, \sigma_n)$. Though our main interest is in homogeneous spaces, it is useful to work out some examples of asymptotic cones of nonhomogeneous spaces in order to appreciate the difference between quasiisometry and $O(u)$ -bilipschitz equivalence.

Examples 2.6. For $i \in \{1, \dots, 4\}$ let P_i be a Riemannian plane with metric $ds^2 = dr^2 + A_i(r)^2 d\theta^2$, where $A_1(r) = 1/r$, $A_2(r) = 1$, $A_3(r) = \log r$ and $A_4(r) = r/2$ for r large enough. See some sketches of P_i on Figure 1, and various cones on Table 1.

Proposition 2.7 (Characterizing quasiisometries I). *Let X and Y be metric spaces, and $\phi : X \rightarrow Y$. Then, ϕ is a quasiisometric embedding if and only if for every (σ_n) such that $\lim_n \sigma_n = +\infty$, it holds:*

$$(I_{\sigma}) \quad \forall (x_n) \in X^{\mathbf{N}}, \forall (x'_n) \in X^{\mathbf{N}}, x_n \sim_{\sigma_n} x'_n \implies \phi(x_n) \sim_{\sigma_n} \phi(x'_n)$$

$$(II_{\sigma}) \quad \forall (x_n) \in X^{\mathbf{N}}, \forall (x'_n) \in X^{\mathbf{N}}, \phi(x_n) \approx_{\sigma} \phi(x'_n) \implies x_n \approx_{\sigma} x'_n$$

and then, given any such σ_n , for all pair $(x_n) \in X^{\mathbf{N}}$ and $(y_n) \in Y^{\mathbf{N}}$, either $\phi(\text{Precone}(X, x_n, \sigma_n)) \cap \text{Precone}(Y, y_n, \sigma_n)$ is empty or for every $\omega \in \beta\mathbf{N} \setminus \mathbf{N}$, ϕ

⁴When $\sigma_n \rightarrow 0$ and x_n is constant, the space $\text{Cone}_{\omega}(X, x_n, \sigma_n)$ is more commonly referred to as a metric tangent. However because our emphasis is on large-scale geometry and moving basepoints, and because the distinction would be artificial here, we denote both by the same name.




FIGURE 1. Sketch view of the four Riemannian planes of Example 2.6 with $U(1) \rtimes \mathbb{Z}/2\mathbb{Z}$ symmetry.

induces a bilipschitz embedding

$$(\text{Cone}) \quad \text{Cone}_\omega(\phi, x_n, y_n, \sigma_n) : \text{Cone}_\omega(X, x_n, \sigma_n) \rightarrow \text{Cone}_\omega(Y, y_n, \sigma_n)$$

whose bilipschitz constant only depends on ϕ . Further, ϕ is a quasiisometry if and only if for every σ_n with $\lim_n \sigma_n = +\infty$, the conditions (I_σ) and (II_σ) hold and in addition

$$(III_\sigma) \quad \forall (y_n) \in Y^\mathbb{N} \exists (x_n) \in X^\mathbb{N} : \phi(x_n) \sim_{\sigma_n} y_n$$

and then for every x_n, y_n, σ_n as above $\text{Cone}_\omega(\phi, x_n, y_n, \sigma_n)$ is either completely undefined or a bilipschitz homeomorphism.

Proof. First, assume that ϕ is not a quasiisometric embedding. Especially it is not a coarse embedding, so there exists an integer $M \geq 0$ and a sequence of points (x_n, x'_n) such that either $d(x_n, x'_n) \leq M$ and $d(\phi(x_n), \phi(x'_n)) \geq \rho(n)$ or $d(x_n, x'_n) \geq \rho(n)$ and $d(\phi(x_n), \phi(x'_n)) \leq M$, where $\rho(n) \rightarrow +\infty$. In the first case, $\text{Precone}(x_n, \rho(n)^{1/2})$ contains (x_n) and (x'_n) , but no $\text{Precone}(y_n, \rho(n)^{1/2})$ will contain $(\phi(x_n))$ and $(\phi(y_n))$ at the same time, contradicting (I_σ) . In the second case, note that $\phi(x_n) \approx_{\rho_n} \phi(x'_n)$, while $x_n \sim_{\rho_n}$ does not hold, contradicting (II_σ) with $\sigma = \rho$. Conversely, assume that ϕ is a (κ, c) -quasiisometric embedding; then $x_n \sim_{\sigma_n} y_n$ means that $d_X(x_n, y_n) \leq C\sigma_n$ for some $C \geq 0$, so that $d_Y(\phi(x_n), \phi(y_n)) \leq \kappa C\sigma_n + c \leq (\kappa C + 1)\sigma_n$ for $n > \sup\{m : \sigma_m \leq c\}$. This proves (I_σ) ; the proof of (II_σ) goes the same way using the left inequality in (2.1) with $\rho_-(r) = \kappa^{-1}r - c$.

Now assume that ϕ is a quasiisometric embedding (or equivalently has (I_σ) and (II_σ)) and let (x_n) and (y_n) be sequences in X and Y . Then $\phi(\text{Precone}(X, x_n, \sigma_n)) \cap \text{Precone}(Y, y_n, \sigma_n)$ equals

$$\begin{cases} \emptyset & y_n \not\sim \phi(x_n) \\ \phi(\text{Precone}(X, x_n, \sigma_n)) & y_n \sim \phi(x_n). \end{cases}$$

If the second case occurs, let κ be the large scale bilipschitz constant of ϕ . For any $\omega \in \beta\mathbb{N} \setminus \mathbb{N}$, $\text{Cone}_\omega(\phi, x_n, \sigma_n)$ exists and is κ -bilipschitz.

To prove that ϕ is a quasiisometry if and only if it has (I_σ) , (II_σ) and (III_σ) for all σ with limit $+\infty$, it remains only to prove that (III_σ) implies coarse surjectivity (the converse being clear). If ϕ is not coarsely surjective, then there exists a sequence (y_n) in (Y_n) and $\rho_n \rightarrow +\infty$ such that $B(y_n, \rho_n) \cap \phi(X) = \emptyset$. This

disproves (III_σ) with $\sigma_n = \sqrt{\rho_n}$. Finally, if ϕ is a quasiisometry, then for every parameters x_n, y_n, σ_n with $\sigma_n \rightarrow +\infty$, $\phi(\text{Precone}(X, x_n, \sigma_n)) \cap \text{Precone}(Y, y_n, \sigma_n)$ becomes

$$\begin{cases} \emptyset & y_n \approx \phi(x_n) \\ \phi(\text{Precone}(Y, y_n, \sigma_n)) & y_n \sim \phi(x_n) \end{cases}$$

and in the latter case, for every $\omega \in \beta\mathbf{N} \setminus \mathbf{N}$, $\text{Cone}_\omega(\phi, x_n, \sigma_n)$ is a bilipschitz homeomorphism, with bilipschitz constant κ independent of ω . $\square$

Proposition 2.8 (Characterizing quasiisometries, II). *Let X and Y be metric spaces and $\phi : X \rightarrow Y$. If for all $(x_n, y_n) \in X^{\mathbf{N}} \times Y^{\mathbf{N}}$ and (σ_n) with limit $+\infty$, either*

$$\phi(\text{Precone}(X, x_n, \sigma_n)) \cap \text{Precone}(Y, y_n, \sigma_n) = \emptyset$$

or $\text{Cone}_\omega(\phi, x_n, y_n, \sigma_n)$ is well-defined and a bilipschitz embedding for all $\omega \in \beta\mathbf{N} \setminus \mathbf{N}$, then ϕ is a quasiisometric embedding.

If for all (x_n, y_n, σ_n) as above, either

$$\phi(\text{Precone}(X, x_n, \sigma_n)) \cap \text{Precone}(Y, y_n, \sigma_n) = \emptyset$$

or $\text{Cone}_\omega(\phi)$ is well-defined and a bilipschitz homeomorphism for all ω , then ϕ is a quasiisometry.

Proof. The first hypothesis implies, for every σ , the conditions (I_σ) and (II_σ) of Proposition 2.7 for ϕ (where the injectivity of the coned map implies (II_σ)). Similarly, the second hypothesis implies, for every σ , (I_σ), (II_σ) and (III_σ). $\square$

The characterization given by Proposition 2.8 may be summarized as follows: a quasiisometry is a map between metric spaces which, when photographed between any pair of asymptotic cones with equal scaling factors, is either completely undefined or induces a bilipschitz homeomorphism.

As mentionned in the introduction, $o(r)$ -bilipschitz equivalences are the maps inducing bilipschitz homeomorphisms between asymptotic cones with fixed base-points. This is less demanding than the previous characterization. We recall Cornulier's characterization below.

Proposition 2.9 (Cornulier). *Let X and Y be metric spaces. Let $\phi : X \rightarrow Y$. The following are equivalent:*

(2.9.1) *ϕ is $o(r)$ -bilipschitz, i.e. There exists $\kappa \geq 1$ and $v : \mathbf{R}_{\geq 0} \rightarrow \mathbf{R}_{\geq 0}$ with $\lim_{r \rightarrow +\infty} v(r)/r = 0$ and for every $(x, x') \in X$ and $y \in Y$,*

$$\begin{aligned} -v(|x| \vee |x'|) + \frac{1}{\kappa} d_X(x, x') &\leq d_Y(\phi(x), \phi(x')) \\ &\leq \kappa d_X(x, x') + v(|x| \vee |x'|) \\ d_Y(y, \phi(x)) &\leq v(|y|), \end{aligned}$$

(2.9.2) *For every sequence (σ_n) of positive real numbers with $\sigma_n \rightarrow +\infty$, there is a well-defined, bilipschitz homeomorphism*

$$(\text{Cone}^\bullet) \quad \text{Cone}_\omega^\bullet(\phi, \sigma_n) : \text{Cone}_\omega^\bullet(X, \sigma_n) \rightarrow \text{Cone}_\omega^\bullet(Y, \sigma_n)$$

Proof. This results from the combination of [Cor11, Propositions 2.4, 2.5, 2.9, 2.12 and 2.13]. There is no sequence σ_n in Cornulier's statement, however the formulations are easily seen to be equivalent to ours. $\square$

In this way, the groupoids of quasiisometries and $o(r)$ -bilipschitz equivalences respectively are the largest groupoids over metric spaces so that the parametrized family of functors Cone and $\text{Cone}^\bullet$ respectively are well defined to the groupoid of metric spaces with bilipschitz homeomorphism. Note that bilipschitzness at the level of asymptotic cone came for free in Proposition 2.7, while it is explicitly required in Proposition 2.9, and there is indeed a strictly larger groupoid, that of isomorphisms in the category of cone-defined maps, whose pictures through $\text{Cone}^\bullet$ only have nonzero and finite local lipschitz and expansion constant at basepoint; see [Cor11, §2.2] for characterizations of this category.

Let us state a refinement of (2.9.1) $\implies$ (2.9.2) in the last Proposition.

Proposition 2.10. *Let X and Y be metric spaces. Let $\phi : X \rightarrow Y$ and assume that (2.9.1) holds for some κ and v , where v is admissible (Definition 2.4). Then for every sequence (σ_n) of positive real numbers and for every $(x_n) \in X^{\mathbf{N}}$ such that $\limsup v(|x_n|)/\sigma_n = 0$, ϕ induces a bilipschitz homeomorphism*

$$(\text{Cone}) \quad \text{Cone}_\omega(\phi, x_n, \sigma_n) : \text{Cone}_\omega(X, x_n, \sigma_n) \rightarrow \text{Cone}_\omega(Y, \phi(x_n), \sigma_n)$$

Proof. This conveniently follows from [KW14], by setting for any $\rho > 0$, $X_n = B(x_n, \rho\sigma_n)$, $t_n = v((1 + \rho)|x_n|)$ and $\phi_n = \phi|_{X_n}$. Since t/σ is infinitesimal, by [KW14, Lemma 1.16] the sequence ϕ_n defines ϕ_ω between the ultralimits of the spaces X_n/σ_n , namely, the ball of radii ρ in the asymptotic cones. $\square$

In Proposition 2.10 the assumption that v be admissible is necessary. Otherwise t_n may not be negligible when compared to σ_n , which is necessary assumption so that the sequence t_n/σ_n defines an infinitesimal number in the real field $\prod_\omega \mathbf{R}$ for every ultrafilter ω .

As an application, we can now distinguish the nonhomogeneous spaces from Examples 2.6:

- None of P_1, P_2, P_3 is $o(r)$ -bilipschitz to P_4 since $\dim \text{Cone}_\omega^\bullet(P_i)$ is 1 for $i = 1, 2, 3$ and 2 for $i = 4$.
- P_3 and P_4 are $O(\log)$ -bilipschitz through the identity map in polar coordinates, but they are not $O(\log^{1-\epsilon})$ -bilipschitz equivalent for any $\epsilon > 0$, since $\dim \text{Cone}_\omega(P_3, x_n, n) = 2$ and $\dim \text{Cone}_\omega(P_3, x_n, n) = 1$ if $|x_n| = e^n$ (See Table 1) and $\log(e^n)^{1-\epsilon} \ll n$.
- P_1 and P_2 are quasiisometric; however they are not $O(u)$ -bilipschitz equivalent for $u \rightarrow 0$.

2.3.1. On cone dimension. We have seen that the covering dimension of (moving) cones is an efficient tool to discriminate between the Examples 2.6 up to quasiisometry or $O(u)$ -bilipschitz equivalence. When X is co-boundedly acted upon, however (which is one case of interest for geometric group theorists) all its asymptotic cones are isometric once the ultrafilter is fixed. Hence, computing $\dim \text{Cone}_\omega$ for fixed ω will provide the same information with respect to QI or SBE.

	$\text{Cone}_\omega(P_1)$	$\text{Cone}_\omega(P_2)$	$\text{Cone}_\omega(P_3)$	$\text{Cone}_\omega(P_4)$
bounded $x_n, \sigma_n \equiv 1$	(P_1, x_ω)	(P_2, x_ω)	(P_3, x_ω)	(P_4, x_ω)
bounded $x_n, \sigma_n \rightarrow +\infty$	$\mathbf{R}_{\geq 0}$	$\mathbf{R}_{\geq 0}$	$\mathbf{R}_{\geq 0}$	$(C, 0)$
bounded $x_n, \sigma_n \rightarrow 0$	$\mathbf{E}^2$	$\mathbf{E}^2$	$\mathbf{E}^2$	$\mathbf{E}^2$
$ x_n = n, \sigma_n = 1/n$	$S^1 \times \mathbf{R}$	$\mathbf{E}^2$	$\mathbf{E}^2$	$\mathbf{E}^2$
$ x_n = n, \sigma_n = 1$	$\mathbf{R}$	$S^1 \times \mathbf{R}$	$\mathbf{E}^2$	$\mathbf{E}^2$
$ x_n = n, \sigma_n = n$	$\mathbf{R}_{\geq -1}$	$\mathbf{R}_{\geq -1}$	$\mathbf{R}_{\geq -1}$	(C, i)
$ x_n = e^n, \sigma_n = 1/n$	$\mathbf{R}$	$\mathbf{E}^2$	$\mathbf{E}^2$	$\mathbf{E}^2$
$ x_n = e^n, \sigma_n = 1$	$\mathbf{R}$	$S^1 \times \mathbf{R}$	$\mathbf{E}^2$	$\mathbf{E}^2$
$ x_n = e^n, \sigma_n = n$	$\mathbf{R}$	$\mathbf{R}$	$S^1 \times \mathbf{R}$	$\mathbf{E}^2$
$ x_n = e^n, \sigma_n = n^2$	$\mathbf{R}$	$\mathbf{R}$	$\mathbf{R}$	$\mathbf{E}^2$

TABLE 1. Various cones on the Riemannian planes P_i from Example 2.6. We provide the cones as pointed metric spaces (on the second line they do not depend on σ_n as soon as it goes to $+\infty$). Here C denotes $\{z \in \mathbf{C} : \Im z \geq 0\}/(x \sim -x)$ with the distance induced from the absolute value.

Beyond geometric models of polynomially growing groups G , it should not be expected that different ultrafilters will yield isometric or even just homeomorphic asymptotic cones; an extensive literature and even the notion of lacunary hyperbolic group on its own have been built over this distinction ([TV00], [KSTT05], [OOS09]). If G is a simply connected, completely solvable Lie group with a completely solvable $\mathfrak{g}$, nevertheless, then for every geometric model X , $\omega \in \beta\mathbf{N} \setminus \mathbf{N}$ and σ_n with $\lim_{\sigma_n} = +\infty$,

$$(\text{conedim}) \quad \dim \text{Cone}_\omega^\bullet(X, \sigma_n) = \dim G^{\text{nil}}$$

where G^{nil} is the largest nilpotent quotient of G [Cor08]. Following Cornulier we denote this integer conedim. This is the first, and perhaps the most natural numerical SBE invariant.

In the special case when G is nilpotent, (conedim) follows from the earlier construction of Pansu, which can be formulated in terms of Gromov-Hausdorff convergence with no reference to a ultrafilter [Pan83]. Beware that this limit is not functorial, however.

When no homogeneity assumption is made, the dimension of the asymptotic cone (even with fixed basepoint) depends not only on the ultrafilter but also on the scaling sequence. One encounters four-dimensional complete Riemannian spaces with positive Ricci curvature and $\text{SU}(2)$ symmetry, for which the covering dimension of the asymptotic cones can be 2 or 4 depending on how one chooses the scaling factors [Per97]. These cones are genuine rescaled Gromov-Hausdorff limits, obtained without passing to a subsequence and thus do not depend on the ultrafilter.

2.4. Coarse structures. In the 1930s, Weil abstracted the notion of a uniform structure from the topology of locally compact groups. Coarse structures are

large-scale counterparts of uniform structures; they were introduced by Roe in the 1990s. We recall below the definition of a coarse space.

Let X be a set. The square $X \times X$ is a groupoid for the composition law $(x_0, x_1) \circ (x_1, x_2) = (x_0, x_2)$ and $(x_0, x_1)^{-1} = (x_1, x_0)$ for $x_0, x_1, x_2 \in X$. For $E, F \subseteq X \times X$, define $E \circ F = \{e \circ f : e \in E, f \in F\}$ and $E^{-1} = \{e^{-1} : e \in E\}$.

Definition 2.11 ([Roe03, Definition 2.3]). A collection $\mathcal{E} \subseteq X \times X$ is called a coarse structure if it contains the diagonal $\Delta_{X \times X}$, is stable by composition, inverse, taking subsets, and taking finite unions; the subsets $E \in \mathcal{E}$ are called entourages.

A coarse structure $\mathcal{E}$ is called monogenic if it is generated by a single entourage, that is if there exists $E \in \mathcal{E}$ such that $\mathcal{E}$ is smallest among all coarse structures containing E . Note that this notion has no analog among uniform structures.

Definition 2.12 (Coarse equivalence). Given two coarse spaces $(X, \mathcal{E}_X)$ and $(Y, \mathcal{E}_Y)$ and a map $\phi : X \rightarrow Y$, we say that ϕ is coarse if

$$(2.12.1) \text{ for all } B \subseteq Y, B \times B \in \mathcal{E}_Y \implies \phi^{-1}(B) \times \phi^{-1}(B) \in \mathcal{E}_X$$

$$(2.12.2) \text{ for all } E \in \mathcal{E}_X, (\phi \times \phi)(E) \in \mathcal{E}_Y, \text{ where } \phi \times \phi(x, y) = (\phi(x), \phi(y)).$$

A pair of coarse maps $\{\phi : X \rightarrow Y, \psi : Y \rightarrow X\}$ realizes a coarse equivalence if the graphs of $\phi \circ \psi$ and $\psi \circ \phi$ are both contained in entourages of the coarse structures.

Proposition 2.13 ($O(u)$ -coarse structure, $o(v)$ -coarse structure). *Let $u : [0, +\infty) \rightarrow (0, +\infty)$ be an admissible function, let v be either an admissible function or $v(r) = r$, and let (X, d_X) be a metric space. Given some $o \in X$, define*

$$(2.3) \quad \mathcal{E}^{O(u)} = \left\{ E \subseteq X \times X : \exists M, \limsup_{(x, x') \in E} \frac{d_X(x, x')}{u(|x|)} \leq M \right\}$$

$$(2.4) \quad \mathcal{E}^{o(v)} = \left\{ E \subseteq X \times X : \limsup_{(x, x') \in E} \frac{d_X(x, x')}{v(|x|)} = 0 \right\}$$

where $|x| = d_X(o, x)$ and $\limsup$ are taken as (x, x') evades every bounded set fixed in advance (for the sup distance in $X \times X$). $\mathcal{E}^{O(u)}$ and $\mathcal{E}^{o(v)}$ define coarse structures on X .

The bounded coarse structure is $\mathcal{E}_X^{O(1)}$, and the coarse equivalences between metric spaces equipped with $\mathcal{E}_X^{O(1)}$ are the coarse equivalences as defined in (2.1). Wright's c_0 coarse structure is $\mathcal{E}^{o(1)}$ [Wri03, Definition 1.1]. Dranishnikov and Smith's sublinear coarse structure is $\mathcal{E}^{o(r)}$ (See §2.5) [DS07].

Proof. We need to check Roe's axioms. In view of (2.3) and (2.4) it is clear that $\mathcal{E}_X^{O(u)}$ and $\mathcal{E}_X^{o(v)}$ are closed under finite union and taking subsets. Possibly left nonobvious is the stability when taking inverses and composing.

Inverses. Fix a basepoint o and take a sequence x_n, x'_n such that $\sup(|x_n|, |x'_n|) \rightarrow +\infty$, with $d_X(x_n, x'_n) \leq Ku(|x_n|)$ for some $K \geq 0$ when n is large enough, resp. $d_X(x_n, x'_n) \leq k_n v(|x_n|)$ where $k_n \rightarrow 0$. We need prove that $d_X(x_n, x'_n) \leq Lu(|x'_n|)$ for some $L \geq 0$, resp. $d_X(x_n, x'_n) \leq \ell_n v(|x'_n|)$ for some $L \geq 0$ when n is large enough.

We claim that

$$(2.5) \quad 0 < \liminf \frac{|x'_n|}{|x_n|} \leq \limsup \frac{|x'_n|}{|x_n|} < +\infty.$$

Indeed, if it were not the case there would be a sequence R_n such that for arbitrarily large values of D , either for arbitrarily large n , $|x_n| \leq R_n \leq DR_n \leq |x'_n|$ or for arbitrarily large n , $|x'_n| \leq R_n \leq DR_n \leq |x_n|$. In the first case, along a sub-sequence, by the triangle inequality $|x'_n| \leq R_n + Ku(R_n)$ (where we may replace u by v and K by some k_{n_0} if necessary) contradicting the hypothesis that $|x'_n| \geq DR_n$ for n large enough (observe that $|x'_n| \rightarrow +\infty$ along that sub-sequence). In the second case, again by the triangle inequality one would have $R_n \geq |x_n| - Ku(|x_n|)$ (or $|x_n| - k_{n_0}|x_n|$ if necessary); but the right-hand side can be assumed greater than $|x_n|/2$ for n large enough if D is set large enough; this is a contradiction. Now from (2.5) and the property that u , resp. v is admissible, we obtain that also

$$0 < \liminf \frac{|u(x'_n)|}{|u(x_n)|} \leq \limsup \frac{|u(x'_n)|}{|u(x_n)|} < +\infty$$

(resp. the same with v replacing u), which provides the requested constant L (resp. ℓ_n) as a function of K (resp. of k_n) and u , resp. v . At this point it is useful to record that we can rewrite $\mathcal{E}$ in a more symmetric way:

$$\mathcal{E}_X^{O(u)} = \left\{ E \subseteq X \times X : \exists r > 0, \sup_{(x,x') \in E \setminus B_r(o) \times B_r(o)} d_X(x,x')/(u(|x|) + u(|x'|)) < +\infty \right\}$$

Composition. Start assuming u nondecreasing; we will explain how to adapt the proof in case it is not the case in the end (this philosophy was alluded to after Lemma 2.5). For every $K, r \geq 0$, introduce

$$E_K^r(X, o) = \{(x, x') : \inf(|x|, |x'|) \geq r, d_X(x, x') \leq K(u(|x|) + u(|x'|))\}.$$

We need to prove that for every K, L there are r, s, t and $\eta(K, L)$ such that

$$(2.6) \quad E_L^s \circ E_K^r \subseteq E_{\eta(K,L)}^t.$$

Let $(x, x'') \in E_L \circ E_K$. By definition, there exists $x' \in X$ such that $d_X(x, x') \leq K(u(|x|) + u(|x'|))$ and $d_X(x', x'') \leq L(u(|x'|) + u(|x''|))$.

Set a radius $R = \sup \{r \geq 0 : u(r) > r/(2K+1)\}$. We claim that

$$(2.7) \quad u(|x'|) \leq \sup(u(3R), u(3|x|))$$

To prove (2.7) we proceed by exhausting all the case arising from the comparison of $|x|$ and $|x'|$ with R .

First, note that either $|x'| \leq R$, or $|x'| > R$ and then $u(|x'|) \leq \frac{|x'|}{2K+1}$. In the second case, by the triangle inequality

$$|x'| \leq |x| + Ku(|x|) + Ku(|x'|) \leq |x| + Ku(|x|) + \frac{|x'|}{2},$$

so that $|x'| \leq 2|x| + 2Ku(|x|)$. So we always have $|x'| \leq \sup(R, 2|x| + 2Ku(|x|))$. Since u has been assumed nondecreasing,

$$u(|x'|) \leq \sup(u(R), u(2|x| + 2Ku(|x|))).$$




FIGURE 2. Some entourages of the $O(u)$ -coarse structure on the half real line $X = [0, +\infty)$, with $u(r) = \sqrt{r}$, $u(r) = 1$ and $u(r) = 1/r$.

Now, either $|x| \leq R$, in which case $u(|x'|) \leq \sup(u(R), u(3|x|))$ and (2.7) holds, or $|x| > R$ and then $2Ku(|x|) \leq |x|$, so $u(|x'|) \leq K \sup(u(3R), u(3|x|))$: (2.7) holds as well. We can now finish the proof using the claim. By the triangle inequality,

$$\begin{aligned} d_X(x, x'') &\leq Ku(|x|) + (K + L)u(|x'|) + Lu(|x''|) \\ &\leq (K + L)[u(|x|) + \sup(u(3R), u(3|x|)) + u(|x''|)] \end{aligned}$$

so we may set $\eta(K, L) = 2(K + L) \limsup_{r \rightarrow +\infty} u(3r)/u(r)$; then for r large enough and arbitrary s , (2.6) holds.

We now resume to the general case when u is not assumed non-decreasing. If $|x'| \leq R$ then there is a uniform bound on $|x|$. If $|x'| > R$ then by the triangle inequality,

$$|x'| \geq |x| - Ku(|x|) - Ku(|x'|) \geq |x| - \frac{|x'|}{2} - Ku(|x|),$$

so that $|x'| \geq 2|x|/3 - 2Ku(|x|)/3$. As soon as $|x| \geq R$, $|x'| \geq |x|/3$. Using the assumption that u is admissible, then, $u(|x'|) \leq Bu(|x|)$ for some $B \geq 1$. Using the same line of reasoning as before, this implies (2.6) with $\eta(K, L) = B(K + L)$. $\square$

Let u_1, u_2 and v be as u and v in the previous proposition. Let X and Y be metric spaces. If $u_1 = O(u_2)$, resp. if $u = o(v)$, then the identity map $(X, \mathcal{E}^{O(u_1)}) \rightarrow (X, \mathcal{E}^{O(u_2)})$ is coarse. Especially, if $\varphi : (X, d_X, \mathcal{E}^{O(u_1)}) \rightarrow (Y, d_Y, \mathcal{E}^{O(u_1)})$ is a coarse equivalence, then $\varphi : (X, d_X, \mathcal{E}^{O(u_2)}) \rightarrow (Y, d_Y, \mathcal{E}^{O(u_2)})$ is a coarse equivalence, as summarized in the diagram below where the arrows are coarse, $u_1 = 1$, $u_2 = \log$ and $v(r) = r$, namely the three most important coarse structures in this paper.

$$\begin{array}{ccccc} (X, \mathcal{E}^{O(1)}) & \xrightarrow{\text{id}_X} & (X, \mathcal{E}^{O(\log)}) & \xrightarrow{\text{id}_X} & (X, \mathcal{E}^{O(r)}) \\ \downarrow \phi & & \downarrow \phi & & \downarrow \phi \\ (Y, \mathcal{E}^{O(1)}) & \xrightarrow{\text{id}_Y} & (Y, \mathcal{E}^{O(\log)}) & \xrightarrow{\text{id}_Y} & (Y, \mathcal{E}^{O(r)}) \end{array}$$

Proposition 2.14. *Let X and Y be metric spaces. Let $u : [0, +\infty) \rightarrow (0, +\infty)$ be a regularly varying function. Let $\phi : X \rightarrow Y$ be a $O(u)$ -bilipschitz equivalence. Then ϕ induces a coarse equivalence $(X, d_X, \mathcal{E}^{O(u)}) \rightarrow (Y, d_Y, \mathcal{E}^{O(u)})$.*

Proof. Let (x_n, x'_n) be a sequence of points with $d(x_n, x'_n) \leq Mu(|x_n|)$ and $|x_n| \rightarrow +\infty$. Then, for n large enough, $|x'_n| \leq 2|x_n|$. Hence

$$d(\phi(x_n), \phi(x'_n)) \leq \kappa Mu(|x_n|) + cu(|x_n| \vee |x'_n|) \leq (\kappa M + c)u(|x_n|)$$

for some $C \geq 1$. But also, for n large enough,

$$(2.8) \quad |\phi(x_n)| \geq |x_n|/(2\kappa).$$

So there exists a constant C' so that $d(\phi(x_n), \phi(x'_n)) \leq C'u(|\phi(x_n)|)$. On the other hand, ϕ has axiom (2.12.1) by (2.8). This proves that ϕ is a coarse map. ϕ has a coarse inverse $\tilde{\phi}$ such that $d(\tilde{\phi} \circ \phi(x), x) \leq c'u(|x|) + c'$ for all $x \in X$ and $d(\phi \circ \tilde{\phi}(y), y) \leq c'u(|y|) + c'$ for all $y \in Y$ [Cor19]. So ϕ is a coarse equivalence. $\square$

Lemma 2.15. *Assume that (X, d_X) is a geodesic metric space. Let u be admissible and unbounded. Then $E_u = \{(x, x') \in X \times X : d_X(x, x') \leq 1 + u(|x| + |x'|)\}$ is a symmetric entourage generating $\mathcal{E}^{O(u)}$ on X . Define $\hat{d}_X$ on X such that*

$$\hat{d}(x, x') = \inf \{n : (x, x') \in E_u^n\}.$$

Then, the identity map $(X, d_X, \mathcal{E}^{O(u)}) \rightarrow (X, \hat{d}_X, \mathcal{E}^{O(1)})$ is a coarse equivalence.

Proof. Let us check first that E_u generates $\mathcal{E}$. Take $E \in \mathcal{E}^{O(u)}$; then by definition

$$\sup \frac{d_X(x, x')}{1 + u(|x|) + u(|x'|)} = M < +\infty.$$

For all (x, x') , and for every segment $\gamma : [0, d_X(x, x')] \rightarrow X$ and set $x_1 = \gamma(1 + u(|x|))$, $x_2 = \gamma(2 + u(|x|) + u(|x_1|))$, ... Let

$$N_\gamma(x, x') = \inf \{n : n + u(|x|) + \dots + u(|x_n|) > d_X(x, x')\}.$$

We claim that $\sup_{(x, x') \in E} \inf_\gamma N < +\infty$. Indeed, if x and x' are far enough there exists some constant $\mu > 0$ such that $u(|x_k|) \geq \mu u(|x|)$ as long as $|x_k| \geq |x|/2$, especially as long as $k + u(|x|) + \dots + u(|x_k|) \leq |x|/2$. So either $N(x, x') \leq \lceil M/\mu \rceil$ or $N + u(|x|) + \dots + u(|x_N|) > |x|/2$. But in the latter case,

$$(2.9) \quad M(1 + u(|x|) + u(|x'|)) \geq d_X(x, x') > \frac{|x|}{2} - 1 - u(|x_N|)$$

where we used the definition of N on the right. To reach a contradiction, note that again by the definition of N , $d(x_N, x') < 1 + u(|x_N|)$, so there exists L such that $d(x_N, x') \leq 1 + Lu(|x'|)$, reproducing the reasoning in the “Inverse” part of the proof of Proposition 2.13. Hence, there exists some constant M' such that if x' is far enough, $u(|x_N|) \leq M'u(|x'|)$. Plugging this in (2.9) yields an inequality of the form $u(|x'|) + u(|x|) \geq \rho|x|$ for some $\rho > 0$, which can only occur if $|x|$ is close to the origin. We conclude that $E \subseteq E_u^{N_{\max}}$, where $N_{\max} = \sup_{(x, x') \in E} \inf_\gamma N$ is a finite integer.

This proves that $(X, d_X, \mathcal{E}^{O(u)}) \rightarrow (X, \hat{d}_X, \mathcal{E}^{O(1)})$ has the axiom (2.12.2) of a coarse map. In order to check (2.12.1) we must prove that if $B \times B$ is in $\mathcal{E}^{O(u)}$ then B is bounded; fixing $x \in B$, by (2.3), for any sequence x'_n that escape to infinity x'_n cannot stay in any entourage of $\mathcal{E}^{O(u)}$ fixed in advance. Conversely, if B is bounded then $B \times B$ is in $\mathcal{E}^{O(u)}$, while axiom (C2) holds for $(X, \hat{d}_X, \mathcal{E}^{O(1)}) \rightarrow (X, d_X, \mathcal{E}^{O(u)})$ by definition of $\hat{d}$. $\square$

The new distance $\widehat{d}$ may be made geodesic as well, by adding metric edges between pairs of point at distance 1. Note however that one may loose properness in this process. Isometric group actions are also lost when passing to $\widehat{d}$, and in fact its main interest is theoretical, and appears in the next Proposition.

Say that a map $\phi : X \rightarrow Y$ between pointed metric spaces is radial if there exists $\kappa \geq 1$ and $R, R' \geq 0$ such that for all $x \in X$,

$$(2.10) \quad \frac{1}{2\kappa} \sup(R, |x|) \leq \sup(R', |\phi(x)|) \leq 2\kappa \sup(R, |x|).$$

Also, call discrete geodesic between x and x' at distance n in X a finite sequence of points x_i with $x = x_0$, $x_n = x'$ and $d(x_i, x_{i+1}) = 1$.

Proposition 2.16. *Let X and Y be geodesic metric spaces, and let $\phi : X \rightarrow Y$ be a $O(\log)$ -coarse equivalence. Then*

- (1) ϕ is radial.
- (2) ϕ is a $O(\log)$ -bilipschitz equivalence.

We need a preliminary Lemma.

Lemma 2.17. *Let t and s be positive real numbers. Then for every $M > 0$, there exists $R \geq 1$ and $M' > 0$ such that*

$$\begin{cases} \frac{t}{\log t} & \leq M \frac{s}{\log s} \\ \inf(s, t) & \geq R \end{cases} \implies t \leq M's$$

Proof. We will prove first a weaker inequality and then self-improve it. Taking logarithms on both sides we get $\log t - \log \log t \leq \log M + \log s - \log \log s$, so for every $\varepsilon > 0$ one has, for s and t large enough, $(1 - \varepsilon/2) \log t \leq (1 + \varepsilon/2) \log s$, and then $t \leq s^{1+\varepsilon}$. Now, assume by contradiction that there is a sequence (s_n, t_n) with $t_n / \log t_n \leq M s_n / \log s_n$, but $q_n = t_n / s_n$ going to infinity. Then $t_n / \log t_n = t_n / (\log s_n + \log q_n)$; but we know that $\log q_n \leq \varepsilon \log s_n$; so $t_n / \log s_n \leq M' s_n / \log s_n$ for some M' , reaching the desired inequality. $\square$

Proof of the Proposition 2.16. Consider the metrics $\widehat{d}_X$ and $\widehat{d}_Y$ provided by Lemma on X and Y . Then $\phi : (X, \widehat{d}_X) \rightarrow (Y, \widehat{d}_Y)$ becomes a $O(1)$ -coarse equivalence. Since $\widehat{d}_X$ and $\widehat{d}_Y$ are geodesic, ϕ is a $\widehat{d}$ -quasiisometry, especially it is $\widehat{d}$ -radial.

Now, we need to compare $\widehat{d}$ and d . Start with (1); for this we need to compare $|x|$ and $\widehat{d}(0, x)$ for all $x \in X$. Let (x_n) be a discrete $\widehat{d}$ -geodesic segment from o (we do not specify an endpoint yet). We claim that $|x_n| \leq 2n \log n + 2n$ for $n > 0$. Let us proceed by induction on n . This holds for $n = 1$. Assume it holds for some $n > 0$. Then,

$$\begin{aligned} |x_{n+1}| &= |x_n| + d(x_n, x_{n+1}) \\ &\leq |x_n| + 1 + \log(|x_n|) \\ &\leq 2n + 2n \log n + 1 + \log 2 + \log n + \log(1 + \log n) \\ &\leq 2n + 2n \log n + 2 + 2 \log n \\ &= (2n + 2) + (2n + 2) \log n \leq (2n + 2) + (2n + 2) \log(n + 1) \end{aligned}$$

where we used $\log 2 < 1$ and $\log n \leq n - 1$. Using this inequality, we deduce

$$(2.11) \quad \widehat{d}_X(o, x) \geq \inf \{n : 2n(1 + \log n) \geq |x|\} \geq \frac{|x|}{1 + 3 \log |x|}$$

Conversely, repeating a construction made in the proof Lemma 2.15, consider a geodesic segment $\gamma : [0, |x|] \rightarrow X$, and a sequence

$$x_0 = o, x_1 = \gamma(2), x_2 = \gamma(1 + \log |x_1|), \dots, x_{i+1} = \gamma(|x_i| + \log |x_i|)$$

and define N such that x_N is the farthest element from o before reaching x ; in this way, $\widehat{d}_X(o, x) \leq N + 1$. By induction on n , we can prove that $|x_n| \geq n \log n$ for all n . So

$$(2.12) \quad \widehat{d}_X(o, x) \leq 1 + \inf \{n : n \log n \geq |x|\} \leq 1 + \frac{|x|}{1 + \log |x|}.$$

We are now ready to prove (1). We know that ϕ is $(\widehat{d}_X, \widehat{d}_Y)$ -radial ; so there exists κ_0 such that

$$(2.13) \quad \frac{|\phi(x)|}{1 + 3 \log |\phi(x)|} \leq \widehat{d}_Y(o, \phi(x)) \leq 2\kappa_0 \left(1 + \frac{|x|}{1 + \log |x|}\right)$$

Combining both inequality, $|\phi(x)|$ and $|x|$ satisfy the hypotheses of t and s in Lemma 2.17. We conclude from the Lemma that ϕ is radial.

The proof of (2) will now rely on (1) together with an estimate akin to (2.11) and (2.12), but where we replace o with $x' \in X$. Let $x, x' \in X$; assume $2 \leq |x| \leq |x'|$, and let γ be a geodesic segment from x to x' . Define $x_0 = x$, $x_{i+1} = \gamma(d(x_0, x_i) + 1 + \log |x_i|)$ as long as it makes sense (let n be the largest one, so that x_n is the closest to x' among all x_i 's). By the triangle inequality, for all i such that $0 \leq i \leq n$,

$$|x_i| \leq |x'| + d(x', x_i) \leq |x| + d(x, x') \leq 2|x'| + |x| \leq 3|x'|.$$

From this inequality, we deduce that

$$\widehat{d}_X(x, x') \geq \text{long}(\gamma)/(2 \log |3x|) \geq \frac{d(x, x')}{4 \log |x'|}$$

Conversely, if $\inf_t |\gamma(t)| \leq |x'|/2$, then $d(x, x') \geq |x'|/2$. So


$$\widehat{d}(x, x') \leq \widehat{d}(x, o) + \widehat{d}(o, x') \leq 2 + \frac{2|x'|}{1 + \log |x'|} \leq 2 + \frac{4d(x, x')}{1 + \log |x'|}.$$

Otherwise, $\inf_t |\gamma(t)| > |x'|/2$, and then $\widehat{d}(x, x') \leq \frac{d(x, x')}{\log(|x'|/2)}$. Combining the previous inequalities, we get that for every pair x, x' with $\sup(|x|, |x'|)$ large enough,

$$(2.14) \quad \frac{1}{\lambda_X} \frac{d_X(x, x')}{\log(\sup(|x|, |x'|))} \leq \widehat{d}_X(x, x') \leq \lambda \frac{d_X(x, x')}{\log(\sup(|x|, |x'|))}$$

for some $\lambda_X > 1$. A similar inequality holds for pairs of points in Y , with a multiplicative factor λ_Y . We are ready to finish the proof. Assume that ϕ is a (κ_0, c_0) quasiisometry with respect to $\widehat{d}_X$ and $\widehat{d}_Y$. Then

$$-c_0 + \frac{1}{\kappa_0} \widehat{d}_X(x, x') \leq \widehat{d}(\phi(x), \phi(x')) \leq \kappa_0 \widehat{d}_X(x, x') + c_0$$

FIGURE 3. Coronae and Gromov boundary for hyperbolic X .

for all x, x' . So, setting $\lambda = \sup(\lambda_X, \lambda_Y)$ and using (2.14) and its counterpart in Y ,

$$-c_0 + \frac{1}{\lambda^2 \kappa_0} \frac{d_X(x, x')}{\log \sup(|x|, |x'|)} \leq \frac{d_Y(\phi(x), \phi(x'))}{\log(\sup(|\phi(x)|, |\phi(x')|))} \leq \lambda^2 \kappa_0 \frac{d_X(x, x')}{\log \sup(|x|, |x'|)} + c_0$$

Using that ϕ is radial, we know that $|\phi(x)|$ and $|\phi(x')|$ are within linear control from $|x|$ and $|x'|$. So we may rewrite the previous estimate as

$$-c_1 + \frac{1}{\kappa_1} \frac{d_X(x, x')}{\log \sup(|x|, |x'|)} \leq \frac{d_Y(\phi(x), \phi(x'))}{\log(\sup(|x|, |x'|))} \leq \kappa_1 \frac{d_X(x, x')}{\log \sup(|x|, |x'|)} + c_1$$

where $\kappa_1 \geq 1$ and $c_1 \geq 0$. Multiplying by $\log \sup(|x|, |x'|)$ on both sides yields the required (1.1). $\square$

Remark 2.18. The assumption $u = \log$ made in Proposition 2.16 is possibly too strong. On the other hand, it is not true that every coarse equivalence between $o(r)$ -coarse structure is a $o(r)$ -bilipschitz equivalence: consider $\phi : \mathbf{R}^n \rightarrow \mathbf{R}^n$ such that $\phi(x) = \|x\|x$. A notable distinction between $\mathcal{E}^{O(\log)}$ and $\mathcal{E}^{o(r)}$ is that the former is monogenic whereas the latter is not. Also, observe that Lemma 2.17 breaks down for $u(t) = t^e$, $e > 0$.

2.5. Invariance of the geometric dimension for connected Lie groups.

Definition 2.19 (sublinear Higson function). Let X be a proper metric space. Define the C^* -algebra $C_{h_L}(X)$ of sublinear Higson functions on X as

$$\left\{ f \in C_b(X, \mathbf{C}) : \forall E \in \mathcal{E}^{o(r)}, \lim_{r \rightarrow +\infty} \sup_{(x, x') \in E, \inf(|x|, |x'|) \geq r} |df(x, x')| = 0 \right\}$$

where $f \in C_b$ means $\sup |f| < +\infty$ and $df(x, x') = f(x) - f(x')$.

Remark 2.20 (Compare Fukaya [Fuk12], 3.1). f is Higson sublinear if and only if there exists $C_f < +\infty$ such that for all x, x' in X and $R > 0$ large enough, if $\inf(|x|, |x'|) \geq R$ and $d_X(x, x') \leq R/2$, then $|f(x) - f(x')| \leq \frac{C_f}{R}$.

The closure $\overline{C_{h_L}(X)}$ is a unital C^* algebra; once modded out by the ideal of functions vanishing at infinity, this algebra is the spectrum of a compact space, the sublinear Higson corona $\nu_L X$ of X [Roe03, Definition 2.37].

Remark 2.21 (See Figure 3). If X is a Gromov-hyperbolic space, then the Gromov functions on X are Higson sublinear. The Higson sublinear functions are Higson. It follows that the sublinear Higson corona sits in between the Higson corona νX and the Gromov boundary $\partial_\infty X$ seen in the topological category.

The following is a generalization of [DS07, Proposition 2.1].

Proposition 2.22. *Let X and Y be metric spaces. Let $\nu_L X$ and $\nu_L Y$ be their sublinear Higson coronae. Then, any $o(r)$ -bilipschitz equivalence $f : X \rightarrow Y$ induces a homeomorphism $\nu_L f : \nu_L X \rightarrow \nu_L Y$.*

Proof. By proposition 2.14, a $o(r)$ -bilipschitz equivalence $X \rightarrow Y$ represents a coarse equivalence $(X, d_X, \mathcal{E}^{o(r)}) \rightarrow (Y, d_Y, \mathcal{E}^{o(r)})$, and then induces a homeomorphism between the Higson coronae [Roe03, Corollary 2.42]. $\square$

Theorem 2.23 ([DS07, Theorem 3.10 and Corollary 3.11]; see also [CDSV11]). *Let X be a proper connected metric space. Assume that $\text{Isom}(X)$ is co-compact on X , and that $\text{asdim}_{\text{AN}}(X) < +\infty$. Then*

$$(2.15) \quad \dim \nu_L X = \text{asdim}_{\text{AN}}(X).$$

Theorem 2.24 ([HP13, Theorem 7.9]). *Let G be a connected Lie group, and let X be any geometric model of G . Then*

$$(2.16) \quad \text{asdim}_{\text{AN}}(X) = \dim G - \dim K.$$

where K is any maximal compact subgroup of G .

Theorem B from the introduction now follows by combining 2.22 with Theorems 2.23 and 2.24.

To the best of the author's knowledge, the only connected Lie group for which some description of the sublinear Higson corona is currently available is $\mathbf{R}^n$: Fukaya proved that $\nu_L \mathbf{R}^n \simeq S^{n-1} \times \nu_L \mathbf{R}$ [Fuk12]. These spaces are “big” and not metrizable, so it seems not easy to extract fine topological invariants from them as one would do for, say, the Gromov boundary.

Question 2.25. Let X be a proper metric space. Is the Čech cohomology group $\check{H}^1(\nu_L X, \mathbf{Q})$ finitely generated?

The answer is known to be negative for the Higson coronae associated to bounded coarse structures [Kee94]; nevertheless Fukaya proves that $\nu_L \phi$ is homotopic to the identity whenever $\phi \in \text{GL}(n, \mathbf{R})$ has positive determinant.

3. REAL HYPERBOLIC SPACES AND THEOREM C

In this section we prove Theorem C on Lie groups $O(u)$ -bilipschitz equivalent to real hyperbolic spaces. §3.1 gathers preliminary results on pinching and conformal dimension, and §3.2 sets the terminology of degenerations and deformations. The equivalences of Theorem C are proved in §3.3.

3.1. Heintze groups, conformal dimension and pinching. In 1955, Jacobson proved that all real Lie algebras who possess a derivation with no purely imaginary eigenvalue are nilpotent [Jac55]. Later Heintze characterized the semidirect products of nilpotent Lie algebra by derivations whose spectrum has positive real part, as the Lie algebras of Lie groups that possess at least one negatively curved left-invariant metric (note that these are centerless) [Hei74]. Most importantly, Heintze showed that the negatively curved metrics on these groups exhaust all the isometrically homogeneous negatively curved manifolds, shedding light on the earlier result of Kobayashi that these spaces had to be simply connected [Kob62].

Definition 3.1 ([CT11]). Let G be a Lie group with finitely many components. Then G is of Heintze type if there exists a simply connected nilpotent N , a derivation $\alpha \in \text{Der}(\mathfrak{n})$ with $\inf \{\Re \lambda : \lambda \in \text{Sp}(\alpha)\} > 0$ and a compact group K with a representation $\rho : K \rightarrow \text{Aut}(N)$ such that

$$(3.1) \quad G = (K \times \mathbf{R}) \ltimes N,$$

where $(k, t).n = \rho(k)(n)e^{\alpha t}n$. A Heintze group is a group of Heintze type with $K = 1$.

By normalized Jordan form of a derivation α as in Definition 3.1, we mean the Jordan form of the unique positive multiple $[\alpha]$ of α such that

$$(3.2) \quad \inf \{\Re \lambda : \lambda \in \text{Spec}([\alpha])\} = 1.$$

Note that $N \rtimes_{\alpha} \mathbf{R} \simeq N \rtimes_{[\alpha]} \mathbf{R}$ (Compare Example 3.12.) The following useful fact is proved in E. Sequeira's thesis using a highest weight argument [SM20, Proposition 5.2.2]⁵.

Proposition 3.2. *Let N be a simply nilpotent Lie group. If the Heintze groups $G_{\alpha} = N \rtimes_{\alpha} \mathbf{R}$ and $G_{\beta} = N \rtimes_{\beta} \mathbf{R}$ are isomorphic, then α and β have the same normalized Jordan form.*

Definition 3.3 (after [EH96, Section 4]). Given two Heintze groups $G = N \rtimes_{\alpha} \mathbf{R}$ and $G' = N' \rtimes_{\alpha'} \mathbf{R}$ and $\lambda > 0$, we write $G \sharp (G')^{\lambda} = (N \times N') \rtimes \mathbf{R}$ where $t.n = (e^{\alpha t}, e^{\lambda \alpha' t})$ with the convention that both α and α' are normalized as in (3.2), and call this group Heintze amalgam of G and G' . The Lie algebra $\text{Lie}(G \sharp (G')^{\lambda})$ we denote $\mathfrak{g} \sharp \lambda \mathfrak{g}'$.

A Heintze group is purely real if it is completely solvable, i.e. if $\text{Sp}(\alpha) \subseteq \mathbf{R}$; every group of Heintze type has a Riemannian model in common with a purely real Heintze group, that we call its shadow (See [Ale75] and §5.1). If G , N , α are as in Definition 3.1 with $K = 1$ and if $\mathfrak{n} = \text{Liespan}(\ker([\alpha] - 1))$, then we say that G , resp. $\mathfrak{g}$ is a Carnot-type Heintze group, resp. algebra. In this case isomorphism type of G does not depend on α , so we abbreviate $G = N \rtimes_{\text{Carnot}} \mathbf{R}$ [Cor16, Proposition 3.5]. Carnot-type Heintze groups are purely real.

⁵[SM20] has the assumption that G_{α} and G_{β} are purely real, but the general proof goes along the same lines.

Example 3.4. Let $\mathbf{K}$ be a division algebra over $\mathbf{R}$ and n a positive integer, $n = 2$ if $\mathbf{K} = \mathbf{Ca}$. $\mathfrak{b}(n, \mathbf{K})$ is the solvable Lie algebra over the vector space $V = \mathbf{K}^{n-1} \oplus \Im \mathbf{K} \oplus \mathbf{R}$ (where $\Im \mathbf{K} = 0$ if $\mathbf{K} = \mathbf{R}$) with Lie bracket

$$[(z_i, \tau, s), (z'_i, \tau', s')] = \left[sz'_i - s'z_i, 2s\tau' - 2s'\tau + \sum_{i=1}^{n-1} \Im(z_i \overline{z'_i}), 0 \right].$$

$\mathfrak{b}(n, \mathbf{K})$ for $\mathbf{K} = \mathbf{R}, \mathbf{C}, \mathbf{H}$ is the maximal completely solvable subalgebra of $\mathfrak{o}(n, 1)$, $\mathfrak{u}(n, 1)$, $\mathfrak{sp}(n, 1)$ respectively.

The Heintze groups with Lie algebra $\mathfrak{b}(n, \mathbf{K})$ are exactly those who carry (rank one) symmetric metrics [Hei74] (for $\mathbf{K} = \mathbf{R}$, all the left-invariant metrics are symmetric, see e.g. [Lau03]).

The topological dimension $\text{Topdim } \partial_\infty$ and conformal dimension $\text{Cdim } \partial_\infty$ are quasisometry invariant of Gromov-hyperbolic locally compact compactly generated groups ([MT10], [CCMT15]). For a group of Heintze type $G = (K \times \mathbf{R}) \ltimes_\alpha N$,

$$(3.3) \quad \text{Topdim } \partial_\infty G = \dim G - \dim K - 1 = \text{geodim } G - 1;$$

$$(3.4) \quad \text{Cdim } \partial_\infty G = \text{Tr}[\alpha].$$

Though not explicitly stated there, the following is a direct consequence of [Pan89a, Section 5].

Theorem 3.5 (After Pansu). *Let (M, g) be a complete, simply connected Riemannian manifold of dimension $n \geq 2$. Let $b \geq 1$. Assume that M is $-1/b^2$ -pinched, i.e. (up to normalization of g) $-b^2 \leq K_g \leq -1$. Then*

$$(3.5) \quad \text{Cdim } \partial_\infty M \leq (n-1)b.$$

Proof. It follows from the lower bound on sectional curvature that $\text{Ric} \geq (n-1)b^2g$. Then, by the Bishop-Gromov inequality

$$\text{vol}(B(x, r)) \leq \text{cst.} \int_0^r \sinh^{n-1}(bt) dt,$$

so that the volume-theoretic entropy $h = \limsup_{r \rightarrow +\infty} r^{-1} \log \text{vol}(B(x, r))$ is bounded above by $(n-1)b$. Pansu proves $\text{Cdim } \partial_\infty S \leq h$ [Pan89a, Lemme 5.2]. Combining these inequalities yields the desired (3.5). $\square$

Corollary 3.6. *Let G be a group of Heintze type; then every Riemannian model of G has a pinching of at least*

$$(3.6) \quad - \left(\frac{\text{geodim } G - 1}{\text{Tr}[\alpha]} \right)^2.$$

The bound (3.6) is not optimal. Building on a theorem of Belegradek and Kapovitch and curvature computations, Healy determined the exact optimal pinching (which is attained) when G is Carnot-type and N has a lattice (equivalently, when $\mathfrak{n}$ has a $\mathbf{Q}$ -form) and found an optimal pinching of $-1/s^2$, where s is the nilpotency step of N [Hea21, Theorem 4.3]. Note that for Carnot type groups, s is the spectral radius of $[\alpha]$ so $\text{Tr}[\alpha] \leq s(\text{Topdim } \partial_\infty G) = s(\text{geodim } G - 1)$.

Corollary 3.7. *Let G be a group of Heintze type. Assume that G has Riemannian models with pinching arbitrarily close to -1 . Then α has all its eigenvalues with the same real part, and N is abelian.*

Proof. Order the eigenvalues of α as $\sigma_1 \leq \dots \leq \sigma_r$. Pansu's theorem forces the equality in

$$\sigma \dim \mathfrak{n} = \sum_{\lambda} \Re \lambda = \operatorname{tr}(\alpha).$$

Denote by $\mathfrak{n}^\lambda$ the generalized eigenspace of α with eigenvalue λ , observe that $[\mathfrak{n}^\lambda, \mathfrak{n}^\mu]$ for any λ, μ . Consequently, if $\bigoplus_{\tau \in \mathbf{R}} \mathfrak{n}^{\sigma+i\tau} = \mathfrak{n}$ for a given positive σ then $[\mathfrak{n}, \mathfrak{n}] \subseteq \bigoplus_{\tau \in \mathbf{R}} \mathfrak{n}^{\sigma+2i\tau}$, and N is abelian. $\square$

Remark 3.8. The conclusion that N is abelian remains if a single left-invariant metric on S is assumed to be strictly more than quarter-pinched, a theorem by Eberlein and Heber, who also characterized the Heintze groups with a quarter-pinched Riemannian metric [EH96].

We note that the converse of Corollary 3.7 also holds.

Proposition 3.9. *Let $S = \mathbf{R}^{n-1} \rtimes_{\alpha} \mathbf{R}$, where $\operatorname{sp}(\alpha) \subseteq \{1 + i\tau : \tau \in \mathbf{R}\}$. Then, S has left invariant Riemannian metrics with pinching arbitrarily close to -1 . Moreover, if K is a compact group of automorphisms of S , then one can assume that those metrics are all K -invariant.*

Proof. Let $\varepsilon > 0$ be a parameter. Let $(e_1, \dots, e_{n-1})$ be a basis of $\mathbf{R}^{n-1}$ in which α appears in real Jordan normal form; group the generalized eigenspaces with non-real eigenvalue first (if there are any), so that there is m such that in the basis

$$\mathcal{F}_{\varepsilon} = (e_1, e_2, \varepsilon e_3, \varepsilon e_4 \dots \varepsilon^{m-1} e_{2m-1}, \varepsilon^{m-1} e_{2m}, e_{2m+1}, \dots, \varepsilon^{n-2m-1} e_n),$$

α has a block upper triangular form with blocks of the form

$$J'_{2d}(1 + i\tau) = \begin{bmatrix} A_{\tau} & \varepsilon I & & \\ & \ddots & \varepsilon I & \\ & & A_{\tau} & \end{bmatrix} \text{ where } A_{\tau} = \begin{pmatrix} 1 & \tau \\ -\tau & 1 \end{pmatrix}$$

and

$$J_d(1) = \begin{bmatrix} 1 & \varepsilon & \\ & \ddots & \varepsilon \\ & & 1 \end{bmatrix}$$

where d denotes the size of the block. Consider the left invariant metric $\langle \cdot, \cdot \rangle_{\varepsilon}$ such that $\mathcal{F}_{\varepsilon}$ is orthonormal and $T \perp [\mathfrak{s}, \mathfrak{s}]$, $\langle T, T \rangle = 1$ for some T such that $\alpha = \operatorname{ad}(T)$. Decompose $\operatorname{ad}(T) = D_{\varepsilon} + S_{\varepsilon}$, D_{ε} is symmetric and S_{ε} is skew-symmetric. To express the Riemann curvature tensor, following Heintze, Eberlein and Heber it is convenient to introduce⁶ $N_{\varepsilon} = D_{\varepsilon}^2 + [D_{\varepsilon}, S_{\varepsilon}]$. For all X, Y, Z in $\mathfrak{s}$,

$$\begin{aligned} R_{X,Y}Z &= -\langle D_{\varepsilon} \underline{Y}, Z \rangle D_{\varepsilon} \underline{X} + \langle D_{\varepsilon} \underline{X}, Z \rangle D_{\varepsilon} \underline{Y} \\ &\quad - \langle \underline{Z}, \langle X, T \rangle N_{\varepsilon} \underline{Y} - \langle Y, T \rangle N_{\varepsilon} \underline{X} \rangle T \\ &\quad + \langle Z, T \rangle (\langle X, T \rangle N_{\varepsilon} \underline{Y} - \langle Y, T \rangle N_{\varepsilon} \underline{X}), \end{aligned}$$

⁶They are denoted D_0, S_0 in [Hei74] and D_0, S_0, N_0 in [EH96].

where $\underline{X}$, $\underline{Y}$ and $\underline{Z}$ are the orthogonal projections of X, Y, Z to $[\mathfrak{s}, \mathfrak{s}]$. (This is differently expressed as in, but still in agreement with, [EH96] who performed a more general computation where $[\mathfrak{s}, \mathfrak{s}]$ is not assumed abelian and provided $R_{X,Y}Z$ for $X, Y, Z \in [\mathfrak{s}, \mathfrak{s}]$ and the sectional curvature of all planes.) Any 2-plane π in $\mathfrak{s}$ can be generated by $u, v \in \mathfrak{s}$ such that $v \in [\mathfrak{s}, \mathfrak{s}]$, so that $v = \underline{v}$. Observe that as $\varepsilon \rightarrow 0$, $D_\varepsilon \rightarrow I$ and $N_\varepsilon \rightarrow I$ so that

$$\begin{aligned} K^\varepsilon(\pi) &= \frac{\langle R^\varepsilon(u, v)v, u \rangle}{\langle u, u \rangle \langle v, v \rangle - \langle u, v \rangle^2} \\ &= \frac{\langle -D_\varepsilon \underline{u}, \underline{u} \rangle \langle D_\varepsilon v, v \rangle + \langle D_\varepsilon \underline{u}, v \rangle^2 - \langle u, T \rangle^2 \langle v, N_\varepsilon v \rangle}{\langle u, u \rangle \langle v, v \rangle - \langle u, v \rangle^2} \\ &\xrightarrow{\varepsilon \rightarrow 0} \frac{-\langle \underline{u}, \underline{u} \rangle \langle v, v \rangle + \langle u, v \rangle^2 - \langle u, T \rangle^2 \langle v, v \rangle}{\langle u, u \rangle \langle v, v \rangle - \langle u, v \rangle^2} = -1, \end{aligned}$$

using that $\langle u, u \rangle = \langle \underline{u}, \underline{u} \rangle + \langle u, T \rangle^2$ and $\langle \underline{u}, \underline{v} \rangle = \langle u, v \rangle$. Finally, the pointwise convergence of a rational function on a Grassmanian implies its uniform convergence, so $\sup K - \inf K$ goes to zero and $\sup K / \inf K$ goes to 1 as $\varepsilon \rightarrow 0$. Let us now prove the “moreover” part. K has a normal connected subgroup K_0 whose Adjoint representation stabilizes the direct sum V of J'_2 and J_1 type blocks, while $\Phi = K/K_0$ is a finite retract of K that permutes the blocks of equal higher size; K splits as $K_0 \rtimes \Phi$. In our construction, $\langle \cdot, \cdot \rangle_\varepsilon$ is already Ad_Φ -invariant. In order to make it Ad_K -invariant, we can average the metric on V using Weyl’s unitary trick, decompose $\mathbf{R}^{n-1} = V \oplus V'$ and reproduce our variation of orthonormal basis $\mathcal{F}_\varepsilon$ only on the complementary subspace V' which is the sum of higher sized Jordan blocks. $\square$

Remark 3.10. Using Eberlein and Heber’s amalgams (Definition 3.3) and curvature estimates would simplify the proof of the first part of Proposition 3.9 (yet not drastically so) by reducing it to the case where α has a single Jordan block as Jordan normal form. See also Remark 3.18.

Question 3.11. Let $G = N \rtimes (K \times R)$ be a group of Heintze type. Is it true that among all negatively curved Riemannian models of G , an optimal pinching is attained if and only if α is diagonalizable over $\mathbf{C}$?

Note that the (Ahlfors-regular) conformal dimension of $\partial_\infty[\mathbf{R}^{n-1} \rtimes_\alpha \mathbf{R}]$ is attained if and only if α is diagonalizable over $\mathbf{C}$ [BK05].

3.2. Degenerations and deformations. We provide more information here than is strictly needed for Theorem C. That will be useful to us in the discussion in §5.1.

3.2.1. Setting. Let $\mathcal{L}_n(\mathbf{R}) \subseteq (\Lambda^2 \mathbf{R}^n)^* \otimes \mathbf{R}^n$ be the subset of Lie algebra laws on $\mathbf{R}^n$. Note that $\mu \in (\Lambda^2 \mathbf{R}^n)^*$ is in $\mathcal{L}_n(\mathbf{R})$ if and only if the Jacobi identity holds in μ , that is, if and only if

$$(3.7) \quad \mu^2(X_1 \wedge X_2 \wedge X_3) = \sum_{\sigma} \mu(\mu(X_{\sigma(1)} \wedge X_{\sigma(2)}) \wedge X_{\sigma(3)}) = 0$$

for every $X_1, X_2, X_3 \in \mathbf{R}^n$, the sum being taken over the three positive permutations σ over $\{1, 2, 3\}$. $\mathcal{L}_n(\mathbf{R})$ has two topologies: the Zariski topology, and the

topology it inherits as a subspace of $\Lambda^2(\mathbf{R}^n)^* \otimes \mathbf{R}^n$ with the operator norm, that we will call the metric topology. It follows from Engel's theorem that the nilpotent laws form a Zariski closed subset $\mathcal{N}_n(\mathbf{R})$.

Let $\lambda \in \mathcal{L}_n(\mathbf{R})$. $\mathbf{R}$, resp. λ , is a λ -module for the trivial, resp. the adjoint representation of λ . Following Chevalley and Eilenberg [CE48, Theorem 10.1] there are differential complexes K_λ and K'_λ on $\Lambda^\bullet(\mathbf{R}^n)^*$ and $\Lambda^\bullet(\mathbf{R}^n)^* \otimes \mathbf{R}^n$ with the following exterior derivatives d_λ , resp. d'_λ on degree q -forms, resp. on λ -valued degree q -forms ω :

$$(3.8) \quad d_\lambda \omega(x_1, \dots, x_{q+1}) = \sum_{k < \ell} (-1)^{k+\ell+1} \omega([x_k, x_\ell], x_1, \dots, \widehat{x}_k, \dots, \widehat{x}_\ell, \dots, x_{q+1})$$

$$(3.9) \quad d'_\lambda \omega(x_1, \dots, x_{q+1}) = \sum_{k < \ell} (-1)^{k+\ell+1} \omega([x_k, x_\ell], x_1, \dots, \widehat{x}_k, \dots, \widehat{x}_\ell, \dots, x_{q+1})$$

$$(3.9) \quad + \sum_k (-1)^{k+1} [x_k, \omega(x_1, \dots, \widehat{x}_k, \dots, x_{q+1})].$$

The group $\mathrm{GL}(n, \mathbf{R})$ acts on $\mathcal{L}_n(\mathbf{R})$ by restricting its natural action on $\Lambda^2(\mathbf{R}^n)^* \otimes \mathbf{R}^n$. We denote the orbit of λ by $O(\lambda)$ or $O_{\mathfrak{g}}$ if $\mathfrak{g}$ is a Lie algebra isomorphic to λ ; it is a smooth submanifold of $\Lambda^2(\mathbf{R}^n)^* \otimes \mathbf{R}^n$ of dimension $n^2 - \dim \mathrm{Der}(\mathfrak{g})$, embedded in $\mathcal{L}_n(\mathbf{R})$. Moreover, $T_\lambda O_{\mathfrak{g}} = B^2(\lambda, \lambda)$, as is most conveniently seen by differentiating the action of $\mathrm{GL}(n, \mathbf{R})$ at λ : for every $\eta \in \mathfrak{gl}(\mathbf{R}^n)$,

$$(3.10) \quad e^\eta \lambda(e^{-\eta} X, e^{-\eta} Y) - \lambda(X \wedge Y) = d'_\lambda \eta(X \wedge Y) + O(\|\eta\|^2).$$

Example 3.12. Let $\mathfrak{g} = \mathfrak{aff}$ be the 2-dimensional affine Lie algebra with basis $[X, T]$ such that $[T, X] = X$ and dual basis (dx, dt) . Then $X \otimes dx \wedge dt \in B^2(\mathfrak{g}, \mathfrak{g})$; in the language of §3.1, $\mathbf{R} \rtimes_{1+\varepsilon} \simeq \mathbf{R} \rtimes_1 \mathbf{R} \simeq \mathfrak{g}$.

Definition 3.13. Let $\mathfrak{g}$ and $\mathfrak{h}$ be Lie algebras of dimension n over $\mathbf{R}$. We say that $\mathfrak{g}$ degenerates to $\mathfrak{h}$, denoted $\mathfrak{g} \rightarrow_{\mathrm{deg}} \mathfrak{h}$, if $\overline{O_{\mathfrak{h}}} \subsetneq \overline{O_{\mathfrak{g}}}$ where the closure is taken for the Zariski topology.

Note that it is equivalent to require a single $\mu \in O_{\mathfrak{h}}$ such that $\mu \in \overline{O_{\mathfrak{g}}}$. Since the metric topology is finer than the Zariski topology, a sufficient condition to have $\mathfrak{g} \rightarrow_{\mathrm{deg}} \mathfrak{h}$ is that there is a sequence $\lambda_0, \dots, \lambda_r$ such that

$$(3.11) \quad \begin{cases} \lambda_0 \in O_{\mathfrak{g}}, \lambda_r \in O_{\mathfrak{h}} \\ \forall X \in \Lambda^2(\mathbf{R}^n), \lim_{t \rightarrow +\infty} (\varphi_{t,i} \cdot \lambda_i)(X) = \lambda_{i+1}(X) \quad i = 0, \dots, r-1. \end{cases}$$

where $\varphi_t \in \mathrm{GL}(n, \mathbf{R})$ is continuous with respect to t .

When $r = 1$, (3.11) amounts to $\mu \in \overline{O(\lambda)}^{\mathrm{met}}$ and is called a contraction (especially, by the physicists). The author does not know whether the existence of a sequence of contractions as in (3.11) is a necessary condition for $\mathfrak{g} \rightarrow_{\mathrm{deg}} \mathfrak{h}$ to hold.

Example 3.14 (Nilpotent Lie algebras). Let $\mathfrak{n}$ be a nilpotent Lie algebra. Let $\mathfrak{n} = \oplus_i V_i$ be a linear splitting such that $V_i \oplus C^{i+1}\mathfrak{n} = C^i\mathfrak{n}$ for all i . For $t \geq 0$, let (φ_t) be the one parameter subgroup of $\mathrm{GL}(\mathfrak{n})$ such that

$$(3.12) \quad \varphi_t(X) = t^i X, \quad X \in V_i.$$

Then, the V_i becomes a Lie algebra grading on $\varphi_t \cdot \mathfrak{n}$ in the limit when $t \rightarrow +\infty$: $\mathfrak{n}$ degenerates metrically to the graded Lie algebra $\mathrm{gr}(\mathfrak{n})$ associated to the central filtration of $\mathfrak{n}$, supporting the asymptotic cone of the simply connected N by [Pan83]. In particular, $\mathfrak{n} \rightarrow_{\mathrm{deg}} \mathrm{gr}(\mathfrak{n})$. (This description of the law in $\mathrm{gr}(\mathfrak{n})$ as a limit is the one given in [CF17, §2.1], who prove a generalization of [Pan83].)

For $\lambda \in \Lambda^2 \mathbf{R}^n \otimes_{\mathbf{R}} (\mathbf{R}[[1/t]])^n$, we denote $(\lambda, t) \mapsto \lambda(t)$ provided that t is in the convergence domain of every coefficient of λ , and $\lambda[1/t^d]$ the monomial of degree d . (The choice of $\mathbf{R}[[1/t]]$ over $\mathbf{R}[[t]]$ is just a peculiarity for our convenience.) We also denote $\lambda(\infty)$ the constant term of λ . If $\lambda(t) \in \mathcal{L}_n(\mathbf{R})$ for all $t \geq 1$, λ is called a formal deformation.

Differentiating (3.7) to express that λ is a formal deformation with $\lambda(\infty) = \mu$ yields an infinite system of equations, the first of which after (3.7) being

$$(3.13) \quad d'_\mu \lambda[1/t] = 0,$$

that is, $\lambda[1/t] \in Z^2(\mu, \mu)$.

Definition 3.15. Let $\mathfrak{g}$ be a Lie algebra over $\mathbf{R}$. Let $\mu \in \mathcal{L}^n(\mathbf{R})$ represent $\mathfrak{g}$, and let $\omega \in H^2(\mathfrak{g}, \mathfrak{g})$ be nonzero. We say that the formal deformation λ integrates the infinitesimal deformation ω at μ if $\lambda(\infty) = \mu$, λ is convergent on $\mathbf{C} \setminus \{0\}$ and $\lambda[1/t] \in Z^2(\mu, \mu)$ represents ω . We say that ω is integrable, resp. linearly expandable (as the authors in [AC01] do) if a formal deformation λ integrates ω , resp. if λ is a formal deformation of ω and $\lambda = \lambda(\infty) + \lambda_1/t$ for some $\lambda_1 \in \Lambda^2 \mathbf{R}^n \otimes \mathbf{R}^n$.

In the last Definition, we insisted more on the cohomology class than on the particular cocycle $\lambda[1/t]$ for the following reason. Two formal deformations λ, λ' of μ are called equivalent if $\lambda(t) = \varphi(t) \cdot \lambda'(t)$ for some $\varphi \in \mathrm{GL}(\mathbf{R}[[t]])$ with $\varphi(\infty) = 1$. If λ and λ' are equivalent then $\lambda[1/t] - \lambda'[1/t] \in B^2(\mu, \mu)$; this is a better version of (3.10), see e.g. Proposition just before §2.5 in [AC01]. In view of (3.10), (3.13) and this, $H^2(\mathfrak{g}, \mathfrak{g})$ encodes the degree to which $\mathfrak{g}$ can be deformed; one should nevertheless beware that infinitesimal deformations are not always integrable (See Remark 3.19).

3.2.2. Degenerations to $\mathfrak{b}(n, \mathbf{R})$. Let $\mathfrak{b}(n, \mathbf{R})$ denote the maximal completely solvable subalgebra of $\mathfrak{o}(n, 1)$, namely $\mathfrak{b}(n, \mathbf{R}) = \mathbf{R}^{n-1} \rtimes_1 \mathbf{R}$, where the adjoint action of $1 \in \mathbf{R}$ on $\mathbf{R}^{n-1}$ is by the identity. The situation of $\mathfrak{b}(n, \mathbf{R})$ with respect to degenerations and deformations is favorable:

Theorem 3.16 (After Lauret). *Let $\mathfrak{g}$ be a completely solvable Lie algebra and $n \geq 2$ an integer. The following are equivalent:*

- (3.16.1) $\mathfrak{g}$ contracts to $\mathfrak{b}(n, \mathbf{R})$.
- (3.16.2) $\mathfrak{g} \rightarrow_{\mathrm{deg}} \mathfrak{b}(n, \mathbf{R})$.
- (3.16.3) $\mathfrak{g}$ decomposes as $\mathbf{R}^{n-1} \rtimes_\nu \mathbf{R}$ where ν is unipotent.

Moreover, under the former conditions there exists $\omega \in H^2(\mathfrak{b}(n, \mathbf{R}), \mathfrak{b}(n, \mathbf{R}))$ linearly expandable into a formal deformation λ such that $\lambda(1) \in \mathcal{O}_{\mathfrak{g}}$ and $\lambda(\infty) \in \mathcal{O}_{\mathfrak{b}(n, \mathbf{R})}$.

Lauret proved (3.16.1) $\iff$ (3.16.3) [Lau03, Theorem 6.2] with no a priori assumption on $\mathfrak{g}$. The core of the proof below uses the same idea. ([Lau03] additionally used bounds on pinching and [EH96] that give constraints a priori on $\mathfrak{g}$).

We need a Lemma which is well-known, however we could only find proofs for the metric topology in the literature.

Lemma 3.17. *Let n be a positive integer and $0 \leq i \leq n$. Then, the following are upper semi-continuous with respect to the Zariski topology on $\mathcal{L}_n(\mathbf{R})$:*

- (a) *The Betti number $b_p(\lambda) = \dim H^p(\lambda, \mathbf{R})$, for all $p \geq 0$.*
- (b) *The dimension of the outer derivations $H^1(\lambda, \lambda) = \text{Der}(\lambda) / \text{InnDer}(\lambda)$*
- (c) *The dimension of the center $\dim Z(\lambda)$.*

Proof. Note that $Z(\lambda) = H^0(\lambda, \lambda)$, so to prove (a), (b) and (c) it is actually sufficient to prove that $\lambda \mapsto b_p(\lambda)$ and $\lambda \mapsto \dim H^p(\lambda, \lambda)$ are upper semicontinuous on $\mathcal{L}_n(\mathbf{R})$. We will prove this by a change of basis argument. Denote by x_{ij}^k the coordinate functions on $\Lambda^2(\mathbf{R}^n)^* \otimes \mathbf{R}^n$, and

let $\mathcal{I}$ be the ideal of $\mathbf{R}[x_{ij}^k]$ generated by the relation (3.7).

Let $A = \mathbf{R}[x_{ij}^k] / \mathcal{I}$. Then A is a noetherian ring by Hilbert's basis theorem, and $\mathcal{L}_n(\mathbf{R})$ with the Zariski topology is a closed subspace of $\text{Spec}(A)$ with the Zariski topology; all the points in $\mathcal{L}_n(\mathbf{R})$ are maximal ideals. Consider the graded A -modules

$$K = \Lambda^\bullet(A^n)^*$$

$$K' = \Lambda^\bullet(A^n)^* \otimes_A A^n.$$

Then, for every $(y_1, y_2) \in \mathbf{R}[x_{ij}^k]^n$, we set $z = [y_1, y_2]$ to be an element of $\mathbf{R}[x_{ij}^k]^n$ such that $z(\lambda) = [y_1(\lambda), y_2(\lambda)]$ for all $\lambda \in \Lambda^2(\mathbf{R}^n)^* \otimes \mathbf{R}^n$. Note that z only depend on y_1 and y_2 modulo $\mathcal{I}^n$, and is well-defined only modulo $\mathcal{I}^n$; thus $[\cdot, \cdot]$ defines an element of $\Lambda^2(A^n)^* \otimes A^n$. The differentials on K and K' are defined as in (3.8) and (3.9).

Then $K_\lambda = K \otimes_A A/\lambda$ and $K'_\lambda = K' \otimes_A A/\lambda$, where A/λ is the residual field of A at the maximal ideal λ . K and K' are flat A -modules, because being flat is preserved by taking exterior and tensor products over the base ring [Laz69, Proposition 2.3]. We may now conclude by applying the following [Gro61, Théorème 7.6.9(i)]: if A is noetherian and K is a differential complex of finitely generated flat modules, then for every $p \geq 0$, the function $y \mapsto H^p(K \otimes_A k(y))$ is upper semi-continuous on $\text{Spec}(A)$, where $k(y)$ denotes the residual field at y . In particular, it is upper semi-continuous on the closed subspace $\mathcal{L}_n(\mathbf{R})$. □

Proof of Theorem 3.16. (1) $\implies$ (2) is clear.

Assume (2). By Lemma 3.17, $b_1(\mathfrak{g}) \leq 1$. If it is zero, then $\mathfrak{g}$ is perfect, especially it is not solvable; hence $b_1 = 1$, and $\mathfrak{g}$ splits as a semidirect product

$$(3.14) \quad [\mathfrak{g}, \mathfrak{g}] \oplus \mathbf{R}A$$

where ad_A is nonsingular in view of the fact that $Z(\mathfrak{g}) = 0$, again by Lemma 3.17. Choosing an adequate representative λ_0 in $O_{\mathfrak{g}}$ and an adequate basis we may as well assume that $[\lambda_0, \lambda_0] = \mathbf{R}^{n-1}$ and $A = (0^{n-1}, 1)$.

The coefficients of the characteristic polynomial $P_{\mu,X}$ of $\text{ad}_X : Y \mapsto \mu(X, Y)$ are polynomial functions on $\mathcal{L}_n(\mathbf{R})$, and for every $\lambda_1 \in O(\lambda)$ the spectrum of $P_{\lambda_1,X}$ is either a nonzero multiple of $\text{Sp}(P_{\lambda_0,A})$, or 0 with multiplicity n , the latter case occurring $X \in [\lambda_1, \lambda_1]$. So, for $\mu \in O(\lambda)$ this holds as well. But for $\mu \in O_{\mathfrak{b}(n,\mathbf{R})}$, this spectrum is always concentrated at one point. So ad_A cannot have two distinct eigenvalues, and then $[\mathfrak{g}, \mathfrak{g}]$ is abelian, which proves (3).

Assume (3). Then $\nu - 1$ is nilpotent; let $X_1, \dots, X_{n-1}$ be a basis of $[\mathfrak{g}, \mathfrak{g}]$ in which it appears in lower-triangular Jordan form, $\nu - 1 = \sum_i \delta_i X_i^* \otimes X_{i+1}$ where $\delta_i \in \{0, 1\}$. One computes that $d(A \wedge X_i^* \otimes X_{i+1}) = 0$ (Lemma A.7; beware that S replaces A there) and that no nonzero linear combination of those is a coboundary (Lemma A.6). Setting μ the law of $\mathfrak{b}(n, \mathbf{R})$ in the basis $(X_1, \dots, X_{n-1}, A)$ and $\omega = A^* \wedge \sum_i \delta_i X_i^* \otimes X_i$ we find that $\lambda_0 = \mu + \omega$. Then μ is the degeneration of λ through (φ_t) , where $\varphi_t A = A$ and $\varphi_t X^i = t^{-i} X_i$ for all t . $\square$


Remark 3.18. A contraction to (a deformation of) $\mathfrak{b}(n, \mathbf{R})$ was already implicitly used in the proof of Proposition 3.9; in accordance with [Lau03], contractions can be considered as limit points in the space of left-invariant Riemannian metrics over a given group.

Remark 3.19. We can additionally check that $H^3(\mathfrak{b}, \mathfrak{b}) = 0$ when $\mathfrak{b} = \mathfrak{b}(2, \mathbf{R})$, though it is unnecessary. This vanishing ensures that the deformation system can be solved and every infinitesimal deformation of $\mathfrak{b}$ is integrable into a formal deformation [NR67, p.98]. For nilpotent Lie algebras $\mathfrak{n}$ that will be discussed more in detail in §5.1; on the other hand, one must beware that $H^3(\mathfrak{n}, \mathfrak{n})$ is large, for instance $\dim H^3(\mathfrak{n}, \mathfrak{n}) \geq 8$ for all the 6-dimensional nilpotent $\mathfrak{n}$ [Mag08, Table 11].

3.3. Groups $O(u)$ -bilipschitz equivalent to $\mathbb{H}_{\mathbf{R}}^n$. We prove here Theorem C. Let us first recall some terminology from [CT11] and [CCMT15].

Definition 3.20. Let G be a connected Lie group with finitely many components. G is of rank-one type if it has a maximal normal compact subgroup W such that G/W is isomorphic to a simple Lie group $G_{\mathbf{R}}$ of real rank one, with $Z(G_{\mathbf{R}}) = 1$.

3.3.1. (C.1) implies (C.3). Let G be a Lie group with finitely many connected components. Assume that G is $O(u)$ -sublinear bilipschitz equivalent to $\mathbb{H}_{\mathbf{R}}^n$ for some n . Then all asymptotic cones of G being $\mathbf{R}$ -trees, G is Gromov-hyperbolic. By Cornulier and Tessera's theorem [CT11], G is either of Heintze or rank-one Lie type. First assume that G is of Heintze type, write $G = (K \times \mathbf{R}) \ltimes N$ and call H the co-compact normal subgroup $\mathbf{R} \ltimes N$ so that G/K is simply transitively acted upon by H . By [Pal20], $\text{Cdim}_{O(u)} \partial_{\infty} H = \text{Cdim}_{O(u)} \partial_{\infty} \mathbb{H}_{\mathbf{R}}^n = n - 1$. By [Cor19], $\text{Topdim} \partial_{\infty} H = n - 1$. So H is metabelian and every eigenvalue of α has real part 1. By Proposition 3.9, (C.3) holds, while by [Cor11, Theorem 1.2], (C.2) holds. If G is of rank-one type, then it acts properly co-compactly by isometries on a rank one symmetric space, which can only be $\mathbb{H}_{\mathbf{R}}^n$ in view of the equality of



conformal dimension and topological dimension of the boundary; especially, (C.2) and (C.3) hold as well.

3.3.2. (C.3) *implies* (C.2). Since it acts geometrically on Gromov-hyperbolic spaces, G is Gromov-hyperbolic. Again by [CT11], it is of Heintze type or rank-one type. If it is rank-one type, then it is quasiisometric to a rank one symmetric space X ; by Pansu's Theorem 3.5, $\text{Cdim}(\partial_\infty G) = \text{Topdim}(\partial_\infty G)$, so $X = \mathbb{H}_{\mathbf{R}}^n$. If it is Heintze-type, then it is $O(\log)$ -bilipschitz equivalent to a purely real Heintze group.

3.3.3. (C.2) *implies* (C.1). $u = \log$ is an admissible function.

3.3.4. *If G is completely solvable then (C.3) implies (C.5).* By [CT11], it is of Heintze type. and by Corollary 3.7, N is abelian and all the eigenvalues of α have real part 1.

3.3.5. (C.4) *and* (C.5) *are equivalent.* This is our version of Lauret's theorem, Theorem 3.16.

3.3.6. (C.4) *implies* (C.3). This is a special case of Proposition 3.9 where all the eigenvalues of ad_A are real.

3.4. **From the pinching condition to Higson coronae.** Corollary D follows by applying (C.3) $\implies$ (C.1) together with Proposition 2.14.

4. PROOF OF THEOREM E

4.1. **Pointed sphere.** We will prove the implication (E.1) $\implies$ (E.3) in Theorem E by establishing a baby case of a variant of Cornuier's pointed sphere conjecture [Cor18, Conjecture 19.104]. Precisely we establish a special case of the conjecture in the setting of sublinear bilipschitz equivalences rather than quasi-isometries for which it is usually formulated. We denote by $\text{SBE}^{O(u)}(X)$ the group of self $O(u)$ -bilipschitz equivalences of the metric space X (modulo the relation of $O(u)$ -closeness). Let us first recall that sublinear bilipschitz equivalences induce homeomorphisms of the compact boundary sphere $\partial_\infty X$ when X is Gromov-hyperbolic [Cor19].

Lemma 4.1. *Let u be an admissible function. Let S be a purely real Heintze group such that $[S, S]$ is abelian, and let Ω be the unique closed orbit of $\text{SBE}^{O(u)}(S)$ acting by homeomorphisms on $\partial_\infty S$. The following are equivalent:*

(4.1.1) α has at least two distinct eigenvalues

(4.1.2) Ω is reduced to a single point.

Proof. The reasoning is inspired from [Pan89a, 6.9 Corollaire]. Let ω be the endpoint of a $\mathbf{R}$ section in S , so that $\partial_\infty S \setminus \{\omega\}$ is simply transitively acted upon by $[S, S]$. Assume (1) and let $\mathcal{F}$ be the foliation on $\partial_\infty S \setminus \{\omega\}$ determined by the cosets of $\ker(\alpha - \lambda)$, where λ is the minimal eigenvalue of α (since $[S, S]$ is abelian, we may identify it with its Lie algebra). Then by [Pal20, Lemma 3.9], for all sublinear bilipschitz equivalence $f : S \rightarrow S$, the boundary map $\partial_\infty f$ preserves $\mathcal{F}$. Now let F be any leaf of $\mathcal{F}$. Then, $\{\omega\}$ can be written as $\overline{F} \setminus F$

Nilradical	Jordan(α)	$\mathbb{H}_{\mathbf{K}}^n$	Nilradical	Jordan(α)	$\mathbb{H}_{\mathbf{K}}^n$
$\mathbf{R}^2$	$\text{diag}(1, \lambda)$		$\mathbf{R}^3$	$\text{diag}(1, 1, 1)$	$\mathbb{H}_{\mathbf{R}}^4$
$\mathbf{R}^2$	$\text{diag}(1, 1)$	$\mathbb{H}_{\mathbf{R}}^3$	$\mathbf{R}^3$	$\text{diag}(1, J_2(1))$	
$\mathbf{R}^2$	$J_2(\lambda)$		$\mathbf{R}^3$	$J_3(1)$	
$\mathbf{R}^3$	$\text{diag}(1, \lambda, \lambda)$		$\mathbf{R}^3$	$\text{diag}(1, \lambda, \mu)$	
$\mathbf{R}^3$	$\text{diag}(1, J_2(\lambda))$		$\mathbf{Heis}_3$	$\text{diag}(1, \lambda, 1 + \lambda)$	
$\mathbf{R}^3$	$\text{diag}(1, 1, \lambda)$		$\mathbf{Heis}_3$	$\text{diag}(1, 1, 2)$	$\mathbb{H}_{\mathbf{C}}^2$
$\mathbf{R}^3$	$\text{diag}(J_2(1), \lambda)$		$\mathbf{Heis}_3$	$\text{diag}(J_2(1), 2)$	

TABLE 2. Purely real Heintze groups of dimension 3 or 4, with parameters $1 < \lambda < \mu$. The plain horizontal lines denote the separations between $O(\log)$ -bilipschitz equivalence classes that can be deduced from [Pal20] and Theorem E. The dash line remains unknown when $\mu = 1 + \lambda$. The isomorphism type of $N \rtimes_{\alpha} \mathbf{R}$ is generally not determined by N and Jordan(α) alone; see the 6-dimensional example after Theorem 1.3 in [CPS17].

or $\overline{(\partial_{\infty} f)F} \setminus (\partial_{\infty} f)F$, so that $\partial_{\infty} f \omega = \omega$. Conversely, if α only has a single eigenvalue, then S is sublinearly bilipschitz equivalent to real hyperbolic space. Since $\text{Isom}(\mathbb{H}_{\mathbf{R}}^n)$ is transitive on $\partial_{\infty} \mathbb{H}_{\mathbf{R}}^n$, $\text{SBE}^{O(u)}(S)$ is transitive on $\partial_{\infty} S$. $\square$

Proposition 4.2. *Let u be an admissible function. Let S be a Heintze group. Assume that S is $O(u)$ -bilipschitz equivalent to $\mathbb{H}_{\mathbf{C}}^2$. Then the shadow of S is isomorphic to $\mathbf{Heis} \rtimes_{\alpha} \mathbf{R}$ where $\mathbf{Heis}$ is the three-dimensional Heisenberg group and*

$$\alpha = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{pmatrix} \quad \text{or} \quad \alpha = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{pmatrix}$$

in a basis (X, Y, Z) of $\mathfrak{heis}$ such that $[X, Y] = Z$.

Proof. Let $S_0 = N \rtimes_{\alpha} \mathbf{R}$ be a semidirect product decomposition of the shadow S_0 of S , where N is three-dimensional and α is normalized so that its lowest eigenvalue is 1. Since S has been assumed $O(u)$ -bilipschitz equivalent to $\mathbb{H}_{\mathbf{C}}^2$, by Theorem B, $\dim S = \text{asdim}_{\text{AN}} \mathbb{H}_{\mathbf{C}}^2 - \text{conedim} \mathbb{H}_{\mathbf{C}}^2 = 3$. So N is isomorphic either to $\mathbf{R}^3$ or to the 3-dimensional Heisenberg group. In the first case, since $\text{Tr}(\alpha) = \text{Cdim}_{O(u)}(S) = 4 > 3$, α has at least two distinct eigenvalues, and by Lemma 4.1, the unique closed orbit of $\text{SBE}^{O(u)}(S_0)$ acting on $\partial_{\infty} S_0$ has only one element (namely, ω from the proof of 4.1). This contradicts the fact that $\text{SBE}^{O(u)}(\mathbb{H}_{\mathbf{C}}^2)$ is transitive on $\partial_{\infty} \mathbb{H}_{\mathbf{C}}^2$, so this cannot be. Consequently, N is isomorphic to the three-dimensional Heisenberg group. Let $1, \lambda, \mu$ be the eigenvalues of α , where μ corresponds to the eigenvector generating the center of $\mathbf{Heis}$, and $1 \leq \lambda \leq \mu$. Necessarily, $1 + \lambda = \mu$ and $1 + \lambda + \mu = 4$, so $2 + 2\lambda = 4$, and then $\lambda = 1$. We deduce from there that α can only be one of the two derivations in the conclusion. $\square$

Proof of (E.1) $\implies$ (E.3). Let G be as in the statement of Theorem E, namely G is a connected Lie group sublinear bilipschitz equivalent to $\mathbb{H}_{\mathbf{C}}^2$. Then G is

commable to a completely solvable group G_0 [Cor08, Lemma 6.7]. Since G_0 is Gromov-hyperbolic, by [CT11] it is a purely real Heintze group [CT11]. We may then apply Proposition 4.2 to G_0 . in the first case where α is diagonalisable, G_0 (hence G) will be commable to $SU(2, 1)$, in the second case it will be commable to S' . $\square$

Let us mention an application to the quasiisometry classification of Heintze groups. The result below also follows from [KDG21, Theorem A] which appeared during the writing of this paper.

Corollary 4.3. *The groups S' and $S'' = \mathbf{R}^3 \rtimes_{\alpha} \mathbf{R}$ where*

$$\alpha = \text{diag}(J_2(1), 2) = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{pmatrix}$$

are not quasiisometric.

Indeed, if S and S' were quasiisometric, they would be $O(\log)$ -bilipschitz equivalent. But S' is $O(\log)$ -bilipschitz equivalent to $\mathbb{H}_{\mathbf{C}}^2$, whereas S'' is not.

See Table 2 for the Heintze groups of dimension at most 4 and the current knowledge on their $O(\log)$ -bilipschitz classification (their quasiisometry classification is known and reduces to isomorphism, see [KDG21, Theorem C]).

Remark 4.4. We can start the same reasoning with $X = \mathbb{H}_{\mathbf{C}}^n$, $n > 2$. By conformal dimension, any purely real Heintze group S that is $O(u)$ -bilipschitz equivalent to X has $[S, S]$ isomorphic to $\mathbf{Heis}^{2k+1} \times \mathbf{R}^{2(n-k)}$ for some $k \in \{0, \dots, k-1\}$, where $\mathbf{Heis}^{2k-1}$ denotes the $2k-1$ -dimensional Heisenberg group for $k \geq 2$ and $H^1 = \mathbf{R}$. Otherwise said, using the amalgam notation (Definition 3.3)

$$\mathfrak{s} = \mathfrak{b}(k, \mathbf{C}) \sharp \mathfrak{b}(2(n-k) + 1, \mathbf{R}),$$

where we recall that $\mathfrak{b}(k, \mathbf{C})$ is the maximal completely solvable subalgebra of $\mathfrak{u}(k, 1)$. But we are only able to prove the pointed sphere conjecture for S when $k = 1$: for $k \geq 2$ the invariant foliation in $\partial_{\infty} S$ provided by [Pal20, Lemma 3.9] becomes a single leaf. The same reasoning also falls short to characterize the triangulable groups S that are $O(u)$ -bilipschitz equivalent to $X = \mathbb{H}_{\mathbf{H}}^2$, for it leaves the possibility that the Lie algebra of their shadow is

$$\mathfrak{s}_0 \in \{\mathfrak{b}(5, \mathbf{R}) \sharp 2\mathfrak{b}(4, \mathbf{R}), \mathfrak{b}(2, \mathbf{C}) \sharp \mathfrak{b}(2, \mathbf{R}) \sharp 2\mathfrak{b}(3, \mathbf{R}), \mathfrak{b}(3, \mathbf{C}) \sharp 2\mathfrak{b}(2, \mathbf{R}), \\ \mathfrak{n}_6 \rtimes_{\text{Carnot}} \mathbf{R} \sharp 2\mathfrak{b}(2, \mathbf{R}), \mathfrak{n}_7 \rtimes_{\text{Carnot}} \mathbf{R}, \mathfrak{b}(4, \mathbf{R}) \sharp \mathfrak{l}_{4,3} \rtimes_{\text{Carnot}} \mathbf{R}, \mathfrak{b}(2, \mathbf{H})\}$$

where $\mathfrak{l}_{4,3}$ denotes the 4-dimensional filiform algebra, $\mathfrak{n}_6$ is among $\mathfrak{l}_{6,8}$, $\mathfrak{l}_{6,22}(-1)$ and $\mathfrak{l}_{6,22}(0)$ (See [dG07] for structure constants), $\mathfrak{n}_7$ is one among the real forms of the 4 complex nilpotent algebras denoted $\mathfrak{g}_{7,3,12}$ (2 real forms), $\mathfrak{g}_{7,3,24}$, $\mathfrak{g}_{7,4,1}$ (2 real forms) or $\mathfrak{g}_{7,4,2}$ in [Mag07]. Using [Pal20] one can only deduce the pointed sphere conjecture (Lemma 4.1) for the first 6 out of these 14 Lie algebras, while it is expected that it holds for all but the last one.

4.2. Degenerations to $\mathfrak{b}(2, \mathbf{C})$. We prove here a variant of Lauret's theorem 3.16.

Lemma 4.5. *Let $\mathfrak{g}$ be a completely solvable Lie algebra of dimension 4. The following are equivalent:*

(4.5.1) $\mathfrak{g}$ contracts to $\mathfrak{b}(2, \mathbf{C})$

(4.5.2) $\mathfrak{g} \xrightarrow{\deg} \mathfrak{b}(2, \mathbf{C})$

(4.5.3) $\mathfrak{g}$ decomposes as $[\mathfrak{g}, \mathfrak{g}] \oplus \mathbf{R}A$, where $[\mathfrak{g}, \mathfrak{g}] = \mathfrak{heis}$ and ad_A is unipotent on $[\mathfrak{g}, \mathfrak{g}]/D^3\mathfrak{g}$.

Proof. As in the proof of Theorem 3.16, the core of the proof is that (4.5.2) implies (4.5.3), so let us focus on this part. Assume that $\mathfrak{g} \xrightarrow{\deg} \mathfrak{b}(2, \mathbf{C})$. Then $b_1(\mathfrak{g}) = 1$ by Lemma 3.17. The ideal $\mathfrak{n} = [\mathfrak{g}, \mathfrak{g}]$ is nilpotent by Lie's theorem, and $\mathfrak{g} = \mathfrak{n} \rtimes_{\beta} \mathbf{R}$ for some nonsingular $\beta \in \text{ad}(\mathfrak{n})$. Without loss of generality we can assume that $\text{Sp}(\beta) = \{1, 2\}$, and that 2 has multiplicity 1. So the nilpotency class of $\mathfrak{n}$ is at most 2, and $\text{codim}[\mathfrak{n}, \mathfrak{n}] = 1$; thus $\mathfrak{n}$ is either $\mathbf{R}^3$ or $\mathfrak{heis}$, and $\mathfrak{g}$ is among the four algebras

$$\mathfrak{b}(2, \mathbf{C}), \mathfrak{b}(3, \mathbf{R}) \sharp 2\mathfrak{b}(2, \mathbf{R}), \mathfrak{s}', \mathfrak{s}'',$$

where we recall that $\mathfrak{s}' = \mathfrak{heis}_{\alpha} \rtimes \mathbf{R}$ with $\alpha = \text{diag}(J_2(1), 2)$ in the basis (X, Y, Z) and $\mathfrak{s}''$ is the Lie algebra of S'' defined in Corollary 4.3. Observe that

$$\dim H^1(\mathfrak{g}, \mathfrak{g}) = \begin{cases} 2 & \mathfrak{g} = \mathfrak{b}(2, \mathbf{C}) \text{ by Proposition A.3} \\ 4 & \mathfrak{g} = \mathfrak{s}'' \text{ by Proposition A.4} \end{cases}$$

Note that $\mathfrak{s}''$ degenerates to $\mathfrak{b}(3, \mathbf{R}) \sharp 2\mathfrak{b}(2, \mathbf{R})$. Hence by Lemma 3.17, $\dim H^1(\mathfrak{g}, \mathfrak{g}) \geq 4$ for $\mathfrak{g} = \mathfrak{b}(3, \mathbf{R}) \sharp 2\mathfrak{b}(2, \mathbf{R})$, which, again by Lemma 3.17, forbids a degeneration of the latter algebra to $\mathfrak{b}(2, \mathbf{C})$. This establishes (4.5.2) $\implies$ (4.5.3). Finally let us prove that $\mathfrak{s}' \xrightarrow{\deg} \mathfrak{b}(2, \mathbf{C})$. Take

$$\varphi_t X = X \quad \varphi_t Y = e^{-t} Y \quad \varphi_t Z = e^{-t} Z \quad \varphi_t A = A.$$

Then $\mathfrak{s}'$ contracts⁷ to $\mathfrak{b}(2, \mathbf{C})$ through (φ_t) . This establishes (4.5.3) $\implies$ (4.5.1). $\square$

The author expects that Lemma 4.5 should hold replacing $\mathfrak{b}(2, \mathbf{C})$ with $\mathfrak{b}(n, \mathbf{C})$ and $\mathfrak{heis}$ with $\mathfrak{heis}^{2n-1}$ in (4.5.3), though generalizing Proposition A.4 to higher dimensional algebras comprises some computational hurdles. The greatest theoretical difficulty in generalizing Theorem E (if it holds) from $\mathbb{H}_{\mathbf{C}}^2$ to $\mathbb{H}_{\mathbf{C}}^n$ with $n > 2$ seems to lie on the analytical side, cf. Remark 4.4 above.

5. SOME REMARKS ON SPACES OTHER THAN $\mathbb{H}_{\mathbf{R}}^n$ AND $\mathbb{H}_{\mathbf{C}}^n$

5.1. Connected Lie groups. In the attempts to relate the large-scale geometry of pairs of connected Lie groups, several sufficient criteria have been found (e.g. for quasiisometry in [Bre14], [Cor11], for sharing simply transitive Riemannian models in [CKD⁺21], and for $O(u)$ -bilipschitz equivalence in [Cor11]). These

⁷This was recorded by Burde and Steinhoff in their list of degenerations between 4-dimensional complex Lie algebras: $\mathfrak{s}' \otimes \mathbf{C}$ is $\mathfrak{g}(1/64, 5/16)$ in [BS99] and $\mathfrak{s}' \otimes \mathbf{C} \xrightarrow{\deg} \mathfrak{b}(4, \mathbf{R}) \otimes \mathbf{C}$ is the case $\gamma = 2$ in Table IV p. 736 op cit.

criteria consist for a large part⁸ in going back to the Lie algebra and simplifying its structure. These criteria can sometimes be formulated using deformations and degenerations of Lie algebras.

- Pansu’s theorem on asymptotic cones: those are degenerations.
- Cornulier’s theorem on asymptotic cones: when the exponential radical is abelian, those are degenerations. (This is the case for the Heintze groups considered in Section 3.)
- Twistings (or normal modifications) introduced by [Ale75] and [GW88] and studied in relation to large-scale geometry in [CKD⁺21]: those are deformations.

5.1.1. *Cornulier’s Theorem.* Let $\mathfrak{g}$ be a completely solvable Lie algebra. Let $\mathfrak{h}$ be a Cartan subalgebra (maximal nilpotent self-normalizing in $\mathfrak{g}$), and let $\mathfrak{r} = \liminf_i C^i \mathfrak{g}$ be the limit of the descending central series of $\mathfrak{g}$. Decompose the adjoint representation of $\mathfrak{h}$ in $\mathfrak{r}$ into primary components,

$$\mathfrak{r} = \bigoplus_{\omega \in \text{Hom}(\mathfrak{h}, \mathbf{R})} \mathfrak{r}^\omega = \bigoplus_{\omega \in \text{Hom}(\mathfrak{h}, \mathbf{R})} \limsup_{i \rightarrow +\infty} \ker(\alpha - \omega)^i$$

where α is the structural morphism $\mathfrak{h} \rightarrow \text{Der}(\mathfrak{r})$. Note that since $\mathfrak{h}$ is nilpotent, its ideal $\mathfrak{w} = \mathfrak{h} \cap \mathfrak{r}$ lies within $\mathfrak{r}^0$. So the semisimple part δ of α factors through $\pi : \mathfrak{h} \rightarrow \mathfrak{h}/\mathfrak{w}$, and the resulting $\mathfrak{h}/\mathfrak{w}$ -module decomposes as

$$(5.1) \quad \mathfrak{r} = \mathfrak{r}^0 \oplus \bigoplus_{\underline{\omega} \in \text{Hom}(\mathfrak{h}/\mathfrak{w}, \mathbf{R}), \underline{\omega} \neq 0} \mathfrak{r}^{\underline{\omega}} = \mathfrak{r}^0 \oplus \bigoplus_{\underline{\omega} \neq 0} \ker(\underline{\delta} - \underline{\omega})$$

where $\delta = \underline{\delta} \circ \pi$ and $\underline{\omega} = \omega \circ \pi$. There is a Lie algebra homomorphism $\underline{\delta}_\infty : \text{gr}(\mathfrak{h}/\mathfrak{w}) \rightarrow \text{Der}(\mathfrak{r})$ and the following diagram:

$$\begin{array}{ccccc} \mathfrak{h} & \xrightarrow{\pi} & \mathfrak{h}/\mathfrak{w} & \xrightarrow{\quad \underline{\delta} \quad} & \text{Der}(\mathfrak{r}) \\ & & \searrow & \nearrow & \\ & & \mathfrak{h}/[\mathfrak{h}, \mathfrak{h}] & \longrightarrow & \text{Der}(\mathfrak{r}) \\ & \nearrow & \nearrow & \nearrow & \\ \text{gr}(\mathfrak{h}) & \longrightarrow & \text{gr}(\mathfrak{h}/\mathfrak{w}) & \xrightarrow{\quad \underline{\delta}_\infty \quad} & \text{Der}(\mathfrak{r}) \end{array}$$

Theorem 5.1 (Cornulier [Cor11]). *Let $\mathfrak{g}$ be a completely solvable Lie algebra. With notation as above, define $\mathfrak{g}_1 = \mathfrak{r} \rtimes_\delta (\mathfrak{h}/\mathfrak{w})$ and $\mathfrak{g}_\infty$ as $\mathfrak{r} \rtimes_{\underline{\delta}_\infty} \text{gr}(\mathfrak{h}/\mathfrak{w})$. Let G, G_1, G_∞ be simply connected with Lie algebras $\mathfrak{g}, \mathfrak{g}_1, \mathfrak{g}_\infty$ respectively. Then*

- G and G_1 are $O(\log)$ -bilipschitz equivalent.*
- If $C^{s+1}\mathfrak{h} = 0$, then G_1 and G_∞ are $O(r^{1-1/s})$ -bilipschitz equivalent.*

⁸Additional subtlety comes from the “medium-scale” topology of the groups when it is non trivial.

Proposition 5.2. *Let $\mathfrak{g}$ be a completely solvable Lie algebra. Assume that $\mathfrak{r} = \liminf C^i \mathfrak{g}$ is abelian. Let $\mathfrak{g}_1, \mathfrak{g}_\infty$ be as in Theorem 5.1. Then*

$$(5.2) \quad \mathfrak{g} \xrightarrow{\deg} \mathfrak{g}_1 \xrightarrow{\deg} \mathfrak{g}_\infty.$$

We already encountered examples of this:

- When $\mathfrak{g}$ is nilpotent, the right degeneration in (5.2) is Example 3.14. Note that $\mathfrak{r} = 0$ in this case.
- When $\mathfrak{g} = [\mathfrak{g}, \mathfrak{g}] \oplus \mathbf{R}A$ and ad_A is unipotent, the left degeneration in (5.2) is the contraction occurring in Theorem 3.16 (2). $\mathfrak{r}$ is abelian and has codimension 1 in this case.

Proof. Start with the decomposition (5.1). Decompose further $\mathfrak{r}$ into $\mathfrak{r}^0$ and a direct sum of subspaces U_i such that

$$(5.3) \quad \bigoplus_{j \geq i} U_j = \bigoplus_{\omega \neq 0} \ker(\alpha - \omega)^i$$

Since $\mathfrak{h}$ is nilpotent, we have that $\mathfrak{w} = \mathfrak{r} \cap \mathfrak{h} \subseteq \mathfrak{r}^0$. Decompose $\text{Vect}(\mathfrak{g})$ into a direct sum

$$(5.4) \quad \mathfrak{g} = \bigoplus_{i \geq 1} U_i \oplus \mathfrak{r}^0 \oplus \mathcal{H}$$

where $\mathcal{H}$ is a linear subspace of $\text{Vect}(\mathfrak{g})$ representing $\mathfrak{h}/\mathfrak{w}$. Denote by μ , resp. μ_1 , resp. μ_∞ the brackets of the three laws on $\text{Vect}(\mathfrak{g})$.

Set $\varphi_t(u) = t^{n-i}u$ for any $u \in U_i$. Then for all $h \in \mathcal{H}$ and $u \in U_i \cap \mathfrak{r}^\omega$,

$$\begin{aligned} \varphi_t \cdot \mu(h, u) &= \varphi_t^{-1} \mu(h, t^i u) = \varphi_t^{-1} t^i (\omega(h)u + v) \quad \text{where } v \in U_{i-1} \\ &= \omega(h)u + t^{-1-i} t^i v \\ &= \omega(h)u + O(t^{-1}), \end{aligned}$$

so μ contracts to μ_1 through φ . □

Remark 5.3. We do not know whether Proposition 5.2 holds in general. This is because the contraction we used in the proof perturbs in general the brackets in $\mathfrak{r}$. We know no obstruction of the kind expressed in Lemma 3.17 for a degeneration from $\mathfrak{g}$ to $\mathfrak{g}_1$.

A question we would like to raise, in view of Remark 5.3 in particular, is whether the group $R = \exp(\mathfrak{r})$ is a large-scale invariant (if the completely solvable G and G' are $O(u)$ -equivalent, does it hold that $\liminf C^i G \simeq \liminf C^i G'$?). This appears quite difficult to determine in general, because this subgroup is exponentially distorted and gets totally disconnected in the asymptotic cones [Gro93]. Nevertheless, it holds by Cornulier's formula (conedim) and Theorem B that the dimension loss

$$(5.5) \quad \dim R = \text{geodim}(G) - \text{conedim}(G)$$

is indeed a $o(r)$ -bilipschitz invariant. When G is of Heintze type, the $o(r)$ -bilipschitz invariance of (5.5) is materialized into the Gromov boundary; note also that the quasiisometry class of R is a quasiisometry invariant of G [KDG21, Theorem A]; but we have no asymptotic invariant in general. We also note that

the nonnegativity $\text{asdim}_{\text{AN}}(X) - \text{conedim } X \geq 0$ holds more generally, a result of Dydak and Higes [DH08].

5.1.2. Shadows and deformations. Let $\mathfrak{g}_0$ be a completely solvable algebra. We call torus an abelian algebra of semisimple derivations of $\mathfrak{g}_0$. A torus $\mathfrak{t}$ is compactly embedded if every $T \in \mathfrak{t}$ has purely imaginary spectrum. Maximal tori are conjugated.

Definition 5.4 (Special case of [GW88, 2.2]). Let $\mathfrak{t}$ be a maximal compactly embedded torus. A modification⁹ of $\mathfrak{g}_0$ is a transversal subalgebra to $\mathfrak{t}$ in $\mathfrak{g}_0 \rtimes \mathfrak{t}$. We call $\mathfrak{g}_0$ the shadow of $\mathfrak{g}$.

The modification $\mathfrak{g}$ is the graph of a linear map $\tau : \mathfrak{g}_0 \rightarrow \mathfrak{t}$, called the modification map: for $X \in \mathfrak{g}_0$, $\tau(X)$ is the only $T \in \mathfrak{t}$ such that $X + T \in \mathfrak{g}$. Note that $\mathfrak{t}$ being abelian, $[\mathfrak{g}, \mathfrak{g}] \subset \mathfrak{g}_0$.

Definition 5.5. Let $\mathfrak{g}$, $\mathfrak{g}_0$ and τ be as above. We say that $\mathfrak{g}$ is a twisting (and τ a twisting map) if in addition $[\mathfrak{g}, \tau(\mathfrak{g}_0)] \subseteq \mathfrak{g}$.

If $\mathfrak{g}_0$ is nilpotent, all its modifications are twistings [GW88].

Early works on modifications ([Ale75], [GW88]) were concerned by the problem of finding adequate data for the classification of solvmanifolds. Modification have attracted the attention more recently because if $\mathfrak{g}$ is a modification of $\mathfrak{g}_0$, then G , G_0 and $G_0 \rtimes T$ (where T is the compact torus of $\text{Aut}(G_0)$ with Lie algebra $\mathfrak{t}$) share a common Riemannian model, especially they are quasiisometric ([Cor08], [CKD⁺21]).

Proposition 5.6. Let $\mathfrak{g}_0$, $\mathfrak{l}$, $\mathfrak{g}$ and τ be as above. Assume that $\mathfrak{g}$ is a twisting. Define $\omega_\tau(X \wedge Y) = [\tau(X), Y] + [X, \tau(Y)] = d\tau(X \wedge Y) + \tau[X, Y]$ for $X, Y \in \mathfrak{g}_0$. Then,

- (1) $\omega_\tau \in Z^2(\mathfrak{g}_0, \mathfrak{g}_0)$, where $\mathfrak{g}_0$ acts in $\mathfrak{g}_0$ through the adjoint representation.
- (2) $[\omega_\tau]$ is a linearly expandable infinitesimal deformation of $\mathfrak{g}_0$. The associated formal deformation goes through $O_{\mathfrak{g}}$.

Proof. (1) By definition,

$$\begin{aligned} d\omega_\tau(X \wedge Y \wedge Z) &= [X, [\tau(Y), Z] + [Y, \tau(Z)]] - [Y, [\tau(X), Z] + [X, \tau(Z)]] \\ &\quad + [Z, [\tau(X), Y] + [X, \tau(Y)]] - [\tau[X, Y], Z] - [[X, Y], \tau(Z)] \\ &\quad + [\tau[X, Z], Y] + [[X, Z], \tau(Y)] - [\tau[Y, Z], X] - [[Y, Z], \tau(X)] \\ &= [\tau[X, Z], Y] - [\tau[Y, Z], X] - [[Y, Z], \tau(X)] \end{aligned}$$

where we used the Jacobi identity in $\mathfrak{g}_0 \rtimes \mathfrak{l}$ three times. If $\mathfrak{g}$ is a twisting then τ is a homomorphism [GW88], hence the remaining terms can be simplified and vanish using the Jacobi identity again.

(2) Set $\mu_0 \in O_{\mathfrak{g}_0}$ and put $\lambda = \mu_0 + \omega_\tau/t$. Then $\lambda(1) \in O_{\mathfrak{g}}$. □

Beware that it is not true that a twisting $\mathfrak{g}$ degenerates to its shadow $\mathfrak{g}_0$. Here is an already encountered example.

⁹Modification is a more general notion, we only consider modifications of completely solvable Lie algebras for our purposes in the present paper.




FIGURE 4. Sketch of $\mathcal{N}_6(\mathbf{R}) \subseteq \mathcal{L}_6(\mathbf{R})$ around the Carnot Lie algebra from Example 5.8, and some deformations.

Example 5.7 (Solvable example). Let $\mathfrak{g}_0 = \mathfrak{h}(3, \mathbf{R})$, with basis (X_1, X_2, T) and brackets

$$(5.6) \quad [X_1, X_2] = 0, [T, X_1] = X_1, [T, X_2] = X_2.$$

Let (dx_1, dx_2, dt) be the dual basis. Then $H^2(\mathfrak{g}_0, \mathfrak{g}_0)$ is 3-dimensional, generated by the classes $\omega_1 = [dt \wedge dx_1 \otimes X_2]$ and $\omega_2 = [dt \wedge dx_2 \otimes X_1]$. ω_1 and ω_2 are linearly expandable into degenerations, but $\omega_1 - \omega_2$ is linearly expandable into a family of twistings that are not degenerations. See Appendix A.1 for a more general computation.

If $\mathfrak{h}$ is a graded Lie algebra and $\mu \in O_{\mathfrak{h}}$, the groups $H^2(\mu, \mu)$ are naturally graded. This is the case, for instance, if $\mathfrak{h}$ is a Carnot-graded group.

Example 5.8 (A nilpotent example). Let G_0 be the simply connected 6-dimensional Lie group having Lie algebra over $X_1, \dots, X_6$ and the nonzero brackets

$$[X_1, X_2] = X_3, [X_1, X_3] = X_4, [X_1, X_4] = X_5.$$

(This algebra is denoted $\mathfrak{l}_{6,7}$ in [dG07].) Note that X_6 generates an abelian direct factor. $\mathfrak{g}_0$ is a Carnot-graded algebra under the grading

$$\langle X_1, X_2, X_6 \rangle \oplus \langle X_3 \rangle \oplus \langle X_4 \rangle \oplus \langle X_5 \rangle.$$

Let $(X^1, \dots, X^6)$ be the dual basis, and denote by μ the law. Consider the following cochains:

$$\omega = X_2^{16} + X_1^{62}; \quad \xi_1 = X_5^{23}; \quad \xi_2 = X_5^{26}; \quad \xi_3 = X_4^{26} + X_5^{36}.$$

where we abbreviate $X_k \otimes X^i \wedge X^j$ into X_k^{ij} . These are cocycles but not coboundaries in $H^2(\mu, \mu)$ (See the computations in Appendix A.2.1). The cohomology classes of ω , ξ_1 , ξ_2 and ξ_3 have weight -1 , 1 , 2 and 1 respectively under the grading. The classes of ξ_1 , ξ_2 , ξ_3 and $\xi_1 + \xi_2$ linearly expand into formal deformations; the corresponding laws are $\mathfrak{l}_{6,6}$, $\mathfrak{l}_{6,12}$, $\mathfrak{l}_{6,13}$ and $\mathfrak{l}_{6,11}$ respectively in [dG07]. All these are also degenerations, entering into the description of Example 3.14 (adding cocycles with positive weights to the Lie algebra law does not change the lower central filtration). For every $k \in \mathbf{R}$, the cohomology class of $k\xi_1 + \omega$ is also integrable into a formal deformation, going through families of twistings of $\mathfrak{g}_0 = \mathfrak{l}_{6,7}$ if $k = 0$, or through families of twistings of $\mathfrak{l}_{6,6}$ if $k \neq 0$.

One can check that all the simply connected solvable Lie groups that are $O(u)$ -bilipschitz equivalent to G_0 appear as deformations of G_0 of the form described above. Let us give a few words on this. If H is such a group, then by [Pan89b] and [Bre14], the shadow $\mathfrak{h}_0$ of its Lie algebra is isomorphic to $\mathfrak{l}_{6,i}$ with $i \in \{6, 7, 11, 12, 13\}$. The three last algebras are irreducible (they have no direct factor), and maximal tori for those are computed in [Mag07]; in this way we check that only $\mathfrak{l}_{6,6}$ and $\mathfrak{l}_{6,7}$ possess derivations with purely imaginary spectra. Thus either $\mathfrak{h}$ is $\mathfrak{l}_{6,11}$, $\mathfrak{l}_{6,12}$, $\mathfrak{l}_{6,13}$ or a twisting of $\mathfrak{l}_{6,7}$ or $\mathfrak{l}_{6,6}$. Note that the quasiisometry classes in this family are not completely known, though it is expected that they are given by the isomorphism type of the shadow [Cor18, Conjecture 19.114]. The real cohomology rings $H^*(\mathfrak{l}_{6,6}, \mathbf{R})$ and $H^*(\mathfrak{l}_{6,7}, \mathbf{R})$ are isomorphic [Cor18, §19.6.6]. However, $b_2(\mathfrak{l}_{6,13})$ is 4 while it is 5 for all the others; and we can check that the rank of $H^2(\mathfrak{l}_{6,i}, \mathbf{R}) \odot H^2(\mathfrak{l}_{6,i}, \mathbf{R}) \rightarrow H^4(\mathfrak{l}_{6,i}, \mathbf{R})$ given by the cup product is 2 for $i = 6$ while it is 3 for $i = 11$ and $i = 12$ (see A.2.2 for some details). All these algebras have $\mathbf{Q}$ -forms. So none of the Lie groups $L_{6,11}$, $L_{6,12}$, $L_{6,13}$ are quasiisometric to G_0 by [Sau06].

It seems natural to expect Carnot graded algebra to have more twistings than their nilpotent deformations (on the other extreme, observe that characteristically nilpotent Lie algebras, which lie “deep down” in $\mathcal{N}_n(\mathbf{R})$, have no twistings). This is indeed the situation on the previous example. Thus we ask:

Question 5.9. Let $\mathfrak{h}$ be a solvable Lie algebra over $\mathbf{R}$, H its associated simply connected Lie group, and let G_0 be a simply connected Carnot-group. Assume that H and G_0 are $O(u)$ -bilipschitz. Is there a formal deformation of $\mathfrak{g}_0$ going through $\mathfrak{h}$?

The author does not know whether the dimension of compactly embedded maximal tori is upper semicontinuous on $\mathcal{N}_n(\mathbf{R})$ (which would hint towards a positive answer to Question 5.9). These tori embed linearly in $H^1(\lambda, \lambda)$ whose dimension we have seen to be upper semicontinuous in Lemma 3.17. However the codimension of the tori may be high (see Table 3).

5.2. Higher-rank symmetric spaces. The real rank of a symmetric space X is $o(r)$ -bilipschitz invariant, as it is the covering dimension of asymptotic cone [Cor08] or, more in line with [KL97, Corollary 6.11], the minimal degree above which all relative homology group of subspaces in $\text{Cone}_\omega^\bullet X$ vanish. This can be refined: the restricted root system is invariant.

$\mathfrak{g}$	$\dim \mathfrak{t}_{\max}^c$	$\dim \mathfrak{t}_{\max}$	$\dim H^1(\mathfrak{g}, \mathfrak{g})$
$\mathfrak{l}_{6,7}$	1	3	9 [Mag08, $\mathfrak{g}_{5,5} \times \mathbf{C}$]
$\mathfrak{l}_{6,6}$	1	2	8 [Mag08, $\mathfrak{g}_{6,5} \times \mathbf{C}$]
$\mathfrak{l}_{6,12}$	0	2 [Mag07, 4.2.5]	7 [Mag08, $\mathfrak{g}_{6,11}$]
$\mathfrak{l}_{6,11}$	0	1 [Mag07, 4.1.1]	6 [Mag08, $\mathfrak{g}_{6,12}$]
$\mathfrak{l}_{6,13}$	0	2 [Mag07, 4.2.6]	5 [Mag08, $\mathfrak{g}_{6,13}$]

TABLE 3. Dimensions of maximal tori, compactly embedded maximal tori, and outer derivation spaces for the nilpotent Lie algebras of Example 5.8.

Proposition 5.10 (After Kleiner and Leeb). *Let $\phi : X \rightarrow Y$ be a sublinear bilipschitz equivalence between irreducible symmetric spaces X of rank ≥ 2 . Then, the restricted root systems associated with X and Y are isomorphic.*

Proof. The spherical Tits building at infinity in $\text{Cone}_\omega(X)$ has the same apartments as the Tits boundary of X [KL97, Theorem 5.2.1]. $\square$

We note that the rank p irreducible symmetric spaces of noncompact type

$$\text{SU}(p, 2q)/S(\mathbf{U}_p \times \mathbf{U}_{2q}) \text{ and } \text{Sp}(p, q)/\text{Sp}(p) \times \text{Sp}(q)$$

have same restricted root system BC_p and same asymptotic Assouad-Nagata dimension $4pq$ [Hel01, Table V p. 518]. Thus, we could not distinguish them with our techniques, and Question 1.2 remains open so far for them.

The author is grateful to P. Pansu and G. Rousseau for bringing these pairs to his attention.

5.3. Right-angled Fuchsian buildings of uniform thickness. Given (p, q) such that $p \geq 5$ and $q \geq 2$, the finitely presented group

$$\Gamma_{p,q} = \langle s_1, \dots, s_p \mid [s_i, s_{i+1}], s_i^q \rangle.$$

has a model $I_{p,q}$ which is a $\text{CAT}(-1)$ cellular complex generalizing the cellular action of the hyperbolic Coxeter group $\Gamma_{p,2}$ on $\mathbb{H}_{\mathbf{R}}^2$ tessellated by right-angled p -gons and, following [Bou97],

$$(5.7) \quad \text{Cdim } \partial_\infty I_{p,q} = \frac{\log \tau(p, q)}{\log \tau(p, 2)} = 1 + \frac{\log(q-1)}{\argch\left(\frac{p-2}{2}\right)}.$$

The conformal dimension of $I_{p,q}$ is not rational unless $q = 2$. It is proven in [Pal20] that $\text{Cdim}_{O(u)} \partial_\infty I_{p,q} = \text{Cdim } \partial_\infty I_{p,q}$, so that it is a $O(u)$ -bilipschitz invariant. Using Poincaré profiles, Hume, Mackay and Tessera proved that there can be no coarse embedding $I_{p,q} \rightarrow I_{p',q'}$ when $\text{Cdim } \partial_\infty I_{p,q} > \text{Cdim } \partial_\infty I_{p',q'}$ [HMT20a, Theorem 13.2]. We found the equality case in (5.7) to be related to the following conjecture.

Conjecture 5.11 (Four exponential conjecture, [Lan66, p.11]). *Let β_1, β_2 be complex numbers, linearly independent over $\mathbf{Q}$, and let z_1, z_2 be complex numbers, also linearly independent over $\mathbf{Q}$. Then, at least one of the numbers $e^{\beta_i z_j}$ is transcendental.*

The analogous statement with two triples $\beta_1, \beta_2, \beta_3, z_1, z_2, z_3$ is known as the six exponentials theorem [Lan66]. The unconditional form of the following Proposition is stated as a conjecture in [Tys99] and [MT10]. (We indicate with an asterisk that our statement is conditional.)

Proposition * 5.12. *Assume that Conjecture 5.11 holds, and let (p, q, p', q') be integers such that $p, p' \geq 5$ and $q, q' \geq 3$. Then the buildings $I_{p,q}$ and $I_{p',q'}$ have equal conformal dimension if and only if there exists positive integers M, N such that*

$$(5.8) \quad (q-1)^N = (q'-1)^M$$

$$(5.9) \quad T_N\left(\frac{p-2}{2}\right) = T_M\left(\frac{p'-2}{2}\right)$$

where T_k is the Tchebychev polynomial of the first kind and degree k .

Proof. Negating the conclusion amounts to assert that there exists an irrational number z and a quadruple (p, q, p', q') such that $z \log(q'-1) = \log(q-1)$ and $z \operatorname{argch}((p-2)/2) = \operatorname{argch}((p'-2)/2)$. Define $\beta_1 = \log q' - 1$,

$$\beta_2 = \operatorname{argch}((p'-2)/2) = \log\left(\frac{p'-2}{2} + \sqrt{p'(p'/4-1)}\right),$$

$z_1 = 1$ and $z_2 = z$. Then, β_2/β_1 is not rational. But $e^{\beta_1}, e^{z\beta_1} = q-1, e^{\beta_2}$ and $e^{z\beta_2} = \frac{p-2}{2} + \sqrt{p(p/4-1)}$ are all algebraic. $\square$

Note that the $I_{p,q}$ are quasiisometrically rigid for $q \geq 3$ [Xie06]; especially they are classified up to quasiisometry by the pair (p, q) for $q \geq 3$.

Proposition * 5.13. *Assume that Conjecture 5.11 holds. If there exists a $O(u)$ -bilipschitz equivalence $\phi : I_{p,q} \rightarrow I_{p',q'}$ then (5.8) and (5.9) hold for some $M, N \geq 1$.*

Proof. This directly follows from Proposition 5.12 and [Pal20]. $\square$

Let us finish with some questions. Though we consider (5.8) and (5.9) perhaps not sufficient for $O(u)$ -equivalence between $I_{p,q}$ and $I_{p',q'}$, we could not distinguish them up to this relation. In a slightly different direction, one can ask:

Question 5.14. Assume that p, q, p', q' are as in (5.8) and (5.9). Are the groups $\Gamma_{p,q}$ and $\Gamma_{p',q'}$ (non)-measure equivalent? If yes, are they L^p -measure equivalent for some $p < +\infty$?

(We recall that a measure equivalence between the finitely generated Γ and Λ is given by a couple of free, commuting, measure preserving actions of Γ and Λ on a Lebesgue space (Ω, m) with Borel fundamental domains of finite measure X and Y , such that the associated cocycles $c : G \times X \rightarrow H$ and $H \times Y \rightarrow G$ have $c(g, \cdot) \in L^p$ for all g .)

Closer to the problems of this paper, the author also believe the following question to remain currently open, and of some interest in view of [HMT20a].

Question 5.15. Assume that p, q, p', q' are as in (5.8) and (5.9), and $p \neq p'$. Is there a coarse embedding $I_{p,q} \rightarrow I_{p',q'}$?

APPENDIX A. METHODS USED FOR THE COHOMOLOGY COMPUTATIONS

The cohomology groups used in this paper are obtained by direct methods (i.e. by somewhat explicit computations of derivative, cocycles and coboundaries). We summarize them below.

A.1. Solvable Lie algebras. Let $\mathfrak{b}(n, \mathbf{K})$ be defined as in Example 3.4, with coordinates (z_α, τ, s) . Decompose $z_\alpha = x_\alpha + iy_\alpha$; for $1 \leq \alpha_1 < \dots < \alpha_s \leq n-1$ and $1 \leq \beta \leq n-1$, denote

$$\begin{aligned} X_\beta^{\alpha_1, \dots, \alpha_s} &= dx_{\alpha_1} \wedge \dots \wedge dx_{\alpha_s} \otimes \frac{\partial}{\partial x_\beta} \\ Y_\beta^{\alpha_1, \dots, \alpha_s} &= dy_{\alpha_1} \wedge \dots \wedge dy_{\alpha_s} \otimes \frac{\partial}{\partial y_\beta} \\ T &= \partial_\tau, \quad T^* = d\tau, \quad S = \partial_s, \quad S^* = ds. \end{aligned}$$

We apply the summation convention where we simplify $\sum_\mu X_\alpha^{\alpha\mu}$ into $X_\alpha^{\alpha\mu}$ in any equality between tensors whenever μ is unbound in the RHS.

The Lie algebra grading $\mathfrak{s}_0 = \langle S \rangle$, $\mathfrak{s}_1 = \langle X_\alpha, Y_\alpha \rangle$ and $\mathfrak{s}_2 = \langle T \rangle$ extends to a grading of the mixed exterior/tensor product so that, say, $X_\beta^{\alpha_1, \dots, \alpha_s} \wedge T$ has weight $1-s+2$. The differentials have degree 0, hence the cohomology groups are graded accordingly. Finally, $\mathfrak{b}(n, \mathbf{C})$ has a preferred complex structure, $JX_\alpha = Y_\alpha$, $JY_\alpha = -X_\alpha$ and $JS = T$. This is because $\mathbf{H}_\mathbf{C}^n$ is Hermitian. J is not an automorphism; nevertheless,

$$\tilde{J}(Z) = \begin{cases} J(Z) & Z \in \mathfrak{s}_1 \\ Z & Z \in \mathfrak{s}_0 \oplus \mathfrak{s}_2 \end{cases}$$

is an automorphism, and we will use it in order to simplify the computations.

A.1.1. Results.

Proposition A.1. $H^1(\mathfrak{b}(n, \mathbf{R}), \mathfrak{b}(n, \mathbf{R})) = \bigoplus_{(\alpha, \beta) \neq (n-1, n-1)} \langle X_\alpha^\beta \rangle$.

Proposition A.2. $H^2(\mathfrak{b}(n, \mathbf{R}), \mathfrak{b}(n, \mathbf{R})) = \bigoplus_{(\alpha, \beta) \neq (n-1, n-1)} \langle [X_\alpha \otimes S^* \wedge X^\beta]_{\alpha \neq \beta} \rangle$.

Since the computation of $H^1(\mathfrak{b}(n, \mathbf{C}), \mathfrak{b}(n, \mathbf{C}))$ proves useful for Proposition A.1 and is not significantly harder than the case $n=2$ used in Section 4, we provide the result for all n and the weight decomposition below.

Proposition A.3. $\dim H^1(\mathfrak{b}(n, \mathbf{C}), \mathfrak{b}(n, \mathbf{C})) = (n-1)^2 + 1$ and

$$H^1(\mathfrak{b}(n, \mathbf{C}), \mathfrak{b}(n, \mathbf{C})) = \text{span} \begin{cases} [T \otimes S^*] & \text{weight } -2 \\ [X_\alpha^\beta - Y_\beta^\alpha]_{1 \leq \alpha, \beta \leq n-1, \alpha \neq \beta} & \text{weight } 0 \\ [X_\alpha \otimes Y^\alpha]_{1 \leq \alpha \leq n-1} & \text{weight } 0. \end{cases}$$

Proposition A.4. Let $\mathfrak{s}''$ be the four-dimensional Lie algebra $\mathbf{R}^3 \rtimes_\alpha \mathbf{R}$, where $\alpha = \text{diag}(J_2(1), 2)$. Then $\dim H^1(\mathfrak{s}'', \mathfrak{s}'') = 4$.

A.1.2. *Method.* In order to gain space for Propositions A.1 to A.3 we gather the computation for $\mathbf{R}$ and $\mathbf{C}$ and then extract the case of $\mathbf{K} = \mathbf{R}$. We abbreviate the derivative of the complex $C^\bullet(\mathfrak{b}(n, \mathbf{K}), \mathfrak{b}(n, \mathbf{K}))$ into $d'_\mathbf{K}$.

Lemma A.5. *For all α, β such that $1 \leq \alpha, \beta \leq n-1$,*

$$\begin{aligned} d'_\mathbf{C} X_\alpha^\beta &= X^\beta \wedge Y^\alpha \otimes T; & d'_\mathbf{C} Y_\alpha^\beta &= X^\alpha \wedge Y^\beta \otimes T; \\ d'_\mathbf{C}(X_\alpha \otimes Y^\beta) &= -2Y^{\alpha\beta} \otimes T; & d'_\mathbf{C}(Y_\alpha \otimes X^\beta) &= 2X^{\alpha\beta} \otimes T; \\ d'_\mathbf{C}(X_\alpha \otimes S^*) &= -Y^\alpha \wedge S^* \otimes T; & d'_\mathbf{C}(Y_\alpha \otimes S^*) &= X^\alpha \wedge S^* \otimes T. \\ d'_\mathbf{C}(T \otimes X^\alpha) &= X^\alpha \wedge S^* \otimes T; & d'_\mathbf{C}(T \otimes Y^\alpha) &= Y^\alpha \wedge S^* \otimes T \\ d'_\mathbf{C}(T \otimes S^*) &= 0 & d'_\mathbf{C}(S \otimes T^*) &= -2S^* \wedge T \otimes S. \end{aligned}$$

$$d'_\mathbf{C}(S \otimes S^*) = -X^\mu \wedge S^* \otimes X_\mu - Y^\mu \wedge S^* \otimes Y_\mu - 2T \wedge S^* \otimes T.$$

$$d'_\mathbf{C}(T \otimes T) = -X^\mu \wedge Y^\mu \otimes T.$$

$$d'_\mathbf{C}(S \otimes X^\alpha) = X_\ell^{\alpha\ell} - X^\alpha \wedge Y^\ell \otimes Y_\ell + X^\alpha \wedge S^* \otimes S - 2X^\alpha \wedge T^* \otimes T;$$

$$d'_\mathbf{C}(S \otimes Y^\alpha) = Y_\ell^{\alpha\ell} - Y^\alpha \wedge X^\ell \otimes X_\ell + Y^\alpha \wedge S^* \otimes S - 2Y^\alpha \wedge T^* \otimes T;$$

$$d'_\mathbf{C}(X_\alpha \otimes T^*) = (X^\mu \wedge Y^\mu - S^* \wedge T) \otimes X_\alpha + Y^\alpha \wedge T^* \otimes T$$

$$d'_\mathbf{C}(Y_\alpha \otimes T^*) = (-Y^\mu \wedge X^\mu - S^* \wedge T) \otimes Y_\alpha - X^\alpha \wedge T^* \otimes T$$

Proof. The whole computation being of little interest, let us explain in detail only how one computes $d_\mathbf{C} X_\alpha^\beta$, $d_\mathbf{C} Y_\alpha^\beta$ and $d(X_\alpha \otimes S^*)$ as a sample of the techniques employed. Applying (3.9),

$$\begin{aligned} d'_\mathbf{C} X_\alpha^\beta(X_{\mu\ell}) &= -X_\alpha^\beta[X_\mu, X_\ell] + [X_\mu, X_\alpha^\beta X_\ell] - [X_\ell, X_\alpha^\beta X_\mu] \\ &= [X_\mu, \delta_{\beta\ell} X_\alpha] - [X_\ell, \delta_{\beta\mu} X_\alpha] = 0; \\ d'_\mathbf{C} X_\alpha^\beta(Y_{\mu\ell}) &= -X_\alpha^\beta[Y_\mu, Y_\ell] + [Y_\mu, X_\alpha^\beta Y_\ell] - [Y_\ell, X_\alpha^\beta Y_\mu] = 0; \\ d'_\mathbf{C} X_\alpha^\beta(X_\mu \wedge Y_\ell) &= -X_\alpha^\beta[X_\mu, Y_\ell] + [X_\mu, X_\alpha^\beta Y_\ell] - [Y_\ell, X_\alpha^\beta X_\mu] \\ &= -\delta_{\mu\ell} X_\alpha^\beta T - \delta_{\beta\mu} [Y_\ell, X_\alpha] = \delta_{\alpha\ell} \delta_{\beta\mu} T; \\ d'_\mathbf{C} X_\alpha^\beta(X_\mu \wedge S) &= -X_\alpha^\beta[X_\mu, S] + [X_\mu, X_\alpha^\beta S] - [S, X_\alpha^\beta X_\mu] \\ &= X_\alpha^\beta X_\mu - [S, \delta_{\beta\mu} X_\alpha] = \delta_{\beta\mu} (X_\alpha - X_\alpha) = 0, \\ d'_\mathbf{C} X_\alpha^\beta(X_\mu \wedge T) &= -X_\alpha^\beta[X_\mu, T] + [X_\mu, X_\alpha^\beta T] - [T, X_\alpha^\beta X_\mu] = -[T, \delta_{\beta\mu} X_\alpha] = 0. \\ d'_\mathbf{C} X_\alpha^\beta(Y_\mu \wedge S) &= -X_\alpha^\beta[Y_\mu, S] + [Y_\mu, X_\alpha^\beta S] - [S, X_\alpha^\beta Y_\mu] = X_\alpha^\beta Y_\mu = 0. \\ d'_\mathbf{C} X_\alpha^\beta(Y_\mu \wedge T) &= -X_\alpha^\beta[Y_\mu, T] + [Y_\mu, X_\alpha^\beta T] - [T, X_\alpha^\beta Y_\mu] = 0; \\ d'_\mathbf{C} X_\alpha^\beta(S \wedge T) &= -X_\alpha^\beta[S, T] + [S, X_\alpha^\beta T] - [T, X_\alpha^\beta S] = -2X_\alpha^\beta T = 0, \end{aligned}$$

which yields the expression of $d_\mathbf{C} X_\alpha^\beta$. Applying $\tilde{J}$ produces $d_\mathbf{C} Y_\alpha^\beta$:

$$d'_\mathbf{C} Y_\alpha^\beta = d_\mathbf{C} \tilde{J} X_\alpha^\beta = \tilde{J} X^\beta \wedge \tilde{J} X^\alpha \otimes T = Y^\beta \wedge (-X^\alpha) \otimes T.$$

Now for $d'_\mathbf{C}(X_\alpha \otimes S^*)(X_\mu \wedge S)$, using the observation that $\ker S^* = \mathfrak{s}_1 \oplus \mathfrak{s}_2 = [\mathfrak{s}, \mathfrak{s}]$, we can reduce the number of terms needed for the computations of $d(X_\alpha \otimes S^*)$: this will evaluate to zero for any bivector where S is not a factor. The remaining

terms are:

$$\begin{aligned} d'_C(X_\alpha \otimes S^*)(X_\mu \wedge S) &= [X_\mu, X_\alpha \otimes S^*S] = 0; \\ d'_C(X_\alpha \otimes S^*)(Y_\mu \wedge S) &= [Y_\mu, X_\alpha \otimes S^*S] = [Y_\mu, X_\alpha] = -\delta_{\alpha\mu}T; \\ d'_C(X_\alpha \otimes S^*)(S \wedge T) &= -[T, X_\alpha \otimes S^*S] + [S, X_\alpha \otimes S^*T] = 0. \end{aligned} \quad \square$$

Lemma A.6. For all α, β such that $1 \leq \alpha, \beta \leq n-1$,

$$\begin{aligned} d'_R X_\alpha^\beta &= 0 \\ d'_R(S \otimes X^\alpha) &= X_\ell^{\alpha\ell} + X^\alpha \wedge S^* \otimes S \\ d'_R(X_\alpha \otimes S^*) &= 0 \\ d'_R(S \otimes S^*) &= -X^\mu \wedge S^* \otimes X^\mu. \end{aligned}$$

Proof. Discard the terms with Y, S, T in the results of Lemma A.5. $\square$

Lemma A.7 (Differentials of 2-cochains).

$$\begin{aligned} d'_R(X_\alpha^{\beta\gamma}) &= -2X_\alpha \otimes X^{\beta\gamma} \wedge S^* \\ d'_R(S \otimes X^{\alpha\beta}) &= -2S \otimes X^{\alpha\beta} \wedge S^* \\ d'_R(X_\alpha \otimes S^* \wedge X^\beta) &= 0 \\ d'_R(S \otimes S^* \wedge X^\alpha) &= 2X_\mu \otimes X^{\mu\alpha} \wedge S^*. \end{aligned}$$

Proof. Let us concentrate on $d'_R(X_\alpha^{\beta\gamma})$. First recall that if $\mathfrak{g}$ is a Lie algebra and ω is a $\mathfrak{g}$ -valued 2-form, then for every $U, V, W \in \mathfrak{g}^3$,

$$\begin{aligned} d\gamma(U \wedge V \wedge W) &= [U, \gamma(V \wedge W)] + [V, \gamma(W, U)] + [W, \gamma(U, V)] \\ &\quad - \gamma([U, V] \wedge W) - \gamma([W, U] \wedge V) - \gamma([V, W] \wedge U). \end{aligned}$$

Applying this, one checks readily that $d'_R(X_\alpha^{\beta\gamma})(X_{\mu\nu\ell}) = 0$ while

$$\begin{aligned} d'_R(X_\alpha^{\beta\gamma})(X_{\mu\nu} \wedge S) &= 0 - X_\alpha^{\beta\gamma}([X_\mu, X_\nu] \wedge S + [S, X_\mu] \wedge X_\nu + [X_\nu, S] \wedge X_\mu) \\ &= (-\delta_{\beta\mu}\delta_{\gamma\nu} + \delta_{\beta\nu}\delta_{\gamma\mu})X_\alpha. \end{aligned} \quad \square$$

Remark A.8. The Lie algebra-valued forms have a wedge product. However we did not find a clear computational advantage in using formulae for the derivative of 2-forms using this wedge product.

Proof of Proposition A.1. By Lemma A.6,

$$Z^1(\mathfrak{b}(n, \mathbf{R}), \mathfrak{b}(n, \mathbf{R})) = \text{span} \{X_\alpha^\beta, X_\alpha \otimes S^*\}_{1 \leq \alpha, \beta \leq n-1},$$

while $d'_R X_\alpha = X_\alpha \otimes S^*$ and $d'_R S = -X_\mu^\mu$. $\square$

Proof of Proposition A.2. By Lemma A.6,

$$B^2(\mathfrak{b}(n, \mathbf{R}), \mathfrak{b}(n, \mathbf{R})) = \text{span} \{X_\ell^{\alpha\ell} + X^\alpha \wedge S^* \otimes S, X_\mu \otimes X^\mu \wedge S^*\}_{\alpha=1, \dots, n-1}$$

while by Lemma A.7,

$$Z^2(\mathfrak{b}(n, \mathbf{R}), \mathfrak{b}(n, \mathbf{R})) = \text{span} \{X_\mu^{\mu\alpha} + S \otimes S^* \wedge X^\alpha, X_\alpha \otimes X^\beta \wedge S^*\}_{\alpha=1, \dots, n-1}.$$

Proof of Proposition A.3. The 1-coboundaries are computed as

$$\begin{aligned} d'_C X_\alpha &= X_\alpha \otimes S^* - T \otimes Y_\alpha \\ d'_C Y_\alpha &= Y_\alpha \otimes S^* + T \otimes X_\alpha \\ d'_C S &= -X_\mu^\mu - Y_\mu^\mu - 2T \otimes T^* \\ d'_C T &= 2T \otimes S^*. \end{aligned}$$

The right-hand side of equations in Lemma A.5 provide the 1-cocycles. $\square$

Proof of Proposition A.4. We recall that $\mathfrak{s}''$ is the Lie algebra over $X_1, \dots, X_4$ with $\mathfrak{s}'' = \langle X_2, X_3, X_4 \rangle \oplus \langle X_4 \rangle$ and

$$\text{ad}(X_4) = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{pmatrix}$$

in the basis (X_2, X_3, X_4) .

Omitting the symbol $\sum_{i < j}$ and using $d'(X_k^\ell)(x^{ij}X_{ij}) = -x^{ij}X_k^\ell[X_i, X_j] + \delta_{j\ell}x^{ij}[X_i, X_k] - \delta_{i\ell}x^{ij}[X_j, X_k]$ one finds

$$\begin{aligned} d'(X_1^1) &= 2X_1^{14} + X_1^{24} & d'(X_1^2) &= -X_1^{24} + X_2^{24} \\ d'(X_1^3) &= -X_1^{12} + 3X_1^{34} & d'(X_1^4) &= 0 \\ d'(X_2^1) &= X_2^{14} & d'(X_2^2) &= X_1^{24} \\ d'(X_2^3) &= -X_1^{34} - 3X_2^{34} & d'(X_2^4) &= 0 \\ d'(X_3^1) &= -X_3^{14} & d'(X_3^2) &= -X_3^{24} \\ d'(X_3^3) &= 0 & d'(X_3^4) &= 0 \\ d'(X_4^1) &= X_4^{14} + X_4^{24} + X_1^{14} & d'(X_4^2) &= X_4^{24} - X_1^{12} + 2X_3^{23} \\ d'(X_4^3) &= 2X_4^{34} - X_1^{13} - X_1^{23} - X_2^{23} & d'(X_4^4) &= -X_1^{14} - X_1^{24} - X_2^{24} - X_3^{34}. \end{aligned}$$

All the nonzero co-boundaries obtained are linearly independent, hence

$$H^1(\mathfrak{s}'', \mathfrak{s}'') = \text{span}(X_3^3, X_1^4, X_2^4, X_3^4). \quad \square$$

A.2. The Lie algebra $\mathfrak{l}_{6,7}$ and its nilpotent deformations. We expand below on the computations needed for Example 5.8.

A.2.1. Adjoint cohomology of $\mathfrak{l}_{6,7}$. One computes the 2-coboundaries as:

$$\begin{aligned} d'_\mu(X_1^1) &= X_3^{12} + X_4^{13} + X_5^{14} & d'_\mu(X_2^1) &= 0 & d'_\mu(X_3^1) &= 0 \\ d'_\mu(X_1^2) &= X_4^{23} + X_5^{24} & d'_\mu(X_2^2) &= X_3^{12} & d'_\mu(X_3^2) &= X_4^{12} \\ d'_\mu(X_1^3) &= -X_3^{23} + X_5^{34} & d'_\mu(X_2^3) &= X_3^{13} & d'_\mu(X_3^3) &= X_4^{13} \\ d'_\mu(X_1^4) &= -X_3^{24} - X_4^{34} & d'_\mu(X_2^4) &= X_3^{14} & d'_\mu(X_3^4) &= X_4^{14} \\ d'_\mu(X_1^5) &= -X_3^{25} - X_4^{35} - X_5^{45} & d'_\mu(X_2^5) &= X_3^{15} & d'_\mu(X_3^5) &= X_4^{15} \\ d'_\mu(X_1^6) &= -X_3^{26} - X_4^{36} - X_5^{46} & d'_\mu(X_2^6) &= X_3^{16} & d'_\mu(X_3^6) &= X_4^{16} \\ d'_\mu(X_4^1) &= 0 & d'_\mu(X_5^1) &= 0 & d'_\mu(X_6^1) &= 0 \\ d'_\mu(X_4^2) &= X_5^{12} & d'_\mu(X_5^2) &= 0 & d'_\mu(X_6^2) &= 0 \\ d'_\mu(X_4^3) &= -X_4^{12} + X_5^{13} & d'_\mu(X_5^3) &= -X_5^{12} & d'_\mu(X_6^3) &= -X_6^{12} \\ d'_\mu(X_4^4) &= -X_4^{14} + X_5^{15} & d'_\mu(X_5^4) &= -X_5^{13} & d'_\mu(X_6^4) &= -X_6^{13} \\ d'_\mu(X_4^5) &= -X_4^{15} + X_5^{15} & d'_\mu(X_5^5) &= -X_5^{14} & d'_\mu(X_6^5) &= -X_6^{14} \\ d'_\mu(X_4^6) &= X_5^{16} & d'_\mu(X_5^6) &= 0 & d'_\mu(X_6^6) &= 0. \end{aligned}$$

This justify the assertion that ω , ξ_1 , ξ_2 , ξ_3 and $\xi_1 + \xi_2$ are not coboundaries. We now check that they are cocycles. In the computation below, we omit the symbols $\sum_{i < j < k}$, and get rid of the terms that can be checked to equal 0 by direct inspection.

$$\begin{aligned} d'_\mu \omega(x^{ijk} X_{ijk}) &= d'_\mu (X_2^{16} + X_1^{62})(x^{ijk} X_{ijk}) \\ &= -x^{ijk} [X_j, X_2^{16} X_{ik}] - x^{ijk} [X_i, X_1^{26} X_{jk}] + x^{ijk} [X_j, X_1^{26} X_{ik}] = 0; \end{aligned}$$

$$\begin{aligned} d'_\mu \xi_1(x^{ijk} X_{ijk}) &= d'_\mu (X_5^{23})(x^{ijk} X_{ijk}) \\ &= -x^{ijk} X_5^{23} ([X_i, X_j] \wedge X_k) + x^{ijk} X_5^{23} ([X_i, X_k] \wedge X_j) \\ &\quad - x^{ijk} X_5^{23} ([X_j, X_k] \wedge X_i) + x^{ijk} [X_i, X_5^{23} X_{jk}] - x^{ijk} [X_j, X_5^{23} X_{ik}] \\ &\quad + x^{ijk} [X_k, X_5^{23} X_{ij}] \\ &= x^{123} [X_1, X_5^{23} X_{23}] + x^{234} [X_4, X_5] + x^{235} [X_5, X_5] = 0; \end{aligned}$$

$$\begin{aligned} d'_\mu \xi_2(x^{ijk} X_{ijk}) &= d'_\mu (X_5^{26})(x^{ijk} X_{ijk}) \\ &= x^{ijk} [X_i, X_5^{26} X_{jk}] - x^{ijk} [X_j, X_5^{26} X_{ik}] + x^{ijk} [X_k, X_5^{26} X_{ij}] = 0; \end{aligned}$$

$$\begin{aligned} d'_\mu \xi_3(x^{ijk} X_{ijk}) &= d'_\mu (X_4^{26} + X_5^{36})(x^{ijk} X_{ijk}) \\ &= x^{126} [X_1, X_4^{26} X_{26}] - x^{126} X_5^{36} ([X_1, X_2] \wedge X_6) \\ &= x^{126} (X_5 - X_5) = 0. \end{aligned}$$

Note that $\dim H^2(\mu, \mu) = 18$, by the computer-produced [Mag08, Table 11].

A.2.2. *Cohomology rings.* Let d_i denote the derivative of $C^*(\mathfrak{l}_{6,i}, \mathbf{R})$. Then

$$\begin{array}{lll} d_7 X^3 = -X^{12} & d_7 X^4 = -X^{13} & d_7 X^5 = -X^{14} \\ d_{11} X^3 = -X^{12} & d_{11} X^4 = -X^{13} & d_{11} X^5 = -X^{14} - X^{23} \\ d_{12} X^3 = -X^{12} & d_{12} X^4 = -X^{13} & d_{12} X^5 = -X^{14} - X^{26}. \end{array}$$

In the notation of [Mag07], $\mathfrak{l}_{6,11} \otimes \mathbf{C}$ is $\mathfrak{g}_{6,12}$ while $\mathfrak{l}_{6,12} \otimes \mathbf{C}$ is $\mathfrak{g}_{6,11}$. We compute that

$$\begin{aligned} H^2(\mathfrak{l}_{6,7}, \mathbf{R}) &= \langle [X^{15}], [X^{16}], [X^{23}], [X^{25} - X^{34}], [X^{26}] \rangle \\ H^2(\mathfrak{l}_{6,11}, \mathbf{R}) &= \langle [X^{13}], [X^{15}], [X^{23}], [X^{16} + X^{25} - X^{34}], [X^{26} - X^{45}] \rangle \\ H^2(\mathfrak{l}_{6,12}, \mathbf{R}) &= \langle [X^{13}], [X^{15}], [X^{16} - X^{34}], [X^{26} - X^{45}], [X^{24}] \rangle \end{aligned}$$

The computations for $\mathfrak{l}_{6,11}$ and $\mathfrak{l}_{6,12}$ can be checked with the help of the derivative d_i written down with computer assistance in [Mag07] on p.44 and p.72 respectively ([Mag07] uses the $\mathfrak{g}_{6,i}$ notation recalled above for the Lie algebras and writes $\omega^{i,j}$ for X^{ij}). Moreover,

$$\begin{aligned} B^4(\mathfrak{l}_{6,7}, \mathbf{R}) &= \langle X^{1234}, X^{1235}, X^{1236}, X^{1245}, X^{1246}, X^{1256}, X^{1356} \rangle \\ B^4(\mathfrak{l}_{6,11}, \mathbf{R}) &= \langle X^{1234}, X^{1235}, X^{1245}, X^{1246}, X^{1236} - X^{1345}, X^{1346}, X^{1256} + X^{2345} \rangle \\ B^4(\mathfrak{l}_{6,12}, \mathbf{R}) &= \langle X^{1234}, X^{1235}, X^{1245}, X^{1246}, X^{1236} - X^{1345}, X^{1346}, X^{1256} + X^{2345} \rangle \end{aligned}$$

If π_i denotes the cup product $H^2(\mathfrak{l}_{6,i}, \mathbf{R}) \times H^2(\mathfrak{l}_{6,i}, \mathbf{R}) \rightarrow H^4(\mathfrak{l}_{6,i}, \mathbf{R})$, then

$$\begin{aligned}\mathrm{Im}(\pi_7) &= \langle [X^{1345}], [X^{2346}] \rangle \\ \mathrm{Im}(\pi_{11}) &= \langle [X^{1236}], [X^{2345}], [X^{1456} + X^{2346}] \rangle \\ \mathrm{Im}(\pi_{12}) &= \langle [X^{1236}], [X^{1456} + 2X^{2346}], [X^{1256}] \rangle\end{aligned}$$

One checks using the co-boundaries spaces B^4 listed above that these vectors are linearly independent, completing the proof that the cohomology rings of $\mathfrak{l}_{6,11}$ and $\mathfrak{l}_{6,12}$ are not isomorphic to that of $\mathfrak{l}_{6,7}$.

REFERENCES

- [AC01] José M. Ancochea and Rutwig Campoamor. Characteristically nilpotent Lie algebras: a survey. *Extracta Math.*, 16(2):153–210, 2001.
- [Ale75] Dmitry Vladimirovich Alekseevskii. Homogeneous Riemannian spaces of negative curvature. *Mat. Sb. (N.S.)*, 96(138):93–117, 168, 1975.
- [BK05] Mario Bonk and Bruce Kleiner. Conformal dimension and Gromov hyperbolic groups with 2-sphere boundary. *Geom. Topol.*, 9:219–246, 2005.
- [Bor95] Armand Borel. Sous-groupes compacts maximaux des groupes de Lie. In *Séminaire Bourbaki, Vol. 1*, pages Exp. No. 33, 271–279. Soc. Math. France, Paris, 1995.
- [Bou97] Marc Bourdon. Immeubles hyperboliques, dimension conforme et rigidité de Mostow. *Geom. Funct. Anal.*, 7(2):245–268, 1997.
- [Bre14] Emmanuel Breuillard. Geometry of locally compact groups of polynomial growth and shape of large balls. *Groups Geom. Dyn.*, 8(3):669–732, 2014.
- [BS99] Dietrich Burde and Christine Steinhoff. Classification of orbit closures of 4-dimensional complex Lie algebras. *J. Algebra*, 214(2):729–739, 1999.
- [CCMT15] Pierre-Emmanuel Caprace, Yves Cornulier, Nicolas Monod, and Romain Tessera. Amenable hyperbolic groups. *J. Eur. Math. Soc. (JEMS)*, 17(11):2903–2947, 2015.
- [CdH16] Yves Cornulier and Pierre de la Harpe. *Metric geometry of locally compact groups*, volume 25 of *EMS Tracts in Mathematics*. European Mathematical Society (EMS), Zürich, 2016. Winner of the 2016 EMS Monograph Award.
- [CDSV11] Matija Cencelj, Jerzy Dydak, Jaka Smrekar, and Aleš Vavpetič. Sub-linear Higson corona and Lipschitz extensions. *Houston J. Math.*, 37(4):1307–1322, 2011.
- [CE48] Claude Chevalley and Samuel Eilenberg. Cohomology theory of Lie groups and Lie algebras. *Trans. Amer. Math. Soc.*, 63:85–124, 1948.
- [CF17] Michael Cantrell and Alex Furman. Asymptotic shapes for ergodic families of metrics on nilpotent groups. *Groups Geom. Dyn.*, 11(4):1307–1345, 2017.
- [CJ94] Andrew Casson and Douglas Jungreis. Convergence groups and Seifert fibered 3-manifolds. *Invent. Math.*, 118(3):441–456, 1994.

- [CKD⁺21] Michael G. Cowling, Ville Kivioja, Enrico Le Donne, Sebastiano Nicolussi Golo, and Alessandro Ottazzi. From homogeneous metric spaces to lie groups, 2021.
- [Cor08] Yves Cornulier. Dimension of asymptotic cones of Lie groups. *J. Topol.*, 1(2):342–361, 2008.
- [Cor11] Yves Cornulier. Asymptotic cones of Lie groups and cone equivalences. *Illinois J. Math.*, 55(1):237–259 (2012), 2011.
- [Cor15] Yves Cornulier. Commability and focal locally compact groups. *Indiana Univ. Math. J.*, 64(1):115–150, 2015.
- [Cor16] Yves Cornulier. Gradings on Lie algebras, systolic growth, and cohopfian properties of nilpotent groups. *Bull. Soc. Math. France*, 144(4):693–744, 2016.
- [Cor18] Yves Cornulier. On the quasi-isometric classification of locally compact groups. In *New directions in locally compact groups*, volume 447 of *London Math. Soc. Lecture Note Ser.*, pages 275–342. Cambridge Univ. Press, Cambridge, 2018.
- [Cor19] Yves Cornulier. On sublinear bilipschitz equivalence of groups. *Ann. Sci. Éc. Norm. Supér. (4)*, 52(5):1201–1242, 2019.
- [CPS17] Matias Carrasco Piaggio and Emiliano Sequeira. On quasi-isometry invariants associated to a Heintze group. *Geom. Dedicata*, 189:1–16, 2017.
- [CT11] Yves Cornulier and Romain Tessera. Contracting automorphisms and L^p -cohomology in degree one. *Ark. Mat.*, 49(2):295–324, 2011.
- [dG07] Willem A. de Graaf. Classification of 6-dimensional nilpotent Lie algebras over fields of characteristic not 2. *J. Algebra*, 309(2):640–653, 2007.
- [DH08] Jerzy Dydak and Jose Higes. Asymptotic cones and Assouad-Nagata dimension. *Proc. Amer. Math. Soc.*, 136(6):2225–2233, 2008.
- [DK18] Cornelia Druţu and Michael Kapovich. *Geometric group theory*, volume 63 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2018. With an appendix by Bogdan Nica.
- [DS07] Alexander. N. Dranishnikov and Justin Smith. On asymptotic Assouad-Nagata dimension. *Topology Appl.*, 154(4):934–952, 2007.
- [EH96] Patrick Eberlein and Jens Heber. Quarter pinched homogeneous spaces of negative curvature. *Internat. J. Math.*, 7(4):441–500, 1996.
- [FN15] David Fisher and Thang Nguyen. Quasi-isometric embeddings of non-uniform lattices. *arXiv preprint arXiv:1512.07285*, 2015.
- [Fuk12] Tomohiro Fukaya. Sublinear Higson corona of Euclidean cone. *Tsukuba J. Math.*, 36(1):67–77, 2012.
- [FW18] David Fisher and Kevin Whyte. Quasi-isometric embeddings of symmetric spaces. *Geom. Topol.*, 22(5):3049–3082, 2018.
- [Gab92] David Gabai. Convergence groups are Fuchsian groups. *Ann. of Math. (2)*, 136(3):447–510, 1992.
- [Gro61] Alexander Grothendieck. Éléments de géométrie algébrique. III. Étude cohomologique des faisceaux cohérents. I. *Inst. Hautes Études Sci.*

- Publ. Math.*, (11):167, 1961.
- [Gro83] M. Gromov. Asymptotic geometry of homogeneous spaces. *Rend. Sem. Mat. Univ. Politec. Torino*, (Special Issue):59–60 (1984), 1983. Conference on differential geometry on homogeneous spaces (Turin, 1983).
 - [Gro84] Mikhael L. Gromov. Infinite groups as geometric objects. In *Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Warsaw, 1983)*, pages 385–392. PWN, Warsaw, 1984.
 - [Gro93] Mikhael L. Gromov. Asymptotic invariants of infinite groups. In *Geometric group theory, Vol. 2 (Sussex, 1991)*, volume 182 of *London Math. Soc. Lecture Note Ser.*, pages 1–295. Cambridge Univ. Press, Cambridge, 1993.
 - [GT87] Mikhael L. Gromov and William P. Thurston. Pinching constants for hyperbolic manifolds. *Invent. Math.*, 89(1):1–12, 1987.
 - [GW88] Carolyn S. Gordon and Edward N. Wilson. Isometry groups of Riemannian solvmanifolds. *Trans. Amer. Math. Soc.*, 307(1):245–269, 1988.
 - [Hea21] Brendan Burns Healy. Pinched curvature in heintze groups of carnot-type, 2021.
 - [Hei74] Ernst Heintze. On homogeneous manifolds of negative curvature. *Math. Ann.*, 211:23–34, 1974.
 - [Hel01] Sigurdur Helgason. *Differential geometry, Lie groups, and symmetric spaces*, volume 34 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2001. Corrected reprint of the 1978 original.
 - [HMT20a] David Hume, John M. Mackay, and Romain Tessera. Poincaré profiles of groups and spaces. *Rev. Mat. Iberoam.*, 36(6):1835–1886, 2020.
 - [HMT20b] David Hume, John M. Mackay, and Romain Tessera. Poincaré profiles of Lie groups and a coarse geometric dichotomy, 2020.
 - [HP13] Jose Higes and Irine Peng. Assouad-Nagata dimension of connected Lie groups. *Math. Z.*, 273(1-2):283–302, 2013.
 - [Jac55] Nathan Jacobson. A note on automorphisms and derivations of Lie algebras. *Proc. Amer. Math. Soc.*, 6:281–283, 1955.
 - [KDG21] Ville Kivioja, Enrico Le Donne, and Sebastiano Nicolussi Golo. Metric equivalences of Heintze groups and applications to classifications in low dimension, 2021.
 - [Kee94] James Keesling. The one-dimensional Čech cohomology of the Higson compactification and its corona. *Topology Proc.*, 19:129–148, 1994.
 - [KL97] Bruce Kleiner and Bernhard Leeb. Rigidity of quasi-isometries for symmetric spaces and Euclidean buildings. *Inst. Hautes Études Sci. Publ. Math.*, (86):115–197, 1997.
 - [Kob62] Shoshichi Kobayashi. Homogeneous Riemannian manifolds of negative curvature. *Tôhoku Math. J. (2)*, 14:413–415, 1962.
 - [KSTT05] Linus Kramer, Saharon Shelah, Katrin Tent, and Simon Thomas. Asymptotic cones of finitely presented groups. *Adv. Math.*, 193(1):142–173, 2005.

- [KW14] Linus Kramer and Richard M. Weiss. Coarse equivalences of Euclidean buildings. *Adv. Math.*, 253:1–49, 2014. With an appendix by Jeroen Schillewaert and Koen Struyve.
- [Lan66] Serge Lang. *Introduction to transcendental numbers*. Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1966.
- [Lau03] Jorge Lauret. Degenerations of Lie algebras and geometry of Lie groups. *Differential Geom. Appl.*, 18(2):177–194, 2003.
- [Laz69] Daniel Lazard. Autour de la platitude. *Bull. Soc. Math. France*, 97:81–128, 1969.
- [Mag07] Louis Magnin. *Adjoint and Trivial Cohomology Tables for Indecomposable Nilpotent Lie Algebras of Dimension less than 7 Over $\mathbf{C}$. Second corrected edition*. Université de Bourgogne, Laboratoire de physique-mathématique, 2007.
- [Mag08] Louis Magnin. Adjoint and trivial cohomologies of nilpotent complex Lie algebras of dimension ≤ 7 . *Int. J. Math. Math. Sci.*, pages Art. ID 805305, 12, 2008.
- [Mar70] Grigory A. Margulis. The isometry of closed manifolds of constant negative curvature with the same fundamental group. *Dokl. Akad. Nauk SSSR*, 192:736–737, 1970.
- [MT10] John M. Mackay and Jeremy T. Tyson. *Conformal Dimension*, volume 54 of *University Lecture Series*. American Mathematical Society, Providence, RI, 2010. Theory and application.
- [NR67] Albert Nijenhuis and Roger W. Richardson, Jr. Deformations of Lie algebra structures. *J. Math. Mech.*, 17:89–105, 1967.
- [OOS09] Alexander Yu. Olshanskii, Denis V. Osin, and Mark V. Sapir. Lacunary hyperbolic groups. *Geom. Topol.*, 13(4):2051–2140, 2009. With an appendix by Michael Kapovich and Bruce Kleiner.
- [Pal20] Gabriel Pallier. Sublinear quasiconformality and the large-scale geometry of Heintze groups. *Conform. Geom. Dyn.*, 24:131–163, 2020.
- [Pan83] Pierre Pansu. Croissance des boules et des géodésiques fermées dans les nilvariétés. *Ergodic Theory Dynam. Systems*, 3(3):415–445, 1983.
- [Pan89a] Pierre Pansu. Dimension conforme et sphère à l’infini des variétés à courbure négative. *Ann. Acad. Sci. Fenn. Ser. A I Math.*, 14(2):177–212, 1989.
- [Pan89b] Pierre Pansu. Métriques de Carnot-Carathéodory et quasiisométries des espaces symétriques de rang un. *Ann. of Math. (2)*, 129(1):1–60, 1989.
- [Per97] Grigory Perelman. A complete Riemannian manifold of positive Ricci curvature with Euclidean volume growth and nonunique asymptotic cone. In *Comparison geometry (Berkeley, CA, 1993–94)*, volume 30 of *Math. Sci. Res. Inst. Publ.*, pages 165–166. Cambridge Univ. Press, Cambridge, 1997.
- [Roe03] John Roe. *Lectures on coarse geometry*, volume 31 of *University Lecture Series*. American Mathematical Society, Providence, RI, 2003.
- [Sau06] R. Sauer. Homological invariants and quasi-isometry. *Geom. Funct. Anal.*, 16(2):476–515, 2006.

- [SM20] Emiliano Sequera Manzano. Relative L^p and Orlicz cohomology and applications to Heintze groups, 2020. Ph.D.Thesis – Universidad de la República and Université de Lille.
- [Tuk86] Pekka Tukia. On quasiconformal groups. *J. Analyse Math.*, 46:318–346, 1986.
- [TV00] Simon Thomas and Boban Velickovic. Asymptotic cones of finitely generated groups. *Bull. London Math. Soc.*, 32(2):203–208, 2000.
- [Tys99] Jeremy T. Tyson. *Geometric and analytic applications of a generalized definition of the conformal modulus*. ProQuest LLC, Ann Arbor, MI, 1999. Thesis (Ph.D.)–University of Michigan.
- [Wri03] Nick Wright. C_0 coarse geometry and scalar curvature. *J. Funct. Anal.*, 197(2):469–488, 2003.
- [Xie06] Xiangdong Xie. Quasi-isometric rigidity of Fuchsian buildings. *Topology*, 45(1):101–169, 2006.
- [Xie14] Xiangdong Xie. Large scale geometry of negatively curved $\mathbb{R}^n \rtimes \mathbb{R}$. *Geom. Topol.*, 18(2):831–872, 2014.

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ DE FRIBOURG, CHEMIN DU MUSÉE, 23
 – CH-1700 FRIBOURG, SWITZERLAND
Email address: gabriel@pallier.org