



**HAL**  
open science

## Models of plant resistance deployment

Loup Rimbaud, Frédéric Fabre, Julien Papaïx, Benoît Moury, Christian Lannou, Luke Barrett, Peter Thrall

► **To cite this version:**

Loup Rimbaud, Frédéric Fabre, Julien Papaïx, Benoît Moury, Christian Lannou, et al.. Models of plant resistance deployment. *Annual Review of Phytopathology*, 2021, 59, pp.125-152. 10.1146/annurev-phyto-020620-122134 . hal-03220213

**HAL Id: hal-03220213**

**<https://hal.science/hal-03220213>**

Submitted on 15 Feb 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Models of plant resistance deployment

Rimbaud Loup<sup>1,2,\*</sup>, Fabre Frédéric<sup>3</sup>, Papaïx Julien<sup>4</sup>, Moury Benoît<sup>1</sup>, Lannou Christian<sup>5</sup>, Barrett Luke G.<sup>2</sup>, Thrall Peter H.<sup>2</sup>

## **Affiliations:**

<sup>1</sup> INRAE, Pathologie Végétale, 84140 Montfavet, France

<sup>2</sup> CSIRO Agriculture and Food, Canberra, ACT 2601, Australia

<sup>3</sup> INRAE, SAVE, Bordeaux Sciences Agro, 33882 Villenave d'Ornon, France

<sup>4</sup> INRAE, BioSP, 84914 Avignon, France

<sup>5</sup> INRAE, BIOGER, 78850 Thiverval Grignon, France

## **Contact information:**

[loup.rimbaud@inrae.fr](mailto:loup.rimbaud@inrae.fr)

[frederic.fabre@inrae.fr](mailto:frederic.fabre@inrae.fr)

[julien.papaix@inrae.fr](mailto:julien.papaix@inrae.fr)

[benoit.moury@inrae.fr](mailto:benoit.moury@inrae.fr)

[christian.lannou@inrae.fr](mailto:christian.lannou@inrae.fr)

[luke.barrett@csiro.au](mailto:luke.barrett@csiro.au)

[peter.thrall@csiro.au](mailto:peter.thrall@csiro.au)

\* corresponding author:

+33 4 32 72 28 75

ORCID: [0000-0002-8098-9984](https://orcid.org/0000-0002-8098-9984)

**Shortened running title:** Models of plant resistance deployment

## **ABSTRACT**

Owing to their evolutionary potential, plant pathogens are able to rapidly adapt to genetically-controlled plant resistance, often resulting in resistance breakdown and major epidemics in agricultural crops. Various deployment strategies have been proposed to improve resistance management. Globally, these rely on careful selection of resistance sources and their combination at various spatio-temporal scales (e.g. via gene pyramiding, crop rotations and mixtures, landscape mosaics). However, testing and optimising these strategies using controlled experiments at large spatio-temporal scales is logistically challenging. Mathematical models provide an alternative investigative tool, and many have been developed to explore resistance deployment strategies under various contexts. This review analyses 67 modelling studies in light of specific model structures (e.g. demographic or demo-genetic, spatial or not), underlying assumptions (e.g. whether pre-adapted pathogens are present before resistance deployment) and evaluation criteria (e.g. resistance durability, disease control, cost-effectiveness). It highlights major research findings and discusses challenges for future modelling efforts.

Key words: adaptation, durability, evolution, host-microbe interaction, immunity, simulation.

## Introduction: Why we need to manage resistance deployment and how models can help

Deployment of plant resistance is a relatively low input, cost-effective way to protect agricultural crops from plant pathogens (67, 125). Plant resistance has been used in plant breeding programs to control diseases of various crops (e.g. (4, 6, 58)). However, pathogens have frequently evolved to quickly break down resistance following field deployment (37, 51, 106), sometimes resulting in catastrophic epidemics and massive use of pesticides. High pathogen evolutionary potential, coupled with the standardisation and intensification of modern agriculture across large cultivated areas has generally led to recurrent cycles of resistance deployment followed by rapid pathogen adaptation, often described as boom and bust cycles (78). When the resistance of a cultivar becomes ineffective, economic losses can be considerable because of the direct impact of epidemics and the cost of alternative control methods. Moreover, breeding for resistance is costly, time consuming and often constrained by the limited availability of genetic resistance sources (40, 147). Resistance genes should therefore be considered as an exhaustible resource deserving careful stewardship.

The design and implementation of strategies that improve **resistance durability** would therefore be of great benefit to agricultural productivity, sustainability and profitability. However, a key point is that **resistance durability** and **epidemiological control** are not necessarily correlated (13, 52). Thus, any strategy designed to control the emergence of **resistance-adapted pathogens** in agro-ecosystems has the potential to conflict with epidemic control both in the short-term (from standing pathogenic variants) and the long-term, should resistance break down.

Pathogen spread and adaptation is favoured by the low host genetic diversity that is representative of intensive agricultural systems (124). Therefore, many proposed strategies rely on the selection of diverse resistance sources and their spatio-temporal deployment to engineer biodiverse cropping systems (9, 13, 77, 84, 125). The goal is to confront pathogens with eco-evolutionary challenges and thus avoid or delay their adaptation to plant resistance, while maintaining effective epidemiological protection (15, 147). Particularly for airborne plant pathogens, deployment strategies are more likely to be effective if implemented across landscapes at large spatial scales (40). However, experimental tests of landscape-based strategies are rarely feasible for obvious practical reasons (but see (26, 57, 150)).

To overcome the difficulties of experimentation with plant resistance deployment at large spatio-temporal scales, numerous mathematical models have been developed (see sidebar: “Complementarity between models, experiments and observations”). However, models are typically faced with the challenge of combining several aspects of resistance deployment (e.g. type of plant-pathogen interaction; spatio-temporal scale of deployment; section 1), incorporating realistic parameters and assumptions with respect to epidemiological and evolutionary processes (section 2), and generating usable outputs (section 3). In this review, we identified 67 studies that used models to assess or optimise

deployment strategies (Table 1 & details in Supplemental Table S1). Our aim is to provide a comprehensive overview of these modelling approaches, noting that the diversity of model structures, assumptions and outputs make direct comparisons difficult. Regardless, we examine their main features from both epidemiological and evolutionary perspectives and highlight major findings of relevance to **resistance durability** and **epidemiological control**.

## 1. Modelling resistance deployment from gene to landscape scales

The main strategies considered in plant resistance deployment act across scales from genes to landscapes. They rely on the appropriate choice of resistance sources, which can be combined in the same cultivar (gene pyramiding), alternated within rotations, mixed within fields, or segregated across a mosaic of fields (Figure 1). Since these have been extensively described in previous reviews (9, 13, 84, 125, 147), we focus on mathematical models developed to assess, compare and improve these strategies (Table 1 & Supplemental Table S1). We note that the evolutionary processes underlying pathogen and pest adaptation to plant resistance are analogous to those associated with the emergence of resistance to chemicals. Therefore, the deployment strategies we consider have counterparts for the management of resistance to pesticides in crops, drugs used for the treatment of animals and humans, as well as vaccines (for reviews see (109, 110, 128, 136)).

### A. [Choosing appropriate resistance sources](#)

The smallest scale that impacts deployment lies in the choice of resistance sources to be deployed. Plant resistance has often been classified as either **qualitative complete** or **quantitative partial**, although considerable empirical evidence suggests that this dichotomy should be revised (see sidebar: “Should the traditional dichotomy of plant resistance be reconsidered?”).

#### Modelling qualitative and quantitative resistance

**Qualitative resistance** usually refers to ‘major resistance genes’, which code for specific host proteins able to recognise a specific pathogen molecular pattern or effector. Such ‘gene-for-gene’ interactions are traditionally modelled using a 2-by-2 matrix describing the occurrence of disease as an outcome of the interaction between host **genotype** (with or without the resistance gene) and pathogen **genotype (non-adapted or adapted)** (34) (Figure 2d). Only **resistance-adapted pathogens** can infect resistant hosts. Depending on whether or not they are present in the initial pathogen population, they may have to appear (e.g. via immigration from distant areas, mutation or recombination, section 2B). In this context, pathogen adaptation leads to **resistance breakdown**, i.e. complete restoration of pathogen infectivity on resistant hosts. However, adaptation is often penalised by a **fitness cost** on susceptible hosts (59, 67, 139), resulting in decreased ability of **resistance-adapted pathogens** to infect susceptible hosts compared to **non-adapted genotypes** (130). Inclusion of a **fitness cost** parameter makes the plant-pathogen interaction matrix (Figure 2b) relevant

for both the ‘gene-for-gene’ (Figure 2d) and ‘matching allele’ (Figure 2e) concepts (1, 116, 129). Indeed, the classic matching allele concept states that pathogen adaptation to a new host makes infection of the other host impossible (i.e. the **fitness** cost is maximal).

**Quantitative resistance** traditionally refers to the additive effects of multiple ‘minor resistance genes’, resulting in a continuous distribution of pathogen adaptation (20, 90, 123). From the plant perspective, pathogen adaptation corresponds to an erosion phenomenon (9, 78, 105) (Figure 2ac). **Quantitative resistance** is classically considered to be **partially efficient**, i.e. resistant hosts can be infected by **maladapted pathogens**, although disease development is reduced. Models with compartmental architecture (section 2A) can disentangle the effect of **partial resistance** on different **pathogenicity traits**: plant infection rate by pathogen propagules (e.g. fungal or bacterial spores, insect vectors carrying viral particles), latent and infectious periods, and propagule production rate (Figure 3a). According to the few modelling studies which have compared the performance of **quantitative resistance** against different **pathogenicity traits**, from a disease management perspective, the most promising target trait for **quantitative resistance** is latent period, followed by infection rate and propagule production rate (72, 113).

However, **qualitative resistance** is not always **complete**, and **quantitative resistance** is not necessarily **partial** (see the sidebar titled “Should the traditional dichotomy of plant resistance be reconsidered?”). For example, many **qualitative resistance** genes actually allow some infection, whether they may be partially broken down, environmentally sensitive, developmentally-regulated, or simply weak (19, 91, 123). In models, inclusion of a resistance efficiency parameter in the plant-pathogen interaction matrix (Figure 2b) allows the representation of **partially efficient qualitative resistance**.

#### Deployed resistance sources must be inherently hard to break down

One of the few models accounting for the molecular mechanisms of pathogen adaptation to resistance highlighted the effect of the number, nature (transition or transversion) and rate of the required mutations and associated **fitness** costs on the **durability** of major resistance genes (31). The impact of mutation rate and **fitness** cost has also been demonstrated in models focusing on phenotypic aspects of **qualitative resistance** ((32, 33, 111); see also section 2B). Similarly, for **quantitative resistance**, models indicate that strong **fitness** costs of pathogen adaptation slow down resistance erosion (72, 101, 103). However, resistance erosion is sensitive to pathogen mutational processes (i.e. the number of mutations required to completely erode **quantitative resistance**) (8). Collectively, these results support the importance of identifying and deploying resistance sources that are inherently hard to break down. These findings have been confirmed experimentally for plant viruses (49, 78, 108).

High resistance efficiency (i.e. the targeted **pathogenicity trait** is drastically reduced for **maladapted pathogens**) reduces pathogen epidemiological impact (104, 113, 118). However, the effect of this parameter on **resistance durability** is less well documented, probably

because it is generally associated with **quantitative resistance**, for which defining **durability** is still a challenge (section 3B). A study on pesticide resistance showed that high application doses slow the appearance of **adapted pathogens**, but hasten their invasion once present (45). These conclusions agree with results obtained for **partially efficient** major genes when **resistance-adapted pathogens** are initially present (21, 107).

## B. [Spatio-temporal strategies for deploying different resistance sources](#)

### Resistance genes can be combined at plant scale

Multiple resistance sources can be stacked into a single plant **genotype**, as a “pyramid” (29, 35, 82). Although different resistance types can be combined, published models have mostly focused on pyramids of major resistance genes (Figure 3b). In line with theoretical predictions (9, 67, 84) and results obtained with single genes (section 1A), models have shown that the **durability** of this strategy increases with the number of mutations required to break down all the genes composing the pyramid, and the strength of associated **fitness** costs (28, 31, 111, 113). However, **durability** of pyramids can be compromised if pyramid gene components are simultaneously deployed individually (73, 117) or if pathogens **adapted** to one or more of these components are already present in the population (73) (see also section 2B). Pyramid cultivars must therefore be deployed carefully, especially because their **breakdown** results in the emergence of **multi-adapted pathogens**.

Extensive empirical evidence has demonstrated the efficacy of combining **qualitative** and **quantitative resistance** in a pyramid (11, 85, 98, 108) but few models have investigated these scenarios. These models have shown that the **breakdown** of a major resistance gene can be delayed when pyramided with the appropriate **quantitative resistance**. Promising target traits include latent period (113) and pathogen effective population size (i.e. the size of an idealized population showing the same degree of randomness in allele frequencies as the real population, noting that small effective sizes amplify genetic drift; section 2B) (114). Several sources of **quantitative resistance** may also be pyramided (60, 123). The only model investigating this scenario highlighted the potential of targeting **pathogenicity traits** whose evolution is constrained by trade-offs (i.e. evolution to improve one trait penalises another trait) (8).

### Resistance genes can be segregated at field and landscape scales

At field scale, varieties carrying different resistance sources (or resistant and susceptible varieties) can be cultivated in the same patch, simultaneously in mixtures or alternating within rotations. Crop varieties can also be segregated into a mosaic of fields within a broader regional strategy (78, 84). Furthermore, varying the different components of mixtures, rotations and mosaics, their relative proportions and their spatial (for mixtures and mosaics) or temporal (for rotations and mosaics) organisation offer a multitude of deployment options (Figure 4).

The value of combining cultivars in mixtures or mosaics is likely to increase with the number and heterogeneity of components. Given **fitness** costs of pathogen adaptation to different hosts, heterogeneous host populations are expected to favour diversifying selection, i.e. to select for higher pathogen specialization (5). This should result in reduced disease spread owing to (i) a dilution effect (i.e. reduced colonization rate of a specialized pathogen in host cultivars to which it is adapted due to increased distance among these hosts, with the remainder of the host population acting as a propagule sink), (ii) a barrier effect (i.e. hosts of a given **genotype** acting as physical barriers to pathogen spread owing to their architecture); and (iii) a competition effect (i.e. different pathogen **genotypes** competing for the same host individual) (14, 54, 143). In accordance with empirical results (38, 48), models confirm the potential for dilution effects by showing that the amount of disease in mixtures is often smaller than the arithmetic mean of the amount of disease obtained from these components in pure stands (50, 53, 65, 66, 118). Assuming **fitness** costs of adaptation to different components, models have also demonstrated that higher numbers of mixture components amplify dilution effects (28, 80) and delay the emergence of **multi-adapted pathogens** through competition with more specialised pathogens (44, 61, 65). In addition to dilution, barrier and competition effects, mixtures and to a lesser extent mosaics may facilitate plant immune system activation via various mechanisms including induced resistance (149). Induced resistance (for any kind of pathogen) or cross protection (for viruses) can occur when a cultivar becomes resistant to a **resistance-adapted pathogen** when previously challenged by a **non-adapted pathogen** coming from another cultivar (148). The only model we are aware of that has investigated induced resistance in mixtures highlighted the impact of the size of the area protected by induced resistance, and of the duration and level of protection on disease reduction (64).

In practice, the spatial organisation of a mixture is often linked to the degree of heterogeneity of its components. Multiline cultivars differing in only a few genes (10), non-fixed populations showing high genetic diversity (e.g. landraces), and mixtures of different cultivars from the same species (143) may be sown after mixing seeds, resulting in completely random mixtures. Conversely, intercropping different species (7), as well as some cultivar mixtures, often require planting in blocks or rows to facilitate crop segregation at harvest (83, 150). Accordingly, mixtures are qualitatively similar to mosaics (hence the term “field mosaic” sometimes used), the main difference being the average distance between components (the size of a genetic unit), noting that this distance must match the scale of pathogen dispersal for effective disease management (41, 119). In fact, results obtained from studies of mixtures and mosaics are likely to be highly correlated (86). This is especially true for non-spatial models where the scale of host population structure could be interchangeably represented as a field or a landscape (Table 1).

Models have found that when **resistance-adapted pathogens** are initially absent, high proportions of resistant hosts in a field or broader region with low aggregation of cultivars (or weak connectivity between hosts of a given cultivar) favour good disease control (39, 47, 72,



96, 99, 103, 104, 111, 118, 122, 127, 138, 146). However, intermediate proportions of resistant cultivars are often preferable when **adapted pathogens** are initially present, with the precise proportion that maximises disease control being sensitive to their initial frequency (32, 94). This U-shape effect is in line with the effect of the proportion of resistance on its **durability** (17, 103, 111, 135). The U shape is attributable to the fact that high proportions of resistance considerably reduce pathogen population size (resulting in a low probability of appearance of **adapted** mutants), while small proportions minimise selection pressure and reduce the probability that an **adapted pathogen** successfully disperses to a resistant field. With respect to spatial aggregation, in contrast to its effect on disease control, well-mixed landscapes (i.e. low aggregation) tend to have a higher probability of **resistance breakdown**, as this increases the interface between resistant and susceptible components, and thus the exposure of resistant hosts to potential **adapted pathogens** emerging from susceptible cultivars (17, 103, 111).

The effect of cultivar rotations on **resistance durability** is less well documented than the effect of mixtures and mosaics, especially in models. Nevertheless, one expectation is that for each rotation cycle, the number of components, their degree of heterogeneity and temporal sequence all affect pathogen survival (equivalent to dispersal in time; (12)) in a conceptually analogous way to pathogen spatial dispersal in mixtures and mosaics. Accordingly, models predict that resistance will be more durable when used in small proportion (92) and with a rapid turnover (111) within the temporal sequence. Furthermore, while most landscape-scale studies use a static landscape (i.e. crop allocation to fields is fixed), some models have demonstrated that resistance deployment can be improved by temporal variation in the proportion of resistance in the landscape (28, 33, 70). This finding emphasises the potential of dynamic mosaics where the number and spatial location of resistant fields vary in time (Figure 4).

In addition to the strategies discussed above, resistance deployment can benefit from complementary control methods (83). Any control method which reduces the effective or census size of the pathogen population (e.g. chemical applications, biological control, removal of infected crop residues and other agronomic practices) should increase **resistance durability**, provided that the pathogen does not adapt to this additional control method (16).

The diversity of resistance types, spatio-temporal scales of deployment and complementary control methods result in an extensive range of possibilities whose combinations merit further investigation. Based on the few results from modelling studies, some combinations are expected to maximize both **epidemiological control** and **resistance durability**, like rotations and mosaics (28, 33, 111) or rotations and mixtures (145). Other combinations are yet to be explored, like those involving different types of resistance (e.g. **complete** and **partial, qualitative** and **quantitative**) at different spatio-temporal scales via pyramids (Figure 3), mosaics, mixtures and rotations (Figure 4). Models are powerful tools to

assess and compare deployment options. Nevertheless, a clear understanding of their structure and assumptions is essential to make appropriate interpretations and comparisons.

## 2. Model structure and assumptions

Following the long-standing tradition of compartmental models in theoretical epidemiology (55), diverse models have been developed to investigate resistance deployment. They are mainly distinguished by how the genetic and spatio-temporal structures of host and pathogen populations are modelled and how epidemiology and evolution are accounted for.

### A. [Modelling epidemics](#)

The basic 'SIR model' consists of a set of coupled non-linear ordinary differential equations (ODE) representing the temporal dynamics of  $S(t)$ ,  $I(t)$  and  $R(t)$ , respectively the numbers of 'susceptible' (i.e. healthy in this context), 'infected' and 'removed' (i.e. epidemiologically inactive) individuals in the host population (75). For pathogens such as viruses that trigger systemic infection, host individuals are often entire plants. For other pathogens such as fungi that trigger localised clonal lesions, individuals may be considered as plant tissue units such as a leaf or part thereof (Supplemental Table S1). If the basic SIR model considers a population of genetically identical hosts, compartmental models can handle several host **genotypes** by including dedicated compartments for each plant cultivar. Such models remain purely demographic from the pathogen perspective, as a single pathogen strain characterised by a unique set of **pathogenicity traits** (e.g. transmission rate, infection rate; see section 1A) is considered (Table 1a). ODE naturally apply to tropical crops with continuous planting and harvest throughout the year. Nevertheless, seasonality of temperate climates (as well as cultivar rotations, section 1B) can be represented in semi-discrete models. They undergo continuous ODE dynamics, and experience discrete dynamics at given time points, typically at pathogen overwintering ((76); see also Supplemental Table S1). Regardless of their structures, ODE rely on a **deterministic** framework where a set of given inputs invariably yields the same outputs. However, epidemic dynamics as well as pathogen demography are impacted by uncertainty affecting individual life events (**demographic stochasticity**) or imposed by the environment (**environmental stochasticity**) (67). **Demographic stochasticity** is typically important during the initial phase of an epidemic. Using **stochastic** SIR models is then a way to account for the probability of epidemic extinction when the number of infected hosts is small.

### B. [Accounting for pathogen evolution](#)

Addressing questions about **resistance durability** requires modelling of pathogen evolution. Historically, theoretical studies of pathogen evolution mainly relied on the framework of adaptive dynamics (25). Assuming that a single pathogen strain is present at its endemic steady state, the method determines under what conditions a second mutant strain, introduced at low frequency, will invade the population. Adaptive dynamics supposes that

epidemiological and evolutionary processes do not interfere with each other or act at different time-scales. It essentially focuses on long-term predictions for endemic diseases and has been used to study pathogen evolution in response to resistance deployment (Table 1b). For example, the effect of resistance genes targeting different **pathogenicity traits** on the evolution of virus multiplication rates has been investigated (134, 137). Through computation of stable evolutionary equilibria, it is possible to resolve evolutionary trade-offs like those between virulence (in this context, virulence refers to host damage caused by pathogen infection) and transmission (36, 131), survival and transmission (131), generalism and specialisation (102). However, in agro-ecosystems, interest is generally focused on short-term epidemics, which impacts both pathogen population size and composition. Day and Proulx (24) and Day and Gandon (23) introduced a framework to simultaneously model the dynamics of epidemics and evolution for any **pathogenicity trait**. This approach, termed evolutionary epidemiology, is inspired by quantitative genetics and is well suited to model the evolution of quantitative traits. To date, it has been applied to study the erosion of **quantitative resistance** targeting different **pathogenicity traits** (27, 72).

As for plant cultivars, multiple pathogen strains are considered in multi-strain compartmental models. By doing so, models explicitly switch from demographic to demographic representations of the pathogen population (Table 1c). Many models solely consider the effect of selection, and assume that all pathogen **genotypes** are initially present, with **adapted genotypes** initially present at low frequencies. These assumptions ignore the time required for the appearance and establishment of adaptive mutations in pathogen populations (135). While this may be an acceptable assumption for pathogens with large population sizes, it can considerably overestimate the speed of adaptation of pathogens with smaller effective population sizes, especially for strategies involving gene pyramids (section 1B).

A further step is to explicitly model the effects of mutation or migration on the establishment of **resistance-adapted pathogens** that are initially absent. Taking into account the classical genetic mutation rate (number of genetic mutations/generation/nucleotide) is difficult as it requires correct identification of the genetic architecture (number, type, location and phenotypic effects of genetic mutations) underlying phenotypic trait variation. Presently, this knowledge is only considered for virus adaptation to major resistance genes using the concept of mutation-selection balance (28, 32, 33). For most pathogens, the links between **genotype** and **phenotype** are ignored. Most models instead consider a phenotypic mutation rate representing a displacement into the pathogen phenotypic space (e.g. (8, 72, 73, 96, 113)). Empirically largely unknown, this rate theoretically integrates the phenotypic effects of genetic mutations into the multi-dimensional space of **pathogenicity traits** as well as potential correlations within and between host **genotypes**. Immigration of pathogen strains from external sources has also been considered as an alternative source of new genetic variants (71, 135). While the rate of appearance of new genetic variants via mutation depends

on the size of the pathogen population, with immigration this rate is generally constant and independent of the epidemiological status of the system.

Besides mutation, genetic recombination (occurring during virus replication, bacterial conjugation, transformation and transduction, or fungal sexual cycle), also generates new variants. For example, recombination can accelerate pathogen adaptation via the re-assortment of adaptive mutations controlled by independent loci. This enables the emergence of **multi-adapted pathogens** and thus favours the **breakdown** of resistance pyramids. Thus, ignoring its role can result in severe bias (2). Only few models (70, 117, 144) included recombination to study the emergence of **multi-adapted genotypes** of a pathogen having one sexual phase at the beginning of each cropping season (Table 1).

**Demographic stochasticity** shapes the effect of these evolutionary processes and can be dealt with **stochastic** models. This is of particular importance when (i) events are inherently rare (e.g. due to low mutation rates), and (ii) pathogen population size is low (e.g. just after a mutant strain appears). Therefore, depending on subtle interactions between effective population size ( $Ne$ ), selection coefficient ( $s$ ) and mutation rate ( $\mu$ ), the evolutionary dynamics of a population can be mainly **deterministic** or **stochastic** (115). For example, if  $Ne \times \mu \ll 1$ , the waiting time of mutations conferring adaptation could be substantial and is thus subjected to large random fluctuations that can hardly be ignored if **adapted pathogens** are initially absent. If  $Ne \times s \ll 1$ , genetic drift generates random fluctuations in pathogen genetic diversity, with the potential to purge variants regardless of their selective value. Most models reviewed here ignore the effect of genetic drift (Table 1). Dealing with such evolutionary force requires specific consideration of  $Ne$  and its variation, especially bottlenecks from which only some individuals will be randomly sampled and survive to reproduce (18). Lo Iacono, et al. (71) used the size of the infected host population as an indirect proxy of such variation, whereas other authors have explicitly modelled bottlenecks occurring at crop harvest and during the off-season (73, 103, 113, 144, 145), during intra-plant movement in viral infection (114) and during viral transmission by insect vectors (31).

### C. [Representing the spatial structure of host populations.](#)

In non-spatial models (Table 1), each infected individual is equally likely to establish an infectious contact with any other healthy individual, regardless of its geographic location. Accounting for space is critical for accurate assessment of deployment strategies relying on the spatial segregation of different cultivars (e.g. mixtures and rotations, section 1B), especially when the pathogen disperses primarily at short distances compared to field size.

New compartments leading to more realistic contact structures between individuals can be introduced in spatially-implicit models. A metapopulation of well-mixed host patches where pathogen spread rates differ within and between patches provides a typical example (42). Addressing this question more deeply requires spatially-explicit approaches, where pathogen dispersal is represented using reaction-diffusion equations (RDE; (46)) or via dedicated kernels (89) within integro-differential equations (IDE). Notably, dispersal kernels

allow explicit representation of long-distance infection events (79). In any case, variables of interest depend on a vector of spatial coordinates  $x$  in addition to time  $t$  (e.g.  $S(t,x)$ ,  $I(t,x)$ ), providing a straightforward way to investigate the effect of habitat geometry (e.g. spatial aggregation, section 1B). **Stochastic** models are often used to randomly segregate pathogen propagules across different landscape elements. They are also used to explicitly consider **environmental stochasticity** associated with random fluctuations of cultivar allocations (73, 145), or pesticide treatments (71, 96) year after year in different parts of the landscape.

### 3. Model outputs: criteria to assess deployment strategies

As described for fungicide application (136), assessment of resistance deployment strategies depends strongly on evaluation criteria. Importantly, the objectives of different stakeholders (e.g. growers, breeders) are not always compatible. Indeed, minimising epidemiological impacts, maximising **resistance durability**, and minimising costs of disease management are all sensible targets that may lead to different optimal strategies (103, 111, 135). This section considers potential key variables used in models (Table 1 & Supplemental Table S1) for assessing resistance deployment strategies from epidemiological, evolutionary and socio-economic perspectives.

#### A. [Epidemiological outputs](#)

Epidemiological outputs characterise the ability of deployment strategies to reduce disease impact in a given region over a given time period (15). In the field, disease impact is classically evaluated using the proportion of infected hosts (disease prevalence; (75)) or a quantitative assessment of symptoms on infected hosts (disease severity; (75)). In models, disease impact is usually measured by the proportion of infected individuals (i.e. infection units, Section 2A), equating to prevalence when an individual is a single plant, or severity when an individual is an infection site. Thus, prevalence and severity can only be distinguished when plant architecture is explicitly modelled in terms of number of infection sites per plant (e.g. (39, 144-146)). Because prevalence and severity change in time, the challenge is to summarize them using representative point estimates to compare different deployment options. One approach focuses on epidemiological status at the end of a given period (118, 122, 127) or at stable evolutionary equilibrium (80, 94, 134, 137). Another approach relies on the integration of prevalence or severity over time into an area under disease progress curve (AUDPC; (28, 32, 33, 111, 113)).

Alternatively, the number of infected hosts can be deduced from the number of healthy hosts (94, 100, 134, 137). Similar to the computation of AUDPC, the dynamic of healthy hosts may be integrated over time into a variable representing the cumulative amount of photosynthetic tissue, assumed to be proportional to crop growth and yield (141). Modelling studies variously refer to this as healthy area duration (HAD; (103)), healthy surface (72, 99, 104), green leaf area (113) or green area canopy duration (107). Depending on the study objectives, AUDPCs and HAD-like variables may be computed for the whole host population

or individually for each cultivar (e.g. (107)), and for the whole simulation run or for specific periods (e.g. before and after **resistance breakdown**; (103, 111)). It may be expressed in absolute value, or relative to a reference context, for example to assess, compared to a fully susceptible landscape, the additional number of healthy hosts resulting from resistance deployment (71, 72, 135) or the intensity of epidemics (28, 32, 33).

Finally, the dynamic of infected hosts in a region can be summarized by other metrics like the speed of the epidemic expansion front (121, 133, 138), the total distance travelled (104), or the final area covered by epidemics (122, 127).

## B. [Evolutionary outputs](#)

Evolutionary outputs mainly assess the ability of deployment strategies to prevent or slow down pathogen adaptation and are often summarized by **resistance durability**. In models where pathogen **genotypes** are classified as **non-adapted** or **adapted** (section 1A), **resistance durability** has been calculated using criteria such as the point in time when resistant hosts become as severely affected as susceptible hosts (107), when **adapted pathogens** first appear (31), or when their prevalence (56, 111, 113) or frequency in the pathogen population (16, 73, 74, 96, 107, 135) exceed an arbitrary threshold. The value of this threshold determines which of the three characteristic phases of **resistance breakdown** is targeted. These phases, analogous to an ecological invasion process of the cropping landscape by **adapted pathogens**, consist of: i) introduction via immigration or appearance (via mutation, recombination, horizontal gene transfer or sexual reproduction), ii) establishment, when extinction becomes unlikely despite potential competition with **non-adapted pathogens**, and iii) spread (or propagation) within the formerly resistant host population, potentially causing severe yield losses. Thus, model results can be significantly impacted by the chosen threshold above which resistance is considered to have been broken down (74), but also by model assumptions (e.g. whether or not **adapted pathogens** are initially present; (73); see also section 2B). When several resistance genes are deployed, and in particular for pyramid cultivars, resistance durability can also be assessed based on the dynamics of the **multi-adapted pathogen** (21, 44, 65, 66, 111).

Measuring pathogen adaptation to resistance when it is continuously distributed is still a challenge. The few approaches that have been proposed rely on the speed of pathogen evolution (102) or of resistance erosion (113), and on the time point when the proportion of healthy hosts drops below an arbitrary threshold (16, 101, 103).

## C. [Socio-economic outputs](#)

Despite the importance of identifying deployment strategies that are not only efficient and durable but also cost-effective and feasible for growers, socio-economic factors are rarely accounted for in resistance deployment models. Although planting density, growth rate and contribution to crop yield are often assumed to be uniform, this is rarely the case in practice. For potato late blight in the Netherlands (96) and blackleg of winter oilseed rape (69, 70), crop yield was computed as a function of crop cultivar, disease severity and climatic variables.

Accounting for yield enables the resolution of trade-offs between the reduction of damage resulting from the use of resistant cultivars and associated costs, like reduced growth (96) or smaller yield (140) of resistant varieties. However, yield data are generally not available and can be challenging to estimate (or predict), particularly the yield of a cultivar carrying a broken down resistance.

Organisational and social aspects (e.g. feasibility criteria accounting for farming practices) of resistance deployment have rarely been considered. Yet, these factors drive farmers' decisions and have considerable impact on the adoption of a deployment strategy, as illustrated for potato late blight (95). Milne, et al. (81) predict that adoption of a resistant maize cultivar producing a *Bacillus-thuringiensis* toxin (Bt, offering better protection against European corn borers but potentially more expensive to grow) by farmers depends on their communication network and sensitivity to risk. In this context, the use of visual interfaces and model-based scenarios can help communicate results to the farmer community and stimulate stakeholder discussions in workshops on plant resistance management (e.g. (97)).

## General conclusions

Models of resistance deployment can be used to deliver insights into the epidemiological, evolutionary and economic performance of deployment strategies (e.g. gene pyramiding, crop rotations, cultivar mixtures, landscape mosaics), at spatio-temporal scales beyond the scope of empirical experimentation. However, it is important to note that the models reviewed here are not designed to precisely predict **resistance durability** or the level of **epidemiological control** in real-world systems. Rather their purpose is to identify key parameters, provide mechanistic insight into the consequences of different deployment strategies and allow decision makers to understand their relative merits (43).

Given this focus, is it possible to use the models to collectively identify a single 'best' strategy for resistance deployment? The diversity of assumptions underlying the different modelling approaches, of scenarios considered and of evaluation criteria used makes it almost impossible to rank strategies. Based on the very few studies that allow such a comparison "all other things being equal", pyramiding seems the most epidemiologically-efficient and evolutionarily-durable strategy to deploy **complete resistance** in the absence of pre-adapted pathogen **genotypes** (73, 111) (Figure 5). However, pyramiding can suffer important limitations as (i) it is challenging to effectively identify the existence of pre-adapted **genotypes** that could be maintained at very low frequencies; (ii) other cultivars or wild relatives can represent evolutionary steps facilitating the adaptation of the pathogen population; and (iii) several precious genetic resources are lost at the same time when a pyramid is overcome. Mosaics and rotations are alternative solutions to cope with these limitations and are predicted to be more efficient (28, 111) and durable (73) when **resistance-adapted genotypes** are present prior to resistance deployment. In addition, by increasing the overall diversity of the crop, these strategies confer a portfolio effect, i.e. increased resilience to other biotic and abiotic constraints. These conclusions should not be overgeneralized to any crop production

situation and pathosystem as they have mostly been obtained in specific contexts. This brings us face-to-face with the modeler's dilemma – it is not possible to maximize generality, realism and precision in the same model (68).

To conclude, we argue that there is no universal strategy. The optimal deployment approach depends on the desired objective, the epidemiological and evolutionary context, and the pathosystem considered, among others. This is in line with what Mundt wrote in a recent review (85): "There are multiple approaches to increasing **durability of resistance** [...], none of which should be considered inherently superior to the other, and all of which likely benefit from being combined". We hope this review paves the way for future modelling investigations towards more efficient, sustainable, cost-effective and feasible deployment of plant resistance.

## Summary points

- During the last twenty years there has been a paradigm shift from the idea of "durable resistance" to "durable management of resistance". This implies that **durability** is no longer considered as an intrinsic property of a resistance gene but rather the result of a clever combination of effects operating at different scales. Accordingly, plant resistance against pathogens must be carefully designed (by breeders) and organised in space and time (by farmers) to be both efficient and durable in spite of pathogen evolutionary potential.
- The multiplicity of deployment strategies, resulting from the huge diversity of choices made from gene to landscape scales (Figure 1), impedes their comparison with empirical experimentation. In this context, models provide powerful tools to explore possible deployment options (Figures 2, 3, 4) and can help identify promising strategies (Figure 5) which, in turn, may be amenable to experimental verification.
- The profusion of modelling approaches offers a wide range of possibilities to model epidemics while accounting for pathogen evolution and spatiotemporal organisation of host plants (Table 1 and Supplemental Table S1). Their epidemiological, evolutionary and socio-economic outputs allow the evaluation of deployment options with respect to multiple criteria.
- There is no 'silver bullet' deployment strategy: optimal deployment approaches will vary depending on the desired objective, the epidemiological and evolutionary context, and the pathosystem considered. Any deployment option will have its own advantages and drawbacks, thus there is likely to be benefit from hybrid strategies.

## Future challenges

- Future models should encompass the diversity of plant immunity mechanisms (including tolerance, which reduces pathogen-induced host damage without affecting pathogen development) and consider the full range of pathogen adaptation mechanisms.



- Recognizing that predictions about real world systems often require relatively complex models, we must take advantage of computers current capacity to model complex resistance deployment scenarios that combine different deployment options and complementary control measures at multiple spatio-temporal scales. Such scenarios should also include adaptive strategies that are continuously updated based on real-time data.
- A unified modelling framework will help compare deployment strategies “all other things being equal”, and understand the impact of ecological, epidemiological, evolutionary and genetic factors for a diversity of pathosystems. This is one of the objectives of the model *landsepi*, freely accessible through an R package (112).
- More efforts should be made to collect and share epidemiological datasets to help calibrate and validate models with empirical data.
- Models must account for socio-economic and organisational constraints of real farming systems in order to identify solutions that are both feasible and likely to be adopted by breeders and growers.
- Results and recommendations need to be communicated to relevant stakeholders in a way that information is accessible and likely to be acted upon. The use of dedicated pedagogical interfaces facilitates such interactions ((97), see also an interface for *landsepi* in Related Resources).

## Acknowledgements

The authors benefited from GRDC grant CSP00192, ANR project ArchiV (2019-2022, grant n°ANR-18-CE32-0004-01), AFB Ecophyto II - Leviers territoriaux project Médée (2020-2022), and the INRA-CSIRO linkage program. Figure 1 was designed by GRAPHIC Business from LR's original drawing.

## References

1. Agrawal A, Lively CM. 2002. Infection genetics: gene-for-gene versus matching alleles models and all points in between. *Evolutionary Ecology Research* 4: 79-90
2. Arenas M, Araujo NM, Branco C, Castelhana N, Castro-Nallar E, Pérez-Losada M. 2018. Mutation and recombination in pathogen evolution: Relevance, methods and controversies. *Infection, Genetics and Evolution* 63: 295-306
3. Bargués-Ribera M, Gokhale CS. 2020. Eco-evolutionary agriculture: Host-pathogen dynamics in crop rotations. *PLoS Computational Biology* 16: e1007546
4. Barker H, Dale MFB. 2006. Resistance to viruses in potato. In *Natural Resistance Mechanisms of Plants to Viruses*, ed. G Loebenstein, JP Carr, pp. 341-66: Springer Netherlands
5. Barrett LG, Kniskern JM, Bodenhausen N, Zhang W, Bergelson J. 2009. Continua of specificity and virulence in plant host–pathogen interactions: causes and consequences. *New Phytologist* 183: 513-29
6. Bonman JM, Khush GS, Nelson RJ. 1992. Breeding rice for resistance to pests. *Annual Review of Phytopathology* 30: 507-28

7. Boudreau MA. 2013. Diseases in intercropping systems. *Annual Review of Phytopathology* 51: 499-519
8. Bourget R, Chaumont L, Durel C-E, Sapoukhina N. 2015. Sustainable deployment of QTLs conferring quantitative resistance to crops: first lessons from a stochastic model. *New Phytologist* 206: 1163-71
9. Brown JKM. 2015. Durable resistance of crops to disease: A darwinian perspective. *Annual Review of Phytopathology* 53: 513-39
10. Browning JA, Frey KJ. 1969. Multiline cultivars as a means of disease control. *Annual Review of Phytopathology* 7: 355-82
11. Brun H, Chèvre A-M, Fitt BDL, Powers S, Besnard A-L, et al. 2010. Quantitative resistance increases the durability of qualitative resistance to *Leptosphaeria maculans* in *Brassica napus*. *New Phytologist* 185: 285-99
12. Buoro M, Carlson SM. 2014. Life-history syndromes: Integrating dispersal through space and time. *Ecology Letters* 17: 756-67
13. Burdon JJ, Barrett LG, Rebetzke G, Thrall PH. 2014. Guiding deployment of resistance in cereals using evolutionary principles. *Evolutionary Applications* 7: 609-24
14. Burdon JJ, Chilvers GA. 1982. Host density as a factor in plant disease ecology. *Annual Review of Phytopathology* 20: 143-66
15. Burdon JJ, Zhan J, Barrett LG, Papaix J, Thrall PH. 2016. Addressing the challenges of pathogen evolution on the world's arable crops. *Phytopathology* 106: 1117-27
16. Carolan K, Helps J, van den Berg F, Bain R, Paveley N, van den Bosch F. 2017. Extending the durability of cultivar resistance by limiting epidemic growth rates. *Proceedings of the Royal Society B: Biological Sciences* 284: 20170828
17. Chabas H, Lion S, Nicot A, Meaden S, van Houte S, et al. 2018. Evolutionary emergence of infectious diseases in heterogeneous host populations. *PLoS Biology* 16: e2006738
18. Charlesworth B. 2009. Effective population size and patterns of molecular evolution and variation. *Nature Reviews Genetics* 10: 195
19. Cole AB, Király L, Lane LC, Wiggins BE, Ross K, Schoelz JE. 2004. Temporal expression of PR-1 and enhanced mature plant resistance to virus infection is controlled by a single dominant gene in a new *Nicotiana* hybrid. *Molecular Plant-Microbe Interactions* 17: 976-85
20. Corwin JA, Kliebenstein DJ. 2017. Quantitative resistance: More than just perception of a pathogen. *The Plant Cell* 29: 655-65
21. Créte R, Pires RN, Barbetti MJ, Renton M. 2020. Rotating and stacking genes can improve crop resistance durability while potentially selecting highly virulent pathogen strains. *Scientific Reports* 10: 19752
22. Crete R, Renton M. 2015. *A new model to investigate whether regional crop rotation strategies can protect crops from fungal pathogens*. Presented at 21st International Congress on Modelling and Simulation, Australia
23. Day T, Gandon S. 2007. Applying population-genetic models in theoretical evolutionary epidemiology. *Ecology Letters* 10: 876-88
24. Day T, Proulx Stephen R. 2004. A general theory for the evolutionary dynamics of virulence. *The American Naturalist* 163: E40-E63
25. Dieckmann U, Metz JAJ, Sabelis MW, Sigmund K. 2002. *Adaptive Dynamics of Infectious Diseases: In Pursuit of Virulence Management*. Cambridge: Cambridge University Press

26. Djian-Caporalino C, Palloix A, Fazari A, Marteu N, Barbary A, et al. 2014. Pyramiding, alternating or mixing: comparative performances of deployment strategies of nematode resistance genes to promote plant resistance efficiency and durability. *BMC Plant Biology* 14: 1-13
27. Djidjou-Demasse R, Ducrot A, Fabre F. 2017. Steady state concentration for a phenotypic structured problem modeling the evolutionary epidemiology of spore producing pathogens. *Mathematical Models and Methods in Applied Sciences* 27: 385-426
28. Djidjou-Demasse R, Moury B, Fabre F. 2017. Mosaics often outperform pyramids: insights from a model comparing strategies for the deployment of plant resistance genes against viruses in agricultural landscapes. *New Phytologist* 216: 239-53
29. Ellis JG, Lagudah ES, Spielmeyer W, Dodds PN. 2014. The past, present and future of breeding rust resistant wheat. *Frontiers in Plant Science* 5: 641
30. Ewald J, Sieber P, Garde R, Lang SN, Schuster S, Ibrahim B. 2020. Trends in mathematical modeling of host–pathogen interactions. *Cellular and Molecular Life Sciences* 77: 467-80
31. Fabre F, Bruchou C, Palloix A, Moury B. 2009. Key determinants of resistance durability to plant viruses: Insights from a model linking within- and between-host dynamics. *Virus Research* 141: 140-49
32. Fabre F, Rousseau E, Mailleret L, Moury B. 2012. Durable strategies to deploy plant resistance in agricultural landscapes. *New Phytologist* 193: 1064-75
33. Fabre F, Rousseau E, Mailleret L, Moury B. 2015. Epidemiological and evolutionary management of plant resistance: optimizing the deployment of cultivar mixtures in time and space in agricultural landscapes. *Evolutionary Applications* 8: 919-32
34. Flor HH. 1955. Host-parasite interaction in flax rust - Its genetics and other implications. *Phytopathology* 45: 680-85
35. Fuchs M. 2017. Pyramiding resistance-conferring gene sequences in crops. *Current Opinion in Virology* 26: 36-42
36. Gandon S, Michalakakis Y. 2000. Evolution of parasite virulence against qualitative or quantitative host resistance. *Proceedings of the Royal Society B: Biological Sciences* 267: 985-90
37. García-Arenal F, McDonald BA. 2003. An analysis of the durability of resistance to plant viruses. *Phytopathology* 93: 941-52
38. Garrett KA, Mundt CC. 1999. Epidemiology in mixed host populations. *Phytopathology* 89: 984-90
39. Gigot C, de Vallavieille-Pope C, Huber L, Saint-Jean S. 2014. Using virtual 3-D plant architecture to assess fungal pathogen splash dispersal in heterogeneous canopies: a case study with cultivar mixtures and a non-specialized disease causal agent. *Annals of botany* 114: 863-75
40. Gilligan CA. 2008. Sustainable agriculture and plant diseases: an epidemiological perspective. *Philosophical Transactions of the Royal Society of London B: Biological Sciences* 363: 741-59
41. Gilligan CA, Truscott JE, Stacey AJ. 2007. Impact of scale on the effectiveness of disease control strategies for epidemics with cryptic infection in a dynamical landscape: an example for a crop disease. *Journal of The Royal Society Interface* 4: 925-34
42. Gilligan CA, van den Bosch F. 2008. Epidemiological models for invasion and persistence of pathogens. *Annual Review of Phytopathology* 46: 385-418

43. Grimm V, Johnston ASA, Thulke HH, Forbes VE, Thorbek P. 2020. Three questions to ask before using model outputs for decision support. *Nature Communications* 11: 4959
44. Groth JV. 1976. Multilines and 'super races'. A simple model. *Phytopathology* 66: 937-39
45. Hobbelen PHF, Paveley ND, van den Bosch F. 2014. The emergence of resistance to fungicides. *PLoS ONE* 9: e91910
46. Holmes EE, Lewis MA, Banks JE, Veit RR. 1994. Partial differential equations in ecology: spatial interactions and population dynamics. *Ecology* 75: 17-29
47. Holt J, Chancellor TCB. 1999. Modelling the spatio-temporal deployment of resistant varieties to reduce the incidence of rice tungro disease in a dynamic cropping system. *Plant Pathology* 48: 453-61
48. Huang C, Sun Z, Wang H, Luo Y, Ma Z. 2012. Effects of wheat cultivar mixtures on stripe rust: A meta-analysis on field trials. *Crop Protection* 33: 52-58
49. Janzac B, Fabre F, Palloix A, Moury B. 2009. Constraints on evolution of virus avirulence factors predict the durability of corresponding plant resistances. *Molecular Plant Pathology* 10: 599-610
50. Jeger MJ, Ellis G, Gareth JD. 1981. Disease progress of non-specialised fungal pathogens in intraspecific mixed stands of cereal cultivars. I. Models. *Annals of Applied Biology* 98: 187-98
51. Johnson R. 1983. Genetic background of durable resistance. In *Durable Resistance in Crops*, ed. F Lamberti, JM Waller, NA Graaff, pp. 5-26. Boston, MA: Springer New York
52. Johnson R. 1984. A critical analysis of durable resistance. *Annual Review of Phytopathology* 22: 309-30
53. Kampmeijer P, Zadoks JC. 1977. *EPIMUL, a Simulator of Foci and Epidemics in Mixtures of Resistant and Susceptible Plants, Mosaics and Multilines*. Wageningen, the Netherlands: Centre for Agricultural Publishing and Documentation. 50 pp.
54. Keesing F, Holt RD, Ostfeld RS. 2006. Effects of species diversity on disease risk. *Ecology Letters* 9: 485-98
55. Kermack WO, McKendrick AG. 1927. A contribution to the mathematical theory of epidemics. *Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences* 115: 700-21
56. Kiyosawa S. 1972. Theoretical comparison between mixture and rotation cultivations of disease-resistant varieties. *Annals of the Phytopathological Society of Japan* 38: 52-59
57. Koller T, Brunner S, Herren G, Hurni S, Keller B. 2018. Pyramiding of transgenic Pm3 alleles in wheat results in improved powdery mildew resistance in the field. *Theoretical and Applied Genetics* 131: 861-71
58. Kolmer JA. 1996. Genetics of resistance to wheat leaf rust. *Annual Review of Phytopathology* 34: 435-55
59. Laine AL, Barrès B. 2013. Epidemiological and evolutionary consequences of life-history trade-offs in pathogens. *Plant Pathology* 62: 96-105
60. Laloï G, Vergne E, Durel CE, Le Cam B, Caffier V. 2017. Efficiency of pyramiding of three quantitative resistance loci to apple scab. *Plant Pathology* 66: 412-22
61. Lannou C. 2001. Intrapathotype diversity for aggressiveness and pathogen evolution in cultivar mixtures. *Phytopathology* 91: 500-10
62. Lannou C. 2012. Variation and selection of quantitative traits in plant pathogens. *Annual Review of Phytopathology* 50: 319-38

63. Lannou C, de Vallavieille-Pope C, Goyeau H. 1994. Host mixture efficacy in disease control: effects of lesion growth analyzed through computer-simulated epidemics. *Plant Pathology* 43: 651-62
64. Lannou C, de Vallavieille-Pope C, Goyeau H. 1995. Induced resistance in host mixtures and its effect on disease control in computer-simulated epidemics. *Plant Pathology* 44: 478-89
65. Lannou C, Mundt CC. 1996. Evolution of a pathogen population in host mixtures: simple race–complex race competition. *Plant Pathology* 45: 440-53
66. Lannou C, Mundt CC. 1997. Evolution of a pathogen population in host mixtures: rate of emergence of complex races. *Theoretical and Applied Genetics* 94: 991-99
67. Leach JE, Vera Cruz CM, Bai J, Leung H. 2001. Pathogen fitness penalty as a predictor of durability of disease resistance genes. *Annual Review of Phytopathology* 39: 187-224
68. Levins R. 1966. The strategy of model building in population biology. *American Scientist* 54: 421-31
69. Lô-Pelzer E, Aubertot JN, Bousset L, Salam MU, Jeuffroy MH. 2010. SIPPOM-WOSR: A Simulator for Integrated Pathogen POpulation Management of phoma stem canker on Winter OilSeed Rape: II. Sensitivity analysis. *Field Crops Research* 118: 82-93
70. Lô-Pelzer E, Bousset L, Jeuffroy MH, Salam MU, Pinochet X, et al. 2010. SIPPOM-WOSR: A Simulator for Integrated Pathogen POpulation Management of phoma stem canker on Winter OilSeed Rape: I. Description of the model. *Field Crops Research* 118: 73-81
71. Lo Iacono G, van den Bosch F, Gilligan CA. 2013. Durable resistance to crop pathogens: An epidemiological framework to predict risk under uncertainty. *PLoS Computational Biology* 9: e1002870
72. Lo Iacono G, van den Bosch F, Paveley N. 2012. The evolution of plant pathogens in response to host resistance: Factors affecting the gain from deployment of qualitative and quantitative resistance. *Journal of Theoretical Biology* 304: 152-63
73. Lof ME, de Vallavieille-Pope C, van der Werf W. 2017. Achieving durable resistance against plant diseases: scenario analyses with a national-scale spatially explicit model for a wind-dispersed plant pathogen. *Phytopathology* 107: 580-89
74. Lof ME, van der Werf W. 2017. Modelling the effect of gene deployment strategies on durability of plant resistance under selection. *Crop Protection* 97: 10-17
75. Madden LV, Hughes G, van den Bosch F. 2017. *The Study of Plant Disease Epidemics: The American Phytopathological Society*. 421 pp.
76. Mailleret L, Lemesle V. 2009. A note on semi-discrete modelling in the life sciences. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* 367: 4779-99
77. McDonald BA. 2014. Using dynamic diversity to achieve durable disease resistance in agricultural ecosystems. *Tropical Plant Pathology* 39: 191-96
78. McDonald BA, Linde C. 2002. Pathogen population genetics, evolutionary potential, and durable resistance. *Annual Review of Phytopathology* 40: 349-79
79. Medlock J, Kot M. 2003. Spreading disease: integro-differential equations old and new. *Mathematical Biosciences* 184: 201-22
80. Mikaberidze A, McDonald BA, Bonhoeffer S. 2015. Developing smarter host mixtures to control plant disease. *Plant Pathology* 64: 996-1004

81. Milne AE, Bell JR, Hutchison WD, van den Bosch F, Mitchell PD, et al. 2016. The effect of farmers? Decisions on pest control with *Bt* Crops: A billion dollar game of strategy. *PLoS Computational Biology* 11: e1004483
82. Mundt CC. 1990. Probability of mutation to multiple virulence and durability of resistance gene pyramids. *Phytopathology* 80: 221-23
83. Mundt CC. 2002. Use of multiline cultivars and cultivar mixtures for disease management. *Annual Review of Phytopathology* 40: 381-410
84. Mundt CC. 2014. Durable resistance: A key to sustainable management of pathogens and pests. *Infection, Genetics and Evolution* 27: 446-55
85. Mundt CC. 2018. Pyramiding for resistance durability: Theory and practice. *Phytopathology* 108: 792-802
86. Mundt CC, Brophy LS. 1988. Influence of number of host genotypes units on the effectiveness of host mixtures for disease control: A modelling approach. *Phytopathology* 78: 1087-94
87. Mundt CC, Leonard KJ. 1986. Analysis of factors affecting disease increase and spread in mixtures of immune and susceptible plants in computer-simulated epidemics. *Phytopathology* 76: 832-40
88. Mundt CC, Leonard KJ, Thal WM, Fulton JH. 1986. Computerized simulation of crown rust epidemics in mixtures of immune and susceptible oat plants with different genotype unit areas and spatial distributions of initial disease. *Phytopathology* 76: 590-98
89. Nathan R, Klein E, Robledo-Arnuncio JJ, Revilla E. 2012. Dispersal kernels: review. In *Dispersal Ecology and Evolution*, ed. J Clobert, M Baguette, TG Benton, JM Bullock, pp. 187-210: Oxford University Press
90. Nelson R, Wiesner-Hanks T, Wisser R, Balint-Kurti P. 2017. Navigating complexity to breed disease-resistant crops. *Nature Reviews Genetics* 19: 21
91. Niks RE, Qi X, Marcel TC. 2015. Quantitative resistance to biotrophic filamentous plant pathogens: Concepts, misconceptions, and mechanisms. *Annual Review of Phytopathology* 53: 445-70
92. Nilusmas S, Mercat M, Perrot T, Djian-Caporalino C, Castagnone-Sereno P, et al. 2020. Multi-seasonal modelling of plant-nematode interactions reveals efficient plant resistance deployment strategies. *Evolutionary Applications* 13: 2206-21
93. Nilusmas S, Mercat M, Perrot T, Touzeau S, Calcagno V, et al. 2017. A multi-seasonal model of plant-nematode interactions and its use to identify durable plant resistance deployment strategies. *Acta Horticulturae* 1182: 211-18
94. Ohtsuki A, Sasaki A. 2006. Epidemiology and disease-control under gene-for-gene plant-pathogen interaction. *Journal of Theoretical Biology* 238: 780-94
95. Pacilly FCA, Hofstede GJ, Lammerts van Bueren ET, Groot JCJ. 2019. Analysing social-ecological interactions in disease control: An agent-based model on farmers' decision making and potato late blight dynamics. *Environmental Modelling & Software* 119: 354-73
96. Pacilly FCA, Hofstede GJ, Lammerts van Bueren ET, Kessel GJT, Groot JCJ. 2018. Simulating crop-disease interactions in agricultural landscapes to analyse the effectiveness of host resistance in disease control: The case of potato late blight. *Ecological Modelling* 378: 1-12
97. Pacilly FCA, Lammerts van Bueren ET, Groot JCJ, Hofstede GJ. 2019. Moving perceptions on potato late blight control: Workshops with model-based scenarios. *Crop Protection* 119: 76-87

98. Palloix A, Ayme V, Moury B. 2009. Durability of plant major resistance genes to pathogens depends on the genetic background, experimental evidence and consequences for breeding strategies. *New Phytologist* 183: 190-99
99. Papaïx J, Adamczyk-Chauvat K, Bouvier A, Kiêu K, S. T, et al. 2014. Pathogen population dynamics in agricultural landscapes: The *Ddal* modelling framework. *Infection, Genetics and Evolution* 27: 509-20
100. Papaïx J, Burdon JJ, Lannou C, Thrall PH. 2014. Evolution of pathogen specialisation in a host metapopulation: joint effects of host and pathogen dispersal. *PLoS Computational Biology* 10: e1003633
101. Papaïx J, Burdon JJ, Zhan J, Thrall PH. 2015. Crop pathogen emergence and evolution in agro-ecological landscapes. *Evolutionary Applications* 8: 385-402
102. Papaïx J, David O, Lannou C, Monod H. 2013. Dynamics of adaptation in spatially heterogeneous metapopulations. *PLoS ONE* 8: e54697
103. Papaïx J, Rimbaud L, Burdon JJ, Zhan J, Thrall PH. 2018. Differential impact of landscape-scale strategies for crop cultivar deployment on disease dynamics, resistance durability and long-term evolutionary control. *Evolutionary Applications* 11: 705-17
104. Papaïx J, Touzeau S, Monod H, Lannou C. 2014. Can epidemic control be achieved by altering landscape connectivity in agricultural systems? *Ecological Modelling* 284: 35-47
105. Pariaud B, Robert C, Goyeau H, Lannou C. 2009. Aggressiveness components and adaptation to a host cultivar in wheat leaf rust. *Phytopathology* 99: 869-78
106. Parlevliet JE. 2002. Durability of resistance against fungal, bacterial and viral pathogens; present situation. *Euphytica* 124: 147-56
107. Pietravalle S, Lemarié S, van den Bosch F. 2006. Durability of resistance and cost of virulence. *European Journal of Plant Pathology* 114: 107-16
108. Quenouille J, Montarry J, Palloix A, Moury B. 2013. Farther, slower, stronger: how the plant genetic background protects a major resistance gene from breakdown. *Molecular Plant Pathology* 14: 109-18
109. REX Consortium. 2013. Heterogeneity of selection and the evolution of resistance. *Trends in Ecology & Evolution* 28: 110-18
110. REX Consortium. 2016. Combining selective pressures to enhance the durability of disease resistance genes. *Frontiers in Plant Science* 7: 1916
111. Rimbaud L, Papaïx J, Barrett LG, Burdon JJ, Thrall PH. 2018. Mosaics, mixtures, rotations or pyramiding: What is the optimal strategy to deploy major gene resistance? *Evolutionary Applications* 11: 1791-810
112. Rimbaud L, Papaïx J, Rey J-F. 2018. landsepi: Landscape Epidemiology and Evolution. *R package version 0.0.3*
113. Rimbaud L, Papaïx J, Rey J-F, Barrett LG, Thrall PH. 2018. Assessing the durability and efficiency of landscape-based strategies to deploy plant resistance to pathogens. *PLoS Computational Biology* 14: e1006067
114. Rousseau E, Bonneault M, Fabre F, Moury B, Mailleret L, Grogard F. 2019. Virus epidemics, plant-controlled population bottlenecks and the durability of plant resistance. *Philosophical Transactions of the Royal Society B: Biological Sciences* 374: 20180263
115. Rouzine IM, Rodrigo A, Coffin JM. 2001. Transition between stochastic evolution and deterministic evolution in the presence of selection: general theory and application to virology. *Microbiology and molecular biology reviews : MMBR* 65: 151-85

116. Sacristán S, García-Arenal F. 2008. The evolution of virulence and pathogenicity in plant pathogen populations. *Molecular Plant Pathology* 9: 369-84
117. Sapoukhina N, Durel C-E, Le Cam B. 2009. Spatial deployment of gene-for-gene resistance governs evolution and spread of pathogen populations. *Theoretical Ecology* 2: 229-38
118. Sapoukhina N, Paillard S, Dedryver F, Vallavieille-Pope C. 2013. Quantitative plant resistance in cultivar mixtures: wheat yellow rust as a modeling case study. *New Phytologist* 200: 888-97
119. Sapoukhina N, Tyutyunov Y, Sache I, Arditi R. 2010. Spatially mixed crops to control the stratified dispersal of airborne fungal diseases. *Ecological Modelling* 221: 2793-800
120. Sisterson MS, Stenger DC. 2018. Modelling effects of vector acquisition threshold on disease progression in a perennial crop following deployment of a partially resistant variety. *Plant Pathology* 67: 1388-400
121. Skelsey P, Rossing WAH, Kessel GJT, Powell J, van der Werf W. 2005. Influence of host diversity on development of epidemics: An evaluation and elaboration of mixture theory. *Phytopathology* 95: 328-38
122. Skelsey P, Rossing WAH, Kessel GJT, van der Werf W. 2010. Invasion of *Phytophthora infestans* at the landscape level: How do spatial scale and weather modulate the consequences of spatial heterogeneity in host resistance? *Phytopathology* 100: 1146-61
123. St Clair DA. 2010. Quantitative disease resistance and quantitative resistance loci in breeding. *Annual Review of Phytopathology* 48: 247-68
124. Stukenbrock EH, McDonald BA. 2008. The origins of plant pathogens in agro-ecosystems. *Annual Review of Phytopathology* 46: 75-100
125. Stuthman DD, Leonard KJ, Miller-Garvin J. 2007. Breeding crops for durable resistance to disease. *Advances in Agronomy* 95: 319-67
126. Sun P, Yang XB. 1999. Two properties of a gene-for-gene coevolution system under human perturbations. *Phytopathology* 89: 811-16
127. Suzuki SU, Sasaki A. 2011. How does the resistance threshold in spatially explicit epidemic dynamics depend on the basic reproductive ratio and spatial correlation of crop genotypes? *Journal of Theoretical Biology* 276: 117-25
128. Tabashnik BE, Brévault T, Carrière Y. 2013. Insect resistance to Bt crops: lessons from the first billion acres. *Nature Biotechnology* 31: 510
129. Thrall PH, Barrett L, Dodds P, Burdon J. 2016. Epidemiological and evolutionary outcomes in gene-for-gene and matching allele models. *Frontiers in Plant Science* 6: 1084
130. Thrall PH, Burdon JJ. 2003. Evolution of virulence in a plant host-pathogen metapopulation. *Science* 299: 1735-37
131. van den Berg F, Bacaer N, Metz JAJ, Lannou C, van den Bosch F. 2011. Periodic host absence can select for higher or lower parasite transmission rates. *Evolutionary Ecology* 25: 121-37
132. van den Berg F, Lannou C, Gilligan CA, van den Bosch F. 2014. Quantitative resistance can lead to evolutionary changes in traits not targeted by the resistance QTLs. *Evolutionary Applications* 7: 370-80
133. van den Bosch F. 1993. Optimal cultivar mixtures in focal plant diseases. *Netherlands Journal of Plant Pathology* 99: 41-50
134. van den Bosch F, Akudibilah G, Seal SUE, Jeger M. 2006. Host resistance and the evolutionary response of plant viruses. *Journal of Applied Ecology* 43: 506-16



135. van den Bosch F, Gilligan CA. 2003. Measures of durability of resistance. *Phytopathology* 93: 616-25
136. van den Bosch F, Gilligan CA. 2008. Models of fungicide resistance dynamics. *Annual Review of Phytopathology* 46: 123-47
137. van den Bosch F, Jeger MJ, Gilligan CA. 2007. Disease control and its selection for damaging plant virus strains in vegetatively propagated staple food crops; a theoretical assessment. *Proceedings of the Royal Society B-Biological Sciences* 274: 11-18
138. van den Bosch F, Verhaar MA, Buiel AAM, Hoogkamer W, Zadoks JC. 1990. Focus expansion in plant disease. IV: Expansion rates in mixtures of resistant and susceptible hosts. *Phytopathology* 80: 598-602
139. Van der Plank JE. 1968. *Disease Resistance in Plants*. London-New York: Academic Press. 225 pp.
140. Vyska M, Cunniffe N, Gilligan C. 2016. Trade-off between disease resistance and crop yield: a landscape-scale mathematical modelling perspective. *Journal of The Royal Society Interface* 13: 20160451
141. Waggoner PE, Berger RD. 1987. Defoliation, disease, and growth. *Phytopathology* 77: 393-98
142. Wingen LU, Shaw MW, Brown JKM. 2013. Long-distance dispersal and its influence on adaptation to host resistance in a heterogeneous landscape. *Plant Pathology* 62: 9-20
143. Wolfe MS. 1985. The current status and prospects of multiline cultivars and variety mixtures for disease resistance. *Annual Review of Phytopathology* 23: 251-73
144. Xu X. 2012. Super-races are not likely to dominate a fungal population within a life time of a perennial crop plantation of cultivar mixtures: a simulation study. *BMC Ecology* 12: 16
145. Xu XM. 2011. A simulation study on managing plant diseases by systematically altering spatial positions of cultivar mixture components between seasons. *Plant Pathology* 60: 857-65
146. Xu XM, Ridout MS. 2000. Stochastic simulation of the spread of race-specific and race-nonspecific aerial fungal pathogens in cultivar mixtures. *Plant Pathology* 49: 207-18
147. Zhan J, Thrall PH, Papaix J, Xie L, Burdon JJ. 2015. Playing on a pathogen's weakness: using evolution to guide sustainable plant disease control strategies. *Annual Review of Phytopathology* 53: 19-43
148. Zhou C, Zhou Y. 2012. Strategies for Viral Cross Protection in Plants. In *Antiviral Resistance in Plants*, ed. J Watson, MB Wang, pp. 69-81. Totowa, NJ: Humana Press
149. Zhu S, Morel J-B. 2019. Molecular mechanisms underlying microbial disease control in intercropping. *Molecular Plant-Microbe Interactions* 32: 20-24
150. Zhu Y, Chen H, Fan J, Wang Y, Li Y, et al. 2000. Genetic diversity and disease control in rice. *Nature* 406: 718-22

## Annotated references

Chabas et al. 2018: predictions based on a theoretical model on pathogen evolutionary emergence are tested experimentally using bacteriophages.

Djian-Caporalino et al. 2014: an experimental comparison of pyramiding, rotation and mixture strategies to deploy pepper resistance against nematodes.

Djidjou-Demasse R, Moury B, Fabre F. 2017: this model shows that mosaics are more efficient than pyramids when resistance-adapted pathogens pre-exist.

Kermack & McKendrick 1927: a seminal paper developing compartmental models in human epidemiology.

Lof ME, de Vallavieille-Pope C, van der Werf W. 2017: this model compares different deployment strategies with and without pre-existence of resistance-adapted pathogens.

Papaïx et al 2018: This model shows that epidemiological control and resistance durability are not necessarily correlated.

Rex Consortium 2013: strategies to manage resistance to xenobiotics are compared based on theoretical models and experimental studies.

Rimbaud L, Papaïx J, Barrett LG, Burdon JJ, Thrall PH. 2018: in this model, pyramids are the most durable strategy when resistance-adapted pathogens do not pre-exist.

Rimbaud L, Papaïx J, Rey J-F. 2018: a freely accessible R package (*landsepi*) proposing a unified modelling framework to compare deployment strategies.

van den Bosch & Gilligan 2003: this model shows that the optimal proportion of resistance depends on how durability is measured.

## Related Resources

1. Prototype of a shiny pedagogical interface for the model *landsepi*:  
[https://shiny.biosp.inrae.fr/app\\_direct/landsepi/](https://shiny.biosp.inrae.fr/app_direct/landsepi/)
2. Tepekule B, Uecker H, Derungs I, Frenoy A, Bonhoeffer S. 2017. Modeling antibiotic treatment in hospitals: A systematic approach shows benefits of combination therapy over cycling, mixing, and mono-drug therapies. *PLoS Comput. Biol.* 13: e1005745

## Terms and definitions

- Resistance-adapted pathogen: a pathogen variant adapted to a plant resistance gene. Also referred to as “infective”, “virulent” and “resistance-breaking” in other contexts.
- Complete resistance: a resistance which completely blocks the infectious cycle of **non-adapted pathogens** (e.g. by preventing infection or the production of propagules).
- Demographic stochasticity: refers to the intrinsic uncertainty associated with the life events of each individual in a population (reproduction, dispersal, mutation, death).
- Deterministic: a framework where a set of given inputs invariably yields the same outputs. Deterministic models inform on mean output tendencies.
- Stochastic: a framework where random events are accounted for. Stochastic models inform on mean tendencies and variability of the outputs.
- Environmental stochasticity: refers to random perturbations imposed to a population by its environment.
- Epidemiological control: reduction in disease impact on the host in a given region and during a given period.
- Fitness: ability to transmit genes to the next generation. Here, pathogen fitness is mostly the relative adaptation to different host **genotypes**.
- Genotype: here, plant genotype refers to the set of susceptibility/resistance genes, and pathogen genotype refers to the set of **pathogenicity** genes.
- Maladapted pathogen: a pathogen variant not fully adapted to plant resistance, i.e. infection is possible although the resulting disease is reduced.
- Multi-adapted pathogen: a pathogen variant adapted to several resistance genes. Also referred to as “multi-virulent” and “superpathogen” in other contexts.
- Non-adapted pathogen: a pathogen variant not adapted to plant resistance. Also referred to as “avirulent” or “wild-type” in other contexts.
- Partial resistance: a resistance for which infection by **maladapted pathogens** is not blocked but reduced, resulting in the attenuation of **pathogenicity traits**.
- Pathogenicity trait: a measure of pathogen ability to develop and spread (e.g. infection, reproduction and survival rates, latent and infectious periods) (62).
- Phenotype: here it refers to the occurrence (and amount) of disease, and is mostly determined by the plant-pathogen-environment interaction.
- Qualitative: distributed in discrete classes (by homology with statistics).
- Quantitative: continuously distributed (by homology with statistics).
- Resistance breakdown: loss of effectiveness (complete or partial, sudden or gradual) of plant resistance as a consequence of pathogen adaptation.
- Resistance durability: Initially defined as “the time during which a given resistance remains effective in spite of an environment favourable to disease” (51, 52); the term has since gained a wide range of accepted usages (section 3B).



## Sidebars

### Sidebar 1: Complementarity between models, experiments and observations.

Numerical experiments performed using computer models represent an alternative way of generating results while circumventing logistical, financial, legal and ethical constraints associated with traditional experiments in laboratory or field conditions (30). Models are powerful tools to predict epidemics and guide disease management. With respect to the use of plant resistance, a huge number of deployment strategies can now be tested via such models thanks to the growing capacity of modern computers, even if the number of possible calculations will always be limited by the “computational cost” (i.e. the time required to perform the calculations). However, we are now confronted with the difficulty of manipulating, analysing and synthesising results obtained from high dimension systems. And more fundamentally, modelling is insufficient on its own. Modellers need empirical knowledge and data acquired from experimental and observational approaches in the laboratory or the field, to develop their model, calibrate its parameters and validate or test its predictions.

### Sidebar 2: Should the traditional dichotomy of plant resistance be reconsidered?

Plant resistance is traditionally classified into two distinct categories. On one hand, **qualitative resistance** is described as monogenic (conferred by a single ‘major’ resistance gene typically coding for an NLR [nucleotide-binding domain leucine-rich repeat containing] protein), **complete**, race-specific (i.e. effective only for some strains of a pathogen species) and often considered as non-durable. On the other hand, **quantitative resistance** is described as polygenic (conferred by the additive action of several ‘minor’ resistance genes), **partial**, race non-specific and often hypothesised as durable. However, over-simplification sometimes leads to erroneous assumptions, and several excellent reviews have pointed out exceptions to this long-standing dichotomy (90, 91, 123), such as the wheat gene *Lr34* against rusts and the barley gene *mlo* against powdery mildew (these genes do not belong to the NLR family, and confer almost **complete**, race non-specific and durable resistances). Consequently, we think it is important to disentangle the **phenotype (complete vs partial)**, specificity, genetic inheritance (monogenic vs polygenic), molecular mechanism and **durability** of a resistance. Furthermore, we argue that pathogen **genotypes** can be structured **qualitatively** (i.e. in discrete classes: either **adapted** or not to the resistance) or **quantitatively** (i.e. continuously: more or less adapted). Corresponding modelling approaches are illustrated in Figure 2.

## Figure legends

[\*\*Note to Annual Reviews: Figure 1 was originally drawn by the first author of this article and was re-designed by GRAPHIC Business at our request. We entirely created figures 2 to 5 for this article. All figures are not based on any previously published image.\*\*]

### Gene scale

*Choice of resistance sources*

### Plant scale


*Targeted pathogenicity traits  
Gene pyramiding*

### Field scale

*Cultivar & crop rotations  
Cultivar & crop mixtures  
Complementation with  
pesticide treatments*


### Landscape scale

*Static & dynamic mosaics*



**Figure 1. Nested scales of plant resistance deployment.** A global deployment strategy is the combination of: 1) the appropriate selection of effective genetic resistance sources (e.g. **qualitative** or **quantitative**, **complete** or **partial**) and 2) their spatio-temporal deployment at


plant, field and landscape scales with the aim of mitigating pathogen spread and evolution. Such deployment may be complemented by agronomic practices, biological control and chemical treatments. Possible options available at each scale are indicated.




**Figure 2. Modelling the phenotype of plant-pathogen interactions for resistance deployment.** The **quantitative** description of plant-pathogen interactions provides a general framework (a). Parameters of interest in plant resistance deployment are indicated in bold blue text. In this depiction, the degree of pathogen adaptation to the resistance source is distributed continuously in the population. Gradual pathogen adaptation via continuous mutation or n discrete adaptive steps (each step involving one or more genetic mutations) results in resistance erosion, with a potential **fitness cost** ( $\theta$ ) paid on susceptible hosts. When the pathogen population is split into two genotypic classes (**adapted** or **non-adapted** to the resistance), resistance is broken down in a single step (i.e.  $n=1$ ), resulting in a simple matrix which describes a gene-for-gene interaction (b). Independent of the distribution of pathogen **genotypes**, resistance can be **partial** (i.e. infection is possible with reduced efficiency  $\rho < 1$ ; a and b) or **complete** (i.e. it provides total immunity,  $\rho=1$ ; c to e). Numbers refer to adaptive

dynamic (*italic*) and demo-genetic (**bold**) models listed in Table 1 that have used these formalisations. Symbol ‘\*’ indicates models where **resistance-adapted pathogens** suffer the **fitness** cost on both susceptible and resistant hosts (21, 72, 92, 93, 107, 117).






**Figure 3. Modelling resistance deployment options at plant scale.** Different resistance types have varying phenotypic effects on plants when challenged by a pathogen: lower infection and propagule production rates (IR and PR respectively), shorter infectious period (IP), or longer latent period (LP). Potentially, all these **pathogenicity traits** could be completely or partially impeded by plant resistance, but in published models **complete resistance** has mostly been associated with infection rate. Different sources of resistance can be stacked into the genome of a single plant as a pyramid. Numbers refer to demographic (normal font), adaptive dynamic (italic) and demo-genetic (bold) SEIR models listed in Table 1 and representing a wide range of pathosystems.



**Figure 4. Modelling resistance deployment options at field and landscape scales.** Field-scale strategies mostly include mixtures and rotations, which can vary with regard to the relative proportion of each component and their degree of aggregation in space (mixtures) or time (rotations). These may be complemented with other control methods (e.g. pesticide treatments). Landscape-scale strategies refer to mosaics of fields where resistance is deployed in controlled proportions and degree of spatio-temporal aggregation. Numbers refer to demographic (normal font), adaptive dynamic (italic) and demo-genetic (bold) models listed in Table 1 that have explored these different options.



**Figure 5. What models tell us about optimal landscape organisations and deployment strategies to manage plant resistance with respect to epidemiological and evolutionary disease control.** Conclusions considerably differ depending on the existence or absence of **resistance-adapted genotypes** in the pathogen population prior to resistance deployment. Illustrations are examples of optimal landscape organisations. PY: gene pyramiding; TO: turnover of cultivars; RO: cultivar rotation; MI: cultivar mixture; MO: landscape mosaic; dynMO: dynamic mosaic; '+' stands for combination. Numbers refer to demographic (normal font) and demo-genetic (bold) models listed in Table 1 and representing different pathosystems.

- 1 **Table 1. List of mathematical models used to test, optimise or compare plant resistance deployment strategies against pathogens (a & c) or predict**  
 2 **their evolutionary consequences (b).** A detailed version is in Supplemental Table S1.

Deployment strategy <sup>a</sup>	Resistance efficiency <sup>b</sup>	Distribution of pathogen genotypes <sup>c</sup>	Model structure <sup>d</sup>	Evolution mechanisms <sup>e</sup>	Pre-existence of adapted pathogens	Host spatial structure <sup>f</sup>	Seasonality <sup>g</sup>	OUTPUTS <sup>h</sup>			References
								Epidemiological	Evolutionary	Socio-Economical	
<b>(a) Purely demographic models</b>											
MI	Non-host		DE		no	SE	no	x			(63)
MI*	Non-host		DE		no	SE	no	x			(53, 86-88)
MI	Non-host		IDE		no	SE	no	x			(121)
MI	Non-host		ODE		no	SE	no	x			(127)
MI	Non-host		other		no		no	x			(138)
MO+RO	Non-host		other		no	SE	yes	x			(70)
MI	Non-host		RDE+STOCH		no	SE	no	x			(119)
MO	Non-host		STOCH		no	SE	no	x			(99)
MI	Non-host or Partial		DE+STOCH		no	SE	no	x			(39)
RO	Non-host or Partial		ODE		no		no	x			(22)
MO	Non-host or Partial		STOCH		no	SE	no	x			(104)
MO or MO+MI	Non-host or Partial		IDE+STOCH		no	SE	no	x			(122)
MI	Partial		ODE		no		no	x			(50)
TO or MI	Partial		ODE		no		yes	x			(56)
MI*	Partial		ODE+STOCH		no		no			x	(140)
MO	Partial		ODE+STOCH		no		no	x		x	(120)
MO	Partial		other		no	SE	yes	x			(47)
<b>(b) Demo-genetic models of adaptive dynamics</b>											
RO	Non-host	Continuous	ODE	S-M	no		yes		x		(131)
Pure stand or MO	Non-host or Partial	Continuous	ODE	S-M	no		no		x		(36)
MO	Partial	Continuous	DE	S-M	no	SE	no		x		(102)
MI*	Partial	Continuous	ODE	S-M	no		no		x		(132)
Pure stand	Partial	Continuous	ODE	S-M	no		no	x	x		(134, 137)

(c) Demo-genetic models											
MO	Non-host or Complete	Classes	ODE	S-M	no	SE	yes	x	x	x	(95, 96)
MO or MO+PY	Non-host or Complete	Classes	ODE	S-M	no	SE	yes	x		x	(97)
TO or MO or PY or MO+PY	Complete	Classes	IDE+STOCH	S-M-D	yes/no	SE	yes		x		(73)
MI* or MI*+PY	Complete	Classes	ODE	S	yes		no	x			(94)
MO	Complete	Classes	ODE	S	yes	SI	yes	x			(32)
MO or dynMO	Complete	Classes	ODE	S	yes	SI	yes	x	x		(33)
MI*	Complete	Classes	ODE	S-I-M	yes/no		no	x	x		(135)
MO or dynMO or PY or dynMO+PY	Complete	Classes	ODE	S-M	yes	SI	yes	x			(28)
RO	Complete	Classes	ODE	S-M	yes		yes	x			(92, 93)
MO+PY	Complete	Classes	ODE (within plant) STOCH (between plant)	S-M-D	no		no		x		(31)
MO or MI	Complete	Classes	ODE+STOCH	S-I-M-D	yes		no	x			(71)
MI	Complete	Classes	other	S	yes		no		x		(44)
TO or MO or PY or MO+PY	Complete	Classes	other	S	yes	SI	yes		x		(74)
MI*	Complete	Classes	other	S-M	yes		no		x		(126)
MI or MI+PY	Complete	Classes	RDE	S-R	only single mutants	SE	yes	x	x		(117)
MO	Complete	Classes	STOCH	S-M	no	SE	no		x		(142)
MO or MI or RO or PY	Complete	Classes	STOCH	S-M-D	no	SE	yes	x	x		(111)
MO	Complete	Continuous	STOCH	S-M-D	no	SE	yes	x	x		(103)
MI	Complete+Induced R	Classes	DE		yes	SE	no	x			(64)
MI	Complete or Partial	Classes	ODE	S	yes		no	x	x		(80)
MI	Complete or Partial	Classes	STOCH	S-M	no	SI	no	x	x		(17)
MO or MI	Complete or Partial	Classes or Continuous	IDE	S-M	no		no	x			(72)
MI or MI+PY	Complete or Complete+Partial	Classes or Classes+Continuous	DE	S	yes		no		x		(61)
MI* or MI*+PY	Complete or Complete+Partial	Classes	STOCH	S-M-D	yes		no	x			(114)
MO or MO+PY	Complete or Partial or Complete+Partial	Classes or Continuous or Classes+Continuous	STOCH	S-M-D	no	SE	yes	x	x		(113)

MI	Non-host or Partial	Classes	STOCH	S	yes/no	SE	no	x			(146)
RO	Non-host or Partial	Continuous	ODE	S-M	no		yes	x	x	x	(3)
MI	Partial	Classes	DE	S	yes	SE	yes	x	x		(65, 66)
MI*	Partial	Classes	ODE	S	yes		yes	x	x		(107)
PY+Fungicides	Partial	Classes	ODE	S-M	no		yes		x		(16)
MI	Partial	Classes	other	S	yes/no		no	x		x	(133)
MI	Partial	Classes	RDE	S	yes	SE	no	x			(118)
Pure stand or RO or RO+PY	Partial	Classes	STOCH	S-R	yes		yes	x	x		(21)
MI or MI+RO	Partial	Classes	STOCH	S-D	yes	SE	yes	x			(145)
PY+MI	Partial	Classes	STOCH	S-R-D	only single mutants	SE	yes		x		(144)
Pure stand	Partial	Continuous	IDE	S-M	no		no		x		(27)
Pure stand or PY	Partial	Continuous	STOCH	S-M	no		no	x	x		(8)
MO	Partial	Continuous	STOCH	S-M-D	no	SE	yes		x		(101)

- 3 <sup>a</sup> TO: turnover, MO: mosaic, dynMO: dynamic mosaic, MI: mixture, RO: rotation, PY: pyramiding. Symbol '+' stands for combination, and symbol '\*' indicates models  
4 where mosaics and mixtures cannot be distinguished.
- 5 <sup>b</sup> Resistance may be completely or partially efficient. 'non-host' is indicated when a cultivar is completely resistant to a pathogen without evolution (i.e. the pathogen  
6 can never adapt and infect such host). Symbol '+' stands for combination.
- 7 <sup>c</sup> A distribution in classes refers to qualitative resistance (to which a pathogen adapts via a sudden breakdown), a continuous distribution refers to quantitative  
8 resistance (to which a pathogen gradually adapts via erosion). Symbol '+' stands for combination.
- 9 <sup>d</sup> DE: difference equations, ODE: ordinary differential equations, IDE: integro-differential equations, RDE: reaction-diffusion equations, STOCH: stochastic equations  
10 (accounting for demographic stochasticity). Symbol '+' stands for combination.
- 11 <sup>e</sup> S: selection, I: immigration, M: mutation, D: genetic drift, R: recombination
- 12 <sup>f</sup> SE: spatially-explicit, SI: spatially-implicit, blank for non spatial models.
- 13 <sup>g</sup> seasonality refers to a periodic host dynamic imposing a recurrent reduction in pathogen population size (but not necessarily with genetic drift).
- 14 <sup>h</sup> Epidemiological, evolutionary and socio-economical outputs used in modelling studies, relative to plant health and pathogen population size, pathogen population  
15 genetic composition, and crop yield or economic profit, respectively.

## 16 SUPPORTING INFORMATION

17 **Table S1.** List of mathematical models used to test, optimise or compare plant resistance  
18 deployment strategies against pathogens of cultivated crops, or predict their evolutionary  
19 consequences. Table S1 is a comprehensive version of Table 1 including additional details,  
20 for every modelling study, on:

- 21 - the pathosystem considered;
- 22 - the deployment strategy (e.g. scale of host population and infection unit, numbers of  
23 host and pathogen genotypes, number of resistance sources)
- 24 - model structure (e.g. continuous or discrete-time, sources of demographic and  
25 environmental stochasticity, state variables, fitness cost of pathogen adaptation and  
26 of plant resistance);
- 27 - model outputs.