

A New Approach to Produce Succinic Acid Through a Co-Culture System

Alaa Salma, Rawa Abdallah, Florence Fourcade, Abdeltif Amrane, Hayet

Djelal

► To cite this version:

Alaa Salma, Rawa Abdallah, Florence Fourcade, Abdeltif Amrane, Hayet Djelal. A New Approach to Produce Succinic Acid Through a Co-Culture System. Applied Biochemistry and Biotechnology, 2021, 193 (9), pp.2872-2892. 10.1007/s12010-021-03572-2 . hal-03223915

HAL Id: hal-03223915 https://hal.science/hal-03223915

Submitted on 18 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A new approach to produce succinic acid through a co-culture system

Alaa Salma¹, Rawa Abdallah², Florence Fourcade¹, Abdeltif Amrane¹ & Hayet Djelal^{3*}

¹Univ Rennes, Ecole Nationale Supérieure de Chimie de Rennes, CNRS, ISCR, UMR6226, F-

3500 Rennes, France

²Universite Libanaise, EDST, Centre Azm pour la Recherche en Biotechnologie et ses

Applications, Rue El Mitein, LBA3B Tripoli, Lebanon

³UniLaSalle-Ecole des Métiers de l'Environnement, Avenue Robert Schuman, Campus de Ker

Lann, 35 170 Rennes, France

*Corresponding author: Unilasalle-Ecole des Métiers de l'Environnement, Campus de Ker Lann, 35170 Bruz,

E-mail addresses: hayet.djelal@unilasalle.fr (H. Djelal)

Abstract

Microorganisms can produce a wide range of bio-based chemicals that can be used in various industrial applications as molecules of interest. In the present work, an analysis of the power production by pure culture, co-culture, and sequential culture was performed. In this study, both the mono-culture and the co-culture strategies of Actinobacillus succinogenes with Saccharomyces cerevisiae as carbon sources to produce succinic acid using glucose and fructose were examined. The cultures were performed in batch mode and a great attention was paid to the co-culture system to improve the biosynthetic pathway between A. succinogenes and S. cerevisiae by combining these two strains in a single fermentation process. Under microaerobic and anaerobic conditions, the process was characterized in terms of sugars concentration, cell density, metabolites, yield (mol-C products/ mol-C sugars), the temperature conditions for productivity and pH. The results showed that the process could consume glucose and fructose and could adapt to different concentrations of the two sugars more quickly than by a single organism and the best results were obtained in a sequential co-culture recording 0.27 mol L⁻¹ of succinic acid concentration and a volumetric productivity of 0.3 g L⁻¹ h⁻¹. Under the investigated operating conditions, the combination of these two strains in a single reactor produced a significant amount of succinic acid (0.70 mol-C SA/ mol-C substrates). A simultaneous and sequential co-culture strategy can be a powerful new approach in the field of bio-based chemical production.

Key-words: Fermentation; Co-culture; sequential culture; bio-based chemicals; Succinic acid.

Introduction

An interest in the bio-based chemicals production from sugars has strongly increased in the last decade, as these sugar-derived products can potentially replace oil-derived chemicals such as succinic acid [1]. Traditionally, succinic acid (SA) is derived from fossil sources via the hydrogenation of maleic anhydride and considered as a key platform chemical as it is used in the production of a wide range of products, from pharmaceuticals to green solvents, fibers and bioplastics [2]. There is a tremendous opportunity for SA biosynthesis: indeed bio-based SA is interesting and advantageous because its production does not contribute to the accumulation of carbon dioxide (CO₂) into the environment and can thus be considered as a "green" chemical [3, 4]. SA fermentation is relatively well studied thanks to an array of microorganisms and under different conditions [5, 6]. The conventional production of bio-SA involves anaerobic bacterial fermentation of pure or mixed sugars with the addition of CO₂ sources, such as Actinobacillus succinogenes [7, 8]. Previous studies showed that the production of SA could be raised by controlling the supply of CO₂ in the fermentation media, which can significantly increase the product cost [9]. A. succinogenes can be used but require an anaerobic environment (nitrogen bubbling), as well as dissolved CO₂ in the fermentation broth, such as MgCO₃, which definitely controls the metabolic flux of carbon. Also the activity of enzymes including phosphoenolpyruvate carboxykinase (PEPCK) represent crucial steps for succinic acid bioproduction by succinate-producing bacteria [10]. To create a potential method, it is important to improve its efficiency and sustainability. To do so, two key tools were used in various fields of study to improve the process output : more precisely to optimize the operating conditions and to manipulate/steer the microbial community, which requires the improvement of the existing knowledge on the relationship between the microbial community structure and operational conditions [11]. Several studies related to the impact on SA outputs of the operating conditions, in different variations and/or combinations, such as the pH [12], temperature [13],

culture conditions [14] and the standardised or various types of substrates [15] were reported, evaluated and investigated. Apart from the effects of the operating parameters, some studies focused on metabolic pathways such as turn on/off gene pathways [16]. Most previous studies focused on finding correlations between the operating parameters and the SA production [4, 7, 10, 13, 15]; however only a few investigated the relationships with the microbial community in the same environment [17]. Different processes have been adopted for SA fermentation: batch fermentation, fed-batch fermentation, free-cell continuous fermentation, continuous fermentation with immobilized cells [5, 7, 18]. Although most products in industrial biotechnology are produced by only one bacterium, mixed culture fermentation has also frequently been employed [17, 19]. Mixed culture biotechnology (MCB) could become an attractive alternative to traditional pure culture-based biotechnology enabling the production of a narrow spectrum of chemicals and/or bioenergy issued from pure or mixed substrates [17, 20]. Mixed-culture fermentations contain inoculum always consisting of the mixture of two or more known or unknown species [21]. They can be the mixed culture of organisms belonging to one microbial group, like bacteria, or they can consist of the mixture of fungi and bacteria organisms or fungi and yeasts organisms or other combinations whose components are quite unrelated [19, 22]. Generally, co-culture was defined as anaerobic or aerobic incubation of different specified microbial strains under aseptic conditions and it seems to be advantageous compared to pure culture fermentation due to the potential for synergistic utilization of the metabolic pathways of all involved microorganisms in a co-culture system [23]. Thus, it may result in increased yield, improved product qualities, and more efficient substrate utilization. It has been widely used in the production of bio-chemicals, biofuel, biopolymers, and enzymes [24]. The main motivation behind employing co-culture in this work is to examine the interactions between microorganisms and to develop new cell approach techniques to produce platform molecules [25]. The microorganisms subsequently consuming sugars must have lower

productivities for the generation of a product than the organisms simultaneously consuming sugars [26].

A. succinogenes was used as succinic acid producer in several studies due to its simple medium requirements for strain culture and relatively high product yield [8, 10, 27]. However, in the context of a large-scale succinic acid production, the use of both MgCO₃ and N₂ gases can become prohibitively expensive due to their high prices, and thus, considered as shortcoming in SA production on high-scale. To overcome these limitations, we suggest a new strategy for an eco-efficient SA production based on a partnership between yeast producing CO₂, *Saccharomyces cerevisiae*, with SA producing bacteria, *A. succinogenes*. In this system, *S. cerevisiae* digested glucose to produce ethanol, glycerol and CO₂ as a carbon source; the CO₂ produced by yeast respiration during fermentation of sugars could be fixed by *A. succinogenes*. Although, *A. succinogenes* can use glycerol as a substrate and in return produce SA. Thus, *S. cerevisiae* is employed in SA/ethanol production and the two species could form an eco-efficient co-culture system, which performed "better together" for biofuel and biochemical generation. It can be a reliable, cheap, easy to manage and sustainable alternative strategy that could effectively replace the addition of MgCO₃ and the N₂ sparging.

The concept focuses on the possibility of the easy "design" of a co-culture approach, in simultaneous and sequential cultures, that could be a bridge between feedstocks and a bio-based production. At this point, we start creating a synthetic co-culture of *A. succinogenes* (DSM 22257) and *S. cerevisiae* (CLIB 95) in order to produce SA by an efficient joint use of sugar mixtures. In this report, the results of this new method producing direct SA from substrates are recorded and Batch data of viability, residual sugars, yield, productivity and SA concentration using the fermentation process of *A. succinogenes* – *S. cerevisiae* co-cultures are discussed.

Materials and Methods

Microorganisms and the inoculum preparation

The Deutsche Sammlung von Mikroorganismen und Zellkulturen (Institute DSMZ-German Collection of Microorganisms and Cell Cultures) issued *A. succinogenes* (DSM 22257) and the *CIRM* (Centre National de Ressources Microbiennes, France) provided yeast, *S. cerevisiae* CLIB 95.

The Bacterial inoculum was prepared as follows: The bacterium was suspended in a Trypton Soya Broth (TSB) and preserved on Tryptic Soya Agar (TSA) plates and slants. Cells were incubated at 37°C for 24-48 hours in an incubator (VWR® INCU- Line), which was the time required for the microorganism to reach the exponential growth phase. As for yeast species, *Saccharomyces cerevisiae* were maintained at 4°C on a Sabouraud glucose agar 4% slant containing (g L⁻¹): 40 g glucose, 10 g peptones, 15 g of agar; then cells were incubated at 28 °C in the incubator for 24 hours. In order to generate bacterium and yeast inocula, culture suspensions were transferred to 50-mL centrifuge tubes, resuspended in KCL 150 mM and incubated in an incubator shaker (New Brunswick, INNOVA 40, NJ, USA) at 37 °C, 150 rpm for 24 and 48 hours for yeast and bacteria respectively.

As for the co-culture, cells were aseptically centrifuged (1800 g, 4°C and 5 min) in a centrifugation (Thermo-Fisher, scientific, HERAEUS Megafuge 16R), suspended in 10 mL KCl (150 mM) at the end of the incubation period. For the preparation of the inoculum, 2 mL of suspended culture cells were transferred into 250 mL Anaerobic bottles containing 200 mL of culture medium. Depending on the experimental set-up, the populations were mixed at a different % v/v ratio of bacterial-yeast populations, respectively (0.25;0.75), (0.5;0.75), (0.75;0.25), with an equal starting optical density. Co-cultures could potentially be carried out in a 200 mL volume range at 37° C and 150 rpm for 48 hours and population densities were

measured at 660 nm. Cells from co-culture were centrifuged and the suspension obtained in 10 mL of KCl 150 mM was used for inoculation.

Fermentation media contained carbon and nutrient sources (per liter): 32 g glucose, 32 g fructose and 0.53 g NH₄Cl, and minor components including inorganic salts, buffers and cofactors (per liter): 3 g K₂HPO₄, 3 g KH₂PO₄, 1.25 g NaCl, 0.3 g MgCl₂·6H₂O, 0.3 g CaCl₂·2H₂O, NaH₂PO₄, Na₂HPO₄. Fermentation media were sterilized by filtration on a 0.22 µm sterile membrane filter (Sartorius), when setting up the fermenters.

Fermentation experiments

Fermentations were achieved under optimized conditions at 37°C in 200 mL of synthetic media in anaerobic conditions, by inoculating different biomass concentrations of *A. succinogenes* and *S. cerevisiae*. The inoculations were performed in the following way: 1) Co-culture (the inoculation of *A. succinogenes* with *S. cerevisiae* simultaneously); 2) Sequential inoculations of *S. cerevisiae*, followed by the inoculation of *A. succinogenes* after 48 hours, were run in a 3L-capacity benchtop reactor with a volume of 1800 mL. *A. succinogenes* and *S. cerevisiae* pure cultures fermentations were also run as controls. As mentioned previously, both strains were pre-cultured and the inoculated volumes were calculated according to the final volume present in anaerobic bottles. A fermentation temperature of 37°C and stirring speed of 150 rpm were maintained throughout the incubation period. During the fermentations, samples were aseptically drawn (5 mL) from the broth at various time points -24, 48, 72 and 94- in order to follow up the microbial growth, sugar consumption and the production or uptake of acids (e.g., succinic, formic, acetic, and lactic acid) and other compounds (e.g., ethanol).

Analyses

Cell growth was detected by measuring the optical density (OD) of the sample at 660nm (λ 660) using a UV-VIS spectrophotometer (PRIM SECOMAM, Milton Roy, Ivyland, PA, USA). OD measurements were carried out immediately after sampling by using 4 ml plastic cuvettes, and dionized water was used as blank (zero absorbance). The sample was homogenized and then analyzed. Samples were collected from the broth every 24 hour and centrifuged at 1800 g for 5 min. The supernatants were then transferred to 15 mL tubes and were analyzed according to previously published methods [28]. Briefly, sugars (glucose, and fructose), organic acids (succinic, acetic acid,) and others were detected by HPLC using BioRad HPX-87H column at 45 °C with 0.01 N H₂SO₄ mobile phases flowing at 0.7 mL/min.

Statistical analyses

The analysis of data was carried out thanks to an ESP script (EXCEL software program 2016) using T-test (p < 0.05) and was reported as a mean and standard deviation.

Results

Comparative analysis of the production of a pure culture of A. succinogenes and S. cerevisiae in 250 mL batch bottles

In this experiment, two cultures were inoculated as pure cultures in separate batch bottles. The readings were carried out for two days. Table 1 shows the potential and current comparison between the *A. succinogenes* and *S. cerevisiae* pure cultures. During fermentation, *A. succinogenes* produced SA under anaerobic conditions and no other acids were produced, in particular lactic acid. This result shows the capacity of *A. succinogenes* to ferment glucose and fructose that are used as carbon sources to synthesize and produce succinic acid. [29] report that *A. succinogenes* prefer consuming glucose, but it can also metabolize glycerol (by-products

of the ethanol fermentation process naturally produced by *S. cerevisiae*). Fermentation performances differ for each and every microbiome. Likewise, *S. cerevisiae* produce glycerol, ethanol and CO₂ in the fermentation media. About 0.2 mol L⁻¹ of SA was produced using *A. succinogenes* after 48 hours, respectively corresponding to a yield of 0.6 mol-C SA/mol-C substrate and 0.49 g L⁻¹ h⁻¹ productivity. However, when *S. cerevisiae* was cultivated, ethanol and glycerol were the metabolites produced by this microorganism recording a value of about 0.07 mol L⁻¹ for each metabolite. Their ability to grow and undergo metabolic processes in both aerobic and anaerobic conditions are remarkable characteristics of *S. cerevisiae*. In aerobic conditions, *S. cerevisiae* performs cellular respiration through the oxidative phosphorylation of pyruvate, producing CO₂. In anaerobic conditions, *S. cerevisiae* turns fermentative metabolism into ethanol and carbon dioxide (as the primary fermentation metabolites) as the cells strive to produce energy and regenerate the NAD⁺ coenzyme [30]. Henceforth, *A. succinogenes* and *S. cerevisiae* were chosen for the fermentation study as co-cultures.

Effect of the inoculation mode on the microbial growth during the fermentation in 250 mL bottles using a % ratio of (0.5 *A. succinogenes*; 0.5 *S. cerevisiae*)

Fig 1 reveals the microbial evolution (OD_{660nm}) of pure cultures and co-cultures with a % volume ratio of (0.5; 0.5) corresponding to 0.04 of *A. succinogenes* and *S. cerevisiae* starter optical density in microaerobic and anaerobic conditions (Fig 1 a- and b-, respectively). A significant difference was found in the cell density between the co-cultures and the *A. succinogenes* single culture (p < 0.05) during fermentation. The results indicate that the growth of *A. succinogenes* was not limited by *S. cerevisiae* regardless of the combination. On the contrary, these were *A. succinogenes* that decreased the growth of *S. cerevisiae* without causing their deaths. The microaerobic co-cultures showed a slight decrease in their evolution during

fermentation in comparison with the *S. cerevisiae* pure culture control. The evolution difference between the co-culture and the *S. cerevisiae* single culture was not obvious, however, and the cell density in *S. cerevisiae* was more than the one obtained by the co-culture (Fig 1-a). During the anaerobic fermentation, when both strains were simultaneously inoculated together (Fig. 1 b), the co-cultures persisted for the first 3 days with an OD higher than the one obtained from the monoculture of *A. succinogenes*, then reducing gradually. *Sequential cultures* attained and retained optical cell densities more important than the ones achieved in the pure culture and in anaerobic co-cultures at the end of the fermentation time. When *A. succinogenes* was sequentially supplied as a starter after 48 hours, the *S. cerevisiae* became the dominant species throughout the fermentation process in comparison with *A. succinogenes*.

Effect of MgCO₃ on a simultaneous co-culture fermentation

To evaluate the effect of MgCO₃ in the fermentation, a simultaneous co-culture was inoculated with a % ratio of (0.5; 0.5) corresponding to 0.04 of *A. succinogenes* and *S. cerevisiae* starter optical density. Fig 2 a- and b represent the sugars and the evolutions of the products profiles with or without the addition of MgCO₃.

As shown in Fig. 2, the use of sugar was not significantly different in (a) and (b). The glucose and fructose levels decreased significantly in the microaerobic co-culture fermentation: more than 91% was consumed. These data confirmed that the co-culture system can consume both glucose and fructose, whereas ethanol and glycerol were produced - about 0.1 mol L⁻¹ and 0.06 mol L⁻¹ respectively - and the produced amounts were nearly similar irrespective of the presence of the MgCO₃ supplement. SA production was monitored and found to remain negligible throughout the culture in the absence of MgCO₃ (Fig 2-a); while 0.04 mol L⁻¹ SA was produced in the presence of MgCO₃ (Fig 2-b), corresponding to a yield of 0.1 mol-C SA/ mol-C substrates and a production of 0.05 g L⁻¹h⁻¹. It confirms the need for a mineral carbon source. The evolution of CO_2 depends on the extent of sugars being metabolized by *S. cerevisiae* and used by *A. succinogenes*. The divergence in the SA concentration observed between its two values could be attributed to the release of CO_2 by *S. cerevisiae*, indicating the stimulation of a CO_2 production and yeast metabolism during the stage of fermentation. Therefore, the assimilation of the inorganic carbon source showed the potential of the two strains to be used as co-cultures in a microaerobic fermentation to increase the persistence of CO_2 and to improve the SA production process quality (supplementary table S1). The CO_2 production in the co-cultures correlates with the transcriptional up-regulation of the glucose fermentation pathway genes from *S. cerevisiae* [31] and the consumption rates correlate to the fixation of CO_2 to produce SA from *A. succinogenes* [32]. The stimulatory effects that occur during fermentation correspond to the direct response of *S. cerevisiae* and *A. succinogenes* to the presence of each other.

Influence of the ratio of *S. cerevisiae*- *A. succinogenes* on the performance of simultaneous microaerobic and anaerobic co-culture during fermentation.

The variation of the biomass ratio in the culture medium may significantly affect the values of the fermentation parameters calculated from the measured concentrations of the substrates and/or products. As shown in the above results, SA could be produced by the co-culture of *S. cerevisiae* and *A. succinogenes*. To evaluate the co-culture fermentation on the SA production, the co-culture of *S. cerevisiae* and *A. succinogenes* were provided to the media at different % v/v ratios (0.25; 0.75; 0.25) corresponding to (0.04; 0.06) starter optical density and compared to their mono-cultures. Fig. 3 reveals the differences between the concentrations of metabolites in the fermentation with a different biomass ratio in the co-culture without MgCO₃. To investigate the impact of biomass ratios on the performance of fermentation, the metabolites in the culture were compared in microaerobic and anaerobic conditions. For both inoculation

ratios, the SA production from co-cultures fermentations was only observed in anaerobic conditions and was significantly reduced in microaerobic conditions (see fig 3 a, and b). The maximum value of the SA was obtained in anaerobic conditions from a co-culture % ratio of (0.25; 0.75) of S. cerevisiae- A. succinogenes, 0.23 mol L^{-1} (Fig 3-a); while only 0.03 mol L^{-1} ¹ SA was produced in microaerobic conditions regardless the biomass ratio corresponding to 0.28 g $L^{-1} h^{-1}$ and 0.037 g $L^{-1} h^{-1}$, respectively of SA productivity, wherein the fermentation activity of the A. succinogenes strain was limited. Regarding the formation of ethanol, there was a substantial reduction in the ethanol concentration from all the co-cultures at the end of the fermentation time (96 hours) compared to that obtained from a co-culture % ratio of (0.75; 0.25) of S. cerevisiae- A. succinogenes. The variation of the ratio of co-culture shows that the total production of (0.75; 0.25) of S. cerevisiae-A. succinogenes was obviously slower than the one obtained by (0.25; 0.75) of S. cerevisiae- A. succinogenes. However, results show that the highest value of SA was obtained in the anaerobic mono-culture of A. succinogenes. Biomass concentrations deduced from Dry cell weight (DCW) decreased at the end of the fermentation (96 hours) for all the co-cultures and recorded a maximum value of 0.40 g L⁻¹, 0.49 g L⁻¹ for figure 3-a, and -b, respectively, as a result of the increased formation of product and substrate limitations which provoked a drastic decrease of the microbial biomass.

Influence of the inoculation mode

To investigate the effect of the inoculation mode on the co-culture fermentation, the two strains were inoculated simultaneously and/or in different orders from an optical cell density of 0.03 starter strains (table 2) with a constant culture time for all the experiments. In the microaerobic co-cultures, significant amounts of SA were produced; while no production was observed in the pure culture of each strain. In addition, the comparison of the simultaneous and the sequential co-cultures showed a slightly higher amount in the case of the sequential co-culture.

As mentioned above, SA was produced by A. succinogenes in anaerobic fermentation while ethanol and glycerol were produced by S. cerevisiae. In all the experiments, the co-culture of A. succinogenes and S. cerevisiae produced more metabolites than in pure cultures. The SA obtained from the sequential 48-hour cultures revealed a remaining in sugar content, while it turns to fall in the co-culture fermentations profile that were analogous in order of metabolites to that obtained from the fermentation of S. cerevisiae pure cultures. Data showed significant production of SA in all the co-cultures, with a decrease of pH, recording a value of 5.56, 5.32 for microaerobic and anaerobic co-cultures and 5.24 for sequential co-cultures. The SA produced from co-cultures in microaerobic and anaerobic conditions was significantly lower when compared with the anaerobic control A. succinogenes, while the glycerol amount produced rose in the sequential fermentations, wherein the fermentation performance of S. cerevisiae strain was important (table 2). Considering the organic acid production, an increase in SA levels was noticed in the sequential culture at 48 h when compared to the *co-culture*, while that from the pure fermentation of A. succinogenes in anaerobic culture was greater when compared to those obtained by the simultaneous and the sequential 48 h aerobic co-cultures. As for the sugars contents present in the fermentations media, the data displays a high

As for the sugars contents present in the fermentations media, the data displays a high consumption of the glucose and fructose; by 85%, 90% for both microaerobic and anaerobic simultaneous co-cultures and by 60% for sequential 48 h co-culture fermentations with respect to the control of *A. succinogenes*. On the other side, the microaerobic co-cultures lead to a decrease in pH, with a succinic acid production, which was significantly detected, even if it remained considerably below the control level of *A. succinogenes* anaerobic fermentation. A development reflecting that the co-culture of *A. succinogenes* with *S. cerevisiae* generally showed the ability to produce SA, even in microaerobic conditions.

Sequential 48-hour cultures led to a $0.14 \text{ g L}^{-1} \text{ h}^{-1}$ SA production rate in a lab-scale (1.8 L), reaching to about an 85% substrate conversion efficiency, recording an overall yield of 0.69

mol-C of added chemical produced/ mol-C sugars consumed, with 54% of carbon content belonging to succinic acid. The results emphasize the efficiency of the co-culture for the biological conversion of sugars to SA and might work in parallel to other succinic acid production processes such as the chemical process, or the anaerobic fermentation. Even though the system differs from earlier studies, the results are consistent with those obtained by [7, 29, 33] in terms of SA concentration, yield and productivity. To sum up, this provides a process for the batchwise fermentation of succinic acid, wherein a carbon dioxide from *S. cerevisiae* is maintained in a fermentation vessel then withdrawing the broth from the vessel and re-filling the vessel with a newly substrate, and preparing it for the reception of a subsequent batch of *A. succinogenes* to produce SA by fermenting the sugars.

Effect of the fermentation temperature

Fig. 4 (a- and b) present the evolution of the concentration of sugar during the fermentations in *S. cerevisiae* and *A. succinogenes* control cultures and in the co-cultures at the different optimal fermentation temperatures of 28 °C and 37 °C, respectively. At both temperatures, the co-culture fermentations evolved more gradually than in the pure strain culture fermentations expected for the *S. cerevisiae at 28*°C. The biomass progression in terms of the OD, pH, residual sugars concentrations and product concentrations in the pure culture and the co-cultures fermentations are given in table 3 at 28 and 37°C. In the pure culture, *S. cerevisiae* start from 0.04 at t0 and achieved its highest optical density value in 16 hours (around 1.2), which remained almost constant till the end of fermentation (4 days), and the remaining sugar was below 0.02 mol L⁻¹. Once the two microbial strains were co-inoculated (*A. succinogenes and S. cerevisiae*), with 0.04 starter OD, the cell concentration was less than that of *S. cerevisiae*, and reached a value of 0.38 and 0.36 at 28°C and 37°C respectively, when the fermentation was almost over (after 96 h). Thus, at these fermentation conditions (28°C), *A. succinogenes*

showed an inhibitory activity in the production of SA even in anaerobic conditions (0.07 mol L⁻¹ SA) when compared to the amounts obtained at 37°C. The compositional profiles of parameters collected in pure and co-culture fermentations at 28°C and 37°C are listed in Table 3. At 37°C and contrarily to 28°C, the co-cultured A. succinogenes and S. cerevisiae showed a decrease in the amount of ethanol and remaining sugar, when compared to the control of S. cerevisiae alone. With respect to S. cerevisiae controls, the co-culture mode indicated a notable increase in the ethanol production at 28°C, whereas a reduction in ethanol was noticed at 37°C. Regarding the total production activity, the levels significantly increased from the coculture at both 28°C and 37 °C, in comparison to their relevant S. cerevisiae controls at 37 °C. This increase in ethanol at 28°C (0.08 mol L⁻¹) was significantly higher than at 37°C (0.05 mol L⁻¹). There was a substantial increase of glycerol formed, when the co-cultures were run at 28°C $(0.06 \text{ mol } L^{-1})$ with regard to the control. The coupling of A. succinogenes -S. cerevisiae in microaerobic co-culture at 37 °C, has a positive impact on the sugar consumption, resulting in SA formation with levels of 0.22 mol L^{-1} , with respect to SA levels obtained from A. succinogenes pure cultures (0.3 mol L⁻¹). This suggests that SA production in microaerobic coculture is primarily impressed by an S. cerevisiae coupling and by the different temperature conditions.

Considering the sequential 48 hour- fermentations (Fig 5-a), no production of SA was observed before the bacterial inoculation and the temperature shift at 37 °C; the final SA value was 0.27 mol L⁻¹ corresponding to a 0.33 g L⁻¹ h⁻¹ productivity. Moreover, there was a general decrease in the profile of volatile compounds production with an increase of succinic acid; when the fermentations were performed at 37°C, with a smaller variation for *S. cerevisiae* and *A. succinogenes* monocultures. Meanwhile, as compared to the respective controls along with the highest SA production, there was a significant glucose consumption with a substantial decrease of glycerol which might be attributed to the use of the yeast/bacteria as carbon source described in many papers [15, 28, 34]. The percentage of Dissolved Oxygen (DO) in the medium Fig 6b was found to have a considerable influence upon the OD value under conditions of rapid fermentation at 28° C. At T0, the DO content in the substrate was 100%, the ethanol concentration reached about 0.05 mol L^{-1} in the substrates containing 43% DO after 48 h, and the glycerol concentration reached about 0.04 mol L^{-1} after 48 h, then decrease to about 0.005 mol L^{-1} after 120 h in the media containing 43% and 24 % DO in the fermentation in which the DO was maintained at about 24%, recording a maximum value of 0.27 mol L-1 of SA and a pH value of 4.6 after 120h.

Discussion

Microbial co-cultures can be used to produce a variety of products ranging from small molecules, such as hydrogen or short chain fatty acids, to more products, such as methane, acetate, butanol, pyruvate, butyrate and alcohols (table 4). The co-culture strategy for SA production has been known for a long time. On the contrary, the concepts of the co-cultures strategies are new. To our knowledge, the co-culture of *A. succinogenes* and *S. cerevisiae* was not investigated beforehand. Here, in the present work, the co-culture was proposed as an alternative inoculation strategy to the traditional fermentation culture to produce succinic acid. It was previously checked that co-cultures of yeasts/bacteria pairs carried out in 250 mL bottles gave similar results as those obtained during pure cultures. Tremendous numbers of data have been published regarding the function relayed on bacteria or yeasts to produce chemicals alone, yet, reports on "if" or "how" they interact mutually together, start to be in the scope of interest recently. In both described co-cultures, the *A. succinogenes* strain was able to grow on in the presence of *S. cerevisiae*, using glucose and fructose as substrates. As observed, the co-culture had only a little effect on the optical cell density compared to the pure cultures; however, *S. cerevisiae* led to a higher cell density than *A. succinogenes* over the whole culture (Fig 1). This

might be due to the fact that *S. cerevisiae* displayed a higher inoculum size which can favourably impact the optical density [35]. The simultaneous co-occurrence of both strains had a smaller impact than the pure cultures in the fermentation systems. Thus, next to providing the co-substrate for the partner strain, the properties of *S. cerevisiae* strain make co-cultivation possible [36]

The study was realised in a synthetic medium to obtain a reproducible medium with a controlled and non-limiting composition. In microaerobic conditions, the advantage of the co-culture strategy over the monoculture could be due to the presence of yeast from the beginning in a medium lacking MgCO₃ unlike the fermented medium used for monocultures (Fig. 2 and Table 2). Some of the yeast inhibitory metabolites produced by the *S. cerevisiae* strain, such as ethanol, were measured at the end of the co-culture (Fig 2, 3 and Table 2). These inhibitory compounds are known to limit bacterial growth and could reduce the ability of bacteria to synthesize SA at different levels depending on their concentrations and the medium composition and pH during co-culture [16, 37–39] Even though these metabolites gradually occurred during the sequential culture, which lasted for 144 h of the co-culture, the bacteria had time to grow and specially to produce SA (Fig. 4 and Table 3). The SA amounts were higher than those obtained in the simultaneous co-culture, which might be referred to the accumulation of CO₂ in the reactor which stimulates the production of succinic acid when applied directly to the bacteria, in addition to the reactor form and volume used in a sequential culture which is different with that used in the simultaneous co-culture.

It is generally recognised that glucose is consumed rapidly by both of *S. cerevisiae* and *A. succinogenes* [8, 40]. Accordingly, in sequential 48h fermentations with *S. cerevisiae* as a starter, it was observed that the residual fructose concentration was substantially greater, with a higher discrepancy between glucose and fructose concentrations (Fig 4). From the collected data, we deduced that a lower fructose amount was used during the sequential culture if

compared to the fructose consumption in pure *S. cerevisiae* and simultaneous co-cultures. As observed, the consumption of both sugars was less than those obtained during the sequential culture.

In our study, considering the simultaneous microaerobic co-culture, besides the ethanol and glycerol productions, a stimulatory effect of *S. cerevisiae* on *A. succinogenes* to produce SA was observed, as drawn from the comparison to the *A. succinogenes* monoculture. Based on a physiological analysis, it was noticed that co-cultivation resulted in significant SA production by *A. succinogenes* during the whole culture.

Interestingly, no major changes were found regarding biomass amounts when comparing monoand co-culture conditions, suggesting that no changes took place on the genetic level. The product shift appears to be caused by changes in the thermodynamics of the system resulting in more CO_2 production as described by [41], inducing favorable conditions for succinate production as a result of an *A. succinogenes* metabolism. An example of a metabolic shift driven by a metabolic interaction is the co-cultivation of *Escherichia coli* grown on glycerol together with the formate-oxidizing methanogen *Methanobacterium formicicum*. Natively, *E. coli* produces mainly ethanol and formate as end-product of glycerol fermentation, with succinate and acetate as byproducts. In co-culture with *M. formicicum*, the spectrum shifts to succinate production instead [42]. A similar interaction also seems to have taken place in the case of the sequential 48 h co-culture, facilitating substrate and CO_2 conversion by the *A. succinogenes*.

For the sequential co-culture fermentation from substrate to succinic acid, the main challenges were the requirements of the involved microorganisms in terms of reactor set-up, medium composition and product synthesis. Optimizing product yield and productivity for a certain process usually addresses the needs of the organism involved. Since sequential co-culture fermentation uses at least two different organisms, the key aspect for sequential co-culture fermentation is either a medium compromise for both organisms or the compatibility of the first (optimized) medium for the second organism in terms of product synthesis. Furthermore, the second organism has to be able to use the product of the first process as a carbon source. The combination of both aspects must be fulfilled to achieve an optimal value added chain from sugars to succinic acid. Exchange of metabolites synthesized in different microbes can take also place. An example of this is the co-culture of *Clostridium ljungdahlii* and *Clostridium acetobutylicum* (table 4). While the products 2,3-butanediol and isopropanol were not formed from glucose by monocultures, the co-cultivation of *C. ljungdahlii* and *C. acetobutylicum* have resulted in the increased formation of 2,3-butanediol and isopropanol [43].

However, co-culture technology can display some drawbacks, which should be taken into consideration, such as the need to build various microbial pools, the requirement to survey and control microbial population and the prospects to engineer a transport model of metabolites between two upstream methods in one system, which, on high- scale, present a roadblock.

Conclusions

Overall, the co-culture approach appears interesting from both application and fundamental points of view. We could successfully show that the production of succinic acid from sequential co-culture is possible by a temperature shift. The advantage of this kind of biotechnological process is the extension of the product portfolio of anaerobic fermentation. The work at hand demonstrates that *A. succinogenes* is able to use glycerol as a substrate for succinic acid formation. Moreover, it shows that it is possible to link aerobic fermentation to succinic acid production using sequential co-cultures of *S. cerevisiae* and *A. succinogenes*. Currently, production rates obtained in our study were about 0.3 g $L^{-1} h^{-1}$ from a sequential 48 co-culture, which is promising in view of the production of SA at a large scale, but they still have to be improved to become commercially interesting. Yet, based on the reported co-culture, as

compared with other strategies traditionally used in SA production research, co-culture can be done and give similar results from microbial pure cultures (co-culture strategy gives the same results from a pure culture without the use of a mineral carbon source such as MgCO₃ and the nitrogen sparging). Additionally, this research shows how important microbial interactions and thermodynamic interactions in a microbial-community are, besides the syntrophic systems described in the literature. Evaluation studies of efficiency, environmental impact and cost assessment will also be very interesting. The field still has several major obstacles to go through before stable scale-up studies can be a reality.

Acknowledgments

The authors thank Ivane LELIEVRE (Unilasalle-EME) for her technical help and Marie-Anne Hairan (UniLasalle-EME) for her helpful comments for improving the English. The authors would like to thank Bebnine municipality, North-Lebanon for financial support through the Program Education of Lebanese student.

Ethical Approval

This article followed the ethical standard of the institute.

Consent to Participate

All authors consented to participation in this research.

Consent to publish

All authors consented to publish this research in this journal.

Authors contribution

Alaa SALMA made substantial contributions to the design of the work, performed the experiments, the analysis and wrote the manuscript. Hayet DJELAL, Abdeltif AMRANE, Florence FOURCADE and Rawa ABDALLAH supervised the work, revised the manuscript critically for important intellectual content and approved the version to be published.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Competing interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this article.

Availability of data and materials

The authors declare that the data and materials are transparent.

References

- 1. Taylor, R. (2015). From the Sugar Platform to biofuels and biochemicals, 183.
- Fu, J., Vasiliadou, E. S., Goulas, K. A., Saha, B., & Vlachos, D. G. (2017). Selective hydrodeoxygenation of tartaric acid to succinic acid. *Catalysis Science & Technology*, 7(21), 4944–4954. https://doi.org/10.1039/C7CY01374D
- Amulya, K., & Mohan, S. V. (2019). Fixation of CO2, electron donor and redox microenvironment regulate succinic acid production in Citrobacter amalonaticus. *Science of The Total Environment*, 695, 133838. https://doi.org/10.1016/j.scitotenv.2019.133838
- 4. Tan, J. P., Luthfi, A. A. I., Manaf, S. F. A., Wu, T. Y., & Jahim, J. Md. (2018). Incorporation of CO2 during the production of succinic acid from sustainable oil palm frond juice. *Journal of CO2 Utilization*, 26, 595–601. https://doi.org/10.1016/j.jcou.2018.06.006
- Bradfield, M. F. A., Mohagheghi, A., Salvachúa, D., Smith, H., Black, B. A., Dowe, N., ... Nicol, W. (2015). Continuous succinic acid production by Actinobacillus succinogenes on xylose-enriched hydrolysate. *Biotechnology for Biofuels*, 8. https://doi.org/10.1186/s13068-015-0363-3
- Pinazo, J., Dómine, M., Parvulescu, V., & Filip, P. (2015). Sustainability metrics for succinic acid production: A comparison between biomass-based and petrochemical routes. *Catalysis Today*, 239, 17–24. https://doi.org/10.1016/j.cattod.2014.05.035
- Almqvist, H., Pateraki, C., Alexandri, M., Koutinas, A., & Lidén, G. (2016). Succinic acid production by Actinobacillus succinogenes from batch fermentation of mixed sugars. *Journal of Industrial Microbiology & Biotechnology*, *43*(8), 1117–1130. https://doi.org/10.1007/s10295-016-1787-x

- Ferone, M., Raganati, F., Ercole, A., Olivieri, G., Salatino, P., & Marzocchella, A. (2018). Continuous succinic acid fermentation by Actinobacillus succinogenes in a packed-bed biofilm reactor. *Biotechnology for Biofuels*, *11*. https://doi.org/10.1186/s13068-018-1143-7
- Zou, W., Zhu, L.-W., Li, H.-M., & Tang, Y.-J. (2011). Significance of CO2 donor on the production of succinic acid by Actinobacillus succinogenes ATCC 55618. *Microbial cell factories*, 10, 87. https://doi.org/10.1186/1475-2859-10-87
- Herselman, J. (2017). Rate and yield dependency of Actinobacillus succinogenes on dissolved CO2 concentration. University of Pretoria. Retrieved from https://repository.up.ac.za/handle/2263/61295
- Atasoy, M., Eyice, O., Schnürer, A., & Cetecioglu, Z. (2019). Volatile fatty acids production via mixed culture fermentation: Revealing the link between pH, inoculum type and bacterial composition. *Bioresource Technology*, 292, 121889. https://doi.org/10.1016/j.biortech.2019.121889
- Ferone, M., Raganati, F., Olivieri, G., & Marzocchella, A. (2019). Bioreactors for succinic acid production processes. *Critical Reviews in Biotechnology*, *39*(4), 571–586. https://doi.org/10.1080/07388551.2019.1592105
- 13. Pinkian, N., Phuengjayaem, S., Tanasupawat, S., & Teeradakorn, S. (2018).
 Optimization of succinic acid production by succinic acid bacteria isolated in Thailand, 10.
- Jiang, M., Ma, J., Wu, M., Liu, R., Liang, L., Xin, F., ... Dong, W. (2017). Progress of succinic acid production from renewable resources: Metabolic and fermentative strategies. *Bioresource Technology*, 245, 1710–1717. https://doi.org/10.1016/j.biortech.2017.05.209

- Sadhukhan, S., Villa, R., & Sarkar, U. (2016). Microbial production of succinic acid using crude and purified glycerol from a Crotalaria juncea based biorefinery. *Biotechnology Reports*, *10*, 84–93. https://doi.org/10.1016/j.btre.2016.03.008
- Ahn, J. H., Jang, Y.-S., & Lee, S. Y. (2016). Production of succinic acid by metabolically engineered microorganisms. *Current Opinion in Biotechnology*, 42, 54– 66. https://doi.org/10.1016/j.copbio.2016.02.034
- Sabra, W., & Zeng, A. (2014). Mixed microbial cultures for industrial biotechnology: Success, chance, and challenges.
- Liu, Y.-P., Zheng, P., Sun, Z.-H., Ni, Y., Dong, J.-J., & Wei, P. (2008). Strategies of pH control and glucose-fed batch fermentation for production of succinic acid by Actinobacillus succinogenes CGMCC1593. *Journal of Chemical Technology & Biotechnology*, 83(5), 722–729. https://doi.org/10.1002/jctb.1862
- Kleerebezem, R., & van Loosdrecht, M. C. M. (2007). Mixed culture biotechnology for bioenergy production. *Current Opinion in Biotechnology*, *18*(3), 207–212. https://doi.org/10.1016/j.copbio.2007.05.001
- Bader, J., Brigham, C. J., Stahl, U., & Popović, M. K. (2019). 3 Fermented Beverages Produced by Mixed Cultures, Pure Cultures, and Defined Cocultures. In A. M. Grumezescu & A. M. Holban (Eds.), *Fermented Beverages* (pp. 67–101). Woodhead Publishing. https://doi.org/10.1016/B978-0-12-815271-3.00003-8
- 21. Shen, N., Dai, K., Xia, X.-Y., Zeng, R., & Zhang, F. (2018). Conversion of syngas (CO and H2) to biochemicals by mixed culture fermentation in mesophilic and thermophilic hollow-fiber membrane biofilm reactors. *Journal of Cleaner Production*, 202. https://doi.org/10.1016/j.jclepro.2018.08.162
- 22. Djelal, H., Chniti, S., Jemni, M., Weill, A., Sayed, W., & Amrane, A. (2017).Identification of strain isolated from dates (Phœnix dactylifera L.) for enhancing very

high gravity ethanol production. *Environmental Science and Pollution Research International*, 24(11), 9886–9894. https://doi.org/10.1007/s11356-016-8018-x

- Jones, J. A., & Wang, X. (2018). Use of bacterial co-cultures for the efficient production of chemicals. *Current Opinion in Biotechnology*, *53*, 33–38. https://doi.org/10.1016/j.copbio.2017.11.012
- Marmann, A., Aly, A. H., Lin, W., Wang, B., & Proksch, P. (2014). Co-Cultivation—A Powerful Emerging Tool for Enhancing the Chemical Diversity of Microorganisms. *Marine Drugs*, 12(2), 1043–1065. https://doi.org/10.3390/md12021043
- Liu, H., Hu, H., Jin, Y., Yue, X., Deng, L., Wang, F., & Tan, T. (2017). Co-fermentation of a mixture of glucose and xylose to fumaric acid by Rhizopus arrhizus RH 7-13-9#. *Bioresource Technology*, 233, 30–33. https://doi.org/10.1016/j.biortech.2017.02.035
- Lu, H., Villada, J. C., & Lee, P. K. H. (2019). Modular Metabolic Engineering for Biobased Chemical Production. *Trends in Biotechnology*, *37*(2), 152–166. https://doi.org/10.1016/j.tibtech.2018.07.003
- Gonzales, T. A., Carvalho Silvello, M. A. de, Duarte, E. R., Santos, L. O., Alegre, R. M., & Goldbeck, R. (2020). Optimization of anaerobic fermentation of Actinobacillus succinogenes for increase the succinic acid production. *Biocatalysis and Agricultural Biotechnology*, 27, 101718. https://doi.org/10.1016/j.bcab.2020.101718
- Djelal, H., Amrane, A., Lahrer, F., & Martin, G. (2005). Effect of medium osmolarity on the bioproduction of glycerol and ethanol by Hansenula anomala growing on glucose and ammonium. *Applied Microbiology and Biotechnology*, 69(3), 341–349. https://doi.org/10.1007/s00253-005-1987-1
- Carvalho, M., Roca, C., & Reis, M. A. M. (2016). Improving succinic acid production by Actinobacillus succinogenes from raw industrial carob pods. *Bioresource Technology*, 218, 491–497. https://doi.org/10.1016/j.biortech.2016.06.140

- Walker, G. M., & Stewart, G. G. (2016). Saccharomyces cerevisiae in the Production of Fermented Beverages. *Beverages*, 2(4), 30. https://doi.org/10.3390/beverages2040030
- Tronchoni, J., Curiel, J. A., Morales, P., Torres-Perez, R., & Gonzalez, R. (2017). Early transcriptional response to biotic stress in mixed starter fermentations involving Saccharomyces cerevisiae and Torulaspora delbrueckii. *International Journal of Food Microbiology*, 241, 60–68. https://doi.org/10.1016/j.ijfoodmicro.2016.10.017
- 32. Cao, W., Wang, Y., Luo, J., Yin, J., Xing, J., & Wan, Y. (2018). Effectively converting carbon dioxide into succinic acid under mild pressure with Actinobacillus succinogenes by an integrated fermentation and membrane separation process. *Bioresource Technology*, 266, 26–33. https://doi.org/10.1016/j.biortech.2018.06.016
- Corona-González, R. I., Bories, A., González-Álvarez, V., & Pelayo-Ortiz, C. (2008). Kinetic study of succinic acid production by Actinobacillus succinogenes ZT-130. *Process Biochemistry*, 43(10), 1047–1053. https://doi.org/10.1016/j.procbio.2008.05.011
- 34. Schindler Bryan David. (2011). Understanding and improving respiratory succinate production from glycerol by Actinobacillus succinogenes. Michigan State University.
- Pereira, A. P., Mendes-Ferreira, A., Oliveira, J. M., Estevinho, L. M., & Mendes-Faia, A. (2013). High-cell-density fermentation of Saccharomyces cerevisiae for the optimisation of mead production. *Food Microbiology*, *33*(1), 114–123. https://doi.org/10.1016/j.fm.2012.09.006
- 36. Reis, V. R., Bassi, A. P. G., Cerri, B. C., Almeida, A. R., Carvalho, I. G. B., Bastos, R. G., & Ceccato-Antonini, S. R. (2018). Effects of feedstock and co-culture of Lactobacillus fermentum and wild Saccharomyces cerevisiae strain during fuel ethanol fermentation by the industrial yeast strain PE-2. *AMB Express*, 8(1), 23. https://doi.org/10.1186/s13568-018-0556-9

- McKinlay, J. B., Zeikus, J. G., & Vieille, C. (2005). Insights into Actinobacillus succinogenes fermentative metabolism in a chemically defined growth medium. *Applied and Environmental Microbiology*, *71*(11), 6651–6656. https://doi.org/10.1128/AEM.71.11.6651-6656.2005
- Nag, A., John, P. C. S., Crowley, M. F., & Bomble, Y. J. (2018). Prediction of reaction knockouts to maximize succinate production by Actinobacillus succinogenes. *PLOS ONE*, *13*(1), e0189144. https://doi.org/10.1371/journal.pone.0189144
- Cheng, K.-K., Wang, G.-Y., Zeng, J., & Zhang, J.-A. (2013). Improved Succinate Production by Metabolic Engineering. *BioMed Research International*. Research article. https://doi.org/10.1155/2013/538790
- Bušić, A., Marđetko, N., Kundas, S., Morzak, G., Belskaya, H., Ivančić Šantek, M., ... Šantek, B. (2018). Bioethanol Production from Renewable Raw Materials and Its Separation and Purification: A Review. *Food Technology and Biotechnology*, *56*(3), 289–311. https://doi.org/10.17113/ftb.56.03.18.5546
- 41. Kimberly Chen, Klous C. Cui, & Jinny Choi. (2013). The effect of temperature on the carbon dioxide production of Saccharomyces cerevisiae as measured by the change in volume of carbon dioxide produced. Retrieved from https://ojs.library.ubc.ca/index.php/expedition/article/view/184143
- 42. Kim, N. Y., Kim, S. N., & Kim, O. B. (2018). Long-term adaptation of Escherichia coli to methanogenic co-culture enhanced succinate production from crude glycerol. *Journal of Industrial Microbiology & Biotechnology*, 45(1), 71–76. https://doi.org/10.1007/s10295-017-1994-0
- Charubin, K., Modla, S., Caplan, J. L., & Papoutsakis, E. T. (2020). Interspecies Microbial Fusion and Large-Scale Exchange of Cytoplasmic Proteins and RNA in a Syntrophic Clostridium Coculture. *mBio*, *11*(5). https://doi.org/10.1128/mBio.02030-20

- 44. Diender, M., Uhl, P. S., Bitter, J. H., Stams, A. J. M., & Sousa, D. Z. (2018). High Rate Biomethanation of Carbon Monoxide-Rich Gases via a Thermophilic Synthetic Coculture. *ACS Sustainable Chemistry & Engineering*, 6(2), 2169–2176. https://doi.org/10.1021/acssuschemeng.7b03601
- 45. Richter, H., Molitor, B., Diender, M., Sousa, D. Z., & Angenent, L. T. (2016). A Narrow pH Range Supports Butanol, Hexanol, and Octanol Production from Syngas in a Continuous Co-culture of Clostridium ljungdahlii and Clostridium kluyveri with In-Line Product Extraction. *Frontiers in Microbiology*, *7*. https://doi.org/10.3389/fmicb.2016.01773
- 46. Diender, M., Parera Olm, I., Gelderloos, M., Koehorst, J. J., Schaap, P. J., Stams, A. J. M., & Sousa, D. Z. (2019). Metabolic shift induced by synthetic co-cultivation promotes high yield of chain elongated acids from syngas. *Scientific Reports*, 9(1), 18081. https://doi.org/10.1038/s41598-019-54445-y
- 47. Moscoviz, R., de Fouchécour, F., Santa-Catalina, G., Bernet, N., & Trably, E. (2017).
 Cooperative growth of Geobacter sulfurreducens and Clostridium pasteurianum with subsequent metabolic shift in glycerol fermentation. *Scientific Reports*, *7*, 44334.
 https://doi.org/10.1038/srep44334
- Salimi, F., & Mahadevan, R. (2013). Characterizing metabolic interactions in a clostridial co-culture for consolidated bioprocessing. *BMC biotechnology*, *13*, 95. https://doi.org/10.1186/1472-6750-13-95
- Zhou, K., Qiao, K., Edgar, S., & Stephanopoulos, G. (2015). Distributing a metabolic pathway among a microbial consortium enhances production of natural products. *Nature Biotechnology*, *33*(4), 377–383. https://doi.org/10.1038/nbt.3095

Figure 1 Optical density evolution during fermentations with different modes of inoculation in microaerobic (a) and anaerobic conditions (b) in the presence of 0.2 mol L⁻¹ MgCO₃ except for the sequential culture. Pure culture fermentations of *A. succinogenes* (- \diamond -) and *S. cerevisiae* (- \Box -). Co-cultures fermentations with a % ratio of (0.5;0.5) of *A. succinogenes* and *S. cerevisiae* respectively. (- \diamond -), and co-culture of *S. cerevisiae* with *A. succinogenes* addition after 48 h (- \bullet -). The arrow (\mapsto) represent the inoculation of *A. succinogenes* in the sequential culture at 37°C.

Figure 2 Fermentation kinetics (as sugars consumption and products formation) showing cocultures fermentation ratio in microaerobic conditions in the absence (a) or in the presence (b) of 0.2 mol L⁻¹ of MgCO₃ during fermentation at 37°C for a % volume ratio of (0.5; 0.5), of *A*. *succinogenes* and *S. cerevisiae*, respectively.

Figure 3 Product time-courses during co-cultures in microaerobic and anaerobic conditions without MgCO₃ during fermentation at 37°C; a- for a % volume ratio of (0.25; 0.75), and b- for a % ratio of (0.75; 0.25) of *S. cerevisiae* and *A. succinogenes* respectively. The filled character represents the products obtained from fermentation in microaerobic conditions (- \diamond -) ethanol; (- \bullet -) glycerol (- Δ -) SA and the empty character represent the products obtained from anaerobic conditions (- \diamond -) ethanol (- \Box -) glycerol (- Δ -).

Figure 4 Fermentation parameters in term of sugars consumption during pure cultures (a) and co- cultures (b) at two different temperatures, 28° C and 37° C with a biomass % volume ratio of (0.5; 0.5) for *A. Succinogenes -S. cerevisiae* respectively. (-**-**-) for *A. succinogenes*, (-**+**-) for *S. cerevisiae*, and (-**-**) for co- cultures.

Figure 5 Sequential 48 h cultures sugars consumption and products formations (a) without MgCO₃ with pH and dissolved oxygen variation (b). The arrow (\mapsto) represent the inoculation of *A. succinogenes* with the temperature shift from 28 °C to 37 °C.















Table 1 Influence of the mode of inoculation on the fermentation profile of *A. succinogenes* and *S. cerevisiae* pure culture at the end of fermentation time (48h). Data are duplicates from two independent experiments represented as a mean \pm standard deviation.

Inoculation	T ₀	Pure culture	Pure culture
modality		AS	SC
Products			
(mol L ⁻¹)			
SA	-	0.21	n.d
		±0.01	
Ethanol	-	n.d	0.07 ±0.008
Glycerol	-	n.d	0.07 ±0.015
Productivity	-	0.49	-
$(SA g L^{-1} h^1)$		±0.009	
YSA	-	0.6	n.d
		±0.005	
Y ethanol	-	n.d	0.38
			± 0.007
Y glycerol	-	n.d	0.28
			±0.003

AS: A. Succinogenes, SC: S. cerevisiae, n.d. none detected; Y: yield mol-C products/ mol-C sugars

Table 2. Influence of the mode of inoculation on the fermentation profile at the end of fermentation time (96h). Data are duplicates from two independent experiments represented as a mean \pm standard deviation.

Fermentatio	n		Microa	aerobic		Anaerob	ic
parameters							
Inoculation	t ₀	Pure	Pure	Simulta	Sequenti	Pure	Simulta
modality		culture	culture	neous	al 48 co-	culture	eous
		AS	SC	Co-	culture	AS	Co-
				culture			culture
OD	0.03	0.39	1.54	1.05	1.32	0.36	0.96
		± 0.002	±0.17	±0.06	±0.01	±0.04	±0.04
pН	6.8	6.15	5.04	5.65	5.24	5.2	5.32
	±0.0	±0.08	±0.07	±0.15	±0.08	±0.02	±0.02
	2						
Substrate							
(mol L ⁻¹)							
Glucose	0.2	0.11	0.01	0.04	0.06	0.02	0.02
	±0.00	±0.005	±0.000	±0.01	±0.009	±0.003	±0.001
	5		1				
Fructose	0.2	0.12	0.014	0.02	0.10	0.03	0.02
	±0.00	±0.0001	±0.000	±0.003	±0.001	±0.005	±0.002
	2	6	6				
Products							
(mol L ⁻¹)							

SA	n d	_	_	0.09	0.12	03	0.17
511	n.u			0.07	0.12	0.5	0.17
				+0.007	+0.0006	+0.01	+0.003
				20.007	10.0000	20.01	20.005
Ethanol	n.d	_	0.147	0.07	0.071	_	0.06
20101101	1110		01117	0107	01071		0.00
			± 0.005	± 0.003	± 0.0007		± 0.01
Glycerol	n.d	_	0.025	0.02	0.013	_	0.04
•							
			± 0.0005	± 0.002	± 0.007		±0.001
Productivit	-	-	-	0.11	0.14	0.3	0.20
У				± 0.008	± 0.0008	± 0.005	± 0.004
$(SA g L^{-1})$							
h ¹)							
YSA	-	-	-	0.41	0.34	0.84	0.44
				± 0.007	± 0.002	± 0.01	± 0.003
			0.51	0.41	0.04		0.10
Y ethanol	-	-	0.51	0.41	0.26	-	0.19
			0.000	0.007	0.005		0.01
			± 0.003	± 0.006	± 0.005		± 0.01
17 1 1			0.06	0.00	0.00		0.10
Y glycerol	-	-	0.06	0.08	0.02	-	0.13
Y glycerol	-	-	0.06	0.08	0.02	-	0.13

AS: A. Succinogenes, SC: S. cerevisiae, n.d. none detected; Y: yield mol-C products/ mol-C

sugars

Table 3. Final parameter values (96 h culture time) during single cultures and co-culture of *A*. *succinogenes* and *S. cerevisiae* at two different temperatures. Data are duplicates from two independent experiments represented as a mean \pm standard deviation.

T° C		/	28			3	37	
Inoculat	Pure	Pure	Co-	Sequen	Pure	Pure	Co-	Sequenti
ion	cultur	cultur	culture	tial 48	culture	culture	culture	al 48 h
Modalit	e AS	e SC		h co-	AS	SC		со-
у				culture				culture
OD	0.09	1.2	0.38	0.47	0.13	0.32±0	0.36	0.42
	± 0.00	±0.2	±0.02	±0.03	±0.007	.05	±0.04	±0.03
	2							
pH	5.9	4.3	5.2	5.4	5.7	4.7	5.5	4.8 ±0.4
	±0.2	±0.4	±0.05	±0.02	±0.04	±0.02	±0.09	
Substrate	e (mole I	- ⁻¹)						
Glucose	0.08	0.02	0.06	0.08	0.05	0.07	0.04	0.006
	±0.00	±0.00	±0.004	±0.001	±0.004	±0.02	±0.009	± 0.0001
	5	7						
Fructos	0.13	0.03	0.07	0.13	0.06	0.1	0.06	0.008
e	±0.00	±0.00	±0.006	± 0.000	±0.007	±0.001	±0.004	± 0.0009
	3	5		2				
Product	(mole L ⁻¹	¹)						
Ethanol	n.d	0.1	0.08	0.06	n.d	0.04	0.05	0.06
		±0.01	±0.006	±0.004		± 0.007	±0.01	±0.003

									_
SA	0.07	n.d	0.09	n.d	0.3	n.d	0.22	0.27	
	±0.01		±0.003		±0.02		±0.01	±0.007	
Glycero	n.d	0.03	0.06	0.06	n.d	0.02	0.03	0.005	
1		±0.00	±0.007	±0.001		± 0.009	±0.005	± 0.0008	
		2							

AS: A. Succinogenes, SC: S. cerevisiae, n.d. non detected, pure culture in this experiment were performed in anaerobic fermentation.

Co-culture	mode	substrate	Products	Ref
Careboxydothermus	Continuous	CO, H ₂	H_2/CO_2 , methane	[44]
hydrogenoformans and				
Methanothermobacter				
thermoautotrophicus				
Clostridiumljungdahlii	Fed batch	CO/	Acetate/ ethanol,	[45]
and Clostridium kluyveri		H_2/CO_2	C4-C8 fatty acid	
			and alcohols	
Clostridium autoethanogenum	Continuous	CO/	Acetate/ ethanol,	[46]
and Clostridium kluyveri	chemostat	H ₂ /CO ₂	C4-C8 fatty acid	
			and alcohols	
A. succinogenes and	Batch	Glucose,	Succinate,	This
S. cerevisiae		Fructose	ethanol, glycerol	study
Escherichia coli	Batch	Glycerol	succinate	[42]
and Methanobacterium				
formicicum				
Geobacter sulfurreducens and	Batch	Glycerol,	propanediol,	[47]
Clostridium pasteurianum		acetate	butyrate	
Clostridium	Batch	Glucose	2,3-butanediol,	[43]
ljungdahlii and Clostridium			isopropanol	
acetobutylicum				
Clostridium acetobutylicum and	Batch	Cellulose	Pyruvate,	[48]
Clostridium cellulolyticum			butanol	
Escherichia	Fed batch	Xylose	oxygenated	[49]
coli and Saccharomyces			taxanes,	
cerevisiae (GMO)			sesquiterpenes,	
			acetate	

Table 4. A selection of microbial co-cultures, their mode cultures, substrates and their products. GMO genetically engineered organism.