



**HAL**  
open science

## **Straightforward synthetic route to gold(I)-thiolato glycoconjugate complexes bearing NHC ligands (NHC = N-heterocyclic carbene) and their promising anticancer activity**

Mauro Safir Filho, Thomas Scattolin, Pascal Dao, Nikolaos V Tzouras, Rachid Benhida, Marina Saab, Kristof van Hecke, Petra Lippmann, Anthony R. Martin, Ingo Ott, et al.

### ► **To cite this version:**

Mauro Safir Filho, Thomas Scattolin, Pascal Dao, Nikolaos V Tzouras, Rachid Benhida, et al.. Straightforward synthetic route to gold(I)-thiolato glycoconjugate complexes bearing NHC ligands (NHC = N-heterocyclic carbene) and their promising anticancer activity. *New Journal of Chemistry*, 2021, 45 (22), pp.9995-10001. 10.1039/D1NJ02117F . hal-03225368

**HAL Id: hal-03225368**

**<https://hal.science/hal-03225368v1>**

Submitted on 12 May 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## ARTICLE

# Straightforward synthetic route to gold(I)-thiolato glycoconjugate complexes bearing NHC ligands (NHC = N-heterocyclic carbene) and their promising anticancer activity

Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Mauro Safir Filho,<sup>a</sup> Thomas Scattolin,<sup>b</sup> Pascal Dao,<sup>a</sup> Nikolaos V. Tzouras,<sup>b</sup> Rachid Benhida,<sup>a,c</sup> Marina Saab,<sup>b</sup> Kristof Van Hecke,<sup>b</sup> Petra Lippmann,<sup>d</sup> Anthony R. Martin,<sup>\*a,†</sup> Ingo Ott<sup>\*d</sup> and Steven P. Nolan<sup>\*b</sup>

A simple and eco-friendly route to gold-NHC complexes bearing different thiosugars is reported. The reaction between [Au(NHC)Cl] precursors, the thiosugars and a weak base proceeds under air using technical grade acetone under mild conditions (60 °C, 1h). The one-pot synthesis of the same complexes starting from the corresponding imidazolium salts was also investigated. The obtained complexes showed strong cytotoxic activities against selected cancer cell lines with IC<sub>50</sub> values in the submicromolar to low micromolar concentration range.

## Introduction

Since the serendipitous discovery of cisplatin, many researchers in the field of coordination and organometallic chemistry of middle and late-transition metals have contributed to the development of new promising anticancer agents.<sup>1</sup> Although the research conducted on platinum complexes has led to the commercialization of the well-known second- and third-generation platinum-based antineoplastic drugs (*e.g.* carboplatin and oxaliplatin),<sup>2</sup> their ineffectiveness against different types of tumours or the development of drug resistance observed in many patients<sup>3</sup> has prompted numerous scientists to explore the anticancer properties of complexes containing metals other than platinum.

In this context, the remarkable *in-vitro*, *ex-vivo* and *in-vivo* anticancer activity exhibited by some categories of mono- and poly-nuclear gold,<sup>4</sup> ruthenium<sup>5</sup> and palladium<sup>6</sup> complexes has proven particularly encouraging. Many of these compounds are considered attractive as they display a completely different mechanism of action compared to classical platinated agents and therefore are often effective toward cisplatin-resistant cancer cells.<sup>7</sup>

Among the various metal derivatives mentioned above, a place of absolute importance in medicinal chemistry is covered

by gold(I) complexes, which are currently used as therapeutic drugs for the treatment of rheumatoid arthritis.<sup>8</sup> Results addressing their activity as antibiotics and chemotherapeutic agents are so encouraging that a number of most promising compounds are currently under clinical trials.<sup>1,4</sup> In-depth studies carried out in recent decades have shown that anticancer gold complexes have a wide range of cellular targets such as mitochondria and proteins playing relevant functional roles (*e.g.* thioredoxin reductase, poly (ADP-ribose) polymerase (PARP-1), cathepsins, proteasome and protein tyrosine phosphatase).<sup>4</sup>

In this respect, the pioneering studies carried out by Mirabelli *et al.* on the 1-thio-β-D-glucopyranose-2,3,4,6-tetraacetato-S)(triethylphosphine) gold(I) complex, which is commonly known as auranofin, has played a crucial role in the explosion of gold anticancer agents.<sup>9</sup> This compound, which was introduced on the market in the 1980s as an anti-arthritis drug, has recently received approval by FDA for clinical phase II for cancer therapy.<sup>10</sup>

Although the mechanism of action of auranofin is still being debated, it appears to involve the inhibition of thioredoxin reductase (Trx), a key protein overexpressed by tumor cells, in which the metal center selectively and covalently bind the selenium atom present in this category of selenocysteine enzymes.<sup>11</sup> From a composition perspective, auranofin presents triethylphosphine as an ancillary ligand and the thio-β-D-glucopyranose-2,3,4,6-tetraacetate actor ligand. The presence of a glycosidic unit is an important factor in the design of new chemotherapeutic agents since the ability of carbohydrates to act as active sites for molecular recognition, their high water-solubility and above all the more efficient uptake in tumor cells with respect to the healthy ones (Warburg's effect) are well-known characteristics of these moieties.<sup>12</sup>

<sup>a</sup> Institut de Chimie de Nice, Université Côte d'Azur, CNRS, UMR7272, Nice, France. E-mail: [anthony.martin@umontpellier.fr](mailto:anthony.martin@umontpellier.fr); ‡ Present address : Institut des Biomolécules Max Mousseron, UMR5247, Université de Montpellier, ENSCM, CNRS, Montpellier, France.

<sup>b</sup> Department of Chemistry and Center for Sustainable Chemistry, Ghent University, Krijgslaan 281 (S-3), 9000, Ghent, Belgium. E-mail: [steven.nolan@ugent.be](mailto:steven.nolan@ugent.be)


<sup>c</sup> Chemical & Biochemical Sciences Green-Process Engineering (CBS-GPE) Mohammed VI Polytechnic University, Lot 660, Hay Moulay Rachid, Benguerir, Morocco.

<sup>d</sup> Institute of Medicinal and Pharmaceutical Chemistry, Technische Universität Braunschweig, Beethovenstr. 55, 38106, Braunschweig, Germany. E-mail: [ingo.ott@tu-bs.de](mailto:ingo.ott@tu-bs.de)

† Electronic Supplementary Information (ESI) available: Synthetic procedure, characterisation and biological data. For ESI See DOI: 10.1039/x0xx00000x




reagents and the disappearance of the indicative signal attributable to the S-H group at 2.28–2.35 ppm. Furthermore, for complex **8c**, X-ray diffraction study of single crystals grown from tetrahydrofuran and pentane, permit the unambiguous atom connectivity to be established.



**Figure 1.** X-ray molecular structure of **8c**, showing thermal displacement ellipsoids at the 50% probability level with hydrogen and solvent atoms omitted for clarity. Selected bond lengths (Å) and angles (°): (C<sub>1</sub>-Au<sub>1</sub> = 2.024(6), Au<sub>1</sub>-S<sub>1</sub> = 2.2826(16), C<sub>1</sub>-Au<sub>1</sub>-S<sub>1</sub> = 174.44(16)). CCDC: 2072449

With the aim of further simplifying the synthetic procedure to gold-NHC thiolates, we tested the direct conversion of the azolium salts (NHC-HCl), which are the classical precursors of NHC ligands, into the desired complexes. Gratifyingly, by first reacting the azolium salt with the [AuCl(DMS)] precursor in the presence of K<sub>2</sub>CO<sub>3</sub> (1 equiv.) for 1h in acetone at 60 °C and subsequently adding 1-thio- $\alpha$ -D-mannose tetraacetate (**c**) as a representative thiosugar and an additional equivalent of base, it was possible to obtain the final complexes in yields comparable to the two-step process (Scheme 2).



**Scheme 2.** One-pot synthesis of gold-NHC thiolato complexes

### Cytotoxicity

Three cancer cell lines representing highly relevant human malignancies, namely A549 lung carcinoma, HT-29 colon adenocarcinoma, and MDA-MB-231 breast carcinoma, were selected to evaluate the anticancer potential of complexes **5-8**. All complexes showed strong antiproliferative effects with IC<sub>50</sub> values in the sub-micromolar to low micromolar range (see Table 1). With IC<sub>50</sub> values in the range of 0.7 to 2.4  $\mu$ M, MDA-

MB-231 cells were most sensitive towards the compounds (IC<sub>50</sub> values in the range of 1.3 to 3.9  $\mu$ M for HT-29 cells and 2.3 to 6.7  $\mu$ M for A-549 cells). The experimental assay conditions required a 96 h incubation period with MDA-MB-231 cells, while for A-549 and HT-29 cells the exposure period was 72 h. The results therefore indicate that the biological activity of the compounds increases over time.

Comparing the results obtained with the individual complexes, a slight preference for glucose as thiosugar ligand was noted. Comparison with results obtained for other gold(I)-NHC complexes in previous reports using similar experimental conditions,<sup>4g,20</sup> confirms the overall improvement of cytotoxic activity exhibited by complexes **5-8**.

**Table 1** Cytotoxic activity of NHC-Au-thiolato glycoconjugate complexes **5-8**

| Complex   | IC <sub>50</sub> ( $\mu$ M) <sup>a</sup> |               |               |
|-----------|--|---------------|---------------|
|           | A-549                                    | HT-29         | MDA-MB-231    |
| <b>5a</b> | 3.0 $\pm$ 0.2                            | 1.6 $\pm$ 0.1 | 0.8 $\pm$ 0.1 |
| <b>5b</b> | 3.7 $\pm$ 0.5                            | 1.3 $\pm$ 0.2 | 0.8 $\pm$ 0.2 |
| <b>5c</b> | 4.5 $\pm$ 0.9                            | 2.3 $\pm$ 0.6 | 1.5 $\pm$ 0.0 |
| <b>6a</b> | 2.2 $\pm$ 0.7                            | 1.0 $\pm$ 0.3 | 0.7 $\pm$ 0.2 |
| <b>6b</b> | 2.9 $\pm$ 0.9                            | 1.9 $\pm$ 0.7 | 1.2 $\pm$ 0.6 |
| <b>6c</b> | 3.4 $\pm$ 0.7                            | 1.8 $\pm$ 0.5 | 1.3 $\pm$ 0.5 |
| <b>7a</b> | 2.3 $\pm$ 0.8                            | 1.2 $\pm$ 0.6 | 1.0 $\pm$ 0.4 |
| <b>7b</b> | 3.7 $\pm$ 0.4                            | 2.0 $\pm$ 0.3 | 1.5 $\pm$ 0.3 |
| <b>7c</b> | 3.4 $\pm$ 0.9                            | 1.6 $\pm$ 0.5 | 1.3 $\pm$ 0.6 |
| <b>8a</b> | 5.8 $\pm$ 0.3                            | 3.4 $\pm$ 0.1 | 2.2 $\pm$ 0.5 |
| <b>8b</b> | 6.7 $\pm$ 0.5                            | 3.9 $\pm$ 0.4 | 2.4 $\pm$ 0.5 |
| <b>8c</b> | 5.7 $\pm$ 1.0                            | 3.6 $\pm$ 0.3 | 2.1 $\pm$ 0.2 |

IC<sub>50</sub> values were obtained from 3 independent experiments. Experimental errors were calculated as standard deviations.

It should be remembered that in a previous study conducted by Riches and Nolan on a panel of cancer cell lines (LNCaP and P21PZ prostate carcinoma, MDA MB231 and B42-Cl16 breast carcinoma, MGH-U1 bladder carcinoma) and normal cells (SUV-HUC-1 and P21TZ), a good cytotoxicity of both **5a** and its chloride precursor **1a** was already observed in all examined cancer cell lines.<sup>16c</sup> More in detail, **5a** was generally more active than **1a** and Au(I) neutral derivatives bearing other biocompatible units.

Although **5a** and **1a** have proved to be cytotoxic also towards normal cells, it is widely recognized that an *in vivo* study is essential to clarify the real toxicity of the compounds and their effectiveness in reducing tumor mass.

Trx inhibition and the mechanism of action of an NHC-Au-thioglyucose complex (NHC = 1,3-dibenzyl-4,5-diphenylimidazolylidene)<sup>19g-h</sup> and other non-glycosidic NHC-Au-thiolates<sup>20</sup> have recently been explored by some research groups.

These studies examined the ability of gold-thiolate complexes to inhibit the mammalian thioredoxin reductase enzyme, showing IC<sub>50</sub> values in the low micromolar range and thus indicating that apoptosis induction through elevated oxidative stress could be their main mechanism of action.

## Conclusions

In summary, we have reported a simple and eco-friendly procedure for the synthesis of 12 gold-NHC thiolate complexes (**5-8**). In particular, these derivatives can be obtained in high yields and purity by reaction of a [Au(NHC)Cl] precursor with a thiosugar in the presence of a weak base such as K<sub>2</sub>CO<sub>3</sub>. All reactions proceed, under air, in technical grade acetone under mild conditions (60 °C, 1h). Interestingly, we have shown that these compounds can be obtained directly from the corresponding azolium salts in a one-pot procedure.

Complexes **5-8** showed strong cytotoxic activities against a panel of selected highly relevant cancers from different tissues (lung, colon, breast). Further evaluation of the anticancer potential of this class of gold organometallics is warranted and is the subject of ongoing studies in our laboratories.

In particular, based on the promising *in vivo* results obtained by Zhu and Tacke with an NHC-Au-thioglucose complex (NHC = 1,3-dibenzyl-4,5-diphenyl-imidazolylidene)<sup>19g-h</sup> we believe that a future work dealing with the *in vivo* efficacy of the most active compounds on xenograft models could give valuable information on the real potential of these gold(I) derivatives.

## Experimental

### Materials and methods

All complex syntheses were performed in air. Solvents and all other reagents were purchased and used as received without further purification unless otherwise stated. 1-thio-β-D-galactose tetraacetate and 1-thio-α-D-mannose tetraacetate were synthesized according to the procedure described by Doyle L. M. et al.<sup>21</sup> <sup>1</sup>H and <sup>13</sup>C{<sup>1</sup>H} Nuclear Magnetic Resonance (NMR) spectra were recorded on a Bruker Advance 400 Ultrashield spectrometer at 298K. Chemical shifts (expressed by parts per million) are referenced to residual solvent peaks.

Elemental analysis was carried out on a Flash EA 1112 series Thermo Finnigan elementary analyzer piloted by Eager Xperience software. The CHN oven was set up at 970°C, and the analysis were performed using helium (140 mL/min) as the vector gas and a catarometer as detector.

X-ray intensity data were collected at 100 K, on a Rigaku Oxford Diffraction Supernova Dual Source diffractometer equipped with an Atlas CCD detector.

### General procedure for complexes 5-8

The precursor [Au(NHC)Cl] **1-4** (100 mg), the thiosugar **a-c** (1 equiv.) and potassium carbonate (1 equiv.) were dissolved in technical grade acetone (0.05 M) and heated at 60°C for 1 hour. Then, the solvent was evaporated under vacuum, the crude product was dissolved in CH<sub>2</sub>Cl<sub>2</sub> and filtered through a pad of silica to remove the inorganic salts. The filtrate was concentrated in vacuo and triturated with pentane to afford the desired complexes as white powders.

### One-pot procedure for complexes 5-8

The NHC-HCl salt (100 mg), chloro(dimethylsulfide)gold(I) ([Au(DMS)Cl], 1 equiv.) and potassium carbonate (1 equiv.) were dissolved in technical grade acetone (0.05 M) and heated at 60°C for 1 hour. Afterwards, the thiosugar **a-c** (1 equiv.) and another portion of potassium carbonate (1 equiv.) were added, and the reaction was heated at 60°C for 1 additional hour.

The solvent was evaporated under vacuum, the crude product was dissolved in CH<sub>2</sub>Cl<sub>2</sub> and filtered through a pad of silica to remove the inorganic salts. The filtrate was concentrated in vacuo and triturated with pentane to afford the respective complexes as white powders.

### Synthesis of [Au(IPr)(1-thio-β-D-glucose tetraacetate)] (**5a**)

Yield general procedure: 89%

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.54 (t, *J* = 7.8 Hz, 1H), 7.34 – 7.31 (m, 4H), 7.15 (s, 2H), 4.93 (t, *J* = 9.3 Hz, 1H), 4.69 (t, *J* = 9.8 Hz, 1H), 4.55 (d, *J* = 9.6 Hz, 1H), 4.42 (t, *J* = 9.4 Hz, 1H), 3.93 (dd, *J* = 11.9, 5.5 Hz, 1H), 3.83 (dd, *J* = 12.0, 2.9 Hz, 1H), 3.37 (ddd, *J* = 10.0, 5.5, 2.9 Hz, 1H), 2.64 – 2.56 (m, 4H), 2.04 (s, 3H), 2.01 (s, 3H), 1.95 (s, 3H), 1.87 (s, 3H), 1.36 (dd, *J* = 6.7, 5.4 Hz, 12H), 1.22 (dd, *J* = 6.8, 5.5 Hz, 12H).

<sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 186.9, 171.1, 170.5, 169.7, 169.3, 146.0, 145.9, 134.3, 130.8, 124.4, 124.3, 122.9, 82.7, 77.4, 75.2, 74.4, 69.7, 63.8, 29.0, 28.9, 24.6, 24.5, 24.3, 24.2, 21.1, 21.0, 20.9, 20.9.

Elemental Analysis for C<sub>41</sub>H<sub>55</sub>AuN<sub>2</sub>O<sub>9</sub>S (%): found C 51.74, H 5.65, N 2.95; calculated: C 51.90, H 5.84, N 2.95.

The data are in agreement with the reported values.<sup>16b-c</sup>

### Synthesis of [Au(IPr)(1-thio-β-D-galactose tetraacetate)] (**5b**)

Yield general procedure: 91%

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.49 (t, *J* = 7.8 Hz, 2H), 7.29 (d, *J* = 7.8 Hz, 4H), 7.16 (s, 2H), 5.27 (d, *J* = 2.6 Hz, 1H), 4.93 (t, *J* = 9.8 Hz, 1H), 4.75 (dd, *J* = 10.1, 3.4 Hz, 1H), 4.36 (d, *J* = 9.5 Hz, 1H), 3.92 (d, *J* = 7.2 Hz, 1H), 3.48 (t, *J* = 7.5 Hz, 1H), 2.61 – 2.54 (m, 4H), 2.06 (s, 3H), 2.04 (s, 3H), 1.92 (s, 3H), 1.87 (s, 3H), 1.34 (d, *J* = 6.8 Hz, 12H), 1.22 (dd, *J* = 6.9, 1.9 Hz, 12H).

<sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 186.5, 170.7, 170.6, 170.5, 169.7, 145.9, 145.9, 134.3, 130.6, 124.3, 124.3, 122.9, 83.7, 73.5, 72.7, 72.5, 67.5, 61.1, 28.9, 24.6, 24.1, 24.1, 21.2, 20.9, 20.8.

Elemental Analysis for C<sub>41</sub>H<sub>55</sub>AuN<sub>2</sub>O<sub>9</sub>S (%): found C 51.92, H 5.80, N 2.96; calculated: C 51.90, H 5.84, N 2.95.

### Synthesis of [Au(IPr)(1-thio-α-D-mannose tetraacetate)] (**5c**)

Yield general procedure: 83%; Yield one-pot procedure: 81%

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.48 (t, *J* = 7.8 Hz, 2H), 7.30 – 7.26 (m, 4H), 7.15 (s, 2H), 5.69 (dd, *J* = 10.0, 3.4 Hz, 1H), 5.24 (d, *J* = 1.5 Hz, 1H), 5.17 (t, *J* = 10.1 Hz, 1H), 5.12 (dd, *J* = 3.4, 1.7 Hz, 1H), 4.48 (dt, *J* = 10.1, 2.6 Hz, 1H), 4.17 (dd, *J* = 12.4, 3.2 Hz, 1H), 3.54 (dd, *J* = 12.3, 2.2 Hz, 1H), 2.58 – 2.51 (m, 4H), 2.08 (s, 6H), 1.95 (s, 3H), 1.88 (s, 3H), 1.32 (dd, *J* = 7.6, 7.2 Hz, 12H), 1.21 (d, *J* = 6.9 Hz, 12H).

<sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 186.4, 171.2, 170.0, 170.0, 169.7, 145.7, 145.7, 134.1, 130.7, 124.3, 124.3, 123.0, 80.7, 75.6, 69.7, 67.2, 66.9, 62.8, 28.9, 24.7, 24.1, 24.1, 21.3, 21.0, 20.9, 20.8.

Elemental Analysis for  $C_{41}H_{55}AuN_2O_9S$  (%): found C 51.94, H 5.83, N 3.00; calculated: C 51.90, H 5.84, N 2.95.

#### Synthesis of [Au(IMes)(1-thio- $\beta$ -D-glucose tetraacetate)] (6a)

Yield general procedure: 71%.

$^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  7.07 (s, 2H), 7.04 (d,  $J$  = 7.1 Hz, 4H), 4.97 (t,  $J$  = 9.3 Hz, 1H), 4.76 (t,  $J$  = 9.7 Hz, 1H), 4.64 (d,  $J$  = 9.5 Hz, 1H), 4.43 (t,  $J$  = 9.4 Hz, 1H), 3.92 (d,  $J$  = 4.2 Hz, 1H), 3.46 (dt,  $J$  = 10.0, 4.2 Hz, 1H), 2.36 (s, 6H), 2.15 (d,  $J$  = 6.7 Hz, 12H), 2.05 (s, 3H), 2.02 (s, 3H), 2.00 (s, 3H), 1.96 (s, 3H).

$^{13}C$  NMR (101 MHz,  $CDCl_3$ )  $\delta$  185.1, 171.1, 170.5, 169.8, 169.4, 139.7, 135.2, 135.1, 134.9, 129.5, 129.4, 121.9, 82.7, 77.3, 75.6, 74.5, 69.7, 63.8, 21.3, 21.2, 21.1, 20.9, 20.9, 17.9, 17.9.

Elemental Analysis for  $C_{37}H_{51}AuN_2O_9S$  (%): found C 49.14, H 4.93, N 3.17; calculated: C 48.61, H 5.01, N 3.24.

#### Synthesis of [Au(IMes)(1-thio- $\beta$ -D-galactose tetraacetate)] (6b)

Yield general procedure: 89%.

$^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  7.08 (s, 2H), 7.00 (d,  $J$  = 5.8 Hz, 4H), 5.31 (d,  $J$  = 2.8 Hz, 1H), 4.99 (t,  $J$  = 9.8 Hz, 1H), 4.82 (dd,  $J$  = 10.1, 3.3 Hz, 1H), 4.48 (d,  $J$  = 9.5 Hz, 1H), 4.07 – 3.94 (m, 2H), 3.56 (dd,  $J$  = 8.7, 5.7 Hz, 1H), 2.34 (s, 6H), 2.12 (s, 12H), 2.08 (s, 3H), 2.03 (s, 3H), 1.94 (s, 3H), 1.94 (s, 3H).

$^{13}C$  NMR (101 MHz,  $CDCl_3$ )  $\delta$  184.7, 170.7, 170.6, 170.5, 169.6, 139.7, 135.0, 134.9, 129.5, 129.4, 122.0, 83.1, 73.8, 73.0, 72.5, 67.6, 61.0, 21.2, 21.2, 20.9, 20.8, 17.9, 17.9.

Elemental Analysis for  $C_{35}H_{43}AuN_2O_9S$  (%): found C 48.28, H 4.90, N 3.16; calculated: C 48.61, H 5.01, N 3.24.

#### Synthesis of [Au(IMes)(1-thio- $\alpha$ -D-mannose tetraacetate)] (6c)

Yield general procedure: 92%; Yield one-pot procedure: 89%

$^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  7.06 (s, 2H), 6.99 (s, 4H), 5.76 (dd,  $J$  = 9.9, 3.4 Hz, 1H), 5.37 (d,  $J$  = 1.5 Hz, 1H), 5.20 (t,  $J$  = 10.0 Hz, 1H), 5.14 (dd,  $J$  = 3.4, 1.7 Hz, 1H), 4.52 (dt,  $J$  = 10.1, 2.7 Hz, 1H), 4.24 (dd,  $J$  = 12.3, 3.4 Hz, 1H), 3.67 (dd,  $J$  = 12.3, 2.1 Hz, 1H), 2.34 (s, 6H), 2.11 – 2.10 (m, 15H), 2.08 (s, 3H), 1.97 (s, 3H), 1.92 (s, 3H).

$^{13}C$  NMR (101 MHz,  $CDCl_3$ )  $\delta$  184.6, 171.1, 170.0, 169.9, 169.7, 139.6, 134.9, 129.6, 129.5, 122.1, 80.5, 75.8, 69.5, 67.4, 67.0, 62.7, 21.3, 21.0, 20.9, 20.9, 17.9, 17.9.

Elemental Analysis for  $C_{35}H_{43}AuN_2O_9S$  (%): found C 48.27, H 4.85, N 3.18; calculated: C 48.61, H 5.01, N 3.24.

#### Synthesis of [Au(SIMes)(1-thio- $\beta$ -D-glucose tetraacetate)] (7a)

Yield general procedure: 95%.

$^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  7.00 (d,  $J$  = 3.8 Hz, 4H), 4.93 (t,  $J$  = 9.3 Hz, 1H), 4.72 (t,  $J$  = 9.7 Hz, 1H), 4.54 (d,  $J$  = 9.5 Hz, 1H), 4.35 (t,  $J$  = 9.4 Hz, 1H), 3.96 (s, 4H), 3.91 (d,  $J$  = 4.2 Hz, 2H), 3.39 (dt,  $J$  = 10.0, 4.2 Hz, 1H), 2.36 (d,  $J$  = 6.8 Hz, 12H), 2.31 (s, 6H), 2.06 (s, 3H), 2.02 (s, 3H), 2.00 (s, 3H), 1.97 (s, 3H).

$^{13}C$  NMR (101 MHz,  $CDCl_3$ )  $\delta$  205.1, 171.1, 170.5, 169.7, 169.4, 138.9, 136.1, 135.9, 134.9, 129.8, 129.7, 82.6, 77.1, 75.5, 74.4, 69.6, 63.8, 50.9, 21.3, 21.1, 21.1, 20.9, 20.8, 18.2, 18.1.

Elemental Analysis for  $C_{35}H_{45}AuN_2O_9S$  (%): found C 48.41, H 5.06, N 3.15; calculated: C 48.50, H 5.23, N 3.23.

#### Synthesis of [Au(SIMes)(1-thio- $\beta$ -D-galactose tetraacetate)] (7b)

Yield general procedure: 91%.

$^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  6.95 (d,  $J$  = 8.2 Hz, 4H), 5.29 (d,  $J$  = 2.4 Hz, 1H), 4.95 (t,  $J$  = 9.8 Hz, 1H), 4.77 (dd,  $J$  = 10.1, 3.4 Hz, 1H), 4.31 (d,  $J$  = 9.5 Hz, 1H), 4.06 – 3.93 (m, 6H), 3.45 (dd,  $J$  = 8.3, 5.8 Hz, 1H), 2.33 – 2.29 (m, 18H), 2.08 (s, 3H), 2.04 (s, 3H), 1.96 (s, 3H), 1.93 (s, 3H).

$^{13}C$  NMR (101 MHz,  $CDCl_3$ )  $\delta$  204.8, 170.7, 170.6, 170.4, 169.6, 138.8, 135.9, 135.9, 134.9, 129.7, 129.6, 83.0, 73.8, 72.8, 72.5, 67.5, 61.0, 50.9, 21.3, 21.1, 20.9, 20.8, 18.2, 18.1.

Elemental Analysis for  $C_{35}H_{45}AuN_2O_9S$  (%): found C 48.28, H 5.10, N 3.25; calculated: C 48.50, H 5.23, N 3.23.

#### Synthesis of [Au(SIMes)(1-thio- $\alpha$ -D-mannose tetraacetate)] (7c)

Yield general procedure: 88%; Yield one-pot procedure: 82%

$^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  6.93 (d,  $J$  = 4.0 Hz, 4H), 5.73 (dd,  $J$  = 9.9, 3.4 Hz, 1H), 5.24 (d,  $J$  = 1.6 Hz, 1H), 5.17 (t,  $J$  = 10.0 Hz, 1H), 5.06 (dd,  $J$  = 3.4, 1.7 Hz, 1H), 4.53 – 4.41 (m, 1H), 4.22 (dd,  $J$  = 12.3, 3.6 Hz, 1H), 3.95 (s, 4H), 3.68 (dd,  $J$  = 12.3, 2.2 Hz, 1H), 2.31 (d,  $J$  = 5.0 Hz, 12H), 2.28 (s, 6H), 2.09 (s, 3H), 2.08 (s, 3H), 1.96 (s, 3H), 1.92 (s, 3H).

$^{13}C$  NMR (101 MHz,  $CDCl_3$ )  $\delta$  204.8, 171.1, 170.0, 169.9, 169.7, 138.8, 135.7, 135.7, 134.8, 129.8, 129.7, 80.2, 75.7, 69.5, 67.5, 67.0, 62.7, 50.9, 21.2, 21.2, 20.9, 20.9, 18.1, 18.0.

Elemental Analysis for  $C_{35}H_{45}AuN_2O_9S$  (%): found C 48.45, H 5.19, N 3.21; calculated: C 48.50, H 5.23, N 3.23.

#### Synthesis of [Au(IAd)(1-thio- $\beta$ -D-glucose tetraacetate)] (8a)

Yield general procedure: 85%.

$^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  7.06 (s, 2H), 5.17 – 5.12 (m, 2H), 5.02 – 5.01 (m, 2H), 4.23 (dd,  $J$  = 12.2, 5.0 Hz, 1H), 4.06 (dd,  $J$  = 12.1, 2.4 Hz, 1H), 3.70 (ddd,  $J$  = 9.4, 4.9, 2.4 Hz, 1H), 2.55 (d,  $J$  = 2.1 Hz, 12H), 2.27 (s, 6H), 2.07 (s, 3H), 2.03 (s, 3H), 2.00 (s, 3H), 1.97 (s, 3H), 1.77 (s, 12H).

$^{13}C$  NMR (101 MHz,  $CDCl_3$ )  $\delta$  179.8, 171.1, 170.6, 169.8, 169.7, 115.3, 83.5, 76.9, 75.8, 74.6, 69.3, 63.3, 59.2, 44.4, 36.0, 30.1, 21.2, 20.9, 20.9, 20.8.

Elemental Analysis for  $C_{41}H_{55}AuN_2O_9S$  (%): found C 49.03, H 5.60, N 3.02; calculated: C 49.55, H 5.73, N 3.12.

#### Synthesis of [Au(IAd)(1-thio- $\beta$ -D-galactose tetraacetate)] (8b)

Yield general procedure: 68%.

$^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  7.06 (s, 2H), 5.40 (d,  $J$  = 2.7 Hz, 1H), 5.21 (t,  $J$  = 9.8 Hz, 1H), 5.03 (d,  $J$  = 9.4 Hz, 1H), 4.99 (dd,  $J$  = 10.1, 3.5 Hz, 1H), 4.18 – 4.08 (m, 2H), 3.91 – 3.87 (m, 1H), 2.55 (d,  $J$  = 2.1 Hz, 12H), 2.26 (s, 6H), 2.11 (s, 3H), 2.08 (s, 3H), 2.00 (s, 3H), 1.96 (s, 3H), 1.76 (s, 12H).

$^{13}C$  NMR (101 MHz,  $CDCl_3$ )  $\delta$  179.8, 170.7, 170.6, 170.5, 169.9, 115.2, 84.4, 74.2, 73.9, 72.6, 67.9, 61.4, 5.2, 44.3, 36.0, 30.1, 21.3, 20.9, 20.9.

Elemental Analysis for  $C_{37}H_{51}AuN_2O_9S$  (%): found C 49.49, H 5.70, N 3.04; calculated: C 49.55, H 5.73, N 3.12.

#### Synthesis of [Au(IAd)(1-thio- $\alpha$ -D-mannose tetraacetate)] (8c)

Yield general procedure: 75%; Yield one-pot procedure: 92%  
 $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.05 (s, 2H), 6.02 (dd,  $J = 9.9, 3.4$  Hz, 1H), 5.75 (d,  $J = 1.4$  Hz, 1H), 5.50 (dd,  $J = 3.3, 1.7$  Hz, 1H), 5.30 (t,  $J = 10.0$  Hz, 1H), 4.83 (ddd,  $J = 10.1, 3.8, 2.5$  Hz, 1H), 4.32 (dd,  $J = 12.2, 4.1$  Hz, 1H), 4.03 (dd,  $J = 12.2, 2.3$  Hz, 1H), 2.56–2.49 (m, 12H), 2.24 (br s, 6H), 2.13 (s, 3H), 2.05 (s, 3H), 2.02 (s, 3H), 1.96 (s, 3H), 1.74 (br s, 12H).

$^{13}\text{C}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  179.0, 171.1, 170.3, 170.2, 169.9, 115.3, 81.5, 75.9, 69.7, 67.8, 67.5, 63.2, 59.2, 44.4, 36.0, 30.1, 21.3, 21.0, 20.9, 20.9.

Elemental Analysis for  $\text{C}_{37}\text{H}_{51}\text{AuN}_2\text{O}_9\text{S}$  (%): found C 49.82, H 5.80, N 3.12; calculated: C 49.55, H 5.73, N 3.12.

### Cell culture and cytotoxicity

A-549, HT-29 and MDA-MB-231 cells were obtained from DSMZ (German Collection of Microorganisms and Cell Cultures GmbH) and maintained by routine procedures. The cytotoxicity of complexes **5–8** was determined according to previously applied procedures.<sup>4f</sup> In short: The test compounds were prepared as stock solutions in dimethylformamide and diluted with cell culture media to graded concentrations. The cells were exposed to the compound containing cell culture media for a period of 72 h (A-549, HT-29) or 96 h (MDA-MB-231), after which the cell biomass was determined by crystal violet staining. The  $\text{IC}_{50}$  values were calculated as the concentrations required to reduce cell proliferation by 50 % compared to the untreated control.

### Conflicts of interest

There are no conflicts to declare.

### Acknowledgements

CNRS, Université Côte d'Azur (UCA, Nice) and UM6P are gratefully acknowledged for financial support. For work conducted in Ghent, we are grateful to the Moonshot D2M project and the BOF (starting and senior grants) to SPN for financial support, as well as to the Research Foundation – Flanders (FWO) for a PhD fellowship to N.V.T (11I6921N). Umicore AG is gratefully acknowledged for generous gift of materials.

### Notes and references

- (a) A. Bergamo, P. J. Dyson, G. Sava, *Coord. Chem. Rev.*, 2018, **360**, 17–33; (b) I. Ott, *Adv. Inorg. Chem.*, 2020, **75**, 121–148.
- N. J. Wheate, S. Walker, G. E. Craig, R. Oun, *Dalton Trans.*, 2010, **39**, 8113–8127.
- R. Oun, Y. E. Moussa, N. J. Wheate, *Dalton Trans.*, 2018, **47**, 6645–6653.
- (a) T. Zou, C. Tung Lum, C.-N. Lok, J.-J. Zhang, C.-M. Che, *Chem. Soc. Rev.*, 2015, **44**, 8786–8801; (b) J.-J. Zhang, M. A. Abu el Maaty, H. Hoffmeister, C. Schmidt, J. K. Muenzner, R. Schobert, S. Wölfl, I. Ott, *Angew. Chem. Int. Ed.*, 2020, **59**, 16795–16800; (c) S. M. Meier-Menches, B. Neuditschko, K. Zappe, M. Schaijer, M. C. Gerner, K. G. Schmetterer, G. Del Favero, R. Bonsignore, M. Cichna-Markl, G. Koellensperger, A. Casini, C. Gerner, *Chem. Eur. J.*, 2020, **26**, 15528–15537; (d) T. A. C. A. Bayrakdar, T. Scattolin, X. Ma, S. P. Nolan, *Chem. Soc. Rev.*, 2020, **49**, 7044–7100; (e) J. R. Stenger-Smith, P. K. Mascharak, *ChemMedChem*, 2020, **15**, 2136–2145; (f) C. Schmidt, L. Albrecht, S. Balasubramanian, R. Misgeld, B. Karge, M. Brönstrup, A. Prokop, K. Baumann, S. Reichl, I. Ott, *Metallomics*, 2019, **11**, 533–545; (g) C. Schmidt, B. Karge, R. Misgeld, A. Prokop, R. Franke, M. Brönstrup, I. Ott, *Chem. Eur. J.*, 2017, **23**, 1869–1880.
- (a) K. Lin, Z.-Z. Zhao, H.-B. Bo, X.-J. Hao, J.-Q. Wang, *Front. Pharmacol.*, 2018, **9**, 1323; (b) A. Gatti, A. Habtemariam, I. Romero-Canelón, J.-I. Song, B. Heer, G. J. Clarkson, D. Rogolino, P. J. Sadler, M. Carcelli, *Organometallics*, 2018, **37**, 891–899; (c) J. Karges, S. Kuang, F. Maschietto, O. Blacque, I. Ciofini, H. Chao, G. Gasser, *Nat. Commun.*, 2020, **11**, 1–13.
- A. R. Kapdi, I. J. S. Fairlamb, *Chem. Soc. Rev.*, 2014, **43**, 4751–4777; (b) T. Scattolin, I. Caligiuri, N. Mouawad, M. El Boustani, N. Demitri, F. Rizzolio, F. Visentin, *Eur. J. Med. Chem.*, 2019, **179**, 325–334; (c) T. Scattolin, E. Bortolamiol, F. Visentin, S. Palazzolo, I. Caligiuri, T. Perin, V. Canzonieri, N. Demitri, F. Rizzolio, A. Togni, *Chem. Eur. J.*, 2020, **26**, 11868–11876; (d) T. Scattolin, E. Bortolamiol, S. Palazzolo, I. Caligiuri, T. Perin, V. Canzonieri, N. Demitri, F. Rizzolio, L. Cavallo, B. Dereli, M. V. Mane, S. P. Nolan, F. Visentin, *Chem. Commun.* 2020, **56**, 12238–12241.
- E. Boros, P. J. Dyson, G. Gasser, *Chem*, 2020, **6**, 41–60.
- L. Messori, G. Marcon, *Met. Ions Biol. Syst.*, 2004, **41**, 279–304.
- C.K. Mirabelli, R.K. Johnson, C.M. Sung, L. Faucette, K. Muirhead, S.T. Crooke, *Cancer Res.*, 1985, **30**, 32–39.
- Hou, G.X., Liu, P.P., Zhang, S., Yang, M., Liao, J., Yang, J., Hu, Y., Jiang, W.Q., Wen, S., Huang, P., *Cell Death Dis.*, 2018, **9**, 89.
- A. Casini, L. Messori, *Curr. Top. Med. Chem.*, 2011, **11**, 2647–2660.
- (a) M. G. Vander Heiden, L. C. Cantley, C. B. Thompson, *Science*, 2009, **324**, 1029–1033; (b) T. Scattolin, E. Bortolamiol, F. Rizzolio, N. Demitri, F. Visentin, *Appl. Organomet. Chem.*, 2020, **34**, e5876; (c) F. Tresin, V. Stoppa, M. Baron, A. Biffis, A. Annunziata, L. D'Elia, D. M. Monti, F. Ruffo, M. Roverso, P. Sgarbossa, S. Bogialli, C. Tubaro, *Molecules*, 2020, **25**, 3850.
- C. Zoppi, L. Messori, A. Pratesi, *Dalton Trans.*, 2020, **49**, 5906–5913.
- E. Vergara, E. Cerrada, C. Clavel, A. Casini, M. Laguna, *Dalton Trans.*, 2011, **40**, 10927–10935.
- (a) T. Marzo, D. Cirri, C. Gabbiani, T. Gamberi, F. Magherini, A. Pratesi, A. Guerri, T. Biver, F. Binacchi, M. Stefanini, A. Arcangeli, L. Messori, *ACS Med. Chem. Lett.*, 2017, **8**, 997–1001; (b) T. Marzo, L. Massai, A. Pratesi, M. Stefanini, D. Cirri, F. Magherini, M. Becatti, I. Landini, S. Nobili, E. Mini, O. Crociani, A. Arcangeli, S. Pillozzi, T. Gamberi, L. Messori, *ACS Med. Chem. Lett.*, 2019, **10**, 656–660.
- (a) M. V. Baker, P. J. Barnard, S. J. Berners-Price, S. K. Brayshaw, J. L. Hickey, B. W. Skelton, A. H. White, *J. Organomet. Chem.*, 2005, **690**, 5625–5635; (b) P. de Frémont, cE. D. Stevens, M. D. Eelman, D. E. Fogg, S. P. Nolan, *Organometallics*, 2006, **25**, 5824–5828; (c) J. Weaver, S. Gaillard, C. Toyé, S. Macpherson, S. P. Nolan, A. Riches, *Chem. Eur. J.*, 2011, **17**, 6620–6624; (d) B. Bertrand, A. de Almeida, E. P. M. van der Burgt, M. Picquet, A. Citta, A. Folda, M. P. Rigobello, P. Le Gendre, E. Bodio, A. Casini, *Eur. J. Inorg. Chem.*, 2014, 4532–4536; (e) R. Visbal, V. Fernández-Moreira, I. Marzo, A. Laguna, M. C. Gimeno, *Dalton Trans.*, 2016, **45**, 15026–15033; (f) N. Estrada-Ortiz, F. Guarra, I. A. M. de Graaf,

- L. Marchetti, M. H. de Jager, G. M. M. Groothuis, C. Gabbiani, A. Casini, *Chem. Med. Chem.*, 2017, **12**, 1429–1435.
- 17 I. Ott, *Adv. Inorg. Chem.*, 2020, **75**, 121–148.
- 18 T. Scattolin, S. P. Nolan, *Trends Chem.*, 2020, **2**, 721–736.
- 19 (a) A. Collado, A. Collado, A. Gómez-Suárez, A. R. Martin, A. M. Z. Slawin, S. P. Nolan, *Chem. Comm.*, 2013, **49**, 5541–5543; (b) N. V. Tzouras, F. Nahra, L. Falivene, L. Cavallo, M. Saab, K. Van Hecke, A. Collado, C. J. Collett, A. D. Smith, C. S. J. Cazin, S. P. Nolan, *Chem. Eur. J.*, 2020, **26**, 4515–4519; (c) N. V. Tzouras, M. Saab, W. Janssens, T. Cauwenbergh, K. Van Hecke, F. Nahra, S. P. Nolan, *Chem. Eur. J.*, 2020, **26**, 5541–5551; (d) T. Scattolin, N. V. Tzouras, L. Falivene, L. Cavallo, S. P. Nolan, *Dalton Trans.*, 2020, **49**, 9694–9700; (e) F. Nahra, N. V. Tzouras, A. Collado, S. P. Nolan, *Nat. Protoc.*, 2021, **16**, 1476–1493; (f) E. A. Martynova, N. V. Tzouras, G. Pisanò, C. S. J. Cazin, S. P. Nolan, *Chem. Commun.*, 2021, DOI: 10.1039/D0CC08149C; (g) O. Dada, G. Sánchez-Sanz, M. Tacke, X. Zhu, *Tetrahedron Lett.*, 2018, **59**, 2904–2908; (h) W. Walther, O. Dada, C. O’Beirne, I. Ott, G. Sánchez-Sanz, C. Schmidt, C. Werner, X. Zhu, M. Tacke, *Lett. Drug Des. Discov.*, 2017, **14**, 125–134; (i) R. Visbal, R. P. Herrera, M. C. Gimeno, *Chem. Eur. J.*, 2019, **25**, 15837–15845.
- 20 R. Rubbiani, E. Schuh, A. Meyer, J. Lemke, J. Wimberg, N. Metzler-Nolte, F. Meyer, F. Mohr, I. Ott, *Med. Chem. Commun.*, 2013, **4**, 942–948.
- 21 L. M. Doyle, S. O’Sullivan, C. Di Salvo, M. McKinney, P. McArdle, P. V. Murphy, *Org. Lett.*, 2017, **19**, 5802–5805.