



On The Relationship Between Differential Algebra and Tropical Differential Algebraic Geometry

François Boulier, Sebastian Falkensteiner, Marc Paul Noordman

► To cite this version:

François Boulier, Sebastian Falkensteiner, Marc Paul Noordman. On The Relationship Between Differential Algebra and Tropical Differential Algebraic Geometry. 2021. hal-03225922v1

HAL Id: hal-03225922

<https://hal.science/hal-03225922v1>

Preprint submitted on 13 May 2021 (v1), last revised 3 Jul 2021 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On The Relationship Between Differential Algebra and Tropical Differential Algebraic Geometry

François Boulier¹, Sebastian Falkensteiner², and Marc Paul Noordman³

¹ Univ. Lille, CNRS, Centrale Lille, Inria, UMR 9189 - CRISTAL - Centre de Recherche en Informatique Signal et Automatique de Lille, F-59000 Lille, France.

<https://pro.univ-lille.fr/francois-boulier>

² Research Institute for Symbolic Computation (RISC), Johannes Kepler University Linz, Austria. <https://risc.jku.at/m/sebastian-falkensteiner>

³ Bernoulli Institute, University of Groningen, The Netherlands.

<https://www.rug.nl/staff/m.p.noordman>

Abstract. This paper presents some relationship between differential algebra and tropical differential algebraic geometry, mostly focusing on the existence problem of formal power series solutions for systems of polynomial ODE and PDE. Moreover, it improves an approximation theorem involved in the proof of the Fundamental Theorem of tropical differential algebraic geometry which permits to improve this latter by dropping the base field uncountability hypothesis used in the original version.

1 Introduction

Differential algebra is an algebraic theory for systems of ordinary or partial polynomial differential equations. It was founded by Ritt in the first half of the former century [11, 12] and developed by Kolchin [10]. Tropical differential algebraic geometry is a much more recent theory, founded by Grigoriev [8] aiming at applying the concepts of tropical algebra (aka min-plus algebra) to the study of formal power series solutions of systems of ODE. Tropical differential algebra obtained an important impulse by the proof of the Fundamental Theorem of tropical differential algebraic geometry [1] which was recently extended to the partial case in [7]. The common topic of both theories is the existence problem of formal power series solutions of polynomial differential equations on which an important paper [6] by Denef and Lipshitz was published in 1984.

In both [1] and [7], the Fundamental Theorem applies to a polynomial differential system Σ with coefficients in formal power series rings $\mathcal{F}[[x]]$ (ordinary case) or $\mathcal{F}[[x_1, \dots, x_m]]$ (partial case) where \mathcal{F} is a characteristic zero differential field of constants which is both algebraically closed and uncountable. In this paper, we prove that the uncountability hypothesis can be dropped. Indeed, we prove that the Fundamental Theorem holds provided that \mathcal{F} is algebraically closed and has countable⁴ transcendence degree over some field of definition \mathcal{F}_0

⁴ In this paper, “countable” stands for “countably infinite”.

of Σ . This improvement of the Fundamental Theorem is achieved by generalizing the proof of a key proposition, which is an approximation theorem. This generalization is achieved in our Theorem 1, which is the main result of our paper. The new versions of the Fundamental Theorem, which follow, are stated in Theorem 2 and Theorem 3.

For the sake of simplicity, the introductory part of our paper focuses on the ordinary case. For completeness, the partial case is covered as well in the more technical sections. It is structured as follows. We recall in Section 2 the basic ideas underlying formal power series solutions of ODE and point out issues and known results, from the differential algebra literature. We state and explain the Fundamental Theorem of tropical differential algebra in Section 3. We provide our new approximation theorem in Section 5 (covering the partial differential case) and show how it is obtained by adapting the corresponding proposition given in [7]. The new version of the Fundamental Theorem, in the ordinary case, is provided in Section 6. In the final Section 7, we give an overview on the generalizations to the partial case, including the partial version of the Fundamental Theorem.

2 Formal Power Series Solutions of ODE

Let us start with a single autonomous ODE (i.e. an ODE the coefficients of which do not depend of the independent variable x) in a single differential indeterminate y (standing for the unknown function $y(x)$).

$$\dot{y}^2 + 8y^3 - 1 = 0.$$

Differentiate it many different times.

$$\begin{aligned} & 2\dot{y}\ddot{y} + 24y^2\dot{y}, \\ & 2\dot{y}y^{(3)} + 2\ddot{y}^2 + 24y^2\ddot{y} + 48y\dot{y}^2, \\ & \vdots \end{aligned}$$

Rename each derivative $y^{(k)}$ as v_k . Solve the obtained polynomial system. The result is a truncated arc \underline{v}

$$(v_0, v_1, v_2, v_3, v_4, v_5, v_6, v_7, \dots) = (0, 1, 0, 0, -24, 0, 0, 2880, \dots).$$

Plug the arc in the generic formula

$$\Psi(\underline{v}) = \sum \frac{v_i}{i!} x^i.$$

One obtains a formal power series solution centered at the origin. Since the ODE is autonomous, the same arc, plugged in the following generic formula

$$\Psi_\alpha(\underline{v}) = \sum \frac{v_i}{i!} (x - \alpha)^i. \quad (1)$$

provides a formal power series solution centered at any expansion point $x = \alpha$.

If the ODE is not autonomous, the arc depends on the expansion point. The process is thus a variant. Consider some non autonomous ODE

$$x \dot{y}^2 + 8 x y^3 - 1 = 0.$$

Differentiate the ODE many different times.

$$\begin{aligned} 2 x \dot{y} \ddot{y} + \dot{y}^2 + 24 x y^2 \dot{y} + 8 y^3, \\ \vdots \end{aligned}$$

Then fix an expansion point α and evaluate the independent variable at $x = \alpha$. Solve the obtained polynomial system. The result is a truncated arc. Plug it in (1) (for the chosen value of α). One gets a formal power series solution centered at $x = \alpha$.

In the above processes, the only issue lies in the polynomial solving step. Indeed, each differentiated equation introduces a new leading derivative. These leading derivatives admit as leading coefficients the initial or the separant of the ODE. If these two polynomials do not vanish at the expansion point and the already secured coordinates of the truncated arc (the initial values, somehow, of the initial value problem), then the formal power series solution exists, is unique and straightforward to compute up to any term. However, if these polynomials vanish, the formal power series solution may fail to exist or be unique.

A device borrowed from [6, page 236] illustrates the issue. It shows how to build an ODE p with coefficients in $\mathbb{Q}[x]$ from a polynomial $f(z)$ in $\mathbb{Q}[z]$. The ODE admits a formal power series solution centered at the origin if and only if the polynomial $f(z)$ has no positive integer solution. In the ordinary case, this device permits to build interesting examples. The approach generalizes to the partial case. It permits to relate the existence problem of formal power series solutions centered at the origin for PDE systems to Hilbert's Tenth Problem and Matiassevich undecidability result [6, Theorem 4.11]. For more details see [3, Sect 1.6].

It is interesting also to observe that any non autonomous ODE can be viewed as an autonomous one by performing a change of independent variable and introducing an extra ODE. Indeed, call ξ the new independent variable. View the former independent variable x as a new differential indeterminate (i.e. as an unknown function $x(\xi)$) and introduce the extra ODE $\dot{x} = 1$. This reduction method only applies to ODE with polynomial coefficients in x . However, if $x = \alpha$ was a problematic expansion point before the reduction then $x(0) = \alpha$ becomes a problematic initial value (hence arc coordinate) after reduction. For more details see [3, Sect 1.4.2].

In his books [11, 12], Ritt implicitly considers autonomous systems (the “autonomous” qualifier does not belong to differential algebra) and we may assume he had in mind the above reduction trick. Though Taylor expansions of solutions are discussed at different places (mostly in a chapter dedicated to PDE), Ritt does not explicitly address the existence problem of formal power series

solutions. However, he pioneered differential elimination methods by means of his theory of characteristic sets (which was much developed afterwards, leading to the theories of regular chains and differential regular chains). This elimination theory solves in particular the following decision problem: given any finite system Σ of ordinary or partial differential polynomial, does $1 \in [\Sigma]$ where $[\Sigma]$ denotes the differential ideal generated by Σ ? This problem is equivalent to the following one, which is thus seen to be decidable: does there exist initial values such that Σ has formal power series solutions⁵?

In the case of systems of non autonomous ODE, thanks to the reduction method to the autonomous case, we can then conclude that the following problem is decidable: given any system Σ , do expansion point and initial values exist such that Σ has formal power series solutions?

3 The Fundamental Theorem of Tropical Differential Algebraic Geometry

In the tropical differential case, the systems under consideration belong to some differential polynomial ring $\mathcal{F}[[x]]\{y_1, \dots, y_n\}$ where \mathcal{F} is a characteristic zero field of constants. Differential polynomials have formal power series coefficients. Thus the reduction trick to the autonomous case does not apply and formal power series solutions are only sought at a fixed expansion point: the origin. More precisely, formal power series solutions are sought in the coefficient ring $\mathcal{F}[[x]]$ of the equations.

The existence problem of such formal power series solutions is much more difficult. An important related paper is [6]. Indeed, [6, Theorem 3.1] claims that, in the case of systems with coefficients in $\mathbb{Q}[x]$, the existence problem of formal power series solutions (with coefficients in \mathbb{C} , \mathbb{R} or \mathbb{Q}_p) is decidable. It is however important to note that, in the same setting, the existence problem of nonzero formal power series solutions is undecidable. See [6, Proposition 3.3] which refers to [14].

In this context, the Fundamental Theorem of tropical differential geometry does not solve any of the left open problems. It only states the following equivalence

$$\text{trop}(\text{sol}(\Sigma)) = \text{sol}(\text{trop}(\Sigma)), \quad (2)$$

where Σ is a differential ideal and the base field \mathcal{F} is both algebraically closed and uncountable (we relinquish this last condition in this paper).

Before entering sophisticated issues, we need to clarify the notations used in (2). The support of a formal power series

$$\varphi = \sum a_i x^i \quad (3)$$

⁵ Indeed, the characteristic sets or regular differential chains computed by differential elimination methods can be viewed as differential systems sufficiently simplified to generalize, for systems of differential equations, the basic methods sketched at the top of the section for computing formal power series solution.

is the set $\{i \in \mathbb{N} \mid a_i \neq 0\}$. The tropicalization $\text{trop}(\varphi)$ of φ is defined as its support.

Since Σ depends on n differential indeterminates y_1, \dots, y_n , its formal power series solutions actually are tuples of n formal power series. One then extends the above definition to tuples of formal power series: the tropicalization of a tuple

$$\varphi = (\varphi_1, \dots, \varphi_n) \quad (4)$$

is defined as the tuple $\text{trop}(\varphi) = (\text{trop}(\varphi_1), \dots, \text{trop}(\varphi_n))$.

On the left hand side of (2), $\text{sol}(\Sigma)$ denotes the set of formal power series solutions of Σ with coefficients in \mathcal{F} . Hence, the left hand side of (2) is the set of the tropicalizations of all the formal power series solutions of Σ . It is a set of tuples of supports.

Let us address now the right hand side of (2). The valuation of a formal power series (3) is defined as ∞ if $\varphi = 0$ and as the smallest $i \in \mathbb{N}$ such that $a_i \neq 0$ otherwise.

$$\begin{array}{ccc} \mathcal{F}[[x]] & \xrightarrow{\text{supp}} & \mathcal{P}(\mathbb{N}) \\ & \searrow \text{valuation} & \downarrow \text{min} \\ & & \mathbb{N} \end{array}$$

Let us now define the tropicalization of a differential monomial

$$m = c v_1^{d_1} \dots v_r^{d_r} \quad (5)$$

at a tuple of supports

$$S = (S_1, \dots, S_n). \quad (6)$$

The coefficient $c \in \mathcal{F}[[x]]$ and each v stands for some derivative of some of the n differential indeterminates y_1, \dots, y_n . Consider any tuple of formal power series (4) whose tropicalization is S . Since m is a monomial, the support of the formal power series $m(\varphi)$ is uniquely defined by S : it does not depend on the actual coefficients of φ . We are led to the following definition⁶:

The tropicalization of a differential monomial m at S is defined as the valuation of $m(\varphi)$ where φ is any tuple of formal power series whose tropicalization is S . Here are a few examples

monomial m	support S	$\text{trop}(m)$ at S
$x^2 y$	$\{0, 1, 2\}$	2
$x^2 y$	$\{2\}$	4
\dot{y}^3	$\{0, 3\}$	6
\dot{y}^3	$\{0, 1\}$	∞

⁶ This is not the definition given in [1, sect. 4] but both definitions are equivalent.

Let us now consider a differential polynomial, expanded as a sum of monomials of the form (5) (if m and m' are two different monomials in the sum (7), we require $v_1^{d_1} \cdots v_r^{d_r}$ to be different from $v_1^{d'_1} \cdots v_{r'}^{d'_{r'}}$):

$$p = m_1 + m_2 + \cdots + m_t. \quad (7)$$

The tropicalization of p at S is defined as

$$\text{trop}(p) = \min_{i=1}^t \text{trop}(m_i). \quad (8)$$

As an example, let us consider the differential polynomial

$$p = \dot{y}^2 - 4y \quad (9)$$

whose solutions are $\varphi = 0$ (support $S = \emptyset$) and $\varphi = (x + c)^2$ where c is an arbitrary constant (supports $S = \{0, 1, 2\}$ and $\{2\}$). The first and second derivatives of p are

$$\dot{p} = 2\dot{y}\ddot{y} - 4\dot{y}, \quad (10)$$

$$\ddot{p} = 2\dot{y}y^{(3)} + 2\ddot{y}^2 - 4\ddot{y}. \quad (11)$$

In the next table, all the considered supports are supports of solutions of the differential polynomials. In the last column, the list of the $\text{trop}(m_i)$ is provided, rather than their minimum. The first row indicates that both monomials of p vanish at $\varphi = 0$. The second row indicates that the two monomials do not vanish but may possibly cancel each other at $\varphi = a_2 x^2$, for some $a_2 \neq 0$ (indeed, they vanish but only for $a_2 = 1$). The third row indicates that, among the three monomials of \ddot{p} , the first one vanishes at any $\varphi = a_2 x^2$ while the two last ones may cancel each other for some $a_2 \neq 0$.

polynomial	support S	list $\text{trop}(m_i)$ at S
p	\emptyset	$[\infty, \infty]$
\dot{p}	$\{2\}$	$[2, 2]$
\ddot{p}	$\{2\}$	$[\infty, 0, 0]$

In the next table, the considered support $S = \{0, 1\}$ is not the support of any solution of p , since p has no solution of the form $\varphi = a_0 + a_1 x$ with $a_0, a_1 \neq 0$. This fact is not observed on the first row, which considers p itself. It is however observed on the second row, which considers the first derivative of p : one of the two monomials vanishes while the second one evaluates to some non zero formal power series.

polynomial	support S	list $\text{trop}(m_i)$ at S
p	$\{0, 1\}$	$[0, 0]$
\dot{p}	$\{0, 1\}$	$[\infty, 0]$

The observed phenomena suggest the following definition, which permits to understand the right hand side of (2).

Let p be a polynomial of the form (7). View $\text{trop}(p)$ as a function of n unknown supports. Then (S_1, \dots, S_n) is said to be a *solution* of $\text{trop}(p)$ if either

1. each $\text{trop}(m_i) = \infty$ or
2. there exists m_i, m_j ($i \neq j$) such that $\text{trop}(m_i) = \text{trop}(m_j) = \min_{k=1}^t (\text{trop}(m_k))$.

Let us conclude this section by a few remarks. In the Fundamental Theorem of tropical differential algebraic geometry, the inclusion $\text{trop}(\text{sol}(\Sigma)) \subset \text{sol}(\text{trop}(\Sigma))$ is easy. The difficult part is the converse inclusion. It requires Σ to be a differential ideal because one may need to consider arbitrary high derivatives of the elements of Σ in order to observe that a given support is not a solution (see the example above). It requires the base field \mathcal{F} to be algebraically closed because of the polynomial system solving step and the fact that solutions are sought in $\mathcal{F}[[x]]$.

Last, its proof relies on an approximation theorem. The two versions of this approximation theorem given in [1, Proposition 7.3] and [7, Proposition 6.3] assume \mathcal{F} to be uncountable. Our new version (Theorem 1) relies on weaker hypotheses.

4 Fields of Definition and Countability

We are concerned with a differential ideal Σ [10, I, sect. 2] in a characteristic zero partial differential polynomial ring $\mathcal{F}[[x_1, \dots, x_m]]\{y_1, \dots, y_n\}$ where \mathcal{F} is an algebraically closed field of constants, the m derivation operators $\delta_1, \dots, \delta_m$ act as $\partial/\partial x_1, \dots, \partial/\partial x_m$ and y_1, \dots, y_n are n differential indeterminates.

Thanks to the Ritt-Raudenbush Basis Theorem (see [4] for details), the differential ideal Σ can be presented by finitely many differential polynomials $g_1, \dots, g_s \in \mathcal{F}[[x_1, \dots, x_m]]\{y_1, \dots, y_n\}$.

The next definition is adapted from [10, I, sect. 5].

Definition 1. *A field of definition of Σ is any subfield $\mathcal{F}_0 \subset \mathcal{F}$ such that $g_1, \dots, g_s \in \mathcal{F}_0[[x_1, \dots, x_m]]\{y_1, \dots, y_n\}$.*

Proposition 1. *Any differential ideal Σ has a countable algebraically closed field of definition \mathcal{F}_0 . Moreover, if \mathcal{F} has countable transcendence degree over \mathcal{F}_0 then \mathcal{F} also is countable.*

Proof. Let S be the family of the coefficients of the formal power series coefficients of any basis of Σ which are transcendental over the field \mathbb{Q} of the rational numbers. The family S is countable. An algebraically closed field of definition \mathcal{F}_0 can be defined as the algebraic closure of $\mathbb{Q}(S)$.

Now, the field \mathbb{Q} is countable. If \mathcal{L} is a countable field and S is a countable family of transcendental elements over \mathcal{L} then $\mathcal{L}(S)$ is countable. Moreover, if \mathcal{L} is countable then its algebraic closure is countable [9, Theorem 65].

The last statement of the proposition follows using the same arguments.

In the sequel, \mathcal{F}_0 denotes an algebraically closed field of definition of Σ .

5 The Approximation Theorem

Denote Θ the commutative semigroup of the derivative operators generated by the derivation operators i.e. $\Theta = \{\delta_1^{a_1} \cdots \delta_m^{a_m} \mid a_1, \dots, a_m \in \mathbb{N}\}$.

Define a one-to-one correspondence between the set of all pairs $(i, \theta) \in [1, n] \times \Theta$ and the set \mathbb{N} of nonnegative integers. This correspondence permits us to enumerate all derivatives θy_i of the differential indeterminates. Fix a correspondence which defines an orderly ranking (derivatives are enumerated by increasing order) [10, chap. I, sect. 8]. The derivatives of the y are denoted v_0, v_1, v_2, \dots .

Define another one-to-one correspondence between the set of all pairs $(i, \theta) \in [1, s] \times \Theta$ and \mathbb{N} . This correspondence permits us to enumerate all derivatives θg_i of the differential polynomials. Again, fix a correspondence which defines an orderly ranking on the derivatives of the g (viewing them as s differential indeterminates). The derivatives of the g , evaluated at $x_1 = \cdots = x_m = 0$, are denoted f_0, f_1, f_2, \dots . The polynomials f thus belong to $\mathcal{F}_0\{y_1, \dots, y_n\}$.

Let k be a positive integer. Denote

$$\begin{aligned}\Sigma_k &= \{f_i \mid 0 \leq i \leq k\}, \\ \Sigma_\infty &= \{f_i \mid i \in \mathbb{N}\}.\end{aligned}$$

Define $\kappa(k) = \kappa$ as the smallest integer such that $\Sigma_k \subset \mathcal{F}_0[v_0, \dots, v_\kappa]$. The index κ exists because the ranking is orderly. Define

$$A_k = \{a \in \mathcal{F}_0^{\kappa+1} \mid f_0(a) = \cdots = f_k(a) = 0\}.$$

Let now S be any subset of \mathbb{N} . Define $A_{k,S}$ as the set of zeros of A_k which are *compatible with S* :

$$A_{k,S} = \{a \in A_k \mid a_i \neq 0 \text{ if and only if } i \in S \cap [0, \kappa]\}.$$

Indeed, thanks to the fixed one-to-one correspondence between the derivatives of the differential indeterminates and the set \mathbb{N} , any such set S encodes a tuple of n supports of formal power series. Define (observe that the coordinates of the elements of A_∞ belong to \mathcal{F})

$$\begin{aligned}A_\infty &= \{a \in \mathcal{F}^\mathbb{N} \mid f_i(a) = 0 \text{ for each } i \in \mathbb{N}\}, \\ A_{\infty,S} &= \{a \in A_\infty \mid a_i \neq 0 \text{ if and only if } i \in S\}.\end{aligned}$$

Theorem 1. *Assume \mathcal{F} has countable transcendence degree over \mathcal{F}_0 . Let S be any subset of \mathbb{N} . If $A_{k,S} \neq \emptyset$ for each $k \in \mathbb{N}$ then $A_{\infty,S} \neq \emptyset$.*

Proof. Let \mathcal{R} be the ring obtained by inversion of all derivatives with indices in S and quotient by the ideal equal to the sum of the ideal generated by Σ_∞ and the ideal generated by the derivatives with indices not in S , i.e.

$$\mathcal{R} = \mathcal{F}_0[v_i, v_j^{-1} \mid i \in \mathbb{N}, j \in S] / (f_i, v_j \mid i \in \mathbb{N}, j \notin S). \quad (12)$$

By Lemma 1, this ring is not the null ring. By Krull's Theorem, it contains a maximal ideal \mathfrak{m} . The field \mathcal{R}/\mathfrak{m} has at most countable transcendence degree over its subfield \mathcal{F}_0 . Thus, it is isomorphic to a subfield \mathcal{F}_1 of \mathcal{F} . In order to see this, consider $\mathcal{F}_0(\bar{v}_0, \bar{v}_1, \dots)$, where $\bar{v}_i \in \mathcal{R}/\mathfrak{m}$ are the images of the v_i . Pick any transcendence basis $B = \{\bar{v}_k \mid k \in I \subset \mathbb{N}\}$ of this field extension. B is at most countable. Map them to the elements of a corresponding transcendence basis of \mathcal{F}_1 over \mathcal{F}_0 (the hypothesis on the countable transcendence degree of \mathcal{F}_1 over \mathcal{F}_0 is used here in order to have one-to-one correspondence between the two bases). Then map the $\bar{v}_i \notin B$ — which are algebraic over $\mathcal{F}_0(B)$ — to corresponding algebraic elements of \mathcal{F}_1 , using the fact that \mathcal{F}_1 is algebraically closed.

Hence, we have a ring homomorphism $\mathcal{F}_0\{y_1, \dots, y_n\} \rightarrow \mathcal{R} \rightarrow \mathcal{R}/\mathfrak{m} \rightarrow \mathcal{F}_1 \subset \mathcal{F}$. The images of the v_i by the above map provide a solution of Σ_∞ in \mathcal{F} .

Though its presentation is different, our proof of Theorem 1 is very close to [7, Proposition 6.3]. One recovers the proof of [7] (mostly due to Marc Paul Noordman and much inspired by methods borrowed from [6]) by changing the last paragraph as follows:

Proof. Recall that in this proof \mathcal{F} is supposed to be uncountable. [...] By Lemma 1, this ring is not the null ring. It thus contains a maximal ideal \mathfrak{m} . The ring \mathcal{R}/\mathfrak{m} is a field. As a \mathcal{F} -algebra, it is countably generated (since \mathcal{R} is). Therefore, it is of countable dimension as a \mathcal{F} -vector space (it is generated, as a \mathcal{F} -vector space, by the products of some set of its generators, as a \mathcal{F} -algebra). If $t \in \mathcal{R}/\mathfrak{m}$ were transcendental over \mathcal{F} then, by the theory of partial fraction decomposition, the elements $1/(t - \alpha)$ for $\alpha \in \mathcal{F}$ would form an \mathcal{F} -linearly independent subset of \mathcal{R}/\mathfrak{m} . This subset would be of uncountable dimension since \mathcal{F} is assumed to be uncountable: a contradiction. Thus \mathcal{R}/\mathfrak{m} is algebraic over \mathcal{F} . Since \mathcal{F} is algebraically closed, \mathcal{R}/\mathfrak{m} is isomorphic to \mathcal{F} .

Lemma 1. *The ring \mathcal{R} defined in (12) is not the null ring.*

Proof. We prove the lemma by showing that Σ_∞ admits a solution in some field \mathcal{F}_0^* (which turns out to be an ultrafield - see Section 5.1) and constructing a map $\mathcal{F}_0\{y_1, \dots, y_n\} \rightarrow \mathcal{F}_0^*$ which factors as $\mathcal{F}_0\{y_1, \dots, y_n\} \rightarrow \mathcal{R} \rightarrow \mathcal{F}_0^*$.

To each $k \in \mathbb{N}$ associate an element $a^k \in A_{k,S}$. We have

$$a^k = (a_0^k, a_1^k, \dots, a_\kappa^k) \in \mathcal{F}_0^\kappa.$$

Fix any non principal ultrafilter \mathcal{D} on \mathbb{N} and consider the ultrafield $\mathcal{F}_0^* = (\prod_{i \in \mathbb{N}} \mathcal{F}_0) / \mathcal{D}$. For each $i \in \mathbb{N}$ define $u_i \in \mathcal{F}_0^*$ by

$$\begin{aligned} u_0 &= (a_0^0, a_0^1, a_0^2, \dots, a_0^k, \dots), \\ u_1 &= (a_1^0, a_1^1, a_1^2, \dots, a_1^k, \dots), \\ &\vdots \\ u_i &= (a_i^0, a_i^1, a_i^2, \dots, a_i^k, \dots), \\ &\vdots \end{aligned}$$

On each column k of the above “array”, the elements a_i^k such that $i > \kappa$ are not defined. Set them to zero. Observe that on each row, there are only finitely many such elements.

We have thus defined a map $v_i \mapsto u_i$.

Let now $i \in \mathbb{N}$ be the index of some polynomial $f_i = f$. Evaluate f to an element of \mathcal{F}_0^* by substituting u_j to v_j for each $j \in \mathbb{N}$. Ultrafield operations are performed componentwise and the zeros of the f appear on the columns in the above array. Thus, f evaluates to zero over the k th coordinate of \mathcal{F}_0^* for all sufficiently large values of k . This set of values of k is cofinite and hence, f evaluates to zero in \mathcal{F}_0^* .

Let now $i \in \mathbb{N}$ be the index of some derivative v_i . By definition of $A_{k,S}$, if $i \notin S$ then all the coordinates of u_i are zero so that u_i is zero in \mathcal{F}_0^* ; if $i \in S$ then the coordinates a_i^k of u_i are nonzero for all sufficiently large values of k and u_i is nonzero in \mathcal{F}_0^* .

The mapping $v_i \mapsto u_i$ thus defines a zero of Σ_∞ which is compatible with S and with coordinates in \mathcal{F}_0^* .

5.1 Basic Notions on Ultraproducts

This section is much inspired by [13, chap. 2] with the notations of [2]. It is provided for the convenience of casual readers.

The set \mathbb{N} is used as an index set on which we fix a (so called) *non-principal ultrafilter* \mathcal{D} . It is by definition a collection of infinite subsets of \mathbb{N} closed under finite intersection, with the property that for any subset $E \subset \mathbb{N}$ either E or its complement $\mathbb{N} \setminus E$ belongs to \mathcal{D} . In particular, the empty set does not belong to \mathcal{D} and, if $E \in \mathcal{D}$ and F is an arbitrary set containing E then also $F \in \mathcal{D}$. Otherwise, $\mathbb{N} \setminus F \in \mathcal{D}$ and therefore, $\emptyset = E \cap (\mathbb{N} \setminus F) \in \mathcal{D}$: a contradiction. Since every set in \mathcal{D} must be infinite, it follows that every set whose complement is finite (such a set is called *cofinite*) belongs to \mathcal{D} .

Let \mathcal{R}_i ($i \in \mathbb{N}$) be a collection of rings. We form the ultraproduct $\mathcal{R}^* = (\prod_{i \in \mathbb{N}} \mathcal{R}_i) / \mathcal{D}$ (or the ultrapower $\mathcal{R}^\mathbb{N} / \mathcal{D}$ if all rings \mathcal{R}_i are the same ring \mathcal{R}) as follows. On the Cartesian product $\prod_{i \in \mathbb{N}} \mathcal{R}_i$ we define the equivalence relation: $a \equiv b$ if and only if the set of indices i such that $a_i = b_i$ belongs to the ultrafilter \mathcal{D} .

We are going to use the following facts.

Consider an element $a \in \mathcal{R}^*$ which has no nonzero coordinates. The set of indices such that $a_i = 0$ is empty. Since the empty set does not belong to the ultrafilter, $a \neq 0$ in \mathcal{R}^* .

Consider an element $a \in \mathcal{R}^*$ which has only finitely many nonzero coordinates. The set of indices such that $a_i = 0$ is cofinite. Thus it belongs to \mathcal{D} . Thus $a = 0$ in \mathcal{R}^* .

An ultraproduct of rings is a ring: addition and multiplication are performed componentwise. Let us prove that an ultraproduct of fields is a field (called an *ultrafield*). Consider some $a \neq 0$ in \mathcal{R}^* . Then the set $E = \{i \in \mathbb{N} \mid a_i \neq 0\}$ does not belong to \mathcal{D} . Thus its complement $F = \mathbb{N} \setminus E$ belongs to \mathcal{D} . Define \bar{a} as follows: for each $i \in \mathbb{N}$, if $a_i = 0$ take $\bar{a}_i = 0$ else take $\bar{a}_i = a_i^{-1}$. Let $u = a \bar{a}$.

The set of indices such that $u_i = 1$ is F , which belongs to \mathcal{D} . Thus a admits an inverse and \mathcal{K}^* is a field.

6 The New Version of the Fundamental Theorem

For completeness, we provide the part of the proof of the Fundamental Theorem which makes use of our Theorem 1. The proof is the same as that of [1, Theorem 8]. We start with an easy Lemma [1, Remark 4.1].

Lemma 2. *Let $S = (S_1, \dots, S_n)$ be a tuple of n supports and $m = cv_1^{d_1} \dots v_r^{d_r}$ be a monomial. Then $\text{trop}(m) = 0$ at S if and only if the valuation of c is zero and each factor $v^d = (y_j^{(k)})^d$ of m is such that $k \in S_j$.*

Before stating the Fundamental Theorem, let us stress that the fields \mathcal{F}_0 and \mathcal{F} mentioned in Theorem 2 can be assumed to be countable, by Proposition 1.

Theorem 2 (Fundamental Theorem for ODE).

Let Σ be a differential ideal of $\mathcal{F}[[x]]\{y_1, \dots, y_n\}$ where \mathcal{F} is an algebraically closed field of constants and \mathcal{F}_0 be an algebraically closed field of definition of Σ . If \mathcal{F} has countable transcendence degree over \mathcal{F}_0 then

$$\text{trop}(\text{sol}(\Sigma)) = \text{sol}(\text{trop}(\Sigma)). \quad (13)$$

Proof. The inclusion \subset is easy. We prove the converse one. We assume that S is not the support of any solution of Σ and we show that S is not a solution of $\text{trop}(\Sigma)$.

For this, we are going to build a differential polynomial $h \in \Sigma$, expanding to a sum of monomials

$$h = m_1 + m_2 + \dots + m_r, \quad (14)$$

such that $\text{trop}(m_1) = 0$ and $\text{trop}(m_i) > 0$ for $2 \leq i \leq r$.

By the Ritt-Raudenbush Basis Theorem (see [4] for details), there exists a finite set g_1, \dots, g_s of differential polynomials of Σ such that the solution set of Σ is the solution set of the differential ideal $[g_1, \dots, g_s]$ generated by the g .

From now on, we use the notations introduced in Section 5. Since $[g_1, \dots, g_s]$ has no solution with support S we have $A_{\infty, S} = \emptyset$ whence, by Theorem 1, there exists some index k such that $A_{k, S} = \emptyset$. Recall that $A_{k, S}$ is a subset of the algebraic variety of some polynomial system obtained by prolonging (and evaluating at $x = 0$) the system of the g up to some order and that the prolonged system belongs to some polynomial ring $\mathcal{F}_0[v_1, \dots, v_\kappa]$.

We are going to seek the differential polynomial h in the ideal (f_0, \dots, f_k) of the Noetherian polynomial ring $\mathcal{F}_0[v_1, \dots, v_\kappa]$. Observe that this polynomial h does not belong to Σ , which is an ideal of $\mathcal{F}_0[[x]]\{y_1, \dots, y_n\}$ because the polynomials f are obtained after evaluation at $x = 0$ of derivatives of the g . This is actually not a problem since all the monomials cancelled by this evaluation have

a coefficient c with positive valuation (Lemma 2). See also [1, Remark 5.1]. More precisely, the existence of a polynomial h in (f_0, \dots, f_k) implies the existence of another polynomial (14) in Σ .

The ideal (f_0, \dots, f_k) has no solution compatible with S . This means that

$$[f_0 = \dots = f_k = 0 \text{ and } v_j = 0 \text{ for all } v_j \text{ s.t. } j \notin S] \Rightarrow v_\ell = 0 \text{ for some } \ell \in S$$

$$\Rightarrow \underbrace{\prod_{\ell \in S, \ell \leq \kappa} v_\ell}_{m_1} = 0.$$

By Lemma 2 we have $\text{trop}(m_1) = 0$ at S . By Hilbert's Nullstellensatz, we have

$$m_1 \in \sqrt{(f_0, \dots, f_k, (v_j)_{j \notin S})}$$

Thus there exists a positive integer d and monomials m_2, \dots, m_r such that $h = m_1^d + m_2 + \dots + m_r$ belongs to the ideal (f_0, \dots, f_k) and for each $2 \leq i \leq r$, there exists some $j \notin S$ for which $\deg(m_i, v_j) > 0$. By Lemma 2, we have $\text{trop}(m_i) > 0$ at S for each $2 \leq i \leq r$.

7 The Partial Differential Case

In this section we give an overview on the generalization to the case of partial differential equations.

We seek for solutions of systems $\Sigma \subset \mathcal{F}[[x_1, \dots, x_m]]\{y_1, \dots, y_m\}$ in the ring of multivariate formal power series $\mathcal{F}[[x_1, \dots, x_m]]$.

In this case, the algorithmic problems are even worse than in the ordinary case. According to [6, Theorem 4.11], there even cannot be an algorithm for deciding solvability of linear systems, a subclass of algebraic differential equations as we consider. Instead of actually computing the solutions of Σ , we again present an equivalent description of the solutions in the form of (2).

As in the ordinary case ($m = 1$), the support of a formal power series

$$\varphi = \sum a_I x^I = \sum a_{(i_1, \dots, i_m)} x_1^{i_1} \cdots x_m^{i_m}$$

is the set $\{I \in \mathbb{N}^m \mid a_I \neq 0\}$ and the tropicalization of φ is defined as its support. Hence, the left hand side of (2) is defined also for $m > 1$.

For the tropicalization of Σ , the generalization can not be done straightforward, since there is no well-defined minimum of elements in \mathbb{N}^m . In [7] is used instead a very specific partial order.

Let $X \subset \mathbb{N}^m$. The *Newton polytope* $\mathcal{N}(X) \subseteq \mathbb{R}_{\geq 0}^m$ of X is defined as the convex hull of $X + \mathbb{N}^m = \{x + n \mid x \in X, n \in \mathbb{N}^m\}$. Moreover, $x \in X$ is called a *vertex* if $x \notin \mathcal{N}(X \setminus \{x\})$, and vert is the set of vertices of X . It follows that $\text{vert}(X)$ is the minimal set in \mathbb{N}^m (with respect to the relation “ \subset ”) generating $\mathcal{N}(X)$. Let us denote all vertex sets as $\mathbb{T}_m = \{\text{vert} \mid X \subset \mathbb{N}^m\}$. Then, the composition

of taking the support and then its vertex set of the formal power series defines a *non-degenerate valuation* such that some ideas of [1] can be recovered.

$$\begin{array}{ccc} \mathcal{F}[[x_1, \dots, x_m]] & \xrightarrow{\text{supp}} & \mathcal{P}(\mathbb{N}^m) \\ & \searrow \text{non-deg. valuation} & \downarrow \text{vert} \\ & & \mathbb{T}_m \end{array}$$

The tropicalization of a differential monomial $m = cv_1^{d_1} \cdots v_r^{d_r}$ at a tuple of supports $S = (S_1, \dots, S_n) \subset (\mathbb{N}^m)^n$ is defined as the non-degenerate valuation of $m(\varphi)$ from above with $\text{supp}(\varphi) = S$. Let us illustrate this.

monomial m	support S	trop(m) at S
$x_1 x_2 y$	$\{(1, 0), (0, 1)\}$	$\{(2, 0), (0, 2)\}$
$(\frac{\partial y}{\partial x_1})^2$	$\{(2, 0), (0, 2)\}$	$\{(2, 0)\}$
$(\frac{\partial^2 y}{\partial x_1 \partial x_2})^2$	$\{(2, 0), (0, 2)\}$	\emptyset

The tropicalization of a differential polynomial p of the form (5) at S is defined as

$$\text{trop}(p) = \text{vert}\left(\bigcup_{i=1}^t \text{trop}(m_i)\right). \quad (15)$$

Let us consider the polynomial

$$p = \frac{\partial y}{\partial x_1} \cdot \frac{\partial y}{\partial x_2} + (-x_1^2 + x_2^2) \frac{\partial^2 y}{\partial x_1 \partial x_2}. \quad (16)$$

polynomial	support S	list trop(m_i) at S
p	$\{(2, 0), (0, 2)\}$	$[\{(1, 1)\}, \emptyset]$
p	$\{(2, 0), (1, 1), (0, 2)\}$	$[\{(2, 0), (1, 1), (0, 2)\}, \{(2, 0), (0, 2)\}]$

Considering $\text{trop}(p)$ as a function of n unknown supports, (S_1, \dots, S_n) is said to be a *solution* of $\text{trop}(p)$ if for every vertex $J \in \text{trop}(p)$ there exists m_i, m_j ($i \neq j$) such that $J \in \text{trop}(m_i) \cap \text{trop}(m_j)$. In the example above, we see that there cannot be a solution of p with support equals $\{(2, 0), (0, 2)\}$, but $\varphi = x_1^2 + 2x_1x_2 - x_2^2$ is indeed a solution.

For more illustrations of the tropicalization of (partial) differential polynomials, see [5].

As in the ordinary case, the inclusion $\text{trop}(\text{sol}(\Sigma)) \subset \text{sol}(\text{trop}(\Sigma))$ in the Fundamental Theorem is relatively easy [7, Proposition 5.7]. The converse inclusion can be shown exactly as in [7, Section 6], except that we replace Proposition 6.3 by the Approximation Theorem 1.

For consistency let us recall the main result here.

Theorem 3 (Fundamental Theorem for PDE). *Let Σ be a differential ideal of $\mathcal{F}[[x_1, \dots, x_m]]\{y_1, \dots, y_n\}$ where \mathcal{F} is an algebraically closed field of constants and \mathcal{F}_0 be an algebraically closed field of definition of Σ . If \mathcal{F} has countable transcendence degree over \mathcal{F}_0 then*

$$\text{trop}(\text{sol}(\Sigma)) = \text{sol}(\text{trop}(\Sigma)). \quad (17)$$

Acknowledgements

The authors would like to thank Zoé Chatzidakis and Mercedes Haiech for their help and their comments. The first author would like to acknowledge the support of the bilateral project ANR-17-CE40-0036 DFG-391322026 SYMBIONT.

References

1. Aroca, F., Garay, C., Toghani, Z.: The Fundamental Theorem of Tropical Differential Algebraic Geometry. *Pacific J. Math.* **283**(2), 257–270 (2016), [arXiv:1510.01000v3](#)
2. Becker, J., Denef, J., Lipshitz, L., van den Dries, L.: Ultraproducts and Approximation in Local Rings I. *Inventiones mathematicae* **51**, 189–203 (1979)
3. Boulier, F.: A Differential Algebra Introduction For Tropical Differential Geometry (2019), lecture Notes for the Workshop on Tropical Differential Geometry, Queen Mary College of the University of London, Dec. 2-7. Available at <https://hal.archives-ouvertes.fr/hal-02378197>
4. Boulier, F., Haiech, M.: The Ritt-Raudenbush Theorem and Tropical Differential Geometry (2019), available at <https://hal.archives-ouvertes.fr/hal-02403365>
5. Cotterill, E., Garay, C., Luviano, J.: Exploring tropical differential equations. <http://arxiv.org/abs/2012.14067> (2020)
6. Denef, J., Lipshitz, L.: Power Series Solutions of Algebraic Differential Equations. *Mathematische Annalen* **267**, 213–238 (1984)
7. Falkensteiner, S., Garay-López, C., Haiech, M., Noordman, M.P., Toghani, Z., Boulier, F.: The Fundamental Theorem of Tropical Partial Differential Algebraic Geometry. In: *Proceedings of ISSAC 2020*. pp. 178–185. Kalamata, Greece (2020)
8. Grigoriev, D.: Tropical Differential Equations. [arXiv:1502.08010v1](#) (2015)
9. Kaplansky, I.: *Fields and Rings*. The University of Chicago, Chicago, second edn. (1972)
10. Kolchin, E.R.: *Differential Algebra and Algebraic Groups*. Academic Press, New York (1973)
11. Ritt, J.F.: *Differential equations from the algebraic standpoint*, American Mathematical Society Colloquium Publications, vol. 14. American Mathematical Society, New York (1932)
12. Ritt, J.F.: *Differential Algebra*, American Mathematical Society Colloquium Publications, vol. 33. American Mathematical Society, New York (1950)
13. Schoutens, H.: *The Use of Ultraproducts in Commutative Algebra*. Springer Verlag (2010)
14. Singer, M.: The Model Theory of Ordered Differential Fields. *Journal of Symbolic Logic* **43**(1), 82–91 (1978)