


# The poor lonesome A subgenome of *Brassica napus* L. may not survive without its mate

**Alexandre Pelé<sup>1</sup>, Gwenn Trotoux<sup>1</sup>, Frédérique Eber<sup>1</sup>, Sylvie Negre<sup>1</sup>, Jérôme Morice<sup>1</sup>, Maryse Lodé-Taburel<sup>1</sup>, Marie-Madeleine Gilet<sup>1</sup>, Cyril Falentin<sup>1</sup>, Mathieu Rousseau-Gueutin<sup>1</sup>, Anne-Marie Chèvre<sup>1</sup>**

<sup>1</sup>IGEPP, INRA, Agrocampus-Ouest, Université de Rennes 1 , Le Rheu, France


# Divergence occurred since *B. napus* formation



(U, 1935 ; Cheung et al., 2009 ; Chalhoub et al., 2014)

# How can we study this evolution?

**Short-term:** Resynthesized *B. napus*




Natural *B. napus* (AACC, 2n=38)

# How can we study this evolution?

## Short-term: Resynthesized *B. napus*

*B. rapa*  
(AA, 2n=20)      *B. oleracea*  
(CC, 2n=18)



### S0 Functional & Epigenetic Changes

(Lukens *et al.*, 2006; Albertin *et al.* 2006; Gaeta *et al.*, 2007 ; Xu *et al.*, 2009)

### S1 Burst of Genetic Changes

(Song *et al.*, 1995; Gaeta *et al.*, 2007 ; Szadkowski *et al.*, 2010;2011; Xiong *et al.*, 2011)

S2

...


Sx

Limited to the first few generations

Natural *B. napus* (AACC, 2n=38)

# How can we study this evolution?

**Long-term:** Comparing *B. napus* to  
the present *B. rapa* / *B. oleracea*




# How can we study this evolution?

**Long-term:** Comparing *B. napus* to the present *B. rapa* / *B. oleracea*

**But progenitors have evolved independently since *B. napus* formation**

**Genic conversion  
Gene Loss / Gain  
Rearrangements**


(Wang *et al.*, 2011 ; Chalhoub *et al.*, 2014, Liu *et al.*, 2014)



**Present *B. napus* AACC,  
*B. rapa* AA and *B. oleracea* CC**

# How can we study this evolution?

## Long-term: Subgenome Extraction



# How can we study this evolution?

Only used in AABBDD Wheat  
for extracting the AABB  
component


(Kerber et al., 1964)

Evolution in polyploid context



Interdependency between each  
original genome?


## Long-term: Subgenome Extraction



Diploid *B. napus* component (AA, 2n=20)

# Extraction of the *B. napus* AA component



## Strategy 1



Theoretical %  
of *B. napus* A  
subgenome in  
AAC hybrids

# Extraction of the *B. napus* AA component

## Strategy 1





Leflon et al., 2006

Theoretical %  
of *B. napus* A  
subgenome in  
AAC hybrids

# Extraction of the *B. napus* AA component

## Strategy 1





Leflon et al., 2006

Theoretical %  
of *B. napus* A  
subgenome in  
AAC hybrids

AA component (2n=20)



# Extraction of the *B. napus* AA component

## Strategy 1



# Extraction of the *B. napus* AA component

## Strategy 1




**Impossibility to obtain the pure AA component of *B. napus*:**

- Rearrangements ?
- Evolution in a polyploid context of the *B. napus* A subgenome ?

# Extraction of the *B. napus* AA component

## Strategy 2

*B. napus* cv. Darmor  
(AACC, 2n=38)



*B. rapa* cv. C1.3  
(AA, 2n=20)




F1 AAC (2n=29)



Cycle 1

Theoretical % of *B. napus* A subgenome in AAC hybrids



# Extraction of the *B. napus* AA component

Strategy 2

Theoretical % of *B. napus* A  
subgenome in AAC hybrids

*B. napus* cv. Darmor  
(AACC, 2n=38)



F1 AAC (2n=29)



*B. rapa* cv. C1.3  
(AA, 2n=20)



AA G1

Cycle 1

X

X

25

# Extraction of the *B. napus* AA component

Strategy 2

Theoretical % of *B. napus* A subgenome in AAC hybrids

*B. napus* cv. Darmor  
(AACC, 2n=38)       X *B. rapa* cv. C1.3  
(AA, 2n=20)      

F1 AAC (2n=29)      

AACC (2n=38)      

AA G1      

AAC (2n=29)      

AACC (2n=38)      

Self fertilization

AAC (2n=29)      

AACC (2n=38)      

AA G4      

AAC (2n=29)      

AA G5      

Cycle 1

Cycle 2,3

Cycle 4

Cycle 5

25

...

68.3


76.3

# Extraction of the *B. napus* AA component

Strategy 2

Theoretical % of *B. napus* A subgenome in AAC hybrids

*B. napus* cv. Darmor  
(AACC, 2n=38)      *B. rapa* cv. C1.3  
(AA, 2n=20)



# Molecular characterization of the G4 and G5 AA plants

% *B. napus* A subgenome extracted

**60 K Illumina Array®**  
(52,127 SNP)

Polymorphism between:  
*B. rapa* (C1.3)  
*B. napus* (Darmor)

7,846 SNP

Genetically & physically  
anchored on A subgenome  
(Darmor/Yudal genetic map)

2,290 SNP

*B. napus* C subgenome introgressions

**60 K Illumina Array®**  
(52127 SNP)

Polymorphism :  
*B. napus* (Darmor) ++  
*B. rapa* (C1.3) --

18,674 SNP

Genetically & physically  
anchored on C subgenome  
(Darmor/Yudal genetic map)

4,807 SNP

# Molecular characterization of the G4 and G5 AA plants

% *B. napus* A subgenome extracted

	% <i>B. napus</i> theoretical	% <i>B. napus</i> observed
<b>AA G4</b>	<b>68%</b>	<b>57%</b>
<b>AA G5 pl1</b>	<b>76%</b>	<b>65%</b>
<b>AA G5 pl2</b>	<b>76%</b>	<b>68%</b>

**Extraction of *B. napus* A subgenome is significantly lower than expected in all AA plants ( $\chi^2$  test,  $\alpha<0.001$ )**

*B. napus* C subgenome introgressions

	C01 start (0.53 Mbp)	C02 start (4.64 Mbp)	C02 end (2.27 Mbp)
<b>AA G4</b>	<b>X</b>	<b>X</b>	
<b>AA G5 pl1</b>		<b>X</b>	
<b>AA G5 pl2</b>			<b>X</b>

**3 introgressions detected and validated (sequencing) with variations between G4 & G5**

# Molecular characterization of the G4 and G5 AA plants

## % *B. napus* A subgenome extracted


	% <i>B. napus</i> theoretical	% <i>B. napus</i> observed
<b>AA G4</b>	<b>68%</b>	<b>57%</b>
<b>AA G5 pl1</b>	<b>76%</b>	<b>65%</b>
<b>AA G5 pl2</b>	<b>76%</b>	<b>68%</b>

**Extraction of *B. napus* A subgenome is significantly lower than expected in all AA plants ( $\chi^2$  test,  $\alpha < 0.001$ )**

Pure *B. napus* AA plant may not survive deprived of C subgenome?


## *B. napus* C subgenome introgressions

	C01 start (0.53 Mbp)	C02 start (4.64 Mbp)	C02 end (2.27 Mbp)
<b>AA G4</b>	<b>X</b>	<b>X</b>	
<b>AA G5 pl1</b>		<b>X</b>	
<b>AA G5 pl2</b>			<b>X</b>




Introgressions of *B. napus* C subgenome don't impact the yield

# Molecular characterization of the G4 and G5 AA plants




# Molecular characterization of the G4 and G5 AA plants

Where are located fixed  
*B. napus* regions in AA  
plants?




# Molecular characterization of the G4 and G5 AA plants

What is unexpected?



# Molecular characterization of the G4 and G5 AA plants

What is unexpected?




# Molecular characterization of the G4 and G5 AA plants

*B. napus* S0

± 7,500 YEARS

1,925 genes lost  
along the A  
subgenome

Natural *B. napus*  
(Chalhoub et al., 2014)




# Molecular characterization of the G4 and G5 AA plants

*B. napus* S0

± 7,500 YEARS


1,925 genes lost  
along the A  
subgenome

Current *B. napus*  
(Chalhoub et al., 2014)



# CONCLUSION

- ❖ We were unable to produce pure *B. napus* AA plants and AA plants generated (G4 & G5) presented a lower proportion of *B. napus* extracted than expected
- ❖ These results were related to the *B. napus* structural evolution in a polyploid context (genes lost)
- ❖ And suggest that only 7,500 years of coevolution leads to interdependency between the original subgenome



**68% of the *B. napus* A subgenome extracted**




How far can we go? New cycle ongoing but:



**595 candidates genes**



Functional characterization of the genes restored by the *B. rapa* genomic regions remaining in AA plants



Low pollen  
fertility


Poor  
Vigor

# ACKNOWLEDGMENT

## Biodiversity & Polyploidy

### Team:

Anne-Marie CHEVRE  
Mathieu ROUSSEAU-GUEUTIN  
Gwenn TROTOUX  
Jérôme MORICE  
Sylvie NEGRE  
Maryse LODE  
Marie GILET  
Olivier CORITON  
Virginie HUTEAU  
Frédérique EBER



**...And thank you for your attention**

