

# Brassica Allotriploids Hybrids: New Way to Exceed Recombination Limits for COs Rate and Distribution



Alexandre Pelé<sup>1</sup>, Gwenn Trotoux<sup>1</sup>, Frédérique Eber<sup>1</sup>, Sylvie Negre<sup>1</sup>, Maryse Lodé<sup>1</sup>, Marie-Madeleine Gilet<sup>1</sup>, Olivier Coriton<sup>1</sup>, Virginie Huteau<sup>1</sup>, Matthieu Falque<sup>2</sup>, Olivier C Martin<sup>2</sup>, Jérôme Morice<sup>1</sup>, Mathieu Rousseau-Gueutin<sup>1</sup>, Anne-Marie Chèvre<sup>1</sup>

<sup>1</sup>IGEPP, INRA, Agrocampus-Ouest, Université de Rennes 1 , Le Rheu, France


<sup>2</sup>GQE-LeMoulon, INRA, Université Paris-Sud, CNRS, AgroParisTech, Université Paris-Saclay, Gif sur Yvette, France

# What is the interest of Crossovers(COs)? What about their frequency and distribution?

## INTEREST IN PLANT BREEDING

Essential for mixing genetic diversity


(Wijnker and De Jong, 2008)



# What is the interest of Crossovers(COs)? What about their frequency and distribution?

## INTEREST IN PLANT BREEDING


Essential for mixing genetic diversity  
(Wijnker and De Jong, 2008)



(Mercier et al., 2014)

## HOWEVER:

- Rarely more than 2 COs per chromosome and per meiosis
- Non homogenously distributed along the chromosomes (Mézard et al., 2007)



CO distribution along wheat 3B chromosome and Arabidopsis 4 chromosome

(Drouaud et al., 2007 ; Saintenac et al., 2007)

Depends of the chromosome conformation  
No CO in the vicinity of centromeric regions

# What is the interest of Crossovers(COs)? What about their frequency and distribution?

## INTEREST IN PLANT BREEDING

Essential for mixing genetic diversity  
(Wijnker and De Jong, 2008)

P<sub>1</sub>      P<sub>2</sub>

## BREEDING CONSTRAINTS:

- Large progenies
- Impossibility to separate some genes

## INTEREST:


To control the number and distribution of COs events

Progenies selection

(Mercier et al., 2014)

## HOWEVER:

- Rarely more than 2 COs per chromosome and per meiosis
- Non homogenously distributed along the chromosomes (Mézard et al., 2007)



CO distribution along wheat 3B chromosome and Arabidopsis 4 chromosome

(Drouaud et al., 2007 ; Saintenac et al., 2007)

Depends of the chromosome conformation  
No CO in the vicinity of centromeric regions

# Why oilseed rape is a good model for studying COs?



- **1<sup>st</sup> oleaginous species in Europe and 4<sup>th</sup> in the world (64,8 Mt)**
- **Use for: Oil production & Seed cake**  
(FAOSTAT, 2012)



***B. rapa* AA,**  
**2n=20**  
(Wang *et al.*, 2011)

X



***B. oleracea***  
**CC, 2n=18**  
(Liu *et al.*, 2014)



***B. napus***  
**AACC, 2n=38**  
(Chalhoub *et al.*, 2014)


# Why oilseed rape is a good model for studying COs?



- 1<sup>st</sup> oleaginous species in Europe and 4<sup>th</sup> in the world (64,8 Mt)
- Use for: Oil production & Seed cake  
(FAOSTAT, 2012)

**Recombination rate between homologous A07 chromosomes increase in AAC (2n=29) compare to AA (2n=20) hybrids**

*B. rapa* AA,  
2n=20  
(Wang et al., 2011)




(Leflon et al., 2010)


## QUESTIONS

- What about the CO frequency and distribution between the 10 homologous A chromosomes?
- What the impact of male and female meiosis?

(Chalhoub et al., 2014)




# F1 hybrids production for recombination analysis on A genome in AA and AAC hybrids



$A_1A_1$  and  $A_2A_2$  represent *B. rapa* cv C1.3 and Chiifu (Sequenced, Wang et al., 2011), respectively.  $C_oC_o$  designates the *B. oleracea* cv (RC34).  $A_D A_D C_D C_D$  represents the *B. napus* cv Darmor.

# F1 hybrids production for recombination analysis on A genome in AA and AAC hybrids



# Selection of polymorphic SNP markers covering the 10 A chromosomes

60 K Illumina Array®  
(52127 SNP)

- Polymorphic SNPs between  $A_2A_2$  and  $A_1A_1$ ,  $A_1A_1C_0C_0$ ,  $A_DA_D$ ,  $A_DA_DC_DC_D$

1980 SNPs

- SNPs Homogenously distributed along the 10 A chromosomes (Wang et al., 2011)
- Adapted to KASPAR®

214 SNPs

- KASPAR® for F<sub>1</sub> progenies
- Genetic mapping at 2pt LOD > 3
- Genetic mapping = physical location

204 SNPs



# Selection of polymorphic SNP markers covering the 10 A chromosomes

60 K Illumina Array®  
(52127 SNP)

- Polymorphic SNPs between  $A_2A_2$  and  $A_1A_1$ ,  $A_1A_1C_0C_0$ ,  $A_D A_D$ ,  $A_D A_D C_0C_0$


1980 SNPs

- SNPs Homogenously distributed along the 10 A chromosomes (Wang et al., 2011)
- Adapted to KASPAR®

214 SNPs


- KASPAR® for  $F_1$  progenies
- Genetic mapping at 2pt LOD > 3
- Genetic mapping = physical location

204 SNPs





1 SNP / 1.25 Mb  
(Var = 0.35 ;  $\sigma$  = 0.59)  
Recovering 94% of A genome

# What about frequency and distribution of COs in AA vs AAC hybrids?



# What about frequency and distribution of COs in AA vs AAC hybrids?





CO frequency always  
AAC >> AA  
between all homologous A  
chromosomes


Genetic background can affect  
recombination rate, particularly in AAC  
hybrids

# What about frequency and **distribution** of COs in AA vs AAC hybrids?


\*  $\chi^2$  test (0.05)



# What about frequency and **distribution** of COs in AA vs AAC hybrids?



\*  $\chi^2$  test (0.05)



**Expected Recombination profile along each homologous A chromosome in AA hybrids**



**Depending on chromosome conformation**

- No CO in pericentromeric regions
- Gradient for CO rate along chromosome arms (Linear regression;  $0.4 < R^2 < 0.98$ ; p-value < 0.05)



**No variation with genetic background**

**What is going on in AAC hybrids?**



# What about frequency and **distribution** of COs in AA vs AAC hybrids?



# What about frequency and **distribution** of COs in AA vs AAC hybrids?



# What about frequency and distribution of COs in AA vs AAC hybrids?




Recombination profiles always  
 $AAC \neq AA$   
between each homologous A chromosome

NOT Depending of chromosome conformation

- CO in pericentromeric regions
- No Gradient for CO rate along chromosome arms

Variation between AAC genetic backgrounds,  
particularly in telomeric regions


# What about interference in AA vs AAC hybrids?




Intensity of interference :  
 $AA \ggg AAC$   
for each homologous A  
chromosome

Are the data equal after female or male meiosis?


# What about frequency and distribution of COs in male and female meiosis in AA vs AAC hybrids?





# What about frequency and distribution of COs in male and female meiosis in AA vs AAC hybrids?





# What about frequency and distribution of COs in male and female meiosis in AA vs AAC hybrids?





# What about frequency and distribution of COs in male and female meiosis in AA vs AAC hybrids?



# What about frequency and distribution of COs in male and female meiosis in AA vs AAC hybrids?



# What about frequency and distribution of COs in male and female meiosis in AA vs AAC hybrids?




**Recombination profiles between each A homologous chromosome in male and female meiosis is:**

- Equal between AA hybrids
- Similar between AAC hybrids but frequently female > male

Always AAC  $\neq$  AA hybrids


# What about interference in male and female meiosis in AA vs AAC hybrids?



Intensity of interference in AA and AAC hybrids:  
male ~ female  
for each homologous A chromosome

But always AA >> AAC

# What about interference in male and female meiosis in AA vs AAC hybrids?



Intensity of interference in AA and AAC hybrids:  
male ~ female  
for each homologous A chromosome

But always AA >> AAC



# Conclusion

## ❖ AAC vs AA hybrids, whatever genetic background and male/female meiosis:

- Increase of CO rate
- Change of CO distribution
- Decrease in interference intensity



By using AAC hybrids, we can break  
the linkage disequilibrium

## ❖ AAC male vs female hybrids:

- Variation in CO rate (female > male)
- Similar CO distribution
- Similar interference intensity



The increase of recombination rate in  
AAC male is due to Type I CO probably  
non interfering


## Prospect:

- What are the mechanisms responsible for modifications of CO rate and distribution? **Epigenetics?** (Melamed-Bessudo and Levy, 2012 ; Kim *et al.*, 2005 ; Gaut *et al.*, 2007 ; Kirik *et al.*, 2008; Tsai *et al.*, 2008; Mirouze *et al.*, 2012; Yelina *et al.*, 2012; Jahns *et al.*, 2014)
- Use in breeding programs

# ACKNOWLEDGMENT

## INRA Le Rheu:

Anne-Marie CHEVRE  
Mathieu ROUSSEAU-GUEUTIN  
Gwenn TROTOUX  
Jérôme MORICE  
Sylvie NEGRE  
Maryse LODE  
Marie GILET  
Olivier CORITON  
Virginie HUTEAU  
Frédérique EBER



## INRA Le Moulon:

Matthieu FALQUE  
Olivier MARTIN

**...And thank you for your attention**

## INRA Clermont-Ferrand:

Charles PONCET

**INRA Versailles:** Christine MEZARD & Eric JENCZEWSKI