
BRASSICA ALLOTRIPOLOIDS, A SOLUTION TO DEAL WITH HAMPERED MEIOTIC RECOMBINATION

Alexandre Pelé¹, Franz Boideau¹, Maryse Lodé-Taburel¹, Gwenn Trotoux¹, Frederique Eber¹, Sylvie Negre¹, Marie-Madeleine Gilet¹, Olivier Coriton¹, Virginie Huteau¹, Matthieu Falque², Olivier C Martin², Jérôme Morice¹, Mathieu Rousseau-Gueutin¹, Anne-Marie Chèvre¹

¹IGEPP, INRA, Agrocampus-Ouest, Université de Rennes 1 , Le Rheu, France

²GQE-LeMoulon, INRA, Université Paris-Sud, CNRS, AgroParisTech, Université Paris-Saclay, Gif sur Yvette, France



21st Crucifer Genetics Conference
Saint-Malo, 01st July 2018



SCIENTIFIC CONTEXT

Meiotic recombination process in brief, from its roles to its limits



- ❖ Ensures the formation of well-balanced gametes





SCIENTIFIC CONTEXT

Meiotic recombination process in brief, from its roles to its limits

- ❖ Ensures the formation of well-balanced gametes & the generation of diversity



Parent 1 Parent 2




SCIENTIFIC CONTEXT

Meiotic recombination process in brief, from its roles to its limits

- ❖ But number and distribution of crossovers are tightly controlled

- ✓ Per pair of homologs, rarely more than 3 COs occur
- ✓ COs distribution is NOT homogenous along the chromosomes





SCIENTIFIC CONTEXT

Meiotic recombination process in brief, from its roles to its limits

- ❖ But number and distribution of crossovers are tightly controlled


- ✓ Per pair of homologs, rarely more than 3 COs occur
- ✓ COs distribution is NOT homogenous along the chromosomes



SCIENTIFIC CONTEXT

How can we enhance meiotic recombination in *Brassica*?


- ❖ By exploiting *Brassica* allotriploids



(Leflon et al., 2010)



B. napus AAC,
2n=4x=38
(Chalhoub et al., 2014)



B. rapa AA,
2n=2x=20
(Wang et al., 2011)




AAC
2n=3x=29

QUESTIONS

- ✓ What about COs rate and distribution between the 10 A homologs in AAC allotriploids?
- ✓ What about CCA allotriploids?

RATE AND DISTRIBUTION OF COs IN AA vs AAC HYBRIDS


Material & Methods



- A_nA_n = A genome of *B. napus* cv. Darmor extracted
- A_rA_r = *B. rapa* cv. Chiifu (Sequenced)
- $A_nA_nC_nC_n$ = *B. napus* cv. Darmor (Sequenced).



RATE AND DISTRIBUTION OF COs IN AA vs AAC HYBRIDS

Material & Methods



RATE AND DISTRIBUTION OF COs IN AA vs AAC HYBRIDS


Material & Methods



1 SNP / 1.25 Mb (SE = 0.04, n = 194)
Covering 94% of the A genome

RATE AND DISTRIBUTION OF COs IN AA vs AAC HYBRIDS

CO rates between the homologous A chromosomes in AA and AAC hybrids








CO rates: AA << AAC

Between all homologous A chromosomes,
extending at the whole A genome the
results of Leflon *et al.* (2010)

RATE AND DISTRIBUTION OF COs IN AA vs AAC HYBRIDS

CO rates between the homologous A chromosomes in AA and AAC hybrids



CO rates: AA << AAC



At the whole A genome the number of COs
is 3.4-fold higher in AAC hybrid

The number of COs is positively correlated
with the A chromosome size

RATE AND DISTRIBUTION OF COs IN AA vs AAC HYBRIDS

COs distribution along the A chromosomes in the AA hybrid

- ❖ CO rates increase gradually from centromeres toward chromosome extremities





CAPTION
 A_nA_r' (2x)

RATE AND DISTRIBUTION OF COs IN AA vs AAC HYBRIDS

COs distribution along the A chromosomes in the AA hybrid

- ❖ ~10% of the A genome is totally deprived of COs, especially close to the centromeres





CAPTION
 A_nA_r' (2x)

RATE AND DISTRIBUTION OF COs IN AA vs AAC HYBRIDS

COs distribution along the A chromosomes in the AA hybrid

- ❖ ~10% of the A genome is totally deprived of COs, especially close to the centromeres



CAPTION
 $A_n A_r$, (2x)

In AA hybrid

Expected Recombination landscapes are
observed along each A chromosome

With:



- ✓ No CO in the vicinity of centromeres
- ✓ Gradient for CO rates along chromosome arms

What is going on in AAC hybrid?

RATE AND DISTRIBUTION OF COs IN AA vs AAC HYBRIDS

COs distribution along the A chromosomes in the AAC hybrid



- ❖ Does CO rates increase proportionally along the A chromosomes compared to the AA hybrid?



RATE AND DISTRIBUTION OF COs IN AA vs AAC HYBRIDS

COs distribution along the A chromosomes in the AAC hybrid



- ❖ Does CO rates increase proportionally along the A chromosomes compared to the AA hybrid? **NO**



RATE AND DISTRIBUTION OF COs IN AA vs AAC HYBRIDS

COs distribution along the A chromosomes in the AAC hybrid

- ❖ CO rates increase significantly in most intervals compared to the AA hybrid





CAPTION

A_nA_r' (2x)
 $A_nA_r'C_n$ (3x)

RATE AND DISTRIBUTION OF COs IN AA vs AAC HYBRIDS

COs distribution along the A chromosomes in the AAC hybrid



- ❖ Even in intervals close or including the **centromeres**, where new recombining regions are formed



RATE AND DISTRIBUTION OF COs IN AA vs AAC HYBRIDS

COs distribution along the A chromosomes in the AAC hybrid

- ❖ Even in intervals close or including the **centromeres**, where new recombining regions are formed



CAPTION

A_nA_r , (2x)
 $A_nA_rC_n$, (3x)

In AAC hybrid



Recombination landscapes along each of the A chromosomes are \neq AA hybrid

- ✓ COs are formed all along the A genome, even in the vicinity of the centromeres
- ✓ No Gradient for CO rates along chromosome arms

RATE AND DISTRIBUTION OF COs IN AA vs AAC HYBRIDS

In conclusion:

- The additional C genome allows to change the rate and distribution of COs between the homologous A chromosomes in AAC allotriploids
- Whatever the sex of meiosis and C genome origin, but with slight variations for COs rate



RATE AND DISTRIBUTION OF COs IN CC vs CCA HYBRIDS

Rate and distribution of COs are also modified along the C homologs in CCA hybrids





B. napus AACC,
 $2n=4x=38$
(Chalhoub et al., 2014)



B. oleracea CC,
 $2n=2x=20$
(Wang et al., 2011)

X

CCA
 $2n=3x=28$




In *Brassica*, it is possible to enhance recombination both between A and C homologs, by using AAC and CCA allotetraploid hybrids, respectively

CONCLUSIONS

Interest for rapeseed breeding programs

- ❖ AAC and CCA hybrids are fertile ($\pm 10\%$ of *B. napus* fertility)
- ❖ Both can produce AACCC plants through intercross with *B. napus*





See the Talk of AM Chèvre

CONCLUSIONS

Interest for rapeseed breeding programs

- ❖ AAC and CCA hybrids allow to generate a larger diversity of gametes
- ❖ Shuffle genomic regions usually deprived of COs, where genes of agronomic interest have been detected




*QTL of yield and seed quality (Bouchet et al., 2016)

CONCLUSIONS

Concerning the speciation success of polyploidy plants


- ❖ *B. napus* may have been formed though a triploid-bridge 7,500 years ago



ACKNOWLEDGMENT

INRA LE RHEU :

ANNE-MARIE CHÈVRE
MATHIEU ROUSSEAU-GUEUTIN
Gwenn TROTOUX
Maryse LODE
Marie GILET
Sylvie NÈGRE
Jérôme MORICE
Gwenaëlle DENIOT
Cyril FALENTIN
Olivier CORITON
Virginie HUTEAU
Frédérique EBER
Thibaut JOUSSEAUME
Benoît MOTAIS (L2)
Guillaume GLAIS (L2)
Sofiane AZOUZ (M2)
Franz BOIDEAU (M1 + PhD)



GENTYANE :

Charles PONCET
Amélie BERTIN

INRA LE MOULON :

Matthieu FALQUE
Olivier MARTIN

INRA VERSAILLES :

ERIC JENCZEWSKI
CHRISTINE MÉZARD

...And thank you for your attention