

# MASER: A Science Ready Toolbox for Low Frequency Radio Astronomy

Baptiste Cecconi, Alan Loh, Pierre Le Sidaner, Renaud Savalle, Xavier Bonnin, Quynh Nhu Nguyen, Sonny Lion, Albert Shih, Stéphane Aicardi, Philippe Zarka, et al.

# ▶ To cite this version:

Baptiste Cecconi, Alan Loh, Pierre Le Sidaner, Renaud Savalle, Xavier Bonnin, et al.. MASER: A Science Ready Toolbox for Low Frequency Radio Astronomy. CODATA Data Science Journal, 2020, 19, 10.5334/dsj-2020-012 . hal-03228972

# HAL Id: hal-03228972 https://hal.science/hal-03228972

Submitted on 19 May 2021  $\,$ 

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



# DATA ARTICLE

# MASER: A Science Ready Toolbox for Low Frequency Radio Astronomy

Baptiste Cecconi<sup>1,2</sup>, Alan Loh<sup>1</sup>, Pierre Le Sidaner<sup>3</sup>, Renaud Savalle<sup>3</sup>, Xavier Bonnin<sup>1</sup>, Quynh Nhu Nguyen<sup>1</sup>, Sonny Lion<sup>1</sup>, Albert Shih<sup>3</sup>, Stéphane Aicardi<sup>3</sup>, Philippe Zarka<sup>1,2</sup>, Corentin Louis<sup>1,4</sup>, Andrée Coffre<sup>2</sup>, Laurent Lamy<sup>1,2</sup>, Laurent Denis<sup>2</sup>, Jean-Mathias Grießmeier<sup>5</sup>, Jeremy Faden<sup>6</sup>, Chris Piker<sup>6</sup>, Nicolas André<sup>4</sup>, Vincent Génot<sup>4</sup>, Stéphane Erard<sup>1</sup>, Joseph N. Mafi<sup>7</sup>, Todd A. King<sup>7</sup>, Jim Sky<sup>8</sup> and Markus Demleitner<sup>9</sup>

- <sup>2</sup> Station de Radioastronomie de Nançay, Observatoire de Paris, CNRS, PSL, Université d'Orléans, Nançay, FR
- <sup>3</sup> DIO, Observatoire de Paris, CNRS, PSL, Paris, FR
- <sup>4</sup> IRAP, CNRS, Université Paul Sabatier, CNES, Toulouse, FR
- <sup>5</sup> LPC2E, CNRS, Université d'Orléans, Orléans, FR
- <sup>6</sup> Dep. Physics and Astronomy, University of Iowa, Iowa City, Iowa, US
- <sup>7</sup> IGPP, UCLA, Los Angeles, California, US
- <sup>8</sup> Radio Sky Publishing, US

<sup>9</sup> Heidelberg Universität, Heidelberg, DE

Corresponding author: Baptiste Cecconi (baptiste.cecconi@observatoiredeparis.psl.eu)

MASER (Measurements, Analysis, and Simulation of Emission in the Radio range) is a comprehensive infrastructure dedicated to time-dependent low frequency radio astronomy (up to about 50 MHz). The main radio sources observed in this spectral range are the Sun, the magnetized planets (Earth, Jupiter, Saturn), and our Galaxy, which are observed either from ground or space. Ground observatories can capture high resolution data streams with a high sensitivity. Conversely, space-borne instruments can observe below the ionospheric cut-off (at about 10 MHz) and can be placed closer to the studied object. Several tools have been developed in the last decade for sharing space physics data. Data visualization tools developed by various institutes are available to share, display and analyse space physics time series and spectrograms. The MASER team has selected a sub-set of those tools and applied them to low frequency radio astronomy. MASER also includes a Python software library for reading raw data from agency archives.

Keywords: Radio astronomy; Tools; Interoperability

# 1. Introduction

Low frequency radio data are providing remote proxies to remotely study energetic and unstable magnetised plasmas. In the solar system, all magnetized plasma environments are emitting radio emissions. The corresponding radio sources are non-thermal emission phenomena, and are not related to atomic and molecular transitions contrarily to electromagnetic emissions at higher frequencies. Their beaming pattern is also strongly anisotropic (Zarka 1998). The *low frequency* radio emissions are observed in the standard VLF (~3 kHz) to VHF (~30 MHz) radio bands. The main radio sources of the solar system are the Sun, Jupiter and Saturn. The Earth, Uranus and Neptune are also hosting natural radio emissions. The planetary radio emissions are linked to the magnetospheric dynamics – i.e., auroral activity, radiation belts, etc. – as well as planetary atmospheres – i.e., lightning electromagnetic pulses.

<sup>&</sup>lt;sup>1</sup> LESIA, Observatoire de Paris, CNRS, PSL, Meudon, FR

The usual data product for low frequency radio emissions observations is a "dynamic spectrum" (a time varying spectrogram). Other products in use are high temporal resolution waveform snapshots (see, e.g., Briand et al 2016) and catalogues of events (with a radio bursts classification, see, e.g. Marques et al 2017). In this frequency range, it is not yet possible to build imaging radio telescopes in space, so that the main source of knowledge is this time-frequency representation of the data. Until recently each low frequency data provider was storing their data products in local formats, or using standard formats with local meta-data dictionaries, which prevented interoperability. The NASA space physics community has promoted the Common Data Format (CDF, http://cdf.gsfc.nasa.gov) format with International Solar Terrestrial Program (ISTP, https://spdf.gsfc.nasa.gov/istp\_guide/istp\_guide.html) guidelines, for day to day usage and archiving. NASA's Planetary Data System (PDS) archive is now accepting CDF/ISTP as an archive format (King & Mafi 2018), and many space mission teams have adopted the same scheme. Ground based observatories are producing data collections reaching several TB per day (Lamy 2017). However, even with a common file format, downloading large data volumes for local processing is not optimal (e.g., long download delays) and should be avoided. There is thus a need for science-ready tools and standards that cover the needs of the low frequency radio astronomy community for time-dependent data.

## 2. The MASER collaboration

MASER (http://maser.lesia.obspm.fr) (Cecconi 2018) is a collaboration of teams throughout the world, whose aim is to facilitate the open access to science ready low frequency radio data. It is led by Observatoire de Paris (ObsParis) in France including people from LESIA (Laboratoire d'Etudes Spatiales et Instrumentation en Astrophysique) and PADC (Paris Astronomical Data Centre). It gathers scientists and software engineers from other space plasma and radio astronomy labs in France (Orléans, Nançay and Toulouse), and in the USA (University of Iowa, and University of California Los Angeles). Regular collaborations also exist with colleagues in Japan (Tohoku University) and Poland.

The MASER team at ObsParis is organized around 4 tasks: (a) data distribution; (b) codes and models; (c) infrastructure and interfaces; and (d) the *MaserLib* open-source repository. Task (a) is covering the full data lifecycle: from the production of the data (for ongoing and future data collection), the preparation of their distribution (data formatting, metadata, previews...), their validation (against the selected standards), the implementation of the access interfaces (web portal, virtual observatory, streaming interface...), as well as the curation of data when applicable. The details are described in a regularly updated Data Management Plan, following the open science policies developed at ObsParis (see **Figure 1**). The other French teams are working with ObsParis to implement similar policies on their data collections.



Figure 1: Synthetic description of the MASER data management plan.

The other tasks deal with infrastructure and software development. Task (b) focusses on modelling codes (e.g., ray tracing code, radio observation modelling..), working with the science team to open the source code, share the simulation runs (through task (a)) and setup run-on-demand capabilities. Task (c) is the development and maintenance of the generic interoperable infrastructures in use to distribute the data collections managed in task (a) and (b). Task (d) is the development of the open source software libraries.

### 3. Data and Metadata Formats

MASER promotes the use of community standards for the data formats and metadata dictionaries. The space physics community is using CDF, whereas the Solar physics remote sensing community is using files formatted in Flexible Image Transport System (FITS, https://fits.gsfc.nasa.gov) (Pence 2010). Interoperability also requires enforcing adoption of standard metadata. MASER thus implements standards from the Heliophysics and Planetary Science communities: Space Physics Archive Search and Extracted (SPASE, http://spase-group.org) and ISTP metadata for Heliophysics; the Virtual European Solar and Planetary Access (VESPA, http://www.europlanet-vespa.eu) metadata for Planetary Sciences (Erard 2018).

#### 4. Tools and Interfaces

The display tools and interfaces selected by MASER have been initially developed for space physics applications, as well as astronomy and solar system sciences. Space physics tools used by MASER have been developed by the University of Iowa: *Autoplot* (http://autoplot.org) (Faden 2010) and *Das2* (https://das2.org) (Piker 2018). We also use more generic technologies, such as VESPA, which provides a search interface that allows the discovery of data of interest for scientific users, and is based on International Virtual Observatory Alliance (IVOA) astronomy standards.

#### 4.1. Data streaming interface

The driving issue of MASER is the *science ready* and *remote access* to low frequency radio astronomy data collections, and more specifically to long time-series or high-resolution datasets.

For instance, each Solar transit observation (8 hours) data from the NewRoutine receiver of the Nancay Decameter Array (NDA) is stored in a 768 MB file, with 57,600 consecutive spectra (1 spectrum every 500 ms) (Lamy 2017). Displaying data on a typical computer screen requires about 2,000 pixels on the horizontal (i.e., temporal) axis, so that displaying the aforementioned Solar transit only requires transferring about 27 MB after temporal resampling (a reduction factor of about 30). As presented by Lamy (2017), some datasets from Nançay are reaching a few milliseconds of temporal resolution, so that the daily file size reaches several TBytes, and the data collection spans over several years. Remote visualisation of such data with adaptive temporal resolution streaming capabilities is thus needed.

The *Das2* data streaming technology allows to visualize data with a server-side time-axis resampling. The transmitted data are adjusted to the client temporal resolution, leading to a reduction of the data transfer over Internet. This reduces the delay for displaying the data, proportionally to the resampling rate, as specified by *das2* clients, such as *Autoplot* or the *das2py* library (https://github.com/das-developers/das2py). The installation and configuration of the *das2* server framework (https://github.com/das-developers/das2-pyserver), is simple and straightforward. Data providers have to develop a data reader script, which writes out the data as a *das2stream* into the local standard output for a given input time interval. The *das2stream* format is documented in its Interface Control Document (ICD) (Piker et al 2017).

In addition to the original das2 servers at University of Iowa, two other das2 servers are running to serve LESIA (http://voparis-das-maser.obspm.fr/das2/server) and Nançay (https://das2server.obs-nancay.fr/das2/server) datasets. Data readers are using the *maser4py* library (see section 5) for reading the data files from the local repositories. **Figure 2** shows a dynamic spectrum of calibrated Cassini/RPWS/HFR data during the Jupiter flyby on December 31<sup>st</sup> 2000 and January 1<sup>st</sup> 2001, using Autoplot and accessing the data through the MASER/LESIA *das2* server.

#### 4.2. Data discovery interface

VESPA is providing a data discovery framework with a metadata dictionary, a query protocol and a registry of services. Each VESPA service consists in a metadata table following the EPNcore metadata dictionary (Erard 2018). Each row contains the metadata corresponding to a single product, including a data access URL. The VESPA services are running over the Table Access Protocol (TAP, http://www.ivoa.net/documents/TAP/) from the IVOA.

MASER teams are currently sharing data files (Raw, CDF or FITS formats) through VESPA. The VESPA services are built upon the Data Centre Helper Suite (DaCHS, http://dachs-doc.readthedocs.io) framework,



**Figure 2:** Time-Frequency spectrogram of radio emissions observed by Cassini/RPWS/HFR during the Jupiter flyby on December 31<sup>st</sup> 2000 and January 1<sup>st</sup> 2001, accessed through Autoplot and the MASER das2server interface.

and the tables are fed directly from reading the CDF or FITS headers. The VESPA main query portal (http:// vespa.obspm.fr) also includes capabilities to interact directly with *Autoplot* (with the Simple Application Messaging Protocol (SAMP, http://www.ivoa.net/documents/SAMP/) of IVOA).

Several TAP servers dedicated to distributing MASER VESPA catalogue tables are available (http://voparistap-maser.obspm.fr at ObsParis, http://vogate.obs-nancay.fr in Nançay). They serve data collections from space mission with radio instruments (Cassini, Voyager, STEREO), from ground instruments (NDA) or modelled data (ExPRES, see section 6).

### 4.3. Run-on-demand interface

The IVOA has developed a computing job management system called Universal Worker Service (UWS, http://www.ivoa.net/documents/UWS/). MASER has implemented an instance of the Observatoire de Paris UWS System (OPUS, https://uws-server.readthedocs.io/en/latest/) available at https://voparis-uws-maser.obspm.fr. This server allows to submit jobs on a local computing cluster, either for automated data production pipelines (through a command line scripting interface), or for external users (through the web interface).

## 5. Maser4py Library

The *maser4py* library (https://github.com/maserlib/maser4py/ is providing data reader modules (for Python 3.6 and up) for legacy and non-standard format radio data collections. It currently includes modules for data collections hosted or produced by LESIA (Cassini/RPWS, Voyager/PRA, Solar Orbiter/RPW), by the Centre de Données de la Physique des Plasmas (CDPP, http://cdpp.eu) (Demeter, Interball, Viking (Swedish auroral mission), ISEE3, Wind), by the Planetary Plasma Interaction (PPI) node of NASA/PDS (Cassini/RPWS, Voyager/PRA), by the Nançay radio telescopes (NDA, NenuFAR), as well as by the radio amateur RadioJOVE project. It also includes generic modules developed for the ground segment of Solar Orbiter/RPW and a query interface for the HELIO-HFC (Heliophysics Integrated Observatory Feature Catalog) (Bonnin 2013). The *maser4py* library is open-source (GPLv3 license).

# 6. Modelling

The *Exoplanetary and Planetary Radio Emission Simulator* (ExPRES) code computes the geometric visibility of modelled auroral planetary radio source (Louis 2017, 2019). ExPRES is based on the Cyclotron Maser Instability (CMI) theory. It needs a planetary magnetic field model as well as parameters of the particle distributions in the modelled radio source. The code outputs time-frequency arrays of the visible auroral planetary radio source parameters (3D locus in the selected planetary frame and other radio source parameters). It is now used routinely to produce modelled daily spectrograms of simulated radio emissions induced by the Jovian Galilean satellites, for various observatory locations (Juno, Earth, STEREO). Precomputed simulation runs

are available are available through different interfaces as defined in the MASER Data Management Plan (see Louis (2019) for more details).

ExPRES is open-source and its code is available at: https://github.com/maserlib/ExPRES. Run-on-demand is also available from the MASER OPUS server (see section 2.3). This computing interface requires an ExPRES JSON input configuration file. Examples of such configuration files are available through the web directory listing or virtual observatory catalogue: each of the precomputed file is provided with its input configuration file. The JSON input files must comply with the ExPRES JSON-schema specification, the current version of which is available at https://voparis-ns.obspm.fr/maser/expres/v1.0/schema#. **Figure 3** shows the run-on-demand web interface where the user can manage his jobs. **Figure 4** shows a simulation run compared to Juno/Waves data (courtesy of C. Louis).

We also plan to distribute the electromagnetic ray tracing code ARTEMIS-P (Gautier 2013) through MASER and UWS.

## 7. Applications

The usage of *das2* server interface with *Autoplot* improves data analysis and processing for low frequency radio astronomy. Within MASER, a few examples can be cited.

- The refurbishment of Voyager/PRA data (Cecconi 2017) has been consolidated by the *das2* server/*Autoplot* setup, allowing efficient and fast data browsing at all temporal scales;
- Distribution of low frequency data sets together with space observations (Lamy 2017);



Figure 3: MASER public run-on-demand interface. A few ExPRES runs are shown here. The user can manage his own jobs.



**Figure 4:** Comparison of ExPRES simulation (top) and Juno/Waves data (bottom). Time-frequency spectrograms of Jovian radio emissions controlled by Io. Figure published in Louis (2017) as Supporting Information.

- The NenuFAR (Zarka 2012) instrument is in commission phase in Nançay, and the team is testing VESPA as an internal data catalogue and *das2* server for fast data access;
- Juno-Ground-Radio (Cecconi 2016) is aggregating ground-based radio data from several observatories (France, USA, Ukraine, Japan, Poland...) and provides data supporting the Juno science team. The data files are distributed through VESPA, using CDF files when possible. *Das2* server interfaces are under study for collaborators in Ukraine and Poland.

## 8. Future Steps

New data readers will be continuously included in the *maser4py* library. The MASER team will also reach out to the community for participation. This requires a consolidation of the *maser4py* interfaces (classes and methods) and tests.

The ExPRES simulations are now used by the Juno/Waves instrument team. Discussions are ongoing with ESA, for using ExPRES as an observation planning support tool for the JUICE mission.

The need for a radio ground support has also been identified by the Solar Orbiter and Parker Solar Probe teams. The MASER tools and data collections are already available and serve those needs.

Finally, there is a growing need for community coordinated open source library and software developments (especially for python-based developments). Several groups are pushing for this, and MASER will participate to these efforts (e.g., http://openplanetary.co for planetary sciences; or the PyHC working group, http://heliopython.org).

In addition to the VESPA access and the *das2* server interface, we will follow the International Heliophysics Data Environment Alliance (https://ihdea.net) recommendation to implement HAPI (Heliophysics API) interfaces (Vandegriff 2018).

#### Acknowledgements

The Europlanet H2020 Research Infrastructure project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 654208. The Europlanet-2024 Research Infrastructure project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 871149. Support from Paris Astronomical Data Centre (PADC) is acknowledged. The teams also received support from Observatoire de Paris, CNES and CNRS/INSU through ASOV.

### **Competing Interests**

The authors have no competing interests to declare.

#### References

Bonnin, X, et al. 2013. HFC Web Services. DOI: https://doi.org/10.5281/zenodo.3557465

- Briand, C, et al. 2016. STEREO database of interplanetary Langmuir electric waveforms. J. Geophys. Res. Space Physics, 121: 1062–1070. DOI: https://doi.org/10.1002/2015JA022036
- **Cecconi, B,** et al. 2016. Sharing Low Frequency Radio Emissions in the Virtual Observatory: Application for JUNO-Ground-Radio Observations Support. *Presented at JpGU 2016*. MGI04: 04. DOI: https://doi.org/10.5281/zenodo.3557279
- **Cecconi, B,** et al. 2017. Re-processing and re-analysis of Planetary Radio Emission (PRA) of Voyager 1 & 2. In: *Planetary, Solar and Heliospheric Radio Emissions (PRE 8).* Graz, Austria. DOI: https://doi.org/10.5281/zenodo.3557372
- **Cecconi, B,** et al. 2018. MASER (Measuring Analyzing & Simulating Emissions in Radio frequencies), a Toolbox for Low Frequency Radio Astronomy. In: *AGU Fall Meeting 2018 posters*. Washington DC, USA. DOI: https://doi.org/10.1002/essoar.10500145.1
- **Erard, S,** et al. 2018. VESPA: A Community-Driven Virtual Observatory in Planetary Science. *Planet. Space Sci.*, 150: 65–85. DOI: https://doi.org/10.1016/j.pss.2017.05.013
- Faden, J, et al. 2010. Autoplot: A Browser for Scientific Data on the Web. *Earth Space Sci. Inform.*, 3: 41–49. DOI: https://doi.org/10.1007/s12145-010-0049-0
- **Gautier, A-L,** et al. 2013. ARTEMIS-P: A General Ray Tracing Code in Anisotropic Plasma for Radioastronomical Applications. In: *Proceedings of the 2013 International Symposium on Electromagnetic Theory*. Hiroshima, Japan.
- King, T and Mafi, J. 2018. Guide to Archiving CDF Files in PDS4. DOI: https://doi.org/10.21978/P8WK8R

- Lamy, L, et al. 2017. 1977–2017: 40 Years of Decametric Observations of Jupiter and the Sun with the Nançay Decameter Array. In: *Planetary, Solar and Heliospheric Radio Emissions (PRE 8)*. Graz, Austria.
- Louis, CK, et al. 2017. Io-Jupiter Decametric Arcs Observed by Juno/Waves Compared to ExPRES Simulations. *Geophys. Res. Lett.*, 44(18): 9225–9232. DOI: https://doi.org/10.1002/2017GL073036
- Louis, CK, et al. 2019. ExPRES: a Tool to Simulate Exoplanetary and Planetary Radio Emissions. *Astron. Astrophys.*, 627: A30. DOI: https://doi.org/10.1051/0004-6361/201935161
- Marques, MS, et al. 2017. Statistical Analysis of 26 Years of Observations of Decametric Radio Emissions from Jupiter. *Astron. Astrophys.*, 604: A17. DOI: https://doi.org/10.1051/0004-6361/201630025
- Pence, WD, et al. 2010. Definition of the Flexible Image Transport System (FITS), Version 3.0. Astron. Astrophys., 524: A42. DOI: https://doi.org/10.1051/0004-6361/201015362
- Piker, C, et al. 2017. Das2 server Interface Control Document. DOI: https://doi.org/10.5281/ zenodo.3588535
- **Piker, C,** et al. 2018. Lightweight Federated Data Networks with Das2 Tools. *AGU Fall Meeting 2018 posters*. Washington DC, USA. DOI: https://doi.org/10.1002/essoar.10500359.1
- **Vandegriff, J,** et al. 2018. Keeping Everyone HAPI: Achieving Interoperability for Heliophysics and Planetary Time Series Data. In: *AGU Fall Meeting 2018 Posters*. Washington DC, USA. DOI: https://doi.org/10.1002/essoar.10500433.1
- Zarka, P. 1998. Auroral radio Emissions at the Outer Planets: Observations and Theories. J. Geophys. Res., 103: 20159–20194. DOI: https://doi.org/10.1016/0273-1177(92)90383-9
- Zarka, P, et al. 2012. LSS/NenuFAR: The LOFAR Super Station project in Nançay. *SF2A-2012: Proc. Annual meeting of the French Society of Astronomy and Astrophysics*. Boissier, S, de Laverny, P, Nardetto, N, Samadi, R, Valls-Gabaud, D and Wozniak, H (eds.), 687–694.

How to cite this article: Cecconi, B, Loh, A, Sidaner, PL, Savalle, R, Bonnin, X, Nguyen, QN, Lion, S, Shih, A, Aicardi, S, Zarka, P, Louis, C, Coffre, A, Lamy, L, Denis, L, Grießmeier, J-M, Faden, J, Piker, C, André, N, Génot, V, Erard, S, Mafi, JN, King, TA, Sky, J and Demleitner, M. 2020. MASER: A Science Ready Toolbox for Low Frequency Radio Astronomy. *Data Science Journal*, 19: 12, pp. 1–7. DOI: https://doi.org/10.5334/dsj-2020-012

Submitted: 06 December 2019 Accepted: 20 December 2019 Published: 18 March 2020

**Copyright:** © 2020 The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC-BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See http://creativecommons.org/ licenses/by/4.0/.



*Data Science Journal* is a peer-reviewed open access journal published by Ubiquity Press.

