



HAL
open science

Synthesis of Substituted α -Trifluoromethyl Piperidinic Derivatives

Sarah Rioton, Domingo Pardo, Janine Cossy

► **To cite this version:**

Sarah Rioton, Domingo Pardo, Janine Cossy. Synthesis of Substituted α -Trifluoromethyl Piperidinic Derivatives. *Molecules*, 2017, 22 (3), pp.483. 10.3390/molecules22030483 . hal-03230702

HAL Id: hal-03230702

<https://hal.science/hal-03230702v1>

Submitted on 9 May 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Review

Synthesis of Substituted α -Trifluoromethyl Piperidinic Derivatives

Sarah Rioton, Domingo Gomez Pardo and Janine Cossy *

Laboratoire de Chimie Organique, Institute of Chemistry, Biology and Innovation (CBI), CNRS UMR 8231, ESPCI Paris, PSL Research University, 10 rue Vauquelin, 75231 Paris CEDEX 05, France; sarah.rioton@espci.fr (S.R.); domingo.gomez-pardo@espci.fr (D.G.P.)

* Correspondence: Janine.cossy@espci.fr; Tel.: +33-140-794-429; Fax: +33-140-794-660

Academic Editor: Margaret A. Brimble CNZM, FRSNZ


Received: 2 March 2017; Accepted: 14 March 2017; Published: 19 March 2017

Abstract: A comprehensive survey of pathways leading to the generation of α -trifluoromethyl monocyclic piperidinic derivatives is provided (65 references). These compounds have been synthesized either from 6-membered rings e.g., pipercolic acid or lactam derivatives by introduction a trifluoromethyl group, from pyridine or pyridinone derivatives by reduction, and from 5-membered rings e.g., prolinol derivatives by ring expansion, from linear amines by cyclization or from dienes/dienophiles by [4 + 2]-cycloaddition.

Keywords: nitrogen heterocycles; fluorine; piperidine; trifluoromethyl group; ring expansion; cyclization; cycloaddition

1. Introduction

Functionalized piperidinic derivatives are among the most ubiquitous heterocyclic cores in natural products and bioactive compounds, therefore a huge number of methods has been developed to prepare piperidinic derivatives [1–7]. With respect to the biologically active targets, there is great interest in introducing substituents that can increase their biological activity which can be related to an increase of the lipophilicity, the bioavailability and the metabolic stability. In this context, the trifluoromethyl group is often used as a bioisostere of a chloride or a methyl group to modulate the steric and electronic properties of a lead compound or to protect a reactive methyl group from metabolic oxidation. This substituent can also increase the lipophilicity of molecules [8].



Scheme 1. Precursors of α -trifluoromethyl piperidinic derivatives.

Herein, we report the synthesis of α -trifluoromethylpiperidinic derivatives of type **A**. These compounds have been synthesized either from 6-membered rings **B** and **C** by introduction of a CF_3 group, from pyridines **D** or pyridinones **E** by reduction, from 5-membered rings **F** by ring expansion, from linear amines **G** by cyclization, from dienes/dienophiles **H/I** by cycloaddition (Scheme 1). We will only report the formation of monocyclic piperidinic derivatives, and of bicyclic derivatives only when they were transformed into the monocyclic derivatives.


2. From Cyclic Substrates

2.1. From 6-Membered Rings

The synthesis of α -trifluoromethylpiperidines **A** has been achieved from pipercolic acid, from δ -lactams, from pyridines and from pyridinones.

2.1.1. From Pipercolic Acid


The first synthesis of 2-(trifluoromethyl)piperidine (**2**) was realized in 1962 by Raash [9], from the sodium salt of pipercolic acid (**1**) that was treated with sulfur tetrafluoride (SF_4) in the presence of HF at 120°C . However, **2** was isolated in a very low yield of 9.6% (Scheme 2).



Scheme 2. Synthesis of 2-trifluoromethyl piperidine from pipercolic acid.

2.1.2. From 2-Trifluoromethylpyridine


One easy access to 2-trifluoromethylpiperidine (**2**) was realized by hydrogenation of the commercially available 2-trifluoromethylpyridine (**3**) in the presence of Pd, Pt or Rh catalysts (Scheme 3) [10–12].



Scheme 3. Hydrogenation of α -trifluoromethylpyridine.

2.1.3. From Trifluoromethyl Pyridinones


Access to trifluoromethylpiperidinones **5** from trifluoromethylpyridinones **4** has been developed by hydrogenation in the presence of Pd/C. Lactams **5** were isolated in 45%–88% yield and subsequent reduction should lead to the 5-amino 2-trifluoromethylpiperidines **6** (Scheme 4) [13].



Scheme 4. Hydrogenation of trifluoromethylpyridinones.

2.1.4. From δ -Lactams


Since the first access to 2-trifluoromethylpiperidine (**2**) different methods have been developed to produce α -trifluoromethylpiperidines of type **A** from δ -lactams **C** via either enamines **C1**, or imines **C2** or **C3** (Scheme 5).



Scheme 5. Synthesis of α -trifluoromethylpiperidines from δ -lactams.

From Enamines **C1**

α -Trifluoromethyl cyclic enamines **C1** can be good precursors of **A** as they are easily obtained from δ -lactams. When lactam **7** was treated with trichloroborane in the presence of an excess of CF_3Br and HEPT, enamine **8** was formed in 30% yield via the formation of iminium species **I** which can be attacked by the CF_3^- anion, generated from CF_3Br . Intermediate **II** was thus produced and after a β -elimination, enamine **8** was formed (Scheme 6) [14]. A reduction of this enamine should lead to 2-trifluoromethylpiperidine **9**. It is worth noting that CF_3Br is not easy to handle as it is a gas and alternatives to synthesize 2-trifluoromethylpiperidines from δ -lactams from imines **C2** and **C3** have been devised.




Scheme 6. Synthesis of α -trifluoromethylpiperidine from δ -lactam via enamine.

From Imines **C2**

Imines of type **C2** were synthesized in four steps from *N*-(diethoxymethyl)piperidin-2-one (**10**). First, **10** was subjected to a Claisen condensation with ethyl trifluoroacetate (NaH , $\text{CF}_3\text{CO}_2\text{Et}$) and,


after deprotection under acidic conditions (HCl), the corresponding fluorinated acyl lactam **11** was isolated as the hydrate, due to the ability of the electron-withdrawing CF_3 group to stabilize the tetrahedral adducts of such type of compounds [15].

The hydrolysis and the decarboxylation of perfluoro acyl lactam **11** under acidic conditions (6 M HCl) and heating at reflux led to **III**. When the decarboxylation was complete, the reaction was carefully made alkaline with KOH and hemiaminal **12** was obtained. After azeotropic removal of water or distillation, **12** was transformed into imine **13** (Scheme 7) [15].




Scheme 7. Synthesis of α -trifluoromethylpiperidine from δ -lactam via imine.

Imine **13** can be involved in a variety of reactions such as reduction, phosphorylation, alkylation under Friedel-Craft conditions, and Ugi-type reactions, producing a diversity of α -trifluoromethyl piperidinic derivatives. Imine **13** was transformed to 2-trifluoromethylpiperidine (**2**) in a yield of 71% by reduction with $\text{NaBH}(\text{OAc})_3$ in MeOH (Scheme 8) [15].




Scheme 8. Synthesis of α -trifluoromethylpiperidine from an imine by reduction.

When **13** was treated with diethyl phosphite in the presence of boron trifluoride etherate ($\text{BF}_3 \cdot \text{OEt}_2$) as the catalyst, 2-(trifluoromethyl)-2-ethylphosphonate piperidine (**14**) was obtained (92%) and easily transformed to the corresponding α -trifluoromethyl substituted cyclic α -aminophosphonic acid **15** by treatment with trimethylbromosilane in CHCl_3 followed by the addition of aqueous MeOH. It is worth pointing out that to avoid the formation of the ammonium salt, due to the liberation of HBr during the hydrolysis of the ester, the addition of propylene oxide was necessary and the free amino acid **15** was isolated (Scheme 9) [16].




Scheme 9. Synthesis of 2-(trifluoromethyl)-2-ethylphosphonate piperidine from imine.



Scheme 11. Synthesis of α -trifluoromethylpiperidines by Friedel-Craft reactions with pyrroles.


Indoles such **25** can also be involved in a Friedel-Craft reaction with imine **13** leading to the piperidines **26** in good yield (Scheme 12) [21].



Scheme 12. Synthesis of α -trifluoromethylpiperidines by a Friedel-Craft reaction with indoles.

From Imines C3

In imines **13**, the CF_3 group is initially located in the α -position and then different R substituents can also be introduced in the α -position. From an imine intermediate, the reverse strategy can be used to access compound **A**, e.g., imines are synthesized from a δ -lactam and then the trifluoromethyl group is introduced. Thus, when imines **27**, prepared from the corresponding δ -lactams, were treated with the Ruppert-Prakash reagent (TMSCF_3) in the presence of TFA and KHF_2 , in MeCN, the α,α -disubstituted piperidines **28** were isolated in good yields. The activation of imines **27** under acidic conditions was necessary to obtain piperidines **28** (Scheme 13) [22].




Scheme 13. Synthesis of α,α -disubstituted piperidines.

It is worth mentioning that all the 2-trifluoromethylpiperidines were obtained in a racemic form, however, recently, we have reported a method that allows the synthesis of optically active substituted 2-trifluoromethylpiperidines by using a stereospecific ring expansion applied to 2'-trifluoromethylprolinols of type F.

2.2. From 5-Membered Rings

The ring expansion of pyrrolidines to piperidines via an aziridinium intermediate under kinetic and thermodynamic conditions is well-established [23–28]. This method was used to access piperidines **30**, **30'** from prolinols **29**, **29'**.

Prolinols **29** and **29'** were prepared from L-proline in a seven-step sequence [29]. When prolinols **29** and **29'** were treated with DAST, I₂/PPh₃ or by Tf₂O, followed by the addition of different nucleophiles (alcohols, amines, thiols, cuprates, cyanides), a diversity of 2-(trifluoromethyl)-3-substituted piperidines **30** and **30'** were obtained with good diastereo- and enantioselectivities (de > 94% and ee > 99%). The regioselective attack of the nucleophiles on the aziridinium intermediates **VII**/**VII'** is due to the presence of the CF₃ group (Scheme 14) [29,30].



Scheme 14. Synthesis of C3-substituted α -trifluoromethylpiperidines from L-proline.

3. From Non-cyclic Substrates

To synthesize 2-trifluoromethyl-substituted piperidines from non-cyclic substrates, a cyclisation step or a cycloaddition is necessary to access the piperidine skeleton.

3.1. Cyclization

3.1.1. By Metathesis

Metathesis reactions have changed the way molecules are constructed due to the commercialization of robust catalysts such as the Grubbs first and second generation catalysts, **G-I** and **G-II**, and the Grubbs-Hoveyda catalysts **GH-II** (Figure 1). Metathesis is a powerful method to access both carbo- and heterocyclic ring systems [31–35] and the reaction is useful to synthesize piperidines and particularly to synthesize 2-trifluoromethylpiperidinic derivatives.





Figure 1. Catalysts for olefin metathesis.


From Dienes

The ammonium salt of the 2-trifluoromethylpiperidine **2·HCl** was formed from **36**, which was obtained from the α -trifluoroaminodiene **35** when treated with **G-I**. This diene has been synthesized in six steps from the hemiacetal **31** which was transformed into imine **32** by treatment with the imine of benzophenone followed by a silylation step (ImSiMe₃). After an allylation under acidic conditions (allylTMS, BF₃·OEt₂), followed by a deprotection/protection sequence, the allyltrifluoroamine **34** was isolated and allylated to produce the piperidinic core precursor **35**. Thus, after treatment of **35** with **G-I** catalyst (1 mol %) in dichloromethane at rt, the unsaturated piperidine **36** was isolated in 91% yield. After hydrogenation (Pd/C, EtOH, rt, 24 h) and treatment with HCl, **2·HCl** was isolated quantitatively (Scheme 15) [36,37]. Compared to the synthesis of **2** from pipercolic acid the synthesis of **2** from **31** is more efficient in terms of yield (47.9% versus 9.6%).




Scheme 15. Synthesis of α -trifluoromethylpiperidine from trifluoromethylhemiacetal.

A shorter synthesis of unsaturated piperidines from imines **37** using a Barbier-type reaction (allylBr, Zn, DMF, rt) followed by a *N*-allylation has been reported. The resulting dienic amino compound **38** was isolated in good yield. After treatment with **G-I** (5–10 mol %) the unsaturated piperidines **39** were isolated (Scheme 16) [38].



Scheme 16. Synthesis of α -trifluoromethylpiperidines from trifluoromethylimines.


Starting from imines, the strategy is very versatile to access a variety of unsaturated piperidines and, particularly, 2-(trifluoromethyl)-2-substituted tetrahydropyridines. Thus, when imines **40** were treated with allylmagnesium bromide (THF, $-78\text{ }^{\circ}\text{C}$) then allylated (NaH, allylbromide) dienes **42** were isolated in good yields and then involved in a ring-closing metathesis (RCM) using different catalysts such as **G-I** or the ruthenium catalyst **J** [39]. It is worth noting that the yields obtained with **G-I** are better than with catalyst **J** (Scheme 17) [40–43].




Scheme 17. Synthesis of α,α -disubstituted unsaturated piperidines from trifluoromethylimines.

It is worth noting that the dienes, precursors of 2-(trifluoromethyl)-unsaturated piperidines, could be synthesized from trifluorodiazido derivatives **44**. Treatment of *N,N*-diallyl-*N*-methylamine by the diazo compound **44** in the presence of the copper catalyst [Cu(F₃-acac)₂], produced ylide **VIII** which, after a [2,3]-sigmatropic rearrangement, led to the dienic derivatives **45**, precursor of the unsaturated piperidines **46** (Scheme 18) [43].

When the ring-opening metathesis (ROM)/ ring-closing metathesis (RCM) was applied to **47**, using the **G-I** catalyst (10 mol %), the unsaturated piperidines **48**, substituted at C4, were isolated in good yields (74%–86%) (Scheme 19) [44].




Scheme 18. Synthesis of α,α -disubstituted piperidines from trifluorodiazo derivatives.




Scheme 19. Synthesis of α -trifluoromethylpiperidines using a ROM/RCM sequence.

From Enynes

It is worth noting that a substituent at C4 could not be introduced on the piperidinic core by using a RCM applied to enyne **49** (Scheme 20, eq. 1). On the contrary, 2-(trifluoromethyl) unsaturated piperidines substituted at C5 can be produced by applying a RCM to enyne **51**. When this latter was treated with **G-I** (5 mol %) the resulting unsaturated piperidine **52** was obtained with an excellent yield (95%) (Scheme 20, eq. 2).



Scheme 20. Cont.



Scheme 20. Synthesis of α -trifluoromethylpiperidines by a RCM to enynes.

The alkyne can be substituted by different alkyl groups (compound **53**), aryl groups (compound **55**) and ketone (compound **57**). In all cases, the corresponding unsaturated piperidinic compounds substituted at C5, were isolated in good yields (Scheme 20, eqs. 3, 4 and 5) [37,38,45].

It is worth mentioning that in the case of **55**, when the aryl group is substituted in the *para* position the yields in **56** are good (60%–74%). On the contrary, when the aryl group is substituted in the *ortho* position, **55** is not transformed into **56** whatever the catalyst used, either the **G-II** or the **GH-II** catalysts [45].


3.1.2. By Cycloaddition

Only the cycloadditions leading to the monocyclic piperidinic core will be reported thus, the [2 + 2]-cycloaddition of allenyne [46] and the Paulson-Khan reaction [38,43,47] will be excluded.


[4 + 2]-Cycloaddition: Aza Diels-Alder

If imines are used as dienophiles, activation of the imine moiety either by electron-withdrawing substituents or by a Lewis acid is required for a successful [4 + 2]-cycloaddition. Imines **59**, possessing three electron-withdrawing groups, are very reactive and the aza Diels-Alder reaction was achieved at low temperature in the presence of different dienes **60**, to produce the corresponding unsaturated piperidines **61** in good yields. It is worth mentioning that phosphonylimines were less reactive than sulfonylimines as the reaction has to be performed at higher temperature and longer reaction times were required (120 h for phosphonylimines versus 14 h for sulfonylimines) (Table 1) [48].

When the silyloxydiene **62** was involved in the aza Diels-Alder reaction, piperidinone **63** was isolated after acidic treatment (Scheme 21) [48]


Table 1. Aza Diels-Alder cycloadditions.

Entry	R	R ¹	R ²	R ³	T (°C)	t (h)	Yield
1	SO ₂ Ph	Me	Me	H	-30 to 20	13	71%
2	SO ₂ Me	Me	Me	H	-30 to 20	14	91%
3	P(O)(OEt) ₂	Me	Me	H	20	120	95%
4	SO ₂ Ph	H	Me	H	-30 to 20	13	81%
5	SO ₂ Me	H	Me	H	-30 to 20	13	97%
6	P(O)(OEt) ₂	H	Me	H	20	150	68%
7	SO ₂ Ph	H	H	Me	4	120	75%
8	P(O)(OEt) ₂	H	H	Me	20	720	89%

**Scheme 21.** Synthesis of piperidinones by an aza Diels-Alder reaction.


Aza Diels-Alder Reaction with 1-Azadiene

The unsaturated piperidine **67** was obtained from an aza Diels-Alder reaction between the *iso*-butyl vinyl ether **66** and the sulfonimine intermediates **IX** generated *in situ* from the trifluoromethyl α,β -unsaturated ketone **64** and the sulfonamine **65**. In this aza Diels-Alder reaction, the HOMO orbital of **66** reacts with the LUMO orbital of the azadienes **IX**. The trifluoromethyl-piperidines **67** were obtained with a good diastereoselectivity in favor of the endo product. In addition, when the reaction was performed with an optically active sulfonamine, the cycloadduct was obtained with a good enantioselectivity (*ee* > 99%) (Scheme 22) [49].

**Scheme 22.** Aza Diels-Alder reaction with 1-azadienes.


3.1.3. 1,4-Addition: Aza-Michael

A Michael acceptor, tethered to a nucleophilic amine, can undergo an intramolecular 1,4-addition with concomitant formation of a piperidinic core. After the preparation of the optically active ω -amino α,β -unsaturated ketone **70** in three steps from the trifluoromethylhemiacetal **68**, the aminoketone **70** was treated under basic conditions to produce the 2-trifluoromethyl-substituted piperidine **71** in 80% yield and with an excellent diastereoselectivity of 94:6 in favor of the *cis*-isomer. This piperidine was transformed to an analog of pinidinol (Scheme 23) [50].



Scheme 23. Synthesis of α -trifluoromethylpiperidine by aza-Michael addition.


2-Trifluoromethylpiperidines with a quaternary center at C2 were synthesized from hemiaminal **72** and an α,β -unsaturated ketone **73** in the presence of a chiral amine ((2*S*,3*S*)-2-amino-*N*-((*S*)-1-hydroxy-3-phenyl-propan-2-yl)-3-methyl-pentanamide) under acidic conditions [*para*-nitrobenzoic acid (PNBA)]. Hemiaminal **72** reacts with the chiral amine to form intermediate **74** which then reacts with the enolized form of **73** to produce intermediate **75**. An intramolecular aza-Michael reaction then occurs and the tricyclic compounds **76** were isolated in good yields and with excellent *dr*(s) and *ee*(s). After reduction of the ketone (NaBH₄) and cleavage of the C–S and N–S bonds (sodium naphthalene), piperidine **77** (R¹ = Ph) was isolated with a good enantiomeric excess (*ee* = 92%) (Scheme 24) [51].



Scheme 24. Synthesis of chiral α -trifluoromethylpiperidines.

3.1.4. Electrophilic-Induced Cyclization


Amino-iodo cyclization has been used to prepare 2-trifluoromethylpipercolic acid derivatives starting from the ω -unsaturated imine **78**. The transformation of **79** to **80** was realized in a one-pot process with an excellent yield of 80%. After treatment of imine **78** with iodine in the presence of NaH, the bicyclic product **79** was formed and, when reacted with I_2 , a migration of the CF_3 group took place leading to the diiodo derivative **80** via an iminium alcoholate intermediate **XI**. The bicyclic compound **80** was then transformed into a variety of 2-trifluoromethylpipercolic acid derivatives **81** in a few steps (Scheme 25) [52,53].



Scheme 25. Synthesis of α -trifluoromethylpipercolic acid derivatives.

If in the electrophilic-induced cyclization of ω -aminoalkenes the CF_3 group was present in the starting material, on the contrary, in the electrophilic-induced cyclization of ω -amino alkynes the CF_3 was introduced after the cyclization.

When the ω -amino alkynes **82** were treated with AgF in the presence of H_2O , a hydroamination takes place to produce the enamine intermediate **XII**, which was reacted with $TMSCF_3$ to produce 2-trifluoromethylpiperidines **83** with excellent yields (88%–92%) (Scheme 26) [54].




Scheme 26. Synthesis of α -trifluoromethylpiperidines by electrophilic-induced cyclization.

3.1.5. Nucleophilic Attack of Iminium Ions


In reaction where iminium ions are involved, the iminium ion is attacked by a nucleophile such as in a Mannich type reaction and a Prins cyclization. A diastereoselective synthesis of 2-trifluoromethylpiperidines was realized by using an intramolecular Mannich reaction starting from the trifluoromethyl amine **84**. After condensation of the latter with aldehydes, the formed imines

85 were transformed to the iminium intermediates **XIII** under acidic conditions (*p*-TsOH, refluxing toluene). A cyclization is taking place to produce the 2-trifluoromethylpiperidinic derivatives **86**. The diastereoselectivity can be explained by the six-membered ring chair transition states **XIV**, in which the steric interactions are minimized (Scheme 27) [55–57].



Scheme 27. Synthesis of α -trifluoromethylpiperidines by an intramolecular Mannich reaction.


A silyl-aza-Prins reaction was also used to prepare highly functionalized 2-trifluoromethylpiperidines. Treatment of the vinyl silyl trifluoromethyl amine **87** with ethyl glyoxylate in the presence of InCl_3 led to **88** via the iminium intermediate **XV**. After an intramolecular attack of the iminium **XV** by the vinyl silane, intermediate **XVI** was formed, a desilylation took place and piperidine **88** was isolated in 65% yield. This piperidine can then be transformed to the highly functionalized 2-trifluoromethylpiperidine **89** (Scheme 28) [58].



Scheme 28. Synthesis of an α -trifluoromethylpiperidine by a silyl-aza-Prins reaction.


3.1.6. Intramolecular Condensation of Aminoketones and Aminoesters

Intramolecular lactamization and hemiaminalization were used to build up the C–N bond of the 2-trifluoromethylpiperidine derivatives. 2-Trifluoromethyl lactam **93**, which can lead to the corresponding piperidine by reduction, was synthesized in three steps from ketoester **90**. After treatment of **90** with aniline (*p*-TsOH, toluene, reflux), imine **91** was reduced to the δ -aminoester **92** which was cyclized to the corresponding lactam **93** under basic conditions (NaH , THF) (Scheme 29) [59].




Scheme 29. Synthesis of a trifluoromethyl lactam from a ketoester.

The α -substituted trifluoromethyl- δ -lactam **96** was prepared from the α -trifluoromethyl nitrosulfinamine **94**, prepared in two steps from trifluoromethylhemiacetal **68**. After a 1,4-addition of **94** to ethyl acrylate under basic conditions, the resulting aminoester **95** was treated with TFA to produce, after deprotection (NH_3 , MeOH), the trifluoromethyl- δ -lactam **96** (Scheme 30) [60].




Scheme 30. Synthesis of a trifluoromethyl lactam from a trifluoromethylhemiacetal.

A lactamization was also utilized to access 2-trifluoromethyl-3-hydroxypiperidine **102**. The latter was synthesized from aldimine **97** and the 2-trimethylsilyloxyfuran **98**, and transformed into the unsaturated lactone **99** after treatment with TBSOTf. After hydrogenation, the obtained lactone **100** was transformed to the δ -lactam **101** under acidic conditions (H_2SO_4 , 80%), and then to the 2,3-disubstituted piperidine **102** by reduction (LiAlH_4 , AlCl_3) (Scheme 31) [61].




Scheme 31. Synthesis of an α -trifluoromethylpiperidine by lactamization.

Due to the high reactivity of trifluoromethyl ketones they can be attacked by weak nucleophiles such as amines to form a hemiaminal. When the carboxylic trifluoromethyl ketones **103** were treated with ammonium carbonate in refluxing toluene, the six-membered ring hemiaminal intermediates **104** were formed and dehydrated under acidic conditions (*p*-TsOH, reflux) to furnish the unsaturated lactams **105** (Scheme 32) [62].



Scheme 32. Synthesis of unsaturated lactams from carboxylic trifluoromethyl ketones.


Highly substituted 2-trifluoromethylpiperidines **112** were obtained from isoxazoles **106**, trifluoromethylketoester **107**, aldehydes **108** and ammonium acetate in ethanol. In this multicomponent reaction, isoxazoles **106** react with the enolized β -ketoester **107**, to form intermediates **109** which can react with imines **110** (obtained by condensation of aldehydes **108** with NH_4OAc) to produce the aminotrifluoromethylketones **111**. The intramolecular attack of the amino group to the highly reactive trifluoromethylketones led to the trifluoromethylpiperidines **112** (Scheme 33) [63].



Scheme 33. Synthesis of highly substituted α -trifluoromethylpiperidines.

3.1.7. Intramolecular Nucleophilic Substitution


The intramolecular nucleophilic displacement of a good leaving group by an amine is one of the most common methods to prepare six-membered nitrogen heterocycles. This method was used to synthesize 2-trifluoromethylpiperidines **114** from the highly reactive aziridines **113**. The high reactivity of **113** is due to the presence of the trifluoromethyl group and the *N*-tosyl group. The attack of aziridine **113** by a variety of nucleophiles was very regioselective and produced intermediates **XVII** that cyclized according to an intramolecular nucleophilic substitution of the chloride to produce **114** (Scheme 34) [64].



Scheme 34. Synthesis of α -trifluoromethylpiperidines from trifluoromethylaziridines.

3.1.8. Intramolecular Nucleophilic Attack of Aldehydes

The annulation of aminoaldehyde **115** was realized by an intramolecular attack of a diphenyl sulfonium species on an aldehyde. When aminoaldehyde **115** was reacted with the trifluoromethylvinyl diphenylsulfonium **116**, the ylide intermediate **XVIII** was formed and a cyclization took place to produce **XIX** which led to the piperidino-epoxide **117**. It is worth noting that **117** was isolated with a good yield (72%) and a good diastereoselectivity (*dr* > 20:1) (Scheme 35) [65].



Scheme 35. Synthesis of α -trifluoromethylpiperidino-epoxides.

4. Conclusions

Despite the considerable synthetic efforts spent to synthesize 2-trifluoromethylpiperidinic derivatives, most of these compounds have been obtained in a racemic form. When they are optically active, the methods involve a chiral auxiliary or the starting material comes from the chiral pool. Thus, there is still a strong demand for catalytic enantioselective methods to efficiently access 2-trifluoromethylpiperidine derivatives in high diastereo- and enantioselectivity.

Acknowledgments: One of us, S. R. thanks GSK for a grant.

Author Contributions: All authors were involved in the preparation of the manuscript: The articles were collected by S. R., the organization and the writing of the manuscript were realized by J. C. and the typing by D. G. P.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Laschat, S.; Dickner, T. Stereoselective synthesis of piperidines. *Synthesis* **2000**, 1781–1813. [[CrossRef](#)]
- Toyooka, N.; Nemoto, H. Application of chiral building blocks to the synthesis of drugs. *Drugs Fut.* **2002**, *27*, 143–158. [[CrossRef](#)]
- Felplin, F.-X.; Lebreton, J. Recent advances in the total synthesis of piperidine and pyrrolidine natural alkaloids with ring-closing metathesis as a key step. *Eur. J. Org. Chem.* **2003**, 3693–3712. [[CrossRef](#)]
- Basra, S.; Blechert, S. Ring rearrangement metathesis—A new concept in piperidine and pyrrolidine synthesis. *Strategies Tactics Org. Synt.* **2004**, *4*, 315–346.
- Pearson, M.S.M.; Mathe-Allainmat, M.; Fargeas, V.; Lebreton, J. Recent advances in the total synthesis of piperidine aza-sugars. *Eur. J. Org. Chem.* **2005**, 2159–2191. [[CrossRef](#)]
- Vicario, J.L.; Badia, D.; Carrillo, L.; Ruiz, N.; Reyes, E. New methods for the asymmetric synthesis of piperidines and pyrrolidines: Chiral auxiliaries and asymmetric organocatalysis. *Targets Heterocycl. Syst.* **2008**, *12*, 302–327. [[CrossRef](#)]
- Amat, M.; Perez, M.; Bosch, J. Stereoselective conjugate addition reactions to phenylglycinol-derived, unsaturated oxazolopiperidone lactams. *Chem. Eur. J.* **2011**, *17*, 7724–7732. [[CrossRef](#)] [[PubMed](#)]
- Meyer, F. Trifluoromethyl nitrogen heterocycles: Synthetic aspects and potential biological targets. *Chem. Commun.* **2016**, *52*, 3077–3094. [[CrossRef](#)] [[PubMed](#)]
- Raasch, M.S. Chemistry of sulfur tetrafluoride. IX. Reaction with amino acids in hydrogen fluoride. *J. Org. Chem.* **1962**, *27*, 1406–1409. [[CrossRef](#)]
- Albrecht, B.K.; Gehling, V.C.; Harmange, J.-C.; Vaswani, R.G. Preparation of Substituted Indolecarboxamides as Modulators of Methyl Modifying Enzymes. WO 2016130396, 18 August 2016.
- Rice, K. Preparation of Benzoxazepines as PI3K/mTOR Inhibitors Useful in the Treatment of Cancer. WO 2012071509, 31 May 2012.

12. Rice, K.; Markby, D. Benzoxazepines as Inhibitors of PI3K/mTOR and Methods of their Use as Antitumor Agents and Manufacture. WO 2012068106, 24 May 2012.
13. Tolmachova, N.A.; Dolovanyuk, V.G.; Gerus, I.I.; Kondratov, I.S.; Polovinko, V.V.; Bergander, K.; Haufe, G. Catalytic hydrogenation of 3-amino-6-(trifluoromethyl)-5,6-dihydropyridin-2(1H)-ones and its use in the synthesis of trifluoromethyl-containing mimetics of ornithine and thalidomide. *Synthesis* **2011**, 1149–1156. [[CrossRef](#)]
14. Bürger, H.; Dittmar, T.; Pawelke, G. A simple one-pot synthesis of α -trifluoromethyl-substituted enamines by C-trifluoromethylation of dialkylamides with $P(\text{NEt}_2)_3/\text{CF}_3\text{Br}$. *J. Fluor. Chem.* **1995**, *70*, 89–93. [[CrossRef](#)]
15. Shevchenko, N.E.; Balenkova, E.S.; Röschenhaler, G.-V.; Nenajdenko, V.G. Practical synthesis of α -perfluoroalkyl cyclic imines and amines. *Synthesis* **2010**, 120–126. [[CrossRef](#)]
16. Odinets, I.L.; Artyushin, O.I.; Lyssenko, K.A.; Shevchenko, N.E.; Nenajdenko, V.G.; Röschenhaler, G.-V. Facile synthesis of cyclic α -perfluoroalkyl- α -aminophosphonates. *J. Fluor. Chem.* **2009**, *130*, 662–666. [[CrossRef](#)]
17. Gulevich, A.V.; Shevchenko, N.E.; Balenkova, E.S.; Röschenhaler, G.-V.; Nenajdenko, V.G. Efficient multicomponent synthesis of α -trifluoromethyl proline, homoproline, and azepan carboxylic acid dipeptides. *Synlett* **2009**, 403–406. [[CrossRef](#)]
18. Shmatova, O.I.; Nenajdenko, V.G. Tetrazole-substituted five-, six-, and seven-membered cyclic amines bearing perfluoroalkyl groups—efficient synthesis by azido-Ugi reaction. *Eur. J. Org. Chem.* **2013**, 6397–6403. [[CrossRef](#)]
19. Shmatova, O.I.; Shevchenko, N.E.; Balenkova, E.S.; Röschenhaler, G.-V.; Nenajdenko, V.G. Highly β -regioselective Friedel-Crafts aminoalkylation of pyrroles with cyclic perfluoroalkylated imines. *Eur. J. Org. Chem.* **2013**, 3049–3058. [[CrossRef](#)]
20. Shmatova, O.I.; Shevchenko, N.E.; Balenkova, E.S.; Röschenhaler, G.-V.; Nenajdenko, V.G. Friedel-Crafts alkylation of natural amino acid-derived pyrroles with CF_3 -substituted cyclic imines. *Mendeleev Commun.* **2013**, *23*, 92–93. [[CrossRef](#)]
21. Shevchenko, N.E.; Shmatova, O.I.; Balenkova, E.S.; Röschenhaler, G.-V.; Nenajdenko, V.G. Aminoalkylation of indoles with α -polyfluoroalkylated cyclic imines. *Eur. J. Org. Chem.* **2013**, 2237–2245. [[CrossRef](#)]
22. Shevchenko, N.E.; Vlasov, K.; Nenajdenko, V.G.; Röschenhaler, G.-V. The reaction of cyclic imines with the Ruppert-Prakash reagent. Facile approach to α -trifluoromethylated nornicotine, anabasine, and homoanabasine. *Tetrahedron* **2011**, *67*, 69–74. [[CrossRef](#)]
23. Cossy, J.; Gomez Pardo, D. Enantioselective ring expansion via aziridinium intermediates. Synthesis of substituted piperidines from substituted pyrrolidines. Synthetic applications. *Targets Heterocycl. Syst.* **2002**, *6*, 1–26. [[CrossRef](#)]
24. Cossy, J.; Gomez Pardo, D.; Dumas, C.; Mirguet, O.; Dechamps, I.; Métro, T.-X.; Burger, B.; Roudeau, R.; Appenzeller, J.; Cochi, A. Rearrangement of β -amino alcohols and application to the synthesis of biologically active compounds. *Chirality* **2009**, *21*, 850–856. [[CrossRef](#)] [[PubMed](#)]
25. Tynoshenko, D.D. Ring expansions of 1-azabicyclo[n.1.0]alkanes. Recent developments. *Arkivoc* **2011**, 329–345. [[CrossRef](#)]
26. Cochi, A.; Gomez Pardo, D.; Cossy, J. Access to optically active 3-aminopiperidines by ring expansion of prolinols: Thermodynamic versus kinetic control. *Eur. J. Org. Chem.* **2012**, 2023–2040. [[CrossRef](#)]
27. Gomez Pardo, D.; Cossy, J. Access to optically active 3-substituted piperidines by ring expansion of prolinols and derivatives. *Chem. Eur. J.* **2014**, *20*, 4516–4525. [[CrossRef](#)] [[PubMed](#)]
28. Dolfen, J.; Yadav, N.N.; De Kimpe, N.; D'hooghe, M.; Ha, H.-J. Bicyclic aziridinium ions in azaheterocyclic chemistry—Preparation and synthetic application of 1-azoniabicyclo[n.1.0]alkanes. *Adv. Synth. Catal.* **2016**, *358*, 3485–3511. [[CrossRef](#)]
29. Rioton, S.; Orliac, A.; Antoun, Z.; Bidault, R.; Gomez Pardo, D.; Cossy, J. Stereoselective rearrangement of (trifluoromethyl)prolinols to enantioenriched 3-substituted 2-(trifluoromethyl)piperidines. *Org. Lett.* **2015**, *17*, 2916–2919. [[CrossRef](#)] [[PubMed](#)]
30. Orliac, A. Synthèse et propriétés physico-chimiques des fluoro-pipéridines. XtalFluor-E et synthèse d'amides. Ph.D. Thesis, Université Pierre et Marie Curie, Paris, France, 13 November 2013.
31. Schmalz, H.-G. Catalytic ring-closing metathesis: a new, powerful technique for carbon-carbon coupling in organic synthesis. *Angew. Chem. Int. Ed.* **1995**, *34*, 1833–1836. [[CrossRef](#)]

32. Grubbs, R.H.; Miller, S.J.; Fu, G.C. Ring-closing metathesis and related processes in organic synthesis. *Acc. Chem. Res.* **1995**, *28*, 446–452. [[CrossRef](#)]
33. Schuster, M.; Blechert, S. Olefin metathesis in organic chemistry. *Angew. Chem. Int. Ed.* **1997**, *36*, 2037–2056. [[CrossRef](#)]
34. Armstrong, S.K. Ring-closing diene metathesis in organic synthesis. *J. Chem. Soc., Perkin Trans. 1: Org. Bio-Org. Chem.* **1998**, 371–388. [[CrossRef](#)]
35. Wright, D.L. Application of olefin metathesis to organic synthesis. *Curr. Org. Chem.* **1999**, *3*, 211–240.
36. Billard, T.; Langlois, B.R. Reactivity of stable trifluoroacetaldehyde hemiaminals. 2. Generation and synthetic potentialities of fluorinated iminiums. *J. Org. Chem.* **2002**, *67*, 997–1000. [[CrossRef](#)] [[PubMed](#)]
37. Gille, S.; Ferry, A.; Billard, T.; Langlois, B.R. Synthesis of α -trifluoromethylated nitrogen heterocycles. *J. Org. Chem.* **2003**, *68*, 8932–8935. [[CrossRef](#)] [[PubMed](#)]
38. Magueur, G.; Legros, J.; Meyer, F.; Ourévitch, M.; Crousse, B.; Bonnet-Delpon, D. A one-pot synthesis of doubly unsaturated trifluoromethyl amines: Easy access to CF_3 -substituted piperidines. *Eur. J. Org. Chem.* **2005**, 1258–1265. [[CrossRef](#)]
39. Fürstner, A.; Picquet, M.; Bruneau, C.; Dixneuf, P.H. Cationic ruthenium allenylidene complexes as a new class of performing catalysts for ring-closing metathesis. *Chem. Commun.* **1998**, 1315–1316. [[CrossRef](#)]
40. Osipov, S.N.; Bruneau, C.; Picquet, M.; Kolomiets, A.F.; Dixneuf, P.H. Synthesis of fluorine-containing cyclic amino acid derivatives via ring-closing olefin metathesis. *Chem. Commun.* **1998**, 2053–2054. [[CrossRef](#)]
41. Osipov, S.N.; Artyushin, O.I.; Kolomiets, A.F.; Bruneau, C.; Dixneuf, P.H. α - CF_3 -substituted phosphorus containing analogs of α -amino acids. Novel six- and seven-membered α -amino phosphonates via ring-closing metathesis with $\text{LnRu}=\text{C}=\text{C}=\text{CR}_2$ precatalyst. *Synlett* **2000**, 1031–1033. [[CrossRef](#)]
42. Osipov, S.N.; Artyushin, O.I.; Kolomiets, A.F.; Bruneau, C.; Picquet, M.; Dixneuf, P.H. Synthesis of fluorine-containing cyclic α -amino acid and α -amino phosphonate derivatives by alkene metathesis. *Eur. J. Org. Chem.* **2001**, 3891–3897. [[CrossRef](#)]
43. Vorobyeva, D.V.; Mailyan, A.K.; Peregodov, A.S.; Karimova, N.M.; Vasilyeva, T.P.; Bushmarinov, I.S.; Bruneau, C.; Dixneuf, P.H.; Osipov, S.N. Synthesis of functionalized CF_3 -containing heterocycles via [2,3]-sigmatropic rearrangement and sequential catalytic carbocyclization. *Tetrahedron* **2011**, *67*, 3524–3532. [[CrossRef](#)]
44. Osipov, S.N.; Kobel'kova, N.M.; Shchetnikov, G.T.; Kolomiets, A.F.; Bruneau, C.; Dixneuf, P.H. Novel synthesis of cyclic α -amino acid esters via ene reaction and ruthenium-catalyzed ring rearrangement. *Synlett* **2001**, 621–622. [[CrossRef](#)]
45. Mailyan, A.K.; Krylov, I.M.; Bruneau, C.; Dixneuf, P.H.; Osipov, S.N. Access to cyclic α - CF_3 -substituted α -amino acid derivatives by ring-closing metathesis of functionalized 1,7-enynes. *Eur. J. Org. Chem.* **2013**, 5353–5363. [[CrossRef](#)]
46. Mailyan, A.K.; Krylov, I.M.; Bruneau, C.; Dixneuf, P.H.; Osipov, S.N. Thermal [2 + 2]-cycloaddition of CF_3 -substituted allenyne: Access to novel cyclobutene-containing α -amino acids. *Synlett* **2011**, 2321–2324. [[CrossRef](#)]
47. Ferry, A.; Billard, T.; Langlois, B.R. Synthesis of α -trifluoromethylated nitrogen bicycles. *Synlett* **2005**, 1027–1029. [[CrossRef](#)]
48. Kobel'kova, N.M.; Osipov, S.N.; Kolomiets, A.F. Aza Diels-Alder reactions of methyl trifluoropyruvate sulfonyl- and phosphorylimines with 1,3-dienes. *Russ. Chem. Bull. Int. Ed.* **2002**, *51*, 1298–1302. [[CrossRef](#)]
49. Li, P.; Liu, L.-J.; Liu, J.-T. Per(poly)fluoroalkanesulfinamides assisted diastereoselective three-component inverse-electron-demand aza Diels-Alder reaction. *Org. Biomol. Chem.* **2011**, *9*, 74–77. [[CrossRef](#)] [[PubMed](#)]
50. Fustero, S.; Monteagudo, S.; Sanchez-Rosello, M.; Flores, S.; Barrio, P.; Del Pozo, C. *N*-Sulfinyl amines as a nitrogen source in the asymmetric intramolecular aza-Michael reaction: Total synthesis of (–)-pinidinol. *Chem. Eur. J.* **2010**, *16*, 9835–9845. [[CrossRef](#)] [[PubMed](#)]
51. Zhang, S.; Cha, L.; Li, L.; Hu, Y.; Li, Y.; Zha, Z.; Wang, Z. Asymmetric formal aza-Diels-Alder reaction of trifluoromethyl hemiaminals with enones catalyzed by primary amines. *J. Org. Chem.* **2016**, *81*, 3177–3187. [[CrossRef](#)] [[PubMed](#)]
52. Fustero, S.; Albert, L.; Aceña, J.L.; Sanz-Cervera, J.F.; Asensio, A. An Efficient entry to optically active anti- and syn- β -amino- α -trifluoromethyl alcohols. *Org. Lett.* **2008**, *10*, 605–608. [[CrossRef](#)] [[PubMed](#)]

53. Fustero, S.; Albert, L.; Mateu, N.; Chiva, G.; Miro, J.; Gonzalez, J.; Aceña, J.L. Stereoselective access to fluorinated and non-fluorinated quaternary piperidines: Synthesis of pipercolic acid and imino-sugar derivatives. *Chem. Eur. J.* **2012**, *18*, 3753–3764. [[CrossRef](#)] [[PubMed](#)]
54. Han, J.; Xu, B.; Hammond, G.B. Synthesis of α -CN and α -CF₃ N-heterocycles through tandem nucleophilic additions. *Org. Lett.* **2011**, *13*, 3450–3453. [[CrossRef](#)] [[PubMed](#)]
55. Bariau, A.; Roblin, J.-P.; Troin, Y.; Canet, J.-L. A simple stereoselective access to α -trifluoromethylated piperidines. *Synlett* **2005**, 1731–1733. [[CrossRef](#)]
56. Bariau, A.; Jatoi, W.B.; Calinaud, P.; Troin, Y.; Canet, J.-L. A simple stereoselective route to α -trifluoromethyl analogues of piperidine alkaloids. *Eur. J. Org. Chem.* **2006**, 3421–3433. [[CrossRef](#)]
57. Ciblat, S.; Besse, P.; Canet, J.-L.; Troin, Y.; Veschambre, H.; Gelas, J. A practical asymmetric synthesis of 2,6-cis-disubstituted piperidines. *Tetrahedron: Asymmetry* **1999**, *10*, 2225–2235. [[CrossRef](#)]
58. Dobbs, A.P.; Parker, R.J.; Skidmore, J. Rapid access to CF₃-containing heterocycles. *Tetrahedron Lett.* **2008**, *49*, 827–831. [[CrossRef](#)]
59. Wan, W.; Hou, J.; Jiang, H.; Yuan, Z.; Ma, G.; Zhao, G.; Hao, J. Reductive amination/cyclization of ω -trifluoromethyl keto esters to trifluoromethylated δ -amino alcohols and lactams. *Eur. J. Org. Chem.* **2010**, 1778–1786. [[CrossRef](#)]
60. García Ruano, J.L.; De Haro, T.; Singh, R.; Belén Cid, M. An efficient method for the synthesis of nitropiperidones. *J. Org. Chem.* **2008**, *73*, 1150–1153. [[CrossRef](#)] [[PubMed](#)]
61. Spanedda, M.V.; Ourévitich, M.; Crousse, B.; Bégué, J.-P.; Bonnet-Delpon, D. Vinylogous Mannich reactions. additions of trimethylsilyloxyfuran to fluorinated aldimines. *Tetrahedron Lett.* **2004**, *45*, 5023–5025. [[CrossRef](#)]
62. Okano, T.; Fumoto, M.; Kusukawa, T.; Fujita, M. Synthesis of optically active trifluoromethylated indolizidine derivatives via stereoselective radical cyclization. *Org. Lett.* **2002**, *4*, 1571–1573. [[CrossRef](#)] [[PubMed](#)]
63. Shi, W.; Wang, Y.; Zhu, Y.; Zhang, M.; Song, L.; Deng, H. A one-pot, multicomponent synthesis of trifluoromethylated spiropiperidines under catalyst-free conditions. *Synthesis* **2016**, *48*, 3527–3536.
64. Dolfen, J.; Kenis, S.; Van Hecke, K.; De Kimpe, N.; D'hooghe, M. Selective synthesis of functionalized trifluoromethylated pyrrolidines, piperidines, and azepanes starting from 1-tosyl-2-(trifluoromethyl)aziridine. *Chem. Eur. J.* **2014**, *20*, 10650–10653. [[CrossRef](#)] [[PubMed](#)]
65. Fritz, S.P.; West, T.H.; McGarrigle, E.M.; Aggarwal, V.K. Diastereoselective synthesis of CF₃-substituted, epoxide-fused heterocycles with β -(trifluoromethyl)vinyliulfonium salts. *Org. Lett.* **2012**, *14*, 6370–6373. [[CrossRef](#)] [[PubMed](#)]



© 2017 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).