



HAL
open science

The course of bipolar disorder as a function of the presence and sequence of onset of comorbid alcohol use disorders in outpatients attending the Fundamental Advanced Centres of Expertise

Romain Icick, Sébastien Gard, Katia M'Bailara, Isabelle Biseul, Ludovic Samalin, Georges Brousse, Valentin Flaudias, Pierre Michel Llorca, Joséphine Loftus, Iréna Cussac, et al.

► To cite this version:

Romain Icick, Sébastien Gard, Katia M'Bailara, Isabelle Biseul, Ludovic Samalin, et al.. The course of bipolar disorder as a function of the presence and sequence of onset of comorbid alcohol use disorders in outpatients attending the Fundamental Advanced Centres of Expertise. *Journal of Affective Disorders*, 2021, 287, pp.196-203. 10.1016/j.jad.2021.03.041 . hal-03230874

HAL Id: hal-03230874

<https://hal.science/hal-03230874>

Submitted on 24 Apr 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **The course of bipolar disorder as a function of the presence and sequence of onset of**
2 **comorbid alcohol use disorders in outpatients attending the Fondamental Advanced**
3 **Centres of Expertise**

4 Romain Icick^{1,2,3*}, Sébastien Gard^{1,4}, Katia M'Bailara^{1,4}, Isabelle Biseul^{1,2}, Ludovic Samalin^{1,5}, Georges
5 Brousse^{1,5}, Valentin Flaudias^{1,5}, Pierre-Michel Llorca^{1,5}, Joséphine Loftus^{1,6}, Iréna Cussac^{1,6}, Valérie
6 Aubin^{1,6}, Raymund Schwan⁷, Paul Roux^{1,8}, Mircea Polosan^{1,9}, Philippe Courtet^{1,10}, Emilie Olié^{1,10},
7 Chantal Henry¹¹, Nicolas Mazer^{1,12}, Emmanuel Haffen^{1,13}, Bruno Etain^{1,2,3}, Marion Leboyer^{1,14}, Frank
8 Bellivier^{1,2,3}, Raoul Belzeaux^{1,15}, Ophélie Godin^{1,14}, Sébastien Guillaume^{1,10}, The BIPADD group of FACE,
9 FondaMental Advanced Centres of Expertise in Bipolar Disorders (FACE-BD) Collaborators.

10

11 ¹Fondation Fondamental, Créteil, France.

12 ²AP-HP, GH Saint-Louis - Lariboisière - Fernand Widal, DMU Neurosciences, Département de
13 psychiatrie et de Médecine Addictologique, Paris, France. Fédération Hospitalo-Universitaire 'FHU
14 NOR-SUD'.

15 ³Université de Paris, UMR-S 1144, Paris, France.

16 ⁴Hôpital Charles Perrens, Centre Expert Trouble Bipolaire, Pôle de Psychiatrie Générale et
17 Universitaire (3/4/7), Bordeaux, France.

18 ⁵Department of Psychiatry, CHU Clermont-Ferrand, University of Clermont Auvergne, EA 7280,
19 Clermont-Ferrand, France;

20 ⁶Pôle de Psychiatrie, Centre Hospitalier Princesse Grace, Monaco.

21 ⁷Université de Lorraine, Inserm U1114, Centre Psychothérapique de Nancy, Nancy, France.

22 ⁸Service Universitaire de Psychiatrie d'Adultes et d'Addictologie, Centre Hospitalier de Versailles, Le
23 Chesnay, Université Paris-Saclay, UVSQ, Inserm, CESP, Team "DevPsy", 94807, Villejuif, France.

24 ⁹Université Grenoble Alpes, CHU de Grenoble et des Alpes, Grenoble Institut des Neurosciences (GIN)
25 Inserm U 1216, Grenoble, France.

26 ¹⁰Department of Emergency Psychiatry and Acute Care, Lapeyronie Hospital, CHU Montpellier,
27 Montpellier, France; PSNREC, Univ Montpellier, INSERM, CHU Montpellier, Montpellier, France.

28 ¹¹Department of Psychiatry, Service Hospitalo-Universitaire, GHU Paris Psychiatrie & Neurosciences,
29 F-75014, Paris, France.

30 ¹²AP-HP, Groupe Hospitalo-universitaire Nord, DMU ESPRIT, service de Psychiatrie et Addictologie.
31 Hopital Louis Mourier, Colombes, Inserm U1266, Faculté de médecine, Université de Paris, France.

32 ¹³ Service de Psychiatrie de l'Adulte, Centres Experts FondaMental, Centre Investigation Clinique
33 1431-INSERM, EA 481 Neurosciences, Université de Franche Comté, Besançon, France.

34 ¹⁴Paris Est Créteil, Inserm U955, IMRB, Laboratoire Neuro-Psychiatrie translationnelle, F-94010,
35 Créteil, France - AP-HP, HU Henri Mondor, Département Medico-Universitaire de Psychiatrie et
36 d'Addictologie (DMU ADAPT), Fédération Hospitalo-Universitaire de Médecine de Precision (FHU
37 IMPACT) F-94010, France.

38 ¹⁵Pôle de Psychiatrie, Assistance Publique Hôpitaux de Marseille, Marseille, France, INT-UMR7289,
39 CNRS Aix-Marseille Université, Marseille, France.

40 ***Corresponding author:** romain.icick@aphp.fr. CSAPA « Espace Murger », Département de
41 Psychiatrie et de Médecine Addictologique, Hôpital Fernand Widal, 200 rue du Faubourg Saint-Denis,
42 75010 Paris, France. Tel: +33140054275 / Fax: +33140054213.

43 **Abstract**

44 *Objectives:* The comorbidity of alcohol use disorder (AUD) and bipolar disorder (BD) has
45 been repeatedly associated with poorer clinical outcomes than BD without AUD. We aimed
46 to extend these findings by focusing on the characteristics associated with the sequence of
47 onset of BD and AUD. *Methods:* 3,027 outpatients from the Fundamental Advanced Centres
48 of Expertise were ascertained for BD-1, BD-2 and AUD diagnoses, including their respective
49 ages at onset (AAOs, N =2,804). We selected the variables associated with both the presence
50 and sequence of onset of comorbid AUD using bivariate analyses corrected for multiple
51 testing to enter a binary regression model with the sequence of onset of BD and AUD as the
52 dependent variable (AUD first - which also included 88 same-year onsets, vs. BD first).
53 *Results:* BD patients with comorbid AUD showed more severe clinical profile than those
54 without. Compared to BD-AUD (N =269), AUD-BD (N =276) was independently associated
55 with a higher AAO of BD (OR =1.1, $p < 0.001$), increased prevalence of comorbid cannabis use
56 disorder (OR =2.8, $p < 0.001$) a higher number of (hypo)manic/mixed BD episodes per year of
57 bipolar illness (OR =3, $p < 0.01$). *Limitations:* The transversal design prevents from drawing
58 causal conclusions. *Conclusion:* Increased severity of BD with AUD compared to BD alone did
59 not differ according to the sequence of onset. A few differences, though, could be used to
60 better monitor the trajectory of patients showing either one of these disorders.

61

62 **Keywords:** bipolar disorder; alcohol use disorder; dual diagnosis; comorbidity; sequence of
63 onset; clinical trajectory

64

65

66

67 **1. Introduction**

68 Bipolar disorder (BD) is a chronic and debilitating mental illness that affects ~2% of the
69 population worldwide (McIntyre et al., 2020). It is the 16th leading cause of disability (Ferrari
70 et al., 2016) and is associated with ~10 years of life lost (Ferrari et al., 2016). Alcohol use
71 disorders (AUDs) are the second most common addiction found in BD individuals after
72 tobacco use disorders (Blanco et al., 2017). BD increases the incidence of AUD as soon as
73 during adolescence (Birmaher et al., 2014), and individuals with AUD also show increased
74 incidence of BD (Bukh et al., 2016). According to numerous studies on the comorbidity
75 between BD and AUD, this condition is overall associated with poorer clinical outcomes than
76 BD alone, especially related to a more severe course of bipolar illness. More precisely, the
77 presence of comorbid AUD has been associated with increased risk of attempted and
78 completed suicide (SA) (Østergaard et al., 2017), increased number of depressive episodes
79 (Janiri et al., 2017), delayed remission from mood episodes (Farren et al., 2013) and
80 increased affective lability (Lagerberg et al., 2017).

81 A common clinical assumption is that people with BD drink alcohol to self-medicate anxiety
82 (Baigent, 2012), depressive symptoms (Meyer et al., 2012), irritability during mixed episodes
83 (Maremmani et al., 2012) and promote sleep (Lorberg et al., 2010), but also to further
84 increase mood during (hypo)manic (Meyer et al., 2012). The few longitudinal studies that
85 have been conducted so far have confirmed that the recovery from mood episodes is slower
86 in BD-AUD than in BD alone (Gold et al., 2018) and that AUD are preferentially associated
87 with bipolar depressive episodes (Simhandl et al., 2015), but rather argue against the self-
88 medication hypothesis in showing that increased alcohol use often precedes BD relapses
89 (Baethge et al., 2008; Strakowski et al., 2005). Interestingly in the study from Strakowski et
90 al., BD patients for whom AUD preceded BD had a quicker recovery from acute mood

91 episodes than both the BD-AUD and the no AUD groups, highlighting the importance of the
92 sequence of onset of BD and AUD in BD-AUD. A larger transversal study of 1090 BD-type 1
93 patients recruited after a manic episode (Azorin et al., 2017) reported that the group with
94 AUD beginning before BD (AUD-BD) showed higher rates of comorbid non-psychiatric,
95 anxiety and addictive disorders with a more frequent depressive polarity at onset than the
96 group where BD occurred before AUD (BD-AUD). The AUD-BD group was further
97 characterized by an irritable temperament.

98 The observed impact of lifetime AUD and its sequence of onset with BD on suicidal risk may
99 be related to factors modified by the comorbidity of AUD in BD (Hunt et al., 2016), such as
100 gender, bipolar subtype, impulsiveness (Etain et al., 2013), affective lability (Ducasse et al.,
101 2017), childhood trauma (Agnew-Blais and Danese, 2016), other comorbid addictions, or
102 decreased adherence to medication (Belzeaux et al., 2015). Extending the findings regarding
103 the relative onsets of BD and AUD using a large, representative and extensively
104 characterized BD sample could further our understanding of comorbid patients' trajectories.
105 This would pave the way to develop targeted prevention strategies aimed at individuals with
106 either BD or AUD, who are at higher risk of developing the comorbid disorder.

107 In order to identify risk factors differentiating bipolar disorder with vs. without alcohol use
108 disorder as a function of their sequence of onset, we analysed data from 2,804 outpatients
109 enrolled in the French network Fondamental Advanced Centres of Expertise – Bipolar
110 Disorders (FACE-BD). We hypothesized (i) that the presence of alcohol use disorder would be
111 associated with more severe clinical profiles of bipolar disorder compared to bipolar
112 disorder without alcohol use disorder but that (ii) several of these risk factors would depend
113 on the sequence of onset between bipolar disorder and alcohol use disorder.

114

115

116 **2. Material and methods**

117 *2.1. Clinical sample*

118 De-identified patient-level data from the FACE-BD national network were used for the
119 current study. The study was carried out in accordance with ethical principles for medical
120 research involving humans (WMA, Declaration of Helsinki). The assessment protocol was
121 approved by the relevant ethical review board (CPP-Ile de France IX, January 18th, 2010).
122 Although the committee waived the requirement for written informed consent, the patients
123 received a letter informing them of the study and asking whether they agreed to participate.
124 Briefly, patients with difficult-to-treat or difficult-to-diagnose BD are referred to one of the
125 12 centres of the network by general practitioners or psychiatrists. Once BD is ascertained,
126 patients undergo a two-days evaluation of their sociodemographic and clinical history by
127 trained nurses, psychologists and M.D.s in order to provide the treating physician and the
128 patient with adequate feedback regarding personalized therapeutic/diagnostic guidance.
129 The FACE-BD network procedures have been described in details elsewhere (Henry et al.,
130 2011). For the current study, we analysed data from 3,027 outpatients with ascertained BD
131 type 1 or 2; 2,804 of whom had available AUD diagnosis and AAOs of both BD and AUD,
132 enrolled between 2009 and August 2020 (date of data extraction from the centralized data
133 management system).

134 We focused on subsamples according to the presence and sequence of onset of BD and
135 AUD: “no AUD”, “BD-AUD” or “AUD-BD”. The “no AUD” group was defined by the absence of
136 lifetime AUD, the “BD-AUD” group by BD preceding AUD, the “AUD-BD” by AUD preceding
137 BD and the “BAUD” group by both disorders occurring the same year.

138 *2.2. Assessments*

139 Throughout the manuscript, the term substance use disorder (SUD) will refer to the
140 presence of abuse or dependence according to the *Diagnostic and Statistical Manual for*
141 *Mental disorders - fourth edition, text revised (DSM-IV, TR)* criteria (American Psychiatric
142 Association, 2000). Data regarding age, gender, educational attainment (years since
143 elementary school) and current living arrangement (living with partner, yes/no) and
144 employment status (currently employed >50%, yes/no) were selected for the current study.
145 History of BD and psychiatric comorbidity, including SUDs, was collected by the *Structured*
146 *Clinical Interview for DSM, axis I of the Diagnostic and Statistical Manual for Mental*
147 *Disorders, fourth edition (SCID-IV)* (Kübler, 2013), which has shown fair to good reliability in
148 SUD samples (Kranzler et al., 1996). Age at onset (AAO) of BD was defined as the age, at
149 which participants reported having their first major mood episode, which was further used
150 to compute BD duration (age at interview – AAO BD). AAO of AUD was considered the age,
151 at which participants first fulfilled DSM-IV criteria for AUD. Polarity at onset referred to the
152 same first BD episode, further dichotomized into “up” ((hypo)manic or mixed] vs. “down”
153 (depressive) categories. The number of mood episodes was also collected, dichotomized on
154 the same basis, and used as a rate per year (density of episodes). The presence of psychotic
155 symptoms during any mood episode occurring during the course of BD (hereafter termed
156 “history of psychosis”) was also collected.

157 Comorbid anxiety disorders, tobacco smoking status (never, former and current smokers)
158 and SUDs were considered on a lifetime basis. Disorders related to cocaine and other
159 stimulants were pooled into a “Stimulant use disorder” (StimUD) variable if any was present.
160 Current functioning was assessed by the *Functional Assessment Short Test (FAST)* (Rosa et
161 al., 2007) that was shown to reliably assess daily impairment in BD individuals. It consists of
162 a composite measure of six specific areas of functioning, of which we retained the total

163 score and the cognitive functioning subscale, due to the well-established links between AUD
164 and cognitive impairment in the general population (Koch et al., 2019). Higher scores
165 indicate poorer functioning. Current depressive symptoms were assessed with the
166 *Montgomery-Asberg Depression Rating Scale* (MADRS) (Montgomery and Asberg, 1979) and
167 current manic symptoms were measured with the *Young Mania Rating Scale* (YMRS) (Young
168 et al., 1978). Non-psychiatric medical history was collected as free text completed by
169 standardized questions about the main categories of diseases. For the current study, we
170 used dummy variables “any vs. none” non-psychiatric conditions from the following
171 categories: any, cardiovascular (hypertension, infarction, coronary disease, arrhythmia,
172 other), digestive (liver cirrhosis, gastric ulcer, drug hepatitis) conditions and dyslipidemia.
173 Body mass index (BMI) was calculated as weight (kg) / height (M²) and waist circumference
174 was measured in centimeters as a marker of abdominal obesity, following a standard
175 procedure (“WHO | Waist circumference and waist–hip ratio,” 2011). Both were actually
176 measured at the baseline visit. Trait impulsiveness was measured using the *Baratt*
177 *Impulsiveness Scale - 10th version, BIS10* and affective lability by the *Affective lability scale,*
178 *ALS* (Baylé et al., 2000). Childhood trauma was assessed by the *Childhood Trauma*
179 *Questionnaire – short form, CTQ* (Bernstein et al., 2003), using only the total score in order
180 to limit multiple testing and since CTQ was not the main focus of the study. These tools have
181 been validated in French (Baylé et al., 2000; Henry et al., 2008; Paquette et al., 2004) and
182 have been widely used in BD (Daruy-Filho et al., 2011; Etain et al., 2013; Lagerberg et al.,
183 2017).

184 Internal consistency was fair to good for the measurement scales, with standardized
185 Cronbach’s α =0.68 for MADRS, 0.72 for YMRS, 0.63 for BIS10, 0.61 for the FAST, 0.68 for the
186 CTQ and 0.57 for ALS.

187 2.3. *Statistical analyses*

188 Continuous variables are described by medians (interquartile range), given that most had
189 non-gaussian distributions (Shapiro-Wilk test p -value <0.05). Categorical data are described
190 by counts (frequency). Firstly, bivariate analyses were used to investigate associations
191 between the collected variables and the presence of lifetime AUD, using Mann-Whitney U
192 for continuous data or Chi^2 for categorical data. This was used to select the variables to test
193 with the sequence of onset of BD and AUD using bivariate analyses. Secondly, we
194 investigated associations of the collected variables and the sequences of onset of BD and
195 AUD, *i.e.*, BD first vs. AUD first. We used Mann-Whitney for continuous data or Chi^2 for
196 categorical data. Finally, a binary logistic regression model was ran with the variables
197 associated with the BD and AUD onset variable showing significance in both the bivariate
198 analyses. The regression model included a random term accounting for centre of inclusion
199 (12 sites). As a sensitivity analysis aimed at checking whether BD and AUD occurring the same
200 year better explained the results than our binary sequence variable, we also ran a
201 multinomial regression comparing BD first vs. AUD first vs. same year onsets with the
202 predictors identified in bivariate analysis. For bivariate analyses, statistical significance was
203 set at (i) $p < 0.05/36 = 0.001389$ (BD with vs. without AUD) and (ii) $p < 0.05/\text{number of}$
204 variables associated with the presence of AUD/2. Raw p -values are shown in all the tables,
205 with significance after Bonferroni correction indicated by*. Multicollinearity was assessed by
206 computing the variance inflation factor (VIF) of each independent variable. A VIF above the
207 “balanced” threshold of 2.5 is problematic and, in that case, the model was run again
208 without the variable, until all VIF values became satisfactory. R version 4.0.2 (R Core Team,
209 2020) through R studio version 1.3.1093 were used to compute statistics under Mac OS
210 X.14.6 system. The R session summary is available as a supplementary online file.

211

212 3. Results

213 The sample was composed of 3,027 outpatients with bipolar disorder. Comorbid AUD was
214 diagnosed in 768 (25%) patient. Mean age of the sample was 41 years (SD =13). There were
215 1,727 (62%) women and 1,351 (48%) patients with BD2 subtype. Other lifetime SUDs were
216 distributed as follows: 600 cannabis use disorders (CUD, 19%), 141 sedative use disorder
217 (SedUD, 5%) and 171 StimUD (6%). There were 545 patients with reliable data on BD and
218 AUD AAOs, including 269 (49%) with bipolar disorder first and AUD in second, 276 (51%)
219 with AUD first and bipolar in second (of which 88 developed BD and AUD the same year),
220 leaving a total sample for the sequence of onset study of 2,804 patients.

221 3.1. *Risk factors of comorbid AUD compared to BD without AUD, regardless of* 222 *their sequence of onset (Table 1)*

223 BD+AUD participants were more likely to be men (48% vs. 36%, $p < 0.001$), with earlier AAO
224 of BD (median =20 vs. 21, $p < 0.001$), higher prevalence of lifetime SA (48% vs. 35%, $p < 0.001$)
225 and of past-year mood episode (84% vs. 78%, $p < 0.001$) but lower prevalence of psychosis
226 history (32% vs. 40%, $p < 0.001$). They also showed increased current mood symptoms
227 according to both the MADRS (median =10 vs. 7) and the YMRS (median = 1 vs. 0) (both p
228 < 0.001), with poorer functioning according to FAST total (median =23 vs. 18, $p < 0.001$) and
229 cognition (mean =4.1 vs. 3.7, $p = 0.004$) scores. They also had a larger waist circumference
230 (median =93 vs. 91, $p < 0.001$) despite similar BMI ($p = 0.511$). They scored higher regarding
231 BIS10 (median =71 vs. 65), ALS (mean =1.5 vs. 1.2) and CTQ (median =20 vs. 21) total scores
232 (all $p < 0.001$). Finally, they exhibited higher rates of comorbid anxiety disorders (53% vs.
233 41%, $p < 0.001$) and lifetime SUDs (all $p < 0.001$): tobacco smoking (66% vs. 40%), sedUD (12%
234 vs. 2%), CUD (41% vs. 13%) and stimUD (16% vs. 2%).

235

236

Table 1 about here

237

238 3.2. *Factors associated with the sequence of onset of BD and AUD in comorbid*
239 *patients (Table 2)*

240 There were 21 significant associations with the presence of comorbid AUD. Thus, the
241 threshold for significance in the “sequence” analysis was $0.05/24 = 0.00208$. When
242 compared to the BD-AUD group, AUD-BD patients in the had a higher AAO of BD (21 vs. 18, p
243 <0.0001) but lower AAO of AUD (18 vs. 27, $p <0.0001$) and lower ALS score (1 vs. 2, p
244 <0.0001). They also reported a higher rate of “up” episodes per year of bipolar illness (0.45
245 vs. 0.32, $p <0.0001$) and increased prevalence of CUD (49% vs. 34%, $p <0.0001$).

246

247

Table 2 about here

248

249 3.3. *Binary logistic regression with the BD-AUD vs. AUD-BD groups as the*
250 *dependent variable (Figure 1)*

251 In the fully adjusted regression model, the sequence of onset of BD and AUD remained
252 independently associated with AAO of BD, which was lower in BD-AUD vs. AUD-BD patients
253 (OR =0.92, 95% CI =0.91-0.96 , $p =0.00002$), with the number of “up” BD episodes per year
254 of bipolar illness, which were less frequent in BD-AUD vs. AUD-BD (OR =0.3, 95% CI =0.22-
255 0.8, $p =0.01614$), with increased prevalence of past-year mood episode (OR =2, $p =0.04$) and
256 with the presence of CUD, which was less frequent in BD-AUD vs. AUD-BD patients (OR
257 =0.39, 95% CI =0.23-0.62, $p =0.00024$). The model was performed with 301 cases, yielding

258 area under curve (AUC) =0.73 (receiver operator curves are available by request from the
259 authors). There was no evidence of multicollinearity (maximum VIF =1.1 for CUD).

260 3.4. *Sensitivity analysis by multinomial regression with the sequence of onset of BD*
261 *and AUD in BD patients with a separate group for those, who began BD and AUD*
262 *during the same year (Supplementary Table 1 & Supplementary Figure 1)*

263 In the multinomial regression model, the significant results from the binary regression
264 remained overall similar as regards the comparison between BD-AUD and AUD-BD patients,
265 except that ALS score was lower in the AUD-BD group and that the rate of (hypo)manic
266 episodes/year of BD in patients, who developed BD and AUD during the same year,
267 appeared preferentially higher than in the BD first group (noticing that the difference with
268 the AUD-BD group yielded a p -value =0.06).

269

270 **Figure 1 about here**

271

272 **4. Discussion**

273 Using extensive data obtained through the comprehensive assessment of a large sample of
274 patients enrolled in the French national FACE-BD network, we first confirmed prior findings
275 that the comorbidity of BD with AUD was associated with a wide range of factors indicating
276 increased BD severity when compared to BD alone. A closer look at the sequence of onset
277 between BD and AUD did not alter this global impression of increased severity. However,
278 having AUD onset prior to BD onset in patients with BD+AUD comorbidity was
279 independently associated with a later AAO of BD, higher rate of “up” episodes per year of
280 bipolar illness and a increased prevalence of cannabis use disorder, as compared to having

281 BD onset prior to AUD onset. To our knowledge, this is the largest published study to date
282 assessing the risk factors of the relative onset of BD and AUD. We relied on standardized and
283 extensive assessment to eventually perform a fully-adjusted regression model. In particular,
284 BD and AUD diagnoses were properly ascertained and we could estimate the effects of key
285 clinical variables, including non-psychiatric comorbidity. The regression model had strong
286 prediction ability power (AUC =0.73) and confidence intervals for odds ratios were overall
287 narrow, indicating good reliability.

288 The first - and probably most expected - main finding was that patients with comorbid BD
289 and AUD had a more severe clinical profile across most of the majors areas known to impact
290 daily functioning and prognosis in BD, namely: current symptoms load, comorbid anxiety
291 disorders and SUD, impulsiveness and affective lability, childhood trauma. This extends on
292 prior literature emphasizing the poor clinical outcomes of individuals with BD and comorbid
293 AUD (Simhandl et al., 2015) and further highlights the need to design targeted and more
294 intensive care for patients with comorbid BD and AUD (Jones et al., 2019).

295 In the current study, we further aimed to characterize these patterns as a function of the
296 sequence of onset of BD and AUD in order to identify more clinically meaningful trajectories
297 that the sole increased severity of the comorbid condition. Firstly, compared to AUD-BD, BD-
298 AUD remained independently associated with an earlier AAO of BD. Individuals with
299 prodromal stages of BD likely drink alcohol prior to AUD diagnosis, attempting to self-
300 regulate prodromal mood symptoms but thereby increasing their risk for first depressive
301 episode (Turner et al., 2018). Since this episode will represent the index episode for dating
302 the beginning of BD, this could lead to both an earlier AAO of BD and increased incidence of
303 later AUD, the latter being associated with increased risk of conversion from unipolar
304 depression to BD (Musliner and Østergaard, 2018). Importantly, this particular finding has to

305 be interpreted in light of the respective ages at onset (AAOs) of BD and AUD in non-
306 comorbid individuals. AAO of BD usually stands around 20 years (Hasin and Grant, 2015),
307 while that of DSM-5 AUD is ~25 . In our sample, the median AAO of BD was 21 years old in
308 the AUD-BD group, which remained in the range of usual AAO of BD and supports the fact
309 that our findings related to the comparison of the BD-AUD and AUD-BD groups reflect at
310 least partially genuine trajectories of the comorbid BD+AUD condition. In the BD-AUD group,
311 AAO of BD was particularly early and that, in the AUD-BD group, AAO of AUD was early as
312 well. Therefore, the characteristics of these trajectories is likely to have been driven by the
313 earlier AAO of the disorder, which began first. In the largest previous study on this issue,
314 Azorin et al. also reported that BD AAO was higher in AUD-BD than in BD-AUD patients
315 (Azorin et al., 2017).

316 Secondly, BD-AUD was also independently associated with a lower number of “up” episodes
317 per year of illness and a lower prevalence of cannabis use disorder. We believe that this
318 reflects previously-reported associations between AUD and increased risk for depressive
319 episodes in BD (Janiri et al., 2017), whereas cannabis use and CUD have been linked to “up”
320 episodes (Lagerberg et al., 2016). In our sample, this is further supported by the fact that the
321 higher prevalence of past-year BD episodes (which shortly failed to reach significance in the
322 sequence analysis, Table 2) in BD first patients was driven by depressive episodes (data not
323 shown) and by a significant association between the presence of CUD and a higher rate of
324 “up” episodes (median =0.29 vs. 0.2, $p < 0.001$). Similar to our sample, there are also reports
325 that “AUD first” BD patients have increased prevalence of CUD (Hunt et al., 2016) compared
326 to BD first cases. In our sample, almost half (49%) of the AUD-BD patients had comorbid
327 CUD, that is, three comorbid SUDs if tobacco use disorder is included. Our tentative
328 interpretation is therefore that BD-AUD patients showing reduced prevalence of “up”

329 episodes due to lower prevalence of CUD. However, there was no difference between BD
330 with or without AUD regarding the rate of depressive episodes in our sample, so that
331 additional factors may be involved in these findings, warranting further investigation.

332 Interestingly, male gender was associated with the comorbid BD+AUD condition compared
333 to BD alone, regardless of the sequence of onset. A reduction in gender differences for
334 comorbid AUDs amongst individuals with BD had yet been previously evidenced (Frye et al.,
335 2003). The lack of association between BD-AUD sequences of onset and comorbid anxiety
336 disorder was unexpected. However, when investigating this issue in light of digestive system
337 disorders and dyslipidemia, we found that the former was almost twice as frequent in cases
338 with a positive history of anxiety disorders compared to those without (6% vs. 4%, $p < 0.001$).
339 These interactions should be investigated in a near future.

340 Finally, the sensitivity analysis aimed at estimating how the “same year” onset category of
341 the sequence variable could have biased our results elicited similar patterns of association as
342 with the dummy sequence variable (supplementary Table and Figure 1). However, a strong
343 separate difference emerged regarding the rate of (hypo)manic episodes per year of bipolar
344 illness, which was especially higher in the “same year” compared to the BD-AUD group. The
345 speed of transition between the disorders could shed further light on these associations.

346 *Limitations*

347 Our findings must be interpreted in light of several limitations. Data regarding lifetime
348 events were retrospective, thus subject to recall or classification bias. Recent substance use
349 was not included in our analyses, and excessive alcohol use - whether AUD is present or not -
350 has also been shown to alter BD course (Gordon-Smith et al., 2020). We merged the
351 category of BD and AUD onset when they occurred the same year with the BD-AUD
352 category, expecting that keeping all three categories would yield too small subgroups for

353 adequate statistical power. Overall, the prevalence of comorbid SUDs is rather lower in the
354 FACE-BD sample than in a recent meta-analysis of BD-AUD in clinical samples (Hunt et al.,
355 2016). We believe this might be due to the FACE-BD network selection bias that is both due
356 to how patients are referred to the network and to the euthymia criterion – a state that is
357 less frequent in BD comorbid with SUDs than without - that is applied so that patients are
358 able to undergo all assessments, which include full neuropsychological testing. The cross-
359 sectional nature of the study meant that we could not address issues of cause and effect.

360 Both the presence and the sequence of onset of the comorbidity between BD and AUD,
361 showed a wide range of associations with important clinical factors indicating increased
362 severity in comorbid patients. For most these factors, the sequence of onset of the disorders
363 was not associated with different association patterns, suggesting that AUD comorbidity in
364 BD has such a devastating effect on illness course that it overcomes the potential effect of its
365 sequence of onset with BD. Having a close look still highlighted that the sequence of onset of
366 BD and AUD had a few, yet relevant clinical implications with regards to AAO of BD, gender,
367 risk of (hypo)manic episodes and comorbidity with CUD. Importantly, our results overall
368 argue against previous findings that BD occurring after SUDs in general could represent a
369 less severe illness presentation as compared to BD alone (Pacchiarotti et al., 2009) (although
370 this may stand for substances, other than alcohol). This should encourage clinicians to
371 consider intensive care for all BD patients with AUD without focusing much on their
372 sequence of onset. Likewise, when considering young alcohol-using BD individuals, who may
373 not have developed AUD yet, full abstinence from alcohol (and possibly other addictive
374 substances as well) should be discussed during the stabilization phase of BD treatment.
375 Similar strategies should also be discussed in order to reduce the risk of developing BD in
376 alcohol-using individuals with prodromal BD symptoms.

377 **Acknowledgements**

378 The FondaMental Foundation (www-fondation-fondamental.org) promotes scientific
379 cooperation in mental health and that is developing a new model for translational research
380 in psychiatry in France. The FondaMental Foundation supports the infrastructure of Centres
381 of Expertise in BD. The authors also thank Hakim Laouamri, and his team (Seif Ben Salem,
382 Karmène Souyris, Victor Barteau, Stephane Beaufort and Mohamed Laaidi) for the
383 development of the FACE-BD computer interface (eBipolar©), data management, quality
384 control and regulatory aspects.

385 **Collaborators**

386 • The BIPolar-ADDiction group (BIPADD) of the Fondamental Advanced Centres of
387 Expertise in Bipolar Disorders (FACE-BD): France; Sébastien Gard and Katia M'Bailara –
388 Bordeaux; Valentin Flaudias, Ludovic Samalin and Georges Brousse - Clermont-Ferrand; Alix
389 de Prémorrel, Nicolas Mazer, Caroline Dubertret and Yann Le Strat- Colombes; Ophélie Godin,
390 Créteil; Sébastien Guillaume – Montpellier, Raoul Belzeaux - Marseille; Romain Icick and
391 Isabelle Biseul – Paris; Valérie Aubin and Joséphine Loftus - Monaco.

392 • FACE-BD Collaborators : B Etain, C Henry, E Olié, M Leboyer, E Haffen, P M Llorca, V
393 Barteau, S Bensalem, O Godin, H Laouamri, K Souryis, H Mondor, S Hotier, A Pelletier, N
394 Drancourt, J P Sanchez, E Saliou, C Hebbache, J Petrucci, L Willaume, E Bourdin, F Bellivier, M
395 Carminati, J Meheust, E Marlinge, M Meyrel, B Antoniol, A Desage, S Gard, A Jutant, K
396 Mbailara, I Minois, L Zanouy, L Bardin, A Cazals, P Courtet, B Deffinis, D Ducasse, M Gachet,
397 A Henrion, F Molière, B Noisette, E Olié, G Tarquini, R Belzeaux, N Correard, F Groppi, A
398 Lefrere, L Lescalier, E Moreau, J Pastol, M Rebattu, B Roux, N Viglianese, R Cohen, Raymund
399 Schwan, Thomas Schwitzer, Dr Grégory Gross, T Bougerol, B Fredembach, A Suisse, B Halili, A
400 Pouchon, M Polosan, A M Galliot, I Grévin, A S Cannavo, N Kayser, C Passerieux, , V Aubin, I

401 Cussac, M A Dupont, J Loftus, I Médecin, C Dubertret, N Mazer, C Portalier, C Scognamiglio,

402 A Bing

403 **Funding sources**

404 This research was supported by the Foundation FondaMental, Institut National de la Santé

405 et de la Recherche Médicale (INSERM), Assistance Publique des Hôpitaux de Paris (AP-HP),

406 and by the Investissements d’Avenir program managed by the Agence National de la

407 Recherche (ANR) under reference ANR-11-IDEX-0004-02 and ANR-10-COHO-10-01. The

408 funding sources had no role in the study design, data collection, analysis, preparation of the

409 manuscript, or decision to submit the manuscript for publication.

410

411 **References**

- 412 Agnew-Blais, J., Danese, A., 2016. Childhood maltreatment and unfavourable clinical
413 outcomes in bipolar disorder: a systematic review and meta-analysis. *The Lancet*
414 *Psychiatry* 3, 342–349. [https://doi.org/10.1016/S2215-0366\(15\)00544-1](https://doi.org/10.1016/S2215-0366(15)00544-1)
- 415 American Psychiatric Association, 2000. *The Diagnostic and Statistical Manual of Mental*
416 *Disorders, Fourth Edition-Text Revised (DSM IV-TR)*. American Psychiatric Association,
417 Washington, DC.
- 418 Azorin, J.-M., Perret, L.C., Fakra, E., Tassy, S., Simon, N., Adida, M., Belzeaux, R., 2017.
419 Alcohol use and bipolar disorders: Risk factors associated with their co-occurrence
420 and sequence of onsets. *Drug Alcohol Depend* 179, 205–212.
421 <https://doi.org/10.1016/j.drugalcdep.2017.07.005>
- 422 Baethge, C., Hennen, J., Khalsa, H.-M.K., Salvatore, P., Tohen, M., Baldessarini, R.J., 2008.
423 Sequencing of substance use and affective morbidity in 166 first-episode bipolar I
424 disorder patients. *Bipolar Disorders* 10, 738–741. <https://doi.org/10.1111/j.1399-5618.2007.00575.x>
- 425
- 426 Baigent, M., 2012. Managing patients with dual diagnosis in psychiatric practice. *Curr Opin*
427 *Psychiatry* 25, 201–205. <https://doi.org/10.1097/YCO.0b013e3283523d3d>
- 428 Baylé, F.J., Bourdel, M.C., Caci, H., Gorwood, P., Chignon, J.M., Adés, J., Lôo, H., 2000. [Factor
429 analysis of french translation of the Barratt impulsivity scale (BIS-10)]. *Can J*
430 *Psychiatry* 45, 156–165. <https://doi.org/10.1177/070674370004500206>
- 431 Belzeaux, R., Boyer, L., Mazzola-Pomietto, P., Michel, P., Correard, N., Aubin, V., Bellivier, F.,
432 Bougerol, T., Olie, E., Courtet, P., Etain, B., Gard, S., Kahn, J.-P., Passerieux, C.,
433 Leboyer, M., Henry, C., Azorin, J.-M., French Advanced Centers of Expertise for
434 Bipolar Disorders Collaborators, 2015. Adherence to medication is associated with
435 non-planning impulsivity in euthymic bipolar disorder patients. *J Affect Disord* 184,
436 60–66. <https://doi.org/10.1016/j.jad.2015.05.041>
- 437 Bernstein, D.P., Stein, J.A., Newcomb, M.D., Walker, E., Pogge, D., Ahluvalia, T., Stokes, J.,
438 Handelsman, L., Medrano, M., Desmond, D., Zule, W., 2003. Development and
439 validation of a brief screening version of the Childhood Trauma Questionnaire. *Child*
440 *Abuse Negl* 27, 169–190. [https://doi.org/10.1016/s0145-2134\(02\)00541-0](https://doi.org/10.1016/s0145-2134(02)00541-0)
- 441 Birmaher, B., Gill, M.K., Axelson, D.A., Goldstein, B.I., Goldstein, T.R., Yu, H., Liao, F., Iyengar,
442 S., Diler, R.S., Strober, M., Hower, H., Yen, S., Hunt, J., Merranko, J.A., Ryan, N.D.,
443 Keller, M.B., 2014. Longitudinal Trajectories and Associated Baseline Predictors in
444 Youths With Bipolar Spectrum Disorders. *AJP* 171, 990–999.
445 <https://doi.org/10.1176/appi.ajp.2014.13121577>
- 446 Blanco, C., Compton, W.M., Saha, T.D., Goldstein, B.I., Ruan, W.J., Huang, B., Grant, B.F.,
447 2017. Epidemiology of DSM-5 bipolar I disorder: Results from the National
448 Epidemiologic Survey on Alcohol and Related Conditions – III. *Journal of Psychiatric*
449 *Research* 84, 310–317. <https://doi.org/10.1016/j.jpsychires.2016.10.003>
- 450 Bukh, J.D., Andersen, P.K., Kessing, L.V., 2016. Rates and predictors of remission, recurrence
451 and conversion to bipolar disorder after the first lifetime episode of depression--a
452 prospective 5-year follow-up study. *Psychol Med* 46, 1151–1161.
453 <https://doi.org/10.1017/S0033291715002676>
- 454 Daruy-Filho, L., Brietzke, E., Lafer, B., Grassi-Oliveira, R., 2011. Childhood maltreatment and
455 clinical outcomes of bipolar disorder. *Acta Psychiatr Scand* 124, 427–434.
456 <https://doi.org/10.1111/j.1600-0447.2011.01756.x>

457 Ducasse, D., Jaussent, I., Guillaume, S., Azorin, J.M., Bellivier, F., Belzeaux, R., Bougerol, T.,
458 Etain, B., Gard, S., Henry, C., Kahn, J.P., Leboyer, M., Loftus, J., Passerieux, C.,
459 FondaMental Advanced Centers of Expertise in Bipolar Disorders (FACE-BD)
460 Collaborators, Olié, E., Courtet, P., 2017. Affect lability predicts occurrence of suicidal
461 ideation in bipolar patients: a two-year prospective study. *Acta Psychiatr Scand* 135,
462 460–469. <https://doi.org/10.1111/acps.12710>

463 Etain, B., Mathieu, F., Liquet, S., Raust, A., Cochet, B., Richard, J.R., Gard, S., Zanouy, L., Kahn,
464 J.P., Cohen, R.F., Bougerol, T., Henry, C., Leboyer, M., Bellivier, F., 2013. Clinical
465 features associated with trait-impulsiveness in euthymic bipolar disorder patients. *J*
466 *Affect Disord* 144, 240–247. <https://doi.org/10.1016/j.jad.2012.07.005>

467 Farren, C.K., Snee, L., Daly, P., McElroy, S., 2013. Prognostic Factors of 2-year Outcomes of
468 Patients with Comorbid Bipolar Disorder or Depression with Alcohol Dependence:
469 Importance of Early Abstinence. *Alcohol Alcohol* 48, 93–98.
470 <https://doi.org/10.1093/alcalc/ags112>

471 Ferrari, A.J., Stockings, E., Khoo, J.-P., Erskine, H.E., Degenhardt, L., Vos, T., Whiteford, H.A.,
472 2016. The prevalence and burden of bipolar disorder: findings from the Global
473 Burden of Disease Study 2013. *Bipolar Disord* 18, 440–450.
474 <https://doi.org/10.1111/bdi.12423>

475 Frye, M.A., Altshuler, L.L., McElroy, S.L., Suppes, T., Keck, P.E., Denicoff, K., Nolen, W.A.,
476 Kupka, R., Leverich, G.S., Pollio, C., Grunze, H., Walden, J., Post, R.M., 2003. Gender
477 Differences in Prevalence, Risk, and Clinical Correlates of Alcoholism Comorbidity in
478 Bipolar Disorder. *AJP* 160, 883–889. <https://doi.org/10.1176/appi.ajp.160.5.883>

479 Gold, A.K., Peters, A.T., Otto, M.W., Sylvia, L.G., Magalhaes, P.V. da S., Berk, M., Dougherty,
480 D.D., Miklowitz, D.J., Frank, E., Nierenberg, A.A., Deckersbach, T., 2018. The impact of
481 substance use disorders on recovery from bipolar depression: Results from the
482 Systematic Treatment Enhancement Program for Bipolar Disorder psychosocial
483 treatment trial. *Aust N Z J Psychiatry* 52, 847–855.
484 <https://doi.org/10.1177/0004867418788172>

485 Gordon-Smith, K., Lewis, K.J.S., Auñón, F.M.V., Florio, A.D., Perry, A., Craddock, N., Jones, I.,
486 Jones, L., 2020. Patterns and clinical correlates of lifetime alcohol consumption in
487 women and men with bipolar disorder: Findings from the UK Bipolar Disorder
488 Research Network. *Bipolar Disorders* 22, 731–738.
489 <https://doi.org/10.1111/bdi.12905>

490 Hasin, D.S., Grant, B.F., 2015. The National Epidemiologic Survey on Alcohol and Related
491 Conditions (NESARC) Waves 1 and 2: Review and summary of findings. *Soc Psychiatry*
492 *Psychiatr Epidemiol* 50, 1609–1640. <https://doi.org/10.1007/s00127-015-1088-0>

493 Henry, C., Etain, B., Mathieu, F., Raust, A., Vibert, J.-F., Scott, J., Leboyer, M., 2011. A French
494 network of bipolar expert centres: a model to close the gap between evidence-based
495 medicine and routine practice. *J Affect Disord* 131, 358–363.
496 <https://doi.org/10.1016/j.jad.2010.11.013>

497 Henry, C., Van den Bulke, D., Bellivier, F., Roy, I., Swendsen, J., M'Bailara, K., Siever, L.J.,
498 Leboyer, M., 2008. Affective lability and affect intensity as core dimensions of bipolar
499 disorders during euthymic period. *Psychiatry Research* 159, 1–6.
500 <https://doi.org/10.1016/j.psychres.2005.11.016>

501 Hunt, G.E., Malhi, G.S., Cleary, M., Lai, H.M.X., Sitharthan, T., 2016. Prevalence of comorbid
502 bipolar and substance use disorders in clinical settings, 1990-2015: Systematic review

503 and meta-analysis. *J Affect Disord* 206, 331–349.
504 <https://doi.org/10.1016/j.jad.2016.07.011>

505 Janiri, D., Di Nicola, M., Martinotti, G., Janiri, L., 2017. Who's the Leader, Mania or
506 Depression? Predominant Polarity and Alcohol/Polysubstance Use in Bipolar
507 Disorders. *Curr Neuropharmacol* 15, 409–416.
508 <https://doi.org/10.2174/1570159X14666160607101400>

509 Jones, S.H., Riste, L., Robinson, H., Holland, F., Peters, S., Hartwell, R., Berry, K., Fitzsimmons,
510 M., Wilson, I., Hilton, C., Long, R., Bateman, L., Weymouth, E., Owen, R., Roberts, C.,
511 Barrowclough, C., 2019. Feasibility and acceptability of integrated psychological
512 therapy versus treatment as usual for people with bipolar disorder and co-morbid
513 alcohol use: A single blind randomised controlled trial. *J Affect Disord* 256, 86–95.
514 <https://doi.org/10.1016/j.jad.2019.05.038>

515 Koch, M., Fitzpatrick, A.L., Rapp, S.R., Nahin, R.L., Williamson, J.D., Lopez, O.L., DeKosky, S.T.,
516 Kuller, L.H., Mackey, R.H., Mukamal, K.J., Jensen, M.K., Sink, K.M., 2019. Alcohol
517 Consumption and Risk of Dementia and Cognitive Decline Among Older Adults With
518 or Without Mild Cognitive Impairment. *JAMA Netw Open* 2, e1910319–e1910319.
519 <https://doi.org/10.1001/jamanetworkopen.2019.10319>

520 Kranzler, H.R., Kadden, R.M., Babor, T.F., Tennen, H., Rounsaville, B.J., 1996. Validity of the
521 SCID in substance abuse patients. *Addiction* 91, 859–868.

522 Kübler, U., 2013. Structured Clinical Interview for DSM-IV (SCID), in: Gellman, M.D., Turner,
523 J.R. (Eds.), *Encyclopedia of Behavioral Medicine*. Springer New York, New York, NY,
524 pp. 1919–1920. https://doi.org/10.1007/978-1-4419-1005-9_66

525 Lagerberg, T.V., Aminoff, S.R., Aas, M., Bjella, T., Henry, C., Leboyer, M., Pedersen, G.,
526 Bellivier, F., Ickick, R., Andreassen, O.A., Etain, B., Melle, I., 2017. Alcohol use disorders
527 are associated with increased affective lability in bipolar disorder. *J Affect Disord* 208,
528 316–324. <https://doi.org/10.1016/j.jad.2016.09.062>

529 Lagerberg, T.V., Ickick, R., Andreassen, O.A., Ringen, P.A., Etain, B., Aas, M., Henry, C., Bjella,
530 T.D., Melle, I., Bellivier, F., 2016. Cannabis use disorder is associated with greater
531 illness severity in tobacco smoking patients with bipolar disorder. *J Affect Disord* 190,
532 286–293. <https://doi.org/10.1016/j.jad.2015.10.023>

533 Lorberg, B., Wilens, T.E., Martelon, M., Wong, P., Parcell, T., 2010. Reasons for Substance
534 Use among Adolescents with Bipolar Disorder. *The American Journal on Addictions*
535 19, 474–480. <https://doi.org/10.1111/j.1521-0391.2010.00077.x>

536 Maremmani, I., Maremmani, A.G.I., Rugani, F., Rovai, L., Pacini, M., Bacciardi, S., Deltito, J.,
537 Dell'osso, L., Akiskal, H.S., 2012. Clinical presentations of substance abuse in bipolar
538 heroin addicts at time of treatment entry. *Ann Gen Psychiatry* 11, 23.
539 <https://doi.org/10.1186/1744-859X-11-23>

540 McIntyre, R.S., Berk, M., Brietzke, E., Goldstein, B.I., López-Jaramillo, C., Kessing, L.V., Malhi,
541 G.S., Nierenberg, A.A., Rosenblat, J.D., Majeed, A., Vieta, E., Vinberg, M., Young, A.H.,
542 Mansur, R.B., 2020. Bipolar disorders. *The Lancet* 396, 1841–1856.
543 [https://doi.org/10.1016/S0140-6736\(20\)31544-0](https://doi.org/10.1016/S0140-6736(20)31544-0)

544 Meyer, T.D., McDonald, J.L., Douglas, J.L., Scott, J., 2012. Do patients with bipolar disorder
545 drink alcohol for different reasons when depressed, manic or euthymic? *J Affect*
546 *Disord* 136, 926–932. <https://doi.org/10.1016/j.jad.2011.09.005>

547 Montgomery, S.A., Asberg, M., 1979. A new depression scale designed to be sensitive to
548 change. *Br J Psychiatry* 134, 382–389.

549 Musliner, K.L., Østergaard, S.D., 2018. Patterns and predictors of conversion to bipolar
550 disorder in 91 587 individuals diagnosed with unipolar depression. *Acta Psychiatrica*
551 *Scandinavica* 137, 422–432. <https://doi.org/10.1111/acps.12869>

552 Østergaard, M.L.D., Nordentoft, M., Hjorthøj, C., 2017. Associations between substance use
553 disorders and suicide or suicide attempts in people with mental illness: a Danish
554 nation-wide, prospective, register-based study of patients diagnosed with
555 schizophrenia, bipolar disorder, unipolar depression or personality disorder.
556 *Addiction* 112, 1250–1259. <https://doi.org/10.1111/add.13788>

557 Pacchiarotti, I., Di Marzo, S., Colom, F., Sánchez-Moreno, J., Vieta, E., 2009. Bipolar disorder
558 preceded by substance abuse: a different phenotype with not so poor outcome?
559 *World J. Biol. Psychiatry* 10, 209–216. <https://doi.org/10.1080/15622970701558488>

560 Paquette, D., Laporte, L., Bigras, M., Zoccolillo, M., 2004. Validation de la version française
561 du CTQ et prévalence de l’histoire de maltraitance. *smq* 29, 201–220.
562 <https://doi.org/10.7202/008831ar>

563 R Core Team, 2020. R: A language and environment for statistical computing. R Foundation
564 for Statistical Computing, Vienna, Austria.

565 Rosa, A.R., Sánchez-Moreno, J., Martínez-Aran, A., Salamero, M., Torrent, C., Reinares, M.,
566 Comes, M., Colom, F., Van Riel, W., Ayuso-Mateos, J.L., Kapczinski, F., Vieta, E., 2007.
567 Validity and reliability of the Functioning Assessment Short Test (FAST) in bipolar
568 disorder. *Clin Pract Epidemiol Ment Health* 3, 5. [https://doi.org/10.1186/1745-0179-](https://doi.org/10.1186/1745-0179-3-5)
569 [3-5](https://doi.org/10.1186/1745-0179-3-5)

570 Simhandl, C., Radua, J., König, B., Amann, B.L., 2015. Prevalence and impact of comorbid
571 alcohol use disorder in bipolar disorder: A prospective follow-up study: *Australian &*
572 *New Zealand Journal of Psychiatry*. <https://doi.org/10.1177/0004867415585855>

573 Strakowski, S.M., DelBello, M.P., Fleck, D.E., Adler, C.M., Anthenelli, R.M., Keck, P.E., Arnold,
574 L.M., Amicone, J., 2005. Effects of Co-occurring Alcohol Abuse on the Course of
575 Bipolar Disorder Following a First Hospitalization for Mania. *Arch Gen Psychiatry* 62,
576 851–858. <https://doi.org/10.1001/archpsyc.62.8.851>

577 Turner, S., Mota, N., Bolton, J., Sareen, J., 2018. Self-medication with alcohol or drugs for
578 mood and anxiety disorders: A narrative review of the epidemiological literature.
579 *Depress Anxiety* 35, 851–860. <https://doi.org/10.1002/da.22771>

580 WHO | Waist circumference and waist–hip ratio [WWW Document], 2011. . WHO. URL
581 http://www.who.int/nutrition/publications/obesity/WHO_report_waistcircumferenc
582 [e_and_waisthip_ratio/en/](http://www.who.int/nutrition/publications/obesity/WHO_report_waistcircumferenc) (accessed 9.21.20).

583 Young, R.C., Biggs, J.T., Ziegler, V.E., Meyer, D.A., 1978. A rating scale for mania: reliability,
584 validity and sensitivity. *Br J Psychiatry* 133, 429–435.

585
586
587
588

589 Table 1: Characteristics of patients in the FACE-BD sample, according to the presence of alcohol use disorder
590 (AUD). N(%) or median (interquartile range, IQR) and raw p-values are shown. * $p < 0.001389$. Unadjusted odds
591 ratios (OR) are displayed for categorical variables. BD, bipolar disorder; MDE, major depressive episode; AAO,
592 age at onset; BIS10, Barratt Impulsiveness Scale - 10th version; ALS, affective lability by the Affective lability
593 scale; CTQ, Childhood Trauma Questionnaire – short form; FAST, Functional Assessment Short Test (FAST);
594 MADRS, Montgomery-Asberg Depression Rating Scale (MADRS); YMRS, Young Mania Rating Scale (YMRS); BMI,
595 Body mass index.

	no AUD N=2259	lifetime AUD N=768	OR	p	N
Gender (W vs. M)	812 (36%)	368 (48%)	1.6 (1.4-1.9)	<0.001*	3025
Age	40 (31-50)	39 (31-48)		0.09	3027
Referral (vs. other modes)				0.215	2628
GP	248 (13%)	66 (10%)	0.9 (0.5-1.8)		
Patient	56 (3%)	25 (4%)	1.5 (0.7-3.3)		
Psychiatrist	1620 (82%)	552 (84%)	1.1 (0.6-2.2)		
Finished high school education	290 (15%)	92 (15%)	1 (0.8-1.3)	0.983	2564
Current relationship status (vs. single)				<0.001*	2586
Married / with partner	1015 (52%)	276 (44%)	0.7 (0.6-0.8)		
Separated	220 (11%)	79 (12%)	0.9 (0.7-1.2)		
Current professional status (vs. other modes)				<0.001*	2549
Student	182 (9%)	57 (9%)	1 (0.7-1.4)		
Inactive	585 (30%)	234 (38%)	1.3 (1.1-1.6)		
Disability leave	77 (4%)	26 (4%)	1.1 (0.7-1.7)		
Retired	146 (8%)	14 (2%)	0.3 (0.2-0.5)		
Family history of BD	470 (21%)	154 (20%)	1.0 (0.8-1.2)	0.693	3027
Family history of AUD	458 (20%)	282 (37%)	2.3 (1.9-2.7)	<0.001*	3027
BD subtype type 2 (vs. 1)	1070 (47%)	394 (51%)	1.2 (1.0-1.4)	0.065	3027
AAO BD	21 (17-28)	20 (17-25)		<0.001*	2905
Polarity at BD onset (depressive vs. “up”)	681 (30%)	208 (27%)	0.9 (0.7-1.0)	0.118	3027
BD duration	15 (7-24)	16 (10-24)		0.013	2903
Lifetime SA	766 (35%)	358 (48%)	1.7 (1.4-2.0)	<0.001*	2958
Past-year BD episode	1598 (77%)	576 (84%)	1.5 (1.2-1.9)	<0.001*	2767
Number of inpatient stays (lifetime)	2 (1-3)	2 (1-4)	1.0 (1.0-1.1)	0.08	2738
Number of MDE per year of bipolar illness ^b	0.3 (0.18-0.5)	0.30 (0.17-0.51)		0.841	2317
Number of “up” episodes per year of bipolar illness ^b	0.19 (0.06-0.4)	0.21 (0.07-0.44)		0.194	2903
History of psychosis	771 (40%)	209 (32%)	0.7 (0.6-0.8)	<0.001*	2597
BIS10 total score	65 (58-73)	71 (64-79)		<0.001*	2862
CTQ total score	38 (31-48)	42 (35-54)		<0.001*	2887
ALS total score ^b	1 (1-2)	2 (1-2)		<0.001*	2845
MADRS total score	7 (2-14)	11 (5-19)		<0.001*	2991
YMRS total score	0 (0-3)	1 (0-4)		<0.001*	2985
FAST total score	18 (8-30)	24 (13-34)		<0.001*	2880
FAST – cognitive subscale	3 (1-6)	3 (1-7)		<0.001*	2925
BMI	24 (22-28)	25 (22-28)		0.533	2859
Waist circumference (cm)	91 (82-101)	93 (83-103)		0.015	2711

	no AUD N=2259	lifetime AUD N=768	OR	p	N
History of any non-psychiatric disorder	1712 (84%)	598 (87%)	1.2 (1.0-1.6)	0.118	2730
History of cardiovascular disorder	262 (12%)	107 (16%)	1.3 (1.0-1.6)	0.045	2796
History of dyslipidemia	313 (15%)	124 (18%)	1.2 (1.0-1.6)	0.071	2843
History of digestive system disorder	86 (4%)	45 (6%)	1.6 (1.1-2.3)	0.014	2790
Comorbid anxiety disorder	904 (41%)	393 (53%)	1.6 (1.4-1.9)	<0.001*	2938
Tobacc smoking (vs. never-smoking)				<0.001*	2871
Former smoking	247 (12%)	103 (14%)	2.9 (2.2-3.9)		
Current smoking	844 (40%)	482 (66%)	4.0 (3.2-4.9)		
Sedative use disorder	51 (2%)	90 (12%)	5.7 (4.0-8.2)	<0.001*	3027
Cannabis use disorder	288 (13%)	312 (41%)	4.7 (3.9-5.7)	<0.001*	3027
Stimulant use disorder	46 (2%)	125 (16%)	9.3 (6.6-13.3)	<0.001*	3027

596

597

Table 2: Characteristics of patients in the FACE-BD sample according to the sequence of onset of alcohol use disorder (AUD) and bipolar disorder (BD). N(%) or median (interquartile range, IQR) and raw p-values are shown. * $p < 0.00208$. ^aAlthough the distribution was highly skewed, we report means (SD) for interpretability. N/A, not applicable; MDE, major depressive episode; AAO, age at onset; BIS10, Barratt Impulsiveness Scale - 10th version; ALS, affective lability by the Affective lability scale; CTQ, Childhood Trauma Questionnaire – short form; FAST, Functional Assessment Short Test (FAST); MADRS, Montgomery-Asberg Depression Rating Scale (MADRS); YMRS, Young Mania Rating Scale (YMRS).

	AUD first N=276	BD first N=269	Odds Ratio (95% CI)	p-value	N
Gender	151 (55%)	112 (42%)	0.6 (0.4-0.8)	0.003	545
Current professional status (vs. other modes)					
Student	27 (12%)	17 (7%)	0.6 (0.3-1.1)	0.3421	456
Inactive Disability leave	83 (37%)	85 (37%)	0.9 (0.6-1.4)		
Retired	7 (3%)	11 (5%)	1.4 (0.5-4.1)		
Current relationship status (vs. single)				0.8344	473
Married / with partner	102 (43%)	107 (45%)	1.1 (0.8-1.6)		
Separated	27 (11%)	29 (12%)	1.1 (0.6-2.1)		
Family history of AUD	105 (38%)	103 (38%)	1.0 (0.7-1.4)	1.0000	545
AAO BD	21 (18-27)	18 (15-22)		<0.0001*	545
AAO AUD	18 (16-20)	27 (20-34)		<0.0001*	545
Past-year BD episode	202 (79%)	226 (89%)	2.2 (1.3-3.7)	0.0026	508
Number of MDE per year of bipolar illness ^a	0.43 (0.34)	0.34 (0.25)		0.0225	417
Number of “up” episodes per year of bipolar illness ^a	0.45 (0.39)	0.32 (0.31)		<0.0001*	321
Lifetime SA	119 (44%)	132 (50%)	1.3 (0.9-1.8)	0.1598	535
History of psychosis	87 (36%)	70 (29%)	0.7 (0.5-1.1)	0.1626	483
BIS10 total score	71 (12)	71 (11)		0.4364	514
CTQ total score	53 (49-58)	54 (50-60)		0.0569	520
ALS total score	1 (1-2)	2 (1-2)	1.6 (1.2-2.1)	<0.0001*	509
MADRS total score	9 (4-16)	12 (5-21)		0.0081	541
YMRS total score	2 (0-4)	1 (0-4)		0.7878	537
FAST total score	21 (11-31)	25 (14-34)		0.0186	515
FAST – cognitive subscale	3 (1-6)	3 (1-7)		0.1283	524
Comorbid anxiety disorder	136 (51%)	158 (60%)	1.5 (1-2.1)	0.0335	530
Tobacc smoking (vs. never-smoking)				0.6024	530
Former smoking	40 (15%)	31 (12%)	0.8 (0.4-1.4)		
Current smoking	174 (64%)	171 (66%)	0.9 (0.6-1.5)		
Sedative use disorder	23 (8%)	36 (13%)	1.7 (1-3)	0.0786	545
Cannabis use disorder	136 (49%)	91 (34%)	0.5 (0.4-0.7)	<0.0001*	545
Stimulant use disorder	48 (17%)	31 (12%)	0.6 (0.4-1)	0.0682	545

Figure 1: Binary logistic regression with the sequence of onset of bipolar disorder (BD) and alcohol use disorder (AUD) as the dependent variable (BD-BD patients are the reference group). Bars represent 95% CI. *p <0.05, ** p <0.001, * p <0.0001. ALS, Affective lability scale; AAO, age at onset; “UP” represents (hypo)manic episodes.**

