



HAL
open science

EXACT DIMENSION OF FURSTENBERG MEASURES

Francois Ledrappier, Pablo Lessa

► **To cite this version:**

Francois Ledrappier, Pablo Lessa. EXACT DIMENSION OF FURSTENBERG MEASURES. 2021.
hal-03231726v2

HAL Id: hal-03231726

<https://hal.science/hal-03231726v2>

Preprint submitted on 27 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

EXACT DIMENSION OF FURSTENBERG MEASURES

FRANÇOIS LEDRAPPIER AND PABLO LESSA

ABSTRACT. For a probability measure μ on $SL_d(\mathbb{R})$, we consider the Furstenberg stationary measure ν on the space of flags. Under general non-degeneracy conditions, if μ is discrete and if $\sum_g \log \|g\| \mu(g) < +\infty$, then the measure ν is exact-dimensional.

1. INTRODUCTION

1.1. Main results. Let μ be a probability measure on the group $SL_d(\mathbb{R})$ of $d \times d$ real matrices with determinant 1. Let \mathcal{F} be the space of complete flags in \mathbb{R}^d ,

$$f \in \mathcal{F} \iff f = U_0 \subset U_1 \subset \dots \subset U_{d-1} \subset U_d,$$

where U_i is a vector space of dimension i in \mathbb{R}^d , $U_0 = \{0\}$, $U_d = \mathbb{R}^d$. $SL_d(\mathbb{R})$ acts naturally on \mathcal{F} . A probability measure ν on \mathcal{F} is called *stationary* if it satisfies

$$\int g_* \nu d\mu(g) = \nu.$$

By compactness, there always exist stationary measures. Understanding stationary measures is central to many studies of linear groups and applications (see [BQ16] for a recent survey).

Let G be a group, μ a probability on G , X a compact G -space and ν a stationary probability measure. We define the *Furstenberg entropy* as the nonnegative number $h(X, \mu, \nu)$ given by

$$h(X, \mu, \nu) := \int_G \int_X \log \frac{dg_* \nu}{d\nu}(x) \frac{dg_* \nu}{d\nu}(x) d\nu(x) d\mu(g),$$

with the convention that $0 \log 0 = 0$ and that the entropy is $+\infty$ if the measure $g_* \nu$ is not absolutely continuous with respect to the measure ν for a set of positive μ -measure of elements $g \in G$.

Let G_0 be the subgroup of $SL_d(\mathbb{R})$ generated by the support of μ . Assume that $\int_{G_0} \log \|g\| d\mu(g) < +\infty$ and ν is extremal among stationary measures. Then, there are d *Lyapunov exponents*

$$\chi_1 \geq \chi_2 \geq \dots \geq \chi_d, \text{ with } \chi_1 + \dots + \chi_d = 0,$$

1991 *Mathematics Subject Classification.* 37C45, 37A99, 28A80.

Key words and phrases. Furstenberg measure, dimension.

FL was partially supported by IFUM; PL thanks CSIC research project 389.

such that for ν -a.e. $f \in \mathcal{F}$, $f = \{\{0\} \subset U_1(f) \subset \dots \subset U_{d-1}(f) \subset \mathbb{R}^d\}$, any $j, j = 1 \dots, d-1$, $\mu^{\otimes \mathbb{N}}$ and almost every sequence g_0, g_1, \dots ,

$$(1) \quad \sum_{i \leq j} \chi_i = \lim_{n \rightarrow \infty} \frac{1}{n} \log |\det_{U_j(f)}(g_{n-1} \dots g_1 g_0)|,$$

where, for any subspace U in \mathbb{R}^d , $|\det_U(g)|$ is the Jacobian of the linear mapping from U to gU , both endowed with the Euclidean metric.

The following inequality is very general (and essentially due to Furstenberg ([F63]))

Theorem 1.1. *Let μ be a probability on $SL_d(\mathbb{R})$ such that $\int_{SL_d(\mathbb{R})} \log \|g\| d\mu(g) < +\infty$. Then there exists a stationary measure ν on \mathcal{F} such that*

$$(2) \quad h(\mathcal{F}, \mu, \nu) \leq \sum_{i, j: i < j} \chi_i - \chi_j.$$

If there is equality in (2), then the measure ν is exact-dimensional with dimension $d(d-1)/2$.

The inequality (2) is proven in Section 5. See the discussion after theorem 1.6 for the equality case.

Remark 1.2. *It follows from theorem 1.1 and its proof, that if $\int_{SL_d(\mathbb{R})} \log \|g\| d\mu(g) < +\infty$, then there exists a stationary measure ν with finite entropy. In particular, for μ -a.e. $g \in G_0$, the measure $g_*\nu$ is absolutely continuous with respect to the measure ν (see Corollary 5.9).*

Let $\mathcal{M}(d)$ be the set of probability measures on $SL_d(\mathbb{R})$ with $\int \log \|g\| d\mu(g) < +\infty$ such that the stationary measure on \mathcal{F} is unique and the Lyapunov exponents are pairwise distinct. For example, if the group G_0 is Zariski dense in $SL(d, \mathbb{R})$ and $\int \log \|g\| d\mu(g) < +\infty$, then $\mu \in \mathcal{M}(d)$ (see [GR89] and [GM89]).

Let (X, ρ) be a metric space, ν a measure on X . The *lower dimension* $\underline{\delta}$ and the *upper dimension* $\bar{\delta}$ of (X, ρ, ν) are defined by

$$\underline{\delta} = \operatorname{ess.\,inf}_{\nu} \liminf_{r \rightarrow 0} \frac{\log \nu(B(x, r))}{\log r}, \quad \bar{\delta} = \operatorname{ess.\,sup}_{\nu} \limsup_{r \rightarrow 0} \frac{\log \nu(B(x, r))}{\log r}.$$

A measure ν on X is called *exact-dimensional with dimension* δ if $\underline{\delta} = \bar{\delta} = \delta$.

If the space (X, ρ) is bilipschitz equivalent to an Euclidean \mathbb{R}^n , $n \geq 1$, and ν is exact-dimensional of dimension δ , then δ is the smallest Hausdorff dimension of sets of positive ν -measure (see e.g. [Y82], Prop 2.1). Our main result is

Theorem 1.3. *Let $\mu \in \mathcal{M}(d)$ be a discrete probability measure. Endow the space \mathcal{F} of flags with the natural Riemannian distance invariant under the action of $SO(d)$. Then the unique stationary probability measure ν on the space \mathcal{F} is exact-dimensional.*

The dimension in Theorem 1.3 is given by a formula involving exponents and some partial entropies (see (8)). This implies an a priori bound on the dimension that we describe now. Denote $\{0 < \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_{d(d-1)/2}\}$ the differences of

exponents $\chi_i - \chi_j$ for all $(i, j), i < j$. Assume μ is discrete and belongs to $\mathcal{M}(d)$. Let ν be the unique stationary measure. We define the continuous, piecewise affine function D_μ on the interval $[0, d(d-1)/2]$ as:

$$D_\mu(0) := h(\mathcal{F}, \mu, \nu) \quad \text{and} \quad D'_\mu(s) = -\lambda_k \text{ for } s \in (k-1, k), \quad k = 1, \dots, d(d-1)/2.$$

Observe that by theorem 1.1, $D_\mu(d(d-1)/2) \leq 0$. Following Kaplan-Yorke ([KY79]) and Douady-Oesterlé ([DO80]), the *Lyapunov dimension* $\dim_{LY}(\mathcal{F}, \mu)$ is the number such that $D_\mu(\dim_{LY}(\mathcal{F}, \mu)) = 0$.

Theorem 1.4. *Let $\mu \in \mathcal{M}(d)$ be discrete. Then, the exact dimension δ of ν satisfies*

$$(3) \quad \delta \leq \dim_{LY}(\mathcal{F}, \mu).$$

We discuss the proof of theorem 1.4 in Section 2.3.

We can prove equality in relation (3) in some examples: in dimension 2, for $\mu \in \mathcal{M}(2)$, we always have $\delta(\nu) = \frac{h(\mathcal{F}, \mu, \nu)}{\chi_1 - \chi_2} = \dim_{LY}(\mathcal{F}, \mu)$ ([Led84], where the formula is proven with a less precise notion of dimension); in section 9, we discuss the terms and the proof of the following

Theorem 1.5. *Let Γ be a cocompact group of isometries of \mathbb{H}^2 , ρ a Hitchin representation of Γ in $PSL_d(\mathbb{R})$ and μ be an adapted probability measure on Γ such that $\sum_g |g| \mu(g) < +\infty$, where $|\cdot|$ is some word metric on Γ . Consider the random walk on $PSL_d(\mathbb{R})$ directed by the probability $\rho_*(\mu)$ and ν the stationary measure on the space \mathcal{F} of flags. Then, ν is exact-dimensional and $\delta(\nu) = \dim_{LY}(\mathcal{F}, \mu)$.*

If $\mu \in \mathcal{M}(d)$, we can also consider a partition $Q = \{0 < q_1 < \dots < q_{\ell-1} < q_\ell = d\}$ of $\{0, 1, \dots, d\}$ into intervals, the group $SL_d(\mathbb{R})$ acts naturally on the space \mathcal{F}_Q of increasing sequences of vector subspaces of \mathbb{R}^d ,

$$\{0\} = U_0 \subset U_1 \subset \dots \subset U_{\ell-1} \subset U_\ell = \mathbb{R}^d,$$

with $\dim U_i = q_i$ for $i = 1, \dots, \ell$. The group G_0 acts naturally on \mathcal{F}_Q and there is a unique stationary probability measure ν_Q for this action. Theorems 1.1, 1.3 and 1.4 will be the particular case ($Q = Q_1 := \{0 < 1 < 2 < \dots < d\}$) of the corresponding results stated and proven for the action of \mathcal{F}_Q , for any partition Q (see respectively theorem 5.1, corollary 2.5 and proposition 2.9).

1.2. Related results. There are many results (and still open questions) related to the regularity of the Furstenberg measure in dimension 2: the space \mathcal{F} is the space of lines in \mathbb{R}^2 and theorem 1.3 holds for a general measure $\mu \in \mathcal{M}$ (Hochman and Solomyak [HS17]). Theorem 1.4 follows with equality in the formula (3). The main result of [HS17] is stronger: it concerns a formula analogous to (3) but with a larger definition of the Lyapunov dimension and is related to the well-known problem of finding conditions under which the Furstenberg measure is absolutely continuous. For any lattice $\Gamma \in SL_2(\mathbb{R})$, by discretizing the Brownian motion on the symmetric space, Furstenberg ([F71]) constructed a probability measure on Γ which belongs to $\mathcal{M}(2)$ and such that the stationary measure is Lebesgue. On the contrary, the stationary ν is singular if μ has finite support on $SL_2(\mathbb{Z})$ (Guivarc'h-Le Jan [GL93]). It is not known whether the same is true for measures with finite

support on uniform lattices. There are examples of measures with finite support and an absolutely continuous stationary measure (Bárány-Pollicott-Simon [BPS] and Bourgain [B12]), but the group generated by the support is dense.

In all dimensions, under exponential moment and proximality conditions, Guivarc'h ([G90]) showed that the unique stationary measure ν has uniformly positive dimension: there is $C, \delta > 0$ such that for all $f \in \mathcal{F}$, all $\varepsilon > 0$, $\nu(B(f, \varepsilon)) \leq C\varepsilon^\delta$. The stationary measure might be absolutely continuous: Furstenberg's discretization works in all dimensions and Benoist-Quint ([BQ18]) found finitely supported measures with absolutely continuous stationary measures. In the other direction, there exist measures supported on a discrete Zariski dense subgroup Γ with arbitrarily small \dim_{LY} (Kaimanovich-Le Prince [KLP11]).

Theorem 1.3 has been recently proven by A. Rapaport ([Rap21]) for the stationary measure on the space $\mathcal{F}_{0 < 1 < d} = \mathbb{RP}^{d-1}$ of directions in \mathbb{R}^d and on the space $\mathcal{F}_{0 < d-1 < d} = \mathcal{G}_{d-1}$ of $(d-1)$ -dimensional hyperplanes in the case when the measure μ is finitely supported. Theorem 1.4 follows with a notion of \dim_{LY} adapted to the projective space (see Corollary 1.7 in [Rap21]). We explain in section 6.2 how to recover [Rap21] from our results when both apply. D. Feng proved a similar result for affine IFSs ([Fen19], see also [BK17]). All these papers (and this one) can be seen as higher dimension extensions of [FH09]. For the flag space (and for the projective space as well), the extension of the equality $\delta(\nu) = \dim_{LY}(\mathcal{F}, \mu)$ to μ discrete in $\mathcal{M}(d)$, $d > 2$ seems to be delicate. The proof of theorem 1.3 may suggest a strategy. This is in accord with the results of [BHR19], [HR19].

1.3. Strategy in dimension 3. The main new feature of our paper is already present for finitely supported measures in the case $d = 3$. The space $\mathcal{F}_{0 < 1 < 3}$ is the space \mathcal{L} of lines in \mathbb{R}^3 , the space $\mathcal{F}_{0 < 2 < 3}$ is the space \mathcal{P} of planes in \mathbb{R}^3 . The stationary measure ν on \mathcal{F} projects on the stationary measures $\nu_{\mathcal{L}}$ and $\nu_{\mathcal{P}}$ and the fibers of the projections are one-dimensional. We know by [Rap21] that $\nu_{\mathcal{L}}$ and $\nu_{\mathcal{P}}$ are exact-dimensional. Moreover, we know by [Les21] that the conditional measures on the fibers are exact-dimensional and [Les21] has formulae for the almost everywhere constant dimensions. This is not enough information to be able to conclude that the measure ν is exact-dimensional and to compute its dimension. Indeed, in the setting of Theorem 1.5 in dimension 3, as soon as the Hitchin representation is not Fuchsian, there exists a probability measure $\mu_0 \in \mathcal{M}(\Gamma)$ for which the dimensions do not add up for the projection from (\mathcal{F}, ν_0) to \mathcal{P} (see proposition 9.1 and theorem 9.2).

We next recall this phenomenon of *dimension conservation* and explain what is the third codimension one projection of \mathcal{F} that we consider and for which we will prove dimension conservation. In the following subsection, we introduce the corresponding formalism in higher dimensions.

Let $(X, \nu), (X', \nu')$ be standard probability spaces, $\pi : (X, \nu) \rightarrow (X', \nu')$ a measure preserving mapping. Recall that a *disintegration* of the measure ν with respect to π is an a.e. defined measurable family of probability measures $x' \mapsto \nu^{x'}$

(or $x' \mapsto \nu_{X'}^{x'}, x' \mapsto \nu_{\pi}^{x'}$) of probability measures on X such that

$$\nu^{x'} \pi^{-1}(x') = 1 \quad \text{and} \quad \nu = \int_{X'} \nu^{x'} d\nu'(x').$$

Two families of disintegrations of the measure ν with respect to π coincide ν' -a.e..

Assume now that $(X, d), (X', d')$ are separable metric spaces and that $\pi : (X, d) \rightarrow (X', d')$ is a Lipschitz mapping. Let ν be a probability measure on X . We say that the projection π is *dimension conserving* for ν if

- (1) the measure ν is exact-dimensional with dimension δ ,
- (2) the measure $\pi_*\nu$ is exact-dimensional with dimension δ' ,
- (3) for $\pi_*\nu$ -a.e. $x' \in X'$, the disintegration $\nu^{x'}$ is exact-dimensional on $\pi^{-1}(x')$ with dimension $\delta - \delta'$.

The definition is adapted from Furstenberg ([F70], [F08]). Dimension conservation occurs often in the presence of iterations or randomness. Classical examples are the results à la Marstrand and Mattila for projections of measures along almost every direction in \mathbb{R}^d ([JM98]), see [JJL04] and the more recent [FFJ15] and [S15] for surveys. On the other hand, it is easy to construct examples (for instance the graphs of the Brownian trajectories or the graph of the Weierstraß function (see [She18])) where 1 and 2 hold, but the conditional measures are Dirac measures with dimension 0 whereas $\delta > \delta'$. See also [Rap17] for an example in a context close to ours.

We will now describe, in dimension 3, a third projection defined on \mathcal{F} with one-dimensional fibers for which we will be able to prove dimension conservation (see corollary 2.3).

We say that two flags $f = \{0\} \subset U_1 \subset U_2 \subset \mathbb{R}^3$ and $f' = \{0\} \subset U'_1 \subset U'_2 \subset \mathbb{R}^3$ are in general position if $U_1 \oplus U'_2 = U_2 \oplus U'_1 = \mathbb{R}^3$. Set $\mathcal{F}^{(2)}$ for the set of pairs of flags in general position. The mapping $F : \mathcal{F}^{(2)} \rightarrow \mathcal{P} \times \mathcal{L} \times \mathcal{F}$ defined by

$$F(f, f') := (U_1 \oplus U'_1, U_2 \cap U'_2, f')$$

has one-dimensional fibers. Indeed, fixing U_1 in $U_1 \oplus U'_1$ determines $U_2 = U_1 \oplus (U_2 \cap U'_2)$; alternatively, choosing $U_2 \supset U_2 \cap U'_2$ determines $U_1 = (U_1 \oplus U'_1) \cap U_2$.

Let μ' be the image of μ under the mapping $g \mapsto g^{-1}$ and let ν' be the μ' -stationary probability measure for the action of G on \mathcal{F} associated by remark 4.3. From our results will follow that, for ν' -a.e. f' , all projections in the following sequence are dimension conserving for $\nu \times \delta_{f'}$

$$\begin{aligned} (f, f') &\mapsto (U_1 \oplus U'_1, U_2 \cap U'_2, f') \mapsto (U_1 \oplus U'_1, f') \mapsto f' & \text{if } \chi_2 \geq 0, \\ (f, f') &\mapsto (U_1 \oplus U'_1, U_2 \cap U'_2, f') \mapsto (U_2 \cap U'_2, f') \mapsto f' & \text{if } \chi_2 \leq 0. \end{aligned}$$

In order to prove theorem 1.3 on \mathcal{F} when $d = 3$, we are reduced to three projections with one dimensional fibers, for which we want to prove exact dimensionality of the conditional measures on the fibers and dimension conservation. Moreover, we have arranged so that the exponents of the dynamics on the fibers are nondecreasing:

$$\begin{aligned} \chi_3 - \chi_1 < \chi_3 - \chi_2 \leq \chi_2 - \chi_1 < 0 & \quad \text{if } \chi_2 \geq 0, \\ \chi_3 - \chi_1 < \chi_2 - \chi_1 \leq \chi_3 - \chi_2 < 0 & \quad \text{if } \chi_2 \leq 0. \end{aligned}$$

Therefore, we can apply the strategy of [Fen19] and [Rap21] (following [LY85], [FH09], [BK17]), and work one exponent at a time. We first generalize the above picture to higher dimensions.

1.4. Topologies, configuration spaces and entropy. A topology T on the set $\{1, \dots, d\}$ will be called admissible if $\{i, i+1, \dots, d\} \in T$ for all i . Given an admissible topology T we denote by $T(i)$ the atom of i i.e. the smallest set in T containing i . Notice that any topology T is determined by listing its atoms $T(1), T(2), \dots, T(d)$. And T is admissible if, and only if, $T(i) \subset \{i, i+1, \dots, d\}$ for all i .

Recall that a topology T is finer than another T' (equivalently, T' is coarser than T), denoted $T \prec T'$, if $T \supset T'$. The coarsest admissible topology T_0 is (defined by the list of atoms)

$$\{1, \dots, d\}\{2, \dots, d\} \dots \{d\},$$

the finest admissible topology T_1 is $\{1\} \dots \{d\}$.

We say an admissible topology T is one step finer than an admissible topology T' (equivalently T' is one step coarser than T), denoted $T \stackrel{1}{\prec} T'$, if there exists a unique $i \in \{1, \dots, d\}$ such that $T(i) \neq T'(i)$ and furthermore $T'(i) \setminus T(i)$ is a singleton. Let j be so that $\{j\} = T'(i) \setminus T(i)$. Then, $j > i$ and $T(i) \cup \{i\} \subset T(i) \subset T(i) \cup \{j\} = T'(i)$. We associate to a pair $T \stackrel{1}{\prec} T'$ its *exponent* $\chi_{T, T'}$

$$(4) \quad \chi_{T, T'} = \chi_i - \chi_j.$$




FIGURE 1. Admissible topologies for $d = 3$, an arrow indicates a topology one step coarser than another, filtered topologies are indicated in gray.

An admissible topology is called filtered if it is generated by the coarsest admissible topology and some subset of $\{\{1\}, \{1, 2\}, \dots, \{1, 2, \dots, d\}\}$. Let Q be a partition of $\{0, 1, \dots, d\}$ into intervals, $Q = \{q_0 = 0 < q_1 < \dots < q_k = d\}$. We associate to it the filtered topology T_Q generated by T_0 and the sets $\{1, 2, \dots, q_j\}, j = 1, \dots, k$. There are exactly 2^{d-1} filtered topologies and they are all obtained that way. The topologies T_1 and T_0 are filtered and correspond to respectively the space of complete flags and the one-point flag space of the trivial partition $Q_0 = \{1, 2, \dots, d\}$.

Figures 1 and 2 represent the graphs of the one-step relations between admissible topologies in dimensions 3 and 4 respectively. In dimension 3, the new fibration correspond to the nonfiltered topology that is one-step finer than T_1 . One sees the two ways of further descending on the graph according to the sign of χ_2 . For a general d , admissible topologies are in one-to-one correspondence with partial orders on $\{1, \dots, d\}$ that are suborders of the natural order. Their number as a function of d is the list A006455 of the Online Encyclopedia of Integer Sequences.¹ In dimension 4, there are indeed 40 admissible topologies and 92 one-step arrows. The non-trivial filtered topologies correspond to the spaces of partial flags with only one level missing or to the Grassmannians of lines, planes or three-dimensional spaces. This correspondence is extended to all admissible topologies by constructing the *configuration spaces* as follows.

Given an admissible topology T we define the configuration space \mathcal{X}_T to be the space of sequences $x = (x_I)_{I \in T}$ indexed on T where

- (1) x_I is a $|I|$ -dimensional subspace of \mathbb{R}^d for each $I \in T$,
- (2) $x_{I \cup J} = x_I + x_J$ for all $I, J \in T$, and
- (3) $x_{I \cap J} = x_I \cap x_J$ for all $I, J \in T$.

¹We thank Yves Coudène for this observation.




FIGURE 2. Admissible topologies for $d = 4$, an arrow indicates a topology one step coarser than another, filtered topologies are indicated in gray.

The configuration space \mathcal{X}_{T_1} is identified with the space of d independent lines in \mathbb{R}^d ; \mathcal{X}_{T_0} is the space \mathcal{F} of complete flags. If T is finer than T' there is a natural projection mapping $\pi_{T,T'} : \mathcal{X}_T \rightarrow \mathcal{X}_{T'}$. In particular, all configuration spaces project onto \mathcal{X}_{T_0} .

We say that two flags $f = \{0\} \subset U_1 \subset \dots \subset \mathbb{R}^d$ and $f' = \{0\} \subset U'_1 \subset \dots \subset \mathbb{R}^d$ are in general position if for all $j, 0 < j < d$, $U_j \oplus U'_{d-j} = \mathbb{R}^d$. Set $\mathcal{F}^{(2)}$ for the set of pairs of flags in general position. Given an admissible topology T , we associate to $(f, f') \in \mathcal{F}^{(2)}$ the configuration $F_T(f, f') \in \mathcal{X}_T$ given, for $I \in T$, by

$$[F_T(f, f')]_I = \bigoplus_{i \in I} (U_i \cap U'_{d-i+1}).$$

Observe that if T is finer than T' , then $F_{T'} = \pi_{T,T'} \circ F_T$. Set, for an admissible topology T and a fixed $f' \in \mathcal{F}$, $\mathcal{X}_T^{f'}$ for the set of $F_T(f, f')$ for all f such that $(f, f') \in \mathcal{F}^{(2)}$. In particular, we identify $\mathcal{X}_{T_0}^{f'}$ with $\{f'\}$, $\mathcal{X}_{T_1}^{f'}$ with the set of flags in general position with respect to $\{f'\}$. If Q is a partition of $\{0, 1, \dots, d\}$ into intervals, the configuration space $\mathcal{X}_{T_Q}^{f'}$ is identified with the set of partial flags \mathcal{F}_Q in general position with respect to f' .

Let μ' be the image of μ under the mapping $g \mapsto g^{-1}$ and let ν' be the μ' -stationary probability measure for the action of G on \mathcal{F} associated by remark 4.3. Endow $\mathcal{F}^{(2)}$ with the measure $\nu \otimes \nu'$ and \mathcal{X}_T with the measure $(\mathcal{F}_T)_*(\nu \otimes \nu')$. Write $f' \mapsto \nu_T^{f'}$ for the ν' -a.e. defined family of disintegrations of $(\mathcal{F}_T)_*(\nu \otimes \nu')$ with respect to the projection on the second coordinate in $\mathcal{F}^{(2)}$. By definition, for ν' -a.e. f' , $\nu_T^{f'}$ is a probability measure supported by $\mathcal{X}_T^{f'}$ and so that $(\mathcal{F}_T)_*(\nu \otimes \nu') = \int_{\mathcal{F}} \nu_T^{f'} d\nu'(f')$.

We define the *entropy* κ_T by

$$(5) \quad \kappa_T := \int \log \frac{dg_* \nu_T^{g^{-1}f'}}{d\nu_T^{f'}}(y) dg_* \nu_T^{g^{-1}f'}(y) d\nu'(f') d\mu(g).$$

We will see in Section 6 that this integral makes sense and can be seen as a conditional mutual entropy $H(gF_T, F_T | f')$. In particular, $\kappa_{T_0} = 0$. If T is filtered and associated to the partition Q , then the mapping of \mathcal{X}_{T_Q} onto \mathcal{F}_Q is a bilipschitz homeomorphism when restricted to each fiber $\mathcal{X}_{T_Q}^{f'}$, and identifies $\nu_{T_Q}^{f'}$ with ν_Q . Therefore,

$$\kappa_{T_Q} = \int_{G \times \mathcal{F}_Q} \log \frac{dg_* \nu_Q}{d\nu_Q}(x) dg_* \nu_Q(x) d\mu(g) = h(\mathcal{F}_Q, \mu, \nu_Q).$$

Assume that the admissible topology T is finer than the admissible topology T' . Then, clearly, for ν' -a.e. $f' \in \mathcal{F}$,

$$(\pi_{T,T'})_* \nu_T^{f'} = \nu_{T'}^{f'}$$

and we set $(\nu_{T,T'}^{x'}, x' \in \mathcal{X}_{T'}^{f'})$ for a family of disintegrations of the measure $\nu_T^{f'}$ with respect to $\pi_{T,T'}$. The *entropy difference*

$$(6) \quad \kappa_{T,T'} := \kappa_T - \kappa_{T'}$$

can be seen as a conditional mutual entropy $H(gF_T, F_T | F_{T'}, f')$ and can be expressed in terms of the measures $\nu_{T,T'}^{x'}$ (see below section 6).

A key step in our proof is

Theorem 1.6. *Fix $\mu \in \mathcal{M}(d)$. Assume T and T' are admissible topologies, with T one step finer than T' . With the above notations, for ν' -a.e. f' , for $\nu_{T'}^{f'}$ -a.e. $x' \in \mathcal{X}_{T'}^{f'}$, the measure $\nu_{T,T'}^{x'}$ is exact-dimensional with dimension $\gamma_{T,T'}$ given by*

$$\gamma_{T,T'} = \frac{\kappa_{T,T'}}{\chi_{T,T'}}.$$

The proof of theorem 1.6 is given in section 7. In dimension $d = 2$, there is only one pair T_1, T_0 , the measure $\nu_{T_1, T_0}^{f'}$ is the constant measure ν and theorem 1.6 comes from [HS17]. In higher dimensions, if $T = T_1$ and T' has one atom of the form $\{i, i+1\}$, then theorem 1.6 is theorem 2 from [Les21]. The general scheme of the proof is the same: we find a one-dimensional parameterization of the support of the measure $\nu_{T,T'}^{x'}$ that is adapted to the G -action, and then use a telescoping argument and the Maker ergodic theorem. It follows from the one-dimensional parameterization that if $T \stackrel{1}{\prec} T'$, then $\gamma_{T,T'} \leq 1$ (see lemma 3.4 below) and this leads to a general estimate of $\kappa_{T,T'}$ in terms of the differences of exponents associated to $\pi_{T,T'}$ (see proposition 2.1.2 below). Inequality (2) in theorem 1.1 corresponds to the particular case of κ_{T_1, T_0} . We provide a direct proof of (2) first, since we use it to ensure that the entropy κ_T defined by equation (5) is finite. Since for all $T \stackrel{1}{\prec} T'$, $\gamma_{T,T'} \leq 1$, if we have equality in (2), all corresponding $\gamma_{T,T'}$ are 1 and the equality case in theorem 1.1 follows from theorem 2.2.1.

In the next section, we reduce the proofs of our results to entropy/dimension statements related to the fine structure of the configuration spaces. A more detailed organization of the remaining proofs is given at the end of the next section.

2. PROOFS

2.1. Proof of theorem 1.3. Observe that for T and T' admissible topologies, $T \prec T'$ if, and only if, for all $i = 1, \dots, d-1$, $T(i) \subset T'(i)$, where $T(i)$ is the atom of T containing $\{i\}$. We denote $D_{T,T'}$ the set of pairs (i, j) such that $j \in T'(i) \setminus T(i)$. The numbers $\chi_i - \chi_j, (i, j) \in D_{T,T'}$ are called the exponents of the pair $T \prec T'$.

Observe that $\{i\} \subset T(i), T'(i) \subset \{i, i+1, \dots, d\}$. Therefore, for $(i, j) \in D_{T,T'}$, $i < j$ and the exponents $\chi_i - \chi_j, (i, j) \in D_{T,T'}$ are positive. Set $N_{T,T'} := \#D_{T,T'} = \sum_{i=1}^d \#(T'(i) \setminus T(i))$.

Proposition 2.1. *Let $T \prec T'$ be a pair of admissible topologies, $N := N_{T,T'}$.*

1. Then, there exists a sequence $T^0 = T', T^1, \dots, T^N = T$ such that T^t is one step finer than T^{t-1} for $t = 1, \dots, N$, and

$$\chi_{T^1, T^0} \leq \chi_{T^2, T^1} \leq \dots \leq \chi_{T^N, T^{N-1}}.$$

2. Moreover, if $\mu \in \mathcal{M}(d)$, $\bar{\delta}_{T,T'} \leq N_{T,T'}$ and $\kappa_{T,T'} \leq \sum_{(i,j) \in D_{T,T'}} (\chi_i - \chi_j)$, where $\bar{\delta}_{T,T'}$ is the essentially constant (in x') value of the upper dimension of the measure $\nu_{T,T'}^{x'}$.

Proposition 2.1.1 is proven in section 3.1. Proposition 2.1.2 is proven at the end of section 3.2 after a more precise description of the metric spaces $(\pi_{T,T'})^{-1}(x')$ for $x' \in \mathcal{X}_{T'}$. For any pair $T \prec T'$ of admissible topologies, we henceforth choose and fix a sequence given by proposition 2.1.1. The entropy $\kappa_{T,T'}$ is the sum of the entropy differences $\kappa_{T^t, T^{t-1}}$.

Theorem 1.6 yields a Ledrappier-Young formula for the entropy $\kappa_{T,T'}$

$$(7) \quad \kappa_{T,T'} = \sum_{t=1}^{N_{T,T'}} \kappa_{T^t, T^{t-1}} = \sum_{t=1}^{N_{T,T'}} \chi_{T^t, T^{t-1}} \gamma_{T^t, T^{t-1}}.$$

The precise form of our main result is the dimension counterpart of (7):

Theorem 2.2. *Let $\mu \in \mathcal{M}(d)$ and $T \prec T'$ be a pair of admissible topologies. With the previous notations, for ν' -a.e. $f' \in \mathcal{F}$, $\nu_T^{f'}$ -a.e. $x' \in \mathcal{X}_{T'}^{f'}$,*

- (1) *the lower dimension $\underline{\delta}_{T,T'}$ of the measure $\nu_{T,T'}^{x'}$ is at least $\sum_{t=1}^{N_{T,T'}} \gamma_{T^t, T^{t-1}}$;*
- (2) *moreover, in the case when μ is discrete, the measure $\nu_{T,T'}^{x'}$ is exact-dimensional, with dimension $\delta_{T,T'}$ given by*

$$\delta_{T,T'} = \sum_{t=1}^{N_{T,T'}} \gamma_{T^t, T^{t-1}}.$$

Theorem 2.2.2 gives a condition for dimension conservation:

Corollary 2.3. *Assume μ is a discrete measure in $\mathcal{M}(d)$ and let $T \prec T' \prec T''$ be admissible topologies. Assume that all differences $\chi_i - \chi_j, (i, j) \in D_{T,T'}$ are at least as large as the differences $\chi_i - \chi_j, (i, j) \in D_{T',T''}$. Then, there is dimension conservation for the projections $\pi_{T,T''} = \pi_{T',T''} \circ \pi_{T,T'}$ of the measure $\nu_T^{f'}$:*

$$\delta_{T,T''} = \delta_{T',T''} + \delta_{T,T'}.$$

Proof. Since differences $\chi_i - \chi_j, (i, j) \in D_{T,T'}$ are at least as large as the differences $\chi_i - \chi_j, (i, j) \in D_{T',T''}$, it is possible to find the sequence associated to $T \prec T''$ by proposition 2.1.1 in such a way that $T^{N_{T',T''}} = T'$. The corollary follows. \square

Observe that there are examples of $T \prec T' \prec T''$ where dimension conservation does not hold (see e.g. proposition 9.1 and theorem 9.2).

In the case when $T' = T_0$, we can identify the measures $\nu_{T^t}^{f'}$ and $\nu_{T^t, T_0}^{f'}$ on $\mathcal{X}_{T^t}^{f'}$ and we obtain, setting $D_T := D_{T, T_0} = \{(i, j), i < j, j \notin T(i)\}$, $N_T := \#D_T$ and, for $(i, j) \in D_T$, $\gamma_{i,j}^T$ for $\gamma_{T^t, T^{t-1}}$ associated by proposition 2.1.1 to (i, j) .

Corollary 2.4. *Assume μ is a discrete measure in $\mathcal{M}(d)$ and let T be an admissible topology. With the previous notations, we have $\kappa_T = \sum_{(i,j) \in D_T} \gamma_{i,j}^T (\chi_i - \chi_j)$ and, for ν' -a.e. $f' \in \mathcal{F}$, the measure $\nu_T^{f'}$ is exact-dimensional, with dimension δ_T given by*

$$\delta_T = \delta_{T, T_0} = \sum_{(i,j) \in D_T} \gamma_{i,j}^T.$$

In particular, if T is a filtered admissible topology associated to a partition Q , then \mathcal{X}_T is identified with \mathcal{F}_Q , the measure $\nu_T^{f'}$ is then identified with the measure ν_Q for almost every f' .

Corollary 2.5. *Let $\mu \in \mathcal{M}(d)$ be a discrete probability measure on $SL_d(\mathbb{R})$. Let Q be a partition of $\{0, 1, \dots, d\}$ into intervals. Then the unique stationary probability measure ν_Q on the space \mathcal{F}_Q of partial flags is exact-dimensional. There are numbers $\gamma_{i,j}^Q := \gamma_{i,j}^{T_Q}$ such that ²*

$$(8) \quad \delta_Q = \sum_{i,j:\ell_Q(i)<\ell_Q(j)} \gamma_{i,j}^Q, \quad h(\mathcal{F}_Q, \mu, \nu_Q) = \sum_{i,j:\ell_Q(i)<\ell_Q(j)} \gamma_{i,j}^Q (\chi_i - \chi_j).$$

Theorem 1.3 is the particular case of corollary 2.5 when $Q = Q_1$.

Proof of Theorem 2.2. We endow $\mathcal{X}_T^{f'}$ with a smooth metric. The following limits are constants for ν' -a.e. $f' \in \mathcal{F}$, $\nu_{T^t}^{f'}$ -a.e. $x' \in \mathcal{X}_{T^t}^{f'}$, $\nu_{T^t}^{x'}$ -a.e. $y' \in \mathcal{X}_{T^t}^{x'}$:

$$\underline{\delta}^t := \liminf_{r \rightarrow 0} \frac{\log \nu_{T^t}^{x'}(B(y', r))}{\log r}, \quad \bar{\delta}^t := \limsup_{r \rightarrow 0} \frac{\log \nu_{T^t}^{x'}(B(y', r))}{\log r}.$$

With this notation, lower and upper dimensions of the measure $\nu_{T^t}^{x'}$ are respectively $\underline{\delta}^0$ and $\bar{\delta}^0$.

For the first part of theorem 2.2, we have that for almost every (f', x', y') , the following relation (9) holds for all $t = N_{T,T'}, N_{T,T'} - 1, \dots, 1$

$$(9) \quad \underline{\delta}^{t-1} \geq \underline{\delta}^t + \gamma_{T^t, T^{t-1}}.$$

Indeed, since $T \succ T^{N_{T,T'}-1}$, (9) holds for $t = N_{T,T'}$ (with $\underline{\delta}^{N_{T,T'}} = 0$) by theorem 1.6. By [LY85] lemma 11.3.1, (9) for $t < N_{T,T'}$ follows from theorem 1.6 as well. Theorem 2.2.1 follows by summing the relations (9) for $t = N_{T,T'}, N_{T,T'} - 1, \dots, 1$.

For the second part of theorem 2.2, it remains to prove

Theorem 2.6. *Assume that the measure $\mu \in \mathcal{M}(d)$ is discrete. With the above notations, for almost every (f', x', y') , for all $t = N_{T,T'}, N_{T,T'} - 1, \dots, 1$,*

$$(10) \quad \bar{\delta}^{t-1} \leq \bar{\delta}^t + \gamma_{T^t, T^{t-1}}.$$

Summing the relations (10) for $t = N_{T,T'}, N_{T,T'} - 1, \dots, 1$ (with $\bar{\delta}^{N_{T,T'}} = 0$) gives $\bar{\delta}_{T,T'} = \bar{\delta}^0 \leq \sum_{t=1}^{N_{T,T'}} \gamma_{T^t, T^{t-1}}$. Comparing with theorem 2.2.1 gives the result. \square

We prove theorem 2.6 in Section 8. The analog of theorem 2.6 in [LY85] is a counting argument (see section (10.2)) that uses partitions with finite entropy in the underlying space. This is not possible here. By working on the trajectory space of the underlying process, [Fen19] and [Rap21] perform this counting procedure in the case when the measure μ has finite support. We follow the same scheme under the hypothesis that μ is discrete.

²Comparing with theorem 1.1, the content of (8) for Q_1 is that the numbers $\gamma_{i,j}^{Q_1}$ are the dimensions of certain conditional measures on specific 1-dimensional leaves in \mathcal{F} .

2.2. Properties of the partial dimensions. The spaces $(\pi_{T^t, T^{t-1}})^{-1}(x)$ and the measures $\nu_{T^t, T^{t-1}}^x$ for a.e. x depend only on the arrow $T^t \xrightarrow{1} T^{t-1}$ and not on its environment $T \prec T'$ such that $T \prec T^t \xrightarrow{1} T^{t-1} \prec T'$. Indeed, the dimension $\gamma_{T^t, T^{t-1}}$ in formula (7) does not depend on the environment $T \prec T'$. But still, there might be several arrows corresponding to the same pair (i, j) , $0 < i < j \leq d$. We can write

$$\begin{array}{ccc} T & \longrightarrow & S \\ \downarrow 1 & & \downarrow 1 \\ T' & \longrightarrow & S' \end{array} \quad \text{commutes and}$$

Lemma 2.7. *Assume the diagram of projections $\downarrow 1$ commutes and i, j are such that $T(i) = T'(i) \setminus \{j\}$ and $S(i) = S'(i) \setminus \{j\}$. Then, $\gamma_{T, T'} \leq \gamma_{S, S'}$.*

Lemma 2.7 is proven in section 7. The example in section 9 shows that, in general, $\gamma_{T, T'} < \gamma_{S, S'}$.

Let i, j satisfy $0 < i < j \leq d$ and let $T_{i,j}$ be the topology defined by

$$T_{i,j}(k) = \{k\} \text{ if } k \neq i, T_{i,j}(i) = \{i, j\}.$$

The topology $T_{i,j}$ is admissible and one step coarser than T_1 .

Corollary 2.8. *Let $T \prec T''$ be admissible topologies. For $T^t \xrightarrow{1} T^{t-1}$ in the decomposition of $T \prec T''$ and (i, j) such that $T^{t-1}(i) = T^t(i) \cup \{j\}$, we have $\gamma_{T_1, T_{i,j}} \leq \gamma_{T^t, T^{t-1}}$.*

Proof. Apply lemma 2.7. to the commutative part of the diagram

$$\begin{array}{ccccc} & & T & & \\ & & \downarrow & & \\ T_1 & \longrightarrow & T^t & & \\ \downarrow 1 & & \downarrow 1 & & \\ T_{i,j} & \longrightarrow & T^{t-1} & \longrightarrow & T'' \end{array}$$

□

In the case when there are several pairs $\{(i_1, j_1), \dots, (i_k, j_k)\}$ with the same difference $\chi_i - \chi_j$, there are different decompositions given by proposition 2.1.1. If the measure μ is discrete, we can apply theorem 2.2.2 to each decomposition and get a common formula by grouping together the terms corresponding to the same difference $\chi_i - \chi_j$. Namely, set $\delta_{T^{t+k}, T^t} := \sum_{\ell=1}^k \gamma_{T^{t+\ell}, T^{t+\ell-1}}$. By theorem 2.2.2 applied to $T^{t+k} \prec T^t$, this number δ_{T^{t+k}, T^t} is the exact dimension of the measures ν_{T^{t+k}, T^t} and thus is independent of the order of the decomposition. We also obtain that $\kappa_{T^{t+k}, T^t} = (\chi_i - \chi_j) \delta_{T^{t+k}, T^t}$.

2.3. Lyapunov Dimension. Theorem 1.4 follows from the previous results. We state and prove the corresponding result for a general partition Q of $\{0, 1, \dots, d\}$ into intervals. Assume $\mu \in \mathcal{M}(d)$ is discrete.

Let $\chi_1 > \dots > \chi_d$ be the Lyapunov exponents of (G, μ) and for $i = 1, \dots, d$, set $\ell_Q(i)$ for the index such that $q_{\ell_Q(i)-1} < i \leq q_{\ell_Q(i)}$. Denote $\{0 < \lambda_1 \leq \lambda_2 \leq \dots \leq$

$\lambda_N\}$ the differences of exponents $\chi_i - \chi_j$ for all (i, j) such that $\ell_Q(i) < \ell_Q(j)$. Define the continuous, piecewise affine function $D_{\mathcal{F}_Q, \mu}$ on the interval $[0, N = \dim \mathcal{F}_Q]$ as:

$$D_{\mathcal{F}_Q, \mu}(0) := h(\mathcal{F}_Q, \mu, \nu_Q) \quad \text{and} \quad D'_{\mathcal{F}_Q, \mu}(s) = -\lambda_k \text{ for } s \in (k-1, k), \quad k = 1, \dots, N.$$

As before, the *Lyapunov dimension* $\dim_{LY}(\mathcal{F}_Q, \mu)$ is such that $D_{\mathcal{F}_Q, \mu}(\dim_{LY}(\mathcal{F}_Q, \mu)) = 0$. Observe that by theorem 5.1, $\dim_{LY}(\mathcal{F}_Q, \mu) \leq \dim \mathcal{F}_Q$.

Proposition 2.9. *Let $\mu \in \mathcal{M}(d)$ be discrete, Q a partition of $\{0, 1, \dots, d\}$ into intervals. Then, the exact dimension δ_Q of ν_Q satisfies*

$$\delta_Q \leq \dim_{LY}(\mathcal{F}_Q, \mu).$$

Proof. The measure ν_Q is exact-dimensional with dimension δ_Q . By equation (8), there are numbers $\gamma_{i,j}^Q$ such that $\delta_Q = \sum_{i,j:\ell_Q(i) < \ell_Q(j)} \gamma_{i,j}^Q$. Moreover, $h(\mathcal{F}_Q, \mu, \nu_Q) = \sum_{i,j:\ell_Q(i) < \ell_Q(j)} \gamma_{i,j}^Q (\chi_i - \chi_j)$. We ordered $\{0 < \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_N\}$ the $\lambda_k := \chi_{i_k} - \chi_{j_k}$ for all (i_k, j_k) such that $\ell_Q(i_k) < \ell_Q(j_k)$. We can define another continuous, piecewise affine Lyapunov function $\underline{D}_{\mathcal{F}_Q, \mu, \nu_Q}$ on the interval $[0, \delta_Q]$ such that $\underline{D}_{\mathcal{F}_Q, \mu, \nu_Q}(0) = h(\mathcal{F}_Q, \mu, \nu_Q)$ and the slope of $\underline{D}_{\mathcal{F}_Q, \mu, \nu_Q}(s)$ on the successive intervals of length γ_{i_k, j_k}^Q is $-\lambda_k$. By (8), $\underline{D}_{\mathcal{F}_Q, \mu, \nu_Q}(\delta_Q) = 0$. On the other hand, since for all (i, j) , $\gamma_{i,j}^Q \leq 1$ (proposition 2.1.2), for all $s \in [0, \delta_Q]$,

$$\underline{D}_{\mathcal{F}_Q, \mu, \nu_Q}(s) \leq D_{\mathcal{F}_Q, \mu}(s).$$

In particular, $D_{\mathcal{F}_Q, \mu}(\delta_Q) \geq \underline{D}_{\mathcal{F}_Q, \mu, \nu_Q}(\delta_Q) = 0$. This shows proposition (2.9). \square

2.4. Content of the paper. Given the previous discussion, we still have to prove theorems 1.1, 1.5, 1.6, 2.6 and 9.2, proposition 2.1 and lemma 2.7. In section 3, we show proposition 2.1 and discuss the geometry of the spaces $\mathcal{X}_T^{f'}$. In particular, lemma 3.4 describes the one-dimensional structure of $\pi_{T, T'}^{-1}(x')$ for $x' \in \mathcal{X}_{T'}^{f'}$ and $T \prec T'$. We also discuss the multidimensional structure of the configuration spaces (proposition 3.2). We recall in section 4 the underlying trajectory space of the associated random walk, and the applications of Oseledets theorem to random walks of matrices. In section 5, we prove theorem 5.1, the generalisation of theorem 1.1. In section 6, we recall the notion of mutual information of random variables and its properties. We prove theorem 1.6, lemma 2.7 in section 7 and theorem 2.6 in section 8. The central arguments in sections 5, 7 and 8 have a long history in ergodic theory. A short comment about background heads each of the corresponding sections. In section 9, we discuss the case of Hitchin representations of cocompact surface groups, exhibit examples of non-conservation of dimension and prove theorems 1.5 and 9.2.

3. PRELIMINARIES

3.1. Topologies and exponents.

Proof of Proposition 2.1.1. Let us order the indices in $D_{T, T'}$ in such a way that

$$\chi_{i_1} - \chi_{j_1} \leq \chi_{i_2} - \chi_{j_2} \leq \dots \leq \chi_{i_{N_{T, T'}}} - \chi_{j_{N_{T, T'}}}.$$

Beginning with $T^0 = T'$ define the sequence inductively so that for each $t = 1, 2, \dots, N_{T, T'}$ the topology T^t is generated by T^{t-1} together with the set $T^{t-1}(i_t) \setminus \{j_t\}$. Since $T^t \prec T^{t-1}$, it follows by induction that T^t is admissible.

We have that $T^t \prec T^{t-1}$; we claim that T^t is one step finer than T^{t-1} . Indeed, if this is not the case, then there exists $j \in T^{t-1}(i)$ such that $i_t < j < j_t$ and $T^t(j) \setminus T^{t-1}(j) = \{j_t\}$. Since $\chi_j - \chi_{j_t} < \chi_{i_t} - \chi_{j_t}$ it must be the case that $j_t \in T(j)$, otherwise j_t would have been removed from $T^{t-1}(j)$ at some previous step. Similarly, since $\chi_{i_t} - \chi_j < \chi_{i_t} - \chi_{j_t}$ we obtain that $j \in T(i_t)$. However, these two facts imply that $j_t \in T(i_t)$ which is a contradiction. It follows that T^t is one step finer than T^{t-1} as claimed.

We finally claim that $T \prec T^t$. To see this, observe that T^t is generated by $\{A(i)\}$, for $i = 1, \dots, d$, where $A(i) = T^{t-1}(i)$ if $i \neq i_t$ and $A(i_t) = T^{t-1}(i_t) \setminus \{j_t\}$. Inductively, if $i \neq i_t$, one has $A(i) = T^{t-1}(i) \supset T(i)$. By construction $j_t \notin T(i_t)$ and therefore $A(i_t) \supset T(i_t)$ as well. Hence, $T \prec T^t$ as claimed. \square

We also record here the following fact about admissible topologies which will be used several times.

Proposition 3.1. *Let $T \stackrel{1}{\prec} T'$ be admissible topologies and i, j be such that $T(i) = T'(i) \setminus \{j\}$. Then both $T'(i) \setminus \{i\}$ and $T'(i) \setminus \{i, j\}$ are in T' .*

Proof. Since T' is admissible we have $T'(i) \subset \{i, i+1, \dots, d\}$, and $\{i+1, i+2, \dots, d\} \in T'$. Therefore, $T'(i) \setminus \{i\} = T'(i) \cap \{i+1, \dots, d\}$ is in T' as claimed.

Repeating the argument for T we obtain that $T'(i) \setminus \{i, j\} = T(i) \setminus \{i\}$ is in T .

Combining this with the fact that $T \stackrel{1}{\prec} T'$ we obtain

$$T'(i) \setminus \{i, j\} = \bigcup_{k \in T'(i) \setminus \{i, j\}} T(k) = \bigcup_{k \in T'(i) \setminus \{i, j\}} T'(k),$$

so that $T'(i) \setminus \{i, j\}$ is in T' as claimed. \square

3.2. Distances on configuration spaces. Let \mathcal{G}_i be the Grassmannian manifold of i -dimensional subspaces of \mathbb{R}^d .

We fix on each \mathcal{G}_i a Riemannian metric which is invariant under the action of orthogonal transformations with the additional property that if $S, S' \in \mathcal{G}_i$ are such that $S + S'$ has dimension $i+1$ then the Riemannian distance satisfies

$$\text{dist}(S, S') = \angle(\pi(S), \pi(S')),$$

where $\pi : S + S' \rightarrow (S + S')/(S \cap S')$ is the projection onto the quotient space $(S + S')/(S \cap S')$, which is endowed with the inner product inherited from \mathbb{R}^d .

From the definition it follows that when $\dim(S + S') = i+1$ one has

$$\text{dist}(S + W, S' + W) \leq \text{dist}(S, S'),$$

for all subspaces W such that $\dim(S + W) = \dim(S' + W)$.

Given an admissible topology T we define the distance on the configuration space \mathcal{X}_T so that

$$\text{dist}((x_I)_{I \in T}, (x'_I)_{I \in T}) = \sum_{I \in T} \text{dist}(x_I, x'_I).$$

Proposition 3.2. *Let $T \prec T'$ be admissible topologies and $x' \in \mathcal{X}_{T'}$. Then, $(\pi_{T, T'})^{-1}(x')$, endowed with the metric dist , is locally bilipschitz homeomorphic to the Euclidean space $\mathbb{R}^{N_{T, T'}}$.*

Proof. By proposition 2.1.1, we take a sequence $T^0 = T', T^1, \dots, T^N = T$ such that, T^t is one step finer than T^{t-1} for $t = 1, \dots, N$. We prove by increasing induction on t that $(\pi_{T^t, T'})^{-1}(x')$, endowed with the metric dist , is locally bilipschitz homeomorphic to the Euclidean space \mathbb{R}^t . This is trivially true for $t = 0$. So, we assume for $t > 0$, that for any $y' \in (\pi_{T^{t-1}, T'})^{-1}(x')$, there is a neighborhood of y' in $(\pi_{T^{t-1}, T'})^{-1}(x')$ which is bilipschitz homeomorphic to the Euclidean space \mathbb{R}^{t-1} . The fibers of the projections $\pi_{T^t, T^{t-1}}$ form a C^∞ foliation of $(\pi_{T^t, T'})^{-1}(x')$. We have to verify that the induced metric by dist on the fibers is bilipschitz equivalent to the Euclidean one dimensional metric, uniformly in the neighborhood of y' .

Let (i, j) such that $T^t(i) = T^{t-1}(i) \setminus \{j\}$, we define a distance on each fiber of the projection $\pi_{T^t, T^{t-1}}$ by setting

$$(11) \quad \text{dist}_{T^t, T^{t-1}}^{x'}(x_1, x_2) = \text{dist}((x_1)_{T(i)}, (x_2)_{T(i)}),$$

for each $x' \in \mathcal{X}_{T^{t-1}}$ and $x_1, x_2 \in \pi_{T^t, T^{t-1}}^{-1}(x')$. These distances are all Lipschitz equivalent on the fibers:

Lemma 3.3. *For all admissible topologies $T \prec T'$, all $x' \in \mathcal{X}_{T'}$ and all $x_1, x_2 \in \pi_{T, T'}^{-1}(x')$, one has*

$$\text{dist}_{T, T'}^{x'}(x_1, x_2) \leq \text{dist}(x_1, x_2) \leq 2^d \text{dist}_{T, T'}^{x'}(x_1, x_2).$$

Proof. The inequality $\text{dist}_{T, T'}^{x'}(x_1, x_2) \leq \text{dist}(x_1, x_2)$ is immediate from the definitions. For the second inequality we assume $x_1 \neq x_2$.

Let i, j be such that $T(i) = T'(i) \setminus \{j\}$.

Since $(x_1)_{T(i)}$ and $(x_2)_{T(i)}$ are distinct codimension one subspaces of $x'_{T'(i)}$ their sum is $x'_{T'(i)}$, and therefore

$$\text{dist}_{T, T'}^{x'}(x_1, x_2) = \text{dist}((x_1)_{T(i)}, (x_2)_{T(i)}) \geq \text{dist}((x_1)_{T(i)} + W, (x_2)_{T(i)} + W),$$

for all subspaces W containing neither $(x_1)_{T(i)}$ nor $(x_2)_{T(i)}$.

For each $I \in T \setminus T'$ one has $i \in I$ and therefore $T(i) \subset I$. Noticing that $J := \bigcup_{k \in I \setminus \{i\}} T(k) = \bigcup_{k \in I \setminus \{i\}} T'(k)$ belongs to T' we obtain

$$\begin{aligned} \text{dist}(x_1, x_2) &= \sum_{I \in T \setminus T'} \text{dist}((x_1)_I, (x_2)_I) \\ &= \sum_{I \in T \setminus T'} \text{dist}((x_1)_{T(i)} + x'_J, (x_2)_{T(i)} + x'_J) \leq 2^d \text{dist}_{T, T'}^{x'}(x_1, x_2). \end{aligned}$$

□

Consider T, T' admissible topologies with $T \stackrel{1}{\prec} T'$, ($i < j$) such that $T'(i) = T(i) \cup \{j\}$. Given $x \in \mathcal{X}_T$, we use the metric (11) to define a bilipschitz homeomorphism $\varphi_x : (-\pi/2, \pi/2) \rightarrow \pi_{T, T'}^{-1}(x')$ where $x' = \pi_{T, T'}(x)$.

For this purpose let $V = x'_{T'(i)} / x'_{T'(i) \setminus \{i, j\}}$ endowed with the inner product inherited from the ambient space \mathbb{R}^d (i.e. the inner product between two classes is calculated by taking representatives perpendicular to $x'_{T'(i) \setminus \{i, j\}}$).

One has that $(\pi_{T, T'})^{-1}(x')$ consists in configurations z with $z_{T'(k)} = x'_{T'(k)}$ for all $k \neq i$ while $z_{T(i)}$ is a codimension one subspace of $x'_{T'(i)}$ which contains $x'_{T'(i) \setminus \{i, j\}}$ and is distinct from $x'_{T'(i) \setminus \{i\}}$. It follows that $(\pi_{T, T'})^{-1}(x')$ endowed with $\text{dist}_{T, T'}^{x'}$ is isometric to the space of one dimensional subspaces of V minus the projection of $x'_{T'(i) \setminus \{i\}}$ with the angle distance.

Let X, Y be a pair of unit vectors in V such that X has a representative in $x_{T(i)}$, Y has a representative in $x'_{T'(i) \setminus \{i\}}$ and such that $\cos \angle(X, Y) > 0$. Define $\varphi_x : (-\pi/2, +\pi/2) \rightarrow \pi_{T, T'}^{-1}(x')$ by associating to $u \in (-\pi/2, +\pi/2)$ the configuration where the corresponding one dimensional subspace of V contains a vector of the form $\cos u X + \sin u Y$. Set $\theta := \angle(X, Y)$.

Lemma 3.4. *In the above context φ_x is a bilipschitz homeomorphism. Moreover,*

$$(12) \quad \left| \tan \frac{\theta}{2} \right| < \text{Lip}(\varphi_x) < \frac{1}{\left| \tan \frac{\theta}{2} \right|}.$$

Proof. Let $e_1 = (1, 0), e_2 = (0, 1)$ be the standard basis of \mathbb{R}^2 and $\mathcal{G}_1(\mathbb{R}^2)$ be the space of one dimensional subspaces with the angle distance. Let L_2 be the subspace generated by e_2 .

The mapping $f : (-\pi/2, \pi/2) \rightarrow \mathcal{G}_1(\mathbb{R}^2) \setminus \{L_2\}$ where $f(u)$ is the subspace generated by $\cos(u)e_1 + \sin(u)e_2$ is an isometry.

By [Les21, Lemma 2], if $\theta \in (0, \pi/2]$ then the mapping $g_\theta : \mathcal{G}_1(\mathbb{R}^2) \setminus \{L_2\}$ induced by the matrix $A = \begin{pmatrix} \sin(\theta) & 0 \\ \cos(\theta) & 1 \end{pmatrix}$ has derivative $|dg_\theta(L)| = \frac{|\det(A)|}{\|A|_L\|^2}$.

This implies for the composition one has

$$|dg_\theta \circ f(u)| = \frac{|\sin(\theta)|}{\cos(u)^2 \sin(\theta)^2 + (\cos(u) \cos(\theta) + \sin(u))^2} = \frac{|\sin(\theta)|}{1 + \sin(2u) \cos(\theta)}.$$

Letting $\theta = \text{dist}(x_{T(i)}, x'_{T'(i) \setminus \{i\}})$ the fiber $\pi_{T, T'}^{-1}(x')$ endowed with $\text{dist}_{T, T'}^{x'}$ is isometric to the projective space $\mathcal{G}_1(\mathbb{R}^2) \setminus \{L_2\}$ where x is identified with the subspace generated by $\sin(\theta)e_1 + \cos(\theta)e_2$.

Under this identification φ_x corresponds to the composition $g_\theta \circ f$, so we have obtained

$$|d\varphi_x(u)| = \frac{|\sin(\theta)|}{1 + \sin(2u) \cos(\theta)}.$$

□

To finish the proof of proposition 3.2, we still have to show that the homeomorphism $\varphi_z, z \in (\pi_{T^t, T'})^{-1}(x')$ depends continuously on z as a bilipschitz homeomorphism. Given the above construction, this is true since we can locally choose in a continuous way parameterizations of the spaces

$$V_z = [\pi_{T^t, T^{t-1}}(z)]_{T^{t-1}(i)} / [\pi_{T^t, T^{t-1}}(z)]_{T^{t-1}(i) \setminus \{i, j\}}$$

and in these spaces vectors X_z, Y_z such that X_z has a representative in $z_{T^t(i)}$, Y_z has a representative in $[\pi_{T^t, T^{t-1}}(z)]_{T^{t-1}(i) \setminus \{i, j\}}$ and $\cos \angle(X_z, Y_z) > 0$. □

Proof of proposition 2.1.2. For $T \prec T'$ admissible topologies, it follows from proposition 3.2 that $\bar{\delta}_{T, T'} \leq N_{T, T'}$. Also since $T^t \stackrel{1}{\prec} T^{t-1}$, we have $\bar{\gamma}_{T^t, T^{t-1}} \leq 1$. By formula (7), we have indeed $\kappa_{T, T'} \leq \sum_{(i, j) \in D_{T, T'}} (\chi_i - \chi_j)$.³ □

3.3. One-dimensional coordinates. Let $i < j$ and consider all arrows T, T' of admissible topologies with $T \stackrel{1}{\prec} T'$, such that $T'(i) = T(i) \cup \{j\}$. Recall that $T_{i, j}$ is the topology defined by

$$T_{i, j}(k) = \{k\} \text{ if } k \neq i, T_{i, j}(i) = \{i, j\}.$$

The topology $T_{i, j}$ is admissible and one step coarser than T_1 .

Lemma 3.5. *Let $T \stackrel{1}{\prec} T'$ be admissible topologies and i, j be such that $T(i) = T'(i) \setminus \{j\}$. For all y', x' such that $\pi_{T_{i, j}, T'} y' = x'$, $\pi_{T_1, T}$ defines a bilipschitz homeomorphism between $(\pi_{T, T'})^{-1}(x')$ and $(\pi_{T_1, T_{i, j}})^{-1}(y')$. The Lipschitz constants depend only on y' .*

Proof. Consider $y_1, y_2 \in \mathcal{X}_{T_1}$, distinct and such that $\pi_{T, T_{i, j}}(y_1) = \pi_{T, T_{i, j}}(y_2) = y'$. Set $x_1 = \pi_{T_1, T}(y_1)$ and $x_2 = \pi_{T_1, T}(y_2)$ and notice that $\pi_{T, T'}(x_1) = \pi_{T, T'}(x_2) = \pi_{T_{i, j}, T'}(y') = x'$. Since $\pi_{T_1, T}$ consists of forgetting some subspaces of each sequence $(x_I)_{I \in T_1}$ it is 1-Lipschitz, and in particular 1-Lipschitz as a mapping between $\pi_{T_1, T_{i, j}}^{-1}(y')$ and $\pi_{T, T'}^{-1}(x')$.

To prove that the inverse is also Lipschitz let $S_1 = (y_1)_{\{i\}}, S_2 = (y_2)_{\{i\}}$. Notice that S_1, S_2 are distinct one dimensional subspaces of the two dimensional subspace $y'_{\{i, j\}}$. Therefore $S_1 + S_2 = y'_{\{i, j\}}$ and $\text{dist}(S_1, S_2) = \angle(S_1, S_2)$.

For each $I \in T_1 \setminus T_{i, j}$ notice that $i \in I$ and $j \notin I$. Setting $W_I = y'_{I \setminus \{i\}}$ we have $(y_1)_I = W_I + S_1$ and $(y_2)_I = W_I + S_2$ from which it follows that

$$\text{dist}((y_1)_I, (y_2)_I) = \angle(\pi_{W_I^\perp}(S_1), \pi_{W_I^\perp}(S_2)) \leq \angle(S_1, S_2),$$

where $\pi_{W^\perp} : \mathbb{R}^d \rightarrow W^\perp$ is the orthogonal projection onto W^\perp . By Lemma 3.3,

$$\text{dist}(y_1, y_2) = \sum_{I \in T_1 \setminus T_{i, j}} \text{dist}((y_1)_I, (y_2)_I) \leq 2^d \angle(S_1, S_2).$$

³Observe that relation (7) depends on theorem 1.6, which will be proven in section 5.

Because y' is a configuration, the minimum over $I \in T_1 \setminus T_{i,j}$ of the angle between W_I and $S_1 + S_2 = y'_{\{i,j\}}$ is positive. It follows that there exists $c > 0$ which depends only on y' such that

$$\text{dist}((y_1)_I, (y_2)_I) = \angle(\pi_{W_I^\perp}(S_1), \pi_{W_I^\perp}(S_2)) \geq c\angle(S_1, S_2),$$

for all $I \in T_1 \setminus T_{i,j}$.

Notice that, since $T_1 \setminus T_{i,j} = \{I : I \ni i \text{ and } I \not\ni j\}$, $T \setminus T' \subset T_1 \setminus T_{i,j}$, and therefore

$$\text{dist}(x_1, x_2) = \sum_{I \in T \setminus T'} \angle(\pi_{W_I^\perp}(S_1), \pi_{W_I^\perp}(S_2)) \geq 2^{-d}c \text{dist}(y_1, y_2).$$

It follows that $\pi_{T_1, T}$ is a bilipschitz homeomorphism between $\pi_{T_1, T_{i,j}}^{-1}(y')$ and $\pi_{T, T'}^{-1}(x')$, as claimed. \square

For $x \in \mathcal{X}_T$, $T \xrightarrow{1} T'$ admissible topologies, lemma 3.4 yields a Lipschitz homeomorphism between $\pi_{T, T'}^{-1}(x')$ where $x' = \pi_{T, T'}(x)$ and $(-\frac{\pi}{2}, \frac{\pi}{2})$ and the Lipschitz constant depends on x . Combining with lemma 3.5 yields

Corollary 3.6. *Assume the diagram of projections*

$$\begin{array}{ccc} T & \longrightarrow & S \\ \downarrow 1 & & \downarrow 1 \\ T' & \longrightarrow & S' \end{array} \quad \text{commutes and}$$

i, j are such that $T(i) = T'(i) \setminus \{j\}$ and $S(i) = S'(i) \setminus \{j\}$. Then, for $x \in \mathcal{X}_T$, there is a bilipschitz homeomorphism between $(\pi_{T, T'})^{-1}(\pi_{T, T'}(x))$ and $(\pi_{S, S'})^{-1}(\pi_{T, S'}(x))$. The Lipschitz constant depends only on x .

4. APPLICATIONS OF OSELEDETS THEOREM

4.1. Oseledets multiplicative ergodic theorem. We review in this section some applications of Oseledets multiplicative ergodic theorem to random walks on matrices. Let μ be a probability measure on the group $G = SL_d(\mathbb{R})$ of $d \times d$ matrices with determinant 1. Let $(\Omega, m) = (G^{\mathbb{Z}}, \mu^{\mathbb{Z}})$ be the probability space of independent trials of elements of G with distribution μ , σ the shift transformation on Ω . Let $\omega \in \Omega$ be the sequence $(g_n)_{n \in \mathbb{Z}}$. We denote g_n the mapping $\omega \mapsto g_n(\omega)$ that associates to $\omega \in \Omega$ its coordinate $g_n \in G$. In particular $g_0(\omega)$ defines a cocycle with values in $SL(d, \mathbb{R})$ over the ergodic system $(\Omega, m; \sigma)$. Oseledets multiplicative ergodic theorem gives

Theorem 4.1 ([O68]). *With the above notations, assume that $\mu \in \mathcal{M}(d)$ and let $\chi_1 > \chi_2 > \dots > \chi_d$ with $\sum_i \chi_i = 0$ be the Lyapunov exponents (cf. equation (1)). Then, for m -a.e. $\omega \in \Omega$, there exists a direct decomposition of \mathbb{R}^d into d one-dimensional spaces*

$$\mathbb{R}^d = E_1(\omega) \oplus E_2(\omega) \oplus \dots \oplus E_d(\omega) \quad \text{such that}$$

(1) *a vector $v \neq 0$ belongs to $E_i(\omega)$ if, and only if,*

$$\lim_{n \rightarrow +\infty} \frac{1}{n} \log |g_{n-1}(\omega) \circ \dots \circ g_0(\omega) v| = \lim_{n \rightarrow -\infty} \frac{1}{n} \log |(g_n(\omega))^{-1} \circ \dots \circ (g_{-1}(\omega))^{-1} v| = \chi_i$$

(2) *for m -a.e. ω , all i , $\lim_{n \rightarrow \pm\infty} \frac{1}{|n|} \log |\sin \angle(E_i(\sigma^n \omega), \sum_{j \neq i} E_j(\sigma^n \omega))| = 0$.*

By our assumption $\mu \in \mathcal{M}(d)$, the exponents are indeed pairwise distinct and the spaces E_i are one-dimensional. The directions $E_i(\omega)$ are defined m -a.e. and are called *Oseledets directions*. For two distinct dimensional subspaces $E, E' \subset \mathbb{R}^d$, $|\sin \angle(E, E')|$ is defined by $|\sin \angle(E, E')| = \inf_{v \neq 0 \in E, v' \neq 0 \in E'} \frac{|v \wedge v'|}{|v||v'|}$.

The set Ω_{reg} of points in Ω such that properties 1 and 2 of Oseledets theorem hold is called the set of regular points. The set Ω_{reg} is σ -invariant, measurable and has full measure in Ω . Observe that by characterization 1, for all i , the mapping $\omega \mapsto E_i(\omega)$ is measurable on Ω_{reg} and we have

$$E_i(\sigma\omega) = g_0 E_i(\omega).$$

Let $i = 1, \dots, d$. We write $U_i(\omega) := \bigoplus_{j=1}^i E_j(\omega)$ for the unstable spaces, $U'_i(\omega) := \bigoplus_{j=d-i+1}^d E_j(\omega)$ for the stable spaces.⁴ Since the exponents are distinct, for $\omega \in \Omega_{reg}$, the flags $E_-(\omega)$ given by $\{0\} = U_0 \subset U_1(\omega) \subset \dots \subset U_d = \mathbb{R}^d$ and $E_+(\omega)$ given by $\{0\} = U'_0 \subset U'_1(\omega) \subset \dots \subset U'_{d-1}(\omega) \subset U'_d = \mathbb{R}^d$ are in general position.

An important classical observation is the following:

Proposition 4.2. *For all i , the mappings $\omega \mapsto U_i(\omega)$ are measurable with respect to the σ -algebra generated by $(g_n)_{n \leq -1}$; for all i' , the mappings $\omega \mapsto U'_{i'}(\omega)$ are measurable with respect to the σ -algebra generated by $(g_n)_{n \geq 0}$. In particular, E_- and E_+ are independent.*

Proof. It suffices to prove that for any i , U_i depends only on $\{g_n\}_{n \leq -1}$. We claim that, for $\omega \in \Omega_{reg}$, $U_i = \{v : \limsup_{n \rightarrow +\infty} \frac{1}{n} \log |g_{-n}^{-1} \circ \dots \circ g_{-1}^{-1} v| \leq -\chi_i\}$. This shows that U_i is completely determined when one knows $\{g_n\}_{n \leq -1}$. To prove the claim, observe that $\{v : \limsup_{n \rightarrow +\infty} \frac{1}{n} \log |g_{-n}^{-1} \circ \dots \circ g_{-1}^{-1} v| \leq -\chi_i\}$ is a vector space that contains $E_j(\omega)$, $j \leq i$ by definition. It is exactly $U_i(\omega)$ since any vector that has a nonzero component in one of the $E_\ell(\omega)$, $\ell > i$ satisfies $\limsup_{n \rightarrow +\infty} \frac{1}{n} \log |g_{-n}^{-1} \circ \dots \circ g_{-1}^{-1} v| \geq -\chi_\ell > -\chi_i$. One verifies in the same way that $U'_{i'}(\omega) = \{v : \limsup_{n \rightarrow +\infty} \frac{1}{n} \log |g_n \circ \dots \circ g_0 v| \leq \chi_{i'}\}$. \square

Let Q be a partition of $\{0, 1, \dots, d\}$ into intervals, $Q = \{q_0 = 0 < q_1 < \dots < q_k = d\}$. Write $U_Q(\omega) \in \mathcal{F}_Q$ for the Q -flag

$$U_Q(\omega) := \{0\} = U_0 \subset U_{q_1}(\omega) \subset \dots \subset U_{q_{k-1}}(\omega) \subset U_d = \mathbb{R}^d.$$

The set \mathcal{G}_i of i -dimensional subspaces is identified with $\mathcal{F}_{\{0 < i < d\}}$.

Since g_0 is independent of $U_i(\omega), U_Q(\omega)$ and the distribution of g_0 is μ , the distribution of U_i (resp. U_Q) is a measure on \mathcal{G}_i (resp. \mathcal{F}_Q) which is stationary under the action of (G, μ) . By uniqueness, the distribution of $U_i(\omega)$ is $\nu_{\{0 < i < d\}}$, the distribution of $U_Q(\omega)$ is ν_Q . Similarly, the distribution of E_- is a stationary measure ν' for the action of μ' on \mathcal{F} . For all partition Q of $\{0, 1, \dots, d\}$ into intervals, the distribution of $U'_Q(\omega)$ is the stationary measure $\nu'_Q := (\pi_Q)_* \nu'$ for the action of μ' on \mathcal{F}_Q .

⁴cf. notations of the Introduction.

Remark 4.3. *We do not know a priori that the measure μ' has a unique stationary measure; in all the paper, we use the distribution ν' of stable flags as stationary measure for the action of μ' on \mathcal{F} .*

Let $\Omega_+ := (g_n)_{n \geq 0}$ (respectively $\Omega_- := (g_n)_{n \leq -1}$) be the space of one-sided sequences of elements of G , m_+ (respectively m_-) the product measure with g_k of distribution μ for all $k \geq 0$ (respectively for all $k < 0$), σ the shift transformation. Then, by proposition 4.2, E_- (respectively E_+) can be seen as a mapping from Ω_- (respectively Ω_+) into \mathcal{F} . By Oseledets theorem 4.1, for almost every ω , the pair $E(\omega) := (E_-(\omega_-), E_+(\omega_+))$ belongs to $\mathcal{F}^{(2)}$.

We recall in our notations the key Furstenberg result

Theorem 4.4 ([F63]). *Assume $\mu \in \mathcal{M}(d)$. Let $f \in \mathcal{F}$, $f = \{\{0\} \subset U_1(f) \subset \dots \subset U_{d-1}(f) \subset \mathbb{R}^d\}$. For m -a.e. $\omega \in \Omega_+$, all $j, j = 1, \dots, d$,*

$$\lim_{n \rightarrow +\infty} \frac{1}{n} \log |\det_{U_j(f)}(g_{n-1} \circ \dots \circ g_0)| = \sum_{i \leq j} \chi_i.$$

Proof. Under our hypotheses, the distributions of all exterior products $\wedge_{i=1}^j g$ satisfy the conditions of Theorem 8.5 in [F63]. \square

We used these relations in the introduction to define the exponents $\chi_j, j = 1, \dots, d$, in general for ν extremal. Since $\mu \in \mathcal{M}(d)$, the stationary ν measure is extremal and thus the skew product $(\Omega_+ \times \mathcal{F}, m_+ \otimes \nu)$ is ergodic for the transformation $\hat{\sigma} : \hat{\sigma}(\omega_+, f) = (\sigma\omega_+, g_0(\omega)f)$. We can write

$$\begin{aligned} \sum_{i \leq j} \chi_i &= \lim_{n \rightarrow +\infty} \frac{1}{n} \log |\det_{U_j(f)}(g_{n-1} \circ \dots \circ g_0)| \\ &= \lim_{n \rightarrow +\infty} \frac{1}{n} \sum_{k=1}^{n-1} \log |\det_{U_j(\hat{\sigma}^k(\omega_+, f))}(g_0(\sigma^k \omega))|. \end{aligned}$$

The last line converges to $\int \log |\det_{U_j(f)}(g)| d\mu(g) d\nu(f)$ by the ergodic theorem, so that

$$(13) \quad \sum_{i \leq j} \chi_i = \int \log |\det_{U_j(f)}(g)| d\mu(g) d\nu(f).$$

We define the Lyapunov exponents by equation (13) when the measure ν is not extremal. When the distribution of $E_-(\omega)$ is ν (in particular, for $\mu \in \mathcal{M}(d)$) and since g_0 is independent of $E_-(\omega)$, then (13) can also be written, for all $j, j = 1, \dots, d$,

$$\sum_{i \leq j} \chi_i = \int_{\Omega} \log (|\det_{U_j(\omega)}(g_0(\omega))|) dm(\omega).$$

Using property 2 in Oseledets multiplicative ergodic theorem 4.1, we get, in that case, for any subset $I \subset \{1, \dots, d\}$, setting $V_I(\omega) = \bigoplus_{k \in I} E_k(\omega)$,

$$(14) \quad \int_{\Omega} \log (|\det_{V_I(\omega)}(g_0(\omega))|) dm(\omega) = \sum_{k \in I} \chi_k.$$

4.2. Approximation of partial Oseledets configurations. Given an admissible topology T we define $E_T(\omega) = F_T(E_-(\omega), E_+(\omega))$ where F_T is defined in section 1.4.

Let $T \prec T'$ be admissible topologies. We will extend the configuration $E_{T'}(\omega)$ to a configuration $\widehat{E}(\omega)$ defined on T by applying a deterministic function to $E_{T'}(\omega)$.

For this purpose, for each $i = 1, \dots, d$, we let $\widehat{E}_i(\omega)$ be the one dimensional subspace of $E_{T'}(\omega)_{T'(i)}$ which is perpendicular to $E_{T'}(\omega)_{T'(i) \setminus \{i\}}$. The configuration $\widehat{E}(\omega)$ is defined by letting

$$\widehat{E}(\omega)_I = \sum_{i \in I} \widehat{E}_i(\omega)$$

for all $I \in T$.

Proposition 4.5. *For m -a.e. ω , and all $I \in T'$ one has $\widehat{E}(\omega)_I = E_{T'}(\omega)_I$.*

Proof. It suffices to verify that $\widehat{E}(\omega)_{T'(i)} = E_{T'}(\omega)_{T'(i)}$ for $i = 1, \dots, d$.

When $i = d$, this is trivial since $E_{T'}(\omega)_{T'(d)} = E_d(\omega)$ and therefore $\widehat{E}(\omega)_{T'(d)} = E_d(\omega)$ as well.

Suppose that the claim is true for $i + 1, \dots, d$, then

$$\widehat{E}(\omega)_{T'(i) \setminus \{i\}} = \sum_{j \in T'(i) \setminus \{i\}} \widehat{E}(\omega)_{T'(j)} = \sum_{j \in T'(i) \setminus \{i\}} E(\omega)_{T'(j)} = E(\omega)_{T'(i) \setminus \{i\}}.$$

It follows that $\widehat{E}_i(\omega)$ is complementary to the codimension one subspace $\widehat{E}(\omega)_{T'(i) \setminus \{i\}}$ within $E(\omega)_{T'(i)}$. Taking the sum one obtains $\widehat{E}(\omega)_{T'(i)} = E_{T'}(\omega)_{T'(i)}$, as claimed. \square

We now show that using the extension above one may approximate $E_T(\omega)$ using only $E_{T'}(\omega)$ and $g_{-1}(\omega), \dots, g_{-n}(\omega)$ up to an error which is exponentially small as $n \rightarrow +\infty$.

Proposition 4.6. *For m -a.e. ω when $n \rightarrow +\infty$ one has*

$$\text{dist}(g_{-1}(\omega) \cdots g_{-n}(\omega) \widehat{E}(\sigma^{-n}(\omega))_T, E_T(\omega)) \leq \exp(-\chi n + o(n)),$$

where $\chi = \min_{(i,j) \in D_{T,T'}} \chi_i - \chi_j$.

Proof. To simplify calculations for each $i = 1, \dots, d$ let $w_i(\omega)$ be a unit vector generating $E_i(\omega)$, and $v_i(\omega)$ be a unit vector generating $\widehat{E}_i(\omega)$. Write

$$v_i(\omega) = \sum_{j \in T'(i)} a_{i,j}(\omega) w_j(\omega).$$

Since $v_i(\sigma^{-n}\omega)$ is a unit vector $|a_{i,j}(\sigma^{-n}\omega)| \leq e^{o(n)}$ for all j , all n , and m -a.e. ω . Furthermore, since the angle between $E_i(\sigma^{-n}\omega)$ and $\sum_{j \in T'(i) \setminus \{i\}} E_j(\sigma^{-n}\omega)$ is at least $e^{-o(n)}$, we obtain $|a_{i,i}(\sigma^{-n}\omega)| \geq e^{-o(n)}$ when $n \rightarrow +\infty$ for m -a.e. ω .

It suffices to show that

$$\text{dist}(g_{-1}(\omega) \cdots g_{-n}(\omega) \widehat{E}(\sigma^{-n}(\omega))_{T(i)}, E_T(\omega)_{T(i)}) \leq \exp(-\chi n + o(n)),$$

when $n \rightarrow +\infty$ for all $i = 1, \dots, d$.

The claim is trivially true when $i = d$.

Suppose that the claim is true for $i + 1, \dots, d$, so in particular one has that

$$\text{dist}(g_{-1}(\omega) \cdots g_{-n}(\omega) \widehat{E}(\sigma^{-n}(\omega))_J, E_T(\omega)_J) \leq \exp(-\chi n + o(n)),$$

as $n \rightarrow +\infty$ where $J = T(i) \setminus \{i\}$.

Let $z_n(\omega)$ be a unit vector in the intersection of $\widehat{E}(\sigma^{-n}(\omega))_{T(i)}$ with the subspace $W_i(\sigma^{-n}(\omega))$, where $W_i(\omega)$ is the space generated by $\{w_j(\omega), j = i \text{ and } j \in T'(i) \setminus T(i)\}$. We have $e^{-o(n)} \leq |\langle z_n, w_i(\sigma^{-n}(\omega)) \rangle|$ and $\|z_n\| \leq 1$. If we write

$$z_n(\omega) = b_{n,i}(\omega)w_i(\sigma^{-n}\omega) + \sum_{j \in T'(i) \setminus T(i)} b_{n,j}(\omega)w_j(\sigma^{-n}(\omega)),$$

we have $|b_{n,i}(\omega)| \geq e^{-o(n)}$ while $|b_{n,j}(\omega)| \leq e^{o(n)}$ for all j .

To conclude notice that

$$g_{-1}(\omega) \cdots g_{-n}(\omega)z_n(\omega) = e^{\chi_i n + o(n)}b_{n,i}(\omega)w_i(\omega) + \sum_{j \in T'(i) \setminus T(i)} e^{\chi_j n + o(n)}b_{n,j}(\omega)w_j(\omega).$$

It follows that the angle between $E_i(\omega)$ and the subspace $L_n(\omega)$ generated by $g_{-1}(\omega) \cdots g_{-n}(\omega)z_n(\omega)$ is at most $e^{-\chi' n + o(n)}$ where $\chi' = \min_{j \in T'(i) \setminus T(i)} \chi_i - \chi_j \geq \chi$. Since $g_{-1}(\omega) \cdots g_{-n}(\omega) \widehat{E}(\omega)_{T(i)} = L_n(\omega) + g_{-1}(\omega) \cdots g_{-n}(\omega) \widehat{E}(\omega)_J$, the claim follows. \square

Assume in the above discussion that $T \stackrel{1}{\prec} T'$ and that $i < j$ is such that $T'(i) = T(i) \cup \{j\}$. Set $x' = E_{T'}(\omega)$. Then the space $W_i(\omega)$, generated by $E_i(\omega), E_j(\omega)$ is a representative of the vector space $V = x'_{T'(i)} / x'_{T'(i) \setminus \{i,j\}}$ discussed in Lemma 3.4. Let $\pi = \pi_{T,T'}$ be the projection from \mathcal{X}_T to $\mathcal{X}_{T'}$ and consider the coordinates $\varphi_{x(\omega)}$ given by lemma 3.4 on $\pi^{-1}(E_{T'}(\omega))$, setting $x(\omega) = E_T(\omega)$ and $x'(\omega) = E_{T'}(\omega)$. The distance $\varphi_{x(\omega)}$ on $W_i(\omega)$ is equivalent to the metric $\text{dist}_{T,T'}^{x'(\omega)}$ (see (11)) on $\pi^{-1}(x'(\omega))$.

Lemma 4.7. *Let $T \stackrel{1}{\prec} T'$ be admissible topologies and $i < j$ such that $T'(i) = T(i) \cup \{j\}$. Fix $\beta > 0$ and let $x_n \in \pi^{-1}(x'(\sigma^{-n}\omega))$ satisfy $\varphi_{x(\sigma^{-n}\omega)}(x_n) \leq \beta$. Then for m -a.e. ω , as $n \rightarrow \infty$, one has*

$$(15) \quad \text{dist}_{T,T'}^{E_{T'}(\omega)}(g_{-1}(\omega) \circ \dots \circ g_{-n}(\omega)(x_n), E_T(\omega)) \leq \exp(-\chi_{T,T'} n + o(n)).$$

Proof. By theorem 4.1.2, the distance from $E_T(\sigma^{-n}\omega)$ to $E_{T'}(\sigma^{-n}\omega)$ is at least $\exp(-o(n))$. Therefore, $\text{dist}_{T,T'}^{E_{T'}(\sigma^{-n}\omega)}((x_n), E_{T'}(\sigma^{-n}\omega)) \geq \exp(-o(n))$ as well. Following the proof of proposition 4.6, we have that the point x_n satisfies

$$\text{dist}_{T,T'}^{E_{T'}(\omega)}(g_{-1}(\omega) \circ \dots \circ g_{-n}(\omega)(x_n), E_T(\omega)) \leq \exp(-\chi_{T,T'} n + o(n)).$$

The lemma follows. \square

5. PROOF OF THEOREM 1.1

Theorem 1.1 states that some entropy is estimated from above by exponents. For random walks on matrices, this is a fundamental observation of Furstenberg ([F63]). Theorem 1.1 and its proof are one more variant of the original proof: one shows equality for a big family of random walks on the same group and one approximates using this family. The exponents are continuous and the entropy has some upper semi-continuous properties. This should be sufficient for the astute reader, but we will give a detailed proof anyway. In particular, it gives some a priori quasi-invariance of stationary measures (see Corollary 5.9). We state and prove the generalisation of theorem 1.1 to the action on \mathcal{F}_Q , for any Q partition de $\{0, 1, \dots, d\}$.

Theorem 5.1. *With the above notations, for any partition Q of $\{0, 1, \dots, d\}$ into intervals, there exists a stationary measure ν_Q on \mathcal{F}_Q such that*

$$(16) \quad h(\mathcal{F}_Q, \mu, \nu_Q) \leq \sum_{i,j: \ell_Q(i) < \ell_Q(j)} \chi_i - \chi_j.$$

If there is equality in (16), then the measure ν_Q is exact dimensional with dimension $\dim \mathcal{F}_Q$.

For $\mu \in \mathcal{M}(d)$, the stationary measure ν_Q is unique and thus satisfies (16).

5.1. Mollification of μ . For each $n = 1, 2, 3, \dots$ fix a probability λ_n with a smooth positive density with respect to Haar measure on the orthogonal group of \mathbb{R}^d , in such a way that $\lim_{n \rightarrow +\infty} \lambda_n = \delta_{Id}$ where δ_{Id} is the point mass at the identity.

Let $\mu_n = \lambda_n * \mu$ so one has, for all continuous $h : G \rightarrow \mathbb{R}$

$$(17) \quad \int_G h(g) d\mu_n(g) = \int_G \int_G h(rg) d\lambda_n(r) d\mu(g).$$

Let η be the orthogonally invariant probability on \mathcal{F} .

Lemma 5.2. *For each n there is a unique μ_n -stationary probability ν_n on \mathcal{F} . Furthermore, ν_n has a continuous positive density with respect to η .*

Proof. For any μ_n -stationary probability we have

$$\nu_n = \mu_n * \nu_n = (\lambda_n * \mu) * \nu_n = \lambda_n * (\mu * \nu_n).$$

Since, for any probability m on \mathcal{F} , the convolution $\lambda_n * m$ has a continuous positive density with respect to η it follows that any μ_n -stationary probability has this property. However, any two distinct extremal μ_n -stationary probabilities must be mutually singular. This implies that ν_n is unique as claimed. \square

Endow $\mathcal{M}(G \times \mathcal{F})$ with the topology of convergence over continuous function φ on $(G \times \mathcal{F})$ with $|\varphi(g, f)| \leq C \log \|g\|$ for some constant C .

Lemma 5.3. *In $\mathcal{M}(G \times \mathcal{F})$, any limit $\lim_{n \rightarrow +\infty} (\mu_n \times \nu_n)$ is of the form $\mu \times \nu$ for some stationary ν on \mathcal{F} .*

Proof. We have $\lim_{n \rightarrow \infty} \mu_n = \mu$ and, since any weak*-limit of ν_n is μ -stationary, $\lim_{n \rightarrow \infty} (\mu_n \times \nu_n)$ is of the form $\mu \times \nu$ for some μ -stationary ν on \mathcal{F} . A priori, we have convergence against only continuous functions with compact support on $G \times \mathcal{F}$. We want to ensure that there is also convergence in $\mathcal{M}(G \times \mathcal{F})$. We may assume that $\{\mu_n \times \nu_n\}_{n \rightarrow \infty}$ is the converging sequence.

By Skorohod's representation theorem, there exists a probability space $(P, \mathcal{E}, \mathbb{P})$ and random elements on this space such that

$$\lim_{n \rightarrow +\infty} (A_n, F_n) = (A, F)$$

almost surely, where (A, F) has distribution $\mu \times \nu$ and (A_n, F_n) has distribution $\mu_n \times \nu_n$ for each n . Let φ be a continuous function φ on $(G \times \mathcal{F})$ with $|\varphi(g, f)| \leq C \log \|g\|$ for some constant C .

With these auxiliary random elements we calculate (denoting integration with respect to \mathbb{P} by $\mathbb{E}(\cdot)$ as usual)

$$\lim_{n \rightarrow +\infty} \int \varphi(g, f) d\mu_n(g) d\nu_n(f) = \lim_{n \rightarrow +\infty} \mathbb{E}(\varphi(A_n, F_n)).$$

Observe that, since left composition with an orthogonal transformation does not change the norm a linear mapping, the random variables $(|\varphi(A_n, F_n)|)_{n=1,2,\dots}$ are uniformly integrable. It follows that

$$\lim_{n \rightarrow +\infty} \mathbb{E}(\varphi(A_n, F_n)) = \mathbb{E}(\varphi(A, F)).$$

□

Lemma 5.4. *For each n and each $i = 1, \dots, d$ let $\chi_{i,n}$ be such that*

$$\sum_{j \leq i} \chi_{j,n} := \int_G \int_{\mathcal{F}} \log(|\det_{U_i(f)}(g)|) d\nu_n(f) d\mu_n(g)$$

. Then one has $\lim_{n \rightarrow +\infty} \chi_{i,n} = \chi_i$ for each $i = 1, \dots, d$.

Proof. Recall the definition of the Lyapunov exponents by equation (13). We may apply the preceding lemma since $|\det_{U_i(f)}(g)| \leq d! |\log g|$. □

5.2. Quasi-independence and mutual information. Let (Ω, \mathbb{P}) be a probability space. Given $X : \Omega \rightarrow \mathcal{X}$ and $Y : \Omega \rightarrow \mathcal{Y}$ taking values in Polish spaces, we say X and Y are quasi-independent if the distribution $\mathbb{P}_{X,Y}$ of X, Y is absolutely continuous with respect to the product $\mathbb{P}_X \otimes \mathbb{P}_Y$ of the distributions of X and Y . We write $f(x, y)$ or $f_{\mathbb{P}}(x, y)$ for the Radon-Nikodym derivative $\frac{d\mathbb{P}_{X,Y}}{d(\mathbb{P}_X \otimes \mathbb{P}_Y)}$. In that case, the disintegration of the measure $\mathbb{P}_{X,Y}$ with respect to the projections over X and Y is respectively $\mathbb{P}^x = f(x, y) \mathbb{P}_Y$ and $\mathbb{P}^y = f(x, y) \mathbb{P}_X$.

For X and Y as above, define the mutual information between X and Y as

$$I(X, Y) = I_{\mathbb{P}}(X, Y) = \sup \sum_{A \in \mathcal{A}} \log \left(\frac{\mathbb{P}_{(X,Y)}(A)}{(\mathbb{P}_X \times \mathbb{P}_Y)(A)} \right) \mathbb{P}_{(X,Y)}(A),$$

where the supremum is over finite Borel partitions \mathcal{A} of $\mathcal{X} \times \mathcal{Y}$ ⁵.

Directly from the definition one sees that $I(X, Y) = I(Y, X)$. By Jensen inequality $0 \leq I(X, Y) \leq +\infty$ with equality to 0 if and only if X and Y are independent. If X takes countably many values and has finite entropy $H(X)$ in the sense of [Sha48] one has $I(X, Y) \leq H(X)$. It was shown in [Dob59] that $I(X, Y)$ is the supremum over any sequence of partitions which generate the Borel σ -algebra in $\mathcal{X} \times \mathcal{Y}$.

It was shown in [GfY] and [Per59] that if $I(X, Y) < +\infty$ then X and Y are quasi-independent and

$$I(X, Y) = \int \log f(x, y) d\mathbb{P}_{X, Y}(x, y) = \int f(x, y) \log f(x, y) d\mathbb{P}_X(x) d\mathbb{P}_Y(y).$$

Let Q be a partition of $\{0, 1, \dots, d\}$ into intervals and $\pi_Q : \mathcal{F} \rightarrow \mathcal{F}_Q$ the projection from \mathcal{F} into the corresponding space of partial flags. Consider the stationary measure $\nu_Q = (\pi_Q)_*(\nu)$ that is a limit of the measures $\nu_{Q, n} = (\pi_Q)_*\nu_n$ as $n \rightarrow \infty$.

We define the probability \mathbb{P} on $G \times \mathcal{F}_Q$ so that

$$\int h(g, f) d\mathbb{P}(g, f) = \int_G \int_{\mathcal{F}_Q} h(g, gf) d\nu_Q(f) d\mu(g)$$

and the mutual information I between the coordinate projections on $G \times \mathcal{F}_Q$ with respect to \mathbb{P} . Inequality (16) for ν_Q and thus theorem 5.1 will follow directly from

Proposition 5.5. *With the above notations,*

$$I \leq \sum_{\ell_Q(i) < \ell_Q(j)} \chi_i - \chi_j.$$

Indeed, by proposition 5.5, the variables G and \mathcal{F}_Q are quasi-independent under \mathbb{P} , with density $\frac{dg_*\nu_Q}{d\nu_Q}(f)$. In particular,

$$h(\mathcal{F}_Q, \mu, \nu_Q) = \int \log \left(\frac{dg_*\nu_Q}{d\nu_Q}(f) \right) dg_*\nu_Q(f) d\mu(g) = I \leq \sum_{\ell_Q(i) < \ell_Q(j)} \chi_i - \chi_j,$$

which is the statement of (16).

Proof of proposition 5.5. We analogously define the probability \mathbb{P}_n on $G \times \mathcal{F}_Q$ so that

$$\int h(g, f) d\mathbb{P}_n(g, f) = \int_G \int_{\mathcal{F}_Q} h(g, gf) d\nu_{Q, n}(f) d\mu_n(g)$$

and the mutual information I_n between the coordinate projections on $G \times \mathcal{F}_Q$ with respect to \mathbb{P}_n .

Lemma 5.6. *In the above context, $I \leq \liminf_{n \rightarrow +\infty} I_n$.*

⁵By convention, $\log \left(\frac{\mathbb{P}_{(X, Y)}(A)}{(\mathbb{P}_X \times \mathbb{P}_Y)(A)} \right) \mathbb{P}_{(X, Y)}(A) = 0$ if $\mathbb{P}_{X, Y}(A) = 0$, the sum is $+\infty$ if there is one $A \in \mathcal{A}$ such that $\mathbb{P}_{X, Y}(A) \neq 0$ and $(\mathbb{P}_X \times \mathbb{P}_Y)(A) = 0$.

Proof. By Dobrushin theorem the supremum may be taken over partitions whose sets belong to any generating set for the Borel σ -algebra. Therefore we may consider only partitions into sets satisfying $\lim_{n \rightarrow +\infty} \mathbb{P}_n(A) = \mathbb{P}(A)$. The inequality follows immediately. \square

Lemma 5.7. *For each n one has $I_n = \sum_{\ell_Q(i) < \ell_Q(j)} \chi_{i,n} - \chi_{j,n}$.*

Proof. Let φ_n be the density of $\nu_{Q,n}$ with respect to the rotationally invariant probability η_Q on \mathcal{F}_Q .

By the Gelfand-Yaglom-Perez theorem one has

$$\begin{aligned} I_n &= \int_{G \times \mathcal{F}_Q} \log \left(\frac{dg_* \nu_{Q,n}}{d\nu_{Q,n}}(f) \right) d\mathbb{P}_n(g, f) \\ &= \int_G \int_{\mathcal{F}_Q} \log \left(\frac{dg_* \nu_{Q,n}}{d\nu_{Q,n}}(gf) \right) d\nu_{Q,n}(f) d\mu_n(g) \\ &= \int_G \int_{\mathcal{F}_Q} \log \left(\frac{\varphi_n(f)}{\varphi_n(gf)} \frac{dg_* \eta_Q}{d\eta_Q}(gf) \right) d\nu_{Q,n}(f) d\mu_n(g) \end{aligned}$$

By μ_n -stationarity of $\nu_{Q,n}$ the integrals involving φ_n cancel, and one obtains

$$I_n = \int_G \int_{\mathcal{F}_Q} \log \left(\frac{dg_* \eta_Q}{d\eta_Q}(gf) \right) d\nu_{Q,n}(f) d\mu_n(g).$$

We have the following explicit formula for $\frac{dg_* \eta_Q}{d\eta_Q}(gf)$

Proposition 5.8. *For $Q = \{q_0 = 0 < q_1 < \dots < q_k = d\}$ and $\eta = \eta_Q$ the rotation invariant measure then*

$$\frac{dg\eta}{d\eta}(gx) = \frac{|\det_{U_{q_1}(x)}(g)|^{q_2} |\det_{U_{q_2}(x)}(g)|^{q_3 - q_1} \dots |\det_{U_{q_{k-1}}(x)}(g)|^{d - q_{k-2}}}{|\det(g)|^{q_{k-1}}}.$$

In particular on the space of full flags one has

$$\frac{dg\eta}{d\eta}(gx) = \frac{|\det_{U_1(x)}(g)|^2 |\det_{U_2(x)}(g)|^2 \dots |\det_{U_{k-1}(x)}(g)|^2}{|\det(g)|^{d-1}}.$$

Proposition 5.8 is proven in the next subsection. Given proposition 5.8, we may write

$$\begin{aligned} I_n &= \int_G \int_{\mathcal{F}_Q} \log \left(\frac{|\det_{U_{q_1}(f)}(g)|^{q_2} |\det_{U_{q_2}(f)}(g)|^{q_3 - q_1} \dots |\det_{U_{q_{k-1}}(f)}(g)|^{d - q_{k-2}}}{|\det(g)|^{q_{k-1}}} \right) d\nu_{Q,n}(f) d\mu_n(g) \\ &= \left(\sum_{j=1}^{k-1} (q_{j+1} - q_{j-1}) \sum_{i=1}^{q_j} \chi_{i,n} \right) - q_{k-1} \sum_{i=1}^d \chi_{i,n} \\ &= \sum_{\ell_Q(i) < \ell_Q(j)} \chi_{i,n} - \chi_{j,n}, \end{aligned}$$

where the last equality follows by direct computation. \square

Using lemma 5.4, proposition 5.5 follows. \square

Corollary 5.9. *Let $\mu \in \mathcal{M}(d), \nu$ the stationary measure on \mathcal{F} . Then,*

$$h(\mathcal{F}, \mu, \nu) \leq \sum_{0 < i < j \leq d} \chi_i - \chi_j < +\infty.$$

In particular, for μ -a. e. g, g_ν is absolutely continuous with respect to ν and the function $\log \frac{dg_*\nu}{d\nu}$ is integrable with respect to $g_*\nu$.*

Indeed, we have $\int_G \int_{\mathcal{F}} \log \left(\frac{dg_*\nu}{d\nu}(f) \right) dg_*\nu(f) d\mu(g) < +\infty$.

5.3. Proof of proposition 5.8.

Lemma 5.10. *Let $Q = \{0 < i < d\}$ and $\eta = \eta_Q$ be the unique rotationally invariant probability on \mathcal{F}_Q the Grassmannian manifold of i -dimensional subspaces of \mathbb{R}^d . Then*

$$\frac{d\eta}{d\eta}(gx) = \frac{|\det_x(g)|^d}{|\det(g)|^i},$$

for all $g \in \text{GL}_d(\mathbb{R})$.

Proof. Let $\pi_x : \mathbb{R}^d \rightarrow x$ be the orthogonal projection onto x and $\pi_{\mathbb{R}^d/x} : \mathbb{R}^d \rightarrow \mathbb{R}^d/x$ the canonical projection. The quotient space \mathbb{R}^d/x is endowed with the inner product such that the projection is an isometry when restricted to the orthogonal complement of x .

Since η is invariant under orthogonal transformations we may compose g with such transformations on both sides. In particular we may assume that $g(x) = x$ and therefore g induces a transformation on \mathbb{R}^d/x which we denote by g as well.

We may further assume that:

- (1) There is an orthogonal basis v_1, \dots, v_i of x and positive eigenvalues $\sigma_1, \dots, \sigma_i > 0$ such that $gv_k = \sigma_k v_k$ for $k = 1, \dots, i$.
- (2) There is an orthogonal basis w_1, \dots, w_{d-i} of \mathbb{R}^d/x and positive eigenvalues $\mu_1, \dots, \mu_{d-i} > 0$ such that $gw_k = \mu_k w_k$ for $k = 1, \dots, d-i$.

Notice that $\sigma_1 \cdots \sigma_i = |\det_x(g)|$ while $\mu_1 \cdots \mu_{d-i} = |\det(g)|/|\det_x(g)|$.

A local parametrization of \mathcal{F}_Q around x is given by identifying each linear mapping $\varphi : x \rightarrow \mathbb{R}^d/x$ with the subspace x_φ such that $v \in x_\varphi$ if and only if $\varphi(\pi_x(v)) = \pi_{\mathbb{R}^d/x}(v)$.

Identify each φ with its matrix $(a_{lk})_{k,l}$ where $\varphi(v_k) = \sum_l a_{lk} w_l$. With this identification, we define a volume form ω on \mathcal{F}_Q such that in any coordinates constructed as above one has $\omega(0) = \pm \bigwedge_{k,l} da_{lk}$. Since ω is orthogonally invariant it must define a volume on \mathcal{F}_Q which is a constant multiple of η .

In this chart the action of g on \mathcal{F}_Q maps φ to $g|_{\mathbb{R}^d/x} \circ \varphi \circ g|_x^{-1}$ so that

$$g|_{\mathbb{R}^d/x} \circ \varphi \circ g|_x^{-1}(v_k) = g\varphi(\sigma_k^{-1}v_k) = \sum_l a_{lk} \frac{\mu_l}{\sigma_k} w_l.$$

It follows that the pull-back under g of ω satisfies $g^*\omega(0) = \prod_{k,l} \frac{\mu_l}{\sigma_k} \omega(0) = \frac{|\det g|^i}{|\det_x g|^d} \omega(0)$,

which implies the desired claim. \square

Corollary 5.11. *If Q is obtained by a splitting an interval $k+1 < k+m$ of Q' into $k+1 < k+i < k+m$ then*

$$\frac{dg\eta_{Q,Q'}^x}{d\eta_{Q,Q'}^{gx}}(gx) = \frac{|\det_{U_{k+i}(x)}(g)|^m}{|\det_{U_k(x)}(g)|^{m-i} |\det_{U_{k+m}(x)}(g)|^i},$$

for all $g \in \mathrm{GL}_d(\mathbb{R})$ and all $x \in \mathcal{F}_Q$.

Proof. The fiber of the projection from \mathcal{F}_Q to $\mathcal{F}_{Q'}$ which contains x is naturally identified with the the i -dimensional Grasmannian of $U_{k+m}(x)/U_k(x)$.

The Jacobian of g as a mapping from $U_{k+m}(x)/U_k(x)$ to its image is $|\det_{U_{k+m}(x)}(g)|/|\det_{U_k(x)}(g)|$. The Jacobian of the restriction of g to the subspace of $U_{k+m}(x)/U_k(x)$ represented by U_{k+i} is $|\det_{U_{k+i}}(g)|/|\det_{U_k}(g)|$.

The result follows replacing these values in the previous lemma. \square

Proposition 5.8 follows from the previous corollary by splitting $\{0 < d\}$ successively into $\{0 < q_{k-1} < d\}$, $\{0 < q_{k-2} < q_{k-1} < d\}$, etc.

6. MUTUAL INFORMATION

6.1. Conditional mutual information. Given $X : \Omega \rightarrow \mathcal{X}, Y : \Omega \rightarrow \mathcal{Y}$ and $Z : \Omega \rightarrow \mathcal{Z}$ taking values in polish spaces, one may define for \mathbb{P}_Z -a.e. $z \in \mathcal{Z}$ the disintegrations $\mathbb{P}_{X \times Y}^z$ with respect to the projections on \mathcal{Z} and the mutual information $I_{\mathbb{P}^z}(X, Y)$ of X and Y given Z using these conditional distributions \mathbb{P}^z of (X, Y) given Z . Define the *conditional mutual information* $I(X, Y|Z)$ of X and Y given Z by

$$I(X, Y|Z) := I_{\mathbb{P}^z}(X, Y)$$

and the *conditional mutual entropy* $H(X, Y|Z)$ of X and Y given Z by

$$H(X, Y|Z) := \int I(X, Y|Z) d\mathbb{P}_Z(z).$$

Observe that X and Y are conditionally independent given Z if, and only if, $I(X, Y|Z) = 0$ almost everywhere, if, and only if, $H(X, Y|Z) = 0$. Observe also that, by definition, $I(X, Y|Z) = I(Y, X|Z)$. If W, X, Y, Z are measurable functions from Ω into Polish spaces then one has the following chain rule

$$H(W, (X, Y)|Z) = H(W, X|Y, Z) + H(W, Y|Z).$$

Let \mathbb{P} be the measure on $G \times \mathcal{F}$ introduced in section 5.2: for any positive measurable function h on $G \times \mathcal{F}$,

$$\int h(g, f) d\mathbb{P}(g, f) = \int h(g, gf) d\mu(g) d\nu(f).$$

We showed that $h(\mathcal{F}, \mu, \nu) = I_{\mathbb{P}}(g, f) < +\infty$.

Lemma 6.1. *The measure \mathbb{P} is the distribution of the variables (g_{-1}, E_-) on (Ω, m) . In particular, $g_{-1}(\omega)$ and $E_-(\omega)$ are quasi-independent.*

Proof. We indeed have, by invariance of m , for any positive h on $G \times \mathcal{F}$

$$\begin{aligned} \int h(g_{-1}(\omega), E_-(\omega)) dm(\omega) &= \int h(g_{-1}(\sigma\omega), E_-(\sigma\omega)) dm(\omega) \\ &= \int h(g_0(\omega), g_0(\omega)E_-(\omega)) dm(\omega) = \int h d\mathbb{P}. \end{aligned}$$

□

Using proposition 4.2, we may write, for m_+ -a.e. ω_+ :

$$h(\mathcal{F}, \mu, \nu) = I(g_{-1}, E_-) = I(g_{-1}, E_- | E_+) = I(g_{-1}, E_- | g_0, g_1, \dots).$$

Let T be an admissible topology. Recall that we defined the entropy κ_T by equation (5)

$$\kappa_T := \int \log \frac{dg_* \nu_T^{g^{-1}f'}}{d\nu_T^{f'}}(g, y) dg_* \nu_T^{g^{-1}f'}(y) d\nu'(f') d\mu(g).$$

For m -a.e. $\omega \in \Omega$ write $E_T(\omega) \in \mathcal{X}_T^{E_+(\omega)}$ for $E_T(\omega) := F_T(E_-(\omega), E_+(\omega))$. The next three propositions give other useful expressions for κ_T .

Proposition 6.2. *With the above notations, we have*

$$(18) \quad \kappa_T = H(g_{-1}, E_T | E_+) < +\infty.$$

Proof. Observe first that the RHS of (18) is finite because

$$H(g_{-1}, E_T | E_+) \leq H(g_{-1}, E_- | E_+) < +\infty.$$

The distribution of $g_{-1}(\omega)$ given $E_+(\omega)$ is μ and the distribution of $E_T(\omega)$ given $E_+(\omega)$ is $\nu_T^{E_+(\omega)}$. Remains to compute the joint distribution of $(g_{-1}(\omega), E_T(\omega))$ given $E_+(\omega)$. We claim that it projects to μ with conditional measures given by $g_{-1}(\omega)_* \nu_T^{g_{-1}(\omega)^{-1}E_+(\omega)}$.

It follows that, for m_+ -a.e. $E_+(\omega_+)$,

$$I(g_{-1}(\omega), E_T(\omega) | E_+(\omega)) = \int \log \frac{dg_* \nu_T^{g^{-1}E_+(\omega)}}{d\nu_T^{E_+(\omega)}}(g, y) dg_* \nu_T^{g^{-1}E_+(\omega)}(y) d\mu(g).$$

By integrating in $E_+(\omega)$, i.e. in f' with respect to the measure ν' , we find that $H(g_{-1}, E_T | E_+)$ is given by

$$H(g_{-1}, E_T | E_+) = \int \log \frac{dg_* \nu_T^{g^{-1}f'}}{d\nu_T^{f'}}(g, y) d\mu(g) dg_* \nu_T^{g^{-1}f'}(y) d\nu'(f'),$$

which is the formula defining κ_T in relation (5).

We prove the claim: firstly, $g_{-1}(\omega)$ is independent of $E_+(\omega)$, so that what remains to compute is the distribution of $E_T(\omega)$ given $((g_{-1}(\omega), E_+(\omega)))$. The distribution of $E_T(\sigma^{-1}\omega)$ given $((g_{-1}(\omega), E_+(\omega)))$ is by definition $\nu_T^{E_+(\sigma^{-1}\omega)}$. Since $E_+(\sigma^{-1}\omega) = g_{-1}(\omega^{-1})E_+(\omega)$, the distribution of $E_T(\omega)$ given $((g_{-1}(\omega), E_+(\omega)))$ is indeed given by $g_{-1}(\omega)_*\nu_T^{g_{-1}(\omega)^{-1}E_+(\omega)}$. \square

Proposition 6.3. *We also have*

$$\kappa_T = H(g_{-1}, E_T(\omega)|(E_+(\omega), g_0, g_1, \dots)) = H(g_{-1}, E_T(\omega)|\omega_+).$$

Proof. By proposition 4.2, the conditional measures on $E_{T_1}(\omega)$ with respect to $E_+(\omega)$ and $E_+(\omega), g_0, g_1, \dots$ coincide with $\nu_{T_1}^{E_+(\omega)}$. We have, for all admissible T , $E_T(\omega) = \pi_{T_1, T} E_{T_1}(\omega)$, so that the conditional measures on $E_T(\omega)$ with respect to $E_+(\omega)$ and $E_+(\omega), g_0, g_1, \dots$ coincide with $(\pi_{T_1, T})_* \nu_{T_1}^{E_+(\omega)} = \nu_T^{E_+(\omega)}$. \square

The projection $\omega \in \Omega \mapsto E_T(\omega) \in \mathcal{X}_T^{E_+(\omega)}$ admits disintegrations that we denote m_T^x . We still denote by m_T^x the projection of m_T^x to Ω_- . For instance, $m_{T_0}^{f'} = m_-$ for ν' -a.e. f' , $m_{T_1}^f$ is the distribution of ω_- given the unstable flag f .

By Proposition 4.2 and (18), the variables g_{-1} and E_T are quasi-independent (Indeed, since E_+ is E_T measurable, $I(g_{-1}, E_T) = I(g_{-1}, E_+) + H(g_{-1}, E_T|E_+)$ equals κ_T and thus is finite). It follows that the density f of the measure $m_T^{E_T(\omega)}$ restricted to $g_{-1}(\omega)$ with respect to μ is given by

$$f(\omega) := \frac{dm_T^{E_T(\omega)}|_{g_{-1}}(g_{-1}(\omega))}{d\mu} = \frac{d(g_{-1}(\omega))_* \nu_T^{(g_{-1}(\omega))^{-1}E_+(\omega)}}{d\nu_T^{E_+(\omega)}}(E_T(\omega)).$$

This yields another formula for κ_T :

$$(19) \quad \kappa_T = \int \left(\log \frac{dm_T^{E_T(\omega)}|_{g_{-1}}(g_{-1}(\omega))}{d\mu} \right) dm(\omega) = \int \log f(\omega) dm(\omega)$$

Proposition 6.4. *We have, for m -a.e. ω ,*

$$\kappa_T = \lim_{n \rightarrow \infty} \frac{1}{n} \log \frac{dm_T^{E_T(\omega)}|_{\{g_{-1}, \dots, g_{-n}\}}(g_{-1}(\omega), \dots, g_{-n}(\omega))}{d \otimes_1^n \mu}$$

Proof. We claim that the ratio $f^{(n)}(\omega) := \frac{dm_T^{E_T(\omega)}|_{\{g_{-1}, \dots, g_{-n}\}}(g_{-1}(\omega), \dots, g_{-n}(\omega))}{d \otimes_1^n \mu}$ is given by $f^{(n)}(\omega) = \prod_{j=0}^{n-1} f(\sigma^{-j}\omega)$. Then $\frac{1}{n} \log f^{(n)}(\omega)$ is an ergodic average of the function $\log f(\omega)$. By the Birkhoff ergodic theorem this average converges to $\int \log f(\omega) dm(\omega) = \kappa_T$ (the function $\log f$ is integrable by Corollary 5.9).

We prove the claim: by the law of composition of conditional probabilities, the ratio $f^{(n)}(\omega)/f^{(n-1)}(\omega)$ is the density with respect to μ of the conditional measures of m relative to $(g_{-1}(\omega), \dots, g_{-n+1}(\omega), E_T(\omega))$ restricted to $g_{-n}(\omega)$. But

the σ -algebra generated by $(g_{-1}(\omega), \dots, g_{-n+1}(\omega), E_T(\omega))$ is the same as the σ -algebra generated by $(E_T(\sigma^{-n+1}\omega), g_0(\sigma^{-n+1}\omega), \dots, g_{n-2}(\sigma^{-n+1}\omega))$. By proposition 6.3 and stationarity, this is the density of the measure $m_T^{E_T(\sigma^{-n+1}\omega)}$ restricted to $g_{-n}(\omega) = g_{-1}(\sigma^{-n+1}\omega)$, that is $f(\sigma^{-n+1}\omega)$.

□

6.2. The case of the projective space \mathbb{RP}^{d-1} . Consider the case of the topology T_Q associated to the partition $Q := 0 < 1 < d$. The space \mathcal{F}_Q is the projective space \mathbb{RP}^{d-1} and the measure ν_Q is the stationary measure considered by [Rap21]. In this section, we compare the formulations of our results in the case when both framework coincide. Namely, from corollary 2.5 applied to the partition Q , we get:

Corollary 6.5. *Let $\mu \in \mathcal{M}(d)$ be a discrete probability measure on $SL_d(\mathbb{R})$. Let Q be the partition $Q := 0 < 1 < d$. Then the unique stationary probability measure ν_Q on the projective space $\mathcal{F}_Q = \mathbb{RP}^{d-1}$ is exact-dimensional. There are numbers γ_j such that*

$$(20) \quad \delta_Q = \sum_{j=1}^{d-1} \gamma_j, \quad h(\mathbb{RP}^{d-1}, \mu, \nu_Q) = \sum_{j=1}^{d-1} \gamma_j (\chi_1 - \chi_{j+1}).$$

Let $T_Q = T^{d-1} \stackrel{1}{\prec} T^{d-2} \stackrel{1}{\prec} \dots \stackrel{1}{\prec} T^j \stackrel{1}{\prec} \dots \stackrel{1}{\prec} T^1 \stackrel{1}{\prec} T^0 = T_0$ be the intermediate topologies from proposition 2.1.1 applied to T_Q . The numbers γ_j are the exact dimensions of the conditional measures $\nu_{T^j, T^{j-1}}$ and we have

$$(21) \quad \gamma_j (\chi_1 - \chi_{j+1}) = \kappa_{T^j, T^{j-1}} = H(g_{-1}, E_{T^j} | E_+) - H(g_{-1}, E_{T^{j-1}} | E_+).$$

Proof. The topology T_Q associated to Q is (defined by its atoms)

$$T_Q = \{1\}, \{2, \dots, d\}, \dots, \{d-1, d\}, \{d\}.$$

Then, $D_{T_Q, T_0} = \{(1, j), 1 < j \leq d\}$ and the set of differences of exponents is $\chi_1 - \chi_{j+1}, 1 \leq j < d$. Therefore, the intermediate topologies from proposition 2.1.1

$$T_Q = T^{d-1} \stackrel{1}{\prec} T^{d-2} \stackrel{1}{\prec} \dots \stackrel{1}{\prec} T^j \stackrel{1}{\prec} \dots \stackrel{1}{\prec} T^1 \stackrel{1}{\prec} T^0 = T_0$$

are given by $T^j = \{1, j+2, \dots, d\}, \{2, \dots, d\}, \dots, \{d-1, d\}, \{d\}$. Since $\chi_{T^j, T^{j-1}} = \chi_1 - \chi_{j+1}$ for $1 \leq j < d$, we indeed have that $\chi_{T^j, T^{j-1}}$ is increasing in j . Relation (20) follows from theorem 2.2 and corollary 2.5, with $\gamma_j = \delta_{T^j, T^{j-1}}$ for $1 \leq j < d$, relation (21) from theorem 1.6 and equation (18). □

If the measure μ has finite support, a similar formula was shown in [Rap21] under the (weaker) hypothesis that the support of μ generates a strongly irreducible and proximal semi-group. Our purpose in this section is to indicate why both formulas are the same.⁶ In our case, all exponents are distinct; so, the $\tilde{\lambda}_j$ in [Rap21] are in our notations,

$$\tilde{\lambda}_j = \chi_{j+1} - \chi_1 \quad \text{for } j = 1, \dots, s = d-1.$$

⁶We will not discuss how, when restricted to T_Q , our arguments would still hold under the hypothesis that the stationary measure is unique on \mathbb{RP}^{d-1} and that $\chi_1 > \chi_2$. This would contain (and be very close to) [Rap21].

Comparing (21) with [Rap21] theorem 1.3, we see that

$$\kappa_{T^j} - \kappa_{T^{j-1}} = \kappa_{T^j, T^{j-1}} = \gamma_j(\chi_1 - \chi_{j+1}) = -\gamma_j \tilde{\lambda}_j = H_{j-1} - H_j,$$

where H_j are the partial entropies from [Rap21] theorem 1.3. Indeed, we now verify that $\kappa_{T^j} = H_0 - H_j$.

Fix $f' \in E_+(\Omega_{reg})$ and write it as the stable flag of the Oseledets decomposition

$$f' = \{0\} \subset U'_1 \subset \dots \subset U'_j \subset \dots \subset U'_{d-1} \subset \mathbb{R}^d.$$

We identify $\mathcal{X}_Q^{f'}$ with the set of directions V_1 in \mathbb{R}^d that do not belong to U'_{d-1} and for $1 \leq j \leq d-1$, $\mathcal{X}_{T^j}^{f'}$ with the set of $(d-j)$ -planes V_{d-j} such that $V_{d-j} \cap U'_{d-1} = U'_{d-j-1}$. The projection π_{T^j, T^j} associates to V_1 the space generated by V_1 and U'_{d-j+1} . For $1 \leq j < d$, we define the partition ζ_j of Ω_{reg} by the mapping $\omega \mapsto E_{T^j}(\omega)$, i.e.

$$\zeta_j(\omega) := \{\omega' \in \Omega_{reg}, U_1(\omega') + U'_{d-j-1}(\omega') = U_1(\omega) + U'_{d-j-1}(\omega)\},$$

where, for $\omega \in \Omega_{reg}$, $U_1(\omega)$ is the first unstable direction of the Oseledets decomposition. Consider the conditional measures $m_j^{E_{T^j}(\omega)}$ of m with respect to the partition ζ_j .

The entropy κ_{T^j} is given by (19):

$$\kappa_{T^j} = \int \left(\log \frac{dm_j^{E_{T^j}(\omega)}|_{g_{-1}}(g_{-1}(\omega))}{d\mu} \right) dm(\omega)$$

and by proposition 6.4, we have, for m -a.e. ω ,

$$\kappa_{T^j} = \lim_{n \rightarrow \infty} \frac{1}{n} \log \frac{dm_j^{E_{T^j}(\omega)}|_{\{g_{-1}, \dots, g_{-n}\}}(g_{-1}(\omega), \dots, g_{-n}(\omega))}{d \otimes_1^n \mu}$$

which coincide with $H_0 - H_j$ in [Rap21], Theorem 1.3. Observe that we have not used the fact that μ is discrete for this last expression. If μ is not discrete, relation (20) is not proven, but this last equation holds as soon as $\int \log \|g\| d\mu(g) < +\infty$.

6.3. Entropy difference. For any pair of admissible topologies $T \prec T'$ we defined $\kappa_{T, T'} = \kappa_T - \kappa_{T'}$. By relation (18), $\kappa_{T, T'} < +\infty$ and if $T \prec T' \prec T''$ one has

$$\kappa_{T, T''} = \kappa_{T, T'} + \kappa_{T', T''}.$$

By the chain rule for conditional mutual entropy, relation(18) and proposition 6.3, we have

$$\kappa_{T, T'} = H(g_{-1}, E_T | E_{T'}) = H(g_{-1}, E_T | (E_{T'}, g_0, g_1, \dots)) = H(g_{-1}, E_T | (E_{T'}, \omega_+)).$$

Recall that in the introduction we defined, for ν' -a.e. $f' \in \mathcal{F}$, $\nu_{T'}^{f'}$ -a.e. $x' \in \mathcal{X}_{T'}^{f'}$, $\nu_{T, T'}^{x'}$ as a family of disintegrations of the measure $\nu_{T'}^{f'}$ with respect to $\pi_{T, T'}$. Then,

$$(22) \quad \frac{dg_* \nu_T^{f'}}{d\nu_T^{g f'}}(y) = \frac{dg_* \nu_{T'}^{x'}}{d\nu_{T, T'}^{g x'}}(y) \frac{dg_* \nu_{T'}^{f'}}{d\nu_{T'}^{g f'}}(x').$$

Proposition 6.6. *If $T \prec T'$ are admissible topologies then*

$$\kappa_{T,T'} = \int_{\Omega} \log \left(\frac{dg_0(\omega) \nu_{T,T'}^{E_{T'}(\omega)}}{d\nu_{T,T'}^{E_{T'}(\sigma(\omega))}}(E_T(\sigma(\omega))) \right) dm(\omega).$$

Proof. We write κ_T as $\kappa_T = \mathbb{E}[I(g_{-1}, E_T|E_+)(\omega)]$. Reporting the explicit expression for $I(g_{-1}(\omega), E_T(\omega)|E_+(\omega))$, we have

$$\begin{aligned} \kappa_T &= \mathbb{E} \left[\log \frac{d(g_{-1}(\omega))_* \nu_T^{(g_{-1}(\omega))^{-1}E_+(\omega)}}{d\nu_T^{E_+(\omega)}}(E_T(\omega)) \right] \\ &= \mathbb{E} \left[\log \frac{d(g_0(\omega))_* \nu_T^{E_+(\omega)}}{d\nu_T^{E_+(\sigma(\omega))}}(E_T(\sigma(\omega))) \right], \end{aligned}$$

where the second line follows by σ -invariance. The proposition follows by making the difference $\kappa_{T,T'} = \kappa_T - \kappa_{T'}$ and applying (22). \square

Fix $T \prec T'$. For a.e. $x \in (\pi_{T,T'})^{-1}(E_{T'}(\omega))$, set $f_\omega(x) := \frac{dg_0(\omega) \nu_{T,T'}^{E_{T'}(\omega)}}{d\nu_{T,T'}^{E_{T'}(\sigma(\omega))}}(x)$. From proposition 6.6, follows

$$(23) \quad \kappa_{T,T'} = \mathbb{E}[\log f_\omega(E_T(\omega))].$$

Recall formula (19) for κ_T and $\kappa_{T'}$. With the same notations, we have

$$\kappa_{T,T'} = \int \left(\log \frac{dm_T^{E_T(\omega)}|_{g_{-1}}(\omega)}{dm_{T'}^{E_{T'}(\omega)}|_{g_{-1}}}(\omega) \right) dm(\omega)$$

and the following corollary of Proposition 6.4

Corollary 6.7. *Assume $T \prec T'$ are admissible topologies. Then, we have, for m -a.e. ω ,*

$$\kappa_{T,T'} = \lim_{n \rightarrow \infty} \frac{1}{n} \log \frac{dm_T^{E_T(\omega)}|_{\{g_{-1}, \dots, g_{-n}\}}}{dm_{T'}^{E_{T'}(\omega)}|_{\{g_{-1}, \dots, g_{-n}\}}}(g_{-1}(\omega), \dots, g_{-n}(\omega)).$$

6.4. Zero entropy difference.

Proposition 6.8. *If $\kappa_{T,T'} = 0$ then $\nu_{T,T'}^{E_{T'}(\omega)}$ is the point mass at $E_T(\omega)$ for m -a.e. $\omega \in \Omega$. In particular, it is exact dimensional with dimension 0.*

Proof. By proposition 2.1, we may assume that $T \stackrel{1}{\prec} T'$ and that $i < j$ are such that $T'(i) = T(i) \cup \{j\}$. By proposition 6.6, we have

$$\kappa_{T,T'} = \int_{\Omega} \log \left(\frac{dg_0(\omega) \nu_{T,T'}^{E_{T'}(\omega)}}{d\nu_{T,T'}^{E_{T'}(\sigma(\omega))}}(E_T(\sigma(\omega))) \right) dm(\omega).$$

Therefore, by Jensen inequality, if $\kappa_{T,T'} = 0$ then for m -almost every ω we have $g_0(\omega) \nu_{T,T'}^{E_{T'}(\omega)} = \nu_{T,T'}^{E_{T'}(\sigma(\omega))}$. Since m is σ -invariant, we obtain that m -almost everywhere one has $\nu_{T,T'}^{E_{T'}(\omega)} = g_{-1}(\omega) \dots g_{-n}(\omega) \nu_{T,T'}^{E_{T'}(\sigma^{-n}(\omega))}$ for all $n \geq 1$.

Observe that, since E_+ and E_- are in general position, one has $E_{T'}(\omega)_{T'(i)\setminus\{i\}} \neq E_T(\omega)_{T(i)}$ for m -almost every $\omega \in \Omega$, and therefore $\nu_{T,T'}^{E_{T'}(\omega)}(\{E_{T'}(\omega)_{T'(i)\setminus\{i\}}\}) = 0$. Using the coordinate $\varphi_{x(\omega)}$ (see lemma 4.7), we have $\nu_{T,T'}^{E_{T'}(\omega)}[\varphi_{x(\omega)}((-\pi/2, \pi/2))] = 1$.

By Poincaré recurrence, for almost every $\omega \in \Omega_-$, there exists an infinite sequence $n_k, k \in \mathbb{N}$ and $\alpha > 0$ such that $\nu_{T,T'}^{E_{T'}(\sigma^{-n_k}(\omega))}$ converges to $\nu_{T,T'}^{E_{T'}(\omega)}$ as $k \rightarrow \infty$ and $\nu_{T,T'}^{E_{T'}(\omega)}((\varphi_{x(\sigma^{-n_k}(\omega))})((-\alpha, \alpha))) > \alpha$.

Hence, for those ω for which lemma 4.7 hold, $g_{-1}(\omega) \dots g_{-n_k}(\omega)$ sends any neighborhood of $E_T(\sigma^{-n_k}(\omega))$ to a small neighborhood of $E_T(\omega)$. This is possible only if $\nu_{T,T'}^{E_{T'}(\omega)}$ is concentrated at $E_T(\omega)$. \square

7. PROOF OF THEOREM 1.6

Theorem 1.6 is an almost everywhere statement. It uses a telescoping argument mixed with weak type (1,1) techniques as in the proof of Shannon-McMillan-Breiman theorem for finite entropy partitions. The proof follows [HS17] and [Les21].

In this section, we assume that $T \stackrel{1}{\prec} T'$ and that $i < j$ are such that $T'(i) = T(i) \cup \{j\}$. Let $\chi := \chi_{T,T'}, \kappa := \kappa_{T,T'}$. we want to show that for ν' -a.e. $f' \in \mathcal{F}$, $\nu_{T'}^{f'}$ -a.e. $x' \in \mathcal{X}_{T'}^{f'}$, the conditional measure $\nu_{T'}^{x'}$ is exact-dimensional with dimension κ/χ .

7.1. Length of stationary neighborhoods. Let $\pi = \pi_{T,T'}$ be the projection from \mathcal{X}_T to $\mathcal{X}_{T'}$ and consider the coordinates given by lemma 3.4 on $\pi^{-1}(E_{T'}(\omega))$, setting $x = E_T(\omega)$ and $x' = E_{T'}(\omega)$.

For each $\alpha, 0 < \alpha < \pi/2$ let $N^\alpha(\omega)$ (respectively $N^{\alpha,+}(\omega), N^{\alpha,-}(\omega)$) the set $\varphi((-\alpha, \alpha))$ (respectively $\varphi([0, \alpha]), \varphi((-\alpha, 0])$). Recall that we denoted by η the rotation invariant probability measure on $\pi^{-1}(E_{T'}(\omega))$. The measure $\varphi_* du$ has a bounded continuous density with respect to η (the bound depends on ω). From lemma 4.7 follows

Lemma 7.1. *For all $\alpha, 0 < \alpha < \pi/2$, for m -a.e. $\omega \in \Omega$ one has*

$$\eta(g_{-1}(\omega) \dots g_{-n}(\omega) N^\alpha(\sigma^{-n}(\omega))) = e^{-\chi n + o(n)} \text{ as } n \rightarrow +\infty,$$

and the same holds for $N^{\alpha,+}$ and $N^{\alpha,-}$.

In the sequel, we set, for each $\alpha, 0 < \alpha < \pi/2$ and for each $n \geq 1$, $N_n^\alpha(\omega)$ for the interval in $\pi^{-1}(E_{T'}(\omega))$ given by

$$N_n^\alpha(\omega) = (g_{-1}(\omega) \dots g_{-n}(\omega))(N^\alpha(\sigma^{-n}(\omega))).$$

$N_n^{\alpha,+}(\omega), N_n^{\alpha,-}(\omega)$ are defined similarly, $N_n^{\alpha,\pm}(\omega)$ is a choice of one of the three intervals.

7.2. Probability of stationary neighborhoods. Theorem 1.6 will follow by comparing lemma 7.1 and the following

Proposition 7.2. *For all $\alpha, 0 < \alpha < \pi/2$, for m -a.e. $\omega \in \Omega$, one has*

$$\nu_{T,T'}^{E_{T'}(\omega)}(N_n^{\alpha,\pm}(\omega)) = e^{-\kappa n + o(n)} \nu_{T,T'}^{E_{T'}(\sigma^{-n}(\omega))}(N_n^{\alpha,\pm'}(\sigma^{-n}(\omega))) \text{ as } n \rightarrow +\infty.$$

In this formula, the sign \pm' is not necessarily the same as \pm .

Proof. Recall that we set $f_\omega(x) = \frac{dg_0(\omega)\nu_{T,T'}^{E_{T'}(\omega)}}{d\nu_{T,T'}^{E_{T'}(\sigma(\omega))}}(x)$, and that by (23),

$$\kappa = \int \log f_\omega(E_T(\omega)) dm(\omega) = \mathbb{E} \left[\int_{\pi^{-1}(E_{T'}(\omega))} \log f_\omega(x) d\nu_{T,T'}^{E_{T'}(\omega)}(x) \right].$$

Since $\kappa < +\infty$, for m -a.e. ω , $\int_{\pi^{-1}(E_{T'}(\omega))} \log f_\omega(x) d\nu_{T,T'}^{E_{T'}(\omega)}(x) < +\infty$.

Set for each $\alpha, 0 < \alpha < \pi/2$, for each $n \geq 1$,

$$f_n^{\alpha,\pm}(\omega) = \frac{\nu_{T,T'}^{E_{T'}(\omega)}(N_n^{\alpha,\pm}(\omega))}{\nu_{T,T'}^{E_{T'}(\sigma(\omega))}(g_0(\omega)N_n^{\alpha,\pm}(\omega))} = \frac{\int f_\omega(x) d\nu_{T,T'}^{E_{T'}(\sigma(\omega))}(x)}{\nu_{T,T'}^{E_{T'}(\sigma(\omega))}(g_0(\omega)N_n^{\alpha,\pm}(\omega))}.$$

Lemma 7.3. *For all $\alpha, 0 < \alpha < \pi/2$, for m -a.e. $\omega \in \Omega$ one has $\lim_{n \rightarrow +\infty} f_n^{\alpha,\pm}(\omega) = f_\omega(E_T(\sigma(\omega)))$. Furthermore $\int \sup_{n \geq 1} |\log(f_n^{\alpha,\pm}(\omega))| dm(\omega) < +\infty$.*

Proof. By lemma 7.1 the intervals $N_n^{\alpha,\pm}(\omega)$ intersect to $E_T(\omega)$ when $n \rightarrow +\infty$ for m -a.e. $\omega \in \Omega$. By the Lebesgue differentiation theorem this implies

$$\lim_{n \rightarrow +\infty} f_n^{\alpha,\pm}(\omega) = f_\omega(g_0(\omega)E_T(\omega)) = f_\omega(E_T(\sigma(\omega))),$$

for m -a.e. $\omega \in \Omega$ as claimed.

For the second claim, let Mf_ω be the maximal function defined by

$$Mf_\omega(x) = \sup_{N \supset \{x\}} \frac{1}{\nu_{T,T'}^{E_{T'}(\sigma(\omega))}(N)} \int_N f_\omega(x) d\nu_{T,T'}^{E_{T'}(\sigma(\omega))}(x),$$

where the supremum is over intervals N starting or finishing at x in $\pi^{-1}(E_{T'}(\sigma(\omega)))$. Since $\pi^{-1}(x')$ is one-dimensional, for all $x' \in \mathcal{X}_{T'}$, the maximal operator is of weak type (1,1) and we have (see [Les21, Lemma 8], [Les21, Lemma 9] for details)

$$(24) \quad \int \log(Mf_\omega(E_T(\sigma(\omega)))) dm(\omega) < +\infty.$$

It remains to show that $\inf_n \log(f_n^{\alpha,\pm}(\omega))$ is m -integrable. See [Les21, Lemma 10] for the argument in a very similar setting. \square

From Birkhoff ergodic theorem we have

$$\begin{aligned}\kappa &= \lim_{n \rightarrow +\infty} \frac{1}{n} \sum_{k=1}^n \log \left(\frac{dg_{-k} \nu_{T,T'}^{E_{T'}(\sigma^{-k}(\omega))}}{d\nu_{T,T'}^{E_{T'}(\sigma^{-k+1}(\omega))}}(E_T(\sigma^{-k+1}(\omega))) \right) \\ &= \lim_{n \rightarrow +\infty} \frac{1}{n} \sum_{k=1}^n \log (f_{\sigma^{-k}(\omega)}(E_T(\sigma(\sigma^{-k}(\omega))))))\end{aligned}$$

To conclude, using Lemma 7.3 and Maker theorem (see e.g. [Rap21, Theorem 5.7]) we obtain

$$\begin{aligned}\kappa &= \lim_{n \rightarrow +\infty} \frac{1}{n} \sum_{k=1}^n \log (f_{n-k}^{\alpha, \pm}(\sigma^{-k}(\omega))) \\ &= \lim_{n \rightarrow +\infty} \frac{1}{n} \sum_{k=1}^n \log \left(\frac{\nu_{T,T'}^{E_{T'}(\sigma^{-k}(\omega))}(N_{n-k}^{\alpha, \pm}(\sigma^{-k}(\omega)))}{\nu_{T,T'}^{E_{T'}(\sigma^{-k+1}(\omega))}(g_{-k}(\omega)N_{n-k}^{\alpha, \pm}(\sigma^{-k}(\omega)))} \right) \\ &= \lim_{n \rightarrow +\infty} \frac{1}{n} \sum_{k=1}^n \log \left(\frac{\nu_{T,T'}^{E_{T'}(\sigma^{-k}(\omega))}(g_{-k-1}(\omega) \dots g_{-n}(\omega)N^{\alpha, \pm}(\sigma^{-n}(\omega)))}{\nu_{T,T'}^{E_{T'}(\sigma^{-k+1}(\omega))}(g_{-k}(\omega) \dots g_{-n}(\omega)N^{\alpha, \pm}(\sigma^{-n}(\omega)))} \right) \\ &= \lim_{n \rightarrow +\infty} \frac{1}{n} \log \left(\frac{\nu_{T,T'}^{E_{T'}(\sigma^{-n}(\omega))}(N^{\alpha, \pm}(\sigma^{-n}(\omega)))}{\nu_{T,T'}^{E_{T'}(\omega)}(N_n^{\alpha, \pm}(\omega))} \right).\end{aligned}$$

□

7.3. Exact dimensionality. We finish the proof of theorem 1.6. Firstly, if $\kappa = 0$, by proposition 6.8, the measure $\nu_{T,T'}^{E_{T'}(\omega)}$ is exact-dimensional with dimension 0. On the other hand, since $\chi > 0$ and $\kappa = 0$, we have $0 = \kappa/\chi$.

So we may assume $\kappa > 0$. By proposition 7.2, for m -a.e. ω , all $\alpha, 0 < \alpha < \pi/2$, n going to infinity,

$$\frac{1}{n} \log \nu_{T,T'}^{E_{T'}(\omega)}(N_n^{\alpha, \pm}(\omega)) = -\kappa + o(1) + \frac{1}{n} \log \nu_{T,T'}^{E_{T'}(\sigma^{-n}\omega)}(N_n^{\alpha, \pm}(\sigma^{-n}\omega)).$$

On the other hand, we have, by lemma 7.1, for m -a.e. ω , all $\alpha, 0 < \alpha < \pi/2$, n going to infinity,

$$B_{\pi^{-1}(E_{T'}(\omega))}^{\pm}(E_T(\omega), e^{-n\chi+o(n)}) \subset N_n^{\alpha, \pm}(\omega) \subset B_{\pi^{-1}(E_{T'}(\omega))}^{\pm}(E_T(\omega), e^{-n\chi+o(n)}),$$

where $B_{\pi^{-1}(E_{T'}(\omega))}^{\pm}(x, r)$ is either of the two intervals of size r based on x .

It follows that for m -a.e. ω , $\nu_{T,T'}^{E_{T'}(\omega)}$ -a.e. x ,

$$\liminf_{r \rightarrow 0} \frac{\log(\nu_{T,T'}^{E_{T'}(\omega)}(B(x, r)))}{\log(r)} \geq \frac{\kappa}{\chi}.$$

We cannot estimate directly $\limsup_{r \rightarrow 0} \frac{\log(\nu_{T,T'}^{E_{T'}(\omega)}(B(x,r)))}{\log(r)}$ in the same way, because we do not know a priori that $\liminf_{n \rightarrow \infty} \frac{1}{n} \log(\nu_{T,T'}^{E_{T'}(\sigma^{-n}\omega)}(N^{\alpha,\pm}(\sigma^{-n}\omega))) = 0$. The observation is that to compute $\limsup_{r \rightarrow 0} \frac{\log(\nu_{T,T'}^{E_{T'}(\omega)}(B(x,r)))}{\log(r)}$, it suffices to consider values of $-\log r$ in \mathbb{N} with a positive density.

We claim that we can take α large enough that one of $\nu_{T,T'}^{E_{T'}(\omega)}(N^{\alpha,+}(\omega))$ and $\nu_{T,T'}^{E_{T'}(\omega)}(N^{\alpha,-}(\omega))$ is at least some $c > 0$ on a set $\Omega' \subset \Omega$ of positive measure. This finishes the proof, because, by Birkhoff ergodic theorem, for m -a.e. ω the sequence n_k such that $\sigma^{-n_k}\omega \in \Omega'$ has positive density.

Remains to prove the claim. We prove it by contradiction: if it is not true, $\nu_{T,T'}^{E_{T'}(\omega)}(N^\alpha(\omega)) = 0$ m -a.e. for all α . But then, the measure $\nu_{T,T'}^{E_{T'}(\omega)}$ is concentrated on $E_{T'}(\omega)_{T'(i) \setminus \{i\}}$, a contradiction with the fact that E_+ and E_- are in general position.

7.4. **Proof of lemma 2.7.** Assume the diagram of projections $\begin{array}{ccc} T & \longrightarrow & S \\ \downarrow 1 & & \downarrow 1 \\ T' & \longrightarrow & S' \end{array}$ commutes and i, j are such that $T(i) = T'(i) \setminus \{j\}$ and $S(i) = S'(i) \setminus \{j\}$.

Then, by Corollary 3.6, for $x' \in \mathcal{X}_{S'}$ and $y' \in (\pi_{T',S'})^{-1}(x')$ there is a bilipschitz homeomorphism between $(\pi_{S,S'})^{-1}(x')$ and $(\pi_{T,T'})^{-1}(y')$. The measure $\nu_{S,S'}^{x'}$ is the average over y' of the measures $(\pi_{T,S})_* \nu_{T,T'}^{y'}$, the average being taken under $d\nu_{T',S'}^{x'}(y')$. Lemma 2.7 then follows from [LY85] Lemma 11.3.2. \square

8. PROOF OF THEOREM 2.6

In this section, we assume that the measure μ is discrete and prove Theorem 2.6.

The general idea of the proof is that entropy conservation implies dimension conservation. In [LY85], dynamical balls are disjoint ellipsoids with exponentially big eccentricities, not suitable for dimension estimates. But they behave very well for entropy estimates, even when considering their slices by invariant foliations. If one takes the slices in increasing order of size, this forces dimension conservation. For IFS or stationary measures, the dynamical balls are not disjoint any more. The idea of [Fen19] is to look at the dynamical balls and the slicing at the level of the invertible dynamics on Ω . Since the measure μ is discrete, working at the level of the space Ω is possible here as well.

8.1. **Setup.** Recall that we consider $T \prec T'$ and the decomposition $T^0, T^1, \dots, T^{N_{T,T'}}$, where $T^0 = T'$ and $T^{N_{T,T'}} = T$ of proposition 2.1 with $T^t \stackrel{1}{\prec} T^{t-1}$ for $t = 1, \dots, N_{T,T'}$.

Recall that for each admissible topology S we set $E_S(\omega) = F_S(E(\omega))$.

We will use x to denote a point in \mathcal{X}_T and always set $x_t = \pi_{T, T^t}(x)$ and $x_{t-1} = \pi_{T, T^{t-1}}(x)$.

We fix t and set $\chi = \chi_{T^t, T^{t-1}}$, $\kappa = \kappa_{T^t, T^{t-1}}$ and $\delta = \overline{\delta^t}$. Then, $\gamma_{T^t, T^{t-1}} = \kappa/\chi$.

We are interested in comparing for $\nu_\omega = \nu_{T, T_0}^{E_+(\omega)}$ -almost every x , the upper dimension at x of $\nu_{T, T^t}^{x_t}$ and $\nu_{T, T^{t-1}}^{x_{t-1}}$.

For this purpose we fix $\varepsilon > 0$ and set $r_n = \exp(-n(\chi - \varepsilon))$ for all n in what follows.

8.2. Approximating configurations.

Lemma 8.1. *There exist functions f_n such that, for m -a.e. ω , setting $E_{T,n}(\omega) = f_n(E_{T^{t-1}}(\omega), g_{-1}(\omega), \dots, g_{-n}(\omega))$ there exists $n(\omega)$ such that for all $n \geq n(\omega)$, $E_{T,n}(\omega) \in B(E_T(\omega), r_n)$.*

Proof. The construction of Proposition 2.1 guarantees that

$$\chi = \min \{ \chi_{T^s, T^{s-1}} : s = t, t+1, \dots, N_{T, T'} \}.$$

Therefore the claim follows directly from proposition 4.6. \square

Corollary 8.2. *For m -a.e. ω , ν_ω -almost every x and $m_{T^t}^{x_t}$ -almost every $(h_n)_{n \leq -1}$ there exists $N(\omega, x_t, (h_n)_{n \leq -1})$ such that, for all $n \geq N(\omega, x_t, (h_n)_{n \leq -1})$.*

$$f_n(x_{t-1}, h_{-1}, \dots, h_{-n}) \in B \left(\lim_{k \rightarrow +\infty} f_k(x_{t-1}, h_{-1}, \dots, h_{-k}), r_n \right).$$

8.3. Approximating conditional probabilities. Let T be an admissible topology. By definition, $E_T(\omega) \in \mathcal{X}_T^{E_+(\omega)}$ and, if $T \prec T'$, $E_{T'}(\omega) = \pi_{T, T'}(E_T(\omega))$. We introduced in section 6.1 the disintegrations of the projection $\omega \in \Omega \mapsto E_T(\omega) \in \mathcal{X}_T^{E_+(\omega)}$ and their restrictions $m_T^{E_T(\omega)}$ to Ω_- . With our notations, we have, for ν' -a.e. f' , $\nu_{T^t}^{f'}$ -a.e. $z \in \mathcal{X}_{T^{t-1}}$,

$$(25) \quad m_{T^{t-1}}^z = \int_{\mathcal{X}_{T^t, T^{t-1}}^z} m_{T^t}^y d\nu_{T^t, T^{t-1}}^z(y).$$

Given $x \in \mathcal{X}_T$, let $A_n(x_{t-1})$ be the set of sequences $(h_n)_{n \leq -1}$ such that, for all $m \geq n$, $f_m(x_{t-1}, h_{-1}, \dots, h_{-m}) \in B \left(\lim_{k \rightarrow +\infty} f_k(x_{t-1}, h_{-1}, \dots, h_{-k}), r_m \right)$. We let $[h_{-n}, \dots, h_{-1}]$ denote the set of sequences $(h'_n)_{n \leq -1}$ such that $h_{-n} = h'_{-n}, \dots, h_{-1} = h'_{-1}$.

Lemma 8.3. *For m -a.e. ω , ν_ω -almost every x and $m_{T^t}^{x_t}$ -almost every $(h_n)_{n \leq -1}$, there exists $N(\omega, x_t, (h_n)_{n \leq -1})$ such that, for all $n \geq N(\omega, x_t, (h_n)_{n \leq -1})$, $m_{T^t}^{x_t}([h_{-n}, \dots, h_{-1}] \cap A_n(x_{t-1})) \leq \exp(n(\kappa + \varepsilon)) m_{T^{t-1}}^{x_{t-1}}([h_{-n}, \dots, h_{-1}] \cap A_n(x_{t-1}))$.*

Proof. For m -a.e. ω , consider the measure \mathbb{P}_ω on the space $\mathcal{X}_T \times \{(h_k)_{k \leq -1}\}$ which projects to ν_ω on \mathcal{X}_T and whose disintegrations on the fibers projecting to $x \in \mathcal{X}_T$ are given by $x \mapsto m_{T^t}^{x_t}$.

By corollary 8.2, for m -a.e. ω , ν_ω -almost every x , $A_n(x_{t-1})$ is increasing with n and $\cup_n A_n(x_{t-1}) = \Omega_-$ up to a set of $m_{T^t}^{x_t}$ -measure 0. By the martingale convergence theorem, the conditional expectation with respect to \mathbb{P}_ω of the indicator of the event $\{(h_k)_{k \leq -1} \in A_n(x_{t-1})\}$ with respect to the σ -algebras generated by $x_t, h_{-1}, \dots, h_{-n}$ (respectively $x_{t-1}, h_{-1}, \dots, h_{-n}$) converge to 1.

The first conditional expectation is given at m -a.e. ω , ν_ω -almost every x and $m_{T^t}^{x_t}$ -almost every $(h_n)_{n \leq -1}$ by

$$\frac{m_{T^t}^{x_t}([h_{-n}, \dots, h_{-1}] \cap A_n(x_{t-1}))}{m_{T^t}^{x_t}([h_{-n}, \dots, h_{-1}])}.$$

The second one by

$$\frac{\int_{\mathcal{X}_{T^t, T^{t-1}}^{x_t, x_{t-1}}} m_{T^t}^y([h_{-n}, \dots, h_{-1}] \cap A_n(x_{t-1})) d\nu_{T^t, T^{t-1}}^{x_t, x_{t-1}}(y)}{\int_{\mathcal{X}_{T^t, T^{t-1}}^{x_t, x_{t-1}}} m_{T^t}^y([h_{-n}, \dots, h_{-1}]) d\nu_{T^t, T^{t-1}}^{x_t, x_{t-1}}(y)}.$$

Using (25), this last expression is

$$\frac{m_{T^{t-1}}^{x_{t-1}}([h_{-n}, \dots, h_{-1}] \cap A_n(x_{t-1}))}{m_{T^{t-1}}^{x_{t-1}}([h_{-n}, \dots, h_{-1}])}$$

Thus, at m -a.e. ω , ν_ω -almost every x and $m_{T^t}^{x_t}$ -almost every $(h_n)_{n \leq -1}$,

$$\lim_{n \rightarrow +\infty} \frac{m_{T^t}^{x_t}([h_{-n}, \dots, h_{-1}] \cap A_n(x_{t-1}))}{m_{T^t}^{x_t}([h_{-n}, \dots, h_{-1}])} = \lim_{n \rightarrow +\infty} \frac{m_{T^{t-1}}^{x_{t-1}}([h_{-n}, \dots, h_{-1}] \cap A_n(x_{t-1}))}{m_{T^{t-1}}^{x_{t-1}}([h_{-n}, \dots, h_{-1}])} = 1.$$

We have shown (in corollary 6.7) that for m -a.e. ω , ν_ω -almost every x and $m_{T^t}^{x_t}$ -almost every $(h_n)_{n \leq -1}$ one has

$$\lim_{n \rightarrow +\infty} \frac{1}{n} \log \left(\frac{m_{T^t}^{x_t}([h_{-n}, \dots, h_{-1}])}{m_{T^{t-1}}^{x_{t-1}}([h_{-n}, \dots, h_{-1}])} \right) = \kappa.$$

Combining these three statements we obtain that for m -a.e. ω , ν_ω -almost every x and $m_{T^t}^{x_t}$ -almost every $(h_n)_{n \leq -1}$ one has

$$\lim_{n \rightarrow +\infty} \frac{1}{n} \log \left(\frac{m_{T^t}^{x_t}([h_{-n}, \dots, h_{-1}] \cap A_n(x_{t-1}))}{m_{T^{t-1}}^{x_{t-1}}([h_{-n}, \dots, h_{-1}] \cap A_n(x_{t-1}))} \right) = \kappa,$$

from which the desired result follows immediately. \square

8.4. Lebesgue Density. Given $x \in \mathcal{X}_T$ let $B_n(x_t)$ denote the set of sequences $(h_n)_{n \leq -1}$ such that

$$m_{T^t}^{x_t}([h_{-n}, \dots, h_{-1}] \cap A_n(x_{t-1})) \leq \exp(n(\kappa + \varepsilon)) m_{T^{t-1}}^{x_{t-1}}([h_{-n}, \dots, h_{-1}] \cap A_n(x_{t-1})).$$

Lemma 8.4. *For m -a.e. ω and ν_ω almost every x there exists $N(\omega, x)$ such that, for all $n \geq N(\omega, x)$,*

$$m_{T^t}^{x_t} \left(\left\{ \lim_{k \rightarrow +\infty} f_k(x_{t-1}, h_{-1}, \dots, h_{-k}) \in B(x, r_n) \right\} \cap A_n(x_{t-1}) \cap B_n(x_t) \right) \geq \exp(-n(\delta + \varepsilon)\chi).$$

Proof. The probability on the left-hand side in the statement is

$$\int_{B(x, r_n)} m_T^y(A_n(x_{t-1}) \cap B_n(x_t)) d\nu_{T, T^t}^{x_t}(y) = \int_{B(x, r_n)} m_T^y(A_n(y_{t-1}) \cap B_n(y_t)) d\nu_{T, T^t}^{x_t}(y).$$

Let $C_n(x) = \bigcap_{m \geq n} B_m(x_t)$ so that $C_n(x)$ is increasing with n .

We have

$$\int_{B(x, r_n)} m_T^y(A_n(y_{t-1}) \cap B_n(y_t)) d\nu_{T, T^t}^{x_t}(y) \geq \int_{B(x, r_n)} m_T^y(A_n(y_{t-1}) \cap C_n(y_t)) d\nu_{T, T^t}^{x_t}(y).$$

By lemma 8.3, for m -a.e. ω , ν_ω -a.e. x , the function $y \mapsto m_T^y(A_n(y_{t-1}) \cap C_n(y_t))$ increases to 1 at $\nu_{T, T^t}^{x_t}$ -a.e. y . Applying the Lebesgue differentiation theorem (justified since configuration spaces are bilipschitz homeomorphic to Euclidean spaces) we obtain a set $L(\omega, x_t)$ of $\nu_{T, T^t}^{x_t}$ -full measure such that for all z in this set

$$\lim_{n \rightarrow +\infty} \frac{1}{\nu_{T, T^t}^{x_t}(B(z, r_n))} \int_{B(z, r_n)} m_T^y(A_n(y_{t-1}) \cap C_n(y_t)) d\nu_{T, T^t}^{x_t}(y) = 1.$$

For m -a.e. ω , ν_ω -a.e. x belongs to $L(\omega, x_t)$. Moreover, by hypothesis, for m -a.e. ω and ν_ω -a.e. x , there exists $N(\omega, x)$ such that for $n \geq N(\omega, x)$, $\nu_{T, T^t}^{x_t}(B(x, r_n)) \geq \exp(-n(\delta + \varepsilon)\chi)$. The result follows. \square

8.5. Proof of the theorem. We now complete the proof of Theorem 2.4.

We begin with Lemma 8.4 and observe that if $\lim_{k \rightarrow +\infty} f_k(x_{t-1}, h_{-1}, \dots, h_{-k}) \in B(x, r_n)$ and $(h_n)_{n \leq -1} \in A_n(x_{t-1})$, then in fact $f_n(x_{t-1}, h_{-1}, \dots, h_{-n}) \in B(x, 2r_n)$.

This implies that for m -a.e. ω , ν_ω -a.e. x and all $n \geq N(\omega, x)$ given by Lemma 8.4 one has

$$\begin{aligned} \exp(-n(\delta + \varepsilon)\chi) &\leq m_{T^t}^{x_t} \left(\left\{ \lim_{k \rightarrow +\infty} f_k(x_{t-1}, h_{-1}, \dots, h_{-k}) \in B(x, r_n) \right\} \cap A_n(x_{t-1}) \cap B_n(x_t) \right) \\ &\leq m_{T^t}^{x_t} (\{f_n(x_{t-1}, h_{-1}, \dots, h_{-n}) \in B(x, 2r_n)\} \cap A_n(x_{t-1}) \cap B_n(x_t)). \end{aligned}$$

Notice that both $D_n(x_{t-1}) := \{(h_n)_{n \leq -1} : f_n(x_{t-1}, h_{-1}, \dots, h_{-n}) \in B(x, 2r_n)\}$ and $B_n(x_t)$ are a union of cylinders $[h_{-n}, \dots, h_{-1}]$. Therefore from the second line above and the definition of $B_n(x_t)$ we obtain

$$\begin{aligned} \exp(-n(\delta + \varepsilon)\chi) &\leq \sum_{[h_{-n}, \dots, h_{-1}] \subset B_n(x_t) \cap D_n(x_{t-1})} m_{T^t}^{x_t}([h_{-n}, \dots, h_{-1}] \cap A_n(x_{t-1})) \\ &\leq \exp(n(\kappa + \varepsilon)) \sum_{[h_{-n}, \dots, h_{-1}] \subset D_n(x_{t-1})} m_{T^{t-1}}^{x_{t-1}}([h_{-n}, \dots, h_{-1}] \cap A_n(x_{t-1})). \end{aligned}$$

For m -a.e. ω , ν_ω -a.e. x and all $n \geq N(\omega, x)$, we have shown that $\exp(-n(\delta + \varepsilon)\chi) \exp(-n(\kappa + \varepsilon)) \leq m_{T^{t-1}}^{x_{t-1}}(\{f_n(x_{t-1}, h_{-1}, \dots, h_{-n}) \in B(x, 2r_n)\} \cap A_n(x_{t-1}))$.

Since whenever $(h_n)_{n \leq -1} \in A_n(x_{t-1})$ we have that $f_n(x_{t-1}, h_{-1}, \dots, h_{-n})$ and $\lim_{k \rightarrow +\infty} f_k(x_{t-1}, h_{-1}, \dots, h_{-k})$ are at distance at most r_n , this implies that for m -a.e. ω , ν_ω -a.e. x and all $n \geq N(\omega, x)$,

$$\begin{aligned} \exp(-n(\delta + \varepsilon)\chi) \exp(-n(\kappa + \varepsilon)) &\leq m_{T^{t-1}}^{x_{t-1}} \left(\left\{ (h_k)_{k \leq -1} : \lim_{k \rightarrow +\infty} f_k(x_{t-1}, h_{-1}, \dots, h_{-n}) \in B(x, 3r_n) \right\} \right) \\ &= \nu_{T, T^{t-1}}^{x_{t-1}}(B(x, 3r_n)). \end{aligned}$$

Since this holds for all $\varepsilon > 0$ it follows that $\overline{\delta^{t-1}} \leq \delta + \frac{\kappa}{\chi} = \overline{\delta^t} + \gamma_{T^t, T^{t-1}}$ as claimed.

9. APPLICATION TO HITCHIN REPRESENTATIONS OF COMPACT SURFACE GROUPS

In this section, we first illustrate by an example the discussion of section 1.2. Take μ discrete in $\mathcal{M}(SL_3(\mathbb{R}))$. Consider the nine arrows of Figure 1 describing projections from $\mathcal{X}_T^{f'}$ to $\mathcal{X}_{T'}^{f'}$, where $T \xrightarrow{1} T'$, in dimension 3. By Theorem 2.2, six of these projections are dimension conserving. There is a natural family of examples of random walks on $SL_3(\mathbb{R})$ for which two of the other projections are not dimension conserving as soon as the middle exponent χ_2 is not 0. Namely, these are the random walks on images of a surface group by a Hitchin representation in $SL_3(\mathbb{R})$. We present these examples and then extend the discussion to Hitchin representations in $PSL_d(\mathbb{R})$, for all $d \geq 3$.

9.1. Hitchin component in dimension 3. Consider a closed surface Σ of genus at least two and the group $\Gamma := \pi_1(\Sigma)$. A representation $\rho : \Gamma \rightarrow PSL_2(\mathbb{R})$ is called *Fuchsian* if it is discrete and cocompact. A representation $\rho : \Gamma \rightarrow SL_3(\mathbb{R})$ is also called *Fuchsian* if it is the composition of a Fuchsian representation and a canonical irreducible representation of $PSL_2(\mathbb{R})$ into $SL_3(\mathbb{R})$. It is called *Hitchin* if it can be obtained by a deformation of a Fuchsian representation.

Hitchin representations have been studied from many points of view, we only list the properties we are going to use. We shall describe points in \mathcal{F} as pairs $(\zeta, \bar{\zeta})$, where ζ is a point in the projective plane \mathbb{RP}^2 and $\bar{\zeta}$ a line in \mathbb{RP}^2 containing ζ . The pairs $(\zeta, \bar{\zeta}), (\eta, \bar{\eta})$ are in general position if, and only if, $\zeta \notin \bar{\eta}, \eta \notin \bar{\zeta}$. By classical results of Koszul [Kos68], Goldman [Gol90] and Choi-Goldman [CG93], if the representation ρ is Hitchin, then there exists a C^1 convex subset $\Delta \subset \mathbb{RP}^2$ invariant under $\rho(\Gamma)$ and a Hölder continuous mapping $(\xi, \bar{\xi}) : \mathbb{S}^1 \rightarrow \mathcal{F}$ such that

- for $s \neq t \in \mathbb{S}^1$, $(\xi, \bar{\xi})(s)$ and $(\xi, \bar{\xi})(t)$ are in general position,
- $\xi(\mathbb{S}^1)$ is the boundary $\partial\Delta$,
- for $t \in \mathbb{S}^1$, $\bar{\xi}(t)$ is the tangent direction to $\partial\Delta$ at $\xi(t)$ and
- the set $\Lambda := (\xi, \bar{\xi})(\mathbb{S}^1)$ is an invariant set for the action of $\rho(\Gamma)$ on \mathcal{F} . The set Λ consists in the tangent elements to $\partial\Delta$.

So, the convex Δ admits a cocompact group of projective mappings, i.e. it is *divisible*. A classical result of Benzécri is that the boundary is of class C^2 if, and only if, Δ is an ellipse, if, and only if, the representation is conjugated to a Fuchsian representation. In that case, $\chi_2 = 0$ and all the dimension questions reduce

to the $PSL_2(\mathbb{R})$ case. Therefore, we may assume that representation ρ is Hitchin but not a Fuchsian representation. Such representations were studied in detail by Y. Benoist. In particular, he showed that

- the boundary $\partial\Delta$ is $C^{1+\beta}$ for some $\beta > 0$, but not $C^{1+abs.cont.}$ ([Be04]),
- the group $\rho(\Gamma)$ is Zariski dense in $SL_3(\mathbb{R})$ ([Be00]).

We denote $\mathcal{M}(\Gamma)$ the set of probability measures μ on Γ such that the group generated by the support of μ is Γ and $\sum_{\gamma} |\gamma| \mu(\gamma) < +\infty$, where $|\cdot|$ is some word metric on Γ . Let $\mu \in \mathcal{M}(\Gamma)$; consider the random walk $(\rho(\Gamma), \rho_*(\mu))$ and the stationary measures ν, ν' on \mathcal{F} . The measures ν and ν' are supported on Λ . For m -a.e. $\omega \in \Omega$, there are $(\xi_+, \bar{\xi}_+)(\omega)$ and $(\xi_-, \bar{\xi}_-)(\omega)$ distinct points in Λ that are the supports of the limit measures of $(g_{-1}(\omega) \dots g_{-n}(\omega))_* \nu$ and, respectively, $(g_0(\omega)^{-1} \dots g_{n-1}(\omega)^{-1})_* \nu'$ as $n \rightarrow +\infty$. The distribution of $(\xi_+, \bar{\xi}_+)(\omega)$ is ν , the distribution of $(\xi_-, \bar{\xi}_-)(\omega)$ is ν' . The point $\xi_+(\omega)$ is the direction of the expanding $E_1(\omega)$, the point $\xi_-(\omega)$ is the direction of the contracting $E_3(\omega)$. The central direction $E_2(\omega)$ is obtained as $\bar{\xi}_+(\omega) \cap \bar{\xi}_-(\omega)$.

Proposition 9.1. *Let ρ be a Hitchin representation of Γ and $\mu \in \mathcal{M}(\Gamma)$. Consider the random walk on $SL_3(\mathbb{R})$ directed by the probability $\rho_*(\mu)$, $\chi_1 > \chi_2 > \chi_3$ its Lyapunov exponents. Let $\mathcal{F}, \mathcal{L}, \mathcal{P}$ be the spaces of flags, lines and planes in \mathbb{R}^3 , $\nu, \nu_{\mathcal{L}}, \nu_{\mathcal{P}}$ the respective stationary measure and $\delta, \delta_{\mathcal{L}}, \delta_{\mathcal{P}}$ their dimensions.*

Assume $\chi_2 > 0$. Then, $\delta_{\mathcal{P}} < \delta_{\mathcal{L}} = \delta$. Moreover, the projections $\nu \rightarrow \nu_{\mathcal{P}}$ are not dimension conserving.

Assume $\chi_2 < 0$. Then, $\delta_{\mathcal{L}} < \delta_{\mathcal{P}} = \delta$. Moreover, the projections $\nu \rightarrow \nu_{\mathcal{L}}$ are not dimension conserving.

Assume $\chi_2 = 0$. Then, $\delta = \delta_{\mathcal{L}} = \delta_{\mathcal{P}}$. All the natural projections are dimension conserving.

Proof. By the above discussion, our results apply in this setting. We can consider

- The distribution ν of $f = (\xi_+, \bar{\xi}_+)(\omega) \in \mathcal{F}$. It has entropy h and dimension δ .
- The distribution $\nu_{\mathcal{L}}$ of $\xi_+(\omega)$. It has entropy $h_{\mathcal{L}}$ and dimension $\delta_{\mathcal{L}}$. Observe that, once one knows $\xi(t) \in \partial\Delta$, $\bar{\xi}(t)$ is the tangent direction to $\partial\Delta$ at $\xi(t)$, so it is uniquely determined. In other words, the projection from ν to $\nu_{\mathcal{L}}$ is a.e. one-to-one. By [Les21] $h_{\mathcal{L}} = h$, but, a priori, there is no dimension conservation and we only get $\delta \geq \delta_{\mathcal{L}}$.
- The distribution $\nu_{\mathcal{P}}$ of $\bar{\xi}_+(\omega)$. It has entropy $h_{\mathcal{P}}$ and dimension $\delta_{\mathcal{P}}$. Observe that, similarly, once one knows $\bar{\xi}(t)$ is a tangent direction to $\partial\Delta$ at some point, then this point is $\xi(t)$, so it is uniquely determined. In other words, the projection from ν to $\nu_{\mathcal{P}}$ is a.e. one-to-one. By [Les21] again, $h_{\mathcal{P}} = h$, but, a priori, there is no dimension conservation and we only get $\delta \geq \delta_{\mathcal{P}}$.

We choose $f' = (\eta, \bar{\eta}) \in \mathcal{F}$. Write $\mathcal{L}^{\mathcal{P}'} := \mathcal{X}_{\{1,3\},\{2\},\{3\}}^{f'}$, $\mathcal{L}' := \mathcal{X}_{\{1,3\},\{2,3\},\{3\}}^{f'}$ and $\mathcal{P}' := \mathcal{X}_{\{1,2,3\},\{2\},\{3\}}^{f'}$. For ν' -a.e. $f' \in \mathcal{F}$, write $\nu_{\mathcal{L}^{\mathcal{P}'}}^{f'}, \nu_{\mathcal{L}'}^{f'}, \nu_{\mathcal{P}'}^{f'}$ for the corresponding conditional measures.

We can also consider the distribution $\nu_{\mathcal{L}^{\mathcal{P}'}}^{f'}$ of the couple made of the point $\bar{\xi}_+ \cap \bar{\eta}$ and the line (η, ξ_+) given $f' = (\eta, \bar{\eta})$. It has entropy $h_{\mathcal{L}^{\mathcal{P}'}}$ and dimension $\delta_{\mathcal{L}^{\mathcal{P}'}}$. Again, this determines $(\xi, \bar{\xi})$ by intersection with $\partial\Delta$ and the dimension on fibers

of the projection from \mathcal{F} to $\mathcal{X}_{\mathcal{L}\mathcal{P}}^{f'}$ is 0. But now, by theorem 2.2, there is dimension conservation, so $h = h_{\mathcal{L}\mathcal{P}}$ and $\delta = \delta_{\mathcal{L}\mathcal{P}}$.

Assume $\chi_2 \geq 0$. We project both $\nu_{\mathcal{L}}$ and $\nu_{\mathcal{L}\mathcal{P}}^{f'}$ to the space \mathcal{L}' of lines going through η by associating the line going through ξ and η in the first case and by forgetting $\bar{\xi} \cap \bar{\eta}$ in the second case. The projection and the image measure $\nu_{\mathcal{L}'}^{f'}$ depend on f' . For ν' -a.e. f' , we have entropy $h_{\mathcal{L}'}$ and dimension $\delta_{\mathcal{L}'}$ on \mathcal{L}' . Both projections have almost everywhere trivial fibers: intersecting the line (η, ξ) with $\partial\Delta$ determines everything. Therefore,

$$h = h_{\mathcal{L}} = h_{\mathcal{L}\mathcal{P}} = h_{\mathcal{L}'}$$

Moreover, by theorem 2.2, both projections have dimension conservation (observe that $\chi_{\mathcal{L},\mathcal{L}'} = \chi_1 - \chi_3 \geq \chi_{\mathcal{L}'} = \chi_1 - \chi_2$). So we obtain

$$\delta = \delta_{\mathcal{L}} = \delta_{\mathcal{L}\mathcal{P}} = \delta_{\mathcal{L}'} = \frac{h}{\chi_1 - \chi_2}.$$

Remain to understand the projections of both $\nu_{\mathcal{P}}$ and $\nu_{\mathcal{L}\mathcal{P}}^{f'}$ on the space \mathcal{P}' of points of $\bar{\eta}$. The projection and the image measure $\nu_{\mathcal{P}'}^{f'}$ depend on f' . For ν' -a.e. f' , write $h_{\mathcal{P}'}$ and $\delta_{\mathcal{P}'}$ for the entropy and the dimension of $\nu_{\mathcal{P}'}^{f'}$. Once more, knowing the point E_2 in $\bar{\eta}$ determines the rest by drawing the unique other tangent to $\partial\Delta$ going through E_2 . So,

$$h_{\mathcal{P}} = h_{\mathcal{P}'} = h_{\mathcal{L}\mathcal{P}}$$

(both $h_{\mathcal{P}}$ and $h_{\mathcal{L}\mathcal{P}}$ are h by the above discussion) and all the entropies are the same h . Moreover, since $\chi_{\mathcal{P},\mathcal{P}'} = \chi_1 - \chi_3 \geq \chi_{\mathcal{P}'} = \chi_2 - \chi_3$,

$$\delta_{\mathcal{P}} = \delta_{\mathcal{P}'} = \frac{h}{\chi_2 - \chi_3}.$$

If $\chi_2 = 0$, then $\chi_1 - \chi_2 = \chi_2 - \chi_3$, $\delta_{\mathcal{L}'} = \delta_{\mathcal{P}'}$, all the dimensions coincide and there is dimension conservation at all the projections of Figure 1.

If $\chi_2 > 0$, then $\chi_2 - \chi_3 > \chi_1 - \chi_2$ and $\delta_{\mathcal{P}'} < \delta_{\mathcal{L}'}$. So $\delta_{\mathcal{P}} < \delta$ and the projection from ν to $\nu_{\mathcal{P}}$ is not dimension conserving. In the same way, $\delta_{\mathcal{P}'} < \delta_{\mathcal{L}\mathcal{P}}$ and, for ν' -a.e. f' , the projection from $\nu_{\mathcal{L}\mathcal{P}}^{f'}$ to $\nu_{\mathcal{P}'}^{f'}$ is not dimension conserving either.

In the case when $\chi_2 \leq 0$, the discussion is the same, exchanging the role of points and lines and of $\chi_2 - \chi_3$ and $\chi_1 - \chi_2$. \square

Proof of theorem 1.5 in dimension $d = 3$. By the above proof, we have $h = \delta(\nu) \min\{(\chi_1 - \chi_2), (\chi_2 - \chi_3)\}$, independently of the sign of χ_2 . Theorem 1.5 follows when $d = 3$. \square

9.2. Rigidity of Hitchin representations. In this section, we prove that the hypotheses of proposition 9.1 are satisfied for some probability measure in $\mathcal{M}(\rho(\gamma))$ if the representation ρ is Hitchin but not Fuchsian, namely that one can find such a measure with $\chi_2 \neq 0$. We have the

Theorem 9.2. *Let ρ be a Hitchin representation of a cocompact surface group in $SL_3(\mathbb{R})$ such that for all probability measures in $\mathcal{M}(\rho(\Gamma))$, $\chi_2 \leq 0$. Then, the representation ρ is Fuchsian.*

Such variational characterizations of Fuchsian representations among Hitchin components have been proven by M. Crampon ([Cr09]) and R. Potrie and A. Sambarino ([PS17]) in greater generality. Theorem 9.2 is a variant of their results adapted to the dimension 3.

Proof. By proposition 9.1, our hypothesis is that for all $\mu \in \mathcal{M}(\Gamma)$, the dimensions $\delta_{\mathcal{L}}, \delta_{\mathcal{P}}$ of the stationary measures on the spaces of lines and planes satisfy

$$(26) \quad \delta_{\mathcal{L}} \leq \delta_{\mathcal{P}}.$$

We are going to use thermodynamical formalism for the geodesic flow on $\rho(\Gamma) \backslash H\Delta$, where $H\Delta$ is the homogeneous tangent bundle to Δ and a construction of [CM07] to obtain, for any Hitchin representation ρ , some $\mu \in \mathcal{M}(\rho(\Gamma))$ such that $\delta_{\mathcal{L}} = 1$. Since $\nu_{\mathcal{P}}$ is also supported on a C^1 circle, (26) implies that $\delta_{\mathcal{P}} = 1$ as well. Using dynamics of the geodesic flow and [Be04], section 6, this will imply that the representation is Fuchsian.

Recall that all matrices $\rho(\gamma), \gamma \in \Gamma, \rho(\gamma) \neq Id$, have three distinct real eigenvalues with absolute values $e^{\ell_1(\gamma)} > e^{\ell_2(\gamma)} > e^{\ell_3(\gamma)}$ ([Lab06]). Let φ be the linear functional on $\Sigma := \{(\ell_1, \ell_2, \ell_3) \in \mathbb{R}^3 : \ell_1 + \ell_2 + \ell_3 = 0\}$ defined by $\varphi := \ell_1 - \ell_2$. Recall that the geodesic flow on $\rho(\Gamma) \backslash H\Delta$ is an Anosov flow. There exists a Hölder continuous function f on $H\Delta$ such that for any $\gamma \in \Gamma, \gamma \neq Id$,

$$\ell_1(\gamma) - \ell_2(\gamma) = \int_{\sigma_\gamma} f,$$

where σ_γ is the periodic orbit associated to γ (see [PS17], sections 2 and 7). Moreover, for any ergodic invariant measure m for the geodesic flow, $\int f dm$ is the positive Lyapunov exponent of the geodesic flow for m ([Be04], Lemma 6.5). In particular, the equilibrium measure m_0 for $-f$ is absolutely continuous along unstable manifolds.

Fix a point $o \in \Delta$. Then, the *Gibbs-Patterson-Sullivan construction* (see e.g. [Led94]) yields an equivariant family of measures ν_0 at the boundary such that for all $\gamma \in \Gamma$, $\frac{d(\rho(\gamma))_* \nu_0}{d\nu_0}(\xi)$ is a Hölder continuous function and with the property that, if a set A of points in $\partial\Delta$ is ν_0 -negligible, then the set of geodesics with end in A is m_0 -negligible. By the absolute continuity of the stable foliation, this implies that ν_0 is absolutely continuous on $\partial\Delta$.

Recall that the limit set Λ of $\rho(\Gamma)$ projects one-to-one in $\partial\Delta$. Denote by ν the lift of the measure ν_0 to Λ .

Next step consists in finding a random walk in $\mathcal{M}(\rho(\Gamma))$ such that ν is the stationary measure on \mathcal{F} or equivalently such that ν_0 is the stationary measure for the action on $\partial\Delta$.

Lemma 9.3. *Let Γ be a co-compact group of isometries of \mathbb{H}^2 , ρ a Hitchin non-Fuchsian representation of Γ in $SL_3(\mathbb{R})$, Δ the open convex proper subset of \mathbb{RP}^2 invariant under $\rho(\Gamma)$, ν_0 be a finite measure on $\partial\Delta$ such that for all $\gamma \in \Gamma$, $\frac{d(\rho(\gamma))_* \nu_0}{d\nu_0}(\xi)$ is a Hölder continuous function. Then there exists a probability measure $\mu_0 \in \mathcal{M}(\Gamma)$ such that ν_0 is $\rho_*(\mu_0)$ -stationary.*

Proof. We can apply [CM07], theorem 1.1, to the action of Γ on the hyperbolic plane with the measure $\nu_{\mathbb{S}} = (\xi^{-1})_*\nu_0$. Since the mapping ξ is Hölder continuous and Γ -equivariant, the measure $\nu_{\mathbb{S}}$ has Hölder continuous Radon-Nikodym derivatives under the action of Γ as well. Let μ_0 be the measure given by [CM07] Theorem 1.1 and such that $\nu_{\mathbb{S}}$ is the stationary measure under μ . The measure μ has whole support on Γ and satisfies $\sum_g \mu_0(g)d(o, go) < +\infty$ ([CM07], page 488). It does indeed belong to $\mathcal{M}(\Gamma)$. \square

To summarize, the measure $\mu := \rho_*\mu_0$ on $\rho(\Gamma)$ has the property that $\nu_{\mathcal{L}}$ is absolutely continuous and is the measure at infinity of the SRB measure of the geodesic flow on $H\Delta$. Moreover, by (26), $\delta_{\mathcal{P}} = 1$.

Consider the dual representation $\rho^*(\gamma) = (\rho(\gamma)^t)^{-1}$ and the measure $\mu^* := (\rho^*)_*\mu_0$. The exponents of the random walk (Γ, μ^*) are the opposite $-\chi_3 > -\chi_2 > -\chi_1$ and we claim that the entropy h^* is the same entropy $h^* = h$. Therefore, the dimension of the stationary measure $\nu_{\mathcal{L}^*}$ is $\delta_{\mathcal{P}} = 1$. By the variational principle again, $\nu_{\mathcal{L}^*}$ is absolutely continuous. By [Be04] Proposition 6.2, the representation is Fuchsian.

To prove the claim, observe that, since $\rho(\Gamma)$ is discrete in $SL_3(\mathbb{R})$, the entropy h is given by the random walk entropy $h = h_{\text{RW}}(\mu) := \lim_n \frac{1}{n} H(\mu^{(n)})$ ([Led85]), which is the same for μ and μ^* . \square

Using the dual representation ρ^* , we also have

Corollary 9.4. *Let ρ be a Hitchin representation of a cocompact surface group in $SL_3(\mathbb{R})$ such that for all probability measures in $\mathcal{M}(\rho(\Gamma))$, $\chi_2 \geq 0$. Then, the representation ρ is Fuchsian.*

9.3. Hitchin components in higher dimensions. Consider the surface group Γ . As before, a representation $\rho : \Gamma \rightarrow PSL_d(\mathbb{R})$ is called *Fuchsian* if it is the composition of a Fuchsian representation and the canonical irreducible representation of $PSL_2(\mathbb{R})$ into $PSL_d(\mathbb{R})$. It is called *Hitchin* if it can be obtained by a deformation of a Fuchsian representation. Geometric properties of Hitchin representations have been studied, notably by F. Labourie (see [Lab06], [Lab07] for history, background, the properties we use below and much more). Let $\rho : \gamma \rightarrow PSL_d(\mathbb{R})$ be a Hitchin representation and denote again by $\rho(\Gamma)$ a lift of the representation to $SL_d(\mathbb{R})$. By [G08], proposition 14, the action of $\rho(\Gamma)$ on \mathbb{R}^d is strongly irreducible: there is no finite union of proper vector subspaces of \mathbb{R}^d that is invariant under $\rho(\Gamma)$. By [Lab06], theorem 1.5, the matrix $\rho(\gamma)$, for γ non-trivial has all eigenvalues real and distinct. In particular, for γ non trivial, there is a unique attracting fixed point γ^+ for the action of $\rho(\gamma)$ on \mathcal{F} . By definition, the *limit set* Λ is the closure of the set of all $\gamma^+, \gamma \neq Id \in \Gamma$. Moreover, the projection from the limit set Λ to \mathbb{RP}^{d-1} is one-to-one ([Lab06] theorem 4.1).

Let $\mu \in \mathcal{M}(\Gamma)$. We claim that $\mu \in \mathcal{M}(d)$: on the one hand, the action is proximal on all exterior products and by [GR89], the exponents $\chi_1 > \dots > \chi_d$ are distinct; on the other hand, since the action on \mathbb{R}^d is strongly irreducible, there is a unique stationary measure ν on \mathbb{RP}^{d-1} . That measure has a unique lift to Λ and therefore,

there is a unique stationary measure on \mathcal{F} . For the same reason, there is a unique stationary measure for μ' on \mathcal{F} . All our discussion and theorem 1.3 apply, the measure ν is exact dimensional with dimension δ and entropy $h := h(\mathcal{F}, \mu, \nu)$.

Proposition 9.5. *Let $\lambda := \inf_{i < j} (\chi_i - \chi_j)$. Then $\delta = h/\lambda$. More generally, let $T \neq T_0$ be an admissible topology, κ_T, δ_T as defined in (5) and corollary 2.4. Let $\lambda_T := \inf_{i < j, j \notin T(i)} (\chi_i - \chi_j)$. Then, $\kappa_T = h$ and $\delta_T = h/\lambda_T$.*

In particular, theorem 1.5 follows in all dimensions. Another consequence is that, as in dimension 3, comparing λ and λ_T is enough to decide whether the projection from \mathcal{F} to $\mathcal{X}_T^{f'}$ is dimension conserving for ν' -a.e. f' .

Corollary 9.6. *Let T be an admissible topology such that $T_1 \stackrel{1}{\prec} T$. Then $\delta_T = \delta$ unless there is a unique i with $\lambda = \chi_i - \chi_{i+1}$ and furthermore $T = \{1\}, \{2\}, \dots, \{i, i+1\}, \dots, \{d\}$.*

Proof of Proposition 9.5. The measure ν is supported by the limit set $\Lambda \subset \mathcal{F}$. Labourie showed that Λ is a *hyperconvex Frenet* curve with *Property (H)*. Namely:

- (1) There is a Hölder continuous Γ -equivariant mapping $\xi : \mathbb{S}^1 \rightarrow \Lambda$,

$$\xi(t) = \{0\} \subset \xi_1(t) \subset \dots \subset \xi_i(t) \subset \dots \subset \xi_d(t) = \mathbb{R}^d.$$

- (2) For any distinct points t_1, \dots, t_ℓ integers d_1, \dots, d_ℓ with $p := \sum_{j=1}^{\ell} d_j$, the following sum is direct

$$\xi_{d_1, \dots, d_\ell}(t_1, \dots, t_\ell) := \xi_{d_1}(t_1) \oplus \dots \oplus \xi_{d_\ell}(t_\ell)$$

and, if the distinct t_1, \dots, t_ℓ all converge to x , then $\xi_{d_1, \dots, d_\ell}(t_1, \dots, t_\ell)$ converge to $\xi_p(x)$.

- (3) for any triple of distinct points (s, t, t') , any integer $i, 0 < i < d$,

$$\xi_i(s) \oplus (\xi_i(t) \cap \xi_{d-i+1}(t')) \oplus \xi_{d-i-1}(t') = \mathbb{R}^d.$$

Property (2) defines a hyperconvex Frenet curve ([Lab06], theorem 1.4) and property (3) is relation (6) in [Lab06] theorem 4.1. (Property (3) is called Property (H) in [Lab06] section 7.1.4.)

Let T^1 be an admissible topology such that $T^1 \stackrel{1}{\prec} T_0$. We claim that there is a unique integer $i, 0 < i < d$, such that

$$T^1(k) = \{k, k+1, \dots, d\} \text{ for } k \neq i, \quad T^1(i) = \{i, i+2, \dots, d\}.$$

Indeed, by definition, there is $i, 0 < i < d$, such that $T^1(k) = T_0(k)$ for $k \neq i$ and $j > i$ such that $T^1(i) = T_0(i) \setminus \{j\}$. By Proposition 3.1, $T^1(i) \setminus \{i, j\} \in T_0$, and this is possible only if $j = i+1$.

Fix $t' \in \mathbb{S}^1$ and set $f' := \xi(t')$. By lemma 3.4, the set $\mathcal{X}_{T^1}^{f'}$ is bilipschitz homeomorphic to an open interval. We associate to $t \in \mathbb{S}^1, t \neq t'$, a configuration $\Psi(t) \in \mathcal{X}_{T^1}^{f'}$ by setting:

$$\begin{aligned} \Psi(t)_{T^1(k)} &= \xi_{d-k+1}(t') \text{ for } k \neq i, \\ \Psi(t)_{T^1(i)} &= (\xi_i(t) \cap \xi_{d-i+1}(t')) \oplus \xi_{d-i-1}(t'). \end{aligned}$$

Lemma 9.7. *The mapping Ψ is an homeomorphism between $\mathbb{S}^1 \setminus \{t'\}$ and its image in $\mathcal{X}_{T^1}^{f'}$.*

Proof. By the hyperconvexity property (2), $\dim(\xi_i(t) \cap \xi_{d-i+1}(t')) = 1$ and that space is in general position with respect to $\xi_{d-i-1}(t')$; therefore the mapping Ψ is continuous. Since Ψ is a mapping between two open intervals, it suffices to show that Ψ is one-to-one. Assume by contradiction that there is $s \neq t, t'$ such that

$$(\xi_i(s) \cap \xi_{d-i+1}(t')) \oplus \xi_{d-i-1}(t') = (\xi_i(t) \cap \xi_{d-i+1}(t')) \oplus \xi_{d-i-1}(t').$$

By property (3), $\xi_i(s)$ should be in direct sum with $(\xi_i(s) \cap \xi_{d-i+1}(t')) \oplus \xi_{d-i-1}(t')$, which is possible only if $\xi_i(s) \cap \xi_{d-i+1}(t') = \{0\}$. This contradicts hyperconvexity. \square

By lemma 9.7, for any $x' \in \mathcal{X}_{T^1}^{f'}$, there at most one point $s \in \mathbb{S}^1$ such that $\pi_{T^1, T^1}(\xi(s)) = x'$, i.e. $(\pi_{T^1, T^1})^{-1}(x')$ is at most one point. So, $\kappa_{T^1, T^1} = 0$. By theorem 1.6, we have $\delta_{T^1} = h/\chi_{T^1, T_0}$.

For a general admissible topology T , we apply proposition 2.1 and obtain a topology T^1 such that $T \prec T^1 \stackrel{1}{\prec} T_0$ and $\chi_{T^1, T_0} = \lambda_T$. Since $T_1 \prec T \prec T^1$, $\kappa_{T, T^1} = 0$ and $\kappa_T = h$. By (7) and corollary 2.4, δ_T is the same as $\delta_{T^1} = h/\lambda_T$. \square

REFERENCES

- [Be00] Yves Benoist. Automorphismes des cônes convexes *Invent. Math.*, 141(1): 149–193, 2000.
- [Be04] Yves Benoist. Convex divisibles, I *Algebraic groups and arithmetic, Tata Inst. Fund. Res. Stud. Math.* 17: 181–237 (2004).
- [BHR19] Balázs Bárány, Michael Hochman and Ariel Rapaport. Hausdorff dimension of planar self-affine sets and measures *Invent. Math.* 216 (3): 601–659, (2019).
- [BK17] Balázs Bárány and Antti Käenmäki. Ledrappier-Young formula and exact dimensionality of self-affine measures *Adv. Math.*, 318: 88–129, (2017).
- [BPS] Balázs Bárány, Mark Pollicott and Károly Simon. Stationary measures for projective transformations *Jour. Stat. Physics*, 148: 393–421, (2012).
- [BQ16] Yves Benoist and Jean-François Quint. Random walks on reductive groups *Ergebnisse der Mathematik und ihrer Grenzgebiete*, vol. 62, Springer 2016.
- [BQ18] Yves Benoist and Jean-François Quint. On the regularity of stationary measures *Israel J. Math.* 226: 1–14 (2018).
- [B12] Jean Bourgain. Finitely supported measures on $SL_2(\mathbb{R})$ which are absolutely continuous at infinity, in *Geometric aspects of functional analysis. Lect. Notes in Math.* 2050: 133–141, Springer, Heidelberg (2012).
- [CG93] Suhyoung Choi and William M. Goldman. Convex real projective structures on surfaces are closed *Proc. Amer. Mat. Soc.* 118: 657–661 (1993).
- [CM07] Chris Connell and Adam Muchnik. Harmonicity of Gibbs measures *Duke Math. J.* 137: 461–509 (2007).
- [Cr09] Mickaël Crampon. Entropies of compact strictly convex projective manifolds *J. of Mod. Dyn.* 3: 511–547 (2009).
- [D80] Yves Derriennic. Quelques applications du théorème ergodique sous-additif *Conference on random walks (Kleebach, 1979), Astérisque*, 74: 283–301 (1980).
- [DO80] Adrien Douady and Joseph Oesterlé. Dimension de Hausdorff des attracteurs *Comptes Rendus Acad. Sci.*, 290: 1136–1138 (1980).
- [Dob59] Roland L. Dobrušin. A general formulation of the fundamental theorem of Shannon in the theory of information *Uspehi Mat. Nauk*, 14: 3–104, (1959).
- [F88] Kenneth J. Falconer. The Hausdorff dimension of self-affine fractals *Math. Proc. Cambridge Phil. Soc.* 103(2): 339–350 (1988).

- [FFJ15] Kenneth Falconer, Jonathan Fraser and Xiong Jin. Sixty years of Fractal Projections *Fractal Geometry and Stochastics V; C. Bandt, K. Falconer and M. Zähle eds*, Progress in Probability, 70, Birkhäuser, Cham (2015), 3–25.
- [Fen19] De-Jun Feng. Dimension of invariant measures for affine iterated function systems, ArXiv 1901.01691.
- [FH09] De-Jun Feng and Huyi Hu. Dimension theory of iterated function systems *Comm. Pure Appl. Math.*, 62: 1435–1500, (2009).
- [F63] Harry Furstenberg. Noncommuting random products *Trans. Amer. Math. Soc.*, 108: 377–428 (1963).
- [F70] Harry Furstenberg. Intersections of Cantor sets and transversality of semi-groups *Problems in Analysis (Symp. Solomon Bochner, Princeton University, Princeton N.J. 1969)* Princeton University Press, Princeton, N.J. (1970), 41–59.
- [F71] Harry Furstenberg. Random walks and discrete subgroups of Lie groups. *Advances in Probability and Related Topics*, 1: 1–63, Dekker, New York (1971).
- [F08] Harry Furstenberg. Ergodic fractal measures and dimension conservation *Ergod. Th. & Dynam. Sys.*, 28: 405–422 (2008).
- [GFY] Israel M. Gel'fand, and Akiva M. Yaglom. Calculation of the amount of information about a random function contained in another such function, *Amer. Math. Soc. Transl. (2)*, 12: 199–246 (1959).
- [GM89] Ilya Ya. Gol'dsheid and Gregory A. Margulis. Lyapunov exponents of a product of random matrices *Russian Math Surveys*, 44: 11–71 (1989).
- [Gol90] William M. Goldman. Convex real projective structures on compact surfaces. *J. Diff. Geom.*, 31: 791–845, 1990
- [G08] Olivier Guichard Composantes de Hitchin et représentations hyperconvexes de surfaces *J. Differential Geom.*, 80: 391–431 (2008).
- [G90] Yves Guivarc'h. Produits de matrices aléatoires et applications *Ergod. Th. Dynam. Sys.*, 10: 483–512 (1990).
- [GL93] Yves Guivarc'h and Yves Le Jan. Asymptotic winding of the geodesic flow on modular surfaces and continued fractions *Ann. Sci. ENS*, 26: 23–50, (1993).
- [GR89] Yves Guivarc'h and Albert Raugi. Propriétés de contraction d'un semi-groupe de matrices inversibles. Coefficients de Liapunoff d'un produit de matrices aléatoires indépendantes *Israel J. Math.*, 65: 165–196 (1989).
- [H14] Michael Hochman. On self-similar sets with overlaps and inverse theorems for entropy *Ann. Math. (2)* 180(2): 773–822, (2014).
- [H15] Michael Hochman. On self-similar sets with overlaps and inverse theorems for entropy in \mathbb{R}^d to appear (*Memoirs of the A.M.S.*) ArXiv 1503.09043.
- [HR19] Michael Hochman and Ariel Rapaport. Hausdorff dimension of planar self-affine sets and measures with overlaps *preprint* ArXiv 1904. 09812.
- [HS17] Michael Hochman and Boris Solomyak. On the dimension of Furstenberg measure for $SL_2(\mathbb{R})$ random matrix products *Invent. Math.*, 210(3): 815–875, (2017).
- [JL04] Esa Järvenpää, Maarit Järvenpää and Marta Llorante. Local dimensions of sliced measures and stability of packing dimensions of sections of sets *Adv. maths.*, 183: 127–154 (2004).
- [JM98] Maarit Järvenpää and Pertti Mattila. Hausdorff and packing dimensions and sections of measures *Mathematika*, 45: 55–77 (1998).
- [KLP11] Vadim A. Kaimanovich and Vincent Le Prince. Matrix random products with singular harmonic measure. *Geom. Dedicata*, 150:257–279, (2011).
- [Kos68] Jean-Louis Koszul. Déformations de connexions localement plates *Ann. Inst. Fourier (Grenoble)*, 18: 103–114, (1968).
- [KV83] Vadim A. Kaimanovich and Anatoly M. Vershik. Random walks and discrete groups: boundary and entropy *Ann. Probab.*, 11: 457–490, (1983).
- [KY79] James L. Kaplan and James A. Yorke. Chaotic behavior of Multidimensional Difference Equation *Springer Lect. Notes*, 730: 204–277 (1979).
- [Lab06] François Labourie. Anosov Flows, Surface Groups and Curves in Projective Space *Invent. Math.* 165: 51–114 (2006).
- [Lab07] François Labourie. Cross ratios, surface groups, $PSL(n, \mathbb{R})$ and diffeomorphisms of the circle *Publ. Math. Inst. Hautes Études Sci.* 106: 139–213 (2007).

- [Led84] François Ledrappier. Quelques propriétés des exposants caractéristiques In *École d'été de probabilités de Saint-Flour, XII—1982*, volume 1097 of *Lecture Notes in Math.*, 305–396. Springer, Berlin, (1984).
- [Led85] François Ledrappier. Poisson boundaries of discrete groups of matrices *Israel J. Math.*, 50: 319–336 (1985).
- [Led94] François Ledrappier. Structure au bord des variétés à courbure négative *Séminaire de théorie spectrale et géométrie de Grenoble* 71: 97–122, (1994/1995)
- [Les21] Pablo Lessa. Entropy and dimension of disintegrations of stationary measures *Trans. Amer. Math. Soc. Ser. B*, 8: 105–129 (2021).
- [LY85] François Ledrappier and Lai-Sang Young. The metric entropy of diffeomorphisms. II. Relations between entropy, exponents and dimension *Ann. of Math. (2)*, 122(3): 540–574, (1985).
- [O68] Valery I. Oseledets. A multiplicative ergodic theorem. Characteristic Ljapunov exponents of dynamical systems *Trudy Moskov. Mat. Obsc* 19:179–210 (1968).
- [Per59] Albert Perez. Information theory with an abstract alphabet. Generalized forms of McMillan's limit theorem for the case of discrete and continuous times *Theor. Probability Appl.*, 4: 99–102 (1959).
- [PS17] Rafael Potrie and Andrés Sambarino. Eigenvalues and entropy of a Hitchin representation *Invent. Math.*, 209: 885–925 (2017).
- [Rap17] Ariel Rapaport. A self-similar measure with dense rotations, singular projections and discrete slices *Adv. Math.* 321: 529–546 (2017).
- [Rap21] Ariel Rapaport. Exact dimensionality and Ledrappier-Young formula for the Furstenberg measure *T.A.M.S.* 374: 5225–5268 (2021).
- [Sha48] Claude E. Shannon. A mathematical theory of communication *Bell System Tech. J.*, 27: 379–423, 623–656 (1948).
- [She18] Weixiao Shen. Hausdorff dimension of the graphs of the classical Weierstrass functions *Math.Z.*, 289: 223–266 (2018).
- [S15] Pablo Shmerkin. Projections of self-similar and related fractals: a survey of recent developments *Fractal Geometry and Stochastics V; C. Bandt, K. Falconer and M. Zähle eds*, Progress in Probability, 70, Birkhäuser, Cham (2015), 33–74.
- [Y82] Lai-Sang Young. Dimension, Entropy and Lyapunov exponents *Ergodic Theory Dynam. Systems.* 2: 109–124 (1982).

FRANÇOIS LEDRAPPIER, UNIVERSITÉ DE PARIS ET SORBONNE UNIVERSITÉ, CNRS, LPSM, BOÎTE COURRIER 158, 4, PLACE JUSSIEU, 75252 PARIS CEDEX 05, FRANCE,

Email address: `fledrapp@nd.edu`

PABLO LESSA, IMERL, FACULTAD DE INGENIERÍA, JULIO HERRERA Y REISSIG 565, 11300 MONTEVIDEO, URUGUAY

Email address: `plessa@fing.edu.uy`