

# **The hyperserial field of surreal numbers**

Vincent Bagayoko, Joris van der Hoeven

## **To cite this version:**

Vincent Bagayoko, Joris van der Hoeven. The hyperserial field of surreal numbers. 2021. hal-03232836

# **HAL Id: hal-03232836 <https://hal.science/hal-03232836>**

Preprint submitted on 22 May 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## The hyperserial field of surreal numbers<sup>\*</sup>

Vincent Bagayoko<sup> $A$ </sup>, UMons, LIX Joris van der Hoeven<sup>B</sup>, CNRS, LIX

For any ordinal  $\alpha > 0$ , we show how to define a hyperexponential  $E_{\omega^{\alpha}}$  and a hyperlogarithm  $L_{\omega^{\alpha}}$  on the class  $\text{No}^{>,\succ}$  of positive infinitely large surreal numbers. Such functions are archetypes of extremely fast and slowly growing functions at infinity. We also show that the surreal numbers form a so-called hyperserial field for our definition.

#### **Introduction**  $\mathbf{1}$

The ordered field  $\bf{No}$  of surreal numbers was introduced by Conway in [11]. Conway originally used transfinite recursion to define both the surreal numbers (henceforth called numbers), the ordering on **No**, and the ring operations. For any two sets L and R of numbers with  $L < R$  (i.e.  $x < y$  for all  $x \in L$  and  $y \in R$ ), there exists a number  $\{L \mid R\}$  with

$$
L \, < \, \{L \, | \, R\} \, < \, R,
$$

and all numbers can be obtained in this way. Given  $x = \{x_L | x_R\}$  and  $y = \{y_L | y_R\}$ , we have

$$
x + y \; := \; \{x_L + y, x + y_L \; | \; x_R + y, x + y_R\}
$$

and similar recursive formulas exist for  $-x$ ,  $x y$  and for deciding whether  $x = y$ ,  $x \le y$ , and  $x < y$ . It is truly remarkable that No turns out to be a totally ordered real-closed field for such "simple" definitions [11]. The bracket  $\{ | \}$  is called the *Conway bracket*. Using this bracket, we obtain a surreal number in any traditional Dedekind cut, which allows us to embed  $\mathbb R$  into No. In addition, No contains all ordinal numbers

$$
0 = \{ | \}, \quad 1 = \{0 | \}, \quad 2 = \{0, 1 | \}, \quad \dots, \quad \omega = \{0, 1, 2, \dots | \}, \quad \omega + 1 = \{0, 1, 2, \dots, \omega | \}, \quad \dots,
$$

so No is actually a proper class.

An interesting question is which other operations from calculus can be extended to the surreal numbers. Gonshor has shown how to extend the real exponential function to the surreal numbers [19] and the resulting exponential field  $(No, \exp)$  turns out to be elementarily equivalent to ( $\mathbb{R}$ , exp) [13]. Berarducci and Mantova recently defined a derivation with respect to  $\omega$  on the surreals [9], again with good model-theoretic properties [2]. In collaboration with Mantova, the authors constructed a surreal solution to the functional equation

$$
E_{\omega}(x+1) = \exp E_{\omega} x,
$$

which is a bijection of  $\mathbf{No}^{>,\geq} := \{x \in \mathbf{No} : x > \mathbb{R}\}\$  onto itself [6]. We call  $E_{\omega}$  a *hyperexponential* and its functional inverse  $L_{\omega}$  a *hyperlogarithm*.

A. vincent.bagayoko@umons.ac.be

B. vdhoeven@lix.polytechnique.fr

<sup>\*.</sup> This article has been written using GNU TEXMACS [27].

The first goal of this paper is to extend the results from [6] to the construction of hyperexponentials  $E_{\omega} \alpha$ : **No**<sup>>,></sup>  $\rightarrow$  **No**<sup>>,></sup> of any ordinal *force*  $\alpha$ , together with their functional inverses  $L_{\omega} \alpha$ . If  $\alpha = \beta + 1$  is a successor ordinal, then  $E_{\alpha}$  satisfies the functional equation

$$
E_{\omega}^{\beta+1}(x+1) = E_{\omega}^{\beta}(E_{\omega}^{\beta+1}(x))
$$

Our second goal is to show that these hyperexponentials are "well-behaved" in the sense that they endow **No** with the structure of a *hyperserial field* in the sense of [5].

#### **Motivation and background**  $1.1$

Whereas it is natural to study surreal exponentiation and differentiation, it may seem more exotic to define and investigate the properties of surreal hyperexponentials and hyperlogarithms. In fact, the main motivation behind our work is a conjecture by the second author  $[26, p. 16]$ and a research program that was laid out in  $[1]$  for proving this conjecture. The ultimate goal is to expose the deep connections between two types of mathematical infinities: numerical infinities and growth rates at infinity. Let us briefly recall the rationale behind this connection.

Cantor's ordinal numbers provide us with a way to count beyond all natural numbers and to keep counting beyond the size of any set. However, ordinal arithmetic is rather poor in the sense that we have no subtraction or division and that addition and multiplication do not satisfy the usual laws of arithmetic, such as commutativity. We may regard Conway's surreal numbers as providing a calculus with Cantor's ordinal numbers which does extend the usual calculus with real numbers. In this sense, Conway managed to construct the ultimate framework for computations with numerical infinities.

Another source for computations with infinitely large quantities stems from the study of growth rates of real functions at infinity. The first major results towards a systematic asymptotic calculus of this kind are due to Hardy in [21, 22], based on earlier ideas by du Bois-Reymond [15, 16, 17]. Hardy defined an *L*-function to be a function constructed from x and the real numbers  $\mathbb R$  using the field operations, exponentiation, and logarithms. He proved that the germs of L-functions at infinity form a totally ordered field. The framework of L-functions is suitable for asymptotic analysis since we have an ordering for comparing the growth at infinity of any two such functions. This is often rephrased by saying that L-functions have a regular growth at infinity.

Hardy also observed [21, p. 22] that "The only scales of infinity that are of any practical importance in analysis are those which may be constructed by means of the logarithmic and exponential functions." In other words, Hardy suggested that the framework of L-functions not only allows for the development of a systematic asymptotic calculus, but that this framework is also sufficient for all "practical" purposes. Alas, there are several "holes". First of all, the framework is not closed under various useful operations such as functional inversion and integration. Secondly, the framework does not contain any functions of extremely fast or slow growth at infinity, like  $E_{\omega}$  and  $L_{\omega}$ , although such functions naturally appear in the analysis of certain algorithms. For instance, the best known algorithm for multiplying two polynomials of degree *n* in  $\mathbb{F}_2[x]$  runs in time  $O(n \log n 4^{L_{\omega}n})$ ; see [23].

This raises the question how to construct a truly universal framework for computations with regular functions at infinity. Our next candidate is the class of transseries. A transseries is a formal object that is constructed from x (with  $x \to \infty$ ) and the real numbers, using exponentiation, logarithms, and *infinite* sums. One example of a transseries is

$$
e^{e^x + e^{x/2} + e^{x/3} + \dots} - 3e^{x^2} + 5(\log x)^{\pi} + 42 + x^{-1} + 2x^{-2} + 6x^{-3} + 24x^{-4} + \dots + e^{-x}.
$$

Depending on conditions satisfied by their supports, there are different types of transseries. The first constructions of fields of transseries are due to Dahn and Göring [12] and Écalle [18]. More general constructions were proposed subsequently by the second author and his former student Schmeling [24, 25, 29]. Clearly, any *L*-function is a transseries, but the class of transseries is also closed under integration and functional inversion, contrary to the class of L-functions.

However, the class of transseries still does not contain any hyperexponential or hyperlogarithmic elements like  $E_{\omega} x$  or  $L_{\omega} x$ . In our quest for a truly universal framework for asymptotic analysis, we are thus lead to look beyond: a *hyperseries* is a formal object that is constructed from  $x$  and the real numbers using exponentiation, logarithms, infinite sums, as well as hyperexponentials  $E_{\omega}$ <sup> $\alpha$ </sup> and hyperlogarithms  $L_{\omega}$ <sup> $\alpha$ </sup> of any force  $\alpha$ . The hyperexponentials  $E_{\omega}$ <sup> $\alpha$ </sup> and the hyperlogarithms  $L_{\omega}$ <sup>*a*</sup> are required to satisfy functional equations

$$
E_{\omega^{\alpha+1}} \circ T_1 = E_{\omega^{\alpha}} \circ E_{\omega^{\alpha+1}} \tag{1.1}
$$

$$
L_{\omega^{\alpha+1}} \circ L_{\omega^{\alpha}} = T_{-1} \circ L_{\omega^{\alpha+1}}, \tag{1.2}
$$

where  $T_s(u) = u + s$ . For  $\gamma = \sum_{i=1}^p \omega^{\alpha_i} n_i$  in Cantor normal form with  $\alpha_1 < \cdots < \alpha_p$ , we also define

$$
L_{\gamma} = L_{\omega^{\alpha_1}}^{\circ n_1} \circ \cdots \circ L_{\omega^{\alpha_p}}^{\circ n_p} \tag{1.3}
$$

and we require that

<span id="page-3-0"></span>
$$
L'_{Y} = \frac{1}{\prod_{\beta < Y} L_{\beta}}.\tag{1.4}
$$

It is non-trivial to construct fields of hyperseries in which these and several other technical properties (see section 4 below) are satisfied. This was first accomplished by Schmeling for hyperexponentials  $E_{\omega^n}$  and hyperlogarithms  $L_{\omega^n}$  of finite force  $n \in \mathbb{N}$ . The general case was tackled in  $[14, 5]$ .

The construction of general hyperseries relies on the definition of an abstract notion of  $hyp$ erserial fields. Whereas the hyperseries that we are really after should actually be hyperseries in an infinitely large variable  $x$ , abstract hyperserial fields potentially contain hyperseries that can not be written as infinite expressions in  $x$ . In the present paper, we define hyperexponentials  $E_{\omega}$  and hyperlogarithms  $L_{\omega}$  on **No** for all ordinals  $\alpha$  and show that this provides **No** with the structure of an abstract hyperserial field. Moreover, any hyperseries  $f$  in  $x$  can naturally be evaluated at  $x = \omega$  to produce a surreal number  $f(\omega)$ . The conjecture from [26, p. 16] states that, for a sufficiently general notion of "hyperseries in  $x$ ", all surreal numbers can actually be obtained in this way. We plan to prove this and the conjecture in a follow-up paper.

#### General overview and summary of our new contributions  $1.2$

Our main goal is to define hyperexponentials  $E_{\alpha} \propto \mathbf{N_0}^{3.5} \rightarrow \mathbf{N_0}^{3.5}$  for any ordinal  $\alpha > 1$  and to show that No is a hyperserial field for these hyperexponentials. Since our construction builds on quite some previous work, the paper starts with three sections of reminders.

In section 2, we recall basic facts about well-based series and surreal numbers. In particular, we recall that any surreal number  $x \in \mathbb{N}$  can be regarded as a well-based series

$$
x = \sum_{\mathfrak{m} \in \mathbf{Mo}} x_{\mathfrak{m}} \mathfrak{m}
$$

with real coefficients  $x_m \in \mathbb{R}$ . The corresponding group of monomials **Mo** consists of those positive numbers  $m \in \mathbb{N} \circ \mathbb{N}$  that are of the form  $m = \{ \mathbb{R}^{\geq} L \mid \mathbb{R}^{\geq} R \}$  for certain subsets L and R of **No** with  $\mathbb{R}^{\geq} L \leq \mathbb{R}^{\geq} R$ .

<span id="page-4-3"></span>Section 3 is devoted to the theory of surreal substructures from [4]. One distinctive feature of the class of surreal numbers is that it comes with a partial, well-founded order  $\sqsubseteq$ , which is called the *simplicity* relation. The Conway bracket can then be characterized by the fact that, for any sets L and R of surreal numbers with  $L < R$ , there exists a unique  $\sqsubseteq$ -minimal number  $\{L | R\}$ with  $L < \{L | R\} < R$ . For many interesting subclasses **S** of **No**, it turns out that the restrictions of  $\leq$  and  $\subseteq$  to **S** give rise to a structure  $(S, \leq, \subseteq)$  that is isomorphic to  $(No, \leq, \subseteq)$ . Such classes **S** are called *surreal substructures* of **No** and they come with their own Conway bracket  $\{ | \}_{S}$ .

In section 4, we recall the definition of hyperserial fields from  $[5]$  and the main results on how to construct such fields. One major fact from [5] on which we heavily rely is that the construction of hyperserial fields can be reduced to the construction of hyperserial skeletons. In the context of the present paper, this means that it suffices to define the hyperlogarithms  $L_{\omega^{\alpha}}$ only for very special, so called  $L_{\leq \omega}$ *a*-atomic elements.

In the case when  $\alpha = 0$ , the L<sub> $\lt 1$ </sub>-atomic elements are simply the monomials in **Mo** and the definition of the general logarithm on  $No<sup>></sup>$  indeed reduces to the definition of the logarithm on **Mo**: given  $x \in \mathbb{N}$ **o**<sup>></sup>, we write  $x = c \cdot m(1 + \varepsilon)$ , where  $c \in \mathbb{R}$ ,  $m \in \mathbb{M}$ **o** and  $\varepsilon$  is infinitesimal, and we take  $\log x = \log m + \log c + \epsilon - \epsilon^2/2 + \epsilon^3/3 + \cdots$ . This very special case will be considered in more detail in section 5.

In the case when  $\alpha = 1$ , the  $L_{\leq \omega}$ -atomic elements of **No**<sup>>,></sup> are those elements  $\alpha \in$ **No**<sup>>,></sup> such that  $L_n$  a is a monomial for every  $n \in \mathbb{N}$ . The construction of  $L_{\omega}$  on **No**<sup>>,></sup> then reduces to the construction of  $L_{\omega}$  on the class  $M\sigma_{\omega}$  of  $L_{<\omega}$ -atomic numbers. This particular case was first dealt with in [6] and this paper can be used as an introduction to the more general results in the present paper.

For general ordinals  $\alpha$ , we say that  $\alpha \in \mathbf{No}^{>,\rangle}$  is  $L_{< \omega}$  atomic if  $L_{\beta} \alpha$  is a monomial for every  $\beta < \alpha$ . The advantage of restricting ourselves to such numbers a when defining hyperlogarithms is that  $L_{\alpha}$  a only needs to verify few requirements with respect to the ordering. This makes it possible to define  $L_{\alpha}$  a using the fairly simple recursive formula

<span id="page-4-0"></span>
$$
L_{\alpha} \mathfrak{a} \; := \; \{ \mathbb{R}, L_{\alpha} \mathfrak{a}' + (L_{<\alpha} \mathfrak{a}')^{-1} \mid L_{\alpha} \mathfrak{a}'' - (L_{<\alpha} \mathfrak{a})^{-1}, L_{<\alpha} \mathfrak{a} \}, \tag{1.5}
$$

where  $\alpha'$ ,  $\alpha''$  range over  $L_{\leq \alpha}$ -atomic numbers with  $\alpha'$ ,  $\alpha'' \sqsubseteq \alpha$  and  $\alpha' < \alpha < \alpha''$ ; see also (7.1).

In section 6, we prove that this definition is warranted and that the resulting functions  $L_{\alpha}$ satisfy the axioms of hyperserial skeletons from [5, Section 3]. Our proof proceeds by induction on  $\alpha$  and also relies on the fact that the class  $Mo_{\alpha}$  of  $L_{\leq \omega}$  atomic numbers actually forms a surreal substructure of No. Our main result is the following theorem:

<span id="page-4-2"></span>THEOREM 1.1. The definition (1.5) gives No the structure of a confluent hyperserial skeleton in the sense of [5]. Consequently, we may uniquely extend  $L_{\alpha\mu}$  to  $\text{No}^{>,>}$  in a way that gives No the structure of a confluent hyperserial field. Moreover, for each ordinal  $\mu$ , the extended function  $L_{\omega}$ ": **No**>>>  $\rightarrow$  **No**>>> is bijective.

Our final section  $7$  is devoted to further identities that illustrate the interplay between the hyperexponential and hyperlogarithmic functions and the simplicity relation  $\subseteq$  on No. We also prove the following more symmetric variant of (1.5):

<span id="page-4-1"></span>
$$
L_{\alpha} \mathfrak{a} = \{ \mathbb{R}, L_{\alpha} \mathfrak{a}' + (L_{<\alpha} \mathfrak{a}')^{-1} \mid L_{\alpha} \mathfrak{a}'' - (L_{<\alpha} \mathfrak{a}'')^{-1}, L_{<\alpha} \mathfrak{a} \},
$$
(1.6)

where  $\alpha'$ ,  $\alpha''$  again range over the  $L_{\leq \alpha}$ -atomic numbers with  $\alpha'$ ,  $\alpha'' \sqsubseteq \alpha$  and  $\alpha' < \alpha < \alpha''$ . An interesting open question is whether there exists an easy argument that would allow us to use (1.6) instead of (1.5) as a definition of  $L_{\alpha}$  a.

#### $\overline{2}$ **Basic notions**

#### Ordered fields of well-based series  $2.1$

#### 2.1.1 Well-based series

Let  $(\mathfrak{M}, \times, 1, \times)$  be a (possibly class-sized) linearly ordered abelian group. We write  $\mathbb{S} = \mathbb{R}[[\mathfrak{M}]]$ for the class of functions  $f: \mathfrak{M} \to \mathbb{R}$  whose support

<span id="page-5-3"></span><span id="page-5-2"></span><span id="page-5-1"></span><span id="page-5-0"></span>
$$
\operatorname{supp} f := \{ \mathfrak{m} \in \mathfrak{M} : f(\mathfrak{m}) \neq 0 \}
$$

is a well-based set, i.e. a set which is well-ordered with respect to the reverse order  $(\mathfrak{M},\geq)$ .

We see elements f of S as formal well-based series  $f = \sum_{m} f_m m$ , where  $f_m$  denotes the coefficient  $f(m) \in \mathbb{R}$  of m in f, for each  $m \in \mathfrak{M}$ . If supp  $f \neq \emptyset$ , then we define  $\delta_f := \max \text{supp } f \in \mathfrak{M}$  to be the *dominant monomial* of f. For  $m \in \mathfrak{M}$ , we let  $f_{\geq m} := \sum_{n > m} f_n n$  and we write  $f_{\geq m} = f_{\geq 1}$ . We say that a series  $g \in \mathbb{S}$  is a *truncation* of f and we write  $g \triangleleft f$  if supp $(f - g) > g$ . The relation  $\triangleleft$ is a well-founded partial order on S with minimum 0.

By [20], the class  $\mathbb S$  is an ordered field under the pointwise sum

<span id="page-5-6"></span><span id="page-5-5"></span><span id="page-5-4"></span>
$$
(f+g) := \sum_{\mathfrak{m}} (f_{\mathfrak{m}}+g_{\mathfrak{m}}) \mathfrak{m},
$$

the Cauchy product

$$
fg = \sum_{\mathfrak{m}} \left( \sum_{\mathfrak{u} \mathfrak{v} = \mathfrak{m}} f_{\mathfrak{u}} g_{\mathfrak{v}} \right) \mathfrak{m},
$$

(where each sum  $\sum_{\mu} f_{\mu} g_{\nu}$  has finite support), and where the positive cone  $\mathbb{S}^{\geq} = \{f \in \mathbb{S} : f > 0\}$ is given by

$$
\mathbb{S}^> := \{ f \in \mathbb{S} : f \neq 0 \land f_{\mathfrak{d}_f} > 0 \}
$$

The identification of  $m \in \mathfrak{M}$  with the formal series  $\sum_{n=m} 1 \cdot n \in \mathbb{S}$  induces an ordered group embedding  $(\mathfrak{M}, \times, \times) \rightarrow (\mathbb{S}^>, \times, \times).$ 

We next define the following asymptotic relations on S:

<span id="page-5-12"></span><span id="page-5-11"></span><span id="page-5-10"></span><span id="page-5-9"></span><span id="page-5-8"></span><span id="page-5-7"></span>
$$
f \prec g \iff \mathbb{R}^> |f| < |g|
$$
\n
$$
f \leq g \iff \exists r \in \mathbb{R}^> , |f| \leq r |g|
$$
\n
$$
f \prec g \iff f \leq g \leq f.
$$

The relation  $\prec$  extends the ordering on  $\mathfrak{M}$ . For non-zero  $f, g \in \mathbb{S}$  we actually have  $f \prec g$  (resp.  $f \preccurlyeq g$ , resp.  $f \preccurlyeq g$ ) if and only if  $\delta_f \preccurlyeq \delta_g$  (resp.  $\delta_f \preccurlyeq \delta_g$ , resp.  $\delta_f = \delta_g$ ). We finally define

$$
S_{>} \coloneqq \{f \in \mathbb{S} : \text{supp} f \subseteq \mathbb{M}^{\geq}\}
$$
  

$$
\mathbb{S}^{\leq} \coloneqq \{f \in \mathbb{S} : \text{supp} f \subseteq \mathbb{M}^{\leq}\} = \{f \in \mathbb{S} : f \prec 1\}
$$
  

$$
\mathbb{S}^{>,>} \coloneqq \{f \in \mathbb{S} : f > \mathbb{R}\} = \{f \in \mathbb{S} : f \geq 0 \land f > 1\}.
$$

Series in  $\mathbb{S}_{>}$  ,  $\mathbb{S}^<$  and  $\mathbb{S}^{>,>}$  are respectively called  $purely$  large, infinitesimal, and positive infinite. 2.1.2 Well-based families

Let  $(f_i)_{i \in I}$  be a family in S, We say that  $(f_i)_{i \in I}$  is well-based if

- i.  $\bigcup_{i \in I}$ supp  $f_i$  is well-based, and
- ii.  $\{i \in I : \mathfrak{m} \in \text{supp } f_i\}$  is finite for all  $\mathfrak{m} \in \mathfrak{M}$ .

In that case, we may define the sum  $\sum_{i \in I} f_i$  of  $(f_i)_{i \in I}$  by

$$
\sum_{i\in I} f_i := \sum_{\mathfrak{m}} \left( \sum_{i\in I} (f_i)_{\mathfrak{m}} \right) \mathfrak{m}.
$$

If  $\mathbb{U} = \mathbb{R}[\lceil \mathfrak{N} \rceil]$  is another field of well-based series and  $\Psi: \mathbb{S} \to \mathbb{U}$  is  $\mathbb{R}$ -linear, then we say that  $\Psi$ is strongly linear if for every well-based family  $(f_i)_{i \in I}$  in S, the family  $(\Psi(f_i))_{i \in I}$  is well-based, with

$$
\Psi\bigg(\sum_{i\in I}f_i\bigg) = \sum_{i\in I}\Psi(f_i).
$$

### 2.2 Surreal numbers

#### 2.2.1 Surreal numbers and simplicity

We denote by **On** the class of ordinal numbers. Following [19], we define **No** to be the class of sign sequences

<span id="page-6-2"></span><span id="page-6-1"></span>
$$
a = (a[\beta])_{\beta < \alpha} \in \{-1, +1\}^{\alpha}
$$

of ordinal length  $\alpha \in \mathbf{On}$ . The terms  $a[\beta] \in \{-1, +1\}$  are called the signs of a and we write  $l_a$  for the length of a. Given two numbers  $a, b \in$  **No**, we define

$$
a \sqsubseteq b \iff l_a \leq l_b \land (\forall \beta < l_a, a[\beta] = b[\beta]).
$$

We call  $\subseteq$  the *simplicity relation* on **No** and note that (**No**,  $\subseteq$ ) is well-founded. See [4, Section 2] for more details about the interaction between  $\sqsubseteq$  and the ordered field structure of No.

Recall that the Conway bracket is characterized by the fact that, for any sets L and R of surreal numbers with  $L < R$ , there exists a unique  $\subseteq$ -minimal number  $\{L | R\}$  with  $L < \{L | R\} < R$ . Conversely, given a number  $a \in \mathbf{No}$ , we define

$$
a_L := \{x \in \mathbf{No} : x \sqsubset a, x < a\}
$$
\n
$$
a_R := \{x \in \mathbf{No} : x \sqsupset a, x > a\}.
$$

Then *a* can canonically be written as

<span id="page-6-0"></span> $a = \{a_I \mid a_R\}.$ 

### 2.2.2 Ordinals as surreal numbers

The structure (No,  $\subseteq$ ) contains an isomorphic copy of (On, <) by identifying each ordinal  $\alpha$  with the constant sequence  $(+1)_{\beta<\alpha}$  of length  $\alpha$ . We will write  $v \le 0$ n to state that v is either an ordinal or the class of ordinals.

For  $\gamma \in \mathbf{On}$ , we write  $\omega^{\gamma}$  for the ordinal exponentiation of  $\omega$  to the power  $\gamma$  and we define

<span id="page-6-5"></span><span id="page-6-4"></span><span id="page-6-3"></span>
$$
\omega^{\mathbf{On}} \coloneqq \{\omega^{\gamma} : \gamma \in \mathbf{On}\}.
$$

If  $\mu \in \mathbf{On}$  is a successor ordinal, then we define  $\mu$  to be the unique ordinal with  $\mu = \mu$  + 1. We also define  $\mu_{-} := \mu$  if  $\mu$  is a limit ordinal. Similarly, if  $\alpha = \omega^{\mu}$ , then we set  $\alpha_{/\omega} := \omega^{\mu_{-}}$ . Recall that every ordinal  $\gamma$  has a unique Cantor normal form

$$
y = \omega^{\eta_1} n_1 + \cdots + \omega^{\eta_r} n_r,
$$

where  $r \in \mathbb{N}$ ,  $n_1, ..., n_r \in \mathbb{N}^{>0}$  and  $\eta_1, ..., \eta_r \in \mathbf{On}$  with  $\eta_1 > \cdots > \eta_r$ .

### 2.2.3 Surreal numbers as well-based series

We define **Mo** to be the class of positive numbers  $m \in \mathbb{N}$  of the form  $m = \{ \mathbb{R}^{\geq} L \mid \mathbb{R}^{\geq} R \}$  for certain subsets L and R of No with  $\mathbb{R}^{\geq} L \lt \mathbb{R}^{\geq} R$ . Numbers in Mo are called *monomials*. It turns out [11, Theorem 21] that the monomials form a subgroup of  $(No^>, \times, \lt)$  and that there is a natural isomorphism between **No** and the ordered field  $\mathbb{R}[\text{[Mo]}]$ . We will identify those two fields and thus see No as a field of well-based series. The ordinal  $\omega$ , seen as a surreal number, is the simplest element, or  $\sqsubseteq$ -minimum, of the class  $\text{No}^{>,\succ}$ .

#### 3 **Surreal substructures**

#### $3.1$ **Surreal substructures**

In [4], we introduced the notion of *surreal substructures*. A surreal substructure is a subclass  $S$ of **No** such that  $(No, \leq, \subseteq)$  and  $(S, \leq, \subseteq)$  are isomorphic. The isomorphism  $No \rightarrow S$  is unique and denoted by  $\Xi$ <sub>S</sub>. Many important subclasses of **No** that are relevant to the study of hyperserial properties of No are surreal substructures. In particular, it is known that the following classes are surreal substructures:

- <span id="page-7-0"></span>• The classes  $\text{No}^>, \text{No}^{>,}$  and  $\text{No}^<$  of positive, positive infinite and infinitesimal numbers.
- The classes  $\text{Mo}$  and  $\text{Mo}^>$  of monomials and infinite monomials.
- The classes  $N\omega$  and  $N\omega$  of purely infinite and positive purely infinite numbers.
- The class  $Mo_{\omega}$  of log-atomic numbers.

If U, V are surreal substructures, then the class  $U \lt V = \Xi_U V$  is a surreal substructure with  $\Xi_{\mathbf{U}\prec\mathbf{V}}=\Xi_{\mathbf{U}}\circ\Xi_{\mathbf{V}}$ .

### 3.2 Cuts

Given a subclass **X** of **No** and  $a \in \mathbf{X}$ , we will write

<span id="page-7-2"></span><span id="page-7-1"></span>
$$
a_L^{\mathbf{X}} \coloneqq \{b \in \mathbf{X} : b < a \land b \sqsubseteq a\} \text{ and } a_R^{\mathbf{X}} \coloneqq \{b \in \mathbf{X} : b > a \land b \sqsubseteq a\},
$$

so that  $a_L = a_L^{\mathbf{N}\mathbf{o}}$  and  $a_R = a_R^{\mathbf{N}\mathbf{o}}$ . We also write  $a_L^{\mathbf{X}} = a_L^{\mathbf{X}} \cup a_R^{\mathbf{X}}$  and  $a_L = a_L^{\mathbf{N}\mathbf{o}}$ .

If **X** is a subclass of **No** and *L*, *R* are subsets of **X** with  $L < S$ , then the class

$$
(L \mid R)_{\mathbf{X}} := \{ a \in \mathbf{X} : (\forall l \in L, l < a) \land (\forall r \in R, a < r) \}
$$

is called a *cut* in **X**. If  $(L | R)$ **x** contains a unique simplest element, then we denote this element by  $\{L \mid R\}$  and say that  $(L, R)$  is a *cut representation* (of  $\{L \mid R\}$ ) in **X**. These notations naturally extend to the case when **L** and **R** are subclasses of **X** with  $L < R$ .

A surreal substructure S may be characterized as a subclass of No such that for all cut representations (L, R) in S, the cut  $(L | R)$ s has a unique simplest element [4, Proposition 4.7].

Let S be a surreal substructure. Note that we have  $a = \{a_L^S \mid a_R^S\}$  for all  $a \in S$ . Let  $a \in S$  and let  $(L, R)$  be a cut representation of a in **S**. Then  $(L, R)$  is cofinal with respect to  $(a_L^S, a_R^S)$  in the sense that L has no strict upper bound in  $a_L^S$  and R has no strict lower bound in  $a_R^S$  [4, Proposition  $4.11(b)$ ].

Given numbers  $a, b \in \mathbb{N}$ o with  $a \leq b$ , the number  $c = \{a_L | b_R\}$  is the unique  $\sqsubseteq$ -maximal number with  $c \sqsubseteq a, b$ . We have  $a \leq c \leq b$ . Let **S** be a surreal substructure. Considering the isomorphism  $\Xi_{\mathbf{S}}: (\mathbf{No}, \leqslant, \sqsubseteq) \to (\mathbf{S}, \leqslant, \sqsubseteq)$ , we see that for all  $a, b \in \mathbf{S}$  with  $a \leqslant b$ , there is a unique  $\sqsubseteq$ -maximal element c of **S** with  $c \subseteq a$ , b, and we have  $a \leq c \leq b$ . In what follows, we will use this basic fact several times without further mention.

#### 3.3 Cut equations

Let  $X \subseteq No$  be a subclass, let T be a surreal substructure and  $F: X \rightarrow T$  be a function. Let  $\lambda, \rho$ be functions defined for cut representations in **X** and such that  $\lambda(L, R)$ ,  $\rho(L, R)$  are subsets of **T** whenever  $(L, R)$  is a cut representation in **X**. We say that  $(\lambda, \rho)$  is a *cut equation* for F if for all  $a \in X$ , we have

 $\lambda(a_1^X, a_2^X) < \rho(a_1^X, a_2^X)$ ,  $F(a) = \lambda(a_1^X, a_2^X) | \rho(a_1^X, a_2^X) |$ 

Elements in  $\lambda(a_L^X, a_R^X)$  (resp.  $\rho(a_L^X, a_R^X)$ ) are called *left* (resp. *right*) *options* of this cut equation at  $a$ .

We say that the cut equation is *uniform* if we have

$$
\lambda(L,R) < \rho(L,R), \quad F(\lbrace L \mid R \rbrace_{\mathbf{X}}) = \lbrace \lambda(L,R) \mid \rho(L,R) \rbrace_{\mathbf{T}}
$$

whenever  $(L, R)$  is a cut representation in **X**. For instance, given  $r \in \mathbb{R}$ , consider the translation  $T_r$ : **No**  $\rightarrow$  **No**;  $a \mapsto a + r$  on **No**. By [\[19](#page-31-1), Theorem 3.2], we have the following uniform cut equation for  $T_r$  on **No**:

<span id="page-8-3"></span><span id="page-8-1"></span>
$$
\forall a \in \mathbf{No}, \quad a + r = \{a_L + r, a + r_L \mid a + r_R, a_R + r\}.
$$
 (3.1)

We will need the following result from [\[4\]](#page-30-4):

*Proposition* 3.1*.* [\[4](#page-30-4), Proposition 4.36] *Let* , *be surreal substructures. Let* Λ *be a function* from S to the class of subsets of T such that for  $x, y \in S$  with  $x < y$ , the set  $\Lambda(y)$  is cofinal with respect to  $\Lambda(x)$ . For  $x \in S$ , let  $\Lambda[x]$  denote the class of elements u of S such that  $\Lambda(x)$  and  $\Lambda(u)$  are *mutually cofinal.* Let  $\{\lambda \mid \rho\}$  *p be a cut equation on* **S** *that is extensive in the sense that* 

$$
\forall x, y \in S, \quad (x \sqsubseteq y \Longrightarrow (\lambda(x_L^S, x_R^S) \subseteq \lambda(y_L^S, y_R^S) \land \rho(x_L^S, x_R^S) \subseteq \rho(y_L^S, y_R^S))).
$$

*Let*  $F: S \to T$  *be strictly increasing with cut equation* 

<span id="page-8-5"></span><span id="page-8-4"></span> $\forall x \in \mathbf{S}, \quad F(x) = {\Lambda(x), \lambda(x_L^{\mathbf{S}}, x_R^{\mathbf{S}}) | \rho(x_L^{\mathbf{S}}, x_R^{\mathbf{S}})}$ .

*Then F induces an embedding*  $(\Lambda[x], \leq, \subseteq) \rightarrow (\mathbf{T}, \leq, \subseteq)$  *for each element x of* **S**.

### 3.4 Convex partitions

One natural way to obtain surreal substructures is *via* convex partitions. If **S** is a surreal substructure, then a*convex partition* of is a partition Π of whose members are convex subclasses of **S** for the order  $\leq$ . We may then consider the class  $\text{Smp}_{\Pi}$  of simplest elements (i.e. ⊑-minima) in each member of  $\Pi$ . Those elements are said  $\Pi$ -simple. For  $a \in S$ , we let  $\Pi[a]$  denote the unique member of  $\Pi$  containing *a*. By [\[4](#page-30-4), Proposition 4.16], the class  $\Pi[a]$  contains a unique Π-simple element, which we denote by *π*Π(*a*). The function *π*Π is a surjective non-decreasing function  $S \rightarrow$  Smp<sub>II</sub> with  $\pi_{\Pi} \circ \pi_{\Pi} = \pi_{\Pi}$ .<br>Given *a*, *b*∈Smp<sub>II</sub>, note that we have *a* < *b* if and only if  $\Pi[a] < \Pi[b]$ . For **X** ⊆ **No**, we write

 $\Pi[X] = \bigcup_{a \in X} \Pi[a]$ . We have the following criterion to characterize elements of  $Smp_{\Pi}$ .

<span id="page-8-2"></span>*Proposition* 3.2*.* [\[4](#page-30-4), Lemma 6.5] *An element a of is* Π*-simple if and only if there is a cut representation* (*L*, *R*) *of a in* **S** *with*  $\Pi[L] < a < \Pi[R]$ *. Equivalently*  $a \in S$  *is*  $\Pi$ *-simple if and only*  $if \Pi[a_L^S] < a < \Pi[a_R^S].$ 

We say that Π is *thin* if each member of Π has acofinal and coinitial subset. We then have:

*Proposition* 3.3*.* [\[4](#page-30-4), Theorem 6.7 and Proposition 6.8] *If* Π *is thin, then the class* Smp<sup>Π</sup> *is a* surreal substructure and  $\Xi_{Smp_{\Pi}}$  has the following uniform cut equation:

<span id="page-8-0"></span>
$$
\forall z \in \mathbf{No}, \quad \Xi_{\mathbf{Smp}_{\Pi}} z = {\Pi[\Xi_{\mathbf{Smp}_{\Pi}} z_L] \mid \Pi[\Xi_{\mathbf{Smp}_{\Pi}} z_R] }_{\mathbf{S}}.
$$

#### 3.5 Function groups

A special type of thin convex partitions is that of partitions induced by function groups acting on surreal substructures. A *function group*  $\mathcal G$  on a surreal substructure **S** is a set-sized group of strictly increasing bijections  $S \rightarrow S$  under functional composition. We see elements  $f, g$  of  $\mathcal G$  as actions on **S** and we sometimes write *fg* and *fa* instead of  $f \circ g$  and  $f(a)$ , where  $a \in S$ .

For such a function group  $\mathcal{G}$ , the collection  $\Pi \mathcal{G}$  of classes

<span id="page-8-6"></span>
$$
\mathcal{G}[a] \coloneqq \{b \in \mathbf{S} : \exists f, g \in \mathcal{G}, fa \leq b \leq ga\}
$$

with  $a\!\in\!{\bf S}$  is a thin convex partition of  ${\bf S}.$  We write  ${\bf Smp}_\mathcal{G} \coloneqq {\bf Smp}_{\Pi_\mathcal{G}}.$  We have the uniform cut equation

$$
\forall z \in \mathbf{No}, \quad \Xi_{\mathbf{Smp}_{\mathcal{G}}} z = \{ \mathcal{G} \Xi_{\mathbf{Smp}_{\mathcal{G}}} z_L \mid \mathcal{G} \Xi_{\mathbf{Smp}_{\mathcal{G}}} z_R \} \mathbf{S}. \tag{3.2}
$$

Consider sets *X*, *Y* of strictly increasing bijections  $S \rightarrow S$ , then we say that *Y* is *pointwise cofinal* with respect to *X*, and we write *X*  $\angle$  *Y*, if we have ∀*f* ∈ *X*, ∀*a* ∈ **S**, ∃*g* ∈ *Y*, *fa* ≤ *ga*. We also define

<span id="page-9-3"></span><span id="page-9-2"></span><span id="page-9-1"></span>
$$
\langle X \rangle \; := \; \{f_0 \circ f_1 \circ \cdots \circ f_n : n \in \mathbb{N}, f_0, \ldots, f_n \in X \cup X^{-1}\}.
$$

It is easy to see that  $\langle X \rangle$  is a function group on **S** and that we have  $\langle X \rangle \leq \langle Y \rangle$  if  $X \leq Y$  or *X*<sup>-1</sup> ∠ *Y*<sup>-1</sup>. The relation  $\langle X \rangle$  ∠  $\langle Y \rangle$  trivially implies  $\text{Smp}_{\langle Y \rangle}$ ⊆  $\text{Smp}_{\langle X \rangle}$ . If  $X \leq Y$  and  $Y \leq X$ , then we say that *X* and *Y* are *mutually pointwise cofinal* and we write  $X \neq Y$ . We then have  $\textbf{Smp}_{\langle X \rangle} = \textbf{Smp}_{\langle Y \rangle}.$ 

We write  $X \le Y$  (resp.  $X \le Y$ ) if we have  $\forall a \in S$ ,  $\forall f \in X$ ,  $\forall g \in Y$ ,  $fa \le ga$  (resp.  $\forall a \in S$ ,  $\forall f \in X$ , *∀g* ∈ *Y*, *fa* < *ga*). We also write *f* < *Y* and *X* < *g* instead of {*f* } < *Y* and *X* < {*g* }.

Given a function group  $\mathcal G$  on S, the relation defined by  $f < g \Leftrightarrow \{f\} < \{g\}$  is a partial order on  $\mathcal G$ . We will frequently rely on the basic fact that  $(\mathcal G, <)$  is *partially bi-ordered* in the sense that

<span id="page-9-4"></span>
$$
\forall f, g, h \in \mathcal{G}, \quad \text{id}_S < g \Longleftrightarrow fh < fgh.
$$

#### 3.6 Remarkable function groups

Each of the examples of surreal substructures from Subsection [3.1](#page-7-1) can be regarded as the classes  $\textbf{Smp}_{\mathscr{G}}$  for actions of the following function groups  $\mathscr{G}$  acting on **No**, **No** $\geq$  or **No** $\geq$ . $\geq$  For  $c \in \mathbb{R}$ and  $r \in \mathbb{R}^>$ , we define

<span id="page-9-7"></span><span id="page-9-6"></span><span id="page-9-5"></span>
$$
T_r := a \mapsto a + c \quad \text{acting on } \text{No or } \text{No}^{>,>}.
$$
  
\n
$$
H_c := a \mapsto ra \quad \text{acting on } \text{No}^{>} \text{ or } \text{No}^{>,>}.
$$
  
\n
$$
P_c := a \mapsto a^r \quad \text{acting on } \text{No}^{>} \text{ or } \text{No}^{>,>}.
$$

<span id="page-9-9"></span><span id="page-9-8"></span>Now consider

<span id="page-9-12"></span><span id="page-9-11"></span><span id="page-9-10"></span><span id="page-9-0"></span>
$$
\mathcal{T} := \{T_c : c \in \mathbb{R}\},
$$
  
\n
$$
\mathcal{H} := \{H_r : r \in \mathbb{R}^>\},
$$
  
\n
$$
\mathcal{P} := \{P_r : r \in \mathbb{R}^>\},
$$
  
\n
$$
\mathcal{E}' := \langle E_n H_r L_n : n \in \mathbb{N}, r \in \mathbb{R}^>\rangle, \text{ and}
$$
  
\n
$$
\mathcal{E}^* := \{E_n, L_n : n \in \mathbb{N}\}.
$$

Then we have the following list of correspondences  $\mathcal{G} \mapsto \text{Smp}_{\mathcal{C}}$ :

- The action of  $\mathcal T$  on No (resp. No<sup>>,></sup>) yields No<sub>></sub> (resp. No>), e.g.  $\text{Smp}_{\mathcal T} = \text{No}_{>}.$
- The action of  $\mathcal{H}$  on No<sup>></sup> (resp. No<sup>>,></sup>) yields Mo (resp. Mo<sup>></sup>).
- The action of  $\mathcal P$  on **No**<sup>></sup>>> yields **Mo**  $\prec$ **Mo** =  $E_1$ **Mo**  $\succeq$ .
- The action of <sup>*%'*</sup> on **No**<sup>>,≻</sup> yields **Mo**<sub>ω</sub>.
- The action of  $\mathscr{E}^*$  on  $\mathbf{No}^{>,\succ}$  yields  $\mathbf{K} := \mathbf{Mo}_{\omega} \prec \mathbf{No}_{\succ}$  (which will coincide with  $E_{\omega} \mathbf{No}_{\succ}$ ).

Generalizations of those function groups will allow us to define certain surreal substructures related to the hyperlogarithms and hyperexponentials on No.

### 4 Hyperserial fields

In this section, we briefly recall the definition of hyperserial fields from [\[5\]](#page-30-2) and how to construct such fields from their hyperserial skeletons.

### 4.1 Logarithmic hyperseries

Let x be a formal, infinitely large indeterminate. The field  $\mathbb{L}$  of *logarithmic hyperseries* of [\[14\]](#page-31-17) is the smallest field of well-based series that contains all ordinal real power products of the hyperlogarithms  $L_{\alpha}$ *x* with  $\alpha \in \mathbf{On}$ . It is naturally equipped with a derivation ∂:  $\mathbb{L} \to \mathbb{L}$  and composition law ∘:  $\mathbb{L} \times \mathbb{L}^{>,} \rightarrow \mathbb{L}$ .

*Definition* Let  $\alpha$  be an ordinal. For each  $\gamma < \alpha$ , we introduce the formal hyperlogarithm  $\ell$ *γ* ≔ *Lγx* and define  $\mathfrak{L}_{<\alpha}$  to be the group of formal power products  $\mathfrak{l} = \prod_{\gamma<\alpha} \ell_{\gamma}^{\mathfrak{l}_{\gamma}}$  with  $\mathfrak{l}_{\gamma} \in \mathbb{R}$ . This group comes with a monomial ordering ≻ that is defined by

<span id="page-10-2"></span><span id="page-10-1"></span>
$$
1 > 1 \Leftrightarrow l_{\beta} > 0 \quad \text{for } \beta = \min\{ \gamma < \alpha : l_{\gamma} \neq 0 \}.
$$

We define  $\mathbb{L}_{<\alpha}$  to be the ordered field of well-based series  $\mathbb{L}_{<\alpha} := \mathbb{R}[[\mathfrak{L}_{<\alpha}]]$ . If  $\alpha, \beta$  are ordinals with  $\beta < \alpha$ , then we define  $\mathfrak{L}_{\beta,\alpha}$  to be the subgroup of  $\mathfrak{L}_{<\alpha}$  of monomials  $\beta$  with  $\beta$ <sub>r</sub> = 0 whenever  $\gamma < \beta$ . As in [\[14\]](#page-31-17), we write

<span id="page-10-3"></span>
$$
\mathbb{L}_{[\beta,\alpha)} := \mathbb{R}[[\mathfrak{L}_{[\beta,\alpha)}]]
$$
  

$$
\mathfrak{L} := \bigcup_{\alpha \in \mathbf{On}} \mathfrak{L}_{<\alpha}
$$
  

$$
\mathbb{L} := \mathbb{R}[[\mathfrak{L}]].
$$

We have natural inclusions  $\mathcal{L}_{\lceil \beta,\alpha \rceil} \subseteq \mathcal{L}_{\leq \alpha} \subset \mathcal{L}$ , hence natural inclusions  $\mathbb{L}_{\lceil \beta,\alpha \rceil} \subseteq \mathbb{L}_{\leq \alpha} \subset \mathbb{L}$ .

*Derivation* on  $\mathbb{L}_{< \alpha}$  The field  $\mathbb{L}_{< \alpha}$  is equipped with a derivation  $\partial: \mathbb{L}_{< \alpha} \to \mathbb{L}_{< \alpha}$  which satisfies the Leibniz rule and which is strongly linear. Write  $\ell^{\dagger}_\gamma := \prod_{\iota \leq \gamma} \ell^{-1}_\iota \in \mathfrak{L}_{<\alpha}$  for all  $\gamma < \alpha$ . The derivative of a logarithmic hypermonomial  $I \in \mathcal{L}_{\leq \alpha}$  is defined by

$$
\partial I := \left( \sum_{\gamma < \alpha} I_{\gamma} \ell_{\gamma}^{\dagger} \right) I.
$$

So  $\partial \ell_{\gamma} = \frac{1}{\prod_{i \leq \gamma} \ell_i}$  for all  $\gamma < \alpha$ . For  $f \in \mathbb{L}_{<\alpha}$  and  $k \in \mathbb{N}$ , we will sometimes write  $f^{(k)} \coloneqq \partial^k f$ .

*Composition on*  $\mathbb{L}_{<\alpha}$  Assume that  $\alpha = \omega^{\nu}$  for a certain ordinal *ν*. Then the field  $\mathbb{L}_{<\alpha}$  is *Composition on*  $\mathbb{L}_{< \alpha}$  Assume that  $\alpha = \omega^{\nu}$  for a certain ordinal  $\nu$ . Then equipped with a composition ∘:  $\mathbb{L}_{< \alpha} \times \mathbb{L}_{< \alpha}^{> \succ} \rightarrow \mathbb{L}_{< \alpha}$  that satisfies in particular:

- For  $g \in \mathbb{L}_{<\alpha}^{>,>}$ , the map  $\mathbb{L}_{<\alpha} \to \mathbb{L}_{<\alpha}$ ;  $f \mapsto f \circ g$  is a strongly linear embedding [\[14](#page-31-17), Lemma 6.6].
- For  $f \in \mathbb{L}_{<\alpha}$  and  $g, h \in \mathbb{L}_{<\alpha}^{>}$ , we have  $g \circ h \in \mathbb{L}_{<\alpha}^{>}$  and  $f \circ (g \circ h) = (f \circ g) \circ h$  [\[14](#page-31-17), Proposition 7.14].
- For  $g \in \mathbb{L}_{<\alpha}^{>,>}$  and successor ordinals  $\mu < \nu$ , we have  $\ell_{\omega^{\mu}} \circ \ell_{\omega^{\mu}} = \ell_{\omega^{\mu}} 1$  [\[14](#page-31-17), Lemma 5.6].

The same properties hold for the composition ∘:  $\mathbb{L} \times \mathbb{L}^{\geq, >} \to \mathbb{L}$  if  $\alpha$  is replaced by **On**. For *γ* <  $\alpha$ , the map  $\mathbb{L}_{<\alpha} \to \mathbb{L}_{<\alpha}$ ;  $f \mapsto f \circ \ell_{\gamma}$  is injective, with image  $\mathbb{L}_{\{ \gamma, \alpha \}}$  [\[14](#page-31-17), Lemma 5.11]. For  $g \in \mathbb{L}_{\{ \gamma, \alpha \}}$ , we define  $g^{\uparrow}$  *γ* to be the unique series in  $\mathbb{L}_{< \alpha}$  with  $g^{\uparrow}$   $\gamma \circ \ell_{\gamma} = g$ .

### <span id="page-10-4"></span>4.2 Hyperserial fields

Let  $\mathfrak{M}$  be an ordered group. A *real powering operation* on  $\mathfrak{M}$  is a law

$$
\mathbb{R} \times \mathfrak{M} \longrightarrow \mathfrak{M}; (r, \mathfrak{m}) \longmapsto \mathfrak{m}^r
$$

of ordered ℝ-vector space on  $\mathfrak{M}$ . Let  $\mathbb{T} = \mathbb{R}[[\mathfrak{M}]]$  be a field of well-based series with  $\mathfrak{M} \neq 1$ , let *ν* ≤ **On**, and let ∘: L × T<sup>>,></sup> → T be a function. For  $\mu \le \nu$  , we define  $\mathfrak{M}_{\omega}$ <sup> $\mu$ </sup> to be the class of series *s*∈ $\mathbb{T}^{>,>}$  with ∀*γ* <  $ω$ <sup>*μ*</sup>,  $\ell$ <sub>*γ*</sub> ∘ *s*∈ $\mathfrak{M}^>$ . We say that ( $\mathbb{T}, \circ$ ) is a *hyperserial field* if

<span id="page-10-0"></span>HF1.  $\mathbb{L} \to \mathbb{T}$ ; *f*  $\mapsto f \circ s$  is a strongly linear morphism of ordered rings for each  $s \in \mathbb{T}^{>}$ .

**HF2.**  $f \circ (g \circ s) = (f \circ g) \circ s$  for all  $f \in \mathbb{L}, g \in \mathbb{L}^{>,}$ , and  $s \in \mathbb{T}^{>,}$ .

**HF3.** 
$$
f \circ (t + \delta) = \sum_{k \in \mathbb{N}} \frac{f^{(k)} \circ t}{k!} \delta^k
$$
 for all  $f \in \mathbb{L}$ ,  $t \in \mathbb{T}^{>,\ge}$ , and  $\delta \in \mathbb{T}$  with  $\delta \prec t$ .

**HF4.**  $\ell_{\alpha\mu}^{\gamma} \circ s < \ell_{\alpha\mu}^{\gamma} \circ t$  for all ordinals  $\mu, \gamma < \omega^{\mu}$ , and  $s, t \in \mathbb{T}^{>,\ge}$  with  $s < t$ .

**HF5.** The map  $\mathbb{R}^>\times\mathfrak{M}^>\to\mathfrak{M}$ ;  $(r,\mathfrak{m})\mapsto\mathfrak{m}^r:=\ell_0^r\circ\mathfrak{m}$  extends to a real powering operation on  $\mathfrak{M}$ .

**HF6.**  $\ell_1 \circ (st) = \ell_1 \circ s + \ell_1 \circ t$  for all  $s, t \in \mathbb{T}^{>,>}$ .

<span id="page-11-0"></span>**HF7.** supp  $\ell_1 \circ \mathfrak{m} > 1$  for all  $\mathfrak{m} \in \mathfrak{M}^>$ ; supp  $\ell_{\omega} \mu \circ \mathfrak{a} > (\ell_{\gamma} \circ \mathfrak{a})^{-1}$  for all  $1 \leq \mu < \nu$ ,  $\gamma < \omega^{\mu}$  and  $\mathfrak{a} \in \mathfrak{M}_{\omega} \mu$ .

For each  $\mu \in \mathbf{On}$ , we define the function  $L_{\omega} \mu : \mathfrak{M}_{\omega} \mu \to \mathbb{T}$ ;  $\mathfrak{a} \mapsto \ell_{\omega} \mu \circ \mathfrak{a}$ . The skeleton of  $(\mathbb{T}, \circ)$  is defined to be the structure  $(\mathbb{T}, (L_{\omega^{\mu}})_{\mu \in \mathbf{On}})$  equipped with the real power operation from **HF5**.

We say that  $(\mathbb{T}, \circ)$  is *confluent* if for all  $\mu \in \mathbf{On}$  with  $\mu \leq \nu$ , we have

<span id="page-11-3"></span> $\forall s \in \mathbb{T}^{>,\succ}, \exists a \in \mathfrak{M}_{\omega^{\mu}}, \exists \gamma < \omega^{\mu}, \quad \ell_{\gamma} \circ s \asymp \ell_{\gamma} \circ a.$ 

In particular  $(\mathbb{L}, \circ)$  is a confluent hyperserial field.

#### 4.3 Hyperserial skeletons

It turns out that each hyperlogarithm  $L_{\alpha^{\mu}}$  on a hyperserial field  $\mathbb T$  can uniquely be reconstructed from its restriction to the subset of  $L_{\leq \omega^{\mu}}$ -atomic hyperseries (here we say that  $f \in \mathbb{T}^{>,>}$  is  $L_{< \omega^{\mu}}$ -atomic if  $L_{\gamma} f \in \mathfrak{M}$  for all  $\gamma < \omega^{\mu}$ ). One of the main ideas behind [14] is to turn this fact into a way to *construct* hyperserial fields. This leads to the definition of a hyperserial skeleton as a field  $\mathbb T$  with partially defined hyperlogarithms  $L_{\alpha}$ <sup> $\mu$ </sup>, which satisfy suitable counterparts of the above axioms HF1 until HF7.

More precisely, let  $\mathbb{T} = \mathbb{R}[[\mathfrak{M}]]$  be a field of well-based series and fix  $v \in \mathbf{On}^> \cup \{\mathbf{On}\}\$ . A hyperserial skeleton on  $\mathbb T$  of force v consists of a family of partial functions  $L_{\alpha^{\mu}}$  for  $\mu < \nu$ , called (hyper)logarithms, which satisfy a list of axioms that we will describe now.

First of all, the domains  $\mathfrak{M}_{\omega^{\mu}}$ := dom  $L_{\omega^{\mu}}$  on which the partial functions  $L_{\omega^{\mu}}$  are defined should satisfy the following axioms:

Domains of definition:

<span id="page-11-4"></span><span id="page-11-1"></span> $DD_0$ . dom  $L_1 = \mathfrak{M}^>$ ; **DD<sub>µ</sub>**. dom  $L_{\omega^{\mu}} = \bigcap_{\eta \leq \mu}$  dom  $L_{\omega^{\eta}}$ , if  $\mu$  is a non-zero limit ordinal; **DD<sub>µ</sub>**. dom  $L_{\omega^{\mu}} = \{s \in \mathbb{T} : L_{\omega^{\mu}}^{\circ n}(s) \in \text{dom } L_{\omega^{\mu-}} \text{ for all } n\}$ , if  $\mu$  is a successor ordinal.

It will be convenient to also define the class  $\mathfrak{M}_{\omega}$ <sup>v</sup> by

$$
\mathfrak{M}_{\omega^{\nu}} := \{ s \in \mathbb{T} : L^{\circ n}_{\omega^{\nu}}(s) \in \mathfrak{M}_{\omega^{\nu}} \text{ for all } n \} \quad \text{if } \nu \text{ is a successor ordinal}
$$
\n
$$
\mathfrak{M}_{\omega^{\nu}} := \bigcap_{\mu < \nu} \mathfrak{M}_{\omega^{\mu}}
$$
\nif  $\nu$  is a non-zero limit ordinal.

Consider an ordinal  $\gamma < \omega^{\nu}$  written in Cantor normal form  $\gamma = \sum_{i=1}^{r} \omega^{\eta_i} n_i$  where  $\eta_1 > \eta_2 > \cdots > \eta_r$ and  $n_1, \ldots, n_r < \omega$ . We denote by  $L<sub>\gamma</sub>$  the partial function

<span id="page-11-2"></span>
$$
L_{\gamma} \coloneqq L_{\omega}^{\circ n_1} \circ \cdots \circ L_{\omega}^{\circ n_r}.\tag{4.1}
$$

It follows from the definition that for all  $\mu \leq \nu$ , the class  $\mathfrak{M}_{\omega}$  consists of those series  $s \in \mathbb{T}^{>,>}$ for which  $s \in \text{dom } L_y$  and  $L_y s \in \mathfrak{M}^>$  for all  $\gamma < \omega^\mu$ . We call such series  $L_{\leq \omega^\mu}$ -atomic.

Secondly, the hyperlogarithms  $L_{\omega^{\mu}}$  with  $\mu < v$  should satisfy the following axioms:

<span id="page-12-9"></span><span id="page-12-8"></span><span id="page-12-7"></span>

<span id="page-12-11"></span><span id="page-12-10"></span><span id="page-12-3"></span><span id="page-12-2"></span><span id="page-12-1"></span><span id="page-12-0"></span>Monotonicity:

<span id="page-12-12"></span><span id="page-12-4"></span> $\mathbf{M}_{\boldsymbol{\mu}}$ .  $\forall \mathfrak{a}, \mathfrak{b} \in \mathfrak{M}_{\beta}, \forall \gamma < \beta, \mathfrak{a} < \mathfrak{b} \Longrightarrow L_{\beta} \mathfrak{a} + (L_{\gamma} \mathfrak{a})^{-1} < L_{\beta} \mathfrak{b} - (L_{\gamma} \mathfrak{b})^{-1}.$ 

Regularity:

<span id="page-12-13"></span><span id="page-12-5"></span>
$$
\mathbf{R}_{\boldsymbol{\mu}} \cdot \forall \mathfrak{a} \in \mathfrak{M}_{\beta}, \forall \mathfrak{y} < \beta, \operatorname{supp} L_{\beta} \mathfrak{a} > (L_{\gamma} \mathfrak{a})^{-1}.
$$

<span id="page-12-14"></span>Finally, for  $\mu \le \nu$  with  $\mu \in \mathbf{On}$ , we also need the following axiom

<span id="page-12-6"></span>

Note that S[L](#page-12-0) and  $\mathbf{R}_0$  $\mathbf{R}_0$  $\mathbf{R}_0$  together imply  $L_1\,\mathfrak{M}^>=\mathbb{T}^>_{>},$  whence  $\mathbf{P}_\mu$  automatically holds. This will in particular be the case for No (see Section [5\)](#page-15-0).

In summary, we have:

 $D$ EFINITION 4.1. [\[5](#page-30-2), Definition 3.3]  $Given$   $\nu$   $\in$   $On$   $\ge$   $\cup$   $\{On\}$ *, we say that*  $(\mathbb{T},(L_{\omega^{\mu}})_{\mu < \nu})$  *is a*  $hyper$ serial skeleton of force v if it satisfies  $DD_{\mu}$  $DD_{\mu}$ ,  $FE_{\mu}$  $FE_{\mu}$ ,  $A_{\mu}$  $A_{\mu}$ ,  $M_{\mu}$  $M_{\mu}$ , and  $R_{\mu}$  $R_{\mu}$  for all  $\mu < \nu$ , as well as  $P_{\mu}$  $P_{\mu}$  for *all ordinals*  $\mu \leq \nu$ *.* 

Assume that  $\mathbb T$  is a hyperserial skeleton of force *ν*. The partial logarithm  $L_1: \mathfrak{M}_1 \to \mathbb T$  extends naturally into a strictly increasing morphism  $(\mathbb{T}^>, \times, <) \rightarrow (\mathbb{T}, +, <)$ , which we call the *logarithm* and denote by  $L_1$  $L_1$  or log [\[5](#page-30-2), Section 4.1]. If  $T$  satisfies SL, then this extended logarithm is actu-ally an isomorphism [\[29](#page-31-16), Proposition 2.3.8]. In that case, for any  $s \in \mathbb{T}^>$  and  $r \in \mathbb{R}$ , we define  $s^r$  = exp( $r \log s$ )  $\in \mathbb{T}^>$ .

#### 4.4 Confluence

 $D$ ΕFINITION 4.2. [\[5](#page-30-2), Definition 3.5] *Given a hyperserial skeleton*  $\mathbb{T} = \mathbb{R}[[\mathfrak{M}]]$  *of force v* ∈ **On**<sup>></sup> *and μ* <*ν, we inductively define the notion of μ-*confluence *in conjunction with the definition of functions*  $\mathfrak{d}_{\omega^{\mu}}: \mathbb{T}^{>,>} \rightarrow \mathfrak{M}_{\omega^{\mu}}$ *, as follows.* 

• *The field*  $\mathbb T$  *is said* 0*-confluent if*  $\mathfrak M$  *is non-trivial. The function*  $\mathfrak d_1$  *maps every positive infinite series*  $s \in \mathbb{T}^{\gt,>}$  *onto its dominant monomial*  $\delta_s$ *. For each*  $s \in \mathbb{T}^{\gt,>}$ *, we write* 

$$
\mathcal{E}_1[s] \; \coloneqq \; \{t \in \mathbb{T}^{>,\succ} : t \approx s\}.
$$

*Let*  $\mu \leq v$  *be such that*  $\mathbb{T}$  *is η*-confluent for all  $\eta < \mu$  and let  $s \in \mathbb{T}^{>,\geq}$ .

• *If μ is a successor ordinal, then we write* ℰ*<sup>ω</sup> <sup>μ</sup>*[*s*] *for the class of series t with*

$$
(L_{\omega^{\mu - \circ}} \delta_{\omega^{\mu -}})^{\circ n}(s) \; \asymp \; (L_{\omega^{\mu - \circ}} \delta_{\omega^{\mu -}})^{\circ n}(t)
$$

*for a certain*  $n \in \mathbb{N}$ *.* 

• *If μ is a limit ordinal, then we write* ℰ*<sup>ω</sup> <sup>μ</sup>*[*s*] *for the class of series t with*

<span id="page-13-4"></span><span id="page-13-1"></span>
$$
L_{\omega^{\eta}}\mathfrak{d}_{\omega^{\eta}}(s) \approx L_{\omega^{\eta}}\mathfrak{d}_{\omega^{\eta}}(t)
$$

<span id="page-13-5"></span>*for a certain*  $\eta < \mu$ .

We say that  $\mathbb T$  is **µ-confluent** if each class  $\mathscr E_{\omega^\mu}[s]$  contains a  $L_{< \omega^\mu}$ -atomic element; we then *define <sup>ω</sup> <sup>μ</sup>*(*s*) *to be this element.*

This inductive definition is sound. Indeed, if  $\mu \le \nu + 1$  and  $\mathbb T$  is *η*-confluent for all  $\eta \le \mu$ , then the functions  $\delta_{\omega}$ <sup>*n*</sup>:  $\mathbb{T}^{>,>}$   $\rightarrow$   $\mathfrak{M}_{\omega}$ <sup>*n*</sup> with *η*<*μ* are well-defined and non-decreasing. Thus, for *η*<*μ*, the collection of  $\mathcal{E}_{\omega}$ <sup>*η*[*s*] with  $s \in \mathbb{T}^{>,>}$  forms a partition of  $\mathbb{T}^{>,>}$  into convex subclasses.</sup>

We say that  $\mathbb T$  is *confluent* if it is *v*-confluent. If  $\mathbb T$  has force **On**, then we say that  $\mathbb T$  is **On**-confluent, or *confluent*, if  $(\mathbb{T},(L_{\omega}\eta)_{\eta<\mu})$  is *μ*-confluent for all  $\mu \in \mathbf{On}$ .

### 4.5 Correspondence between fields and skeletons

**Proposition** 4.3. [\[5](#page-30-2), Theorem 1.1] *If*  $(\mathbb{T}, (L_{\omega^{\mu}})_{\mu \in \mathbf{On}})$  *is a confluent hyperserial skeleton, then there is a unique function* ∘:  $\mathbb{L} \times \mathbb{T} > \rightarrow \mathbb{T}$  *with* 

<span id="page-13-3"></span>
$$
\forall \mu \in \mathbf{On}, \forall \mathfrak{a} \in \mathfrak{M}_{\omega^{\mu}}, \quad \ell_{\omega^{\mu}} \circ \mathfrak{a} = L_{\omega^{\mu}} \mathfrak{a}
$$

*such that*  $(T, \circ)$  *is a confluent hyperserial field.* 

Assume now that  $\mathbb T$  is only a hyperserial skeleton of force  $v \in \text{On}^> \cup \{\text{On}\}$  and that  $\mu$  is an ordinal with  $0 < \mu < \nu$  such that  $(\mathbb{T}, (L_{\omega^{\eta}})_{\eta < \mu})$  is  $\mu$ -confluent. Let  $\beta = \omega^{\mu}$ . By [\[5](#page-30-2), Definition 4.11 and Lemma 4.12], the partial function  $L_\beta$  naturally extends into a function  $\mathbb{T}^{\geq, >} \to \mathbb{T}^{\geq, >}$  that we still denote by *Lβ* . This extended function is strictly increasing, by' [\[5](#page-30-2), Corollary 4.17]. If *μ* is a successor ordinal, then it satisfies the functional equation

<span id="page-13-2"></span>
$$
\forall s \in \mathbb{T}^{>,>} , L_{\beta} L_{\beta/\omega} s = L_{\beta} s - 1,
$$
\n(4.2)

by [\[5](#page-30-2), Proposition 4.13]. For  $\gamma < \beta$ , we have a strictly increasing function  $L_{\gamma}: \mathbb{T}^{>,>} \to \mathbb{T}^{>,>}$ obtained as a composition of functions  $L_{\omega^{\eta}}$  with  $\eta<\mu$ , as in [\(4.1\)](#page-11-2). By [\[5](#page-30-2), Proposition 4.7], we have

$$
\mathcal{E}_{\beta}[s] = \{t \in \mathbb{T}^{>,>} : \exists \gamma < \beta, L_{\gamma} t \approx L_{\gamma} s\}.
$$

#### 4.6 Hyperexponentiation

In a traditional transseries field  $\mathbb T$ , the transmonomials are characterized by the fact that, for any  $f \in \mathbb{T}^>$ , we have

<span id="page-13-0"></span>
$$
f \in \mathfrak{M} \iff \text{supp}\log f > 1. \tag{4.3}
$$

In particular, the logarithm log:  $\mathbb{T}^{\geq} \to \mathbb{T}$  is surjective as soon as  $\exp \varphi$  is defined for all  $\varphi \in \mathbb{T}$ with supp  $\varphi > 1$ . In hyperserial fields, similar properties hold for  $L_{\leq \varphi}$  atomic elements with respect to the hyperexponential  $E_{\alpha}$ <sup>n</sup>, as we will recall now.

Given  $v \in \mathbf{On}^> \cup \{\mathbf{On}\}\$ , let  $\mathbb T$  be a confluent hyperserial skeleton  $\mathbb T$  of force  $v$ . By [5, Theorem 4.1], we have a composition  $\circ: L_{< \omega} v \times T^{>} \to T$ . Given  $\eta < v$ , the extended function  $L_{\omega}$ n:  $\mathbb{T}^{>,>} \to \mathbb{T}^{>,>}$  is strictly increasing and hence injective. Consequently,  $L_{\omega}$ <sup>n</sup> has a partially defined functional inverse that we denote by  $E_{\omega}$ .

The characterization  $(4.3)$  generalizes as follows:

DEFINITION 4.4. [5, Definition 7.10] We say that  $\varphi \in \mathbb{T}^{>,\geq}$  is **1-truncated** if

<span id="page-14-4"></span>supp  $\varphi > 1$ .

Given  $0 < \eta < \nu$ , we say that a series  $\varphi \in \mathbb{T}^{>,\ge}$  is  $\omega^{\eta}$ -truncated if

 $\forall m \in \text{supp }\varphi, \quad m \prec 1 \Longrightarrow (\forall \gamma < \omega^{\eta}, \varphi < \ell^{\uparrow} \gamma_{\eta} \circ m^{-1}).$ 

For any  $\beta = \omega^{\eta} < \omega^{\nu}$ , we write  $\mathbb{T}_{>.\beta}$  for the class of  $\beta$ -truncated series in  $\mathbb{T}$ .

PROPOSITION 4.5. [5, Corollary 7.21] For  $f \in \mathbb{T}^{>,\ge}$  and  $\beta = \omega^{\eta} < \omega^{\nu}$ , we have

<span id="page-14-5"></span>
$$
f \in \mathfrak{M}_{\beta} \iff L_{\beta} f \in \mathbb{T}_{>,\beta}.
$$

In general, we have  $\mathbb{T}_{>.\beta}$  +  $\mathbb{R}^{\geq} \subseteq \mathbb{T}_{>.\beta}$ . Whenever  $\eta$  is a successor ordinal, we even have

<span id="page-14-3"></span>
$$
\mathbb{T}_{\geq,\beta} + \mathbb{R} = \mathbb{T}_{\geq,\beta} \tag{4.4}
$$

Let  $\varphi$  be a series such that  $E_{\beta}\varphi$  is defined. By [5, Lemma 7.14], the series  $\varphi$  is  $\beta$ -truncated if and only if

<span id="page-14-7"></span><span id="page-14-2"></span>
$$
\forall \gamma < \beta
$$
, supp  $\varphi > (L_{\gamma} E_{\beta} \varphi)^{-1}$ .

For  $\mu < v$ , the axiom  $\mathbf{R}_{\mu}$  is therefore equivalent to the inclusion  $L_{\omega^{\mu}} \mathfrak{M}_{\omega^{\mu}} \subseteq \mathbb{T}_{\geq \omega^{\mu}}$ . For  $s \in \mathbb{T}^{>,},$ there is a unique  $\leq$ -maximal truncation  $\sharp_{\beta}(s)$  of s which is  $\beta$ -truncated. By [5, Propositions 6.16 and 6.17], the classes

<span id="page-14-6"></span>
$$
\mathcal{L}_{\beta}[s] \; \coloneqq \; \left\{ t \in s + \mathbb{T}^{1} : t = s, \text{ or } \exists \gamma < \beta, \left( t < \ell_{\beta}^{\uparrow \gamma} \circ |s - t|^{-1} \right) \right\} \tag{4.5}
$$

with  $s \in \mathbb{T}^{>,>}$  form a partition of  $\mathbb{T}^{>,>}$  into convex subclasses. Moreover, the series  $\sharp_{\beta}(s)$  is both the unique  $\beta$ -truncated element and the  $\triangleleft$ -minimum of  $\mathcal{L}_{\beta}[s]$ . If  $E_{\beta}s$  is defined, then we have the following simplified definition [5, Proposition 7.19] of the class  $\mathcal{L}_{\beta}[s]$ :

<span id="page-14-1"></span><span id="page-14-0"></span>
$$
\mathcal{L}_{\beta}[s] \; := \; \bigg\{ t \in \mathbb{T}^{>,>} : \exists \, \gamma < \beta, \, t - s < \frac{1}{L_{\gamma} E_{\beta} s} \bigg\}.\tag{4.6}
$$

The following shows that the existence of  $E_\beta$  on  $\mathbb{T}^{\geq,>}$  is essentially equivalent to its existence on  $\mathbb{T}_{\geq \beta}$ .

PROPOSITION 4.6. [5, Corollary 7.24] Let  $\mu \leq v$  and assume that for  $\eta < \mu$ , the function  $E_{\alpha}$ <sup>n</sup> is defined on  $\mathbb{T}_{\geq \omega}$ <sup>n</sup>. Then each hyperlogarithm  $L_{\omega}$ <sup>n</sup> for  $\eta < \mu$  is bijective.

If Proposition 4.6 holds, then we say that  $\mathbb T$  is a (confluent) hyperserial field of force  $(v, \mu)$ . Since every function  $L_y$ ,  $\gamma < \omega^{\mu}$  is then a strictly increasing bijection  $\mathbb{T}^{>,>} \to \mathbb{T}^{>,>}$ , we obtain

$$
\mathcal{E}_{\lambda}[s] = \{t \in \mathbb{T}^{>,>} : \exists \gamma < \lambda, \exists r_0, r_1 \in \mathbb{R}^>, E_{\gamma}(r_0 L_{\gamma} s) < t < E_{\gamma}(r_1 L_{\gamma} s)\},\tag{4.7}
$$

for each ordinal  $\lambda = \omega^t$  with  $\iota \leq \mu$ . By [5, Corollary 7.23], for all  $s \in \mathbb{T}^{>,}$ , we have

<span id="page-15-2"></span>
$$
E_{\beta}(\sharp_{\beta}(s)) = \delta_{\beta}(E_{\beta}s). \tag{4.8}
$$

#### The transseries field No 5

Recall that **No** is identified with the ordered field of well-based series  $\mathbb{R}[[M\mathbf{o}]]$ . In this section, we describe, in the first level  $v = 1$  of our hierarchy, the properties of **No** equipped with the Kruskal-Gonshor logarithm.

#### **Surreal exponentiation**  $5.1$

In [19, Chapter 10], Gonshor defines the exponential function  $\exp:\mathbf{No} \rightarrow \mathbf{No}^>$ , relying on partial Taylor sums of the real exponential function. For  $a \in \mathbb{N}$  and  $n \in \mathbb{N}$ , write

<span id="page-15-0"></span>
$$
[a]_n := \sum_{k \leq n} \frac{a^k}{k!}.
$$

We then have the recursive definition

$$
\forall a \in \mathbf{No}, \quad \exp a \coloneqq \left\{ \exp(a_L) \left[ a - a_L \right]_\mathbb{N}, \exp(a_R) \left[ a - a_R \right]_{2\mathbb{N}+1} \middle| \frac{\exp a_R}{\left[ a_R - a \right]_\mathbb{N}}, \frac{\exp a_L}{\left[ a_L - a \right]_{2\mathbb{N}+1}} \right\}.
$$

We will sometimes write  $e^a$  instead of exp a. The function exp:  $(No, +, <) \rightarrow (No^>, \times, <)$  is a bijective morphism [19, Corollary 10.1, Corollary 10.3], which satisfies:

- exp coincides with the natural exponential on  $\mathbb{R} \subseteq$  **No** [19, Theorem 10.2].
- $e^{N_0} = Mo$  [19, Theorems 10.7, 10.8 and 10.9].

We define  $\log : \mathbf{No}^> \to \mathbf{No}$  to be the functional inverse of exp, and we set  $L_1 = \log \{1\}$  . Given an ordinal  $\alpha$ , we understand that  $\omega^{\alpha}$  still stands for the  $\alpha$ -th ordinal power of  $\omega$  from section 2.2.2 and warn the reader that  $\omega^{\alpha}$  does not necessarily coincide with  $e^{\alpha \log \omega}$ .

Together, the above facts imply that  $L_1$  satisfies the axioms  $FE_0$ ,  $A_0$ ,  $M_0$ ,  $R_0$  and SL. Therefore, (No, L<sub>1</sub>) is a hyperserial skeleton of force 1. The extension of L<sub>1</sub> to No<sup>></sup> from section 4.5 coincides with log. It was shown in [13] that  $(No, +, \times, <, exp)$  is an elementary extension of  $(\mathbb{R}, +, \times, <, exp)$ . See [28, 7, 8] for more details on exp and log.

#### 5.2 No as a transseries field

Berarducci and Mantova identified the class  $Mo_{\omega}$  of log-atomic numbers as  $Mo_{\omega} = Smp_{\mathcal{E}}$  [9, Corollary 5.17] and showed that  $(No, L_1)$  is 1-confluent [9, Corollary 5.11]. Thus  $(No, L_1)$  is a confluent hyperserial skeleton of force  $(1,1)$ . Thanks to [5, Theorem 1.1], it is therefore equipped with a composition law  $\mathbb{L}_{<\omega} \times \mathbf{No}^{>\succ} \to \mathbf{No}$ . See [29, 10] for further details on extensions of this composition law to exponential extensions of  $\mathbb{L}_{\leq \omega}$ .

<span id="page-15-3"></span>Berarducci and Mantova also proved [9, Theorem 8.10] that **No** is a field of transseries in the sense of [24, 29], i.e. that  $(No, L_1)$  satisfies the axiom **T4** of [29, Definition 2.2.1]. We plan to prove in subsequent work that  $(No,(L_o)^{\mu})_{\mu \in \mathbf{On}}$  satisfies a generalized version of T4.

#### <span id="page-15-1"></span>**Hyperserial structure on No** 6

We have seen in section 5 that  $(No, L_1)$  is a confluent hyperserial skeleton of force  $(\nu, \nu)$ for  $v = 1$ . The aim of this section is to extend this result to any ordinal v. More precisely, we will define a sequence  $(L_{\omega}\mu)_{\mu\in\mathbf{On}}$  of partial functions on **No** such that for each ordinal v, the structure (No,  $(L_{\alpha^{\mu}})_{\mu < \nu}$ ) is a confluent hyperserial skeleton of force (v, v), and  $L_1$  coincides with Gonshor's logarithm.

#### **Remarkable group actions on No**  $6.1$

Assume for the moment that we can define  $L<sub>y</sub>$  and  $E<sub>y</sub>$  as bijective strictly increasing functions on No<sup>>,></sup> for all ordinals  $\gamma$ . This is the case already for  $\gamma < \omega$ . Let us introduce several useful groups that act on No, as well as several remarkable subclasses of No.

Given an ordinal v, we write  $\alpha = \omega^{\nu}$  and we consider the function groups

$$
\mathcal{E}'_{\alpha} = \langle E_{\gamma} H_r L_{\gamma} : \gamma < \alpha \land r \in \mathbb{R}^{\geq} \rangle
$$
\n
$$
\mathcal{E}_{\alpha}^{*} = \langle E_{\gamma}, P_r : \gamma < \alpha \land r \in \mathbb{R}^{\geq} \rangle.
$$

where  $E_{\gamma}$ ,  $H_{s}$ ,  $P_{s}$  and  $L_{\gamma}$  act on **No**<sup>>,></sup>. We also define

<span id="page-16-4"></span><span id="page-16-3"></span><span id="page-16-2"></span><span id="page-16-1"></span><span id="page-16-0"></span>
$$
\mathcal{L}'_{\alpha} = L_{\alpha} \mathcal{E}'_{\alpha} E_{\alpha}
$$

$$
\mathcal{L}^*_{\alpha} = L_{\alpha} \mathcal{E}^*_{\alpha} E_{\alpha}
$$

We write  $L_{< \lambda} := \{L_y : y < \lambda\}$  and  $E_{< \lambda} := \{E_y : y < \lambda\}$  for each  $\lambda \leq \alpha$ . In the case when  $\alpha = 1$ , note that

$$
\mathcal{E}'_1 = \mathcal{H}
$$
  

$$
\mathcal{E}_1^* = \mathcal{P}
$$
  

$$
\mathcal{L}_1' = \mathcal{T}
$$
  

$$
\mathcal{L}_1^* = \mathcal{H}.
$$

By Proposition 3.3 and the fact the set-sized function groups  $\mathscr{E}'_{\alpha}$ ,  $\mathscr{E}'_{\alpha}$ ,  $\mathscr{E}'_{\alpha}$ , and  $\mathscr{L}^*_{\alpha}$  induce thin partitions of  $No^{>,\lambda}$ , we may define the following surreal substructures

<span id="page-16-8"></span><span id="page-16-7"></span><span id="page-16-6"></span><span id="page-16-5"></span>
$$
Mo'_{\alpha} := \mathrm{Smp}_{\mathscr{E}'_{\alpha}}\nMo^*_{\alpha} := \mathrm{Smp}_{\mathscr{E}^*_{\alpha}}\nTr_{\alpha} := \mathrm{Smp}_{\mathscr{L}'_{\alpha}}\nTr^*_{\alpha} := \mathrm{Smp}_{\mathscr{E}^*_{\alpha}}
$$

Here we note that **Mo**<sup>1</sup> corresponds to the class **Mo**<sup> $>$ </sup> = **Mo**<sub>1</sub> of infinite monomials in **No** and  $\pi'_{\mathcal{E}_1}$ maps positive infinite numbers to their dominant monomial. Similarly,  $Tr_1$  coincides with  $No$ . and  $\pi_{\mathcal{L}_1}$  maps  $a \in \mathbf{No}^{>,\rangle}$  to  $a_{\succ}$ . In sections 6 and 7, we will prove the following identities.



The first and third identities imply in particular that the classes  $M\omega_{\alpha}$  and  $N\omega_{\alpha}$  from section 4 are in fact surreal substructures, when regarding No as a hyperserial field.

#### 6.2 Inductive setting

 $\forall$ 

For the definition of the partial hyperlogarithm  $L_{\omega^{\mu}}$ , we will proceed by induction on  $\mu$ . Let  $\mu$ be an ordinal. Until the end of this section we make the following induction hypotheses:

### **Induction hypotheses**

<span id="page-17-0"></span> $I_{1,\mu}$ . For  $\eta < \mu$ , the partial hyperlogarithm  $L_{\omega^{\eta}}$  is defined; we have  $L_1 = \log 1$  Mo<sup>></sup> and  $(\mathbf{No}, (L_{\omega^{\eta}})_{n \leq \mu})$  is a confluent hyperserial skeleton of force  $(\mu, \mu)$ .

<span id="page-17-1"></span> $\mathbf{I}_{2,\mu}$ . For  $r, s \in \mathbb{R}$  with  $1 < s$  and for  $\gamma, \rho < \omega^{\mu}$  with  $\gamma < \rho$ , we have

$$
\forall a \in \mathbf{No}^{>,\succ}, \quad E_Y(rL_Ya) < E_\rho(sL_\rho a).
$$

<span id="page-17-2"></span> $\mathbf{I}_{3,\mu}$ . For  $\eta \leq \mu$ , the class  $\mathbf{Mo}'_{\omega^{\eta}}$  is that of  $L_{< \omega^{\eta}}$ -atomic surreal numbers, i.e.  $\mathbf{Mo}'_{\omega^{\eta}} = \mathbf{Mo}_{\omega^{\eta}}$ .

These induction hypotheses require a few additional explanations. Assuming that  $I_{1,\mu}$  holds, the partial functions  $L_{\omega}$ <sup>n</sup> with  $\eta < \mu$  extend into strictly increasing bijections  $L_{\omega}$ <sup>n</sup>: No<sup>>,></sup>  $\rightarrow$  $\text{No}^{>,\text{>}}$ , by the results from section 4. Using (1.3), this allows us to define a strictly increasing bijection  $L_v$ : **No**<sup>>,></sup>  $\rightarrow$  **No**<sup>>,></sup> for any  $\gamma < \mu$  and we denote by  $E_\gamma$  its functional inverse. In particular, this ensures that the hypotheses  $I_{2,\mu}$  and  $I_{3,\mu}$  make sense.

Remark 6.1. In addition to the above induction hypotheses, we will implicitly assume that our hyperlogarithms  $L_{\alpha}$ <sup>n</sup> for  $\eta < \mu$  are always defined by (6.1) below. In particular, our construction of  $L_{\omega^{\mu}}$  is not relative to any potential construction of the preceding hyperlogarithms  $L_{\omega}$ <sup>n</sup> with  $\eta < \mu$  that would satify the induction hypotheses  $I_{1,\mu}$ ,  $I_{2,\mu}$ , and  $I_{3,\mu}$ . Instead, we define one specific family of functions  $(L_{\omega} \eta)_{n \in \mathbf{On}}$  that satisfy our requirements, as well as the additional identities listed in subsection 6.1.

<span id="page-17-3"></span>PROPOSITION 6.2. The axioms  $I_{1,1}$ ,  $I_{2,1}$  and  $I_{3,1}$  hold for (No, L<sub>1</sub>).

PROOF. Section 5 shows that  $I_{1,1}$  holds. Consider  $r, s \in \mathbb{R}^>$  with  $s > 1$ . On  $\text{No}^{>,\,}$ , we have  $T_{\log r}$  < H<sub>s</sub>, hence  $H_r = E_1 T_{\log r} L_1$  <  $E_1 H_s L_1$ . It follows that we have  $E_n H_r L_n$  <  $E_{n+1} H_s L_{n+1}$ on No<sup>>,></sup> for all  $n \in \mathbb{N}$ . This implies that  $I_{2,1}$  holds. Finally,  $I_{3,1}$  is valid because of the relation  $M\mathbf{o}_{\omega} = \mathbf{Smp}_{\mathcal{E}}$ .  $\Box$ 

PROPOSITION 6.3. Let v be a limit ordinal and assume that  $I_{1,\mu}$ ,  $I_{2,\mu}$ , and  $I_{3,\mu}$  hold for all  $\mu < \nu$ . Then  $I_{1,v}$ ,  $I_{2,v}$ , and  $I_{3,v}$  hold.

PROOF. The statement  $I_{2, \nu}$  follows immediately by induction. Towards  $I_{3, \nu}$ , note that we have  $\mathbf{M}\mathbf{o}_{\alpha} = \bigcap_{\eta < \nu} \mathbf{M}\mathbf{o}_{\omega^{\eta}} = \bigcap_{\eta < \nu} \mathbf{M}\mathbf{o}'_{\omega^{\eta}}$  by  $\mathbf{I}_{1,\eta}$  (and thus  $\mathbf{D}\mathbf{D}_{\eta}$ ) and  $\mathbf{I}_{3,\eta}$  for all  $\eta < \nu$ . By [4, Proposition 6.28], we have  $\mathbf{Mo}'_{\alpha} = \bigcap_{n \leq v} \mathbf{Mo}'_{\alpha} = \mathbf{Mo}_{\alpha}$ . So  $\mathbf{I}_{3,v}$  holds.

By  $I_{1,n}$  for all  $\eta < \nu$ , we need only justify that  $(No,(L_o \eta)_{\eta < \nu})$  is  $\nu$ -confluent to deduce that  $I_{1,\nu}$ holds. For  $a \in \mathbb{N}0^{>,>}$ , by  $\mathbf{I}_{2,\nu}$ , there are a  $\alpha \in \mathbb{M}0'_{\alpha} = \mathbb{M}0_{\alpha}$  and a  $\beta = \omega^{\eta} < \alpha$  with  $E_{\beta}(1/2L_{\beta}a) \leq \alpha \leq \pi$  $E_{\beta}(2L_{\beta}a)$ . We deduce that  $L_{\beta}a \approx L_{\beta}a$ , thus  $a \in \mathcal{E}_{\beta}[a]$ . This concludes the proof.  $\Box$ 

From now on, we assume that  $I_{1,\mu}$ ,  $I_{2,\mu}$ , and  $I_{3,\mu}$  are satisfied for  $\mu \ge 1$  and we define

<span id="page-17-4"></span>
$$
\nu := \mu + 1
$$
  

$$
\alpha := \omega^{\nu}
$$
  

$$
\beta := \omega^{\mu}.
$$

The remainder of the section is dedicated to the definition of  $L_{\beta}$  and the proof of the inductive hypotheses  $I_{1,v}$ ,  $I_{2,v}$ , and  $I_{3,v}$  for v. In combination with Propositions 6.2 and 6.3, this will complete our induction and the proof of Theorem 1.1.

#### Defining the hyperlogarithm 6.3

Recall that we have  $\mathbf{Mo}_{\beta}' = \mathbf{Mo}_{\beta}$  by  $\mathbf{I}_{3,\mu}$ . In particular  $\mathbf{Mo}_{\beta}$  is a surreal substructure. Consider  $\eta < \nu$ . The skeleton (No,  $(L_{\omega})_{\nu < \eta}$ ) is a confluent hyperserial skeleton of force  $(\eta, \eta)$  by  $I_{1,\mu}$ . So for  $a \in \mathbf{No}^{>,\,}$ , (4.7) and  $\mathbf{I}_{2,\mu}$  yield  $\mathcal{E}_{\omega} \eta[a] = \mathcal{E}'_{\omega} \eta[a]$ .

In view of  $A_\mu$  $A_\mu$  and  $M_\mu$  $M_\mu$ , the simplest way to define  $L_\beta$  is *via* the cut equation:

$$
\forall \mathfrak{a} \in \mathbf{Mo}_{\beta}, \quad L_{\beta} \mathfrak{a} := \left\{ \mathbb{R}, L_{\beta} \mathfrak{a}' + \frac{1}{L_{<\beta} \mathfrak{a}'} \colon \mathfrak{a}' \in \mathfrak{a}_L^{\mathbf{Mo}_{\beta}} \middle| L_{\beta} \mathfrak{a}_R^{\mathbf{Mo}_{\beta}} - \frac{1}{L_{<\beta} \mathfrak{a}}, L_{<\beta} \mathfrak{a} \right\}.
$$
 (6.1)

Note the asymmetry between left and right options  $L_{\beta} \mathfrak{a}' + (L_{<\beta} \mathfrak{a}')^{-1}$  and  $L_{\beta} \mathfrak{a}'' - (L_{<\beta} \mathfrak{a})^{-1}$ (instead of  $L_{\beta} a'' - (L_{<\beta} a'')^{-1}$ ) for generic  $a' \in a_L^{\mathbf{M}\mathbf{o}_{\beta}}$  and  $a'' \in a_R^{\mathbf{M}\mathbf{o}_{\beta}}$ . In Corollary [7.4](#page-27-0) below, we will derive a more symmetric but equivalent cut equation for *Lβ* , as promised in the intro- duction. For now, we prove that [\(6.1\)](#page-18-0) is warranted and that  $A_\mu$  $A_\mu$ ,  $M_\mu$  $M_\mu$ , and  $R_\mu$  $R_\mu$  hold.

 $P$ ROPOSITION 6.4. *The function*  $L_{\beta}$  *is well-defined on*  $\mathbf{Mo}_{\beta}$  *and, for*  $\mathfrak{a}\in \mathbf{Mo}_{\beta}$ *, we have* 

$$
\mathbf{H}_{\mathfrak{a}}: \ \ \left(\forall \mathfrak{a}' \in \mathfrak{a}_L^{\mathbf{MO}_{\beta}}, L_{\beta} \mathfrak{a}' + \frac{1}{L_{<\beta} \mathfrak{a}'} < L_{\beta} \mathfrak{a} - \frac{1}{L_{<\beta} \mathfrak{a}}\right) \ and \ \left(\forall \mathfrak{a}'' \in \mathfrak{a}_R^{\mathbf{MO}_{\beta}}, L_{\beta} \mathfrak{a} + \frac{1}{L_{<\beta} \mathfrak{a}} < L_{\beta} \mathfrak{a}'' - \frac{1}{L_{<\beta} \mathfrak{a}''}\right).
$$

*Proof.* We prove this by induction on ( $\mathbf{Mo}_{\beta}$ , ⊑). Let a ∈  $\mathbf{Mo}_{\beta}$  such that  $\mathbf{H}_\mathfrak{b}$  holds for all  $\mathfrak{b} \in$  $\alpha_{\square}^{\mathbf{M}\mathbf{o}_{\beta}}$ . Let  $\alpha' \in \alpha_{L}^{\mathbf{M}\mathbf{o}_{\beta}}$  and  $\alpha'' \in \alpha_{R}^{\mathbf{M}\mathbf{o}_{\beta}}$ . We have  $\alpha' \in (\alpha')_{L}^{\mathbf{M}\mathbf{o}_{\beta}}$  or  $\alpha'' \in (\alpha')_{R}^{\mathbf{M}\mathbf{o}_{\beta}}$ , so  $\mathbf{H}_{\alpha'}$  or  $\mathbf{H}_{\alpha''}$  yields

<span id="page-18-0"></span>
$$
L_{\beta} \mathfrak{a}' + \frac{1}{L_{<\beta} \mathfrak{a}'} < L_{\beta} \mathfrak{a}'' - \frac{1}{L_{<\beta} \mathfrak{a}''}.
$$

For *γ* < *β*, we have  $\ell_{\gamma+1}$  <  $\frac{1}{2}$   $\ell_{\gamma}$  and  $\frac{1}{L_{\gamma} \mathfrak{a}'} > \frac{1}{L_{\gamma} \mathfrak{a}'}, \frac{1}{L_{\gamma} \mathfrak{a}}$ , v  $\frac{1}{L_\gamma \mathfrak{a}''}, \frac{1}{L_\gamma \mathfrak{a}}, \text{whence}$ 

$$
\frac{1}{L_{\gamma+1} \, \mathfrak{a}'} \; > \; \frac{2}{L_{\gamma} \, \mathfrak{a}'} \; > \; \frac{1}{L_{\gamma} \, \mathfrak{a}'} + \frac{1}{L_{\gamma} \, \mathfrak{a}''} + \frac{1}{L_{\gamma} \, \mathfrak{a}},
$$

for all *γ* <*β*. Hence,

$$
L_{\beta} \mathfrak{a}' + \frac{1}{L_{<\beta} \mathfrak{a}'} < L_{\beta} \mathfrak{a}'' - \frac{1}{L_{<\beta} \mathfrak{a}}.
$$

We clearly have  $L_{\beta} \mathfrak{a}'' - \frac{1}{L_{<\beta} \mathfrak{a}} \asymp L_{\beta} \mathfrak{a}'' > \mathbb{R}$ . Finally,

$$
L_{\beta} \mathfrak{a}' + \frac{1}{L_{<\beta} \mathfrak{a}'} \approx L_{\beta} \mathfrak{a}' \prec L_{<\beta} \mathfrak{a}',
$$

so  $L_{\beta}$   $\mathfrak{a}' + \frac{1}{L_{<\beta} \mathfrak{a}'} < L_{<\beta} \mathfrak{a}$ . This shows that  $L_{\beta} \mathfrak{a}$  is defined and

$$
L_{\beta} \mathfrak{a}' + \frac{1}{L_{<\beta} \mathfrak{a}'} \ < \ L_{\beta} \mathfrak{a} \ < \ L_{\beta} \mathfrak{a}'' - \frac{1}{L_{<\beta} \mathfrak{a}}.
$$

Since  $a' < a < a''$ , it follows that

$$
L_{\beta} \mathfrak{a}' + \frac{1}{L_{<\beta} \mathfrak{a}'} \ < L_{\beta} \mathfrak{a} \pm \frac{1}{L_{<\beta} \mathfrak{a}} \ < L_{\beta} \mathfrak{a}'' - \frac{1}{L_{<\beta} \mathfrak{a}}.
$$

By induction, this proves  $H_a$  for all  $a \in Mo_\beta$ . . □

*Proposition* 6.5*. The axiom* [M](#page-12-4)*μ holds.*

Proof. Let a, b∈**Mo***β* with a < b. Since **Mo***β* is a surreal substructure, there is a ເ∈**Mo***β* with  $\mathfrak{c} \sqsubseteq \mathfrak{a}$ ,  $\mathfrak{b}$  and  $\mathfrak{a} \leq \mathfrak{c} \leq \mathfrak{b}$ . If  $\mathfrak{a} < \mathfrak{c}$ , then we have  $L_{\beta} \mathfrak{a} + (L_{<\beta} \mathfrak{a})^{-1} < L_{\beta} \mathfrak{c} - (L_{<\beta} \mathfrak{c})^{-1}$  by  $\mathbf{H}_{\mathfrak{a}}$ . If  $\mathfrak{c} < \mathfrak{b}$ , then we have *L<sub>β</sub>*  $\epsilon$  + (*L*<sub><β</sub> $\epsilon$ )<sup>−1</sup> < *L<sub>β</sub>*  $\beta$  − (*L*<sub><β</sub> $\epsilon$ )<sup>−1</sup> by **H**<sub>b</sub>. We cannot have both α =  $\epsilon$  and  $\epsilon$  =  $\epsilon$ , so this proves that *L<sub>β</sub>* α + (*L*<*β* α)<sup>−1</sup><*L<sub>β</sub>*  $\mathfrak{b}$  −(*L*<*β*  $\mathfrak{b}$ )<sup>−1</sup>. Therefore  $\mathbf{M}_{\mu}$  $\mathbf{M}_{\mu}$  $\mathbf{M}_{\mu}$  holds. □

*Proposition* 6.6*. The axiom* [A](#page-12-3)*μ holds.*

*Proof*. The rightmost options in [\(6.1\)](#page-18-0) directly yield [A](#page-12-3)*μ*. □

*Proposition* 6.7*. The axiom* [R](#page-12-5)*μ holds.*

*Proof.* Let  $a \in Mo_{\beta}$  and write  $\varphi := L_{\beta}a$ . Let  $m \in \text{supp }\varphi$  with  $m < 1$ . We have  $\varphi < L_{<\beta}a$  and  $\varphi$ <sub>>m</sub>  $\leq \varphi$  so  $\varphi$ <sub>>m</sub>  $\lt L$ <sub> $lt$  $\beta$ </sub> a. Moreover  $\varphi$ <sub>>m</sub> is positive infinite. The number  $\varphi$ <sub>>m</sub> is strictly simpler than  $\varphi$ , so  $\varphi_{\geq m}$  does not lie in the cut which defines  $L_\beta$  a in [\(6.1\)](#page-18-0). Therefore, there is an  $\mathfrak{a}' \in \mathfrak{a}_L^{\mathbf{Mo}_\beta}$ or an  $\alpha'' \in \alpha_R^{\mathbf{Mo}_{\beta}}$  and an ordinal  $\gamma < \beta$  with  $\varphi_{> \mathfrak{m}} \leqslant L_{\beta} \alpha' + (L_{\gamma} \alpha')^{-1}$  or  $\varphi_{> \mathfrak{m}} \geqslant L_{\beta} \alpha'' - (L_{\gamma} \alpha)^{-1}$ . Consider the first case. We have  $L_{\beta} \alpha' + (L_{< \beta} \alpha')^{-1} \lt \varphi \leq \varphi_{\geq m} + \varphi_m \mathfrak{m} + \delta$  for a certain  $\delta \lt \mathfrak{m}$ . So  $\varphi_{\rm m}$  > 0 and

$$
\frac{1}{L_{<\beta}\mathfrak{a}'} < \frac{1}{L_{\gamma}\mathfrak{a}'} + \varphi_{\mathfrak{m}}\mathfrak{m}.
$$

For  $\rho < \beta$  with  $\gamma < \rho$ , we have  $(L_{\rho} \alpha')^{-1} > (L_{\gamma} \alpha')^{-1}$  so  $(L_{\rho} \alpha')^{-1} - (L_{\gamma} \alpha')^{-1} \asymp (L_{\rho} \alpha')^{-1}$ . We deduce that  $(L_\rho \alpha')^{-1}$  ≤ m for all such *ρ*. It follows that  $(L_\rho \alpha)^{-1}$  ≤ m for all *ρ* < *β*. In the second case, we directly get  $m > (L_\gamma \mathfrak{a})^{-1}$ . This proves that we always have  $m > (L_{<\beta} \mathfrak{a})^{-1}$ . In other words  $\supp \varphi > (L_{< \beta} \mathfrak{a})^{-1}$ , whence  $\mathbf{R}_{\mu}$  $\mathbf{R}_{\mu}$  $\mathbf{R}_{\mu}$  holds. □

*Proposition* 6.8*. If μ is a successor ordinal, then the cut equation* [\(6.1\)](#page-18-0) *is uniform.*

*Proof.* Let  $(\mathfrak{L_0}, \mathfrak{R_0})$  be a cut representation in  $\mathbf{Mo}_\beta$  and write  $\mathfrak{a} \coloneqq \{\mathfrak{L_0} \mid \mathfrak{R_0}\}_{\mathbf{Mo}_\beta}$ . For  $\mathfrak{l} \in \mathfrak{L_0}$ , we have  $L_{\beta}$   $\ell < L_{\beta}$   $\alpha < L_{<\beta}$  a so  $L_{\beta}$   $\ell < L_{<\beta}$   $\alpha$ . For  $r \in \mathfrak{R}_\alpha$ , we have  $L_{\beta}$   $\ell + (L_{<\beta} \ell)^{-1} < L_{\beta}$  r by  $M_{\mu}$  $M_{\mu}$ . Since  $1 < a$ , it follows that  $L_{\beta}$  Ι +  $(L_{<\beta}$  Ι)<sup>−1</sup> <  $L_{\beta}$  r −  $(L_{<\beta} a)$ <sup>−1</sup>. We may thus define the number

<span id="page-19-0"></span>
$$
\varphi := \left\{ \mathbb{R}, L_{\beta} \mathfrak{l} + \frac{1}{L_{<\beta} \mathfrak{l}} : \mathfrak{l} \in \mathfrak{L}_{\mathfrak{a}} \middle| L_{\beta} \mathfrak{R}_{\mathfrak{a}} - \frac{1}{L_{<\beta} \mathfrak{a}}, L_{<\beta} \mathfrak{a} \right\}.
$$

In order to show that [\(6.1\)](#page-18-0) is uniform, we need to prove that  $L_{\beta} \mathfrak{a} = \varphi$ , for any choice of the cut representation ( $\mathfrak{L}_{\mathfrak{a}}$ ,  $\mathfrak{R}_{\mathfrak{a}}$ ). We will do so by proving that  $L_{\beta} \mathfrak{a} \sqsubseteq \varphi$  and  $\varphi \sqsubseteq L_{\beta} \mathfrak{a}$ .

Recall that  $(\mathfrak{L}_\mathfrak{a}, \mathfrak{R}_\mathfrak{a})$  is cofinal with respect to  $(\mathfrak{a}_L^{\mathbf{M}\mathbf{o}_\beta} | \; \mathfrak{a}_R^{\mathbf{M}\mathbf{o}_\beta})$  and that  $L_\beta$  is strictly increasing. Consequently, we have

$$
\varphi < L_{\beta} \mathfrak{a}_R^{\mathbf{Mo}_{\beta}} - (L_{< \beta} \mathfrak{a})^{-1}.
$$

Given  $\alpha' \in \alpha_L^{\mathbf{MO}\beta}$  $\alpha' \in \alpha_L^{\mathbf{MO}\beta}$  $\alpha' \in \alpha_L^{\mathbf{MO}\beta}$ , there is an  $\mathfrak{l} \in \mathfrak{L}_\mathfrak{a}$  with  $\alpha' \leqslant \mathfrak{l}$ . By  $\mathbf{M}_\mu$ , we have  $L_\beta \alpha' + (L_\gamma \alpha')^{-1} \leqslant L_\beta \mathfrak{l} + (L_\gamma \mathfrak{l})^{-1}$ for all *γ* < *β*, so  $\varphi$  > {L<sub>β</sub> α' + (L<sub><β</sub> α')<sup>-1</sup>: α' ∈ α $_L^{Mof}$ β. This proves that  $\varphi$  lies in the cut defining L<sub>β</sub> α as per [\(6.1\)](#page-18-0), whence  $L_\beta \mathfrak{a} \subseteq \varphi$ .

Conversely, in order to prove that  $\varphi \sqsubseteq L_{\beta} \mathfrak{a}$ , it suffices to show that  $L_{\beta} \mathfrak{a}$  lies in the cut

$$
\left(L_{\beta}\mathfrak{l} + \frac{1}{L_{<\beta}\mathfrak{l}}\mathfrak{l} : \mathfrak{l} \in \mathfrak{L}_{\mathfrak{a}} \middle| L_{\beta}\mathfrak{R}_{\mathfrak{a}} - \frac{1}{L_{<\beta}\mathfrak{a}}\right).
$$

Let  $I \in \mathcal{L}_{\mathfrak{a}}$  and let  $\mathfrak{b} \in \mathbf{Mo}_{\beta}$  $\mathfrak{b} \in \mathbf{Mo}_{\beta}$  $\mathfrak{b} \in \mathbf{Mo}_{\beta}$  be  $\sqsubseteq$ -maximal with  $\mathfrak{b} \sqsubseteq I$ ,  $\mathfrak{a}$ . We have  $I \leq \mathfrak{b} \leq \mathfrak{a}$ , whence  $L_{\beta}\mathfrak{b} \leq L_{\beta}\mathfrak{a}$ , by  $M_{\mu}$ . If  $\mathfrak{b} \subset \mathfrak{l}$ , then  $\mathfrak{b} \in \mathfrak{l}_R^{\mathbf{M}\mathbf{o}_\beta}$ , so  $\mathbf{H}_\mathfrak{l}$  yields  $L_\beta\mathfrak{l} + (L_{<\beta}\mathfrak{l})^{-1} < L_\beta\mathfrak{b}$  and  $L_\beta\mathfrak{l} + (L_{<\beta}\mathfrak{l})^{-1} < L_\beta\mathfrak{a}$ . Otherwise  $I = b ∈ a_L^{\mathbf{MO}\beta}$ , so **H**<sub>a</sub> yields  $L_{\beta}I + (L_{<\beta}I)^{-1} < L_{\beta}$  a. This proves that  ${L_{\beta}I + (L_{<\beta}I)^{-1} : I ∈ \mathfrak{L}_\alpha} < L_{\beta}$  a.

Let  $\mathfrak{r} \in \mathfrak{R}_{\mathfrak{a}}$  and let  $\mathfrak{c} \in \mathbf{Mo}_{\beta}$  be ⊑-maximal with  $\mathfrak{c} \sqsubseteq \mathfrak{r}, \mathfrak{a}.$  As above, if  $\mathfrak{c} \sqsubset \mathfrak{a},$  then  $\mathfrak{c} \in \mathfrak{a}_R^{\mathbf{Mo}_{\beta}}$  so  $\mathbf{H}_{\mathfrak{a}}$ yields  $L_{\beta}$  α <  $L_{\beta}$  c − ( $L_{<\beta}$  α)<sup>−1</sup>, whence  $L_{\beta}$  α <  $L_{\beta}$  r − ( $L_{<\beta}$  α)<sup>−1</sup>. Otherwise α = c ∈ r $_{L}^{\mathbf{Mo}_{\beta}}$  so  $\mathbf{H}_{\mathrm{r}}$  yields  $L_{\beta}$  **r** >  $L_{\beta}$  α + ( $L_{<\beta}$  α)<sup>−1</sup>. Hence  $L_{\beta}$  α <  $L_{\beta}$   $\Re$  α − ( $L_{<\beta}$  α)<sup>−1</sup> and we conclude by induction. □

#### 6.4 Functional equation

In this subsection we derive  $FE_{\mu}$  $FE_{\mu}$ , under the assumption that  $\mu$  is a successor ordinal. We start with the following inequality.

<span id="page-20-0"></span>LEMMA 6.9. If  $\mu$  > 1, then we have  $E_{< \beta_{/\omega}} < E_{\beta_{/\omega}} H_2 L_{\beta_{/\omega}}$  on  $\text{No}^{>,>}$ .

PROOF. For  $\gamma < \beta_{/\omega}$ , there are  $\eta < \mu$  and  $n < \omega$  with  $\gamma < \omega^{\eta} n$ . We have

$$
E_{\gamma} < E_{\omega} \eta_n = E_{\omega} \eta + 1 \sum_{\mu} \sum_{\mu} \eta_{\mu} + 1 < E_{\omega} \eta + 1 H_2 L_{\omega} \eta_{\mu} + 1
$$

on  $\text{No}^{>,>}$  by (4.2). Note that  $\eta + 1 \leq \mu - \leq \mu$ , so  $I_{2,\mu}$  yields

$$
E_{\omega^{\eta+1}}H_2L_{\omega^{\eta+1}}\ \leq\ E_{\beta_{/\omega}}H_2L_{\beta_{/\omega}},
$$

whence  $E_{\gamma} < E_{\beta_{/\omega}} H_2 L_{\beta_{/\omega}}$ .

Let  $a \in Mo_{\beta}$ . Since  $Mo_{\beta}$  is a surreal substructure, we may consider the  $L_{<\beta}$ -atomic number

$$
\mathfrak{b} \; := \; \{ L_{\beta_{/\omega}} \mathfrak{a}_L^{\mathbf{Mo}_{\beta}} \, | \, L_{\beta_{/\omega}} \mathfrak{a}_R^{\mathbf{Mo}_{\beta}}, \mathfrak{a} \}_{\mathbf{Mo}_{\beta}}
$$

We claim that  $\mathfrak{b} = L_{\beta/\omega} \mathfrak{a}$ . Assume that  $\mu = 1$  and write  $\mathfrak{a} = \Xi_{\mathbf{Mo}_{\omega}} a$ . We have

$$
\log a = \Xi_{\mathbf{Mo}_{\omega}}(a-1)
$$
\n
$$
= \Xi_{\mathbf{Mo}_{\omega}}\{a_{L} - 1 \mid a_{R} - 1, a\}
$$
\n
$$
= \{\Xi_{\mathbf{Mo}_{\omega}}(a_{L} - 1) \mid \Xi_{\mathbf{Mo}_{\omega}}(a_{R} - 1), \Xi_{\mathbf{Mo}_{\omega}}a\}_{\mathbf{Mo}_{\omega}}
$$
\n
$$
= \{\log \Xi_{\mathbf{Mo}_{\omega}}a_{L} \mid \log \Xi_{\mathbf{Mo}_{\omega}}a_{R}, \Xi_{\mathbf{Mo}_{\omega}}a\}_{\mathbf{Mo}_{\omega}}
$$
\n
$$
= \{\log a_{L}^{\mathbf{Mo}_{\omega}} \mid \log a_{R}^{\mathbf{Mo}_{\omega}}, a\}_{\mathbf{Mo}_{\omega}}
$$
\n
$$
= b.
$$
\n
$$
(by [2, Proposition 2.5])
$$

Assume now that  $\mu > 1$ . The function  $L_{\beta/\omega}$  is strictly increasing with  $L_{\beta/\omega} < id_{\mathbf{No} > \cdot}$ . Therefore

$$
L_{\beta/\omega} \mathfrak{a} \in (L_{\beta/\omega} \mathfrak{a}_L^{\mathbf{M}\mathbf{0}_\beta} | L_{\beta/\omega} \mathfrak{a}_R^{\mathbf{M}\mathbf{0}_\beta, \mathfrak{a}})_{\mathbf{M}\mathbf{0}_\beta},
$$

so  $\mathfrak{b} \sqsubseteq L_{\beta_{/\omega}} \mathfrak{a}$ . Since  $\mathfrak{a} \in \mathbf{Mo}_{\beta}$ , the cut equation (6.1) for  $\mu$ -yields

$$
L_{\beta_{/\omega}}\mathfrak{a} = \{ \mathbb{R}, L_{\beta_{/\omega}}\mathfrak{a}' + (L_{<\beta}\mathfrak{a}')^{-1} : \mathfrak{a}' \in \mathfrak{a}_L^{\mathbf{MO}_{\beta_{/\omega}}} \mid L_{\beta_{/\omega}}\mathfrak{a}_R^{\mathbf{MO}_{\beta_{/\omega}}} - (L_{<\beta}\mathfrak{a})^{-1}, L_{<\beta_{/\omega}}\mathfrak{a} \}.
$$
 (6.2)

Given  $\alpha' \in \alpha_L^{\mathbf{Mo}_{\beta/\omega}}$ , we have  $\delta_{\beta}(\alpha') \in \alpha_L^{\mathbf{Mo}_{\beta}}$  and  $\alpha' \in \mathcal{E}_{\beta}[\delta_{\beta}(\alpha')]$ . We deduce that

$$
L_{\beta/\omega} \mathfrak{a}' \in L_{\beta/\omega} \mathcal{E}_{\beta}[\mathfrak{d}_{\beta}(\mathfrak{a}')] = \mathcal{E}_{\beta}[L_{\beta/\omega} \mathfrak{d}_{\beta}(\mathfrak{a}')] .
$$

Moreover, by definition, we have

<span id="page-20-2"></span>
$$
\mathfrak{b} > \mathcal{E}_{\beta}'[L_{\beta_{/\omega}} \mathfrak{d}_{\beta}(\mathfrak{a}')] = \mathcal{E}_{\beta}[L_{\beta_{/\omega}} \mathfrak{d}_{\beta}(\mathfrak{a}')] .
$$

so  $\mathfrak{b}>L_{\beta/\omega}$  a'. Symmetric arguments yield  $\mathfrak{b} < L_{\beta/\omega}$  a<sup>Mo<sub> $\beta/\omega$ </sub>. Lemma 6.9 implies that  $L_{<\beta/\omega}$  a  $\subseteq \mathscr{E}_{\beta}[\alpha]$ ,</sup> whence  $\delta_{\beta}(L_{<\beta/\omega}\mathfrak{a}) = {\mathfrak{a}}$ . We get  $\mathfrak{b} < \mathscr{E}_{\beta}\mathfrak{d}_{\beta}(L_{<\beta/\omega}\mathfrak{a})$ , whence  $\mathfrak{b} < L_{<\beta/\omega}\mathfrak{a}$ . Thus  $\mathfrak{b}$  lies in the cut defining  $L_{\beta_{\ell_{\omega}}}$  a in (6.2), so  $L_{\beta_{\ell_{\omega}}}$  a  $\subseteq$  b. This proves our claim that

<span id="page-20-3"></span>
$$
\forall \mathbf{a} \in \mathbf{Mo}_{\beta}, \quad L_{\beta_{/\omega}} \mathbf{a} = \{L_{\beta_{/\omega}} \mathbf{a}_{L}^{\mathbf{Mo}_{\beta}} \mid L_{\beta_{/\omega}} \mathbf{a}_{R}^{\mathbf{Mo}_{\beta}}, \mathbf{a}\}_{\mathbf{Mo}_{\beta}}.
$$
\n(6.3)

We now derive  $\mathbf{FE}_{\mu}$ .

PROPOSITION 6.10. For  $a \in \mathbf{Mo}_{\beta}$ , we have  $L_{\beta}L_{\beta}$   $a = L_{\beta}a - 1$ .

PROOF. We prove this by induction on  $(Mo_\beta, \subseteq)$ . Let  $a \in Mo_\beta$  be such that the result holds on  $\mathfrak{a}^{\textbf{Mo}}_{\square}$ . By (6.3), we have

$$
L_{\beta/\omega} \mathfrak{a} = \{L_{\beta/\omega} \mathfrak{a}_L^{\mathbf{MO}_{\beta}} \mid L_{\beta/\omega} \mathfrak{a}_R^{\mathbf{MO}_{\beta}} \mathfrak{a}_R^{\mathbf{MO}_{\beta}} \mathfrak{a}_R \}
$$

<span id="page-20-1"></span> $\Box$ 

Let a' and a'' range in  $\alpha_L^{\text{Mo}_{\beta}}$  and  $\alpha_R^{\text{Mo}_{\beta}}$  respectively. Proposition 6.8 and our induction hypothesis yield:

$$
L_{\beta}L_{\beta_{/\omega}}\mathfrak{a} = \left\{\mathbb{R}, L_{\beta}L_{\beta_{/\omega}}\mathfrak{a}' + \frac{1}{L_{<\beta}L_{\beta_{/\omega}}\mathfrak{a}'} \middle| L_{\beta}L_{\beta_{/\omega}}\mathfrak{a}'' - \frac{1}{L_{<\beta}L_{\beta_{/\omega}}\mathfrak{a}}, L_{\beta}\mathfrak{a} - \frac{1}{L_{<\beta}\mathfrak{a}}, L_{<\beta}L_{\beta_{/\omega}}\mathfrak{a} \right\}
$$
  
= 
$$
\left\{\mathbb{R}, L_{\beta}\mathfrak{a}' - 1 + \frac{1}{L_{<\beta}\mathfrak{a}'} \middle| L_{\beta}\mathfrak{a}'' - 1 - \frac{1}{L_{<\beta}\mathfrak{a}}, L_{\beta}\mathfrak{a} - \frac{1}{L_{<\beta}\mathfrak{a}}, L_{<\beta}\mathfrak{a} \right\}.
$$

On the other hand, we have

$$
L_{\beta} \mathfrak{a} - 1 = \left\{ \mathbb{R} - 1, L_{\beta} \mathfrak{a}' + \frac{1}{L_{<\beta} \mathfrak{a}'} - 1 \middle| L_{\beta} \mathfrak{a}'' - \frac{1}{L_{<\beta} \mathfrak{a}} - 1, L_{<\beta} \mathfrak{a} - 1, L_{\beta} \mathfrak{a} \right\}
$$
  
= 
$$
\left\{ \mathbb{R}, L_{\beta} \mathfrak{a}' + \frac{1}{L_{<\beta} \mathfrak{a}'} - 1 \middle| L_{\beta} \mathfrak{a}'' - \frac{1}{L_{<\beta} \mathfrak{a}} - 1, L_{\beta} \mathfrak{a}, L_{<\beta} \mathfrak{a} \right\}.
$$

In order to conclude that  $L_{\beta}L_{\beta}$   $\alpha = L_{\beta} \alpha - 1$ , it remains to show that  $L_{\beta} \alpha - 1 < L_{\beta} \alpha - (L_{<\beta} \alpha)^{-1}$ and that  $L_{\beta}L_{\beta}L_{\beta}$   $\alpha < L_{\beta}$  a. The first inequality holds because  $(L_{<\beta} \alpha)^{-1}$  is a set of infinitesimal numbers. An easy induction shows that  $L_{\beta/\omega} a < a$  for all  $a \in \mathbb{N}e^{>,>}$ . The second inequality follows, because  $L_{\beta}$  is strictly increasing on  $\mathbf{Mo}_{\beta}$ . This completes our inductive proof.  $\Box$ 

Combining our results so far, we have proved that  $(No,(L_o \eta)_{n\lt v})$  is a hyperserial skeleton of force  $\nu$ .

#### 6.5 Confluence

We next prove that  $(No, (L_o \eta)_{n \lt v})$  is *v*-confluent.

<span id="page-21-0"></span>LEMMA 6.11. If  $\mu$  is a non-zero limit ordinal, then the function groups  $\mathscr{E}'_B$  and  $\mathscr{E}_B^*$  are mutually pointwise cofinal. In particular, we have  $\mathbf{Mo}_{\beta} = \mathbf{Mo}_{\beta}^*$  and  $\mathbf{Tr}_{\beta} = \mathbf{Tr}_{\beta}^*$ .

PROOF. For  $\gamma \in (0, \beta)$  and  $r \in \mathbb{R}^>$ , we have  $E_Y H_Y L_Y \lt E_Y$  since  $H_Y \lt E_Y$ . We have

$$
\{L_{\rho}, E_{\rho} : \rho \in (0, \beta)\} \leq \mathcal{E}_{\beta}^*,
$$

whereas  $I_{2,u}$  yields

$$
\{E_{\rho}H_{r}L_{\rho}\,:\,\rho\in(0,\beta)\}\ \leq\ \mathcal{E}_{\beta}
$$

Therefore  $\mathcal{E}'_{\beta} \leq \mathcal{E}_{\beta}^*$ . For  $\rho < \beta$ , there is  $\eta < \mu$  with  $\rho < \omega^{\eta}$ . By (4.2), we have

$$
E_{\rho} < E_{\omega^{\eta}} = E_{\omega^{\eta+1}} T_1 L_{\omega^{\eta+1}} < E_{\omega^{\eta+1}} H_2 L_{\omega^{\eta+1}},
$$

which proves the inequality  $\mathscr{E}_{\beta}^* \leq \mathscr{E}_{\beta}$ .

<span id="page-21-2"></span>LEMMA 6.12. For each  $a \in No^{>,>}$ , any  $\sqsubseteq$ -minimal element of  $\mathcal{E}_{\alpha}[a]$  is  $L_{< \alpha}$ -atomic.

PROOF. Let 2l denote the class of numbers  $a \in No^{>}\rangle$  that are  $\subseteq$ -minimal in  $\mathscr{E}_{\alpha}[\alpha]$ . Any such  $\subseteq$ -minimal number a is also  $\subseteq$ -minimal in  $\mathscr{C}'_\beta[\mathfrak{a}] = \mathscr{C}_\beta[\mathfrak{a}] \subseteq \mathscr{C}_\alpha[\mathfrak{a}]$ , hence  $L_{<\beta}$ -atomic. Thus  $L_\beta$  is defined on  $\mathfrak{A}$ . It is enough to prove that  $\mathfrak{A}$  is closed under  $L_{\beta}$  in order to obtain that  $\mathfrak{A} \subseteq Mo_{\alpha}$ .

Consider  $a \in \mathfrak{A}$ , and recall that we have

$$
L_{\beta} \mathfrak{a} = \left\{ \mathbb{R}, L_{\beta} \mathfrak{a}' + \frac{1}{L_{<\beta} \mathfrak{a}'}; \mathfrak{a}' \in \mathfrak{a}_L^{\mathbf{MO}_{\beta}} \middle| L_{\beta} \mathfrak{a}_R^{\mathbf{MO}_{\beta}} - \frac{1}{L_{<\beta} \mathfrak{a}}, L_{<\beta} \mathfrak{a} \right\}.
$$
 (6.4)

Assume for contradiction that  $L_{\beta}$  a is not  $\subseteq$ -minimal in  $\mathcal{E}_{\alpha}[L_{\beta} \cap \mathcal{E}]$ . So there is a  $\mathfrak{b} \in \mathcal{E}_{\alpha}[L_{\beta} \cap \mathcal{E}]$ with  $\mathfrak{b} \subset L_{\beta} \mathfrak{a}$ . This implies that  $\mathfrak{b}$  lies outside the cut defining  $L_{\beta} \mathfrak{a}$ , so  $\mathfrak{b}$  is larger than a right option of  $(6.4)$  or smaller than a left option of  $(6.4)$ .

<span id="page-21-1"></span> $\Box$ 

Assume first that  $b < L_{\beta} a$ . So there is an  $a' \in a_L^{\mathbf{M}\mathbf{o}_\beta}$  with  $b \leq L_{\beta} a'$ . We have  $\delta_\alpha(a) = \delta_\alpha(b)$  so there is an  $n \in \mathbb{N}$  with

$$
(L_{\beta}\circ \delta_{\beta})^{\circ n}(\mathfrak{b}) \approx (L_{\beta}\circ \delta_{\beta})^{\circ n}(L_{\beta}\mathfrak{a}).
$$

Thus

$$
(L_{\beta} \circ \delta_{\beta})^{\circ (n+1)}(\alpha') \approx (L_{\beta} \circ \delta_{\beta})^{\circ (n+1)}(\alpha).
$$

This contradicts the  $\sqsubseteq$ -minimality of  $\alpha$ .

Now consider the other case when  $b > L_\beta a$ . In particular,  $b$  must be larger than a right option of [\(6.4\)](#page-21-1). Symmetric arguments imply that we cannot have  $\delta \ge L_\beta \mathfrak{a}''$  for some  $\mathfrak{a}'' \in \mathfrak{a}_R^{\mathbf{M}\mathbf{o}_\beta}$ . So there must exist a  $\gamma < \beta$  with  $\delta \ge L_\gamma \mathfrak{a}$ . If  $\mu$  is a limit ordinal, then  $\gamma < \mu_-$  so Lemma [6.11](#page-21-0) yields *δ*<sub>β</sub>(*Lγ*α) = α, whence *δ*<sub>β</sub>(*b*) ≽ α. If *μ* is a successor ordinal, then there is a *k* ∈ ℕ with *γ* ≤ *β*<sub>/ω</sub>*k*, so

$$
\delta_{\beta}(\mathfrak{b}) \geq \delta_{\beta}(L_{(\beta_{/\omega})k}\mathfrak{a}) = L_{(\beta_{/\omega})k}\mathfrak{a}
$$

and Proposition [6.1](#page-20-3)0 yields  $L_{\beta} \delta_{\beta}(b) \ge L_{\beta} \mathfrak{a} - k \ge L_{\beta} \mathfrak{a}$ . In both cases, we thus have  $L_{\beta} \delta_{\beta}(b) \ge L_{\beta} \mathfrak{a}$ . For any integer *n*>1, we deduce that

$$
(L_{\beta}\circ \mathfrak{d}_{\beta})^{\circ n}(\mathfrak{b}) \geq (L_{\beta}\circ \mathfrak{d}_{\beta})^{\circ n}(\mathfrak{a}) > (L_{\beta}\circ \mathfrak{d}_{\beta})^{\circ (n+1)}(\mathfrak{a}) = (L_{\beta}\circ \mathfrak{d}_{\beta})^{\circ n}(L_{\beta}\mathfrak{a}).
$$

This contradicts the fact that  $\delta$  lies in  $\mathcal{E}_{\alpha}[L_{\beta} \mathfrak{a}].$ 

We have shown that the cases  $b < L_\beta a$  and  $b > L_\beta a$  both lead to a contradiction. Consequently, *L*<sup>β</sup>  $\alpha$  is ⊆-minimal in  $\mathcal{E}_{\alpha}[L_{\beta} \alpha]$  and we conclude that  $L_{\beta} \mathfrak{A} \subseteq \mathfrak{A}$ , as claimed. □

<span id="page-22-1"></span>*COROLLARY* 6.13*.* ( $\mathbf{No}$ ,  $(L_{\omega} \eta)_{\eta < \nu}$ ) *is v*-confluent.

*Proof.* We already know that  $(No,(L_{\omega}\eta)_{\eta<\mu})$  is  $\mu$ -confluent by  $I_{1,\mu}$  $I_{1,\mu}$ . Recall that  $(No, \sqsubseteq)$  is wellfounded, so each class  $\mathscr{E}_{\alpha}[a]$  for  $a{\in}\mathbf{No}^{>,\succ}$  contains a ⊑-minimal element. Lemma [6.12](#page-21-2) therefore implies that No is *ν*-confluent. □

The corollary implies that  $({\bf No},(L_{\omega}\eta)_{\eta<\nu})$  is a confluent hyperserial skeleton of force *ν*. Moreover, the class  ${\rm No}_{\gt,\beta}$  is that of  $\triangleleft$ -minima and thus ⊑-minima in the convex classes

$$
\mathscr{L}_{\beta}[a] = \{b \in a + \mathbf{N} \mathbf{o}^{\prec} : b = a \vee (\exists \gamma < \beta, a < \ell_{\beta}^{\uparrow \gamma} \circ |a - b|^{-1})\},\
$$

for  $a \in \mathbf{No}^>,\rangle$ . In other words, we have  $\mathbf{No}_{>,\beta} = \mathbf{Smp}_{\mathscr{B}_{\beta}^*}$ . In order to conclude that  $\mathbf{No}_{>,\beta}$  is a surreal substructure, we still need to prove that the convex partition  $\mathscr{L}_{\beta}$  is thin. This will be done at the end of section [6.6](#page-23-0) below.

*Proposition* 6.14*. The cut equation* [\(6.1\)](#page-18-0) *is uniform.*

*P*roof. Let ( $\mathfrak{L}_\mathfrak{a}, \mathfrak{R}_\mathfrak{a}$ ) be a cut representation in  $\mathbf{Mo}_\beta$  and write α ≔{ $\mathfrak{L}_\mathfrak{a} \mid \mathfrak{R}_\mathfrak{a} \}$ Mo $_\beta$ . We have

<span id="page-22-0"></span>
$$
\mathscr{L}_{\beta}[L_{\beta} \mathfrak{L}_{\mathfrak{a}}] < \mathscr{L}_{\beta}[L_{\beta} \mathfrak{a}] < \mathscr{L}_{\beta}[L_{\beta} \mathfrak{R}_{\mathfrak{a}}].
$$

By  $(4.6)$ , this shows that

$$
L_{\beta} \mathfrak{a} \in \left( \mathbb{R}, L_{\beta} \mathfrak{l} + \frac{1}{L_{<\beta} \mathfrak{l}} : \mathfrak{l} \in \mathfrak{L}_{\mathfrak{a}} \middle| L_{\beta} \mathfrak{R}_{\mathfrak{a}} - \frac{1}{L_{<\beta} \mathfrak{a}}, L_{<\beta} \mathfrak{a} \right).
$$

In particular, the number

$$
\varphi := \left\{ \mathbb{R}, L_{\beta} \mathfrak{l} + \frac{1}{L_{<\beta} \mathfrak{l}} : \mathfrak{l} \in \mathfrak{L}_{\mathfrak{a}} \middle| L_{\beta} \mathfrak{R}_{\mathfrak{a}} - \frac{1}{L_{<\beta} \mathfrak{a}}, L_{<\beta} \mathfrak{a} \right\}
$$

is well-defined, with  $\varphi \subseteq L_\beta$  a. As in the proof of Proposition [6.8](#page-19-0), we have  $L_\beta$  a  $\subseteq \varphi$ , whence  $\varphi = L_{\beta}$  a. We conclude that the cut equation [\(6.1\)](#page-18-0) is uniform. □

### 6.6 Hyperexponentials

We have shown that  $({\bf No},(L_\omega\eta)_{\eta<\nu})$  is a hyperserial skeleton of force  $(\nu,\mu)$ . In order to prove that (No,  $(L_{\omega}$ η)<sub>η $lt$ ν</sub>) has force (*ν*, *ν*), it remains to prove that every β-truncated number *φ* has a hyperexponential *Eβφ*. This is the purpose of this subsection.

Proposition 6.15. We have  $L_{\beta}M{\bf o}_{\beta}$  =  $N{\bf o}_{\succ,\beta}$ , and  $E_{\beta}$  has the following cut equation on  $N{\bf o}_{\succ,\beta}$ :

<span id="page-23-4"></span><span id="page-23-3"></span>
$$
\forall \varphi \in \mathbf{No}_{>,\beta}, \quad E_{\beta}\varphi = \left\{ E_{< \beta}\varphi, \, E_{< \beta} \left( \frac{1}{\rho_R \mathbf{No}_{>,\beta} - \varphi} \right), \mathcal{E}_{\beta}' E_{\beta}\varphi_L^{\mathbf{No}_{>,\beta}} \right\} \mathcal{E}_{\beta}' E_{\beta}\varphi_R^{\mathbf{No}_{>,\beta}} \right\}.
$$

*Proof.* We prove the result by induction on ( $\mathbf{No}_{\gt,\beta}$ , ⊑). Let  $\varphi \in \mathbf{No}_{\gt,\beta}$  such that  $E_\beta$  is defined on  $\varphi_{\Box}^{\mathbf{N}\mathbf{o}}$ <sub> $\sim$ </sub>*β* with the given equation. We will first show that the number

$$
\mathfrak{a} \; := \; \left\{ E_{< \beta} \varphi, E_{< \beta} \left( \frac{1}{\varphi_R^{\mathbf{No}_{\gt, \beta} - \varphi}} \right), \mathcal{E}_{\beta}' E_{\beta} \varphi_L^{\mathbf{No}_{\gt, \beta}} \right\} \; \mathcal{E}_{\beta}' E_{\beta} \varphi_R^{\mathbf{No}_{\gt, \beta}} \right\} \tag{6.6}
$$

is well-defined. We will then prove that  $L_{\beta} \mathfrak{a} = \varphi$ .

Let  $\varphi' \in \varphi_L^{\mathbf{No}_{>,\beta}}$  and  $\varphi'' \in \varphi_R^{\mathbf{No}_{>,\beta}}$ . If  $\varphi' \in (\varphi'')_L^{\mathbf{No}_{>,\beta}}$ , then  $E_\beta \varphi'' > \mathscr{E}'_\beta E_\beta \varphi'$  by the definition of  $E_\beta\varphi''$ . So  $\mathscr{E}'_\beta E_\beta\varphi' < \mathscr{E}'_\beta E_\beta\varphi''$ . Otherwise, we have  $\varphi'' \in (\varphi')_R^{\mathbf{N}\mathbf{o}} \times \beta$ , whence  $\mathscr{E}'_\beta E_\beta\varphi'' > E_\beta\varphi'$  by definition of *Eβφ*′, so ℰ*<sup>β</sup>* ′*Eβφ*′<ℰ*<sup>β</sup>* ′*Eβφ*′′. So we always have

<span id="page-23-2"></span><span id="page-23-0"></span>
$$
\mathcal{E}_{\beta}'E_{\beta}\varphi_L^{\mathbf{No}_{>,\beta}} < \mathcal{E}_{\beta}'E_{\beta}\varphi_R^{\mathbf{No}_{>,\beta}}.
$$

We also have  $E_{<\beta}\varphi'' < E_{\beta}\varphi''$ , so  $E_{<\beta}\varphi < \mathcal{E}'_{\beta}E_{\beta}\varphi''$ . This proves that  $E_{<\beta}\varphi < \mathcal{E}'_{\beta}E_{\beta}\varphi_R^{\mathbf{No}_{>,\beta}}$ . It remains to show that

$$
E_{<\beta}\bigg(\frac{1}{\varphi_R^{\mathbf{N}\mathbf{o}_{>\beta}-\varphi}}\bigg)\;\leq\;\mathscr{E}'_{\beta}E_{\beta}(\varphi_R^{\mathbf{N}\mathbf{o}_{>\beta}}).
$$

Note that  $\varphi^{\mathbf{No}_>,\beta}_R\!>\!\mathscr{L}_\beta[\varphi],$  so by the definition of  $\mathscr{L}_\beta[\varphi],$  we have

<span id="page-23-1"></span>
$$
L_{\beta}^{\uparrow < \beta} \left( \frac{1}{\varphi_R^{\mathbf{No}_{>,\beta}} - \varphi} \right) < \varphi < \varphi_R^{\mathbf{No}_{>,\beta}}.
$$
\n(6.7)

.

Hence  $E_{<\beta}((\varphi_R^{\mathbf{N}\mathbf{o}_{>,\beta}}-\varphi)^{-1})< E_{\beta}\varphi_R^{\mathbf{N}\mathbf{o}_{>,\beta}},$  which completes the proof that  $\frak a$  is well-defined.

Let us now prove that  $L_\beta a = \varphi$ . Note that  $a \in \mathbf{Mo}_\beta$  by Proposition [3.2.](#page-8-2) First assume that  $\mu$  is a limit ordinal. Lemma [6.11](#page-21-0) yields  $\langle E_{\lt}\beta \rangle \neq \mathscr{E}_{\beta}$ , so we may write

$$
\mathbf{a} = \left\{ \delta \beta(\varphi), \delta \beta \left( \frac{1}{\varphi_R^{\mathbf{No}_{>,\beta}} - \varphi} \right), E_{\beta} \varphi_L^{\mathbf{No}_{>,\beta}} \middle| E_{\beta} \varphi_R^{\mathbf{No}_{>,\beta}} \right\}_{\mathbf{Mo}_{\beta}}.
$$

By [\(4.6\)](#page-14-2), for  $b \in \mathbf{No}^{>,>}$  the classes that  $\mathscr{L}_{\beta}[L_{\beta}b]$  and  $L_{\beta}b \pm (L_{<\beta}b)^{-1}$  are mutually cofinal and coinitial. Moreover, we have  $L_{\beta} E_{\beta} \psi = \psi$  for all  $\psi \in \varphi_{\square}^{\mathbf{N} \mathbf{o}_{>,\beta}}$ , by our hypothesis on  $\varphi$ . Hence, Proposition [6.14](#page-22-0) and [\(4.6\)](#page-14-2) imply

$$
L_{\beta} \mathfrak{a} = \left\{ \mathbb{R}, \mathscr{L}_{\beta}[L_{\beta} \mathfrak{d}_{\beta}(\varphi)], \mathscr{L}_{\beta}\left[L_{\beta} \mathfrak{d}_{\beta}\left(\frac{1}{\varphi_{R}^{\mathbf{N}\mathbf{o}_{>,\beta}} - \varphi}\right)\right], \mathscr{L}_{\beta}[\varphi_{L}^{\mathbf{N}\mathbf{o}_{>,\beta}}] \middle| \varphi_{R}^{\mathbf{N}\mathbf{o}_{>,\beta}} - \frac{1}{L_{<\beta} \mathfrak{a}} \right\}.
$$

Note that  $L_{\beta}$  a  $\in$  (  $\varphi^{\mathbf{No}_>,\beta}_{L} |$   $\varphi^{\mathbf{No}_>,\beta}_{R}$ ) $_{\mathbf{No}_{>,\beta}}$  so  $\varphi$   $\sqsubseteq$   $L_{\beta}$  a. Now  $L_{\beta}$   $\mathfrak{d}_{\beta}(\varphi)$   $\in$   $\mathscr{L}_{\beta}[L_{\beta}\varphi]$   $<$   $\varphi$ . We also have

$$
L_{\beta} \delta_{\beta} \left( \frac{1}{\varphi_R^{\mathbf{No}_{>,\beta-\varphi}}} \right) \in L_{\beta} \mathcal{E}'_{\beta} \left[ \frac{1}{\varphi_R^{\mathbf{No}_{>,\beta-\varphi}}} \right],
$$

where

$$
L_{\beta} \mathcal{E}_{\beta}^{\prime} \left[ \frac{1}{\varphi_R^{\mathbf{No}_{>,\beta-\varphi}}} \right] = L_{\beta} \mathcal{E}_{\beta}^{*} \left[ \frac{1}{\varphi_R^{\mathbf{No}_{>,\beta-\varphi}}} \right]
$$
 (by Lemma 6.11)  

$$
\leq L_{\beta}^{\uparrow} \leq \beta \left( \frac{1}{\varphi_R^{\mathbf{No}_{>,\beta-\varphi}}} \right)
$$
  

$$
< \varphi.
$$
 (by (6.7))

 $\int$ So *L<sub>β</sub>*  $\delta$ *β*( $\phi$ *R*<sup>0</sup>>, $\beta$  −  $\phi$ )<sup>-1</sup>< φ. Since  $\phi$  ∈ No<sub>≻,α</sub>, the inequality  $\mathscr{L}_{\beta}[\phi^{\mathbf{No}_{>},\beta}]$ < φ follows from Propo-sition [3.2.](#page-8-2) Finally, we have by definition that  $a > E  $\beta((\varphi_R^{\mathbf{N_o}}, \beta - \varphi)^{-1})$ , so  $\varphi_R^{\mathbf{N_o}}, \beta - (L  $\beta a)^{-1} > \varphi$ .$$ This proves that  $L_\beta \mathfrak{a} \subseteq \varphi$ , so  $L_\beta \mathfrak{a} = \varphi$ .

Assume now that  $\mu$  is a successor ordinal. For all  $b \in \mathbf{No}^{>,>}$ , the sets  $E_{< \beta} \varphi$ ,  $E_{< \beta} \varphi_{\beta}(\varphi)$ , and *Eβ*/*ω*<sup>ℕ</sup> *<sup>β</sup>* (*φ*) are mutually cofinal. So we can rewrite [\(6.6\)](#page-23-2) as

$$
\alpha = \left\{ E_{\beta_{/\omega} \mathbb{N}} \, \delta_{\beta}(\varphi), E_{\beta_{/\omega} \mathbb{N}} \, \delta_{\beta} \left( \frac{1}{\varphi_{R}^{\mathbf{N} \mathbf{o}_{>,\beta} - \varphi}} \right), \mathcal{E}_{\beta}^{\prime} E_{\beta} \varphi_{L}^{\mathbf{N} \mathbf{o}_{>,\beta}} \right\} \mathcal{E}_{\beta}^{\prime} E_{\beta} \varphi_{R}^{\mathbf{N} \mathbf{o}_{>,\beta}} \right\}
$$
\n
$$
= \left\{ E_{\beta_{/\omega} \mathbb{N}} \, \delta_{\beta}(\varphi), E_{\beta_{/\omega} \mathbb{N}} \, \delta_{\beta} \left( \frac{1}{\varphi_{R}^{\mathbf{N} \mathbf{o}_{>,\beta} - \varphi}} \right), E_{\beta} \varphi_{L}^{\mathbf{N} \mathbf{o}_{>,\beta}} \right\} E_{\beta} \varphi_{R}^{\mathbf{N} \mathbf{o}_{>,\beta}} \right\}
$$

As in the limit case, Proposition [6.14](#page-22-0) yields

$$
L_{\beta} \mathfrak{a} = \left\{ \mathbb{R}, \mathscr{L}_{\beta} \big[ L_{\beta}^{\uparrow < \beta} \mathfrak{d}_{\beta}(\varphi) \big], \mathscr{L}_{\beta} \bigg[ L_{\beta}^{\uparrow < \beta} \mathfrak{d}_{\beta} \bigg( \frac{1}{\varphi_R^{\mathbf{No}_{>, \beta} - \varphi}} \bigg) \right], \mathscr{L}_{\beta} \big[ \varphi_L^{\mathbf{No}_{>, \beta}} \big] \bigg| \varphi_R^{\mathbf{No}_{>, \beta} - \frac{1}{L_{< \beta} \mathfrak{a}}} \right\}.
$$

Let *γ* < *β*. There is an  $n \in \mathbb{N}$  with  $\gamma < \beta_{/\omega} n$ . Since  $L_{\beta} \varphi < \varphi - (n+1)$ , we have

$$
\varphi > L_{\beta}^{\uparrow \beta/\omega(n+1)} \delta_{\beta}(\varphi) \geq L_{\beta}^{\uparrow \gamma} \delta_{\beta}(\varphi) + 1.
$$

In particular  $φ > ∅_β[L_β^γ ∳_β(φ)]$ . We saw in [\(6.7\)](#page-23-1) that  $L_β^γ ∗_β((φ_R^N)^{−, β} - φ)^{-1}) < φ$ , whence  $\mathscr{L}_{\beta}$ [*L*<sub>β</sub><sup>↑</sup>γ<sub>β</sub>((*φ*<sub>R</sub><sup>No</sup>>,*β*−*φ*)<sup>-1</sup>)] < *φ*. We also obtain the inequalities

<span id="page-24-0"></span>
$$
\mathcal{L}_{\beta}[\varphi_L^{\mathbf{N}\mathbf{o}_{>,\beta}}] < \varphi < \varphi_R^{\mathbf{N}\mathbf{o}_{>,\beta}} - (L_{<\beta}\mathfrak{a})^{-1}
$$

in a similar way as in the limit case.

We conclude that  $\varphi = L_{\beta}$  a holds in general. It follows by induction that the formula for  $E_{\beta}$  is valid. In particular *L<sub>β</sub>*: **Mo**<sub>*β*</sub> → **No**<sub>></sub><sub>,*β*</sub> is surjective. □

With Proposition [6.15](#page-23-3), we have completed the proof of  $\mathbf{I}_{1,\nu}$ . By [\(4.7\)](#page-14-1), we have  $\mathcal{E}_{\beta\omega}[a] = \mathcal{E}_{\beta\omega}'[a]$ for all  $a \in \mathbf{No}^{>,>}$ . Given  $a \in \mathbf{No}_{>,\beta}$ , we also deduce from [\(4.6\)](#page-14-2) that the set  $a \pm (L_{<\beta} E_{\beta} a)^{-1}$  is cofinal and coinitial in ℒ*β* [*a*]. The convex partition defined by ℒ*<sup>β</sup>* is thus thin. By Proposi-tion [3.3](#page-8-0), the class  $\mathbf{No}_{>,\beta}$  is a surreal substructure with uniform cut equation

<span id="page-24-1"></span>
$$
\forall a \in \mathbf{No}, \quad \Xi_{\mathbf{No}_{>,\beta}} a = \{ \mathbb{R}, \mathcal{L}_{\beta}[\Xi_{\mathbf{No}_{>,\beta}} a_L] \mid \mathcal{L}_{\beta}[\Xi_{\mathbf{No}_{>,\beta}} a_R] \} \tag{6.8}
$$

For  $a \in \mathbf{No}$ , we have  $\mathscr{L}_{\beta}[\Xi_{\mathbf{No}_{>,\beta}}a] < \Xi_{\mathbf{No}_{>,\beta}}a_R$ , so  $\Xi_{\mathbf{No}_{>,\beta}}a < \Xi_{\mathbf{No}_{>,\beta}}a_R - (L_{<\beta}E_{\beta}\Xi_{\mathbf{No}_{>,\beta}}a)^{-1}$ . We deduce that the following equivalent is equivalent to [\(6.8\)](#page-24-0):

$$
\Xi_{\mathbf{No}_{>,\beta}}a = \left\{ \mathbb{R}, \Xi_{\mathbf{No}_{>,\beta}}a' + \frac{1}{L_{<\beta}E_{\beta}\Xi_{\mathbf{No}_{>,\beta}}a'} : a' \in a_L \; \middle| \; \Xi_{\mathbf{No}_{>,\beta}}a_R - \frac{1}{L_{<\beta}E_{\beta}\Xi_{\mathbf{No}_{>,\beta}}a} \right\}.
$$
 (6.9)

#### 6.7 End of the inductive proof

We now prove  $I_{2,\nu}$ ,  $I_{3,\nu}$  and Theorem [1.1.](#page-4-2)

LEMMA 6.16. *If*  $\mu$  *is a limit ordinal, then we have*  $E_\beta T_1 L_\beta$ *>E<sub><* $\beta$ *</sub> on No<sup>>,></sup>.* 

*P*ROOF. Let *a*∈**No**<sup>>,></sup>. We have  $\sharp_{\beta}(L_{\beta}a+1) > \sharp_{\beta}(L_{\beta}a)$ , so [\(4.8\)](#page-15-2) yields

<span id="page-25-2"></span>
$$
\delta_{\beta}(E_{\beta}(L_{\beta}a+1)) = E_{\beta}(\sharp_{\beta}(L_{\beta}a+1)) > E_{\beta}(\sharp_{\beta}(L_{\beta}a)) = \delta_{\beta}(a).
$$

We deduce that  $E_\beta(L_\beta a + 1) > \mathcal{E}_\beta a$  so  $E_\beta(L_\beta a + 1) > E_{\leq \beta} a$  by Lemma [6.11.](#page-21-0)

PROPOSITION 6.17. For  $r, s \in \mathbb{R}$  with  $s > 1$  and  $\gamma < \rho < \alpha$ , we have  $E_{\gamma} H_r L_{\gamma} < E_{\rho} H_s L_{\rho}$  on  $\mathbf{No}^{>,>}$ , *i.e.*  $\mathbf{I}_{2,v}$  *holds.* 

Proof. Throughout this proof, we consider inequalities and equalities of functions on No<sup>>,≻</sup>. Write  $γ = βm + ρ$  and  $ρ = βn + θ$  where  $m, n < ω$  and  $ι, θ < β$ . We have

$$
E_Y H_r L_Y = E_{\beta m} E_t H_r L_t L_{\beta m}
$$
 and  

$$
E_{\rho} H_r L_{\rho} = E_{\beta n} E_{\theta} H_s L_{\theta} L_{\beta n}.
$$

[I](#page-17-1)f  $m = n$ , then  $\iota < \theta$ , so  $I_{2,\mu}$  yields  $E_t H_r L_i < E_\theta H_s L_\theta$ , whence  $E_\gamma H_r L_\gamma < E_\rho H_s L_\rho$ . Assume that  $m < n$ . [I](#page-17-1)f *μ*− is a successor ordinal, then there is *p* < ω with *ι* < *β*/*ωp*. By I<sub>2,*μ*</sub>, we have  $E_{\theta}H_{s}L_{\theta} \ge H_{s} > T_{p}$ . So  $E_\beta(E_\theta H_s L_\theta) L_\beta > E_\beta T_p L_\beta = E_{\beta_{/\omega} p}$ . We conclude by noting that  $E_{\beta_{/\omega} p} > E_\iota > E_\iota H_r L_\iota$ . If  $\mu_-$  is a limit ordinal, then  $E_{\theta}H_{s}L_{\theta} > T_1$  so  $E_{\beta}(E_{\theta}H_{s}L_{\theta})L_{\beta} > E_{\iota} > E_{\iota}H_{r}L_{\iota}$  by Lemma [6.16.](#page-25-2) It follows that for  $k \in \mathbb{N}^>$ , we have  $E_{\beta(k+1)} E_{\theta} H_s L_{\theta} L_{\beta(k+1)} > E_{\beta k} E_t H_r L_t L_{\beta k}$ . An easy induction on k yields the result. the result. □

<span id="page-25-1"></span>*PROPOSITION 6.18.*  $\mathbf{Mo}_{\alpha}$ <sup>*is*</sup> *the class of*  $L_{\leq \alpha}$ *-atomic numbers, i.e.*  $\mathbf{I}_{3,y}$  *holds.* 

*Proof.* Let  $a \in \mathbf{No}^{>,>}$ . By Corollary [6.13](#page-22-1), the simplest element of  $\mathcal{E}_{\alpha}[a]$  is  $L_{< \alpha}$ -atomic. Since  $\mathscr{E}_{\alpha}[a] = \mathscr{E}_{\alpha}'[a]$ , we deduce that  $\mathbf{Mo}_{\alpha} \subseteq \mathbf{Mo}_{\alpha}$ .

Conversely, given  $a \in Mo_{\alpha}$ , we have  $b := \pi g_{\alpha}^{\prime}(a) \in Mo_{\alpha}^{\prime} \subseteq Mo_{\alpha}$ . Now  $b \in \mathcal{C}_{\alpha}^{\prime}[a]$ , so by  $I_{2,\nu}$ , there are  $r, s \in \mathbb{R}^>$  and  $\gamma < \alpha$  with  $E_\gamma(rL_\gamma a) < b < E_\gamma(sL_\gamma a)$ . Hence,  $L_\gamma b \approx L_\gamma a$ ,  $L_\gamma b = L_\gamma a$  and  $b = a$ . We conclude that  $\alpha \in \mathbf{Mo}_{\alpha}^{\prime}$ . □

In particular, the class  $M\mathbf{o}_\alpha$  is a surreal substructure. We have proved  $\mathbf{I}_{1,\nu}, \mathbf{I}_{2,\nu}$ , and  $\mathbf{I}_{3,\nu}$ , so we obtain the following by induction:

*Theorem* 6.19*. The field* (No,(*L<sup>ω</sup> <sup>η</sup>*)*η*∈On) *is a confluent hyperserial skeleton of force* (On,On)*.*

Combining this with Propositions [4.3](#page-13-3) and [4.6](#page-14-0), we obtain Theorem [1.1.](#page-4-2) Let us finally show that (No, ∘) contains only one *L*<On-atomic element.

*Proposition* 6.20*. The number*  $\omega$  *is the only*  $L_{**On***-atomic* element *in* **No***. For all*  $a \in \mathbf{No}^{>,\lambda}$ ,$ *there is*  $\gamma \in \mathbf{On}$  *with*  $L_{\gamma} a \asymp L_{\gamma} \omega$ .

*Proof*. The number *ω* lies in Mo*ω <sup>μ</sup>* for all *μ*∈On, so it is *L*<On-atomic. For *ν*∈On, the number  $E_{\omega^{\nu}}\omega = \{E_{<\omega^{\nu}}\omega \mid \emptyset\}$  is an ordinal. As a sign sequence, the number  $L_{\omega^{\nu}}\omega = \{\emptyset \mid L_{<\omega^{\nu}}\omega\}_{\mathbf{No}}$ >,> is  $\omega$ followed by a string containing only minuses [\[2](#page-30-0), Lemma 2.6]. Since the sequences  $(E_{\omega^{\nu}}\omega)_{\nu\in\mathbf{On}}$ and  $(L_{\omega^{\nu}}\omega)_{\nu \in \mathbf{On}}$  are strictly increasing and strictly decreasing respectively, the classes  $\{E_{\omega^{\nu}}\omega\}$ : *ν*∈On} and {*L*<sub>ω</sub>*ν*ω: *ν*∈On} are respectively cofinal and coinitial in No<sup>>,></sup> = {*a*∈No: ω⊑ *a*}. Thus for  $a \in \mathbf{No}^{>,>}$ , there is  $v \in \mathbf{On}$  with  $E_{\omega}v \omega > a > L_{\omega}v \omega$ , whence  $L_{\omega}v+1 \omega \approx L_{\omega}v+1 a$ .

### <span id="page-25-0"></span>7 Remarkable identities

In this section, we give various identities regarding the function groups introduced in Sec-tion [6.1.](#page-16-0) In what follows, *v* is a non-zero ordinal and  $\alpha = \omega^{\nu}$ .

#### Simplified cut equations for  $L_{\alpha}$  and  $E_{\alpha}$  $7.1$

Given  $\varphi \in \mathbf{No}^{>,\rangle}$ , let  $E_{\lhd \alpha} = \{E_{(\alpha_{/\omega})n} \varphi : n \in \mathbb{N}\}\$ if  $\nu$  is a successor ordinal and  $E_{\lhd \alpha} \varphi = \{\varphi\}\$ if  $\nu$  is a limit ordinal. In this subsection, we will derive the following simplified cut equations for  $L_{\alpha}$ on  $\mathbf{Mo}_{\alpha}$  and  $E_{\alpha}$  on  $\mathbf{No}_{\geq,\alpha}$ :

$$
\forall \mathfrak{a} \in \mathbf{Mo}_{\alpha}, \ L_{\alpha} \mathfrak{a} = \{L_{\alpha} \mathfrak{a}_{L}^{\mathbf{Mo}_{\alpha}} | L_{\alpha} \mathfrak{a}_{R}^{\mathbf{Mo}_{\alpha}}, L_{<\alpha} \mathfrak{a}\}_{\mathbf{No}_{>,\alpha}}
$$
\n
$$
\left(\sum_{\alpha} \mathfrak{a}_{\alpha}^{\alpha}, \sum_{\alpha} \mathfrak{a}_{\alpha}^{\alpha} | L_{\alpha} \mathfrak{a}_{R}^{\alpha}\right) = \left(\sum_{\alpha} \mathfrak{a}_{\alpha}^{\alpha} | L_{\alpha} \mathfrak{a}_{R}^{\alpha}\right) \tag{7.1}
$$

$$
= \left\{ \mathbb{R}, L_{\alpha} \mathfrak{a}' + \frac{1}{L_{<\alpha} \mathfrak{a}'}; \mathfrak{a}' \in \mathfrak{a}_L^{\mathbf{MO}_{\alpha}} \middle| L_{\alpha} \mathfrak{a}'' - \frac{1}{L_{<\alpha} \mathfrak{a}''}, L_{<\alpha} \mathfrak{a}: \mathfrak{a}'' \in \mathfrak{a}_L^{\mathbf{MO}_{\alpha}} \right\},\tag{7.2}
$$

$$
\forall \varphi \in \mathbf{No}_{>,\alpha}, \ E_{\alpha} \varphi = \{ E_{\lhd \alpha} \delta_{\alpha}(\varphi), E_{\alpha} \varphi_{L}^{\mathbf{No}_{>,\alpha}} \mid E_{\alpha} \varphi_{R}^{\mathbf{No}_{>,\alpha}} \mid \mathbf{Mo}_{\alpha} \tag{7.3}
$$

<span id="page-26-3"></span><span id="page-26-2"></span><span id="page-26-1"></span><span id="page-26-0"></span>
$$
= \{ E_{<\alpha} \varphi, \mathcal{E}_{\alpha} E_{\alpha} \varphi_L^{\mathbf{N}\mathbf{o}_{>\alpha}} \mid \mathcal{E}_{\alpha} E_{\alpha} \varphi_R^{\mathbf{N}\mathbf{o}_{>\alpha}} \}.
$$
\n(7.4)

For all  $a \in \mathbb{N}e^{>,\rangle}$ , the set  $E_{\lhd \alpha} \delta_{\alpha}(a)$  contains only  $L_{\lhd \alpha}$ -atomic numbers, so (7.3) is indeed a cut equation of the form  $\{\rho \mid \lambda\}$ <sub>Mo</sub>.

Remark 7.1. The changes with respect to (6.1) and (6.5) lie in the occurrence of  $\alpha''$  instead of  $\alpha$ in (7.2) and the (related) absence of the left option  $E_{< \alpha}((\varphi_R^{\mathbf{No}_{\geq,\alpha}} - \varphi)^{-1})$  in (7.4). So (7.2) and (7.4) give lighter sets of conditions than those in (6.1) and (6.5) to define  $L_{\alpha}$  and  $E_{\alpha}$ . This seemingly meager simplification will be crucial in further work. Indeed, combined with Proposition 3.1, this allows one to determine large classes of numbers *a*, *b* with  $a \subseteq b \Longrightarrow E_{\alpha} a \subseteq E_{\alpha} b$ .

First note that the cut equations (7.1) and (7.3) if they hold are uniform (see [6, Remark 1]). Moreover, we claim that  $(7.1, 7.2)$  are equivalent and that  $(7.3, 7.4)$  are equivalent. Indeed, recall that for a thin convex partition  $\Pi$  of a surreal substructure S and any cut representation  $(L, R)$ in  $Smp_{\Pi}$ , one has

$$
{L \mid R}_{\text{Smp}_{\Pi}} = {\Pi[L] \mid \Pi[R]}\rangle_{\text{S}}
$$

For  $\alpha' \in \alpha_L^{\mathbf{Mo}_\alpha}$  and  $\alpha'' \in \alpha_R^{\mathbf{Mo}_\alpha}$  the classes  $L_\alpha \alpha' + (L_{< \alpha} \alpha')^{-1}$  and  $\mathcal{L}_\alpha[L_\alpha \alpha']$  are mutually cofinal by (4.6). Similarly,  $L_{\alpha}$  a''  $-L_{\leq \alpha}$  a'')<sup>-1</sup> and  $\mathcal{L}_{\alpha}[L_{\alpha}$  a''] are mutually coinitial. By Lemma 6.11, the classes  $E_{\leq \alpha} \varphi$  and  $\mathcal{E}_{\alpha}[E_{\leq \alpha} \delta_{\alpha}(\varphi)]$  are mutually cofinal. So it is enough to prove that (7.1) and (7.3) are valid cut equations for  $L_{\alpha}$  and  $E_{\alpha}$  respectively.

LEMMA 7.2. If v is a successor ordinal, then the identities  $(7.1)$  and  $(7.3)$  hold.

PROOF. Let  $a \in Mo_{\alpha}$  and set

$$
\varphi := \{L_{\alpha} \mathfrak{a}_{L}^{\mathbf{MO}_{\alpha}} \mid L_{\alpha} \mathfrak{a}_{R}^{\mathbf{MO}_{\alpha}}, L_{<\alpha} \mathfrak{a}\}_{\mathbf{NO}_{>,\alpha}}= \{R, L_{\alpha} \mathfrak{a}' + \frac{1}{L_{<\alpha} \mathfrak{a}'} : \mathfrak{a}' \in \mathfrak{a}_{L}^{\mathbf{MO}_{\alpha}} \mid L_{\alpha} \mathfrak{a}'' - \frac{1}{L_{<\alpha} \mathfrak{a}''}, L_{<\alpha} \mathfrak{a} : \mathfrak{a}'' \in \mathfrak{a}_{R}^{\mathbf{MO}_{\alpha}}\}.
$$

We have  $\mathscr{L}_{\alpha}[L_{\alpha} \mathfrak{a}_L^{\mathbf{Mo}_{\alpha}}] < \varphi < L_{<\alpha} \mathfrak{a}$  so in view of (6.1), it is enough to prove that  $\varphi < L_{\alpha} \mathfrak{a}_R^{\mathbf{Mo}_{\alpha}}$  $(L <sub>$\alpha$</sub>  \mathfrak{a})^{-1}$  to conclude that  $\varphi = L_{\alpha} \mathfrak{a}$ . Let  $\mathfrak{a}'' \in \mathfrak{a}_R^{\mathbf{Mo}_{\alpha}}$ . If  $\mathfrak{a}'' \in \mathcal{E}_{\alpha}^*[\mathfrak{a}]$ , then the inequality  $\varphi < L_{\alpha} \mathfrak{a}''$ entails  $\varphi < \mathcal{L}_{\alpha}[L_{\alpha} \alpha'']$  whence  $\varphi < L_{\alpha} \alpha'' - (L_{<\alpha} \alpha'')^{-1}$  and  $\varphi < L_{\alpha} \alpha'' - (L_{<\alpha} \alpha)^{-1}$ . Otherwise, we have  $a < L_{<\alpha} a''$ , so  $L_{\alpha} a < L_{\alpha} a'' - 2$ , and  $L_{\alpha} a'' - (L_{<\alpha} a)^{-1} > L_{\alpha} a + 1$ . It is enough to prove that  $L_{\alpha}$   $\alpha + 1 \ge \varphi$ . Recall that

$$
L_{\alpha} \mathfrak{a} + 1 = \left\{ L_{\alpha} \mathfrak{a}, L_{\alpha} \mathfrak{a}' + \frac{1}{L_{<\alpha} \mathfrak{a}'} + 1 : \mathfrak{a}' \in \mathfrak{a}_L^{\mathbf{M}\mathbf{0}_\alpha} \mid L_{\alpha} \mathfrak{a}_R^{\mathbf{M}\mathbf{0}_\alpha} - \frac{1}{L_{<\alpha} \mathfrak{a}} + 1, L_{<\alpha} \mathfrak{a} \right\}
$$

by (3.1). We see that  $L_{\alpha} a' + \frac{1}{L_{<\alpha} a'} < L_{\alpha} a + 1$  for all  $a' \in a_L^{\text{Mo}\alpha}$ . We have  $1 - \frac{1}{L_{<\alpha} a} > \frac{1}{L_{<\alpha} a_R^{\text{Mo}\alpha}}$  so  $L_{\alpha} a_R^{\text{Mo}\alpha} - \frac{1}{L_{<\alpha} a} + 1 > \varphi$ . Thus  $\varphi \le L_{\alpha} a + 1$ . So (7.1) holds.

The hyperserial field of surreal numbers 27 and 27 and

Now let  $\psi \in \mathbf{No}_{\succ,\alpha}$  and set

$$
\mathfrak{b} \; := \; \big\{ E_{\alpha_{/\omega} \mathbb{N}} \, \mathfrak{d}_{\alpha} (\psi), E_{\alpha} \, \psi_L^{\mathbf{No}_{>,\alpha}} \, | \; E_{\alpha} \, \psi_R^{\mathbf{No}_{>,\alpha}} \big\}_{\mathbf{Mo}_{\alpha}}.
$$

By uniformity of [\(7.1\)](#page-26-0), we have

$$
L_{\alpha} \mathfrak{b} = \{ L_{\alpha} E_{\alpha_{/\omega} \mathbb{N}} \mathfrak{d}_{\alpha}(\psi), \psi_L^{\mathbf{N}\mathbf{o}_{>,\alpha}} | \psi_R^{\mathbf{N}\mathbf{o}_{>,\alpha}}, L_{<\alpha} \mathfrak{b}_{\mathbf{N}\mathbf{o}_{>,\alpha}},
$$

whence  $L_{\alpha}$   $\mathfrak{b} \supseteq {\psi_L^{\mathbf{No}_{>},\alpha} \mid \psi_R^{\mathbf{No}_{>},\alpha}} \models \psi$ . Conversely,  $\mathfrak{b} > E_{\alpha/\omega} \mathfrak{N} \mathfrak{d}(\psi)$  and  $\mathfrak{b} > E_{<\alpha} \psi$ , so  $\psi < L_{< \alpha}$  b. We have  $L_{\alpha} E_{\alpha_{/\omega} N}$   $\delta_{\alpha}(\psi) = L_{\alpha} \delta_{\alpha}(\psi) + N$ . Since  $L_{\alpha} \delta_{\alpha}(\psi) < L_{\alpha_{/\omega}} \delta_{\alpha}(\psi) < \psi$ , this yields  $L_{\alpha} E_{\alpha_{/\omega} N} \delta_{\alpha}(\psi) < \psi$ . This proves that  $\psi$  lies in the cut defining  $L_{\alpha} \mathfrak{b}$ . We conclude that  $\psi = L_{\alpha}$  b, hence [\(7.3\)](#page-26-1) holds. □

We now assume that *v* is a limit ordinal. For  $z \in \textbf{No}$ , define

$$
F(z) := \{b_{\alpha}(\Xi_{\mathbf{No}_{>,\alpha}}z), F(z_L) \mid F(z_R)\}\mathbf{Mo}_{\alpha}, \text{ and}
$$
  
\n
$$
\Xi z := \{\mathbb{R}, \Xi z' + (L_{<\alpha}F(z'))^{-1} : z' \in z_L \mid \Xi z_R - (L_{<\alpha}F(z))^{-1}\}.
$$

*LEMMA* 7.3*. For all*  $z \in$  **No**, we have

$$
F(z) \t is defined \t(7.5)
$$

<span id="page-27-4"></span><span id="page-27-3"></span><span id="page-27-2"></span><span id="page-27-1"></span>
$$
\Xi z \text{ is defined} \tag{7.6}
$$

$$
\Xi z = \Xi_{\mathbf{No}_{>,\alpha}} z \tag{7.7}
$$

$$
F(z) = E_{\alpha} \Xi z \tag{7.8}
$$

*Proof.* We prove the result by induction on  $(No, \subseteq)$ . Let  $z \in No$  be such that [\(7.5\)](#page-27-1), [\(7.6\)](#page-27-2), [\(7.7\)](#page-27-3) and [\(7.8\)](#page-27-4) hold for all  $\gamma \in \mathbb{N}$ **o** with  $\gamma \subset z$ .

For  $z'' \in z_R$  and  $z' \in z_L$ , we have  $\delta_\alpha(\Xi_{N\mathbf{0}_{\geq \alpha}}z) \leq \delta_\alpha(\Xi_{N\mathbf{0}_{\geq \alpha}}z'') < F(z'')$ . We have  $F(z') < F(z'')$ by definition of  $F(z'')$  if  $z' \in (z'')_L$  and by definition of  $F(z')$  if  $z'' \in (z')_R$ . This proves that  $F(z)$ is defined.

Let *z*′ ∈ *z<sub>L</sub>* and *z*<sup>′′</sup> ∈ *z<sub>R</sub>*. If *z*<sup>′</sup> ∈ (*z*<sup>′′</sup>)<sub>*L*</sub>, then we have  $\Xi$ *z*<sup>′′</sup> >  $\Xi$ *z*<sup>′</sup> + (*L*<sub>≤α</sub>*F*(*z*<sup>′</sup>))<sup>−1</sup> by definition of  $\Xi z''$ . Since  $F(z') < F(z)$  and  $F(z), F(z') \in M_0$ , we have  $L_y F(z') < L_y F(z)$  for all  $\gamma$ *α*. We deduce that  $\Xi z'' - (L_{< a}F(z))^{-1} > Ξ z' + (L_{< a}F(z'))^{-1}$ . If  $z'' ∈ (z')_L$ , then  $\Xi z' < Ξ z'' (L <sub>$\alpha$</sub>  F(z'))<sup>-1</sup>$  by definition of  $\Xi z'$ . Since  $F(z') < F(z)$ , we obtain  $\Xi z'' - (L <sub>$\alpha$</sub>  F(z))<sup>-1</sup> > \Xi z' +$  $(L_{<\alpha} F(z'))^{-1}.$  This proves that  $\Xi z$  is defined.

Since [\(7.7\)](#page-27-3) and [\(7.8\)](#page-27-4) hold on  $z_{\square}$ , we have

$$
\Xi z = \{ \mathbb{R}, \Xi_{\mathbf{N}\mathbf{o}_{>,\alpha}} z' + (L_{< \alpha} E_{\alpha} \Xi_{\mathbf{N}\mathbf{o}_{>,\alpha}} z')^{-1} : z' \in z_L \mid \Xi_{\mathbf{N}\mathbf{o}_{>,\alpha}} z_R - (L_{< \alpha} E_{\alpha} \Xi_{\mathbf{N}\mathbf{o}_{>,\alpha}} z)^{-1} \}
$$

By [\(6.9\)](#page-24-1), this yields  $\Xi z = \Xi_{NQ} z$ , so [\(7.7\)](#page-27-3) holds for *z*.

From [\(7.7\)](#page-27-3), we get  $\delta_{\alpha}(\Xi_{NQ_{\alpha},\alpha}z) = \delta_{\alpha}(\Xi z)$ . By Proposition [6.14](#page-22-0) and our assumption that [\(7.8\)](#page-27-4) holds on  $z_\sqsubset$ , we have

$$
L_{\alpha}F(z) = \{ \mathbb{R}, \mathcal{L}_{\alpha}[L_{\alpha} \mathfrak{d}_{\alpha}(\Xi z)], \mathcal{L}_{\alpha}[L_{\alpha}F(z_L)] \mid L_{\alpha}F(z_R) - (L_{<\alpha}F(z))^{-1}, L_{<\alpha}F(z) \} = \{ \mathbb{R}, \mathcal{L}_{\alpha}[L_{\alpha} \mathfrak{d}_{\alpha}(\Xi z)], \mathcal{L}_{\alpha}[\Xi z_L] \mid \Xi z_R - (L_{<\alpha}F(z))^{-1}, L_{<\alpha}F(z) \}.
$$

Recall that  $\Xi z$  = { $\mathbb{R}$  ,  $\mathscr{L}_{\alpha}[\Xi z_{L}]$  |  $\Xi z_{R}-(L_{<\alpha}F(z))^{-1}$ }. Therefore it suffices to show that  $\Xi z$  lies in the cut  $(\mathcal{L}_{\alpha}[L_{\alpha} \delta_{\alpha}(\Xi z)]] | L_{<\alpha} F(z)$  to conclude that  $L_{\alpha} F(z) = \Xi z$  and thus that  $F(z) = E_{\alpha} \Xi z$ . Now  $L_{\alpha} \mathfrak{d}_{\alpha}(\Xi z) < \mathcal{E}_{\alpha}^*[\Xi z]$  so  $L_{\alpha} \mathfrak{d}_{\alpha}(\Xi z) < \Xi z$  and  $\mathcal{L}_{\alpha}[L_{\alpha} \mathfrak{d}_{\alpha}(\Xi z)] < \Xi z$ . We have  $F(z) > \mathfrak{d}_{\alpha}(\Xi z)$ , where *F*(*z*)∈Mo<sub>*α*</sub>. Since *v* is a limit ordinal, Lemma [6.11](#page-21-0) implies that  $F(z) > E_{< \alpha} \Xi z$ , so  $\Xi z < L_{< \alpha} F(z)$ . This completes the proof that  $F(z) = E_\alpha \Xi z$ .

<span id="page-27-0"></span>*Corollary* 7.4*. The identities* [\(7.1\)](#page-26-0)*,* [\(7.2\)](#page-26-2), [\(7.3\)](#page-26-1)*, and* [\(7.4\)](#page-26-3) *all hold.*

*Proof*. It is enough to prove [\(7.1\)](#page-26-0) and [\(7.3\)](#page-26-1). The identity [\(7.3\)](#page-26-1) follows from [\(7.7\)](#page-27-3) and [\(7.8\)](#page-27-4). In order to obtain [\(7.1\)](#page-26-0), we consider  $\mathfrak{a} \in \mathbf{Mo}_{\alpha}$ , set  $\psi \coloneqq \{ L_{\alpha} \mathfrak{a}^{\mathbf{Mo}_{\alpha}}_{L} \mid L_{\alpha} \mathfrak{a}^{\mathbf{Mo}_{\alpha}}_{R}, L_{<\alpha} \mathfrak{a} \}_{\mathbf{No}_{>,\alpha}}$  and we show that  $a = E_\alpha \psi$ . Since [\(7.3\)](#page-26-1) is uniform, we have

$$
E_{\alpha}\psi = \{ \delta_{\alpha}(\psi), E_{\alpha}L_{\alpha} \mathfrak{a}_{L}^{\mathbf{M}\mathbf{o}_{\alpha}} | E_{\alpha}L_{\alpha} \mathfrak{a}_{R}^{\mathbf{M}\mathbf{o}_{\alpha}}, E_{\alpha}L_{<\alpha} \mathfrak{a} \}^{\mathbf{M}\mathbf{o}_{\alpha}}= \{ \delta_{\alpha}(\psi), \mathfrak{a}_{L}^{\mathbf{M}\mathbf{o}_{\alpha}} | \mathfrak{a}_{R}^{\mathbf{M}\mathbf{o}_{\alpha}}, E_{\alpha}L_{<\alpha} \mathfrak{a} \}^{\mathbf{M}\mathbf{o}_{\alpha}}= \{ \delta_{\alpha}(\psi), \mathfrak{a}_{L}^{\mathbf{M}\mathbf{o}_{\alpha}} | \mathfrak{a}_{R}^{\mathbf{M}\mathbf{o}_{\alpha}}, E_{\alpha}L_{<\alpha} \mathfrak{a} \}^{\mathbf{M}\mathbf{o}_{\alpha}} \}
$$

We have  $\delta_{\alpha}(\psi) < \alpha$  because  $\psi < L_{<\alpha} \alpha$ , and  $E_{\alpha}L_{<\alpha} \alpha > \alpha$  because  $E_{\alpha} > E_{<\alpha}$  on No<sup>>,></sup>. Since  $\alpha =$  $\{a_L^{\mathbf{M}\mathbf{o}_\alpha}\}\mathbf{M}\mathbf{o}_{\alpha}, \text{ we deduce that } E_\alpha \psi = \mathfrak{a}.$ 

*Remark* 7.5. The simplified cut equations for  $E_\alpha$ ,  $L_\alpha$  can be viewed as alternative definitions for those functions, since they hold inductively on their domain of definition. It is unclear how to develop our theory directly upon these alternative definitions. In particular, does there exists a direct way to see that the cut equation [\(7.2\)](#page-26-2) is warranted, and that the corresponding function satisfies  $\mathbf{R}_{\mu}$  $\mathbf{R}_{\mu}$  $\mathbf{R}_{\mu}$  and  $\mathbf{M}_{\mu}$  $\mathbf{M}_{\mu}$  $\mathbf{M}_{\mu}$ ?

### 7.2 Identities involving  $\text{Tr}_{\alpha}$  and  $\text{Tr}_{\alpha}^*$ .

<span id="page-28-0"></span>*PROPOSITION* 7.6*. Defining*  $\text{Tr}_{\alpha}$  =  $\text{Smp}_{\beta}$ *as in Section [6.1,](#page-16-0) we have*  $\text{Tr}_{\alpha}$  =  $\text{No}_{\geq, \alpha}$ *.* 

*Proof.* Let  $\varphi \in \mathbf{No}_{\geq \alpha}$ . We have  $E_{\alpha} \mathcal{L}_{\alpha}[\varphi] = \mathcal{E}_{\alpha}[E_{\alpha} \varphi]$  by [\[5](#page-30-2), Proposition 7.22]. Recall that  $\mathcal{E}_{\alpha}[a] = \mathcal{E}_{\alpha}'[a]$  for all  $a \in \mathbb{N}o^{>,>}$ . Now  $\mathcal{E}_{\alpha}' \circ E_{\alpha} = E_{\alpha} \circ \mathcal{L}_{\alpha}'$  by definition of  $\mathcal{L}_{\alpha}'$ , so  $E_{\alpha} \mathcal{L}_{\alpha}[\varphi] =$ *E*<sub>α</sub>  $\mathscr{L}_{\alpha}[\varphi]$  and  $\mathscr{L}_{\alpha}[\varphi] = \mathscr{L}_{\alpha}'[\varphi]$ . By definition of  $\text{Tr}_{\alpha}$ , we conclude that  $\text{Tr}_{\alpha} = \text{Smp}_{\mathscr{L}_{\alpha}} = \text{No}_{\succ,\alpha}$ .

Assume that *v* is a successor ordinal. Then we have  $N\mathbf{o}_{\geq,\alpha} = N\mathbf{o}_{\geq,\alpha} + \mathbb{R}$  by [\(4.4\)](#page-14-3), so the functions  $T_r \Xi_{NQ_{>a}}$  and  $\Xi_{NQ_{>a}} T_r$  are both strictly increasing bijections from No onto No<sub>>,*α*</sub>.

<span id="page-28-1"></span>LEMMA 7.7. Assume that v is a successor ordinal. Then for  $r \in \mathbb{R}$ , we have  $T_r \Xi_{NQ_{>0}} = \Xi_{NQ_{>0}} T_r$ *on* No*.*

*Proof.* Let us abbreviate  $\Xi = \Xi_{\mathbf{No}_{\succ,\alpha}}$ . We prove the lemma by induction on  $(\mathbf{No}, \sqsubseteq) \times (\mathbb{R}, \sqsubseteq)$ . Let  $(z, r) \in$ **No** × ℝ with

$$
\Xi y + s = \Xi (y + s)
$$

whenever  $(y, s) ∈ \mathbf{No} × \mathbb{R}$  is strictly simpler than  $(z, r)$ . We let  $z', z'', r', r''$  denote generic elements of  $z_L$ ,  $z_R$ ,  $r_L$ ,  $r_R$  and we note that  $r'$ ,  $r'' \in \mathbb{R}$ . By [\(6.8\)](#page-24-0), we have

$$
\Xi(z+r) = \left\{ \Xi(z'+r) + \frac{1}{L_{<\alpha}E_{\alpha}\Xi(z'+r)}, \ \Xi(z+r') + \frac{1}{L_{<\alpha}E_{\alpha}\Xi(z+r')} \right\}
$$

$$
\Xi(z+r'') - \frac{1}{L_{<\alpha}E_{\alpha}\Xi(z+r'')}, \ \Xi(z''+r) - \frac{1}{L_{<\alpha}E_{\alpha}\Xi(z''+r)} \right\}_{\text{No}>>}
$$

$$
= \left\{ T_r \Xi z' + \frac{1}{L_{<\alpha}E_{\alpha}T_r \Xi z'}, \ T_{r'} \Xi z + \frac{1}{L_{<\alpha}E_{\alpha}T_{r'} \Xi z} \right\}_{\text{To}>>}
$$

$$
T_{r''} \Xi z - \frac{1}{L_{<\alpha}E_{\alpha}T_{r''} \Xi z}, \ T_r \Xi z'' - \frac{1}{L_{<\alpha}E_{\alpha}T_r \Xi z''} \right\}_{\text{No}>>}
$$

Recall that *ν* is a successor ordinal. Since [\(4.2\)](#page-13-2) holds for all  $a \in \mathbf{No}^{>,>}$  the sets  $L_{<\alpha} E_\alpha \mathscr{T} a$  and  $L <sub>α</sub>E<sub>α</sub> a$  are mutually cofinal and coinitial. Moreover  $T<sub>s</sub>(z + b) = T<sub>s</sub> z + b$  for all  $s \in \mathbb{R}$  and  $b \in \mathbb{N}$ o, so

$$
\Xi(z+r) = \left\{ T_r \left( \Xi z' + \frac{1}{L_{<\alpha} E_{\alpha} \Xi z'} \right), T_{r'} \left( \Xi z + \frac{1}{L_{<\alpha} E_{\alpha} \Xi z} \right) \right\}
$$

$$
T_{r''} \left( \Xi z - \frac{1}{L_{<\alpha} E_{\alpha} \Xi z} \right), T_r \left( \Xi z'' - \frac{1}{L_{<\alpha} E_{\alpha} \Xi z''} \right) \right\}
$$
<sub>No>></sub>

The hyperserial field of surreal numbers 29

By  $(3.1)$ , we have

$$
T_r \Xi z = \left\{ T_r \left( \Xi z' + \frac{1}{L_{<\alpha} E_{\alpha} \Xi z'} \right), T_{r'} \Xi z \mid T_{r''} \Xi z, T_r \left( \Xi z'' - \frac{1}{L_{<\alpha} E_{\alpha} \Xi z''} \right) \right\}_{\mathbf{No}^{>,\geq}.
$$

The numbers  $T_r \, \Xi z$ ,  $T_{r'} \, \Xi z$  and  $T_{r''} \, \Xi z$  are  $\alpha$ -truncated so  $T_r \, \Xi z$  lies in the cut

$$
\left(\bigcup_{r'} T_{r'}\left(\Xi z + \frac{1}{L_{<\alpha}E_{\alpha}\Xi z}\right)\bigg|\bigcup_{r''} T_{r''}\left(\Xi z - \frac{1}{L_{<\alpha}E_{\alpha}\Xi z}\right)\right)_{\text{No}^{>}}.
$$

We deduce that  $T_r \Xi z = \Xi T_r z$ . The result follows by induction. □

<span id="page-29-0"></span>Lemma 7.8. *If v is a successor ordinal, then we have*  $\mathcal{T}\neq\mathscr{L}_{\alpha}^*$  *on*  $\mathbf{No}^{>,>}$ *. Consequently,*  $\mathbf{Tr}_{\alpha}^*=\mathbf{No}^{>}_{>}$ *.* 

*Proof.* The set  $E_{< \alpha}$  is pointwise cofinal in  $\mathcal{E}_\alpha^*$ . So  $L_\alpha E_{< \alpha} E_\alpha$  is pointwise cofinal in  $\mathcal{L}_\alpha^*$ . For *γ* < *α*, there is *n* ∈ ℕ such that *γ* ≤  $\alpha$ /*ωn*. We have

$$
L_{\alpha}E_{\gamma}E_{\alpha} \leq L_{\alpha}E_{\alpha_{/\omega}n}E_{\alpha} = (L_{\alpha}E_{\alpha_{/\omega}}E_{\alpha})^{\circ n} = (L_{\alpha}E_{\alpha}T_{1})^{\circ n} = T_{1}^{\circ n} = T_{n} \in \mathcal{F}.
$$

We deduce that  $\mathscr{T} \neq \mathscr{L}_{\alpha}^*$  on  $\mathbf{No}^{>,\succ}$ , whence  $\mathbf{Tr}_{\alpha}^* = \mathbf{Smp}_{\mathscr{T}} = \mathbf{No}^>_{\succ}$ .

### 7.3 Identities involving  $Mo_{\alpha}$  and  $Mo_{\alpha}^*$ .

<span id="page-29-2"></span>*Lemma* 7.9*. If v is a successor ordinal, then for*  $z \in$  **No** *we have* 

$$
\Xi_{\mathbf{Mo}_{\alpha}}(z-1) = L_{\alpha_{/\omega}} \Xi_{\mathbf{Mo}_{\alpha}} z.
$$

*Proof*. This can be seen as a converse to the proof of the identity [\(6.3\)](#page-20-2). We proceed by induction on (N**o**,  $\subseteq$ ). Let *z* be such that the relation holds on *z*<sub>⊏</sub>. By [\(6.3\)](#page-20-2), we have

$$
L_{\alpha_{/\omega}} \Xi_{\mathbf{M}\mathbf{o}_{\alpha}} z = \{L_{\alpha_{/\omega}} (\Xi_{\mathbf{M}\mathbf{o}_{\alpha}} z)_{L}^{\mathbf{M}\mathbf{o}_{\alpha}} | L_{\alpha_{/\omega}} (\Xi_{\mathbf{M}\mathbf{o}_{\alpha}} z)_{R}^{\mathbf{M}\mathbf{o}_{\alpha}}, \Xi_{\mathbf{M}\mathbf{o}_{\alpha}} z\}_{\mathbf{M}\mathbf{o}_{\alpha}}
$$
  
\n
$$
= \{L_{\alpha_{/\omega}} \Xi_{\mathbf{M}\mathbf{o}_{\alpha}} z_L | L_{\alpha_{/\omega}} \Xi_{\mathbf{M}\mathbf{o}_{\alpha}} z_R, \Xi_{\mathbf{M}\mathbf{o}_{\alpha}} z\}_{\mathbf{M}\mathbf{o}_{\alpha}}
$$
  
\n
$$
= \{\Xi_{\mathbf{M}\mathbf{o}_{\alpha}} (z_L - 1) | \Xi_{\mathbf{M}\mathbf{o}_{\alpha}} (z_R - 1), \Xi_{\mathbf{M}\mathbf{o}_{\alpha}} z\}_{\mathbf{M}\mathbf{o}_{\alpha}}
$$
 (by the inductive hypothesis)  
\n
$$
= \Xi_{\mathbf{M}\mathbf{o}_{\alpha}} \{z_L - 1 | z_R - 1, z\}
$$
  
\n
$$
= \Xi_{\mathbf{M}\mathbf{o}_{\alpha}} (z - 1) \text{ by (3.1)}.
$$

We conclude by induction.  $\Box$ 

Noting that  $E_{\alpha/\omega} = E_{\alpha} T_1 L_{\alpha}$  on  $\text{No}^{>,>}$ , the previous relation further generalizes as follows.

*Proposition* 7.10*. Assume that ν is a successor ordinal and let r* ∈ℝ*. Then*

<span id="page-29-1"></span>
$$
\Xi_{\mathbf{M}\mathbf{o}_{\alpha}}T_{r} = E_{\alpha}T_{r}L_{\alpha}\Xi_{\mathbf{M}\mathbf{o}_{\alpha}} \tag{7.9}
$$

**PROOF.** We proceed by induction. Let  $(z, r) \in \mathbf{No} \times \mathbb{R}$  be such that

<span id="page-29-3"></span>
$$
\Xi_{\mathbf{M}\mathbf{o}_{\alpha}}T_{s}\mathbf{y} = E_{\alpha}T_{s}L_{\alpha}\Xi_{\mathbf{M}\mathbf{o}_{\alpha}}\mathbf{y}
$$

for all strictly simpler  $(y, s) \in \mathbb{N} \circ \mathbb{R}$  with respect to the product order  $\subseteq \times \subseteq$ . For  $s \in \mathbb{R}$ , let  $\phi_s$ be the function  $b \mapsto E_{\alpha} T_s L_{\alpha} b$  on  $N\sigma^{>,>}$  and let  $\alpha = \Xi_{M\omega_{\alpha}} z$ . By [\(3.1\)](#page-8-1) and [\(3.2\)](#page-9-1), we have

$$
\Xi_{\mathbf{Mo}_{\alpha}}(z+r) = \{ \mathbb{R}, \mathcal{E}_{\alpha} \Xi_{\mathbf{Mo}_{\alpha}}(z_{L}+r), \mathcal{E}_{\alpha} \Xi_{\mathbf{Mo}_{\alpha}}(z+r_{L}) \mid \mathcal{E}_{\alpha} \Xi_{\mathbf{Mo}_{\alpha}}(z_{R}+r), \mathcal{E}_{\alpha} \Xi_{\mathbf{Mo}_{\alpha}}(z+r_{R}) \} \n= \{ \mathbb{R}, \mathcal{E}_{\alpha} \phi_{r}(\alpha_{L}^{\mathbf{Mo}_{\alpha}}), \mathcal{E}_{\alpha} \phi_{r}(\alpha) \mid \mathcal{E}_{\alpha} \phi_{r}(\alpha_{R}^{\mathbf{Mo}_{\alpha}}), \mathcal{E}_{\alpha} \phi_{r}(\alpha) \}.
$$

By  $(7.1)$ , Lemma [7.7](#page-28-1) and  $(3.1)$ , we have:

$$
T_rL_\alpha\mathfrak{a} = \{T_rL_\alpha\mathfrak{a}_L^{\mathbf{Mo}_\alpha}, T_{r_L}L_\alpha\mathfrak{a} \mid T_{r_R}L_\alpha\mathfrak{a}, T_rL_\alpha\mathfrak{a}_R^{\mathbf{Mo}_\alpha}, L_{<\alpha}\mathfrak{a}\}_{\mathbf{Tr}_\alpha}.
$$

<sup>&</sup>gt;. □

We deduce that

$$
\phi_r(\mathfrak{a}) = \{E_{<\alpha} T_r L_\alpha \mathfrak{a}, \mathcal{E}_\alpha \phi_r(\mathfrak{a}_L^{\mathbf{MO}\alpha}), \mathcal{E}_\alpha \phi_{r_L}(\mathfrak{a}) \mid \mathcal{E}_\alpha \phi_{r_R}(\mathfrak{a}), \mathcal{E}_\alpha \phi_r(\mathfrak{a}_R^{\mathbf{MO}\alpha}), \mathcal{E}_\alpha E_\alpha L_{<\alpha} \mathfrak{a}\}
$$
\n
$$
= \{E_{<\alpha} L_\alpha \mathfrak{a}, \mathcal{E}_\alpha \phi_r(\mathfrak{a}_L^{\mathbf{MO}\alpha}), \mathcal{E}_\alpha \phi_{r_L}(\mathfrak{a}) \mid \mathcal{E}_\alpha \phi_{r_R}(\mathfrak{a}), \mathcal{E}_\alpha \phi_r(\mathfrak{a}_R^{\mathbf{MO}\alpha}), E_\alpha L_{<\alpha} \mathfrak{a}\}.
$$

It is enough to prove that  $E_{<\alpha}L_{\alpha}$   $\alpha \leq \Xi_{\mathbf{Mo}_{\alpha}}(z+r) \leq E_{\alpha}L_{<\alpha}$  a to conclude that  $\phi_r(\mathfrak{a}) = \Xi_{\mathbf{Mo}_{\alpha}}(z+r)$ . Towards this, fix an *n*∈ℕ with −*n*⩽*r* ⩽*n*. Lemma [7.9](#page-29-2) yields

$$
\Xi_{\mathbf{Mo}_{\alpha}}(z+r) \leq \Xi_{\mathbf{Mo}_{\alpha}}(z+n) = E_{\alpha_{/\omega^n}} \mathfrak{a} < E_{\alpha} L_{<\alpha} \mathfrak{a}
$$
\n
$$
\Xi_{\mathbf{Mo}_{\alpha}}(z+r) \geq \Xi_{\mathbf{Mo}_{\alpha}}(z-n) = L_{\alpha_{/\omega^n}} \mathfrak{a} > E_{<\alpha} L_{\alpha} \mathfrak{a}.
$$

We conclude by induction that  $(7.9)$  holds.  $\Box$ 

*Remark* 7.11. For  $r, s \in \mathbb{R}$ , we have  $\phi_{r+s} = \phi_r \circ \phi_s$ , and  $\phi_1 = E_{\alpha_{/\omega}}$ . Therefore we can see  $(\phi_r)_{r \in \mathbb{R}}$ as a system of fractional and real iterates of the hyperexponential function  $E_{\alpha_{/\omega}}$  on  ${\rm No}^{>,>}$ . The previous proposition shows that the action of those iterates on  $L_{\leq \alpha}$ -atomic numbers reduces to translations, modulo the parametrization  $\Xi_{Mo_\alpha}$ . In particular, one can compute the functional square root of exp on  $Mo_{\omega}$  in terms of sign sequences using the material from [\[3\]](#page-30-8).

<span id="page-30-6"></span>*PROPOSITION 7.12. If v is a successor ordinal, then*  $M\sigma_{\alpha}^* = M\sigma_{\alpha} \cdot N\sigma_{\alpha}$ .

*Proof.* For  $\theta \in \mathbf{No}$ , we have  $\theta_L + \mathbb{N} < \theta < \theta_R - \mathbb{N}$ . By Lemma [7.9](#page-29-2), it follows that  $E_{\alpha/\alpha} \mathbb{E} \mathbf{No}$   $\theta_L <$  $\Xi_{\mathbf{M}\mathbf{o}_{\alpha}}\theta < L_{\alpha_{/\omega}N} \Xi_{\mathbf{M}\mathbf{o}_{\alpha}}\theta_R$ . This implies that  $\mathscr{E}_{\alpha}^* \Xi_{\mathbf{M}\mathbf{o}_{\alpha}}\theta_L < \Xi_{\mathbf{M}\mathbf{o}_{\alpha}}\theta < \mathscr{E}_{\alpha}^* \Xi_{\mathbf{M}\mathbf{o}_{\alpha}}\theta_R$ , so  $\Xi_{\mathbf{M}\mathbf{o}_{\alpha}}\theta$  is  $\mathscr{E}_{\alpha}^*$ simple.

Conversely, consider  $\theta \in \mathbf{No}^{\geq},\geq$  such that  $\Xi_{\mathbf{Mo}_{\alpha}}\theta$  is  $\mathscr{E}_{\alpha}^*$ -simple. We have  $\Xi_{\mathbf{Mo}_{\alpha}}\theta_L \subseteq (\Xi_{\mathbf{Mo}_{\alpha}}\theta)_L$ and  $\Xi_{\mathbf{M}\mathbf{o}_{\alpha}}\theta_R \subseteq (\Xi_{\mathbf{M}\mathbf{o}_{\alpha}}\theta)_R$ , whence  $E_{\alpha_{/\omega}N} \Xi_{\mathbf{M}\mathbf{o}_{\alpha}}\theta_L < \Xi_{\mathbf{M}\mathbf{o}_{\alpha}}\theta < L_{\alpha_{/\omega}N} \Xi_{\mathbf{M}\mathbf{o}_{\alpha}}\theta_R$ . We obtain  $\theta_L + N <$  $\theta$  < *θ*<sub>*R*</sub>− <sup>N</sup>, which proves that *θ* ∈ **No**<sub>≻</sub>. □

*Proposition* 7.13*. We have*  $E_{\alpha}$   $\mathbf{Tr}_{\alpha}^{*} = \mathbf{Mo}_{\alpha}^{*}$ .

*Proof.* Let  $\varphi \in \mathbf{Tr}^*_{\alpha}$ . So  $\varphi \in \mathbf{Tr}_{\alpha}$ . By Proposition [3.1](#page-8-3), the number  $E_{\alpha}\varphi$  is simplest in

<span id="page-30-7"></span>
$$
E_{\alpha}(\mathscr{E}_{\alpha}^{*}[\varphi] \cap \mathrm{Tr}_{\alpha}) = \mathscr{E}_{\alpha}^{*}[E_{\alpha}\varphi] \cap \mathrm{Mo}_{\alpha}.
$$

Since  $\mathbf{Mo}_{\alpha}^* \subseteq \mathbf{Mo}_{\alpha}$ , we have  $E_{\alpha} \varphi \sqsubseteq \mathcal{E}_{\alpha}^*[E_{\alpha} \varphi] \cap \mathbf{Mo}_{\alpha}^*$  so  $E_{\alpha} \varphi \sqsubseteq \mathfrak{d}_{\alpha}^*(E_{\alpha} \varphi)$ . We deduce that  $E_{\alpha} \varphi =$  $\delta^*_\alpha(E_\alpha \varphi)$ , so  $E_\alpha \varphi$  is  $\mathscr{E}^*_\alpha$ -simple. Conversely, let  $\mathfrak{a} \in \mathbf{Mo}^*_\alpha$ . By Proposition [3.1](#page-8-3) the number  $L_\alpha \mathfrak{a}$  is  $\sup_{\alpha}$  simplest in  $L_{\alpha}(\mathscr{E}_{\alpha}^{*}[a] \cap \mathbf{Mo}_{\alpha}) = \mathscr{E}_{\alpha}^{*}[L_{\alpha}a] \cap \mathbf{No}_{\alpha,\alpha}$ . Since  $\mathrm{Tr}_{\alpha}^{*} \subseteq \mathbf{No}_{\alpha,\alpha}$ , we have  $L_{\alpha}a \sqsubseteq \mathscr{E}_{\alpha}^{*}[L_{\alpha}a] \cap \mathbf{No}_{\alpha,\alpha}$  $\operatorname{Tr}^*_{\alpha}$  so  $L_{\alpha}$  a  $\subseteq$  # $^*_{\alpha}(L_{\alpha}$  a). We deduce that  $L_{\alpha}$  a  $\subseteq$  # $^*_{\alpha}(L_{\alpha}$  a) is  $\mathscr{L}^*_{\alpha}$ -simple. □

*C*OROLLARY 7.14*. If v is a successor ordinal, then*  $M\sigma_{\alpha}^{*} = E_{\alpha}N\sigma_{\beta}^{2}$ .

#### BIBLIOGRAPHY

- <span id="page-30-3"></span>[1] M. Aschenbrenner, L. van den Dries, and J. van der Hoeven. On numbers, germs, and transseries. In *Proc. Int. Cong. of Math. 2018*, volume 1, pages 1–24. Rio de Janeiro, 2018.
- <span id="page-30-0"></span>[2] M. Aschenbrenner, L. van den Dries, and J. van der Hoeven. The surreal numbers as a universal *H*-field. *J. Eur. Math. Soc.*, 21(4):1179–1199, 2019.
- <span id="page-30-8"></span>[3] V. Bagayoko. Sign sequences of log-atomic numbers. Hal-02952455, 2020.
- <span id="page-30-4"></span>[4] V. Bagayoko and J. van der Hoeven. Surreal substructures. HAL-02151377 (pre-print), 2019.
- <span id="page-30-2"></span>[5] V. Bagayoko, J. van der Hoeven, and E. Kaplan. Hyperserial fields. (notes non publiées), 2020.
- <span id="page-30-1"></span>[6] V. Bagayoko, J. van der Hoeven, and V. Mantova. Defining a surreal hyperexponential. HAL-02861485 (pre print), 2020.
- <span id="page-30-5"></span>[7] A. Berarducci. Surreal numbers, exponentiation and derivations. <https://arxiv.org/abs/2008.06878>, 08 2020.

#### The hyperserial field of surreal numbers

- <span id="page-31-20"></span>A. Berarducci, S. Kuhlmann, V. Mantova, and M. Matusinski. Exponential fields and Conway's omega-map. [8] Proceedings of the American Mathematical Society, 2019.
- <span id="page-31-3"></span>A. Berarducci and V. Mantova. Surreal numbers, derivations and transseries. *JEMS*, 20(2):339-390, 2018. [9]
- <span id="page-31-21"></span>[10] A. Berarducci and V. Mantova. Transseries as germs of surreal functions. Trans. of the AMS, 371:3549-3592, 2019.
- <span id="page-31-0"></span>[11] J. H. Conway. On numbers and games. Academic Press, 1976.
- <span id="page-31-12"></span>[12] B. I. Dahn and P. Göring. Notes on exponential-logarithmic terms. Fundamenta Mathematicae, 127:45-50, 1986.
- <span id="page-31-2"></span>[13] L. van den Dries and Ph. Ehrlich. Fields of surreal numbers and exponentiation. Fundamenta Mathematicae, 167(2):173-188, 2001.
- <span id="page-31-17"></span>[14] L. van den Dries, J. van der Hoeven, and E. Kaplan. Logarithmic hyperseries. Trans. of the AMS, 372(7):5199-5241, 2019.
- <span id="page-31-8"></span>[15] P. du Bois-Reymond. Sur la grandeur relative des infinis des fonctions. Annali di Matematica Pura ed Applicata  $(1867-1897), 4(1):338-353, 1870.$
- <span id="page-31-9"></span>[16] P. du Bois-Reymond. Über asymptotische Werte, infinitäre Approximationen und infinitäre Auflösung von Gleichungen. Math. Ann., 8:363-414, 1875.
- <span id="page-31-10"></span>[17] P. du Bois-Reymond. Über die Paradoxen des Infinitärscalcüls. Math. Ann., 11:149-167, 1877.
- <span id="page-31-13"></span>[18] J. Écalle. Introduction aux fonctions analysables et preuve constructive de la conjecture de Dulac. Hermann, collection: Actualités mathématiques, 1992.
- <span id="page-31-1"></span>[19] H. Gonshor. An Introduction to the Theory of Surreal Numbers. Cambridge Univ. Press, 1986.
- <span id="page-31-18"></span>[20] H. Hahn. Über die nichtarchimedischen Größensysteme. Sitz. Akad. Wiss. Wien, 116:601-655, 1907.
- <span id="page-31-6"></span>[21] G. H. Hardy. Orders of infinity. Cambridge Univ. Press, 1910.
- <span id="page-31-7"></span>[22] G.H. Hardy. Properties of logarithmico-exponential functions. Proceedings of the London Mathematical Society,  $10(2):54-90, 1911.$
- <span id="page-31-11"></span>[23] D. Harvey and J. van der Hoeven. Faster polynomial multiplication over finite fields using cyclotomic coefficient rings. Accepted for publication in J. of Complexity, 2019.
- <span id="page-31-14"></span>[24] I. van der Hoeven. Automatic asymptotics. PhD thesis. École polytechnique. Palaiseau. France. 1997.
- <span id="page-31-15"></span>[25] J. van der Hoeven. Transséries fortement monotones. Chapter 1 of unpublished CNRS activity report, https:// www.texmacs.org/joris/schmeling/rap1-2000.pdf.2000.
- <span id="page-31-5"></span>[26] J. van der Hoeven. Transseries and real differential algebra, volume 1888 of Lecture Notes in Mathematics. Springer-Verlag, 2006.
- <span id="page-31-4"></span>[27] J. van der Hoeven. The Jolly Writer. Your Guide to GNU TeXmacs. Scypress, 2020.
- <span id="page-31-19"></span>[28] V. Mantova and M. Matusinski. Surreal numbers with derivation, Hardy fields and transseries: a survey. Contemporary Mathematics, pages 265-290, 2017.
- <span id="page-31-16"></span>[29] M. C. Schmeling. Corps de transséries. PhD thesis, Université Paris-VII, 2001.

### **GLOSSARY**



