



HAL
open science

PreCEE (Pregnancy and Combined Environmental Exposure), a research program in two middle-sized cities - Part 2, Results and discussion

Frédéric Mauny, Alice Brembilla, Sophie Pujol, Paul Sagot, Didier Riethmuller, Gérard Thiriez, Hélène Houot, Anne-Sophie Mariet, Alice Barba-Vasseur, Jérôme Defrance, et al.

► To cite this version:

Frédéric Mauny, Alice Brembilla, Sophie Pujol, Paul Sagot, Didier Riethmuller, et al.. PreCEE (Pregnancy and Combined Environmental Exposure), a research program in two middle-sized cities - Part 2, Results and discussion. Forum Acusticum, Dec 2020, Lyon, France. pp.3353-3355, 10.48465/fa.2020.1107 . hal-03233779

HAL Id: hal-03233779

<https://hal.science/hal-03233779>

Submitted on 26 May 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

PreCEE (PREGNANCY AND COMBINED ENVIRONMENTAL EXPOSURE) A RESEARCH PROGRAM IN TWO MIDDLE-SIZED CITIES - PART 2, RESULTS AND DISCUSSION

Frédéric Mauny^{1,2,3} Alice Brembilla^{1,2,3} Sophie Pujol^{1,2,3} Paul Sagot⁴
Didier Riethmuller¹ Gérard Thiriez¹ Hélène Houot⁵
Anne-Sophie Mariet^{2,4} Marie Barba-Vasseur^{1,2} Jérôme Defrance⁶ Vinh-Phuc Luu¹
Eric Benzenine⁴ Catherine Quantin⁴ Nadine Bernard²

¹Centre hospitalier universitaire de Besançon, Besançon, France

²UMR Chrono-environnement, CNRS/université Bourgogne Franche-Comté, Besançon, France

³INSERM CIC 1431, uMETH, Besançon, France

⁴Centre Hospitalier Universitaire de Dijon, Dijon, France

⁵UMR ThéMA, CNRS /université Bourgogne Franche-Comté, Besançon, France

⁶Centre Scientifique et Technique du Bâtiment (CSTB), 24 rue Joseph Fourier, 38400 Saint Martin D'Hères, France

Correspondence frederic.mauny@univ-fcomte.fr

ABSTRACT

The PreCEE (Pregnancy And Combined Environmental Exposure) research program aims to assess the relationships between environmental exposure to noise and air pollution and the occurrence of adverse pregnancy outcomes (APO). Data from nearly 11000 deliveries that occurred between 2005 and 2009 at the Besançon or Dijon (France) were collected, including medical, obstetrical, socio-economic and environmental conditions. Air pollution, proximity to green spaces and noise exposure were quantified at the living neighbourhood scale of each residential building. A deprivation index was calculated for the two cities. Several approaches were conducted to investigate the relationships between the occurrence of APO (preterm birth, fetal growth disorders, hypertensive disorders) and the urban environment of two middle-sized cities. The relationship between APO and the accumulation of vulnerability markers during pregnancy (medico-obstetrical, demographic, behavioral and environmental dimensions) was also investigated. The risk of APO increased significantly with the cumulative number of vulnerabilities.

In women with uncomplicated pregnancy, no significant differences in pollutant exposure levels were found between cases and controls.

In pregnancies with associated comorbidities, low levels of air pollution do not seem to affect preterm delivery. However, moderate noise exposure could potentially affect pregnancy when associated with vulnerability factors.

pregnancy outcomes (APO) are associated with many and various factors related to the course of pregnancy, the mother's characteristics, and her socio-economic and professional conditions [3]. Beyond the well-known risk factors, additional environmental factors, such as air pollution or noise, are suspected to play a role.

In urban areas, the density of pollution sources combined with a high number of residents, constitute the optimal conditions for simultaneous exposure of pregnant women to many environmental pollutants. The link between environment and pregnancy outcome has been highlighted over the past 10 years. Most of studies on air pollution exposure to carbon monoxide (CO), nitrogen dioxide (NO₂) or particulate matter (PM₁₀, PM_{2.5}) reported a higher risk of fetal growth restriction [4–7]. Effects on preterm birth appear to be more discussed [4–6,8]. Recent studies have suspected adverse associations with environmental noise, especially for low birth weight [9,10]. A benefic effect of maternal proximity to green spaces on birth weight was highlighted by few studies [11,12].

The aim of the PreCEE (Pregnancy And Combined Environmental Exposure) research program is to study the influence of environmental exposures to noise and air pollution on the occurrence of APO. More specifically, preterm birth, fetal growth disorders (fetal growth restriction / small for gestational age), hypertensive disorders of pregnancy, and gestational diabetes will be investigated.

1. INTRODUCTION

The occurrence of complications during pregnancy - such as hypertensive disorders, premature delivery, fetal growth disorders - can have serious consequences on the health of the mother and / or the newborn, in particular on children's physical and intellectual development [1-2]. Adverse

2. POPULATION AND METHODS

This section is described in the previous paper (FA2020/1106): PreCEE (Pregnancy and Combined Environmental Exposure), a research program in two middle-sized cities - Part 1, Population and methods. Logistic regression analyses were performed using classical and multilevel models. Many sensitivity analyses

were conducted to explore the potential influence of the retained definition of exposure (pollutant, living area, time exposure window), adverse pregnancy outcomes (birth weight standards...) and missing data treatment.

2.1 Vulnerability accumulation during pregnancy

A first approach based on the accumulation of vulnerability markers during pregnancy was conducted. Fifteen vulnerability markers regrouped in six dimensions (maternal age, smoking, body mass index, socio-economic, medico-obstetrical and environmental vulnerabilities) were analyzed in comparison with two APO (preterm birth, low birth weight) and two pregnancy complications (preeclampsia and vaginal bleeding during the second and third pregnancy trimesters).

2.2 Preterm birth

Two case-control studies were conducted on preterm delivery on single pregnancies. First, only pregnancies without any associated comorbidities were considered (N=1511). Secondly, only pregnancies with associated comorbidities were considered (N=2503).

3. RESULTS

3.1 Vulnerability accumulation during pregnancy

Among the 3686 women included in this approach, 21% were not exposed to any marker and 19% accumulated three or more markers among the six dimensions. The risk of APO increased significantly with the cumulative number of vulnerabilities.

3.2 Case-control studies

3.2.1 Pregnancies without any associated comorbidities

The correlation between noise and NO₂ indices ranged from 0.41 to 0.59. The mean NO₂ level ranged between 24 and 25 µg/m³ and the mean L_{night} ranged between 51 and 52 dB(A). No significant differences in pollutant exposure levels were found between cases and controls in this study. The adjusted odds ratios (OR) ranged between 0.96 and 1.08. Sensitivity analysis conducted using different temporal and spatial exposure windows demonstrated the same results.

3.2.2 Pregnancies with associated comorbidities

The OR associated with NO₂ > 40µg/m³ was 0.83 (95%CI: 0.61-1.13) and the OR associated with L_{night} ≥ 55dB(A) was 1.23 (95%CI: 0.97-1.56). A strong and monotonic association between the cumulative number of comorbidities during pregnancy and preterm delivery was also identified.

4. DISCUSSION

By combining medical, behavioral, socio-economic and environmental markers, this study shows that vulnerabilities tend to accumulate in a part of the pregnant women. This accumulation seems to represent a particular risk of adverse pregnancy outcome or complication during pregnancy. This study confirms the need to identify vulnerable women as early as possible during pregnancy to adapt their pregnancy monitoring to avoid complications.

Low levels of air pollution do not seem to affect preterm delivery. However, moderate noise exposure could potentially affect pregnancy when associated with vulnerability factors.

5. REFERENCES

- [1] C. Osmond & D. J. Barker: Fetal, infant, and childhood growth are predictors of coronary heart disease, diabetes, and hypertension in adult men and women. *Environ. Health Perspect.* 108(Suppl 3), 545–553 (2000).
- [2] S. R. Dalziel, V. Parag, A. Rodgers, & J. E. Harding: Cardiovascular risk factors at age 30 following preterm birth. *Int. J. Epidemiol.* 36, 907–915 (2007).
- [3] A. Brembilla, N. Bernard, S. Pujol, A.L. Parmentier, A. Eckman, A.S. Mariet, H. Houot, Q. Tenailleau, G. Thiriez, D. Riethmuller, M. Barba-Vasseur, F. Mauny. Pregnancy vulnerability in urban areas: a pragmatic approach combining behavioral, medico-obstetrical, socio-economic and environmental factors. *Sci Rep.* 2019 Dec 11;9(1):18878. doi: 10.1038/s41598-019-55005-0 (2019)
- [4] Shah, P. S. & Balkhair, T. & Knowledge Synthesis Group on Determinants of Preterm/LBW births. Air pollution and birth outcomes: a systematic review. *Environ. Int.* 37, 498–516 (2011).
- [5] Stieb, D. M., Chen, L., Eshoul, M. & Judek, S. Ambient air pollution, birth weight and preterm birth: a systematic review and metaanalysis. *Environ. Res.* 117, 100–111 (2012).
- [6] Nieuwenhuijsen, M. J., Dadvand, P., Grellier, J., Martinez, D. & Vrijheid, M. Environmental risk factors of pregnancy outcomes: a summary of recent meta-analyses of epidemiological studies. *Environ. Health Glob. Access Sci. Source* 12, 6 (2013).
- [7] Mariet, A.-S. et al. Multiple pregnancies and air pollution in moderately polluted cities: Is there an association between air pollution and fetal growth? *Environ. Int.* 121, 890–897 (2018).
- [8] Barba-Vasseur, M. et al. Does low to moderate environmental exposure to noise and air pollution

influence preterm delivery in medium-sized cities? *Int. J. Epidemiol.* 46, 2017–2027 (2017).

- [9] Ristovska, G., Laszlo, H. E. & Hansell, A. L. Reproductive outcomes associated with noise exposure - a systematic review of the literature. *Int. J. Environ. Res. Public. Health* 11, 7931–7952 (2014).
- [10] Nieuwenhuijsen, M. J., Ristovska, G. & Dadvand, P. WHO Environmental Noise Guidelines for the European Region: A Systematic Review on Environmental Noise and Adverse Birth Outcomes. *Int. J. Environ. Res. Public. Health* 14 (2017).
- [11] Agay-Shay, K. et al. Green spaces and adverse pregnancy outcomes. *Occup. Environ. Med.* 71, 562–569 (2014).
- [12] Dadvand, P. et al. Green space, health inequality and pregnancy. *Environ. Int.* 40, 110–115 (2012).