



**HAL**  
open science

# Bridging the Multiscale Hybrid-Mixed and Multiscale Hybrid High-Order methods

Théophile Chaumont-Frelet, Alexandre Ern, Simon Lemaire, Frédéric Valentin

► **To cite this version:**

Théophile Chaumont-Frelet, Alexandre Ern, Simon Lemaire, Frédéric Valentin. Bridging the Multiscale Hybrid-Mixed and Multiscale Hybrid High-Order methods. 2021. hal-03235525v1

**HAL Id: hal-03235525**

**<https://hal.science/hal-03235525v1>**

Preprint submitted on 25 May 2021 (v1), last revised 7 Jan 2022 (v3)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# BRIDGING THE MULTISCALE HYBRID-MIXED AND MULTISCALE HYBRID HIGH-ORDER METHODS

THÉOPHILE CHAUMONT-FRELET<sup>†</sup>, ALEXANDRE ERN<sup>‡</sup>, SIMON LEMAIRE<sup>♯</sup>,  
AND FRÉDÉRIC VALENTIN<sup>‡</sup>

ABSTRACT. We establish the equivalence between the Multiscale Hybrid-Mixed (MHM) and the Multiscale Hybrid High-Order (MsHHO) methods for a variable diffusion problem with piecewise polynomial source terms. Under the idealized assumption that the local problems defining the multiscale basis functions are exactly solved, we prove that the equivalence holds for general polytopal (coarse) meshes and arbitrary approximation orders. Also, we leverage the interchange of properties to perform a unified convergence analysis, as well as to improve on both methods.

## 1. INTRODUCTION

The tremendous development of massively parallel architectures in the last decade has led to a revision of what is expected from computational simulators, which must embed asynchronous and communication-avoiding algorithms. In such a scenario where precision and robustness remain fundamental properties, but algorithms must take full advantage of the new architectures, numerical methods built upon the "divide-and-conquer" philosophy fulfill simulator imperatives better than standard methods operating in a monolithic fashion on the different scales of the problem at hand. Among the vast literature on the subject, driven by domain decomposition methodologies (see, e.g., [40] for a survey), multiscale numerical methods emerge as an attractive option to efficiently handle problems with highly heterogeneous coefficients, as well as multi-query scenarios in which the problem solution must be computed for a large number of source terms. These scenarios may arise when considering highly oscillatory, nonlinear, time-dependent models, or within optimization algorithms when solving problems featuring PDE-based constraints, or in models including stochastic processes, to cite a few.

The development of multiscale methods started with the seminal work [6]. Important advances were then provided in [31, 32] (cf. also [8, 10], and the unifying viewpoint of [9]) and in [28, 29], laying the ground, respectively, for the Variational Multiscale method, and for the Multiscale Finite Element Method (MsFEM). Overall, the common idea behind these multiscale methods is to consider basis functions especially designed so as to upscale to an overlying coarse mesh the sub-mesh variations of the model. Particularly appealing is the fact that the multiscale basis functions are defined by entirely independent problems. From this viewpoint, multiscale numerical methods may also be seen as a (non-iterative) domain decomposition technique [24]. Since the pioneering works on multiscale methods, a large number of improvements and new approaches have been proposed. In the MsFEM context (see [21] for a survey), one can cite the oversampling technique of [22], as well as the Petrov–Galerkin variant of [30] (see also [3]), the high-order method of [2], or the multipoint flux mixed finite element method of [42]. More recent research directions focus on reducing and possibly eliminating the cell resonance error. In that vein, one can cite the Generalized MsFEM [20], or the Local Orthogonal Decomposition approach [27, 36]. Based on a quite different paradigm, one can also cite the Heterogeneous Multiscale Method [19, 1].

---

(†) INRIA, UNIV. CÔTE D'AZUR, CNRS, UMR 7351 - LABORATOIRE J. A. DIEUDONNÉ, F-06000 NICE, FRANCE

(‡) CERMICS, ÉCOLE DES PONTS, F-77455 MARNE-LA-VALLÉE CEDEX 2 & INRIA PARIS, F-75589 PARIS, FRANCE

(♯) INRIA, UNIV. LILLE, CNRS, UMR 8524 - LABORATOIRE PAUL PAINLEVÉ, F-59000 LILLE, FRANCE

(‡) LNCC, PETRÓPOLIS - RJ, BRAZIL & INRIA, F-06000 NICE, FRANCE

*E-mail addresses:* (†) [theophile.chaumont@inria.fr](mailto:theophile.chaumont@inria.fr), (‡) [alexandre.ern@enpc.fr](mailto:alexandre.ern@enpc.fr), (♯) [simon.lemaire@inria.fr](mailto:simon.lemaire@inria.fr), (‡) [valentin@lncc.br](mailto:valentin@lncc.br).

Hybridization has also been used recently in the context of (multiscale) Discontinuous Galerkin methods, leading to the multiscale Hybridizable Discontinuous Galerkin method of [23]; cf. also the Weak Galerkin method of [37], devised in the spirit of the Generalized MsFEM.

Recently, two families of hybrid multiscale numerical methods that are applicable on general meshes have been proposed, namely the Multiscale Hybrid-Mixed (MHM) and the Multiscale Hybrid High-Order (MsHHO) methods. The MHM method has been first introduced in [25], and further analyzed in [4, 38, 7] (see also [26] for an abstract setting), whereas the MsHHO method has been proposed in [12, 13], as an extension of the HHO method first introduced in [16, 15] (cf. also [17]). The MHM method relates to the MsFEM proposed in [11], as well as to the subgrid upscaling method of [5] (see [26, Section 5.1.2] for further details). The MsHHO method generalizes to arbitrary polynomial orders the low-order nonconforming multiscale methods of [33, 34]. Note in passing that similar ideas have also been developed in a conforming framework in the context of BEM-based FEM [14, 41]. The MHM and MsHHO methods substantially differ in their construction. Picking the Poisson equation as an example, the MHM method hinges on the primal hybrid formulation analyzed in [39]. As a consequence, whereas the local problems are defined as coercive Neumann problems, the global upscaled linear system is of saddle-point type, involving face unknowns that are the normal fluxes through the mesh faces (also the Neumann data for the local problems, up to the sign), plus one degree of freedom per mesh cell that enforces a local balance between the normal fluxes and the source term. Notice that the (global) saddle-point structure of the MHM method can be equivalently replaced by a sequence of positive-definite linear systems as shown recently in [35]. On the other hand, the MsHHO method is directly built upon the primal formulation of the problem. As a consequence, the local (Neumann) problems are defined as constrained minimization problems, and as such exhibit a saddle-point structure. At the opposite, the global upscaled linear system is coercive, and only involves face unknowns that are the traces of the solution at interfaces. Notice that, as opposed to the MHM method, the MsHHO method also uses cell unknowns (that are locally eliminable from the global upscaled linear system), which are associated with basis functions solving local problems with nonzero source terms. As such, the MsHHO method is naturally suited to deal with multi-query scenarios.

In this work, we revisit the MHM and MsHHO methods and we prove an equivalence result between their solutions. Notice that such a relationship is not straightforward since, at first glance, the two methods exhibit structures that are genuinely different. Nonetheless, we demonstrate that such an equivalence holds under the assumption that the source term of the continuous problem is piecewise polynomial (cf. Theorem 5.1). For this equivalence to hold, we make the idealized assumption that the local problems defining the multiscale basis functions are exactly solved. The corresponding methods are then referred to as *one-level*. Leveraging this equivalence result, the present work also contributes to derive, in a unified fashion, an energy-norm error estimate that is valid for both methods (cf. Theorem 6.3). More specifically,

- in the MHM framework, this result is a refined version (especially in the tracking of the dependency with respect to the diffusion coefficient) of the results in [4];
- in the MsHHO framework, this result is new and is complementary to the homogenization-based error estimate derived in [12].

We also explore these stimulating results to transfer properties proved for one method to the other, and to reveal how the interplay between the methods can guide advances for both. Notably, we show that

- the MHM method can be adapted to deal with multi-query scenarios (cf. Section 7.2.1);
- the MsHHO method can be recast as a purely face-based method, in the sense that it can be alternatively defined without using cell unknowns (cf. Remark 7.5).

The outline of the article is as follows. Section 2 introduces the model problem, the partition, the notation and a number of useful tools. We present the MHM method in Section 3, and the MsHHO method in Section 4. The equivalence result is stated in Section 5, along with some further properties and remarks. The energy-norm error estimate is proved in Section 6. The solution strategies for both methods are discussed in Section 7, leveraging the equivalence result

at hand to propose enhancements for both methods. Finally, some conclusions are drawn in Section 8.

## 2. SETTING

In this section, we present the setting, introduce the partition, and define useful broken spaces on this partition.

**2.1. Model problem.** We consider an open polytopal domain  $\Omega \subset \mathbb{R}^d$ ,  $d = 2$  or  $3$ , with boundary  $\partial\Omega$ . Given  $f : \Omega \rightarrow \mathbb{R}$ , we seek a function  $u : \Omega \rightarrow \mathbb{R}$  such that

$$(2.1) \quad \begin{cases} -\nabla \cdot (\mathbb{A} \nabla u) = f & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

We assume that the diffusion coefficient  $\mathbb{A} \in L^\infty(\Omega; \mathbb{R}^{d \times d})$  is symmetric and uniformly elliptic, and that the source term  $f$  is in  $L^2(\Omega)$ . Problem (2.1) can be equivalently recast into the following weak form: find  $u \in H_0^1(\Omega)$  such that

$$(2.2) \quad (\mathbb{A} \nabla u, \nabla v)_\Omega = (f, v)_\Omega \quad \text{for all } v \in H_0^1(\Omega),$$

where  $(\cdot, \cdot)_D$  denotes the  $L^2(D; \mathbb{R}^\ell)$ ,  $\ell \in \{1, d\}$ , inner product for any measurable set  $D \subset \bar{\Omega}$ . It is well-established that Problem (2.2) admits a unique solution.

**2.2. Partition.** The domain  $\Omega$  is partitioned into a (coarse) mesh  $\mathcal{T}_H$  that consists of polytopal (open) cells  $K$  with diameter  $H_K$ , and we set  $H := \max_{K \in \mathcal{T}_H} H_K$ . In practice, both the MHM and MsHHO methods consider a fine submesh (characterized by a mesh-size  $h \ll H$ ) to compute the local basis functions, but this finer mesh is not needed in the present discussion since we will assume that the local problems defining the basis functions are exactly solved. We collect the mesh faces  $F$  of  $\mathcal{T}_H$  in the set  $\mathcal{F}_H$ , and split this set into the subset of internal faces (or interfaces)  $\mathcal{F}_H^{\text{int}}$  and the subset of boundary faces  $\mathcal{F}_H^{\text{bnd}}$ . The mesh faces are defined to be planar, i.e., every mesh face  $F \in \mathcal{F}_H$  is supported by an affine hyperplane  $\mathcal{H}_F$  (recall that the mesh cells have planar faces since they are polytopes). For a face  $F \in \mathcal{F}_H^{\text{int}}$ , we have

$$(2.3) \quad F = \partial K_+ \cap \partial K_- \cap \mathcal{H}_F,$$

for two cells  $K_\pm \in \mathcal{T}_H$ , whereas for a boundary face  $F \in \mathcal{F}_H^{\text{bnd}}$ , we have

$$(2.4) \quad F = \partial K \cap \partial\Omega \cap \mathcal{H}_F,$$

for a cell  $K \in \mathcal{T}_H$ . For simplicity, the definitions (2.3) and (2.4) do not allow, respectively, for the case of several coplanar faces that would be shared by the same two cells, or for the case of several coplanar boundary faces in a boundary cell, but of course these cases could be seamlessly handled. We denote by  $\partial\mathcal{T}_H$  the skeleton of the mesh  $\mathcal{T}_H$ , defined by  $\partial\mathcal{T}_H := \bigcup_{K \in \mathcal{T}_H} \partial K$ . Given  $K \in \mathcal{T}_H$ , we denote by  $\mathcal{F}_K$  the set of its faces, and by  $\mathbf{n}_K$  the unit outward normal vector to its boundary (whose restriction to the face  $F \in \mathcal{F}_K$  is the constant vector denoted by  $\mathbf{n}_{K,F}$ ). We associate with each face  $F \in \mathcal{F}_H$  a unit normal vector  $\mathbf{n}_F$  whose orientation is fixed, with the convention that  $\mathbf{n}_F := \mathbf{n}_{\Omega|F}$  if  $F \in \mathcal{F}_H^{\text{bnd}}$ , where  $\mathbf{n}_\Omega$  is the unit outward normal vector to  $\partial\Omega$ .

**2.3. Infinite-dimensional broken spaces.** We first define the broken space of piecewise smooth functions on  $\mathcal{T}_H$ :

$$(2.5) \quad H^1(\mathcal{T}_H) := \{v \in L^2(\Omega) : v_K \in H^1(K) \quad \forall K \in \mathcal{T}_H\},$$

where we let  $v_D := v|_D$ . For any  $v \in H^1(\mathcal{T}_H)$ , we define the jump  $[[v]]_F$  of  $v$  across  $F \in \mathcal{F}_H$  by

$$(2.6) \quad [[v]]_F := v_{K_+|F} (\mathbf{n}_{K_+,F} \cdot \mathbf{n}_F) + v_{K_-|F} (\mathbf{n}_{K_-,F} \cdot \mathbf{n}_F)$$

if  $F \subseteq \partial K_+ \cap \partial K_-$  is an interface, and simply by

$$(2.7) \quad [[v]]_F := v_{K|F}$$

if  $F \subseteq \partial K \cap \partial\Omega$  is a boundary face. We also define the broken gradient operator  $\nabla_H : H^1(\mathcal{T}_H) \rightarrow L^2(\Omega; \mathbb{R}^d)$  such that, for any  $v \in H^1(\mathcal{T}_H)$ ,

$$(2.8) \quad (\nabla_H v)|_K := \nabla v_K \quad \text{for all } K \in \mathcal{T}_H.$$

We next introduce the space of piecewise smooth functions on  $\mathcal{T}_H$  whose broken (weighted) flux belongs to  $\mathbf{H}(\text{div}, \Omega)$ :

$$(2.9) \quad \mathcal{V}(\mathcal{T}_H; \text{div}, \Omega) := \{v \in H^1(\mathcal{T}_H) : \mathbb{A} \nabla_H v \in \mathbf{H}(\text{div}, \Omega)\}.$$

Notice that  $\mathcal{V}(\mathcal{T}_H; \text{div}, \Omega) \not\subset H^1(\Omega)$ . We now define the two ‘‘skeletal’’ spaces

$$(2.10) \quad \Sigma_0(\partial\mathcal{T}_H) := \left\{ z := (z_{\partial K})_{K \in \mathcal{T}_H} \in \prod_{K \in \mathcal{T}_H} H^{1/2}(\partial K) \mid \begin{array}{l} \exists w(z) \in H_0^1(\Omega) \text{ s.t.} \\ z_{\partial K} = w_K(z)|_{\partial K} \quad \forall K \in \mathcal{T}_H \end{array} \right\},$$

and

$$(2.11) \quad \Lambda(\partial\mathcal{T}_H) := \left\{ \mu := (\mu_{\partial K})_{K \in \mathcal{T}_H} \in \prod_{K \in \mathcal{T}_H} H^{-1/2}(\partial K) \mid \begin{array}{l} \exists \boldsymbol{\sigma}(\mu) \in \mathbf{H}(\text{div}, \Omega) \text{ s.t.} \\ \mu_{\partial K} = \boldsymbol{\sigma}_K(\mu)|_{\partial K} \cdot \mathbf{n}_K \quad \forall K \in \mathcal{T}_H \end{array} \right\}.$$

Letting  $\langle \cdot, \cdot \rangle_{\partial K}$  stand for the duality pairing between  $H^{-1/2}(\partial K)$  and  $H^{1/2}(\partial K)$ , we define the following pairing, for all  $\mu \in \prod_{K \in \mathcal{T}_H} H^{-1/2}(\partial K)$  and all  $z \in \prod_{K \in \mathcal{T}_H} H^{1/2}(\partial K)$ ,

$$(2.12) \quad \langle \mu, z \rangle_{\partial\mathcal{T}_H} := \sum_{K \in \mathcal{T}_H} \langle \mu_{\partial K}, z_{\partial K} \rangle_{\partial K},$$

so that for all  $\mu \in \Lambda(\partial\mathcal{T}_H)$  and all  $z \in \Sigma_0(\partial\mathcal{T}_H)$ , recalling that  $\boldsymbol{\sigma}(\mu) \in \mathbf{H}(\text{div}, \Omega)$  and  $w(z) \in H_0^1(\Omega)$ , we have

$$(2.13) \quad \langle \mu, z \rangle_{\partial\mathcal{T}_H} = \sum_{K \in \mathcal{T}_H} \left( (\nabla \cdot \boldsymbol{\sigma}(\mu), w(z))_K + (\boldsymbol{\sigma}(\mu), \nabla w(z))_K \right) = 0.$$

**2.4. Finite-dimensional broken spaces.** Let  $q \in \mathbb{N}$  denote a given polynomial degree. The space of piecewise ( $d$ -variate) polynomial functions on  $\mathcal{T}_H$  of total degree up to  $q$  is denoted by

$$(2.14) \quad \mathbb{P}^q(\mathcal{T}_H) := \{v \in L^2(\Omega) : v_K \in \mathbb{P}^q(K) \quad \forall K \in \mathcal{T}_H\},$$

whereas the space of piecewise ( $(d-1)$ -variate) polynomial functions on  $\mathcal{F}_H$  of total degree up to  $q$  is denoted by

$$(2.15) \quad \mathbb{P}^q(\mathcal{F}_H) := \left\{ v \in L^2 \left( \bigcup_{F \in \mathcal{F}_H} \overline{F} \right) : v_F \in \mathbb{P}^q(F) \quad \forall F \in \mathcal{F}_H \right\},$$

and its subset incorporating homogeneous boundary conditions by

$$(2.16) \quad \mathbb{P}_0^q(\mathcal{F}_H) := \{v \in \mathbb{P}^q(\mathcal{F}_H) : v_F = 0 \quad \forall F \in \mathcal{F}_H^{\text{bnd}}\}.$$

For all  $K \in \mathcal{T}_H$ , we also define the local space of piecewise ( $(d-1)$ -variate) polynomial functions on  $\mathcal{F}_K$  of total degree up to  $q$  as follows:

$$(2.17) \quad \mathbb{P}^q(\mathcal{F}_K) := \{v \in L^2(\partial K) : v_F \in \mathbb{P}^q(F) \quad \forall F \in \mathcal{F}_K\}.$$

We consider the following finite-dimensional subspace of  $\Lambda(\partial\mathcal{T}_H)$ :

$$(2.18) \quad \Lambda^q(\partial\mathcal{T}_H) := \{\mu \in \Lambda(\partial\mathcal{T}_H) : \mu_{\partial K} \in \mathbb{P}^q(\mathcal{F}_K) \quad \forall K \in \mathcal{T}_H\} \subsetneq \Lambda(\partial\mathcal{T}_H).$$

Notice that for every interface  $F \in \mathcal{F}_H^{\text{int}}$  with  $F \subseteq \partial K_+ \cap \partial K_-$ , as a consequence of (2.13), we have  $\mu_{\partial K_+|F} + \mu_{\partial K_-|F} = 0$  for all  $\mu \in \Lambda^q(\partial\mathcal{T}_H)$ . We also define, for any integer  $m \geq 0$ , the spaces

$$(2.19) \quad \begin{cases} \mathcal{U}^{m,q}(K) := \{v \in H^1(K) : \nabla \cdot (\mathbb{A} \nabla v) \in \mathbb{P}^m(K), \quad \mathbb{A} \nabla v|_{\partial K} \cdot \mathbf{n}_K \in \mathbb{P}^q(\mathcal{F}_K)\}, \quad \forall K \in \mathcal{T}_H, \\ \mathcal{U}^{m,q}(\mathcal{T}_H) := \{v \in H^1(\mathcal{T}_H) : v_K \in \mathcal{U}^{m,q}(K) \quad \forall K \in \mathcal{T}_H\}. \end{cases}$$

To alleviate the notation, we shall drop the superscript  $m$  when considering  $m = q - 1$  for  $q \geq 1$ , and write  $\mathcal{U}^q(K)$  and  $\mathcal{U}^q(\mathcal{T}_H)$  in place of  $\mathcal{U}^{q-1,q}(K)$  and  $\mathcal{U}^{q-1,q}(\mathcal{T}_H)$ , respectively.

We finally introduce the space of ‘‘discretely  $H_0^1(\Omega)$ ’’ functions on  $\mathcal{T}_H$ :

$$(2.20) \quad \tilde{H}_0^{1,q}(\mathcal{T}_H) := \{v \in H^1(\mathcal{T}_H) : ([v]_F, p)_F = 0 \quad \forall p \in \mathbb{P}^q(F), \quad \forall F \in \mathcal{F}_H\}.$$

Equivalently, we have

$$(2.21) \quad \tilde{H}_0^{1,q}(\mathcal{T}_H) = \{v \in H^1(\mathcal{T}_H) : \langle \mu, v \rangle_{\partial\mathcal{T}_H} = 0 \quad \forall \mu \in \Lambda^q(\partial\mathcal{T}_H)\}.$$

### 3. THE MHM METHOD

Let us first set

$$(3.1) \quad \begin{cases} H^1(K)^\perp := \{v \in H^1(K) : (v, 1)_K = 0\}, & \forall K \in \mathcal{T}_H, \\ H^1(\mathcal{T}_H)^\perp := \{v \in H^1(\mathcal{T}_H) : (v_K, 1)_K = 0 \quad \forall K \in \mathcal{T}_H\}. \end{cases}$$

For integers  $m, q \in \mathbb{N}$ , we also define the subspaces  $\mathcal{U}^{m,q}(K)^\perp := \{v \in \mathcal{U}^{m,q}(K) : (v, 1)_K = 0\}$  for all  $K \in \mathcal{T}_H$  and  $\mathcal{U}^{m,q}(\mathcal{T}_H)^\perp := \{v \in \mathcal{U}^{m,q}(\mathcal{T}_H) : (v_K, 1)_K = 0 \quad \forall K \in \mathcal{T}_H\}$ .

Let  $K \in \mathcal{T}_H$ , and consider the two local operators

$$(3.2) \quad T_K^N : H^{-\frac{1}{2}}(\partial K) \rightarrow H^1(K)^\perp, \quad T_K^S : L^2(K) \rightarrow H^1(K)^\perp.$$

For all  $\mu_{\partial K} \in H^{-\frac{1}{2}}(\partial K)$  and all  $g_K \in L^2(K)$ ,  $T_K^N(\mu_{\partial K})$  and  $T_K^S(g_K)$  are the unique elements in  $H^1(K)^\perp$  such that

$$(3.3) \quad \begin{cases} (\mathbb{A} \nabla T_K^N(\mu_{\partial K}), \nabla v)_K = \langle \mu_{\partial K}, v \rangle_{\partial K}, \\ (\mathbb{A} \nabla T_K^S(g_K), \nabla v)_K = (g_K, v)_K, \end{cases} \quad \forall v \in H^1(K)^\perp.$$

The superscripts in the operators indicate that  $T_K^N$  lifts a (Neumann) normal flux and  $T_K^S$  lifts a source term. Elementary arguments show that

$$(3.4a) \quad -\nabla \cdot (\mathbb{A} \nabla T_K^N(\mu_{\partial K})) = -\frac{1}{|K|} \langle \mu_{\partial K}, 1 \rangle_{\partial K} \text{ in } K, \quad \mathbb{A} \nabla T_K^N(\mu_{\partial K}) \cdot \mathbf{n}_K = \mu_{\partial K} \text{ on } \partial K,$$

$$(3.4b) \quad -\nabla \cdot (\mathbb{A} \nabla T_K^S(g_K)) = g_K - \frac{1}{|K|} (g_K, 1)_K \text{ in } K, \quad \mathbb{A} \nabla T_K^S(g_K) \cdot \mathbf{n}_K = 0 \text{ on } \partial K.$$

It is convenient to define the following global versions of the above lifting operators:

$$(3.5) \quad T^N : \Lambda(\partial\mathcal{T}_H) \rightarrow H^1(\mathcal{T}_H)^\perp, \quad T^S : L^2(\Omega) \rightarrow H^1(\mathcal{T}_H)^\perp.$$

For all  $\mu \in \Lambda(\partial\mathcal{T}_H)$  and all  $g \in L^2(\Omega)$ , we set

$$(3.6) \quad T^N(\mu)|_K := T_K^N(\mu_{\partial K}), \quad T^S(g)|_K := T_K^S(g_K).$$

Equivalently, we have

$$(3.7) \quad \begin{cases} (\mathbb{A} \nabla_H T^N(\mu), \nabla_H v)_\Omega = \langle \mu, v \rangle_{\partial\mathcal{T}_H}, \\ (\mathbb{A} \nabla_H T^S(g), \nabla_H v)_\Omega = (g, v)_\Omega, \end{cases} \quad \forall v \in H^1(\mathcal{T}_H)^\perp,$$

which results from summing (3.3) cell-wise. We remark that the solution  $u \in H_0^1(\Omega)$  to Problem (2.2) satisfies

$$(3.8) \quad u = u^0 + T^N(\lambda) + T^S(f),$$

where  $(u^0, \lambda) \in \mathbb{P}^0(\mathcal{T}_H) \times \Lambda(\partial\mathcal{T}_H)$  solve

$$(3.9a) \quad \langle \lambda, v^0 \rangle_{\partial\mathcal{T}_H} = -(f, v^0)_\Omega \quad \forall v^0 \in \mathbb{P}^0(\mathcal{T}_H),$$

$$(3.9b) \quad \langle \mu, u^0 \rangle_{\partial\mathcal{T}_H} + \langle \mu, T^N(\lambda) \rangle_{\partial\mathcal{T}_H} = -\langle \mu, T^S(f) \rangle_{\partial\mathcal{T}_H} \quad \forall \mu \in \Lambda(\partial\mathcal{T}_H).$$

Notice that, owing to (3.7) and to the fact that  $\mathbb{A}$  is symmetric, we have  $\langle \mu, T^S(f) \rangle_{\partial\mathcal{T}_H} = (f, T^N(\mu))_\Omega$ .

Let  $k \in \mathbb{N}$  be a given polynomial degree. The MHM method [4] reads as follows: Find  $(u_H^0, \lambda_H) \in \mathbb{P}^0(\mathcal{T}_H) \times \Lambda^k(\partial\mathcal{T}_H)$  such that

$$(3.10a) \quad \langle \lambda_H, v_H^0 \rangle_{\partial\mathcal{T}_H} = -(f, v_H^0)_\Omega \quad \forall v_H^0 \in \mathbb{P}^0(\mathcal{T}_H),$$

$$(3.10b) \quad \langle \mu_H, u_H^0 \rangle_{\partial\mathcal{T}_H} + \langle \mu_H, T^N(\lambda_H) \rangle_{\partial\mathcal{T}_H} = -\langle \mu_H, T^S(f) \rangle_{\partial\mathcal{T}_H} \quad \forall \mu_H \in \Lambda^k(\partial\mathcal{T}_H),$$

and the MHM solution is then defined by

$$(3.11) \quad u_H^{\text{MHM}} := u_H^0 + T^N(\lambda_H) + T^S(f).$$

The well-posedness of Problem (3.10) is established in [4, Theorem 3.2]. Notice that we also have, on the discrete level,  $\langle \mu_H, T^s(f) \rangle_{\partial \mathcal{T}_H} = (f, T^N(\mu_H))_\Omega$ .

**Lemma 3.1** (Characterization of the MHM solution (3.11)). *Let  $u_H^{\text{MHM}}$  be defined by (3.11). Then,* (i)  $(\mathbb{A} \nabla_H u_H^{\text{MHM}}|_{\partial K}) \cdot \mathbf{n}_K \in \mathbb{P}^k(\mathcal{F}_K)$  for all  $K \in \mathcal{T}_H$  and  $u_H^{\text{MHM}} \in \tilde{H}_0^{1,k}(\mathcal{T}_H)$ ; (ii)  $u_H^{\text{MHM}} \in \mathcal{V}(\mathcal{T}_H; \text{div}, \Omega)$  and  $-\nabla \cdot (\mathbb{A} \nabla_H u_H^{\text{MHM}}) = f$  in  $\Omega$ .

*Proof.* By (3.11) and (3.4), we infer that for all  $K \in \mathcal{T}_H$ ,

$$(3.12) \quad \mathbb{A} \nabla_H u_H^{\text{MHM}}|_{\partial K} \cdot \mathbf{n}_K = \mathbb{A} \nabla T_K^N(\lambda_H|_{\partial K}) \cdot \mathbf{n}_K + \mathbb{A} \nabla T_K^s(f_K) \cdot \mathbf{n}_K = \lambda_H|_{\partial K} \in \mathbb{P}^k(\mathcal{F}_K).$$

That  $u_H^{\text{MHM}} \in \tilde{H}_0^{1,k}(\mathcal{T}_H)$  follows from the characterization (2.21) of  $\tilde{H}_0^{1,k}(\mathcal{T}_H)$  and (3.10b). Now, to prove that  $u_H^{\text{MHM}} \in \mathcal{V}(\mathcal{T}_H; \text{div}, \Omega)$ , we need to show that  $\mathbb{A} \nabla_H u_H^{\text{MHM}} \in \mathbf{H}(\text{div}, \Omega)$ . Owing to (3.4), we infer that for all  $K \in \mathcal{T}_H$ , we have

$$(3.13) \quad \begin{aligned} \nabla \cdot (\mathbb{A} \nabla_H u_H^{\text{MHM}})|_K &= \nabla \cdot (\mathbb{A} \nabla T_K^N(\lambda_H|_{\partial K})) + \nabla \cdot (\mathbb{A} \nabla T_K^s(f_K)) \\ &= \frac{1}{|K|} (\lambda_{\partial K}, 1)_{\partial K} - f_K + \frac{1}{|K|} (f_K, 1)_K = -f_K \in L^2(K), \end{aligned}$$

where the last equality follows from (3.10a). This shows that  $\mathbb{A} \nabla_H u_H^{\text{MHM}}|_K \in \mathbf{H}(\text{div}, K)$  for all  $K \in \mathcal{T}_H$ . Moreover, (3.12) shows that  $\mathbb{A} \nabla_H u_H^{\text{MHM}}|_{\partial K} \cdot \mathbf{n}_K$  can be localized to each face of  $K$  and since for every interface  $F \subseteq \partial K_+ \cap \partial K_-$ ,  $\lambda_{\partial K_+|F} + \lambda_{\partial K_-|F} = 0$ , we infer that  $[\mathbb{A} \nabla_H u_H^{\text{MHM}}]_F \cdot \mathbf{n}_F = 0$  on  $F$ . It results that  $\mathbb{A} \nabla_H u_H^{\text{MHM}} \in \mathbf{H}(\text{div}, \Omega)$ . Finally,  $-\nabla \cdot (\mathbb{A} \nabla_H u_H^{\text{MHM}}) = f$  in  $\Omega$  follows from (3.13) since  $K \in \mathcal{T}_H$  is arbitrary.  $\square$

Let us take a closer look at the MHM method (3.10)-(3.11). First, we observe that since  $T^N(\lambda_H) \in \mathcal{U}^{0,k}(\mathcal{T}_H)^\perp$ , this function is computable from a finite-dimensional calculation. The same holds for the right-hand side of (3.10b) since  $\langle \mu_H, T^s(f) \rangle_{\partial \mathcal{T}_H} = (f, T^N(\mu_H))_\Omega$ . However, the situation is different in (3.11) for  $T^s(f)$ . One needs indeed to define, so as to fully explicit the (one-level) method, an approximation of this function that is also computable from a finite-dimensional calculation. For this reason, the original MHM method defined by (3.10)-(3.11) can be viewed as *semi-explicit*, whereas a *fully explicit* version of it is obtained after approximating  $T^s(f)$ . Among various possibilities, perhaps the simplest one is to choose an integer  $m \geq 0$ , project  $f \in L^2(\Omega)$  onto the finite-dimensional subspace  $\mathbb{P}^m(\mathcal{T}_H)$ , and compute  $T^s(\Pi_H^m(f))$ , where  $\Pi_H^m$  is the  $L^2$ -orthogonal projector onto  $\mathbb{P}^m(\mathcal{T}_H)$ . This leads to the fully explicit MHM solution

$$(3.14) \quad u_H^{\text{MHM}} := u_H^0 + T^N(\lambda_H) + T^s(\Pi_H^m(f)),$$

where  $(u_H^0, \lambda_H) \in \mathbb{P}^0(\mathcal{T}_H) \times \Lambda^k(\partial \mathcal{T}_H)$  now solve

$$(3.15a) \quad \langle \lambda_H, v_H^0 \rangle_{\partial \mathcal{T}_H} = -(f, v_H^0)_\Omega \quad \forall v_H^0 \in \mathbb{P}^0(\mathcal{T}_H),$$

$$(3.15b) \quad \langle \mu_H, u_H^0 \rangle_{\partial \mathcal{T}_H} + \langle \mu_H, T^N(\lambda_H) \rangle_{\partial \mathcal{T}_H} = -(\Pi_H^m(f), T^N(\mu_H))_\Omega \quad \forall \mu_H \in \Lambda^k(\partial \mathcal{T}_H).$$

We notice in particular that in (3.14) we have  $T^N(\lambda_H) \in \mathcal{U}^{0,k}(\mathcal{T}_H)^\perp \subseteq \mathcal{U}^{m,k}(\mathcal{T}_H)^\perp$  and  $T^s(\Pi_H^m(f)) \in \mathcal{U}^{m,0}(\mathcal{T}_H)^\perp \subseteq \mathcal{U}^{m,k}(\mathcal{T}_H)^\perp$ . Thus, all the quantities involved in (3.14)-(3.15) are members of the space  $\mathcal{U}^{m,k}(\mathcal{T}_H)$ . We refer to Remark 5.3 for an example of an alternative definition for the fully explicit method. In what follows, we exclusively focus on (the semi-explicit method (3.10)-(3.11) and on) the fully explicit method defined by (3.14)-(3.15). Adapting the arguments of the proof of Lemma 3.1 leads to the following result.

**Lemma 3.2** (Characterization of the MHM solution (3.14)). *Let  $u_H^{\text{MHM}}$  be defined by (3.14). Then,* (i)  $u_H^{\text{MHM}} \in \mathcal{U}^{m,k}(\mathcal{T}_H) \cap \tilde{H}_0^{1,k}(\mathcal{T}_H)$ ; (ii)  $u_H^{\text{MHM}} \in \mathcal{V}(\mathcal{T}_H; \text{div}, \Omega)$  and  $-\nabla \cdot (\mathbb{A} \nabla_H u_H^{\text{MHM}}) = \Pi_H^m(f)$  in  $\Omega$ .

#### 4. THE MSHHO METHOD

Let again  $k \in \mathbb{N}$  be a given polynomial degree, and let  $m \geq 0$  be an integer. The MSHHO method hinges on the following set of discrete unknowns:

$$(4.1) \quad \hat{U}_H^{m,k} := \mathbb{P}^m(\mathcal{T}_H) \times \mathbb{P}^k(\mathcal{F}_H),$$

which is composed of cell and face degrees of freedom (one can also consider the case  $m = -1$ , so that the method is based on face unknowns only; cf. Remark 5.4). The standard MSHHO method,

referred to as mixed-order MsHHO method in [12], corresponds to the case  $m = k - 1$  for  $k \geq 1$ . For all  $K \in \mathcal{T}_H$ , we let  $\widehat{v}_K := (v_K, v_{\mathcal{F}_K}) \in \widehat{U}_K^{m,k} := \mathbb{P}^m(K) \times \mathbb{P}^k(\mathcal{F}_K)$  denote the local counterpart of  $\widehat{v}_H := (v_{\mathcal{T}_H}, v_{\mathcal{F}_H}) \in \widehat{U}_H^{m,k}$ . For all  $F \in \mathcal{F}_H$ ,  $v_F \in \mathbb{P}^k(F)$  is defined by  $v_F := v_{\mathcal{F}_H|F}$ . Notice that  $v_F = v_{\mathcal{F}_{K_+}|F} = v_{\mathcal{F}_{K_-}|F}$  if  $F \subseteq \partial K_+ \cap \partial K_-$  is an interface, and  $v_F = v_{\mathcal{F}_K|F}$  if  $F \subseteq \partial K \cap \partial \Omega$  is a boundary face.

The MsHHO method is based on the following local reconstruction operator: For all  $K \in \mathcal{T}_H$  and all  $\widehat{v}_K \in \widehat{U}_K^{m,k}$ , there exists a unique function  $r_K(\widehat{v}_K) \in \mathcal{U}^{m,k}(K)$  (recall that  $\mathcal{U}^{m,k}(K)$  is defined in (2.19)) such that

$$(4.2a) \quad (\mathbb{A} \nabla r_K(\widehat{v}_K), \nabla w)_K = -(v_K, \nabla \cdot (\mathbb{A} \nabla w))_K + (v_{\mathcal{F}_K}, \mathbb{A} \nabla w \cdot \mathbf{n}_K)_{\partial K} \quad \forall w \in \mathcal{U}^{m,k}(K),$$

$$(4.2b) \quad (r_K(\widehat{v}_K), 1)_{\partial K} = (v_{\mathcal{F}_K}, 1)_{\partial K}.$$

Notice that the usual choice of closure relation for  $r_K(\widehat{v}_K)$  is  $(r_K(\widehat{v}_K), 1)_K = (v_K, 1)_K$ . The operator  $r_K$  is the (local) reconstruction operator associated with the finite element

$$(4.3) \quad \left( K, \mathcal{U}^{m,k}(K), \widehat{\Sigma}_K \right),$$

with the set of degrees of freedom  $\widehat{\Sigma}_K : \mathcal{U}^{m,k}(K) \rightarrow \widehat{U}_K^{m,k}$  such that  $\widehat{\Sigma}_K(v) := (\Pi_K^m(v), \Pi_{\mathcal{F}_K}^k(v))$  for all  $v \in \mathcal{U}^{m,k}(K)$ , where  $\Pi_K^m$  and  $\Pi_{\mathcal{F}_K}^k$  are the  $L^2$ -orthogonal projectors onto, respectively,  $\mathbb{P}^m(K)$  and  $\mathbb{P}^k(\mathcal{F}_K)$ . For further use, we also define  $\Pi_F^k$  to be the  $L^2$ -orthogonal projector onto  $\mathbb{P}^k(F)$  for all  $F \in \mathcal{F}_H$ . Importantly, the reconstruction operator  $r_K$  satisfies  $\widehat{\Sigma}_K(r_K(\widehat{v}_K)) = \widehat{v}_K$  for all  $\widehat{v}_K \in \widehat{U}_K^{m,k}$ , i.e.,

$$(4.4a) \quad (r_K(\widehat{v}_K), q)_K = (v_K, q)_K \quad \forall q \in \mathbb{P}^m(K),$$

$$(4.4b) \quad (r_K(\widehat{v}_K), q)_{\partial K} = (v_{\mathcal{F}_K}, q)_{\partial K} \quad \forall q \in \mathbb{P}^k(\mathcal{F}_K).$$

In the MsHHO method, the essential boundary conditions can be enforced strongly by considering the subspace

$$(4.5) \quad \widehat{U}_{H,0}^{m,k} := \mathbb{P}^m(\mathcal{T}_H) \times \mathbb{P}_0^k(\mathcal{F}_H).$$

The MsHHO method for Problem (2.2) reads as follows: Find  $\widehat{u}_H \in \widehat{U}_{H,0}^{m,k}$  such that

$$(4.6) \quad \sum_{K \in \mathcal{T}_H} (\mathbb{A} \nabla r_K(\widehat{u}_K), \nabla r_K(\widehat{v}_K))_K = \sum_{K \in \mathcal{T}_H} (f_K, v_K)_K \quad \forall \widehat{v}_H \in \widehat{U}_{H,0}^{m,k}.$$

The approximate MsHHO solution  $u_H^{\text{HHO}} \in \mathcal{U}^{m,k}(\mathcal{T}_H)$  is then defined by:

$$(4.7) \quad u_H^{\text{HHO}}|_K := r_K(\widehat{u}_K) \quad \forall K \in \mathcal{T}_H.$$

The existence and uniqueness of  $\widehat{u}_H \in \widehat{U}_{H,0}^{m,k}$  solution to (4.6) is established in [12, Lemma 5.3] in the case  $m = k - 1$  for  $k \geq 1$ , but the arguments directly extend to a generic  $m \in \mathbb{N}$ . Moreover, it is easy to see that the function  $u_H^{\text{HHO}}$  defined in (4.7) actually sits in  $\widetilde{H}_0^{1,k}(\mathcal{T}_H)$ . Indeed, owing to (4.4b), for any interface  $F \in \mathcal{F}_H^{\text{int}}$  such that  $F \subseteq \partial K_+ \cap \partial K_-$ , one has for all  $q \in \mathbb{P}^k(F)$ ,

$$\begin{aligned} (\llbracket u_H^{\text{HHO}} \rrbracket_F, q)_F &= (r_{K_+}(\widehat{u}_{K_+}) (\mathbf{n}_{K_+,F} \cdot \mathbf{n}_F), q)_F + (r_{K_-}(\widehat{u}_{K_-}) (\mathbf{n}_{K_-,F} \cdot \mathbf{n}_F), q)_F \\ &= (u_{\mathcal{F}_{K_+}} (\mathbf{n}_{K_+,F} \cdot \mathbf{n}_F), q)_F + (u_{\mathcal{F}_{K_-}} (\mathbf{n}_{K_-,F} \cdot \mathbf{n}_F), q)_F \\ &= (u_F (\mathbf{n}_{K_+,F} \cdot \mathbf{n}_F), q)_F + (u_F (\mathbf{n}_{K_-,F} \cdot \mathbf{n}_F), q)_F = 0. \end{aligned}$$

For boundary faces, one uses again (4.4b) along with the fact that  $\widehat{u}_H \in \widehat{U}_{H,0}^{m,k}$ . A crucial observation made in [12, Remark 5.4] is that the MsHHO method can be equivalently reformulated as follows: Find  $u_H^{\text{HHO}} \in \mathcal{U}^{m,k}(\mathcal{T}_H) \cap \widetilde{H}_0^{1,k}(\mathcal{T}_H)$  such that

$$(4.8) \quad (\mathbb{A} \nabla_H u_H^{\text{HHO}}, \nabla_H v_H)_\Omega = (\Pi_H^m(f), v_H)_\Omega \quad \forall v_H \in \mathcal{U}^{m,k}(\mathcal{T}_H) \cap \widetilde{H}_0^{1,k}(\mathcal{T}_H),$$

where, for any  $K \in \mathcal{T}_H$ ,  $\Pi_H^m(f)|_K := \Pi_K^m(f_K)$ .

**Lemma 4.1** (Characterization of the MsHHO solution). *Let  $u_H^{\text{HHO}}$  solve (4.8). Then, (i)  $u_H^{\text{HHO}} \in \mathcal{U}^{m,k}(\mathcal{T}_H) \cap \widetilde{H}_0^{1,k}(\mathcal{T}_H)$ ; (ii)  $u_H^{\text{HHO}} \in \mathcal{V}(\mathcal{T}_H; \text{div}, \Omega)$  and  $-\nabla \cdot (\mathbb{A} \nabla_H u_H^{\text{HHO}}) = \Pi_H^m(f)$  in  $\Omega$ .*



*Proof.* We have already shown above that  $u_H^{\text{HHO}} \in \mathcal{U}^{m,k}(\mathcal{T}_H) \cap \tilde{H}_0^{1,k}(\mathcal{T}_H)$ . Let us now show that  $\mathbb{A}\nabla_H u_H^{\text{HHO}} \in \mathbf{H}(\text{div}, \Omega)$ . Since  $u_H^{\text{HHO}} \in \mathcal{U}^{m,k}(\mathcal{T}_H)$ , we already know that  $\nabla \cdot (\mathbb{A}\nabla_H u_H^{\text{HHO}})|_K \in \mathbb{P}^m(K) \subset L^2(K)$  and  $\mathbb{A}\nabla_H u_H^{\text{HHO}}|_{\partial K} \cdot \mathbf{n}_K \in \mathbb{P}^k(\mathcal{F}_K)$  for all  $K \in \mathcal{T}_H$ . Moreover, owing to (4.6), (4.7), and the definition (4.2), we infer that

$$(4.9) \quad - \sum_{K \in \mathcal{T}_H} (\nabla \cdot (\mathbb{A}\nabla_H u_H^{\text{HHO}}), v_K)_K + \sum_{F \in \mathcal{F}_H^{\text{int}}} ([\mathbb{A}\nabla_H u_H^{\text{HHO}}]_F \cdot \mathbf{n}_F, v_F)_F = \sum_{K \in \mathcal{T}_H} (\Pi_K^m(f_K), v_K)_K,$$

for all  $v_K \in \mathbb{P}^m(K)$  and all  $K \in \mathcal{T}_H$ , and for all  $v_F \in \mathbb{P}^k(F)$  and all  $F \in \mathcal{F}_H^{\text{int}}$  (notice that we have used that  $v_F = 0$  for all  $F \in \mathcal{F}_H^{\text{bnd}}$  for  $\hat{v}_H \in \hat{U}_{H,0}^{m,k}$ ). This readily implies that

$$-\nabla \cdot (\mathbb{A}\nabla_H u_H^{\text{HHO}})|_K = \Pi_K^m(f_K) \quad \text{for all } K \in \mathcal{T}_H,$$

and that

$$[\mathbb{A}\nabla_H u_H^{\text{HHO}}]_F \cdot \mathbf{n}_F = 0 \quad \text{for all } F \in \mathcal{F}_H^{\text{int}}.$$

It follows that  $\mathbb{A}\nabla_H u_H^{\text{HHO}} \in \mathbf{H}(\text{div}, \Omega)$  and that  $-\nabla \cdot (\mathbb{A}\nabla_H u_H^{\text{HHO}}) = \Pi_H^m(f)$  in  $\Omega$ .  $\square$

## 5. MAIN EQUIVALENCE RESULT AND FURTHER COMMENTS

The following result, which is a consequence of Lemma 3.1, Lemma 3.2, and Lemma 4.1, summarizes our main result on the equivalence between the MHM and MsHHO methods.

**Theorem 5.1** (Equivalence between MHM and MsHHO). *Let  $m, k \in \mathbb{N}$ . The following holds true:*

- (i) *Let  $u_H^{\text{MHM}}$  be the (original, semi-explicit) MHM solution defined by (3.11) using  $k \geq 0$ . Let  $u_H^{\text{HHO}}$  be the MsHHO solution solving (4.8) using  $m, k \geq 0$ . Then,  $u_H^{\text{MHM}} = u_H^{\text{HHO}}$  if  $f \in \mathbb{P}^m(\mathcal{T}_H)$ .*
- (ii) *Let  $u_H^{\text{MHM}}$  be the (fully explicit) MHM solution defined by (3.14) using  $m, k \geq 0$ . Let  $u_H^{\text{HHO}}$  be the MsHHO solution solving (4.8) using  $m, k \geq 0$ . Then,  $u_H^{\text{MHM}} = u_H^{\text{HHO}}$  for all  $f \in L^2(\Omega)$ .*

We now collect several remarks providing further insight into the above equivalence result.

**Remark 5.2** (Comparison of heuristic viewpoints). *Using the infinite-dimensional spaces introduced in Section 2.3, it is possible to sketch the two complementary visions behind the MHM and MsHHO methods. In the MHM method, the general thread is to search for the exact solution among the members of the affine functional space*

$$\{\bar{u} \in \mathcal{V}(\mathcal{T}_H; \text{div}, \Omega) : -\nabla \cdot (\mathbb{A}\nabla_H \bar{u}) = f \quad \text{in } \Omega\},$$

and to enforce that  $\bar{u} \in H_0^1(\Omega)$  by requiring that

$$\langle \mu, \bar{u} \rangle_{\partial \mathcal{T}_H} = 0 \quad \text{for all } \mu \in \Lambda(\partial \mathcal{T}_H).$$

In the MsHHO method, the general thread is to search for the exact solution among the members of the affine functional space

$$\{\bar{u} \in H_0^1(\Omega) : -\nabla \cdot (\mathbb{A}\nabla_H \bar{u}_K) = f_K \quad \text{in } K \quad \forall K \in \mathcal{T}_H\},$$

and to enforce that  $\bar{u} \in \mathcal{V}(\mathcal{T}_H; \text{div}, \Omega)$  by requiring that

$$\langle \mathbb{A}\nabla_H \bar{u} \cdot \mathbf{n}, z \rangle_{\partial \mathcal{T}_H} = 0 \quad \text{for all } z \in \Sigma_0(\partial \mathcal{T}_H).$$

**Remark 5.3** (Modification of the right-hand side). *It is observed in [12, Remark 5.8] that a variant of the MsHHO method is obtained by searching  $u_H^{\text{HHO}} \in \mathcal{U}^{m,k}(\mathcal{T}_H) \cap \tilde{H}_0^{1,k}(\mathcal{T}_H)$  such that*

$$(5.1) \quad (\mathbb{A}\nabla_H u_H^{\text{HHO}}, \nabla_H v_H)_\Omega = (f, v_H)_\Omega \quad \forall v_H \in \mathcal{U}^{m,k}(\mathcal{T}_H) \cap \tilde{H}_0^{1,k}(\mathcal{T}_H).$$

One advantage of (5.1) is that the source term  $f$  is now seen through its  $L^2$ -orthogonal projection onto  $\mathcal{U}^{m,k}(\mathcal{T}_H)$  instead of its projection onto the smaller space  $\mathbb{P}^m(\mathcal{T}_H)$  as in (4.8). However, if  $u_H^{\text{HHO}}$  solves (5.1),  $\mathbb{A}\nabla_H u_H^{\text{HHO}}$  slightly departs from  $\mathbf{H}(\text{div}, \Omega)$ , i.e., we no longer have  $u_H^{\text{HHO}} \in \mathcal{V}(\mathcal{T}_H; \text{div}, \Omega)$  as for the solution to (4.8). This modified MsHHO solution can be bridged to the fully explicit MHM solution obtained by approximating the lifting  $T^s$  by the operator  $T_H^s : L^2(\Omega) \rightarrow \mathcal{U}^{m,k}(\mathcal{T}_H)^\perp$  such that, for all  $g \in L^2(\Omega)$ ,  $T_H^s(g) \in \mathcal{U}^{m,k}(\mathcal{T}_H)^\perp$  solves

$$(\mathbb{A}\nabla_H T_H^s(g), \nabla_H v)_\Omega = (g, v), \quad \forall v \in \mathcal{U}^{m,k}(\mathcal{T}_H)^\perp.$$

Indeed, the modified MsHHO solution solving (5.1) coincides with the fully explicit MHM solution

$$u_H^{\text{MHM}} := u_H^0 + T^{\text{N}}(\lambda_H) + T_H^{\text{S}}(f),$$

where  $(u_H^0, \lambda_H) \in \mathbb{P}^0(\mathcal{T}_H) \times \Lambda^k(\partial\mathcal{T}_H)$  now solve

$$\begin{aligned} \langle \lambda_H, v_H^0 \rangle_{\partial\mathcal{T}_H} &= -(f, v_H^0)_\Omega & \forall v_H^0 \in \mathbb{P}^0(\mathcal{T}_H), \\ \langle \mu_H, u_H^0 \rangle_{\partial\mathcal{T}_H} + \langle \mu_H, T^{\text{N}}(\lambda_H) \rangle_{\partial\mathcal{T}_H} &= -\langle \mu_H, T_H^{\text{S}}(f) \rangle_{\partial\mathcal{T}_H} & \forall \mu_H \in \Lambda^k(\partial\mathcal{T}_H). \end{aligned}$$

**Remark 5.4** (Variant with no cell unknowns (case  $m = -1$ )). *It is possible to consider the case  $m = -1$  in the above MHM and MsHHO settings, leading to a MsHHO formulation without cell unknowns. The spaces  $\mathcal{U}^{m,q}(K)$  and  $\mathcal{U}^{m,q}(\mathcal{T}_H)$  can still be defined by (2.19) when  $m = -1$ , with the convention that  $\mathbb{P}^{-1}(K) := \{0\}$ . The fully explicit MHM method is still defined as in Section 3. An important modification in the analysis is that the last statement in Lemma 3.2 now becomes  $-\nabla \cdot (\mathbb{A} \nabla_H u_H^{\text{MHM}}) = \Pi_H^0(f)$  in  $\Omega$ . Notice also that (3.14) becomes  $u_H^{\text{MHM}} = u_H^0 + T^{\text{N}}(\lambda_H)$  and that the right-hand side of (3.15b) is zero. Actually, since  $T^{\text{S}}(c_H) = 0$  for any  $c_H \in \mathbb{P}^0(\mathcal{T}_H)$  owing to (3.4b), we infer that the (fully explicit) MHM method for  $m = -1$  coincides with the (fully explicit) MHM method for  $m = 0$ . Concerning the MsHHO method, the variant (5.1) has to be adopted in the case  $m = -1$ . Finally, we observe that in the case  $m = -1$ , the MHM and MsHHO solutions do not coincide.*

## 6. UNIFIED CONVERGENCE ANALYSIS

We derive, in a unified fashion, an energy-norm error estimate that is valid for both the (fully explicit) MHM and MsHHO methods.

**6.1. Setting.** Let  $\mathcal{T}_H$  be a given (coarse) polytopal mesh of the domain  $\Omega$  in the sense of Section 2.2. Since we are interested in deriving a quantitative estimate on the discretization error for the MHM/MsHHO methods, we need to define a measure of regularity for the mesh at hand. To do so, and as classical in the HHO context, we assume that the mesh  $\mathcal{T}_H$  admits a matching simplicial submesh  $\mathcal{S}_H$ , and that there exists some real parameter  $0 < \rho_H < 1$  such that, for all  $K \in \mathcal{T}_H$ , and all  $T \in \mathcal{S}_H$  such that  $T \subseteq K$ , (i)  $\rho_H H_T \leq R_T$  where  $R_T$  denotes the inradius of the simplex  $T$ , and (ii)  $\rho_H H_K \leq H_T$ . The parameter  $\rho_H$  measures the regularity of the mesh  $\mathcal{T}_H$ . When studying a convergence process in which the meshes of some given sequence  $(\mathcal{T}_H)_{H \in \mathcal{H}}$  are successively refined, we shall assume that the mesh sequence  $(\mathcal{T}_H)_{H \in \mathcal{H}}$  is uniformly regular, in the sense that there exists  $0 < \rho < 1$  such that, for all  $H \in \mathcal{H}$ ,  $\rho \leq \rho_H$ . Standard local Poincaré-Steklov and (continuous) trace and inverse inequalities, as well as (optimal) approximation properties for local  $L^2$ -orthogonal projectors, then hold on each cell  $K \in \mathcal{T}_H$  for any  $H \in \mathcal{H}$ , with multiplicative constants only depending on  $\rho$ . In what follows, we use the symbol  $\lesssim$  to denote an inequality that is valid up to a multiplicative constant only depending on the discretization through the parameter  $\rho$ .

In order to track the dependency of the error estimates with respect to the diffusion coefficient, for any  $K \in \mathcal{T}_H$ , we denote by  $a_{b,K} > 0$  the local smallest eigenvalue of the coefficient  $\mathbb{A}$  in the cell  $K$ , in such a way that  $\mathbb{A}(\mathbf{x})\boldsymbol{\xi} \cdot \boldsymbol{\xi} \geq a_{b,K} |\boldsymbol{\xi}|^2$  for all  $\boldsymbol{\xi} \in \mathbb{R}^d$  and almost every  $\mathbf{x} \in K$ .

Finally, given any measurable set  $D \subset \bar{\Omega}$ , and any integer  $s \geq 0$ , we respectively denote by  $|\cdot|_{s,D}$  and  $\|\cdot\|_{s,D}$  the standard seminorm and norm in  $H^s(D; \mathbb{R}^\ell)$ , for  $\ell \in \{1, d\}$ . We also define  $H^s(\mathcal{T}_H; \mathbb{R}^\ell)$  as the space of piecewise  $\mathbb{R}^\ell$ -valued  $H^s$  functions on the partition  $\mathcal{T}_H$ , with the convention that  $H^s(\mathcal{T}_H; \mathbb{R})$  is simply noted  $H^s(\mathcal{T}_H)$ .

**6.2. Local approximation.** Let  $m, k \in \mathbb{N}$  be given. Let  $K \in \mathcal{T}_H$ , and recall the definition (2.19) of the space  $\mathcal{U}^{m,k}(K)$ .

**Lemma 6.1** (Approximation in  $\mathcal{U}^{m,k}(K)$ ). *Let  $v \in H^1(K)$ , and set  $g := -\nabla \cdot (\mathbb{A} \nabla v)$  in  $K$ . Assume that  $g \in H^{m+1}(K)$  and that  $\mathbb{A} \nabla v \in H^{k+1}(K; \mathbb{R}^d)$ . There exists  $\pi_K^{m,k}(v) \in \mathcal{U}^{m,k}(K)$  such that*

$$(6.1) \quad \|\mathbb{A}^{1/2} \nabla (v - \pi_K^{m,k}(v))\|_{0,K} \lesssim a_{b,K}^{-1/2} (H_K^{m+2} |g|_{m+1,K} + H_K^{k+1} |\mathbb{A} \nabla v|_{k+1,K}).$$

*Proof.* Define  $\pi_K^{m,k}(v) \in \mathcal{U}^{m,k}(K)$  such that

$$(6.2) \quad -\nabla \cdot (\mathbb{A} \nabla \pi_K^{m,k}(v)) = \Pi_K^m(g) \quad \text{in } K, \quad \mathbb{A} \nabla \pi_K^{m,k}(v) \cdot \mathbf{n}_K = \Pi_{\mathcal{F}_K}^k(\mathbb{A} \nabla v \cdot \mathbf{n}_K) \quad \text{on } \partial K.$$

Since  $g = -\nabla \cdot (\mathbb{A} \nabla v)$ , we easily check that  $(\Pi_K^m(g), 1)_K + (\Pi_{\mathcal{F}_K}^k(\mathbb{A} \nabla v \cdot \mathbf{n}_K), 1)_{\partial K} = 0$ ; hence, the data of the Neumann problem (6.2) are compatible, and  $\pi_K^{m,k}(v)$  is well-defined (up to an additive constant). Multiplying (6.2) by  $w \in H^1(K)$ , integrating by parts, and using the compatibility of the data, yields

$$(6.3) \quad \begin{aligned} (\mathbb{A} \nabla \pi_K^{m,k}(v), \nabla w)_K &= (\Pi_K^m(g), w)_K + (\Pi_{\mathcal{F}_K}^k(\mathbb{A} \nabla v \cdot \mathbf{n}_K), w)_{\partial K} \\ &= (\Pi_K^m(g), w - \Pi_K^0(w))_K + (\Pi_{\mathcal{F}_K}^k(\mathbb{A} \nabla v \cdot \mathbf{n}_K), w - \Pi_K^0(w))_{\partial K}. \end{aligned}$$

By definition of  $g$ , we also have

$$(6.4) \quad \begin{aligned} (\mathbb{A} \nabla v, \nabla w)_K &= (g, w)_K + (\mathbb{A} \nabla v \cdot \mathbf{n}_K, w)_{\partial K} \\ &= (g, w - \Pi_K^0(w))_K + (\mathbb{A} \nabla v \cdot \mathbf{n}_K, w - \Pi_K^0(w))_{\partial K}. \end{aligned}$$

Subtracting (6.3) to (6.4), we obtain, for any  $w \in H^1(K)$ ,

$$(6.5) \quad \begin{aligned} (\mathbb{A} \nabla(v - \pi_K^{m,k}(v)), \nabla w)_K &= (g - \Pi_K^m(g), w - \Pi_K^0(w))_K \\ &\quad + (\mathbb{A} \nabla v \cdot \mathbf{n}_K - \Pi_{\mathcal{F}_K}^k(\mathbb{A} \nabla v \cdot \mathbf{n}_K), w - \Pi_K^0(w))_{\partial K}. \end{aligned}$$

Applying the Cauchy–Schwarz inequality together with a local Poincaré–Steklov inequality for the first term in the right-hand side of (6.5), and the Cauchy–Schwarz inequality combined with a (continuous) trace inequality and a local Poincaré–Steklov inequality for the second, we infer

$$(6.6) \quad \begin{aligned} (\mathbb{A} \nabla(v - \pi_K^{m,k}(v)), \nabla w)_K &\lesssim \|g - \Pi_K^m(g)\|_{0,K} H_K |w|_{1,K} \\ &\quad + \|\mathbb{A} \nabla v - \Pi_{\mathcal{F}_K}^k(\mathbb{A} \nabla v)\|_{0,\partial K} H_K^{1/2} |w|_{1,K}, \end{aligned}$$

where we have also used the fact that  $\Pi_{\mathcal{F}_K}^k(\mathbb{A} \nabla v \cdot \mathbf{n}_K) = \Pi_{\mathcal{F}_K}^k(\mathbb{A} \nabla v) \cdot \mathbf{n}_K$  since the mesh faces are planar, combined with the fact that  $\mathbf{n}_K$  is unitary, to handle the boundary term. By definition of  $L^2$ -orthogonal projectors, we have

$$(6.7) \quad \|\mathbb{A} \nabla v - \Pi_{\mathcal{F}_K}^k(\mathbb{A} \nabla v)\|_{0,\partial K} = \min_{\mathbf{p} \in \mathbb{P}^k(\mathcal{F}_K; \mathbb{R}^d)} \|\mathbb{A} \nabla v - \mathbf{p}\|_{0,\partial K} \leq \|\mathbb{A} \nabla v - \Pi_K^k(\mathbb{A} \nabla v)\|_{0,\partial K}.$$

By standard approximation properties of  $L^2$ -orthogonal projectors, we finally obtain from (6.6) and (6.7),

$$\sup_{w \in H^1(K) \setminus \{0\}} \frac{(\mathbb{A} \nabla(v - \pi_K^{m,k}(v)), \nabla w)_K}{|w|_{1,K}} \lesssim H_K^{m+2} |g|_{m+1,K} + H_K^{k+1} |\mathbb{A} \nabla v|_{k+1,K}.$$

The conclusion follows choosing  $w = v - \pi_K^{m,k}(v)$ , and since  $|w|_{1,K}^2 \leq a_{b,K}^{-1} \|\mathbb{A}^{1/2} \nabla w\|_{0,K}^2$ .  $\square$

**Remark 6.2** (Case  $m = -1$ ). Recall that  $\mathbb{P}^{-1}(K) := \{0\}$ . The result of Lemma 6.1 remains valid as it is in the case  $m = -1$  (for  $g \in L^2(K)$ ). The proof needs just be slightly adapted with respect to the general case  $m \geq 0$ . The interpolant  $\pi^{-1,k}(v) \in \mathcal{U}^{-1,k}(K)$  is defined as follows:

$$-\nabla \cdot (\mathbb{A} \nabla \pi_K^{-1,k}(v)) = 0 \quad \text{in } K, \quad \mathbb{A} \nabla \pi_K^{-1,k}(v) \cdot \mathbf{n}_K = \Pi_{\mathcal{F}_K}^k(\mathbb{A} \nabla v \cdot \mathbf{n}_K) + \frac{1}{|\partial K|} (g, 1)_K \quad \text{on } \partial K.$$

The identity (6.5) becomes

$$\begin{aligned} (\mathbb{A} \nabla(v - \pi_K^{-1,k}(v)), \nabla w)_K &= (g, w - \Pi_K^0(w))_K - \frac{1}{|\partial K|} (g, 1)_K (w - \Pi_K^0(w), 1)_{\partial K} \\ &\quad + (\mathbb{A} \nabla v \cdot \mathbf{n}_K - \Pi_{\mathcal{F}_K}^k(\mathbb{A} \nabla v \cdot \mathbf{n}_K), w - \Pi_K^0(w))_{\partial K}. \end{aligned}$$

The conclusion then follows from the same arguments, using in addition that  $\frac{|K|}{|\partial K|} \lesssim H_K$  under our mesh regularity assumptions to handle the second term in the first line of the right-hand side.

**6.3. Energy-norm error estimate.** Let  $m, k \in \mathbb{N}$  be given. We introduce, for any  $K \in \mathcal{T}_H$ , the (local, canonical) interpolation operator  $\mathcal{I}_K : H^1(K) \rightarrow \mathcal{U}^{m,k}(K)$  associated with the finite element (4.3) such that  $\mathcal{I}_K := r_K \circ \widehat{\Sigma}_K$ . Using the definition (4.2) of the reconstruction operator, as well as the definition of the reduction operator  $\widehat{\Sigma}_K$ , we infer that, for any  $v \in H^1(K)$ ,

$$(6.8a) \quad (\mathbb{A} \nabla \mathcal{I}_K(v), \nabla w)_K = (\mathbb{A} \nabla v, \nabla w)_K \quad \forall w \in \mathcal{U}^{m,k}(K),$$

$$(6.8b) \quad (\mathcal{I}_K(v), 1)_{\partial K} = (v, 1)_{\partial K}.$$

Hence,  $\mathcal{I}_K(v) \in \mathcal{U}^{m,k}(K)$  is the ( $\mathbb{A}$ -weighted) elliptic projection of  $v \in H^1(K)$  onto  $\mathcal{U}^{m,k}(K)$ . As such, we have

$$(6.9) \quad \|\mathbb{A}^{1/2} \nabla (v - \mathcal{I}_K(v))\|_{0,K} = \min_{w \in \mathcal{U}^{m,k}(K)} \|\mathbb{A}^{1/2} \nabla (v - w)\|_{0,K}.$$

**Theorem 6.3** (Energy-norm error estimate). *Recall that  $u \in H_0^1(\Omega)$  is the unique solution to (2.2). Let  $u_H \in \mathcal{U}^{m,k}(\mathcal{T}_H) \cap \widetilde{H}_0^{1,k}(\mathcal{T}_H)$  denote either the (fully explicit) MHM solution (3.14) to Problem (3.15), or the MsHHO solution (4.7) to Problem (4.6). Assume that  $f \in H^{m+1}(\mathcal{T}_H)$  and that  $\mathbb{A} \nabla u \in H^{k+1}(\mathcal{T}_H; \mathbb{R}^d)$ . Then, we have*

$$(6.10) \quad \|\mathbb{A}^{1/2} \nabla_H (u - u_H)\|_{0,\Omega} \lesssim \left( \sum_{K \in \mathcal{T}_H} a_{b,K}^{-1} \left( H_K^{2(m+2)} |f|_{m+1,K}^2 + H_K^{2(k+1)} |\mathbb{A} \nabla u|_{k+1,K}^2 \right) \right)^{1/2}.$$

*Proof.* First, by Theorem 5.1, we know that the fully explicit MHM and MsHHO solutions coincide for all  $f \in L^2(\Omega)$ . We here consider the characterization (4.8) of  $u_H$ . Let  $\mathcal{I}_H : H^1(\mathcal{T}_H) \rightarrow \mathcal{U}^{m,k}(\mathcal{T}_H)$  denote the global interpolation operator such that, for all  $v \in H^1(\mathcal{T}_H)$ ,  $\mathcal{I}_H(v)|_K := \mathcal{I}_K(v_K)$  for all  $K \in \mathcal{T}_H$ . Remark that, since  $u \in H_0^1(\Omega)$ ,  $\mathcal{I}_H(u) \in \mathcal{U}^{m,k}(\mathcal{T}_H) \cap \widetilde{H}_0^{1,k}(\mathcal{T}_H)$ . By the triangle inequality, we split the discretization error as follows:

$$(6.11) \quad \|\mathbb{A}^{1/2} \nabla_H (u - u_H)\|_{0,\Omega} \leq \|\mathbb{A}^{1/2} \nabla_H (u - \mathcal{I}_H(u))\|_{0,\Omega} + \|\mathbb{A}^{1/2} \nabla_H (\mathcal{I}_H(u) - u_H)\|_{0,\Omega}.$$

The first term in the right-hand side of (6.11) is an approximation error, and is estimated using the optimality property (6.9) combined with the local approximation properties in  $\mathcal{U}^{m,k}(\mathcal{T}_H)$  of Lemma 6.1. Letting, for all  $v \in H^1(\mathcal{T}_H)$ ,  $\pi_H^{m,k}(v) \in \mathcal{U}^{m,k}(\mathcal{T}_H)$  be the global interpolate such that  $\pi_H^{m,k}(v)|_K = \pi_K^{m,k}(v_K)$  for all  $K \in \mathcal{T}_H$ , we infer

$$(6.12) \quad \begin{aligned} \|\mathbb{A}^{1/2} \nabla_H (u - \mathcal{I}_H(u))\|_{0,\Omega} &= \min_{w_H \in \mathcal{U}^{m,k}(\mathcal{T}_H)} \|\mathbb{A}^{1/2} \nabla_H (u - w_H)\|_{0,\Omega} \\ &\leq \|\mathbb{A}^{1/2} \nabla_H (u - \pi_H^{m,k}(u))\|_{0,\Omega} \\ &\lesssim \left( \sum_{K \in \mathcal{T}_H} a_{b,K}^{-1} \left( H_K^{2(m+2)} |f|_{m+1,K}^2 + H_K^{2(k+1)} |\mathbb{A} \nabla u|_{k+1,K}^2 \right) \right)^{1/2}. \end{aligned}$$

The second term in the right-hand side of (6.11) is the consistency error of the method, which satisfies, since  $(\mathcal{I}_H(u) - u_H) \in \widetilde{\mathcal{U}}_0^{m,k}(\mathcal{T}_H) := \mathcal{U}^{m,k}(\mathcal{T}_H) \cap \widetilde{H}_0^{1,k}(\mathcal{T}_H)$ ,

$$(6.13) \quad \|\mathbb{A}^{1/2} \nabla_H (\mathcal{I}_H(u) - u_H)\|_{0,\Omega} = \max_{\substack{v_H \in \widetilde{\mathcal{U}}_0^{m,k}(\mathcal{T}_H), \\ \|\mathbb{A}^{1/2} \nabla_H v_H\|_{0,\Omega} = 1}} (\mathbb{A} \nabla_H (\mathcal{I}_H(u) - u_H), \nabla_H v_H)_\Omega.$$

Let  $v_H \in \widetilde{\mathcal{U}}_0^{m,k}(\mathcal{T}_H)$  such that  $\|\mathbb{A}^{1/2} \nabla_H v_H\|_{0,\Omega} = 1$ . Since  $u_H$  solves (4.8), we infer

$$(6.14) \quad \begin{aligned} (\mathbb{A} \nabla_H (\mathcal{I}_H(u) - u_H), \nabla_H v_H)_\Omega &= (\mathbb{A} \nabla_H \mathcal{I}_H(u), \nabla_H v_H)_\Omega - (\Pi_H^m(f), v_H)_\Omega \\ &= (\mathbb{A} \nabla_H \mathcal{I}_H(u), \nabla_H v_H)_\Omega + (\nabla \cdot (\mathbb{A} \nabla u), v_H)_\Omega + (f - \Pi_H^m(f), v_H)_\Omega \\ &= (\mathbb{A} \nabla_H (\mathcal{I}_H(u) - u), \nabla_H v_H)_\Omega + \sum_{K \in \mathcal{T}_H} \sum_{F \in \mathcal{F}_K} (\mathbb{A} \nabla u_K \cdot \mathbf{n}_{K,F}, v_K)_F \\ &\quad + (f - \Pi_H^m(f), v_H)_\Omega \\ &= \sum_{F \in \mathcal{F}_H} (\mathbb{A} \nabla u \cdot \mathbf{n}_F, \llbracket v_H \rrbracket_F)_F + (f - \Pi_H^m(f), v_H)_\Omega =: \mathfrak{I}_1 + \mathfrak{I}_2, \end{aligned}$$

where we have added/subtracted  $(f, v_H)_\Omega$  and used that  $f = -\nabla \cdot (\mathbb{A} \nabla u)$  in  $\Omega$  to pass from the first to the second line, we have performed cell-by-cell integration by parts to pass from the second to the third line, and finally used the local orthogonality property (6.8a) as well as the fact that  $\llbracket \mathbb{A} \nabla u \rrbracket_F \cdot \mathbf{n}_F = 0$  for all  $F \in \mathcal{F}_H^{\text{int}}$  as a consequence of the fact that  $\mathbb{A} \nabla u \in \mathbf{H}(\text{div}, \Omega) \cap H^1(\mathcal{T}_H; \mathbb{R}^d)$  to pass from the third to the fourth line. To estimate  $\mathfrak{T}_1$ , we remark that, since  $v_H \in \tilde{H}_0^{1,k}(\mathcal{T}_H)$ ,  $\Pi_F^k(\llbracket v_H \rrbracket_F) = 0$  for all  $F \in \mathcal{F}_H$ . We thus have

$$\begin{aligned} \mathfrak{T}_1 &= \sum_{F \in \mathcal{F}_H} (\mathbb{A} \nabla u \cdot \mathbf{n}_F - \Pi_F^k(\mathbb{A} \nabla u \cdot \mathbf{n}_F), \llbracket v_H - \Pi_F^0(v_H) \rrbracket_F)_F \\ &= \sum_{K \in \mathcal{T}_H} \sum_{F \in \mathcal{F}_K} ((\mathbb{A} \nabla u_K - \Pi_F^k(\mathbb{A} \nabla u_K)) \cdot \mathbf{n}_{K,F}, v_K - \Pi_F^0(v_K))_F. \end{aligned}$$

By two successive applications of the Cauchy–Schwarz inequality, we infer

$$\mathfrak{T}_1 \leq \left( \sum_{K \in \mathcal{T}_H} a_{b,K}^{-1} H_K \|\mathbb{A} \nabla u_K - \Pi_{\mathcal{F}_K}^k(\mathbb{A} \nabla u_K)\|_{0,\partial K}^2 \right)^{1/2} \left( \sum_{K \in \mathcal{T}_H} a_{b,K} H_K^{-1} \|v_K - \Pi_{\mathcal{F}_K}^0(v_K)\|_{0,\partial K}^2 \right)^{1/2}.$$

The first factor in the right-hand side is estimated using (6.7) and standard approximation properties of  $L^2$ -orthogonal projectors. The second factor is estimated by adding/subtracting  $\Pi_K^0(v_K)$ , using a triangle inequality combined with the  $L^2(\partial K)$ -stability of  $\Pi_{\mathcal{F}_K}^0$ , and concluding by the use of a (continuous) trace inequality combined with a local Poincaré–Steklov inequality. We obtain

$$\mathfrak{T}_1 \lesssim \left( \sum_{K \in \mathcal{T}_H} a_{b,K}^{-1} H_K^{2(k+1)} |\mathbb{A} \nabla u|_{k+1,K}^2 \right)^{1/2} \left( \sum_{K \in \mathcal{T}_H} a_{b,K} |v_K|_{1,K}^2 \right)^{1/2}.$$

Recalling that  $\|\mathbb{A}^{1/2} \nabla_H v_H\|_{0,\Omega} = 1$ , and since  $a_{b,K} |v_K|_{1,K}^2 \leq \|\mathbb{A}^{1/2} \nabla v_K\|_{0,K}^2$ , we finally infer that

$$(6.15) \quad \mathfrak{T}_1 \lesssim \left( \sum_{K \in \mathcal{T}_H} a_{b,K}^{-1} H_K^{2(k+1)} |\mathbb{A} \nabla u|_{k+1,K}^2 \right)^{1/2}.$$

The term  $\mathfrak{T}_2$  is, in turn, easily estimated upon writing

$$\mathfrak{T}_2 = (f - \Pi_H^m(f), v_H - \Pi_H^0(v_H))_\Omega,$$

and invoking the Cauchy–Schwarz inequality, a local Poincaré–Steklov inequality, and standard approximation properties of  $L^2$ -orthogonal projectors to conclude. We obtain

$$(6.16) \quad \begin{aligned} \mathfrak{T}_2 &\lesssim \left( \sum_{K \in \mathcal{T}_H} a_{b,K}^{-1} H_K^{2(m+2)} |f|_{m+1,K}^2 \right)^{1/2} \left( \sum_{K \in \mathcal{T}_H} a_{b,K} |v_K|_{1,K}^2 \right)^{1/2} \\ &\lesssim \left( \sum_{K \in \mathcal{T}_H} a_{b,K}^{-1} H_K^{2(m+2)} |f|_{m+1,K}^2 \right)^{1/2}, \end{aligned}$$

where we used again that  $\|\mathbb{A}^{1/2} \nabla_H v_H\|_{0,\Omega} = 1$  to pass from the first to the second line. Finally, plugging (6.15)-(6.16)-(6.14)-(6.13) and (6.12) into (6.11) proves (6.10).  $\square$

**Remark 6.4** (Case  $m = -1$ ). *We know from Remark 5.4 that the (fully explicit) MHM method for  $m = -1$  coincides with the (fully explicit) MHM method for  $m = 0$ . As far as the MsHMO method is concerned, in the case  $m = -1$ , one adopts the variant (5.1) of the method, and the a priori estimate of Theorem 6.3 remains valid as it is (for  $f \in L^2(\Omega)$ ). The proof actually simplifies with respect to the general case  $m \geq 0$ , since the term  $\mathfrak{T}_2$  can be discarded. The conclusion follows from Lemma 6.1 and Remark 6.2.*

**Remark 6.5** (Case  $m = k - 1$ ). *In the case  $m = k - 1$ , the result (6.10) (see Remark 6.4 for the case  $k = 0$  and  $m = -1$ ) simplifies since  $|f|_{k,K} \leq \sqrt{d} |\mathbb{A} \nabla u|_{k+1,K}$  for all  $K \in \mathcal{T}_H$ . Under the sole*

assumption that  $\mathbb{A}\nabla u \in H^{k+1}(\mathcal{T}_H; \mathbb{R}^d)$ , we have

$$\|\mathbb{A}^{1/2}\nabla_H(u - u_H)\|_{0,\Omega} \lesssim \left( \sum_{K \in \mathcal{T}_H} a_{\mathfrak{b},K}^{-1} H_K^{2(k+1)} |\mathbb{A}\nabla u|_{k+1,K}^2 \right)^{1/2}.$$

In the MHM setting, when  $k = 0$  (then one can discard the contribution given by the operator  $T^{\mathfrak{S}}$ ), we obtain an optimal error estimate under the sole assumption on the source term that  $f \in L^2(\Omega)$ , which improves on [4, Corollary 4.2] where more regularity is needed.

**Remark 6.6** (Link with previous results). *In the MHM framework, the error estimate of Theorem 6.3 is a refined version of [4, Theorem 4.1] (for the original, semi-explicit MHM method), both in terms of regularity assumptions and in terms of tracking of the dependency of the multiplicative constants with respect to the diffusion coefficient. In the MsHHO framework, such an error estimate is new, and is complementary to the homogenization-based error estimate of [12, Theorem 5.6] (such a homogenization-based analysis is also available in the MHM setting; cf. [38]). The a priori estimate of [12, Theorem 5.6] is robust in highly oscillatory diffusion regimes but is suboptimal for mildly varying diffusion. The present result fills this gap.*

## 7. BASIS FUNCTIONS AND SOLUTION STRATEGIES

We address the decomposition of the MHM and MsHHO solutions in terms of multiscale basis functions and highlight the impact of such a decomposition on the possible organization of the computations using an offline-online strategy. Let  $k \geq 1$  be a given integer. In what follows, to keep the presentation simple, we consider for a polynomial degree  $k$  on the faces the polynomial degree  $m := k - 1 \geq 0$  in the cell, so that we can simply write  $\mathcal{U}^k(K)$  in lieu of  $\mathcal{U}^{k-1,k}(K)$  for all  $K \in \mathcal{T}_H$ . The key observation is that there are two possible constructions of basis functions for the local space  $\mathcal{U}^k(K)$ . Both sets of basis functions are composed of cell-based and face-based functions. The construction of the two sets is however different. The first construction, referred to as *primal set*, will prove to be relevant for the MHM method, whereas the second construction, referred to as *dual set*, will prove to be relevant for the MsHHO method.

### 7.1. Basis functions.

**7.1.1. Polynomial basis functions.** Let  $q \in \mathbb{N}$ . We denote by  $n_l^q$  the dimension of the vector space of  $l$ -variate polynomial functions of total degree up to  $q$ . For any cell  $K \in \mathcal{T}_H$ , let  $\{\psi_i^{q,K}\}_{1 \leq i \leq n_d^q}$  be a basis of  $\mathbb{P}^q(K)$ , and for any face  $F \in \mathcal{F}_H$ , let  $\{\psi_j^{q,F}\}_{1 \leq j \leq n_{d-1}^q}$  be a basis of  $\mathbb{P}^q(F)$ . With the choice we have made of degree  $q := k - 1$  in the cell and degree  $q := k$  on the faces, we henceforth drop the corresponding superscripts in the polynomial basis functions to alleviate the notation. For convenience, we assume that  $\psi_1^K \equiv 1$ ; this assumption will be useful in the MHM setting.

**7.1.2. Primal basis functions.** For  $K \in \mathcal{T}_H$ , we locally construct the set of primal basis functions for  $\mathcal{U}^k(K)$ . Regarding the cell-based basis functions, we set  $\phi_1^{p,K} \equiv 1$ , and for all  $2 \leq i \leq n_d^{k-1}$ , we define  $\phi_i^{p,K}$  as the unique function in  $H^1(K)^\perp$  solving the following well-posed Neumann problem:

$$(7.1) \quad \begin{cases} -\nabla \cdot (\mathbb{A}\nabla \phi_i^{p,K}) = \psi_i^K - \Pi_K^0(\psi_i^K) \text{ in } K, \\ \mathbb{A}\nabla \phi_i^{p,K} \cdot \mathbf{n}_K = 0 \text{ on } \partial K. \end{cases}$$

Concerning the face-based basis functions, for all  $F \in \mathcal{F}_K$  and all  $1 \leq j \leq n_{d-1}^k$ , we define  $\phi_{F,j}^{p,K}$  as the unique function in  $H^1(K)^\perp$  solving the following well-posed Neumann problem:

$$(7.2) \quad \begin{cases} -\nabla \cdot (\mathbb{A}\nabla \phi_{F,j}^{p,K}) = -\frac{1}{|K|}(\psi_j^F, 1)_F \text{ in } K, \\ \mathbb{A}\nabla \phi_{F,j}^{p,K} \cdot \mathbf{n}_{K,F} = \psi_j^F \text{ on } F \quad \text{and} \quad \mathbb{A}\nabla \phi_{F,j}^{p,K} \cdot \mathbf{n}_{K,\sigma} = 0 \text{ on } \sigma \in \mathcal{F}_K \setminus \{F\}. \end{cases}$$

Then, for all  $v \in \mathcal{U}^k(K)$ , setting

$$(i) \quad -\nabla \cdot (\mathbb{A}\nabla v) := g_K = g_{K,1} + \sum_{i=2}^{n_d^{k-1}} g_{K,i} \psi_i^K \in \mathbb{P}^{k-1}(K) \quad (\text{recall that } \psi_1^K \equiv 1),$$

- (ii)  $\mathbb{A}\nabla v|_{\partial K} \cdot \mathbf{n}_K := \mu_{\mathcal{F}_K} \in \mathbb{P}^k(\mathcal{F}_K)$  with  $\mu_{\mathcal{F}_K}|_F = \sum_{j=1}^{n_d^k-1} \mu_{F,j} \psi_j^F$  for all  $F \in \mathcal{F}_K$ ,  
(iii)  $\Pi_K^0(v) := v_K^0 \in \mathbb{P}^0(K)$ ,

with  $(g_K, 1)_K + (\mu_{\mathcal{F}_K}, 1)_{\partial K} = 0$ , we have

$$(7.3) \quad v = v_K^0 + \sum_{F \in \mathcal{F}_K} \sum_{j=1}^{n_d^k-1} \mu_{F,j} \phi_{F,j}^{\text{p},K} + \sum_{i=2}^{n_d^k-1} g_{K,i} \phi_i^{\text{p},K}.$$

A set of global basis functions for the space  $\mathcal{U}^k(\mathcal{T}_H) \cap \mathcal{V}(\mathcal{T}_H; \text{div}, \Omega)$  is given by

$$\{\tilde{\phi}_i^{\text{p},K}\}_{K \in \mathcal{T}_H, 1 \leq i \leq n_d^{k-1}} \cup \{\tilde{\phi}_j^{\text{p},F}\}_{F \in \mathcal{F}_H, 1 \leq j \leq n_d^{k-1}},$$

where for each  $K \in \mathcal{T}_H$ ,

$$(7.4) \quad \tilde{\phi}_i^{\text{p},K}|_K = \phi_i^{\text{p},K} \quad \text{and} \quad \tilde{\phi}_i^{\text{p},K}|_{\Omega \setminus \bar{K}} = 0,$$

for each interface  $F \subseteq \partial K_+ \cap \partial K_-$ ,

$$(7.5) \quad \tilde{\phi}_j^{\text{p},F}|_{K_{\pm}} = (\mathbf{n}_{K_{\pm},F} \cdot \mathbf{n}_F) \phi_{F,j}^{\text{p},K_{\pm}} \quad \text{and} \quad \tilde{\phi}_j^{\text{p},F}|_{\Omega \setminus \overline{K_+ \cup K_-}} = 0,$$

and for each boundary face  $F \subseteq \partial K \cap \partial \Omega$ ,

$$(7.6) \quad \tilde{\phi}_j^{\text{p},F}|_K = \phi_{F,j}^{\text{p},K} \quad \text{and} \quad \tilde{\phi}_j^{\text{p},F}|_{\Omega \setminus \bar{K}} = 0.$$

**Remark 7.1** (Link to lifting operators). *Recall the local lifting operators  $T_K^{\text{N}}, T_K^{\text{S}}$  and their global counterparts  $T^{\text{N}}, T^{\text{S}}$  introduced in Section 3. For all  $K \in \mathcal{T}_H$ , one readily verifies that*

$$(7.7) \quad \phi_i^{\text{p},K} = T_K^{\text{S}}(\psi_i^K), \quad \phi_{F,j}^{\text{p},K} = T_K^{\text{N}}(E_F^{\partial K}(\psi_j^F)),$$

where the first identity holds for all  $2 \leq i \leq n_d^{k-1}$  and the second identity holds for all  $F \in \mathcal{F}_K$  and all  $1 \leq j \leq n_d^{k-1}$ , where  $E_F^{\partial K}$  denotes the zero-extension operator from  $F$  to  $\partial K$ . For the global basis functions, we have

$$(7.8) \quad \tilde{\phi}_i^{\text{p},K} = T^{\text{S}}(E_K^{\Omega}(\psi_i^K)), \quad \tilde{\phi}_j^{\text{p},F} = T^{\text{N}}(E_F^{\partial \mathcal{T}_H}(\psi_j^F)),$$

where  $E_K^{\Omega}$  denotes the zero-extension operator from  $K$  to  $\Omega$ , and  $E_F^{\partial \mathcal{T}_H}(\psi_j^F)|_{\partial K} := E_F^{\partial K}(\psi_j^F)(\mathbf{n}_{K,F} \cdot \mathbf{n}_F)$  if  $F \in \mathcal{F}_K$  and  $E_F^{\partial \mathcal{T}_H}(\psi_j^F)|_{\partial K} := 0$  otherwise, for all  $K \in \mathcal{T}_H$ .

**Remark 7.2** (Energy minimization). *Consider the local energy functional  $J_K : H^1(K) \rightarrow \mathbb{R}_+$  such that  $\varphi \mapsto \frac{1}{2}(\mathbb{A}\nabla \varphi, \nabla \varphi)_K$ . Then, one can characterize  $\phi_i^{\text{p},K}$  for all  $2 \leq i \leq n_d^{k-1}$  as follows:*

$$(7.9) \quad \phi_i^{\text{p},K} = \arg \min_{\varphi \in H^1(K)^{\perp}} (J_K(\varphi) - (\psi_i^K - \Pi_K^0(\psi_i^K), \varphi)_K),$$

and one can characterize  $\phi_{F,j}^{\text{p},K}$  for all  $F \in \mathcal{F}_K$  and all  $1 \leq j \leq n_d^{k-1}$  as follows:

$$(7.10) \quad \phi_{F,j}^{\text{p},K} = \arg \min_{\varphi \in H^1(K)^{\perp}} \left( J_K(\varphi) - (\psi_j^F, \varphi)_F + \frac{1}{|K|} (\psi_j^F, 1)_F (\varphi, 1)_K \right),$$

where we recall that  $H^1(K)^{\perp} := \{v \in H^1(K) : (v, 1)_K = 0\}$ .

**7.1.3. Dual basis functions.** For  $K \in \mathcal{T}_H$ , we locally construct the set of dual basis functions for  $\mathcal{U}^k(K)$ . For this purpose, we rely on the fact that the triple  $(K, \mathcal{U}^k(K), \widehat{\Sigma}_K)$  is a finite element (see (4.3)). For all  $1 \leq i \leq n_d^{k-1}$ , the cell-based basis functions  $\phi_i^{\text{d},K} \in \mathcal{U}^k(K)$  are obtained by requiring that

$$(7.11) \quad \Pi_K^{k-1}(\phi_i^{\text{d},K}) = \psi_i^K, \quad \Pi_{\mathcal{F}_K}^k(\phi_i^{\text{d},K}) = 0,$$

that is, we have  $\phi_i^{\text{d},K} := r_K((\psi_i^K, 0))$ . Moreover, for all  $F \in \mathcal{F}_K$  and all  $1 \leq j \leq n_d^{k-1}$ , the face-based basis functions  $\phi_{F,j}^{\text{d},K} \in \mathcal{U}^k(K)$  are obtained by requiring that

$$(7.12) \quad \Pi_K^{k-1}(\phi_{F,j}^{\text{d},K}) = 0, \quad \Pi_F^k(\phi_{F,j}^{\text{d},K}) = \psi_j^F, \quad \Pi_{\sigma}^k(\phi_{F,j}^{\text{d},K}) = 0 \text{ for all } \sigma \in \mathcal{F}_K \setminus \{F\},$$

that is, we have  $\phi_{F,j}^{\text{d},K} := r_K((0, E_F^{\partial K}(\psi_j^F)))$ . Then, for all  $v \in \mathcal{U}^k(K)$ , setting

$$(i) \quad \Pi_K^{k-1}(v) := v_K = \sum_{i=1}^{n_d^{k-1}} v_{K,i} \psi_i^K \in \mathbb{P}^{k-1}(K),$$

(ii)  $\Pi_{\mathcal{F}_K}^k(v) := v_{\mathcal{F}_K} \in \mathbb{P}^k(\mathcal{F}_K)$  with  $v_{\mathcal{F}_K|F} = \sum_{j=1}^{n_d^{k-1}} v_{F,j} \psi_j^F$  for all  $F \in \mathcal{F}_K$ , we have

$$(7.13) \quad v = \sum_{i=1}^{n_d^{k-1}} v_{K,i} \phi_i^{d,K} + \sum_{F \in \mathcal{F}_K} \sum_{j=1}^{n_d^k} v_{F,j} \phi_{F,j}^{d,K}.$$

Notice that we also have  $v = r_K(\widehat{v}_K)$  where  $\widehat{v}_K := (v_K, v_{\mathcal{F}_K}) \in \widehat{U}_K^k$ .

A set of global basis functions for the space  $\mathcal{U}^k(\mathcal{T}_H) \cap \widetilde{H}_0^{1,k}(\mathcal{T}_H)$  is given by

$$(7.14) \quad \{\widetilde{\phi}_i^{d,K}\}_{K \in \mathcal{T}_H, 1 \leq i \leq n_d^{k-1}} \cup \{\widetilde{\phi}_j^{d,F}\}_{F \in \mathcal{F}_H^{\text{int}}, 1 \leq j \leq n_d^{k-1}},$$

where for each  $K \in \mathcal{T}_H$ ,

$$(7.15) \quad \widetilde{\phi}_i^{d,K}|_K = \phi_i^{d,K} \quad \text{and} \quad \widetilde{\phi}_i^{d,K}|_{\Omega \setminus \overline{K}} = 0,$$

and for each interface  $F \subseteq \partial K_+ \cap \partial K_-$ ,

$$(7.16) \quad \widetilde{\phi}_j^{d,F}|_{K_{\pm}} = \phi_{F,j}^{d,K_{\pm}} \quad \text{and} \quad \widetilde{\phi}_j^{d,F}|_{\Omega \setminus \overline{K_+ \cup K_-}} = 0.$$

**Remark 7.3** (Energy minimization). *Recall the local energy functional  $J_K$  defined in Remark 7.2. Then, one can characterize  $\phi_i^{d,K}$  for all  $1 \leq i \leq n_d^{k-1}$  as follows:*

$$(7.17) \quad \phi_i^{d,K} := \arg \min_{\varphi \in H_i^K} J_K(\varphi),$$

where  $H_i^K := \{v \in H^K : \Pi_K^{k-1}(v) = \psi_i^K\}$  is a nonempty, convex, closed subset of the Hilbert space  $H^K := \{v \in H^1(K) : \Pi_{\mathcal{F}_K}^k(v) = 0\}$ . This means that  $\phi_i^{d,K} \in H^1(K)$  is obtained by solving the following saddle-point problem with dual unknowns  $\gamma_i^K \in \mathbb{P}^{k-1}(K)$  and  $\mu_i^{\partial K} \in \mathbb{P}^k(\mathcal{F}_K)$  such that  $(\gamma_i^K, 1)_K + (\mu_i^{\partial K}, 1)_{\partial K} = 0$ :

$$(7.18) \quad \begin{cases} -\nabla \cdot (\mathbb{A} \nabla \phi_i^{d,K}) = \gamma_i^K & \text{in } K, \quad \mathbb{A} \nabla \phi_i^{d,K} \cdot \mathbf{n}_K = \mu_i^{\partial K} & \text{on } \partial K, \\ \Pi_K^{k-1}(\phi_i^{d,K}) = \psi_i^K, \quad \Pi_{\mathcal{F}_K}^k(\phi_i^{d,K}) = 0. \end{cases}$$

Similarly, one can characterize  $\phi_{F,j}^{d,K}$  for all  $F \in \mathcal{F}_K$  and all  $1 \leq j \leq n_d^{k-1}$  as follows:

$$(7.19) \quad \phi_{F,j}^{d,K} := \arg \min_{\varphi \in H_{F,j}^K} J_K(\varphi),$$

where  $H_{F,j}^K := \{v \in H_F^K : \Pi_F^k(v) = \psi_j^F\}$  is a nonempty, convex, closed subset of the Hilbert space  $H_F^K := \{v \in H^1(K) : \Pi_K^{k-1}(v) = 0 \text{ and } \Pi_{\sigma}^k(v) = 0 \quad \forall \sigma \in \mathcal{F}_K \setminus \{F\}\}$ . This means that  $\phi_{F,j}^{d,K} \in H^1(K)$  is obtained by solving the following saddle-point problem with dual unknowns  $\gamma_{F,j}^K \in \mathbb{P}^{k-1}(K)$  and  $\mu_{F,j}^{\partial K} \in \mathbb{P}^k(\mathcal{F}_K)$  such that  $(\gamma_{F,j}^K, 1)_K + (\mu_{F,j}^{\partial K}, 1)_{\partial K} = 0$ :

$$(7.20) \quad \begin{cases} -\nabla \cdot (\mathbb{A} \nabla \phi_{F,j}^{d,K}) = \gamma_{F,j}^K & \text{in } K, \quad \mathbb{A} \nabla \phi_{F,j}^{d,K} \cdot \mathbf{n}_K = \mu_{F,j}^{\partial K} & \text{on } \partial K, \\ \Pi_K^{k-1}(\phi_{F,j}^{d,K}) = 0, \quad \Pi_F^k(\phi_{F,j}^{d,K}) = \psi_j^F, \quad \Pi_{\sigma}^k(\phi_{F,j}^{d,K}) = 0 & \text{for all } \sigma \in \mathcal{F}_K \setminus \{F\}. \end{cases}$$

**7.2. Offline-online strategy.** In view of Section 7.1, primal basis functions, as they globally span  $\mathcal{U}^k(\mathcal{T}_H) \cap \mathcal{V}(\mathcal{T}_H; \text{div}, \Omega)$ , appear to be naturally suited to the MHM framework. On the other hand, dual basis functions, as they globally span  $\mathcal{U}^k(\mathcal{T}_H) \cap \widetilde{H}_0^{1,k}(\mathcal{T}_H)$ , appear to be naturally suited to the MsHHO framework (cf. Remark 5.2). In this section, we detail how the MHM and MsHHO computations can be optimally organized using an offline-online strategy. This type of organization of the computations is particularly relevant in a multi-query context, in which the solution has to be computed for a large amount of data, so that it is crucial to pre-process as many data-independent quantities as possible in an offline stage, while keeping the size of the online system to its minimum. We focus in the sequel on the situation where many instances of the source term  $f$  are considered (we could also consider the case of multiple boundary data).



7.2.1. *The MHM case.* By Remark 7.1, the (fully explicit) MHM solution  $u_H^{\text{MHM}} \in \mathcal{U}^k(\mathcal{T}_H) \cap \mathcal{V}(\mathcal{T}_H; \text{div}, \Omega)$  defined by (3.14) with  $m := k - 1$ , where the pair  $(u_H^0, \lambda_H) \in \mathbb{P}^0(\mathcal{T}_H) \times \Lambda^k(\partial\mathcal{T}_H)$  solves (3.15), writes

$$(7.21) \quad u_H^{\text{MHM}} = \sum_{K \in \mathcal{T}_H} u_K^0 \tilde{\phi}_1^{\text{p},K} + \sum_{F \in \mathcal{F}_H} \sum_{j=1}^{n_d^{k-1}} \lambda_{F,j} \tilde{\phi}_j^{\text{p},F} + \sum_{K \in \mathcal{T}_H} \sum_{i=2}^{n_d^{k-1}} f_{K,i} \tilde{\phi}_i^{\text{p},K},$$

where  $u_K^0 := u_{H|K}^0 = \Pi_K^0(u_H^{\text{MHM}})$  for any  $K \in \mathcal{T}_H$ ,  $\lambda_{F,j}$  is defined, for any  $F \in \mathcal{F}_H$ , as the  $j^{\text{th}}$  coefficient of  $\lambda_{H|F}$  on the basis  $\{\psi_j^F\}_{1 \leq j \leq n_d^{k-1}}$ , and  $f_{K,i}$  stands for the  $i^{\text{th}}$  coefficient of  $\Pi_K^{k-1}(f_K)$  on the basis  $\{\psi_i^K\}_{1 \leq i \leq n_d^{k-1}}$ . This motivates the following offline-online decomposition of the computations:

Offline stage: For each  $K \in \mathcal{T}_H$ :

- (1) Compute the basis functions  $\phi_i^{\text{p},K}$  from (7.1), for all  $i = 2, \dots, n_d^{k-1}$ ;
- (2) Compute the basis functions  $\phi_{F,j}^{\text{p},K}$  from (7.2), for all  $F \in \mathcal{F}_K$  and all  $j = 1, \dots, n_d^{k-1}$ .

Online stage:

- (3) Compute the vector  $(f_{K,i})_{K \in \mathcal{T}_H}^{i=1, \dots, n_d^{k-1}}$  by solving the local symmetric positive-definite (SPD) systems

$$\sum_{i=1}^{n_d^{k-1}} f_{K,i} (\psi_i^K, \psi_j^K)_K = (f_K, \psi_j^K)_K,$$

for all  $j = 1, \dots, n_d^{k-1}$ , and all  $K \in \mathcal{T}_H$ ;

- (4) Compute the vectors  $(u_K^0)_{K \in \mathcal{T}_H}$  and  $(\lambda_{F,j})_{F \in \mathcal{F}_H}^{j=1, \dots, n_d^{k-1}}$  by solving the global saddle-point problem

$$\sum_{F \in \mathcal{F}_K} \sum_{j=1}^{n_d^{k-1}} \lambda_{F,j} (\psi_j^F, 1)_F = -(f_K, 1)_K,$$

for all  $K \in \mathcal{T}_H$ , and (recall that  $\phi_1^{\text{p},K} \equiv 1$  and that  $(\phi_{F',j'}^{\text{p},K}, 1)_K = 0$ )

$$\sum_{K \in \mathcal{T}_{F'}} u_K^0 (\psi_{j'}^{F'}, 1)_{F'} + \sum_{K \in \mathcal{T}_{F'}} \sum_{F \in \mathcal{F}_K} \sum_{j=1}^{n_d^{k-1}} \lambda_{F,j} (\psi_j^{F'}, \tilde{\phi}_j^{\text{p},F})_{F'} = - \sum_{K \in \mathcal{T}_{F'}} \sum_{i=2}^{n_d^{k-1}} f_{K,i} (\psi_i^K, \phi_{F',j'}^{\text{p},K})_K,$$

for all  $j' = 1, \dots, n_d^{k-1}$ , and all  $F' \in \mathcal{F}_H$  with  $\mathcal{T}_{F'} := \{K_+, K_-\}$  if  $F' \in \mathcal{F}_H^{\text{int}}$  and  $\mathcal{T}_{F'} := \{K\}$  if  $F' \in \mathcal{F}_H^{\text{bnd}}$ ;

- (5) Form  $u_H^{\text{MHM}}$  using (7.21).

**Remark 7.4** (Mono-query case). *In a mono-query scenario, in which the solution to the discrete problem is only needed for one (or a few) source term(s), one can advantageously consider an amended version of (7.21), where the last term in the decomposition is simply replaced by  $T^{\text{s}}(\Pi_H^{k-1}(f))$ . From a practical point of view, the step (1) above can be bypassed, and replaced by solving, inbetween steps (3) and (4), Problem (7.1) for all  $K \in \mathcal{T}_H$  with right-hand side  $\Pi_K^{k-1}(f_K)$  (in lieu of  $\psi_i^K$ ), whose solution is precisely  $T_K^{\text{s}}(\Pi_K^{k-1}(f_K))$ .*

7.2.2. *The MsHHO case.* The solution  $u_H^{\text{HHO}} \in \mathcal{U}^k(\mathcal{T}_H) \cap \tilde{H}_0^{1,k}(\mathcal{T}_H)$  to Problem (4.8) writes

$$(7.22) \quad u_H^{\text{HHO}} = \sum_{K \in \mathcal{T}_H} \sum_{i=1}^{n_d^{k-1}} u_{K,i} \tilde{\phi}_i^{\text{d},K} + \sum_{F \in \mathcal{F}_H^{\text{int}}} \sum_{j=1}^{n_d^{k-1}} u_{F,j} \tilde{\phi}_j^{\text{d},F},$$

where  $u_{K,i}$  is defined as the  $i^{\text{th}}$  coefficient of  $u_K := \Pi_K^{k-1}(u_H^{\text{HHO}})$  on the basis  $\{\psi_i^K\}_{1 \leq i \leq n_d^{k-1}}$  for any  $K \in \mathcal{T}_H$ , and  $u_{F,j}$  as the  $j^{\text{th}}$  coefficient of  $u_F := \Pi_F^k(u_H^{\text{HHO}})$  on the basis  $\{\psi_j^F\}_{1 \leq j \leq n_d^{k-1}}$  for any  $F \in \mathcal{F}_H^{\text{int}}$  (recall that  $\Pi_F^k(u_H^{\text{HHO}}) = 0$  for all  $F \in \mathcal{F}_H^{\text{bnd}}$ ). This, combined with the equivalent

formulation (4.9) of the MsHHO method, and Remark 7.3 (recall, in particular, the notation introduced therein), motivates the following offline-online decomposition of the computations:

Offline stage: For each  $K \in \mathcal{T}_H$ :

- (1) Compute the basis functions  $\phi_i^{d,K}$  from (7.18), for all  $i = 1, \dots, n_d^{k-1}$ ;
- (2) Compute the basis functions  $\phi_{F,j}^{d,K}$  from (7.20), for all  $F \in \mathcal{F}_K$  and all  $j = 1, \dots, n_{d-1}^k$ ;

Define

- the  $n_d^{k-1} \times n_d^{k-1}$  matrix  $\mathbb{G}^{KK}$ , whose column  $1 \leq i \leq n_d^{k-1}$  is formed by the  $n_d^{k-1}$  coefficients of the decomposition of  $\gamma_i^K \in \mathbb{P}^{k-1}(K)$  on the basis  $\{\psi_{i'}^K\}_{1 \leq i' \leq n_d^{k-1}}$ ;
  - for each  $F \in \mathcal{F}_K$ , the  $n_d^{k-1} \times n_{d-1}^k$  matrix  $\mathbb{G}^{KF}$ , whose column  $1 \leq j \leq n_{d-1}^k$  is formed by the  $n_d^{k-1}$  coefficients of the decomposition of  $\gamma_{F,j}^K \in \mathbb{P}^{k-1}(K)$  on the basis  $\{\psi_i^K\}_{1 \leq i \leq n_d^{k-1}}$ ;
  - for each  $F \in \mathcal{F}_K$ , the  $n_{d-1}^k \times n_{d-1}^k$  matrix  $\mathbb{M}^{FK}$ , whose column  $1 \leq i \leq n_{d-1}^k$  is formed by the  $n_{d-1}^k$  coefficients of the decomposition of  $\mu_{iF}^{\partial K} \in \mathbb{P}^k(F)$  on the basis  $\{\psi_j^F\}_{1 \leq j \leq n_{d-1}^k}$ ;
  - for each  $F, F' \in \mathcal{F}_K$ , the  $n_{d-1}^k \times n_{d-1}^k$  matrix  $\mathbb{M}^{F'F}$ , whose column  $1 \leq j \leq n_{d-1}^k$  is formed by the  $n_{d-1}^k$  coefficients of the decomposition of  $\mu_{F,j|F'}^{\partial K} \in \mathbb{P}^k(F')$  on the basis  $\{\psi_{j'}^{F'}\}_{1 \leq j' \leq n_{d-1}^k}$ ;
- (3) Invert the matrix  $\mathbb{G}^{KK}$ .

Online stage:

- (4) Compute the vectors  $(\mathbf{f}_K)_{K \in \mathcal{T}_H} := (f_{K,i})_{K \in \mathcal{T}_H}^{i=1, \dots, n_d^{k-1}}$  by solving the local SPD systems

$$\sum_{i=1}^{n_d^{k-1}} f_{K,i} (\psi_i^K, \psi_j^K)_K = (f_K, \psi_j^K)_K,$$

for all  $j = 1, \dots, n_d^{k-1}$ , and all  $K \in \mathcal{T}_H$ ;

- (5) Compute the vectors  $(\mathbf{u}_F)_{F \in \mathcal{F}_H^{\text{int}}} := (u_{F,j})_{F \in \mathcal{F}_H^{\text{int}}}^{j=1, \dots, n_{d-1}^k}$  by solving the global SPD problem

$$\sum_{K \in \mathcal{T}_{F'}} \sum_{F \in \mathcal{F}_K \cap \mathcal{F}_H^{\text{int}}} (\mathbb{M}^{F'F} - \mathbb{M}^{F'K} [\mathbb{G}^{KK}]^{-1} \mathbb{G}^{KF}) \mathbf{u}_F = - \sum_{K \in \mathcal{T}_{F'}} \mathbb{M}^{F'K} [\mathbb{G}^{KK}]^{-1} \mathbf{f}_K,$$

for all  $F' \in \mathcal{F}_H^{\text{int}}$ ;

- (6) Reconstruct locally the vectors  $(\mathbf{u}_K)_{K \in \mathcal{T}_H} := (u_{K,i})_{K \in \mathcal{T}_H}^{i=1, \dots, n_d^{k-1}}$ : for all  $K \in \mathcal{T}_H$ ,

$$\mathbf{u}_K = [\mathbb{G}^{KK}]^{-1} \left( \mathbf{f}_K - \sum_{F \in \mathcal{F}_K \cap \mathcal{F}_H^{\text{int}}} \mathbb{G}^{KF} \mathbf{u}_F \right);$$

- (7) Form  $u_H^{\text{HHO}}$  using (7.22).

**Remark 7.5** (Purely face-based version). *Using the (primal-dual) local set of basis functions for  $\mathcal{U}^k(K)$ ,  $K \in \mathcal{T}_H$ , introduced in [12, Section 4.1] (but not fully exploited therein), the MsHHO method can be alternatively defined as a purely face-based method, i.e. without using cell unknowns. To see this, let  $K \in \mathcal{T}_H$ , and recall the local energy functional  $J_K$  defined in Remark 7.2. Define  $\phi_i^K$  for all  $1 \leq i \leq n_d^{k-1}$  as follows:*

$$(7.23) \quad \phi_i^K := \arg \min_{\varphi \in H^K} (J_K(\varphi) - (\psi_i^K, \varphi)_K),$$

where the space  $H^K$  is defined in Remark 7.3. Equivalently,  $\phi_i^K \in H^1(K)$  is obtained by solving the following saddle-point problem with dual unknown  $\mu_i^{\partial K} \in \mathbb{P}^k(\mathcal{F}_K)$  such that  $(\psi_i^K, 1)_K + (\mu_i^{\partial K}, 1)_{\partial K} = 0$ :

$$(7.24) \quad \begin{cases} -\nabla \cdot (\mathbb{A} \nabla \phi_i^K) = \psi_i^K & \text{in } K, \quad \mathbb{A} \nabla \phi_i^K \cdot \mathbf{n}_K = \mu_i^{\partial K} & \text{on } \partial K, \\ \Pi_{\mathcal{F}_K}^k(\phi_i^K) = 0. \end{cases}$$

Similarly, define  $\phi_{F,j}^K$  for all  $F \in \mathcal{F}_K$  and all  $1 \leq j \leq n_{d-1}^k$  as follows:

$$(7.25) \quad \phi_{F,j}^K := \arg \min_{\varphi \in H_{F,j}^K} J_K(\varphi),$$

where  $H_{F,j}^K := \{v \in H_F^K : \Pi_F^k(v) = \psi_j^F\}$  as in Remark 7.3, but now we set  $H_F^K := \{v \in H^1(K) : \Pi_\sigma^k(v) = 0 \ \forall \sigma \in \mathcal{F}_K \setminus \{F\}\}$ . Equivalently,  $\phi_{F,j}^K \in H^1(K)$  is obtained by solving the following saddle-point problem with dual unknown  $\mu_{F,j}^{\partial K} \in \mathbb{P}^k(\mathcal{F}_K)$  such that  $(\mu_{F,j}^{\partial K}, 1)_{\partial K} = 0$ :

$$(7.26) \quad \begin{cases} -\nabla \cdot (\mathbb{A} \nabla \phi_{F,j}^K) = 0 \text{ in } K, & \mathbb{A} \nabla \phi_{F,j}^K \cdot \mathbf{n}_K = \mu_{F,j}^{\partial K} \text{ on } \partial K, \\ \Pi_F^k(\phi_{F,j}^K) = \psi_j^F, & \Pi_\sigma^k(\phi_{F,j}^K) = 0 \text{ for all } \sigma \in \mathcal{F}_K \setminus \{F\}. \end{cases}$$

For all  $v \in \mathcal{U}^k(K)$ , setting

$$(i) \quad -\nabla \cdot (\mathbb{A} \nabla v) := g_K = \sum_{i=1}^{n_d^{k-1}} g_{K,i} \psi_i^K \in \mathbb{P}^{k-1}(K),$$

$$(ii) \quad \Pi_{\mathcal{F}_K}^k(v) := v_{\mathcal{F}_K} \in \mathbb{P}^k(\mathcal{F}_K) \text{ with } v_{\mathcal{F}_K|F} = \sum_{j=1}^{n_{d-1}^k} v_{F,j} \psi_j^F \text{ for all } F \in \mathcal{F}_K,$$

we then have

$$(7.27) \quad v = \sum_{i=1}^{n_d^{k-1}} g_{K,i} \phi_i^K + \sum_{F \in \mathcal{F}_K} \sum_{j=1}^{n_{d-1}^k} v_{F,j} \phi_{F,j}^K.$$

As we did for the dual set of basis functions in (7.14)–(7.15)–(7.16), we can easily construct a set of global basis functions  $\{\tilde{\phi}_i^K\}_{K \in \mathcal{T}_H, 1 \leq i \leq n_d^{k-1}} \cup \{\tilde{\phi}_j^F\}_{F \in \mathcal{F}_H^{\text{int}}, 1 \leq j \leq n_{d-1}^k}$  for the space  $\mathcal{U}^k(\mathcal{T}_H) \cap \tilde{H}_0^{1,k}(\mathcal{T}_H)$ .

The solution  $u_H^{\text{HHO}} \in \mathcal{U}^k(\mathcal{T}_H) \cap \tilde{H}_0^{1,k}(\mathcal{T}_H)$  to Problem (4.8) then writes

$$(7.28) \quad u_H^{\text{HHO}} = \sum_{K \in \mathcal{T}_H} \sum_{i=1}^{n_d^{k-1}} f_{K,i} \tilde{\phi}_i^K + \sum_{F \in \mathcal{F}_H^{\text{int}}} \sum_{j=1}^{n_{d-1}^k} u_{F,j} \tilde{\phi}_j^F,$$

where  $f_{K,i}$  is defined as the  $i^{\text{th}}$  coefficient of  $\Pi_K^{k-1}(f_K)$  on the basis  $\{\psi_i^K\}_{1 \leq i \leq n_d^{k-1}}$  for any  $K \in \mathcal{T}_H$ , and  $u_{F,j}$  as the  $j^{\text{th}}$  coefficient of  $u_F := \Pi_F^k(u_H^{\text{HHO}})$  on the basis  $\{\psi_j^F\}_{1 \leq j \leq n_{d-1}^k}$  for any  $F \in \mathcal{F}_H^{\text{int}}$ . The new decomposition (7.28) leads to a simplification of the offline-online solution strategy. In the offline stage, the static condensation step (3) can be bypassed. Also, the steps (1) and (2), which consist in solving saddle-point problems of the form (7.24) and (7.26), are a bit less expensive than before, as the number of Lagrange multipliers is decreased. In the online stage, the reconstruction step (6) can be bypassed, and the global problem to solve in the step (5) simplifies to finding  $(\mathbf{u}_F)_{F \in \mathcal{F}_H^{\text{int}}} := (u_{F,j})_{F \in \mathcal{F}_H^{\text{int}}, j=1, \dots, n_{d-1}^k}$  such that

$$(7.29) \quad \sum_{K \in \mathcal{T}_{F'}} \sum_{F \in \mathcal{F}_K \cap \mathcal{F}_H^{\text{int}}} \mathbb{M}^{F'F} \mathbf{u}_F = - \sum_{K \in \mathcal{T}_{F'}} \mathbb{M}^{F'K} \mathbf{f}_K,$$

for all  $F' \in \mathcal{F}_H^{\text{int}}$ .

**Remark 7.6** (Mono-query case). The purely face-based version of the MsHHO method is particularly suited to the mono-query context. In that case, the step (1) can be bypassed, and replaced by solving, inbetween steps (4) and (5), Problem (7.24) for all  $K \in \mathcal{T}_H$  with right-hand side  $\Pi_K^{k-1}(f_K)$  (in lieu of  $\psi_i^K$ ), whose solution is denoted  $\phi_{f_K}^K$ . Letting  $\mu_{f_K}^{\partial K}$  be the corresponding dual unknown, one must then replace in (7.29) the vector  $\mathbb{M}^{F'K} \mathbf{f}_K$  by the vector  $\mu_{f_K, F'}^{\partial K} \in \mathbb{R}^{n_{d-1}^k}$  formed by the coefficients of the decomposition of  $\mu_{f_K|F'}^{\partial K} \in \mathbb{P}^k(F')$  on the basis  $\{\psi_j^{F'}\}_{1 \leq j \leq n_{d-1}^k}$ . The MsHHO solution in that context is finally given by

$$(7.30) \quad u_H^{\text{HHO}} = \sum_{K \in \mathcal{T}_H} \tilde{\phi}_{f_K}^K + \sum_{F \in \mathcal{F}_H^{\text{int}}} \sum_{j=1}^{n_{d-1}^k} u_{F,j} \tilde{\phi}_j^F,$$

in lieu of (7.28).

7.2.3. *Summary.* The following table summarizes the main computational aspects, in a multi-query context, for both the (fully explicit) MHM and MsHHO methods based on  $\mathcal{U}^k(\mathcal{T}_H)$ ,  $k \geq 1$ , in both the offline and online stages, so as to provide to the reader a one-glance comparison of the two methods. For simplicity, we assume that all the mesh cells have the same number of faces, denoted by  $n_\partial$ .

MHM	offline	local SPD systems	$n_d^{k-1} - 1 + n_{d-1}^k n_\partial$ problems per cell
	online	global saddle-point problem	$\#\mathcal{T}_H + n_{d-1}^k \#\mathcal{F}_H$ unknowns
MsHHO	offline	local saddle-point systems	$n_d^{k-1} + n_{d-1}^k n_\partial$ problems per cell
	online	global SPD problem	$n_{d-1}^k \#\mathcal{F}_H^{\text{int}}$ unknowns

TABLE 1. Comparison of MHM and MsHHO on the main computational aspects

The offline stage is of course performed once and for all, independently of the data (here, the source term). In practice, the approximation of the oscillatory basis functions solution to the local problems can be computationally costly, but the fact that all problems are local makes of the offline stage an embarrassingly parallel task. The offline stage can hence naturally benefit from parallel architectures. In the online stage, the linear systems to solve (for the different data) only attach unknowns to the coarse mesh at hand, hence the computational burden remains limited.

**Remark 7.7** (Other boundary conditions). *The MHM and MsHHO methods easily adapt to the case of (nonhomogeneous) mixed Dirichlet–Neumann boundary conditions. If  $\mathcal{F}_H^{\text{D}} \cup \mathcal{F}_H^{\text{N}}$  forms a (disjoint) partition of  $\mathcal{F}_H^{\text{bnd}}$  into, respectively, Dirichlet and Neumann boundary faces, then the size of the online linear systems in the MHM method becomes  $\#\mathcal{T}_H + n_{d-1}^k \#(\mathcal{F}_H^{\text{int}} \cup \mathcal{F}_H^{\text{D}})$ , whereas that for the MsHHO method becomes  $n_{d-1}^k \#(\mathcal{F}_H^{\text{int}} \cup \mathcal{F}_H^{\text{N}})$ .*

**Remark 7.8** (Second-level discretization). *We have shown that, under the idealized assumption that the local problems defining the multiscale basis functions are exactly solved, the MHM and MsHHO solutions satisfy  $\mathbb{A}\nabla_H u_H \in \mathbf{H}(\text{div}, \Omega)$ . Since the local Neumann problems defining the multiscale basis functions feature polynomial data, we expect such a conformity to be inherited by the two-level method as soon as a mixed method is used to approximate the local second-level problems (on a fine submesh with mesh-size  $h \ll H$ ). This is the approach pursued in [18], but not in [13] and [7] where a primal method is used instead.*

**Remark 7.9** (Strongly varying source terms). *If one wants to be able to deal with strongly varying source terms in a multi-query context, a possibility is to replace the local space  $\mathbb{P}^{k-1}(K)$  by a richer finite-dimensional space, for example  $\mathbb{P}^{k-1}(K_\mathfrak{h})$ , where  $K_\mathfrak{h}$  denotes a mesh (of size  $h \leq \mathfrak{h} \ll H$ ) of the coarse cell  $K$  (see also [35, Section 4] for a basis defined from element-wise generalized eigenvalue problems).*

## 8. CONCLUSION

Although they originate from entirely different constructions, we have proved that the one-level (original) semi-explicit MHM and MsHHO methods provide the same numerical solution when the source term is piecewise polynomial on the (coarse) mesh, and this is also the case for the fully explicit MHM and MsHHO methods for any source term in  $L^2(\Omega)$ . As a byproduct, we have proposed a unified convergence analysis, as well as improved versions of the two methods. More precisely, we have introduced a version of the MHM method that is prompt to be used in a multi-query context, and a version of the MsHHO method that only uses face unknowns.

## REFERENCES

1. A. Abdulle, W. E. B. Engquist, and E. Vanden-Eijnden, *The Heterogeneous Multiscale Method*, Acta Numerica **21** (2012), 1–87.
2. G. Allaire and R. Brizzi, *A multiscale finite element method for numerical homogenization*, SIAM Multiscale Model. Simul. **4** (2005), no. 3, 790–812.

3. R. Araya, G. R. Barrenechea, L. P. Franca, and F. Valentin, *Stabilization arising from PGEM: a review and further developments*, Appl. Numer. Math. **59** (2009), no. 9, 2065–2081.
4. R. Araya, C. Harder, D. Paredes, and F. Valentin, *Multiscale Hybrid-Mixed method*, SIAM J. Numer. Anal. **51** (2013), no. 6, 3505–3531.
5. T. Arbogast and K. J. Boyd, *Subgrid upscaling and mixed multiscale finite elements*, SIAM J. Numer. Anal. **44** (2006), no. 3, 1150–1171.
6. I. Babuška and E. Osborn, *Generalized finite element methods: Their performance and their relation to mixed methods*, SIAM J. Numer. Anal. **20** (1983), no. 3, 510–536.
7. G. R. Barrenechea, F. Jaillet, D. Paredes, and F. Valentin, *The multiscale hybrid mixed method in general polygonal meshes*, Numer. Math. **145** (2020), 197–237.
8. F. Brezzi, M. O. Bristeau, L. P. Franca, M. Mallet, and G. Rogé, *A relationship between stabilized finite element methods and the Galerkin method with bubble functions*, Comput. Methods Appl. Mech. Engrg. **96** (1992), no. 1, 117–129.
9. F. Brezzi, L. P. Franca, T. J. R. Hughes, and A. Russo,  $b = \int g$ , Comput. Methods Appl. Mech. Engrg. **145** (1997), no. 3-4, 329–339.
10. F. Brezzi and A. Russo, *Choosing bubbles for advection-diffusion problems*, Math. Models Methods Appl. Sci. **4** (1994), no. 4, 571–587.
11. Z. Chen and T. Y. Hou, *A mixed multiscale finite element method for elliptic problems with oscillating coefficients*, Math. Comp. **72** (2002), no. 242, 541–576.
12. M. Cicuttin, A. Ern, and S. Lemaire, *A Hybrid High-Order method for highly oscillatory elliptic problems*, Comput. Methods Appl. Math. **19** (2019), no. 4, 723–748.
13. ———, *On the implementation of a multiscale Hybrid High-Order method*, Numerical Mathematics and Advanced Applications—ENUMATH 2017, Lect. Notes Comput. Sci. Eng., vol. 126, Springer, Cham, 2019, pp. 509–517.
14. D. Copeland, U. Langer, and D. Pusch, *From the Boundary Element Domain Decomposition Methods to Local Trefftz Finite Element Methods on Polyhedral Meshes*, Domain Decomposition Methods in Science and Engineering XVIII, Lect. Notes Comput. Sci. Eng., vol. 70, Springer, Berlin, 2009, pp. 315–322.
15. D. A. Di Pietro and A. Ern, *A Hybrid High-Order locking-free method for linear elasticity on general meshes*, Comput. Methods Appl. Mech. Engrg. **283** (2015), 1–21.
16. D. A. Di Pietro, A. Ern, and S. Lemaire, *An arbitrary-order and compact-stencil discretization of diffusion on general meshes based on local reconstruction operators*, Comput. Methods Appl. Math. **14** (2014), no. 4, 461–472.
17. ———, *A review of Hybrid High-Order methods: formulations, computational aspects, comparison with other methods*, Building Bridges: Connections and Challenges in Modern Approaches to Numerical Partial Differential Equations (G. R. Barrenechea, F. Brezzi, A. Cangiani, and E. H. Georgoulis, eds.), Lecture Notes in Computational Science and Engineering, vol. 114, Springer, 2016, pp. 205–236.
18. O. Durán, P. R. B. Devloo, S. M. Gomes, and F. Valentin, *A multiscale hybrid method for Darcy’s problems using mixed finite element local solvers*, Comput. Methods Appl. Mech. Engrg. **354** (2019), 213–244.
19. W. E and B. Engquist, *The Heterogeneous Multiscale Methods*, Commun. Math. Sci. **1** (2003), no. 1, 87–132.
20. Y. Efendiev, J. Galvis, and T. Y. Hou, *Generalized Multiscale Finite Element Methods (GMsFEM)*, J. Comput. Phys. **251** (2013), 116–135.
21. Y. Efendiev and T. Y. Hou, *Multiscale Finite Element Methods - Theory and Applications*, Surveys and Tutorials in the Applied Mathematical Sciences, vol. 4, Springer-Verlag, New York, 2009.
22. Y. Efendiev, T. Y. Hou, and X.-H. Wu, *Convergence of a nonconforming multiscale finite element method*, SIAM J. Numer. Anal. **37** (2000), no. 3, 888–910.
23. Y. Efendiev, R. Lazarov, and K. Shi, *A multiscale HDG method for second order elliptic equations. Part I: Polynomial and homogenization-based multiscale spaces*, SIAM J. Numer. Anal. **53** (2015), no. 1, 342–369.
24. R. Glowinski and M. F. Wheeler, *Domain decomposition and mixed finite element methods for elliptic problems*, First International Symposium on Domain Decomposition Methods for Partial Differential Equations (Philadelphia) (R. Glowinski, G. H. Golub, G. A. Meurant, and J. Périaux, eds.), SIAM, 1988, pp. 144–172.
25. C. Harder, D. Paredes, and F. Valentin, *A family of multiscale hybrid-mixed finite element methods for the Darcy equation with rough coefficients*, J. Comput. Phys. **245** (2013), 107–130.
26. C. Harder and F. Valentin, *Foundations of the MHM method*, Building Bridges: Connections and Challenges in Modern Approaches to Numerical Partial Differential Equations (G. R. Barrenechea, F. Brezzi, A. Cangiani, and E. H. Georgoulis, eds.), Lecture Notes in Computational Science and Engineering, Springer, 2016.
27. P. Henning and D. Peterseim, *Oversampling for the multiscale finite element method*, SIAM Multiscale Model. Simul. **11** (2013), no. 4, 1149–1175.
28. T. Y. Hou and X.-H. Wu, *A multiscale finite element method for elliptic problems in composite materials and porous media*, J. Comp. Physics **134** (1997), 169–189.
29. T. Y. Hou, X.-H. Wu, and Z. Cai, *Convergence of a multiscale finite element method for elliptic problems with rapidly oscillating coefficients*, Math. Comp. **68** (1999), no. 227, 913–943.
30. T. Y. Hou, X.-H. Wu, and Y. Zhang, *Removing the cell resonance error in the multiscale finite element method via a Petrov-Galerkin formulation*, Commun. Math. Sci. **2** (2004), no. 2, 185–205.

31. T. J. R. Hughes, *Multiscale phenomena: Green's functions, the Dirichlet-to-Neumann formulation, subgrid scale models, bubbles and the origins of stabilized methods*, *Comput. Methods Appl. Mech. Engrg.* **127** (1995), 387–401.
32. T. J. R. Hughes, G. R. Feijó, L. M. Mazzei, and J.-B. Quincy, *The variational multiscale method - a paradigm for computational mechanics*, *Comput. Methods Appl. Mech. Engrg.* **166** (1998), 3–24.
33. C. Le Bris, F. Legoll, and A. Lozinski, *MsFEM à la Crouzeix–Raviart for highly oscillatory elliptic problems*, *Chinese Annals of Mathematics, Series B* **34** (2013), no. 1, 113–138.
34. ———, *An MsFEM-type approach for perforated domains*, *SIAM Multiscale Model. Simul.* **12** (2014), no. 3, 1046–1077.
35. A. L. Madureira and M. Sarkis, *Hybrid localized spectral decomposition for multiscale problems*, *SIAM J. Numer. Anal.* **59** (2021), no. 2, 829–863.
36. A. Målqvist and D. Peterseim, *Localization of elliptic multiscale problems*, *Math. Comp.* **83** (2014), no. 290, 2583–2603.
37. L. Mu, J. Wang, and X. Ye, *A Weak Galerkin generalized multiscale finite element method*, *J. Comp. Appl. Math.* **305** (2016), 68–81.
38. D. Paredes, F. Valentin, and H. M. Versieux, *On the robustness of Multiscale Hybrid-Mixed methods*, *Math. Comp.* **86** (2017), no. 304, 525–548.
39. P.-A. Raviart and J.-M. Thomas, *Primal hybrid finite element methods for 2nd order elliptic equations*, *Math. Comp.* **31** (1977), no. 138, 391–413.
40. A. Toselli and O. Widlund, *Domain Decomposition Methods - Algorithms and Theory*, Springer Series in Computational Mathematics, vol. 34, Springer-Verlag, Berlin, 2005.
41. S. Weißer, *BEM-based Finite Element Approaches on Polytopal Meshes*, *Lect. Notes Comput. Sci. Eng.*, vol. 130, Springer, Cham, 2019.
42. M. F. Wheeler, G. Xue, and I. Yotov, *A multiscale mortar multipoint flux mixed finite element method*, *Math. Models Methods Appl. Sci.* **46** (2012), 759–796.