



HAL
open science

Nycthemeral Movements of Wintering Shorebirds Reveal Important Differences in Habitat Uses of Feeding Areas and Roosts

C. Jourdan, Jérôme Fort, D. Pinaud, P. Delaporte, J. Gernigon, N.
Lachaussée, J.-C. Lemesle, C. Pignon-Mussaud, P. Pineau, F. Robin, et al.

► To cite this version:

C. Jourdan, Jérôme Fort, D. Pinaud, P. Delaporte, J. Gernigon, et al.. Nycthemeral Movements of Wintering Shorebirds Reveal Important Differences in Habitat Uses of Feeding Areas and Roosts. *Estuaries and Coasts*, 2021, 44 (5), pp.1454-1468. 10.1007/s12237-020-00871-5 . hal-03240728

HAL Id: hal-03240728

<https://hal.science/hal-03240728>

Submitted on 22 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Nycthemeral movements of wintering shorebirds reveal important differences in habitat uses of feeding areas and roosts

C. Jourdan, J. Fort, D. Pinaud, P. Delaporte, J. Gernigon, N. Lachaussée, J.-C. Lemesle, C. Pignon-Mussaud, P. Pineau, F. Robin, P. Rousseau, and P. Bocher

C. Jourdan, J. Fort, N. Lachaussée, C. Pignon-Mussaud, P. Pineau, and P. Bocher, Laboratory Littoral Environnement et Sociétés UMR LIENSs 7266 CNRS- La Rochelle University, La Rochelle, France. – D. Pinaud Centre d'Etudes Biologiques de Chizé (CEBC), UMR 7372, CNRS-La Rochelle Université, 79360, Villiers en Bois, France. – P. Delaporte and P. Rousseau, National Nature Reserve of Moëze-Oléron, LPO Ligue pour la Protection des Oiseaux, Plaisance, Saint-Froult, France. – J. Gernigon and J.-C. Lemesle, National Nature Reserve of Lileau des niges, LPO, Les Portes en Ré, France. – F. Robin, LPO, Fonderies Royales, Rochefort, France

Corresponding author: C. Jourdan – clement.jourdan1@univ-lr.fr

ORCID iD of the corresponding author: 0000-0002-5477-7494

Abstract

Most shorebirds depend on coastal habitats for much of their life cycle. The quality and diversity of feeding areas during the wintering period directly condition their winter survival, subsequent migration, and breeding success. During their wintering in France, shorebirds use intertidal areas for feeding, both in daylight and at night, depending on the availability of mudflats during the tidal cycle. In this context, we studied whether the bar-tailed godwit (*Limosa lapponica*) shows contrasting foraging behaviors and distributions between day and night in response to differences in visual capacities, prey availability, potential predation risk, and human activities. We carried out a fine-scale GPS tracking of birds at one of their main wintering sites along the French Atlantic coast. We predicted smaller foraging home ranges at night because of limits for godwits to detect prey visually, suggesting more sediment probing and less movement. Godwits used the entire time window when they have access to intertidal areas, but they faithfully selected distinct diurnal and nocturnal feeding areas using a low number of patches. This variability in space use highlights differences in selection of habitats, such as seagrass beds selected by most of the tracked godwits by day and used much less at night. In addition, distinct feeding distributions of monitored birds revealed interindividual variability in habitat selection, even more by night, most likely to reduce intraspecific competition. We therefore urge greater consideration of the night distribution of birds, rarely evaluated in shorebirds studies, to define areas and habitats of importance to future management and conservation measures.

Keywords: Coastal ecology, coastal habitats, intertidal mudflats, GPS tracking, nocturnal foraging, *Limosa lapponica*

1 **Introduction**

2 According to the principle of optimal foraging, animals tend to maximize their net energy intake
3 per unit time (Stephens and Krebs 1986) and thus forage in the best conditions to catch maximal
4 prey with minimal energy lost. In most birds, vision is the main sense used both day and night
5 (Martin 2012). Therefore, the majority of bird species are active during daylight, that is, when
6 visual conditions are best for foraging (Martin 1990). However, foraging may also be
7 influenced by, for instance, predation risk, competition, food availability, weather conditions,
8 and human disturbance, and birds have to adapt their behavior to devise the optimum strategy
9 toward survival (McNamara and Houston 1980; Abrahams and Dill 1989).

10 In coastal shorebirds feeding in intertidal areas, the quality of wintering sites, and
11 especially of feeding areas, directly affects their winter survival, subsequent migration, and
12 breeding success (Pitelka 1979; Piersma et al. 1993; Gunnarsson et al. 2005; West et al. 2005;
13 Gunnarsson et al. 2006; Morrison et al. 2007). The spatiotemporal distribution of birds in
14 intertidal areas therefore depends on prey diversity and abundance (Kelsey and Hassall 1989;
15 VanDusen et al. 2012) as well as their availability (Colwell and Landrum 1993) to maximize
16 the rate of energy intake over the exposure period in such areas (van Gils et al. 2003; Goss-
17 Custard et al. 2007; Quaintenne et al. 2010). Studies on wintering shorebird activity showed a
18 tidal pattern of space use linked with the tidal effect on the availability of intertidal trophic
19 resources (Puttick 1984; Colwell and Landrum 1993; Granadeiro et al. 2006). Thus, shorebirds
20 wintering on sites subject to the tidal cycle will use intertidal foraging areas when available at
21 low tides, that is, both during daylight and night (McNeil et al. 1992; McNeil and Rodriguez
22 1996). This sustained feeding behavior during nocturnal low tides could allow the birds to (1)
23 benefit from better feeding opportunities (prey activity and availability), according to the
24 *preference hypothesis* (Dugan 1981; Evans 1987; Mouritsen 1994), or (2) meet daily energetic
25 requirements not satisfied in daylight, according to the *supplementary hypothesis* (McNeil and

26 Rodriguez 1996; Smith et al. 1999; Sitters 2000). However, in addition to different diurnal and
27 nocturnal foraging conditions, bird visual capacities, such as prey availability or predation risk
28 (Dugan 1981; Evans 1987), could constrain bird foraging patterns and behaviors. Consequently,
29 they should adjust their foraging activity, technique, and space use between daytime and
30 nighttime (Mouritsen 1993; Rojas et al. 1999; Kuwae 2007). Species that can switch from sight
31 to tactile feeding, such as the dunlin (*Calidris alpina*), the Eurasian oystercatcher (*Haematopus*
32 *ostralegus*), the black-tailed godwit (*Limosa limosa*), or the *Tringa* species (Mouritsen 1994;
33 McNeil and Rodriguez 1996; Lourenço et al. 2008) could take advantage of nocturnal feeding.
34 Sight-feeding shorebirds could also benefit from night feeding because of the moonlight
35 (McNeil et al. 1992) and/or physiological adaptations improving their nocturnal vision
36 (Pienkowski 1983; Rojas et al. 1999) to maintain a high intake rate during nighttime
37 (Pienkowski 1983; Lourenço et al. 2008). Furthermore, the higher activity and accessibility of
38 prey at night in intertidal mudflats (McNeil et al. 1995; Esser et al. 2008), combined with the
39 aforementioned bird capacities, could explain such an interest in nocturnal foraging behavior
40 to satisfy wintering energetic requirements.

41 Both diurnal and nocturnal foraging areas are therefore important for wintering
42 shorebirds. However, most studies focused on their daylight activity and distribution,
43 disregarding possible important areas used only during nighttime yet also essential for adapted
44 conservation and management measures to protect these vulnerable species. In that context, we
45 studied the diurnal and nocturnal spatial distributions of the bar-tailed godwit (*Limosa*
46 *lapponica lapponica*) wintering on Ré Island (Pertuis Charentais, Atlantic French coast), a site
47 of national importance for the species. The population of *L. l. lapponica* is estimated at 120,000
48 (Delany et al. 2009), and France sees 5.8% of wintering godwits, among which about 650 (>1%
49 of national numbers) are recorded on Ré Island in January (J-C. Lemesle, *Pers. Com.*). The bar-
50 tailed godwit is characterized by increasing trends of the wintering population at the European

51 scale (BirdLife International 2015), and its numbers have been stable in France in the last 12
52 years (Schmaltz et al. 2019). Nonetheless, large fluctuations of the population have been
53 observed in recent decades in France, including significant decreases during the 1980s and
54 1990s (Triplet et al. 2010) likely due to degraded wintering habitats and increasing human
55 disturbance (Goeldner-Gianella 2005; Delany et al. 2009; BirdLife International 2017). In this
56 context, we tested the hypothesis that the bar-tailed godwit shows contrasting foraging
57 behaviors and distributions between day and night. Indeed, visual capacities, prey availability,
58 predation risk, and human activities are all factors influencing the behavior of birds, and their
59 variability between day and night could lead to differences in their distribution. We predicted
60 smaller foraging home ranges at night because of birds' difficulty in visually detecting prey
61 (Turpie and Hockey 1993), suggesting more sediment probing and therefore less movement
62 (Pienkowski 1983; Lourenço et al. 2008). To achieve these objectives, we conducted a fine-
63 scale GPS tracking of wintering habitat use and investigated the birds' (1) spatial distribution
64 and foraging home ranges, (2) feeding movements, (3) habitat selection, and (4) their variation
65 between daytime and nighttime.

66

67 **Methods**

68 **Study site**

69 Fieldwork was carried out in the Pertuis Charentais, which includes the largest area of intertidal
70 mudflats on estuarine systems in France (Goeldner-Gianella 2005), during the wintering period
71 (from November to March) of shorebirds in 2015–2016 and 2016–2017 (Fig. 1). We focused
72 on Ré Island, one of the main wintering sites for bar-tailed godwits in the Pertuis Charentais.
73 On Ré Island, godwits are known to roost in the former saltpans of the National Nature Reserve
74 of Lilleau des Niges and use two main feeding areas on soft substrates (Aubouin 2014; Duijns
75 et al. 2014). The first feeding site is a mudflat inside a semienclosed bay classified according
76 to the Ramsar Convention since 2003, the “Fier d’Ars” (46°13'18"N; 1°30'29"W) (Fig. 1). A
77 part of the “Fier d’Ars” (the western part of the bay and the high-tide roosts in saltmarshes) is
78 also included in the Nature Reserve. The second main feeding site is “La Loge” (46°14'25"N;
79 1°28'42"W), a sandflat on the exposed coast north of the Island and where the foreshore remains
80 uncovered by the tide for about four hours only, thus reducing food availability (Fig. 1). “La
81 Loge” is part of the Ramsar labeled site “Marshes of the Fier d’Ars” but does not benefit from
82 any protection status, and recreational activities on the beach in the summer, as well as in the
83 winter to a lesser extent, are frequent during the day.

84 **Capturing and tracking godwits**

85 Bar-tailed godwits were captured using mist nets during nonmoonlight nights inside the Nature
86 Reserve. Godwits were immediately marked with a metal ring and a unique color ring
87 combination, and body mass (accuracy within 1.0 g), wing length (nearest 1 mm), tarsus length
88 (nearest 0.5 mm) and bill length (nearest 0.5 mm) were measured using standard methods
89 (Prater et al. 1977; Evans 1986). When possible, the sex (bill length: 69–90 mm for males and
90 86–110 mm for females; mass: 230–383 g for males and 280–455 g for females) and age (fringe

91 of median covert feathers is pale with brown tips in juveniles, white in adults; primary flight
92 feathers are worn in juveniles, new in adults) of individuals were determined (Pyle 2008;
93 Demongin 2020). The heaviest godwits captured, which were all females, were fitted with a
94 GPS-VHF logger (STERNA VHF-SRD with solar charger, Ecotone, Gdansk, Poland; 35 × 16
95 × 10 mm, 7.5 g) when the mass of the tag remained less than 3% of the bird body mass. In total,
96 15 individuals (11 in 2015 and 4 in 2016) were equipped with a GPS logger. Juvenile females,
97 potentially not faithful to the study site during winter, were not selected. Tags (accuracy of ±10
98 m) were fixed on birds with a 2 mm Teflon harness (on the lower back) according to the “leg-
99 loop” method (Mallory and Gilbert 2008) and were programmed to record positions every 30
100 minutes. Six of the 15 individuals provided a sufficient number of positions over the winter and
101 were retained for analyses. Three individuals provided data during both winters, two others
102 during winter 2015–2016, and a last one during winter 2016–2017. For individuals with two
103 winter surveys, we retained the data of the winter with the best balance of GPS fixed numbers
104 between day and night. Thus, we used the data of winter 2015–2016 for BTG01, BTG02,
105 BTG04, and BTG05, and the data of winter 2016–2017 for BTG03 and BTG06. The monitoring
106 period extended from November to March for four individuals and from December to March
107 for the other two (Table 1). The data were stored and processed from a PostgreSQL/PostGIS
108 database.

109 **Habitat mapping**

110 For each bird monitored, sediment core sampling was performed around the centroid of several
111 feeding areas defined by GPS locations collected during the first two months after the birds
112 were fitted with transmitters (see below for details). This sampling method, carried out on a
113 grid of 9 cores spaced by 10 meters and arranged around the central reference point of the
114 station, should describe the availability and quality of trophic resources (distribution and
115 density of benthic macrofauna) at the feeding areas used by godwits. The samples were

116 collected by foot at low tide according to methods described in Bocher et al. (2007) and
117 Bijleveld et al. (2012). Each sample consisted of a 15 cm diameter sediment core (0.01 m²),
118 with a depth of 15 cm (maximum depth reached by foraging birds according to mean bill length
119 of females), sieved over a 1 mm mesh size on site. Annelids were immediately preserved in
120 70% ethanol, and mollusks were stored at -20°C until sorted at the laboratory. Mudsnaails
121 (*Hydrobia ulvae*), a potentially small and abundant prey, were sampled through an additional
122 core (70 mm diameter) of 0.0037 m², to a depth of 5 cm, and sieved in the laboratory over a 0.5
123 mm mesh. A larger sediment core sampling for mudsnails would be a time-consuming process.
124 In the laboratory, the organisms were identified to the species level as often as possible, and
125 mollusks were measured within a 0.1 mm accuracy.

126 Mean grain size (mm) and percentage of silt (fraction < 63 µm) of a sediment core for each
127 sampling station (depth of 5 cm) was determined using a Malvern Mastersizer 2000 diffraction
128 laser (particle sizes analyzed from 0.04 to 2,000 µm) to characterize the granulometry of the
129 substrate. The results of the sediment's particle size characteristics helped define each sampling
130 station in terms of habitat type and thus build a habitat map for the two study areas.

131 Finally, the benthic macrofauna data, sediment characteristics, and field observations allowed
132 us to build a fine-scale map of intertidal habitat typology in the study site using the European
133 Nature Information System (EUNIS) classification of coastal habitats as a reference (Bajjouk
134 et al. 2015). Seagrass beds were delimited on site using GPS Trimble GeoXH during summer
135 2012.

136 **Home ranges and habitat selection**

137 The estimation of the utilization distribution (UD), that is, the probability of finding each bird
138 at any location (Calenge 2015), was used to analyze diurnal and nocturnal space use by godwits.
139 Kernel density estimates (KDE) (Van Winkle 1975; Worton 1989) were used to describe UD

140 (Laver and Kelly 2008) with the “kernelUD” function from the R package “adehabitatHR”
141 (Calenge 2015). The spatial distribution of bird locations and the corresponding time spent in
142 an area are considered by KDE method to estimate the home ranges (at 95% isopleth) and core
143 areas (at 50% isopleth) of each individual (Worton 1989). To estimate the kernel home ranges,
144 a compromise between the undersmoothing of least-square cross-validation (LSCV) and the
145 over-smoothing of reference bandwidth (ad hoc) was used: a smoothing factor of 70% of the
146 minimum reference value obtained by the “ad hoc” bandwidth (Kie et al. 2010; Schuler et al.
147 2014). Using a single smoothing factor ($h = 80$), calculated on the set of GPS data and used to
148 estimate bird kernels, allowed us to compare UD_s between individuals. Considering the
149 accuracy of the GPS positions (± 10 m), the grid size was set to 20 m. After estimating global
150 UD_s, the specific foraging distribution and resting distribution were computed. Field
151 observations allowed to define a spatial delimitation on upper intertidal areas above which birds
152 were mainly at rest and not foraging. Thus, foraging distribution was estimated from the GPS
153 positions of birds located, at low tide, on intertidal areas below an elevation of 3.3 m relative to
154 the hydrographic zero. In the same way, resting distribution was computed from the GPS
155 positions of birds located on saltpans, ponds in marshes, or upper intertidal areas above an
156 elevation of 3.3 m. Sunrise and sunset data from the R package “GeoLight” were used to
157 distinguish diurnal and nocturnal positions and allowed for the computation of birds’ foraging
158 and resting distribution by differentiating day and night.

159 UD_s were then used to investigate the importance of foraging habitats through habitat selection
160 analysis. For this, we considered a second-order selection (design II), that is, a same availability
161 of habitats for all birds and an analysis of habitat selection at the individual scale (Johnson
162 1980). A minimum convex polygon (MCP) of foraging points from all monitored birds
163 provided an estimation of available foraging habitats for godwits. Individual kernel home
164 ranges and the habitat typology of the study area were then superimposed to perform habitat

165 selection analysis through Manly's selection ratios (Manly et al. 2002) and using the R package
166 "adehabitatHS" (Calenge 2011). We explored the interindividual variability of habitat selection
167 through an Eigen analysis of selection ratios (Calenge and Dufour 2006), a multifactorial
168 method. These analyses and other statistical tests (ANOVA, paired t-tests and chi-squared tests)
169 were performed with the software R (3.6.1).

170

171 **Results**

172 **Day and night feeding areas**

173 Godwits showed an extreme fidelity to their feeding grounds along the wintering period (from
174 November to March), with a limited number of prospected areas per individual, both during
175 daylight and night. The mean sizes of diurnal feeding home ranges (dFHR, kde 95%) and
176 diurnal feeding core areas (dFCA, kde 50%) were 224 ± 77 ha (min–max: 125–321 ha) and 45
177 ± 18 ha (22–68 ha), respectively (Table 1, Fig. 2). The mean sizes of nocturnal feeding home
178 ranges (nFHR, kde 95%) and nocturnal feeding core areas (nFCA, kde 50%) were, compared
179 with diurnal ones, much smaller with 128 ± 72 ha (36–226 ha) and 23 ± 12 ha (8–34 ha),
180 respectively (dFCA/nFCA: $t = 6.535$, $df = 5$, $p\text{-value} = 0.001$; dFHR/nFHR: $t = 9.0391$, $df = 5$,
181 $p\text{-value} < 0.001$). In addition, the overlaps between diurnal and nocturnal home ranges were
182 low at an individual scale (Table 1). The mean individual day–night overlap was $35 \pm 14\%$
183 (min–max: 12%–54%) between dFHR and nFHR and $18 \pm 11\%$ (min–max: 5%–34%) between
184 dFCA and nFCA (Table 1).

185 **Rhythm of feeding activity**

186 The recorded bird foraging positions indicated that godwits could start foraging mainly between
187 2 and 3 hours before the low tide and 3 and 4 hours after the low tide, both during daylight and
188 night (Fig. 3). From 2 hours before and 3 hours after low tide, birds spent about 95%–100% of
189 their time on foraging areas during daylight and 90%–95% during the night except BTG05 with
190 80%–85% and 75%–80%, respectively. The proportion of time spent on the foraging area
191 during the ebb tide (*c.* 6 hours) increased continuously during daylight, from 5%–10% to 10%
192 and 15% during the first 2 hours, to 50% during the third hour. At night, godwits spent a steady
193 proportion of time on the foraging area (around 0%–5%) during the first 2 hours of the ebbing
194 tide but suddenly went up to 40% between the third and fourth hours. On average, godwits

195 therefore joined the foraging areas earlier by daylight, spending significantly more time on
196 intertidal areas during the day than at night during the fourth and third hour before the low tide,
197 with an average of 11% and 12% of additional time (Fig. 3). Similarly, godwits remained longer
198 on mudflats at the end of the daylight rising tide compared to the nighttime, with 6% and 7%
199 additional time spent on the foraging areas over the third and fourth hour after low tide,
200 respectively, although differences were not significant (Fig. 3). Finally, considering the entire
201 tidal cycle (from -6 hours to 6 hours around low tide), bar-tailed godwits spent a higher percent
202 of their time on foraging grounds by day than by night (paired t-test: $t = 5.35$, $df = 5$, $p\text{-value} =$
203 0.003).

204 **Daylight and night local movements**

205 The distances measured between the two successive recorded locations on the feeding areas
206 were shorter at night than during daylight for all individuals ($t\text{-value} = -3.82$, $p\text{-value} < 0.001$;
207 $\text{meanNight} = 298$ m, $\text{meanDay} = 411$ m). In proportion, godwits systematically made more
208 small movements (or displacements) at night than during the daylight. For nocturnal
209 movements, 0–50 m distance class was the most represented, and most direct distances recorded
210 between two consecutive locations (30 min) were less than 100 m (Fig. 4). By day, most
211 distances between points on the feeding area exceeded 100 m, and the most represented distance
212 class was 100–150 m (Fig. 4).

213 **Habitat diversity and structure**

214 In total, 11 locations, as potential feeding habitats, were identified in the intertidal areas of the
215 study site (Fig. 5). The “Fier d’Ars” area included a large diversity of habitats, with a clear
216 dominance of seagrass beds (A2.6111 – 27% of the total intertidal site surface), bare mudflat
217 (A2.313 – 19%), and oyster parks (A2.32 – 8%) on the muddy foreshore at the center of the
218 bay (Fig 5.b). Habitat A2.6111 was characterized by a fine muddy sand substrate with an
219 abundance of *Zostera noltei* and infaunal species dominated by polychaetes (*Scoloplos armiger*

220 and *Arenaria marina*), oligochaetes, and mollusks (*Cerestoderma edule*, *Macoma balthica*, and
221 *Hydrobia ulvae*). Habitat A2.313 corresponded to a littoral sandy mud, mainly characterized by
222 polychaetes (*Hediste diversicolor*) and bivalves (*M. balthica* and *Scrobicularia plana*). A2.32
223 was a muddy substrate mainly characterized by a low diversity of polychaetes and oligochaetes,
224 covered with artificial oyster grow-out tables. Another habitat well represented in the “Fier
225 d’Ars” was an intertidal muddy sandflat dominated by polychaetes (*Eteone longa*, *S. armiger*)
226 and capitellidae combined with the bivalve *C. edule* (A2.242 – 6%). Saltmarsh creeks in the
227 A2.5 coastal saltmarshes (A2.325 – 17%) surrounded the muddy bay while rocky elements, that
228 is, bedrock and boulders dominated by mussels and/or barnacle communities (A1.11 – 1%) and
229 habitats with littoral rock features (A1.41 – 6%) characterized the bay entrance.

230 The site of “La Loge” contained specific habitats not found on “Fier d’Ars” with clearly sandy
231 characteristics (Fig. 5.c). These habitats were A2.221 (17% of the total study intertidal site
232 surface), a barren coarse sandbank that can only shelter an extremely small number of
233 oligochaetes because of the constant mobility of the coarse sediment, and A2.231 (38%), a
234 littoral fine sand dominated by polychaetes including *Nephtys cirrosa* and *S. armiger*. The
235 northern part of “La Loge” was described by habitats A1.41 (composed of rocky elements and
236 sheltering littoral rockpool communities – 21%) and A2.242 (dominated by polychaetes *E.*
237 *longa*, *S. armiger*, and capitellidae, combined with the bivalve *C. edule* – 18%).

238 **Foraging habitat selection**

239 During daylight, godwits did not use available foraging habitats randomly but showed a
240 significant habitat selection, both at an individual scale ($\chi^2 = 47563.98$, $df = 9.0$, $p < 0.001$) and
241 when combining the six birds monitored ($\chi^2 = 51039.79$, $df = 54.0$, $p < 0.001$). In the diurnal
242 feeding core areas, the seagrass bed habitat (A2.6111) was selected by four godwits (BT01,
243 BTG02, BTG03, and BTG04) (Fig. 6). For BTG05 and BTG06, the most represented habitats
244 were the sandflat dominated by *C. edule* and polychaetes (A2.242) and the mudflat dominated

245 by *H. diversicolor* and *S. plana* (A2.313), respectively. Habitat A2.313 was also a prevailing
246 habitat in the core feeding areas of BTG02 and BTG05. Hence, habitat A2.6111 appeared
247 preferentially used by godwits on average, followed by A2.242 and A2.313 (global selection
248 ratios > 1) (Fig. 7). Conversely, habitat A1.31 (fucoids on sheltered shores) was globally
249 avoided by the studied birds (global selection ratios < 1) although standard errors indicate their
250 marginal use by some birds. The six remaining habitats were clearly avoided (global selection
251 ratios and SE < 1).

252 During nighttime, the birds also showed significant habitat selection when considering all birds
253 ($\chi^2 = 38081.9$, $df = 54.0$, $p < 0.001$) and individuals independently ($\chi^2 = 31774.09$, $df = 9.0$, p
254 < 0.001). Habitats in the nocturnal feeding core areas were more specific, with strong individual
255 patterns. The feeding core areas of BTG01, BTG02, BTG03, BTG04, BTG05, and BTG06 were
256 mainly composed of habitats of oyster parks (A2.32), bare mudflat (A2.313), intertidal muddy
257 sandflat (A2.242), littoral fine sand (A2.231), bare mudflat (A2.313), and seagrass beds
258 (A2.6111), respectively, highlighting a strong interindividual variability (Fig. 6). This
259 observation was confirmed by the result of the Manly selectivity measure, which did not allow
260 a clear identification of a habitat preferentially used by the six birds. Indeed, habitats of littoral
261 fine sand (A2.231), intertidal muddy sandflat (A2.242), bare mudflat (A2.313), oyster parks
262 (A2.32), and seagrass beds (A2.6111) all appeared selected (global selection ratios > 1) but
263 with large standard errors that well illustrate the nonidentical use of habitats by all birds ($\chi^2 =$
264 941.584 ; $df = 45.0$; $p < 0.001$) (Fig. 7). Although seagrass beds (A2.6111) remain globally the
265 most selected habitat at night (Fig. 7), we note that all individuals showed a significant decrease
266 of its use between day and night, with a proportion decreasing from an average of 48% to 19%,
267 except BTG06, which showed an opposite trend (Fig. 6).

268 The Eigen analysis emphasized these differences between diurnal and nocturnal habitat
269 selection. During daytime, the reported positions of four individuals (BTG01, BTG02, BTG03,

270 and BTG04) in the space described by habitats highlighted their preferential use of seagrass
271 beds (A2.6111) (Supplementary Material Fig. S1). The other two birds stood out from this first
272 group, with BTG06 clearly selecting the sandflat habitat dominated by *C. edule* and polychaetes
273 (A2.242) and BTG05 in an intermediate position between A2.6111 and A2.242. At night, each
274 individual selected a specific foraging habitat with five different habitats, thus preferentially
275 used by godwits at night (A2.242, A2.231, A2.313, A2.6111, and A2.32).

276 **Roost selection**

277 Godwits' use of roosts differed markedly between day and night. The number of roosts used by
278 individuals was not different between day and night, with one to three main roosts according to
279 the estimation of roost core areas (isopleth 50) (Fig. 8). However, the location clearly varied
280 between diurnal and nocturnal stages with a mean overlapping rate of 32% for roosting home
281 ranges (RHRs) and 8% for roosting core areas (RCAs) (Supplementary Material Table S1). By
282 day, during high neap tide, birds preferentially roosted on the upper foreshore and used less
283 significantly the saltpan roosts beyond dikes. At night, individuals clearly selected the saltpans
284 of the Nature Reserve as roosts and avoided free intertidal areas during high neap tide. During
285 spring tide, they could not stay on flooded intertidal areas and returned on saltpan roosts either
286 by day or night.

287

288 **Discussion**

289 By analyzing the nycthemeral use of habitats by wintering bar-tailed godwits at an
290 extremely fine spatial scale, the present study revealed distinct foraging activities in shorebirds
291 between day and night. Previous studies showed that shorebirds could use different feeding and
292 roosting areas depending on the time of day and according to predation risk, disturbance, and
293 density/activity of preys (Burton and Armitage 2005; Piersma et al. 2006). Here, we went
294 further and showed that bar-tailed godwits faithfully selected distinct diurnal and nocturnal
295 feeding areas using a low number of main feeding patches (CA), both during daylight and
296 nighttime, with a relatively small feeding area by individuals. Moreover, birds moved less when
297 prospecting at night, resulting in smaller nocturnal feeding home ranges. This variability in
298 space use underlined differences in habitat selection, such as seagrass beds strongly selected by
299 most of the tracked godwits by day and much less used at night. In addition, the distinct feeding
300 distributions of the monitored females revealed interindividual variability in habitat selection.
301 We noted that all our tracked birds were females, and there could be large differences in diet
302 and patterns of space use between males and females during day and night.

303 Our study highlights that bar-tailed godwits spent as much time on feeding areas at night as
304 during the day, devoting most of it to feeding activities from two hours before to three hours
305 after low tide, that is, during the entire time window of mudflat availability. As observed in
306 black-tailed godwits (Lourenço et al. 2008), this result highlights the importance of nocturnal
307 feeding in the acquisition of daily energy needs by wintering bar-tailed godwits. Night foraging
308 activity has been reported in most shorebird species wintering on tidal coasts although the
309 relative importance of nocturnal activity depends on the species (Dugan 1981; McNeil and
310 Rodriguez 1996). Many species favor feeding activity during the day, with visual capabilities
311 allowing for higher intake rate and consequently providing the major part of the daily energy
312 needs for birds (Dodd and Colwell 1996; Kam et al. 2004; Lourenço et al. 2008). Night foraging

313 was therefore mainly considered as “supplementary” in previous studies, that is, used when
314 diurnal prey intake is not enough to offset the increased energy demands of less favorable
315 wintering conditions (McNeil and Rodriguez 1996; Smith et al. 1999; Sitters 2000). However,
316 some shorebirds could fill a significant part of their energy requirements during the night
317 (Turpie and Hockey 1993; Lourenço et al. 2008) because of physiological adaptations such as
318 better nocturnal vision for *Charadriidae* (high rod–cone ratio in the eyes) (De Azuje et al. 1993;
319 Rojas et al. 1999) or higher tactile sensitivity for *Scolopacidae* as in godwits (Cunningham et
320 al. 2013), which facilitate nocturnal foraging.

321 In our study, the space used by wintering godwits differed between day and night, both
322 during feeding and resting periods, as reported for other shorebirds in previous studies (Rompré
323 and McNeil 1996; Piersma et al. 2006; Rogers et al. 2006). Birds were highly faithful to their
324 daylight feeding grounds, with extremely restricted feeding core areas recorded over the winter.
325 Their nocturnal feeding core areas were even more restricted and mostly spatially distinct from
326 the daytime ones. These results contrast with the observations of Burton and Armitage (2005)
327 on redshanks (*Tringa tetanus*), which also used separate diurnal and nocturnal foraging areas
328 but with larger core areas and home ranges at night. In addition, the distinct distributions of bar-
329 tailed godwits between day and night were associated with a change in habitat selection, notably
330 with a strong decrease in the use of seagrass beds at night, as well as an increased use of oyster
331 parks and sandflats. Various factors that we consider below, mainly related to the characteristics
332 of wintering habitats, could explain these differences in space use between day and night.

333 Contrary to studies that highlighted an avoidance of nearshore areas by shorebirds during
334 their nocturnal foraging in response to increased predation risks (Sitters et al. 2001; Burton and
335 Armitage 2005; Piersma et al. 2006), we did not find that upper intertidal areas were less used
336 by godwits at night. Conversely, a portion of monitored birds tended to focus on strategic points
337 of the upper foreshore during nocturnal feeding, near anthropogenic illuminations. Studies

338 highlighted a positive effect of artificial light on the nocturnal foraging of shorebirds by
339 recreating full-moon conditions across the nearby intertidal areas that allow birds to maintain a
340 sight-based foraging behavior at night and improve their prey intake rate (Santos et al. 2010;
341 Dwyer et al. 2013). On Ré Island, public lighting points at the edge of the bay increase the level
342 of ambient light across mudflat areas and could attract birds by allowing them to feed by sight
343 (Lourenço et al. 2008; Santos et al. 2010).

344 Recreational and harvesting activities represent another form of disturbance related to
345 anthropogenic activities, which has already been shown to affect the habitat use and foraging
346 activity of wintering or migrating shorebirds (Lafferty 2001; Colwell et al. 2003; Navedo and
347 Masero 2007; Burger and Niles 2013). For instance, studies emphasized that recreational
348 activities on the foreshore, as well as shellfish harvesting activities, could induce a reduction of
349 foraging time, flight behavior, or even a complete avoidance of the disturbed foraging areas
350 (De Boer and Longamane 1996; Thomas et al. 2003; Navedo and Masero 2007; Burger and
351 Niles 2013). This pattern is what we observed for some godwits on Ré Island, which avoided
352 oyster parks (A2.32) and beaches (A2.231) during daytime while they commonly used these
353 habitats at night. With a high predominance of polychaetes in the macrobenthic community,
354 these habitats constitute attractive feeding areas for godwits, but the presence of oyster farmers,
355 shell fishers, and walkers during the day constitutes a significant level of disturbance (Burton
356 and Armitage 2005; Dias et al. 2008).

357 Contrasting foraging methods between day and night were also found to affect the spatial
358 distribution of shorebirds. For instance, Mouritsen (1993) found that the dunlin (*C. alpina*)
359 switched from visual feeding during the day to tactile feeding during the night, as did black-
360 tailed godwits which showed a higher rate of stitches and sweeps at night and a majority of
361 pecks during the day (Lourenço et al. 2008). Generally, the bill of the *Scolopacidae* species
362 benefits from a high number of touch-sensitive nerve endings which favors tactile feeding (De

363 Azuaje et al. 1993). Hence, we can assume that bar-tailed godwits switch from sight feeding
364 during the day to probing at night and, as observed in dunlins (*C. alpina*) by Mouritsen
365 (Mouritsen 1994), this change in behavior may induce the use of distinct feeding habitats based
366 on prey availability. This could be related to the dominance of polychaetes, more active and
367 likely more available at night (Last and Olive 2004; Kuwae 2007) in habitats A2.32 and A2.231,
368 which could explain the nocturnal use of these habitats by godwits. We did not have available
369 data to compare diurnal and nocturnal prey density, but studies underlined that polychaete
370 worms, the favored preys of bar-tailed godwits (Duijns et al. 2013), may be closer to the
371 sediment surface at night (McNeil et al. 1992; Esser et al. 2008). Such a density of prey
372 available in the top layer of the sediment could explain the nocturnal preference of polychaete-
373 dominated habitats by godwits, as observed in dunlins and redshanks (Mouritsen 1994; Burton
374 and Armitage 2005). Beyond their influence on the location of godwits' feeding areas, the
375 differences in prey availability and foraging methods between day and night also seemed to
376 affect the size of foraging areas.

377 On night foraging sites, higher prey availability could lead to better intake rates (Zharikov
378 and Skilleter 2003), resulting in an area-restricted search for tactile predators through increased
379 spatial turning rates (Dias et al. 2009). Godwits could thus use sinuous low-speed searches in
380 more profitable patches at night (Nolet and Mooij 2002) and prefer faster direct searches with
381 more step rates during the day as observed in black-tailed godwits (Lourenço et al. 2008). This
382 pattern of feeding behavior could explain the differences observed in our study in the distances
383 separating two successive foraging locations between day and night.

384 At high tide, results showed marked differences in the use of roosts between day and night
385 during neap and spring tides. Indeed, at daytime, godwits selected upper mudflat roosts as soon
386 as they were available, that is, during neap tides. However, at night, birds always used saltpans
387 in whatever tidal heights. In a previous study based on daytime observations, Rosa et al. (2006)

388 showed that wintering shorebirds, including bar-tailed godwits, preferentially select mudflats
389 to roost, and move to saltpan roosts when the upper foreshore became unavailable during
390 highest tides. This study also highlighted the effect of both raptor presence, higher in salt pans
391 than in mudflats, and visibility, lower in salt pans, on daytime shorebird roost choice (Rosa et
392 al. 2006). Others studies reported the anti-predator strategy of shorebirds avoiding some
393 specific roosting areas by nighttime because of higher predation risks (Hilton et al. 1999;
394 Rogers et al. 2006). Hence, in addition to their availability, the roost choice by shorebirds
395 between day and night in the present study could be influenced by the predation risk (Handel
396 and Gill 1992; Rohweder 2001). Birds would favor mudflats at day, less prone to overflying
397 raptors and offering better visibility of approaching predators (Rosa et al. 2006), and salt pans
398 at night, with a water barrier against nocturnal predators such as foxes or mustelids (Cramp et
399 al. 1983; Sitters et al. 2001). Further studies are nonetheless needed to validate this hypothesis
400 and better explain the pattern of use of roost sites on Ré Island, including the monitoring of
401 predator activities.

402 Our results showed that all monitored bar-tailed godwits used the entire time window during
403 which they have access to foraging grounds, both during the day and the night. Nocturnal
404 foraging thus appears crucial along the winter to allow godwits to meet their daily energy
405 requirements and is not only a “supplement” for diurnal foraging. However, nocturnal foraging
406 is probably not preferential since we did not observe more use of feeding areas at night than
407 during the day. We hypothesize that differences in biotic and abiotic environmental conditions
408 (human disturbance, predation risk, feeding methods, and prey availability) between day and
409 night all together incite birds to develop specific feeding strategies and behaviors, including the
410 selection of contrasting habitats. Bar-tailed godwits also tended to specialize in their habitat use
411 and thus in prey at an individual scale, even more by night, most likely to reduce intraspecific
412 competition. Roost choice also depended on the time of day. The visibility of approaching

413 predators and proximity to feeding sites, when possible, seemed to be important to the choice
414 of daytime roosts. Finally, these results provide an important knowledge of the nonbreeding
415 survival strategies of bar-tailed godwits on the French Atlantic coast. We therefore urge greater
416 consideration of the night distribution of birds, rarely evaluated in shorebird studies, to define
417 areas and habitats of importance in management and conservation. The day–night connectivity
418 in shorebirds’ space use needs to be integrated into all spatial management plans where human
419 activities can deal with natural protected areas or their proximities.

420 **Acknowledgments** – We thank Jérôme Jourde for his help in determining the benthic
421 macrofauna, Tania Damany and Sébastien Comarmond for their help with sample processing,
422 and Françoise Amélineau and Chloé Tanton for their help during bird capture/marketing sessions.
423 We also thank all the volunteers who participated in bird captures and sediment sampling. This
424 work has been supported by the ECONAT project funded by the Contrat de Plan Etat-Région
425 and the CNRS and the European Regional Development Fund (QUALIDRIS project). This
426 work was also funded by the ANR Pampas (ANR-18 CE32-0006) and the Ligue pour la
427 Protection des Oiseaux. We thank Christine Dupuy and Christel Lefrançois for the logistic
428 support. All work adheres to the legal requirements of the countries in which it was carried out
429 and meets ethical and animal welfare guidelines.

430

431 **References**

- 432 Abrahams, M. V., and L. M. Dill. 1989. A determination of the energetic equivalence of the
 433 risk of predation. *Ecology* 70: 999–1007. <https://doi.org/10.2307/1941368>.
- 434 Aubouin, Naïs. 2014. *Etude de la distribution et des stratégies de survie hivernale de*
 435 *populations migratrices d’oiseaux limicoles en vue d’actions de conservation en*
 436 *région Poitou Charentes: exemple de la Barge rousse Limosa lapponica*. Msc Thesis
 437 64p.
- 438 De Azuaje, L. M. R., S. Tai, and R. McNeil. 1993. Comparison of rod/cone ratio in three
 439 species of shorebirds having different nocturnal foraging strategies. *Auk*: 141–145.
- 440 Bajjouk, T., B. Guillaumont, N. Michez, B. Thouin, C. Croguennec, J. Populus, J. Louvel-
 441 Glaser, et al. 2015. *Classification EUNIS, Système d’information européen sur la*
 442 *nature : Traduction française des habitats benthiques des Régions Atlantique et*
 443 *Méditerranée. Vol. 1. Habitats Littoraux*.
- 444 Bijleveld, A. I., J. A. van Gils, J. van der Meer, A. Dekinga, C. Kraan, H. W. van der Veer,
 445 and T. Piersma. 2012. Designing a benthic monitoring programme with multiple
 446 conflicting objectives. *Methods in Ecology and Evolution* 3: 526–536.
 447 <https://doi.org/10.1111/j.2041-210X.2012.00192.x>.
- 448 BirdLife International. 2015. *European red list of birds*.
- 449 BirdLife International. 2017. *Limosa lapponica* (amended version of 2016 assessment). *The*
 450 *IUCN Red List of Threatened Species 2017*.
- 451 Bocher, P., T. Piersma, A. Dekinga, C. Kraan, M. G Yates, T. Guyot, E. Folmer, and G.
 452 Radenac. 2007. Site-and species-specific distribution patterns of molluscs at five
 453 intertidal soft-sediment areas in northwest Europe during a single winter. *Marine*
 454 *Biology* 151: 577–594. <https://doi.org/10.1007/s00227-006-0500-4>.
- 455 De Boer, W. F., and F. A. Longamane. 1996. The exploitation of intertidal food resources in
 456 Inhaca Bay, Mozambique, by shorebirds and humans. *Biological Conservation* 78:
 457 295–303. [https://doi.org/10.1016/S0006-3207\(96\)00050-X](https://doi.org/10.1016/S0006-3207(96)00050-X).
- 458 Burger, J., and L. Niles. 2013. Shorebirds and stakeholders: Effects of beach closure and
 459 human activities on shorebirds at a New Jersey coastal beach. *Urban Ecosystems* 16:
 460 657–673. <https://doi.org/10.1007/s11252-012-0269-9>.
- 461 Burton, N. H. K., and M. J. S. Armitage. 2005. Differences in the diurnal and nocturnal use of
 462 intertidal feeding grounds by Redshank *Tringa totanus*. *Bird Study* 52: 120–128.
 463 <https://doi.org/10.1080/00063650509461381>.
- 464 Calenge, C., and A. B. Dufour. 2006. Eigenanalysis of selection ratios from animal radio-
 465 tracking data. *Ecology* 87: 2349–2355. [https://doi.org/10.1890/0012-9658\(2006\)87\[2349:EOSRFA\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2006)87[2349:EOSRFA]2.0.CO;2).
- 467 Calenge, C. 2011. Exploratory analysis of the habitat selection by the wildlife in R: the
 468 adehabitatHS Package. *Office national de la chasse et de la faune sauvage, Saint*
 469 *Benoist, 78610*.
- 470 Calenge, C. 2015. Home range estimation in R: the adehabitatHR package. *Office national de*
 471 *la classe et de la faune sauvage: Saint Benoist, Auffargis, France*.
- 472 Colwell, M. A., T. Danufsky, N. W. Fox-Fernandez, J. E. Roth, and J. R. Conklin. 2003.
 473 Variation in shorebird use of diurnal, high-tide roosts: How consistently are roosts
 474 used? *Waterbirds* 26: 484–493. [https://doi.org/10.1675/1524-4695\(2003\)026\[0484:VISUOD\]2.0.CO;2](https://doi.org/10.1675/1524-4695(2003)026[0484:VISUOD]2.0.CO;2).
- 476 Colwell, M. A., and S. L. Landrum. 1993. Nonrandom shorebird distribution and fine-scale
 477 variation in prey abundance. *The Condor* 95: 94–103.
 478 <https://doi.org/10.2307/1369390>.

479 Cramp, S., K. E. L. Simmons, D. C. Brooks, N. J. Collar, E. Dunn, R. Gillmor, P. a. D.
480 Hollom, et al. 1983. *Handbook of the birds of Europe, the Middle East and North*
481 *Africa. The birds of the Western Palearctic: 3. Waders to gulls.*

482 Cunningham, S. J., J. R. Corfield, A. N. Iwaniuk, I. Castro, M. R. Alley, T. R. Birkhead, and
483 S. Parsons. 2013. The Anatomy of the bill Tip of Kiwi and Associated Somatosensory
484 Regions of the Brain: Comparisons with Shorebirds. *PloS One* 8. Public Library of
485 Science: e80036. <https://doi.org/10.1371/journal.pone.0080036>.

486 Delany, S., D. Scott, T. Dodman, and D. Stroud. 2009. *The Wader atlas: An atlas of Wader*
487 *populations in Africa and Western Eurasia.*

488 Demongin, L. 2020. *Guide d'identification des oiseaux en main : Les 250 espèces les plus*
489 *baguées en France: identification, mensurations, variations géographiques, mue, sexe*
490 *et âge.* L. Demongin.

491 Dias, M. P., J. P. Granadeiro, and J. M. Palmeirim. 2009. Searching behaviour of foraging
492 waders: Does feeding success influence their walking? *Animal Behaviour* 77: 1203–
493 1209. <https://doi.org/10.1016/j.anbehav.2009.02.002>.

494 Dias, M. P., F. Peste, J. P. Granadeiro, and J. M. Palmeirim. 2008. Does traditional
495 shellfishing affect foraging by waders? The case of the Tagus estuary (Portugal). *Acta*
496 *Oecologica* 33: 188–196. <https://doi.org/10.1016/j.actao.2007.10.005>.

497 Dodd, S. L., and M. A. Colwell. 1996. Seasonal variation in diurnal and nocturnal
498 distributions of nonbreeding shorebirds at North Humboldt Bay, California. *The*
499 *Condor* 98: 196–207.

500 Dugan, P. J. 1981. The importance of nocturnal foraging in shorebirds: A consequence of
501 increased invertebrate prey activity. In *Feeding and Survival Strategies of Estuarine*
502 *Organisms*, ed. N. V. Jones and W. J. Wolff, 251–260. Boston, MA: Springer US.
503 https://doi.org/10.1007/978-1-4613-3318-0_19.

504 Duijns, S., N. A. Hidayati, and T. Piersma. 2013. Bar-tailed godwits *Limosa l. lapponica* eat
505 polychaete worms wherever they winter in Europe. *Bird Study* 60: 509–517.
506 <https://doi.org/10.1080/00063657.2013.836153>.

507 Duijns, S., J. A. Van Gils, B. Spaans, J. ten Horn, M. Brugge, and T. Piersma. 2014. Sex-
508 specific winter distribution in a sexually dimorphic shorebird is explained by resource
509 partitioning. *Ecology and Evolution* 4: 4009–4018. <https://doi.org/10.1002/ece3.1213>.

510 Dwyer, R. G., S. Bearhop, H. A. Campbell, and D. M. Bryant. 2013. Shedding light on light:
511 benefits of anthropogenic illumination to a nocturnally foraging shorebird. *Journal of*
512 *Animal Ecology* 82: 478–485. <https://doi.org/10.1111/1365-2656.12012>.

513 Esser, W., S. Vöge, and K.-M. Exo. 2008. Day-night activity of intertidal invertebrates and
514 methods to estimate prey accessibility for shorebirds. *Senckenbergiana maritima* 38:
515 115–122. <https://doi.org/10.1007/BF03055286>.

516 Evans, A. 1987. Relative availability of the prey of wading birds by day and by night. *Marine*
517 *Ecology Progress Series* 37: 103–107.

518 Evans, P. R. 1986. Correct measurement of the wing-length of waders. *Wader Study Group*
519 *Bull* 48:11.

520 van Gils, Jan A., I. W. Schenk, O. Bos, and T. Piersma. 2003. Incompletely informed
521 shorebirds that face a digestive constraint maximize net energy gain when exploiting
522 patches. *The American Naturalist* 161: 777–793. <https://doi.org/10.1086/374205>.

523 Goeldner-Gianella, L. 2005. Verger (F.) - *Marais maritimes et estuaires du littoral français.*
524 Paris, Belin, 2005, 335 p. *Norois. Environnement, aménagement, société:* 138–139.

525 Goss-Custard, J. D., A. D. West, M. G. Yates, R. W. G. Caldow, R. A. Stillman, L. Bardsley,
526 J. Castilla, et al. 2007. Intake rates and the functional response in shorebirds
527 (Charadriiformes) eating macro-invertebrates. *Biological Reviews* 81: 501–529.
528 <https://doi.org/10.1111/j.1469-185X.2006.tb00216.x>.

- 529 Granadeiro, J. P., M. P. Dias, R. C. Martins, and J. M. Palmeirim. 2006. Variation in numbers
530 and behaviour of waders during the tidal cycle: Implications for the use of estuarine
531 sediment flats. *Acta Oecologica* 29: 293–300.
532 <https://doi.org/10.1016/j.actao.2005.11.008>.
- 533 Gunnarsson, T. G., J. A. Gill, J. Newton, P. M. Potts, and W. J. Sutherland. 2005. Seasonal
534 matching of habitat quality and fitness in a migratory bird. *Proceedings of the Royal
535 Society B: Biological Sciences* 272: 2319–2323.
536 <https://doi.org/10.1098/rspb.2005.3214>.
- 537 Gunnarsson, T. G., J. A. Gill, P. W. Atkinson, G. Gélinaud, P. M. Potts, R. E. Croger, G. A.
538 Gudmundsson, G. F. Appleton, and W. J. Sutherland. 2006. Population-scale drivers
539 of individual arrival times in migratory birds. *Journal of Animal Ecology* 75: 1119–
540 1127.
- 541 Handel, C. M., and R. E. Gill. 1992. Roosting behavior of premigratory dunlins (*Calidris
542 alpina*). *The Auk* 109: 57–72. <https://doi.org/10.2307/4088266>.
- 543 Hilton, G. M., G. D. Ruxton, and W. Cresswell. 1999. Choice of foraging area with respect to
544 predation risk in redshanks: The effects of weather and predator activity. *Oikos* 87:
545 295–302. <https://doi.org/10.2307/3546744>.
- 546 Johnson, D. H. 1980. The comparison of usage and availability measurements for evaluating
547 resource preference. *Ecology* 61: 65–71. <https://doi.org/10.2307/1937156>.
- 548 Kam, J. van de, B. Ens, T. Piersma, and L. Zwarts. 2004. *Shorebirds: An Illustrated
549 Behavioural Ecology*. Brill.
- 550 Kelsey, M. G., and M. Hassall. 1989. Patch selection by dunlin on a heterogeneous mudflat.
551 *Ornis Scandinavica (Scandinavian Journal of Ornithology)* 20: 250–254.
552 <https://doi.org/10.2307/3676488>.
- 553 Kie, J. G., J. Matthiopoulos, J. Fieberg, R. A. Powell, F. Cagnacci, M. S. Mitchell, J.-M.
554 Gaillard, and P. R. Moorcroft. 2010. The home-range concept: Are traditional
555 estimators still relevant with modern telemetry technology? *Philosophical
556 Transactions of the Royal Society B: Biological Sciences* 365: 2221–2231.
557 <https://doi.org/10.1098/rstb.2010.0093>.
- 558 Kuwae, T. 2007. Diurnal and nocturnal feeding rate in Kentish plovers *Charadrius
559 alexandrinus* on an intertidal flat as recorded by telescopic video systems. *Marine
560 Biology* 151: 663–673.
- 561 Lafferty, K. D. 2001. Birds at a Southern California beach: Seasonality, habitat use and
562 disturbance by human activity. *Biodiversity & Conservation* 10: 1949–1962.
- 563 Last, K. S., and P. J. W. Olive. 2004. Interaction between photoperiod and an endogenous
564 seasonal factor in influencing the diel locomotor activity of the benthic polychaete
565 *Nereis virens* Sars. *The Biological Bulletin* 206: 103–112.
566 <https://doi.org/10.2307/1543541>.
- 567 Laver, P. N., and M. J. Kelly. 2008. A critical review of home range studies. *Journal of
568 Wildlife Management* 72: 290–298. <https://doi.org/10.2193/2005-589>.
- 569 Lourenço, P. M., A. Silva, C. D. Santos, A. C. Miranda, J. P. Granadeiro, and J. M.
570 Palmeirim. 2008. The energetic importance of night foraging for waders wintering in a
571 temperate estuary. *Acta Oecologica* 34: 122–129.
572 <https://doi.org/10.1016/j.actao.2008.04.005>.
- 573 Mallory, M. L., and C. D. Gilbert. 2008. Leg-loop harness design for attaching external
574 transmitters to seabirds. *Marine Ornithology* 36: 183–188.
- 575 Manly, B. F., L. McDonald, D. L. Thomas, T. L. McDonald, and W. P. Erickson. 2002.
576 *Resource selection by animals: Statistical design and analysis for field studies*.
577 Springer Science & Business Media. Springer Science & Business Media.
- 578 Martin, G. 1990. *Birds by Night*. Poyser.

- 579 Martin, G. R. 2012. Through birds' eyes: Insights into avian sensory ecology. *Journal of*
580 *Ornithology* 153: 23–48. <https://doi.org/10.1007/s10336-011-0771-5>.
- 581 McNamara, J., and A. Houston. 1980. The application of statistical decision theory to animal
582 behaviour. *Journal of Theoretical Biology* 85: 673–690. [https://doi.org/10.1016/0022-](https://doi.org/10.1016/0022-5193(80)90265-9)
583 [5193\(80\)90265-9](https://doi.org/10.1016/0022-5193(80)90265-9).
- 584 McNeil, R., O. Díaz Díaz, I. Liñero A., and J. R. Rodríguez S. 1995. Day- and night-time prey
585 availability for waterbirds in a tropical lagoon. *Canadian Journal of Zoology* 73: 869–
586 878. <https://doi.org/10.1139/z95-102>.
- 587 McNeil, R., P. Drapeau, and J. D. Goss-Custard. 1992. The occurrence and adaptive
588 significance of nocturnal habits in waterfowl. *Biological Reviews* 67: 381–419.
589 <https://doi.org/10.1111/j.1469-185X.1992.tb01188.x>.
- 590 McNeil, R., and J. R. Rodríguez. 1996. Nocturnal foraging in shorebirds. *International Wader*
591 *Studies* 8: 114-121.
- 592 Morrison, G., N. C. Davidson, and J. Wilson. 2007. Survival of the fattest: Body stores on
593 migration and survival in red knots *Calidris canutus islandica*. *Journal of Avian*
594 *Biology* 38: 479-487.
- 595 Mouritsen, K. N. 1993. Diurnal and nocturnal prey detection by dunlins *Calidris alpina*. *Bird*
596 *Study* 40: 212–215. <https://doi.org/10.1080/00063659309477185>.
- 597 Mouritsen, K. N. 1994. Day and night feeding in dunlins *Calidris alpina*: Choice of habitat,
598 foraging technique and prey. *Journal of Avian Biology* 25: 55-62.
599 <https://doi.org/10.2307/3677294>.
- 600 Navedo, J. G., and J. A. Masero. 2007. Measuring potential negative effects of traditional
601 harvesting practices on waterbirds: A case study with migrating curlews. *Animal*
602 *Conservation* 10: 88–94. <https://doi.org/10.1111/j.1469-1795.2006.00076.x>.
- 603 Nolet, B. A., and W. M. Mooij. 2002. Search paths of swans foraging on spatially
604 autocorrelated tubers. *Journal of Animal Ecology* 71: 451–462.
605 <https://doi.org/10.1046/j.1365-2656.2002.00610.x>.
- 606 Pienkowski, M. W. 1983. Changes in the foraging pattern of plovers in relation to
607 environmental factors. *Animal Behaviour* 31: 244–264. [https://doi.org/10.1016/S0003-](https://doi.org/10.1016/S0003-3472(83)80195-X)
608 [3472\(83\)80195-X](https://doi.org/10.1016/S0003-3472(83)80195-X).
- 609 Piersma, T., R. E. Gill, P. de Goeij, A. Dekinga, M. L. Shepherd, D. Ruthrauff, and L.
610 Tibbitts. 2006. Shorebird avoidance of nearshore feeding and roosting areas at night
611 correlates with presence of a nocturnal avian predator. *Bulletin-Wader Study Group*
612 109: 73–76.
- 613 Piersma, T., P. de Goeij, and I. Tulp. 1993. An evaluation of intertidal feeding habitats from a
614 shorebird perspective: Towards relevant comparisons between temperate and tropical
615 mudflats. *Netherlands Journal of Sea Research* 31: 503–512.
616 [https://doi.org/10.1016/0077-7579\(93\)90062-W](https://doi.org/10.1016/0077-7579(93)90062-W).
- 617 Pitelka, F. A. 1979. *Shorebirds in marine environments*. Cooper Ornithological Society.
- 618 Prater, A. J., J. H. Marchant, and J. Vourinen. 1977. *Guide to the identification and ageing of*
619 *Holarctic waders*. Tring, Herts. BTO Guide No. 17.
- 620 Puttick, G. M. 1984. Foraging and activity patterns in wintering shorebirds. *Behavior of*
621 *Marine Animals: Current Perspectives in Research*.
- 622 Pyle, P. 2008. *Identification guide to North America birds: Part II*. Slate Creek Press.
- 623 Quaintenne, G., J. A. van Gils, P. Bocher, A. Dekinga, and T. Piersma. 2010. Diet selection in
624 a molluscivore shorebird across Western Europe: Does it show short- or long-term
625 intake rate-maximization? *Journal of Animal Ecology* 79: 53–62.
626 <https://doi.org/10.1111/j.1365-2656.2009.01608.x>.
- 627 Rogers, D. I. 2003. High-tide roost choice by coastal waders. *Bulletin-Wader Study Group*
628 100: 73-79.

- 629 Rogers, D. I., P. F. Battley, T. Piersma, J. A. Van Gils, and K. G. Rogers. 2006. High-tide
630 habitat choice: Insights from modelling roost selection by shorebirds around a tropical
631 bay. *Animal Behaviour* 72: 563–575. <https://doi.org/10.1016/j.anbehav.2005.10.029>.
- 632 Rohweder, D. A. 2001. Nocturnal roost use by migratory waders in the Richmond River
633 Estuary, northern New South Wales, Australia. *Stilt*: 23–28.
- 634 Rojas, L. M., R. McNeil, T. Cabana, and P. Lachapelle. 1999. Diurnal and nocturnal visual
635 capabilities in shorebirds as a function of their feeding strategies. *Brain, Behavior and*
636 *Evolution* 53: 29–43. <https://doi.org/10.1159/000006580>.
- 637 Rompré, G., and R. McNeil. 1996. Variability in day and night feeding habitat use in the
638 Willet *Catoptrophorus semipalmatus* during the non-breeding season in northeastern
639 Venezuela. *Wader Study Group Bull*: 82–87.
- 640 Rosa, S., A. L. Encarnação, J. P. Granadeiro, and J. M. Palmeirim. 2006. High water roost
641 selection by waders: Maximizing feeding opportunities or avoiding predation? *Ibis*
642 148: 88–97. <https://doi.org/10.1111/j.1474-919X.2006.00497.x>.
- 643 Santos, C. D., A. C. Miranda, J. P. Granadeiro, P. M. Lourenço, S. Saraiva, and J. M.
644 Palmeirim. 2010. Effects of artificial illumination on the nocturnal foraging of waders.
645 *Acta Oecologica* 36: 166–172. <https://doi.org/10.1016/j.actao.2009.11.008>.
- 646 Schmaltz, L., G. Quaintenne, L. Couzi, and J. Dupuy. 2019. *Comptage des oiseaux d'eau à la*
647 *mi-janvier en France. Résultats 2019 du comptage Wetlands International*. LPO.
- 648 Schuler, K. L., G. M. Schroeder, J. A. Jenks, and J. G. Kie. 2014. Ad hoc smoothing
649 parameter performance in kernel estimates of GPS-derived home ranges. *Wildlife*
650 *Biology* 20: 259–266. <https://doi.org/10.2981/wlb.12117>.
- 651 Sitters, H. P., P. M. Gonzalez, T. Piersma, A. J. Baker, D. J. Price, H. P. Sitters, P. M.
652 González, A. J. Baker, and D. J. Price. 2001. Day and night feeding habitat of red
653 knots in Patagonia: Profitability versus safety? *Journal of Field Ornithology* 72: 86–
654 95. <https://doi.org/10.1648/0273-8570-72.1.86>.
- 655 Sitters, H. P. 2000. *The role of night-feeding in shorebirds in an estuarine environment with*
656 *specific reference to mussel-feeding Oystercatchers*. University of Oxford.
- 657 Smith, K. W., J. M. Reed, and B. E. Trevis. 1999. Nocturnal and diurnal activity patterns and
658 roosting sites of green sandpipers *Tringa ochropus* wintering in southern England.
659 *Ringing & Migration* 19: 315–322. <https://doi.org/10.1080/03078698.1999.9674200>.
- 660 Stephens, D., and J. R. Krebs. 1986. *Foraging theory*. Vol. 1. Princeton University Press.
661 <https://doi.org/10.2307/1381654>.
- 662 Thomas, K., R. G. Kvitek, and C. Bretz. 2003. Effects of human activity on the foraging
663 behavior of sanderlings *Calidris alba*. *Biological Conservation* 109: 67–71.
664 [https://doi.org/10.1016/S0006-3207\(02\)00137-4](https://doi.org/10.1016/S0006-3207(02)00137-4).
- 665 Triplet, P., S. Le Dréan Quéneec'hdu, and R. Mahéo. 2010. La Barge rousse *Limosa lapponica*
666 hivernant en France (1977-2009). *Alauda* 78: 207–216.
- 667 Turpie, J. K., and P. a. R. Hockey. 1993. Comparative diurnal and nocturnal foraging
668 behaviour and energy intake of premigratory grey plovers *Pluvialis squatarola* and
669 whimbrels *Numenius phaeopus* in South Africa. *Ibis* 135: 156–165.
670 <https://doi.org/10.1111/j.1474-919X.1993.tb02827.x>.
- 671 Van Winkle, W. 1975. Comparison of several probabilistic home-range models. *The Journal*
672 *of Wildlife Management*: 118–123. <https://doi.org/10.2307/3800474>.
- 673 VanDusen, B. M., S. R. Fegley, and C. H. Peterson. 2012. Prey distribution, physical habitat
674 features, and guild traits interact to produce contrasting shorebird assemblages among
675 foraging patches. *PloS One* 7. <https://doi.org/10.1371/journal.pone.0052694>.
- 676 West, A. D., J. D. Goss-Custard, S. E. A. Le V. dit Durell, and R. A. Stillman. 2005.
677 Maintaining estuary quality for shorebirds: Towards simple guidelines. *Biological*
678 *Conservation* 123: 211–224. <https://doi.org/10.1016/j.biocon.2004.11.010>.

679 Worton, B. J. 1989. Kernel methods for estimating the utilization distribution in home-range
680 studies. *Ecology* 70: 164–168. <https://doi.org/10.2307/1938423>.
681 Zharikov, Y, and G. Skilleter. 2003. Depletion of benthic invertebrates by bar-tailed godwits
682 *Limosa lapponica* in a subtropical estuary. *Marine Ecology Progress Series* 254: 151–
683 162. <https://doi.org/10.3354/meps254151>.
684

685 **Table 1** Size of diurnal and nocturnal feeding areas (FCA: feeding core areas 50%, FHR: feeding home ranges 95%) and their overlap rate

Logger	Mass (mm)	Bill length (mm)	Tracking period	Number of days	DAY			NIGHT			Overlap dFCA/nFCA (%)	Overlap dFHR/nFHR (%)
					GPS fixes	dFCA (ha)	dFHR (ha)	GPS fixes	nFCA (ha)	nFHR (ha)		
BTG01	301	91.5	20 Nov. - 17 Mar.	118	557	68.3	321.8	474	34.5	226.3	17.2	39.4
BTG02	335	105	31 Dec. - 31 Mar.	90	489	31.2	212.5	418	8.7	72.9	5.9	12.2
BTG03	325	99	31 Dec. - 15 Mar.	74	138	36.3	148.7	125	24.9	89.6	17.2	32.4
BTG04	332	101	11 Nov. - 26 Mar.	136	1125	53.1	276.1	1046	29.6	168.4	5.4	43.1
BTG05	326	95	11 Nov. - 29 Mar.	139	951	22.5	125.0	872	7.9	36.0	33.6	28.4
BTG06	316	101	01 Nov. - 17 Mar.	136	1687	58.4	262.4	1809	30.8	171.8	28.3	53.4
686												
687												
688												

Figure captions

Fig. 1 Map of the Pertuis Charentais (Central French Atlantic coast) and localization of the study areas on Ré Island. Dark gray corresponds to the mainland and light gray to the intertidal area

Fig. 2 Diurnal and nocturnal feeding home ranges of six bar-tailed godwits during the nonbreeding period on Ré Island. Feeding home ranges were calculated as 50% (core areas) and 95% (home ranges) kernel density contours

Fig. 3 Mean proportion of time spent by six Bar-tailed godwits on mudflats, i.e. foraging, by day and by night during the tide cycle. Significant differences between Day and Night, for each slot time around the low tide, are illustrated by the symbols ‘*’ when $p < 0.05$ and ‘***’ when $p < 0.001$.

Fig. 4 Day and night distribution of distances between two successive locations (30 min.)

Fig. 5 Habitat typology at (a) Ré Island, (b) “La Loge,” and (c) “Fier d’Ars”

Fig. 6 Day and night proportion of habitat types on feeding core areas (50% kernel density contour) of each bar-tailed godwit

Fig. 7 Results of the selection ratio analysis highlighting habitat selection by six bar-tailed godwits on 10 intertidal habitat types by (a) daytime and (b) nighttime

Fig. 8 Diurnal and nocturnal roosting home ranges of six bar-tailed godwits during the nonbreeding period on Ré Island. Roosting home ranges were calculated as 50% (core areas) and 95% (home ranges) kernel density contours