



HAL
open science

K-NARAYANA SEQUENCE SELF-SIMILARITY

Engin Özkan, Bahar Kuloğlu, James F Peters

► **To cite this version:**

Engin Özkan, Bahar Kuloğlu, James F Peters. K-NARAYANA SEQUENCE SELF-SIMILARITY. 2021. hal-03242990

HAL Id: hal-03242990

<https://hal.science/hal-03242990>

Preprint submitted on 31 May 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

K-NARAYANA SEQUENCE SELF-SIMILARITY

ENGIN ÖZKAN, BAHAR KULOĞLU, AND JAMES F. PETERS

ABSTRACT. This paper introduces self-similarity inherent in planar Milich centered flip graphs derived from the Narayana sequence. We show that self-similarity found in a Narayana sequence yields a connected spanning subgraph with a centered flip. This paper has several main results (1) Every Narayana sequence constructs a flip graph, (2) Every Narayana sequence is self-similar and (3) Every Pascal 3-triangle has a free group presentation.

CONTENTS

1. Introduction	1
2. Preliminaries	2
3. The k -Narayana numbers and properties	5
4. The Pascal 3-triangle	7
5. Free Abelian Group Presentation of a Pascal 3-Triangle	7
6. Views and Properties of the Pascal 3-triangle	10
6.1. Some Properties of the Pascal 3-triangle Diagonals	11
6.2. Properties of the rows the Pascal 3-triangle	12
7. Expression of k^n as a k -Narayana number	12
8. Catalan transform of the k -Narayana sequence	13
9. Hankel Transform	14
10. Conclusion	15
Appendix A. Cell Complexes	15
Appendix B. Shape Complexes	16
Appendix C. Finite 1-cycles and their Cyclic Group Presentation	17
References	19

1. INTRODUCTION

In this paper, both population growth as well as population decline are represented by the bipolar form of the Narayana sequence, namely,


$$\dots, -13, -9, -6, -4, -3, -2, -1, -1, -1, 0, 1, 1, 1, 2, 3, 4, 6, 9, 13, \dots$$

There is an inherent self-similarity in Narayana sequences (see, *e.g.*, the partial plot of three different forms of this sequence shown in Fig. 1.

For the k^{th} member N_k of a population, the Narayana growth model is defined by the linear recurrence sequence $N_k = N_{k-1} + N_{k-3}$, $k \geq 3$, starting with $N_0, N_1, N_2, N_3 =$

2010 *Mathematics Subject Classification.* 11B39 (Fibonacci numbers); 05C10 (planar graphs).

Key words and phrases. Flip Graphs, k -Narayana sequence, Pascal 3-triangle, Planar Graph, Self-Similarity.

FIGURE 1. Narayana population growth & decline plot over $[-13, 13]$

$0, 1, 1, 1$ representing 3 generations that starts growing at $N_3 = N_2 + N_0 = 1 + 1 = 2$. This population growth model was introduced 1356 by Nārāyana Paṇḍita [14]. Similarly, this sequence has a population decline model defined by $N_k = N_{k-1} + N_{k-3}, k \leq -3$, starting with $N_0, N_{-1}, N_{-2}, N_{-3} = 0, -1, -1, -1$ representing 3 declining generations that starts delining at $N_{-4} = N_{-1} + N_{-3} = -1 + (-1) = -2$.

The self-similarity in Fig. 1 is a result of considering the generalized Narayana sequence [18] defined by

$$\begin{aligned} N_{k,n} &= kN_{k,n-1} + N_{k,n-3}, n \geq 0, k \geq 3 \\ &= 0, 1, 1, 1, 2, 3, 4, 6, 9, 13, \end{aligned}$$

with the initial values $N_{k,0} = 0, N_{k,1} = 1$ and $N_{k,2} = k$, yielding the sequence $0, 1, k, k^2, k^3 + 1, k^4 + 2k, k^5 + 3k^2$. In the study of Narayana sequence self-similarity, we consider $\pm N_{k,n}, k \geq 3$, representing population growth as well as population decline.

It is well-known graph self-similarity such that shown for $\pm N_{k,n}$ in Fig. 1 leads to generalizations of self-similarity [5, §4.4, p. 120]. From the edges in graph similarity, it is possible to construct examples of border (outline) fractals [5, §6.1]. The self-similarity of the Narayana sequence $\pm N_{k,n}$ leads to the following result.

Theorem Every Narayana sequence is self-similar.

A flip graph is a form of undirected, simple graph, which we consider next in a closer look at Narayana sequences and their self-similarities.

2. PRELIMINARIES

In this section the construction of self-similar graphs via bipolar Narayana sequences. A **bipolar Narayana sequence** $\pm N_{k,n}$ represents two extremes, namely, the growth of a population at one extreme and the decline of a population at the other extreme, representable in a planar self-similar graph.

This paper also considers Pascal-like triangles[2] with corresponding finite cyclic group presentations derived from k -Narayana sequences.

Definition 1. [11, 10] A **planar flip graph** is a 2D graph whose vertexes represent combinatorial objects of interest and an edge between any two objects that differ only by a small change operation called a flip. ■

Example 1. A flip graph representing the beginning of a population and its decline is shown in Fig. 2.1. This flip graph exhibits the simplest form self-similarity centered at the origin, i.e., $-2, -1, -1, 0, 1, 1, 1, 2$. The altitude 2 to the right of the




FIGURE 2. Narayana flip graphs

origin represents initial of a population. The altitude 2 to the left of the origin represents initial decline of the same population. ■

Example 2. A flip graph representing the beginning of a population and its decline is shown in Fig. 2.2. This flip graph exhibits 4th generation self-similarity centered at the origin, i.e., $-6,-4,-3,-2,-1,-1,0,1,1,1,2,3,4,6$. The altitudes 2,3,4,6 to the right of the origin represent initial population growth. The altitudes 2,3,4,6 to the left of the origin represents initial decline of the same population. ■

Remark 1. Positive altitudes to the left of the origin in Fig. 2.1 and Fig. 2.2 represents count of those generations in a population that failed to appear. ■

Lemma 1. A bipolar Narayana sequence constructs a flip graph.

Proof. Let $\pm N_{k,n}$ be a bipolar Narayana sequence. We start with $k = \pm 3$ with three vertexes, namely, vertexes at $\pm 1, \pm 3$ on the horizontal axis and a vertexes at $(3,2), (-3,2)$, which we triangulate. The edge from $(1,0)$ to $(3,2)$ represents a small change (a flip) in population growth, which contrasts another flip along the edge

from $(-1,0)$ to $(-3,2)$. Continuing this triangulation on additional triple vertexes derived from a Narayana sequence constructs a flip graph. \square

Theorem 1. *Every bipolar Narayana sequence is self-similar.*

Proof. From what has been observed about the relationship between self-similarity and flip graphs and from Lemma 1, the result follows. \square

Remark 2. *From Theorem 1, we obtain a natural basis for a free group presentation of the flip graph for a Narayana sequence. That is, a flip graph viewed as a pair of cycles along the edges of the triangles in the graph make it possible to travel in either the positive or negative direction. In effect, every $+ve$ traversal a to the right in a Narayana flip graph has a corresponding $-ve$ traversal $-a$ to the left, giving us inverse elements, which sum to zero, $a - a = 0$. For more about this, see Section 5 and Appendix C. \blacksquare*

The k -Narayana numbers are given by Binet's formula.

$$(1) \quad N_{k,n} = \frac{\alpha_k^{n+1}}{(\alpha_k - \beta_k)(\alpha_k - \gamma_k)} + \frac{\beta_k^{n+1}}{(\beta_k - \alpha_k)(\beta_k - \gamma_k)} + \frac{\gamma_k^{n+1}}{(\gamma_k - \beta_k)(\gamma_k - \alpha_k)}$$

where

$$\alpha_k = \frac{1}{3}(k + k^2 \sqrt[3]{\frac{2}{27 + 2k^3 + 3\sqrt{81 + 12k^3}}} + \sqrt[3]{\frac{27 + 2k^3 + 3\sqrt{81 + 12k^3}}{2}}),$$

$$\beta_k = \frac{1}{3}(k - wk^2 \sqrt[3]{\frac{2}{27 + 2k^3 + 3\sqrt{81 + 12k^3}}} + w^2 \sqrt[3]{\frac{27 + 2k^3 + 3\sqrt{81 + 12k^3}}{2}}),$$

$$\gamma_k = \frac{1}{3}(k + w^2k^2 \sqrt[3]{\frac{2}{27 + 2k^3 + 3\sqrt{81 + 12k^3}}} + w \sqrt[3]{\frac{27 + 2k^3 + 3\sqrt{81 + 12k^3}}{2}}).$$

$w = \frac{1+i\sqrt{3}}{2}$ is the primitive cube root of unity [18]. Characteristic equation of these numbers is

$$(2) \quad x^3 + kx^2 - 1 = 0$$

For any integer number $k, k \neq 0$, we have

$$Q_k = \begin{pmatrix} k & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$$

and for $n \geq 3$,

$$(Q_k)^n = \begin{pmatrix} N_{k,n+1} & N_{k,n-1} & N_{k,n} \\ N_{k,n} & N_{k,n-2} & N_{k,n-1} \\ N_{k,n-1} & N_{k,n-3} & N_{k,n-2} \end{pmatrix}$$

Then Q_k is a generating matrix of the k -Narayana sequence [18].

3. THE k -NARAYANA NUMBERS AND PROPERTIES

Theorem 2. [Cassini's Identity]

$$N_{k,n-1}N_{k,n+1} - N_{k,n}^2 = -N_{k,-n-1}.$$

Proof. By using the Binet's formula, we get

$$\begin{aligned} & \left(\frac{\alpha_k^n}{(\alpha_k - \beta_k)(\alpha_k - \gamma_k)} + \frac{\beta_k^n}{(\beta_k - \alpha_k)(\beta_k - \gamma_k)} + \frac{\gamma_k^n}{(\gamma_k - \beta_k)(\gamma_k - \alpha_k)} \right) \\ & \left(\frac{\alpha_k^{n+2}}{(\alpha_k - \beta_k)(\alpha_k - \gamma_k)} + \frac{\beta_k^{n+2}}{(\beta_k - \alpha_k)(\beta_k - \gamma_k)} + \frac{\gamma_k^{n+2}}{(\gamma_k - \beta_k)(\gamma_k - \alpha_k)} \right) \\ & - \left(\frac{\alpha_k^{n+1}}{(\alpha_k - \beta_k)(\alpha_k - \gamma_k)} + \frac{\beta_k^{n+1}}{(\beta_k - \alpha_k)(\beta_k - \gamma_k)} + \frac{\gamma_k^{n+1}}{(\gamma_k - \beta_k)(\gamma_k - \alpha_k)} \right)^2 \\ & = \frac{\alpha_k^n \beta_k^{n+2} - 2\alpha_k^{n+1} \beta_k^{n+1} + \alpha_k^{n+2} \beta_k^n}{(\alpha_k - \beta_k)(\alpha_k - \gamma_k)(\beta_k - \alpha_k)(\beta_k - \gamma_k)} \\ & + \frac{\alpha_k^n \gamma_k^{n+2} - 2\alpha_k^{n+2} \gamma_k^n + \alpha_k^{n+2} \gamma_k^n}{(\alpha_k - \beta_k)(\alpha_k - \gamma_k)(\gamma_k - \alpha_k)(\gamma_k - \beta_k)} \\ & + \frac{\beta_k^n \gamma_k^{n+2} - 2\beta_k^{n+1} \gamma_k^{n+1} + \beta_k^{n+2} \gamma_k^n}{(\beta_k - \alpha_k)(\beta_k - \gamma_k)(\gamma_k - \alpha_k)(\gamma_k - \beta_k)} \\ & = \frac{\alpha_k^n \beta_k^n (\alpha_k - \beta_k)^2}{(\alpha_k - \beta_k)(\alpha_k - \gamma_k)(\beta_k - \alpha_k)(\beta_k - \gamma_k)} \\ & + \frac{\alpha_k^n \gamma_k^n (\alpha_k - \gamma_k)^2}{(\alpha_k - \beta_k)(\alpha_k - \gamma_k)(\gamma_k - \alpha_k)(\gamma_k - \beta_k)} \\ & + \frac{\beta_k^n \gamma_k^n (\beta_k - \gamma_k)^2}{(\gamma_k - \alpha_k)(\gamma_k - \beta_k)(\beta_k - \alpha_k)(\beta_k - \gamma_k)} \\ & = - \left(\frac{\alpha_k^n \beta_k^n}{(\alpha_k - \gamma_k)(\beta_k - \gamma_k)} + \frac{\alpha_k^n \gamma_k^n}{(\alpha_k - \beta_k)(\gamma_k - \beta_k)} + \frac{\beta_k^n \gamma_k^n}{(\beta_k - \alpha_k)(\gamma_k - \alpha_k)} \right) \\ & = - \left(\frac{\gamma_k^{-n}}{(\alpha_k - \gamma_k)(\beta_k - \gamma_k)} + \frac{\beta_k^{-n}}{(\alpha_k - \beta_k)(\gamma_k - \beta_k)} + \frac{\alpha_k^{-n}}{(\beta_k - \alpha_k)(\gamma_k - \alpha_k)} \right) \\ & = -N_{k,-n-1} \end{aligned}$$

□

Remark 3. A 3D flip graph exhibiting the self-similarity of the Narayana sequence using Binet's formula is shown in Fig. 3. ■

Theorem 3.

$$\lim_{n \rightarrow \infty} \frac{N_{k,n}}{N_{k,n-1}} = \alpha_k$$

Proof. Symmetric with the proof of Theorem 2. □

Theorem 4. For any integer $n \geq 2$, we have

$$\begin{aligned} \alpha_k^{n+3} &= k\alpha_k^{n+2} + 1. \\ \beta_k^{n+3} &= k\beta_k^{n+2} + 1. \\ \gamma_k^{n+3} &= k\gamma_k^{n+2} + 1. \end{aligned}$$

where $\alpha_k, \beta_k, \gamma_k$ are the roots of the characteristic equation 2.




FIGURE 3. 3D flip graph of Narayana population growth & decline

Proof. Since $\alpha_k, \beta_k, \gamma_k$ are the roots of the characteristic equation 2 then

$$\begin{aligned}\alpha_k^{n+3} &= k\alpha_k^{n+2} + 1 \\ \beta_k^{n+3} &= k\beta_k^{n+2} + 1 \\ \gamma_k^{n+3} &= k\gamma_k^{n+2} + 1\end{aligned}$$

multiplying both sides of these equations by α_k^n, β_k^n and γ_k^n respectively, we obtain the desired result. \square

Theorem 1. For any integer $n \geq 0$, we write

$$\sum_{i=0}^n k^i N_{k,2i} = 0.$$

Proof. Symmetric with the proof of Theorem 2. \square

Theorem 5. For $n \geq 3$, we have

$$\begin{vmatrix} N_{k,n+1} & N_{k,n-1} & N_{k,n} \\ N_{k,n} & N_{k,n-2} & N_{k,n-1} \\ N_{k,n-1} & N_{k,n-3} & N_{k,n-2} \end{vmatrix} = 1$$

Proof. If we are making some algebraic operation:

$$\begin{aligned}N_{k,n+1}^3 + N_{k,n}^3 + N_{k,n-1}^3 - 2N_{k,n+1}^2 N_{k,n} + N_{k,n}^2 N_{k,n+1} + N_{k,n+1} N_{k,n-1}^2 \\ + N_{k,n}^2 N_{k,n-1} - 3N_{k,n-1} N_{k,n+1} N_{k,n} = 1\end{aligned}$$

We get the equation.

For example, $n = 3$, we have

$$\begin{vmatrix} N_{k,4} & N_{k,2} & N_{k,3} \\ N_{k,3} & N_{k,1} & N_{k,2} \\ N_{k,2} & N_{k,0} & N_{k,1} \end{vmatrix} = \begin{vmatrix} k^3 + 1 & k & k^2 \\ k^2 & 1 & k \\ k & 0 & 1 \end{vmatrix} = 1$$

□

Similarly, we obtain the following result.

Theorem 6 (Sum of first terms). . *Let $S_{k,n}$ be the sum of the first $(n + 1)$ terms of the k -Narayana sequence, that is*

$$S_{k,n} = \sum_{i=0}^n N_{k,i}.$$

Then

$$S_{k,n} = \frac{N_{k,1} + N_{k,n} + N_{k,n+1} + N_{k,n+2}}{k} - N_{k,n+1}.$$

4. THE PASCAL 3-TRIANGLE

We deduce the value of any k -Narayana number by simple substitution on the corresponding $N_{k,n}$. The sequence $N_{1,n}$ is the classical Narayana sequence. By the definition of the k -Narayana numbers [18], the first of them are:

$$\begin{aligned} N_{k,1} &= 1 \\ N_{k,2} &= k \\ N_{k,3} &= k^2 \\ N_{k,4} &= k^3 + 1 \\ N_{k,5} &= k^4 + 2k \\ N_{k,6} &= k^5 + 3k^2 \\ N_{k,7} &= k^6 + 4k^3 + 1 \\ N_{k,8} &= k^7 + 5k^4 + 3k \\ N_{k,9} &= k^8 + 6k^5 + 6k^2 \\ N_{k,10} &= k^9 + 7k^6 + 10k^3 + 1 \end{aligned}$$


It is worth to be noted that the obtained coefficients can be written in a triangular position, in this way that every side of the triangle is triple, and for this reason this triangle has been called Pascal 3-triangle. See table 1.

5. FREE ABELIAN GROUP PRESENTATION OF A PASCAL 3-TRIANGLE

This section introduces free finitely-generated (fg) Abelian group presentations that correspond to Pascal 3-triangles in a planar CW space. Briefly, a planar CW space is a collection of cells in a finite, bounded region of the Euclidean plane. A *cell* in the Euclidean plane is either a 0-cell (vertex K_0) or 1-cell (edge K_1) or 2-cell (filled triangle K_2). A **cell complex** is a collection cells attached to each other by 0- or 1-cells or by having one or more common cells. A nonvoid collection of cell complexes K has a *Closure finite Weak* (CW) topology, provided K is Hausdorff (every pair of distinct cells is contained in disjoint neighbourhoods [12, §5.1, p. 94]) and the collection of cell complexes in K satisfy the Whitehead [24, pp. 315-317] conditions, namely, containment (the closure of each cell complex is in K) and intersection (the nonempty intersection of cell complexes is in K) (for the details, see App. A).

TABLE 1. The Pascal 3-triangle

				1				
				1				
				1				
			1		1			
			1		2			
			1		3			
		1		4		1		
		1		5		3		
		1		6		6		
	1		7		10		1	
	1		8		15		4	
	1		9		21		10	
...	

FIGURE 4. Pascal 3-triangle Nested Cycle Shapes $\text{sh}E, \text{sh}E'$

Pascal 3-triangles like those in Table 1 form collections of nested cycles.

Example 3. A pair of Pascal 3-triangle cycles are shown in Fig. 4, namely,
 1° Pascal 3-triangle in Fig. 4.1 with 3 nested cycles: $1, 1; 1, 2; 1, 3$. The leftmost single vertex labelled 1 in Fig. 4.1 is a cycle that spins on itself (this single vertex represents traversal back to 1 for each of the three cycles shown). The Pascal 3-triangle $1; 1, 1$ in Fig. 4.1 forms a triangle cycle. The Pascal 3-triangle $1; 1, 1; 1, 2; 1, 3$ in Fig. 4.1 forms a collection of nested cycles.
 2° Next, the Pascal 3-triangle in Fig. 4.1 is attached to $1, 4, 1; 1, 5, 3'$ to form an expanded collection of nested cycles shown in Fig. 4.2. ■

Notice that the pair of collections of nested cycles derived from Pascal 3-triangles form a pair of distinct shapes, namely, shape $\text{sh}E$ in Fig. 4.1 and shape $\text{sh}E'$ in Fig. 4.2. The boundary of $\text{sh}E$ is denoted by $\text{bdy}(\text{sh}E)$ (represented by its vertexes).

Each of these shapes has a distinct boundary, which is a source of generators of a free fg Abelian group presentation derived from a Pascal 3-triangle. For the details concerning CW shape complexes, see App. B.

Example 4. *The pair of Pascal 3-triangle nested cycles form planar shapes as shown in Fig. 4, namely,*

1° *Shape shE in Fig. 4.1 with*

Boundary in Fig. 4.1: $bdy(shE) = \{1, 1', 2, 3\}$, limited to the upper boundary vertexes of shE.

2° *Shape shE' in Fig. 4.2 with*

Boundary in Fig. 4.2: $bdy(shE') = \{1, 1', 2, 3, 4, 5, 3'\}$, limited to the upper boundary vertexes of shE'. ■




FIGURE 5. Pascal 3-triangle Cycles cycE1, cycE'2

Theorem 7. *Every Pascal 3-triangle has a free group presentation.*

Proof. From Theorem 8 in App. C, every shape boundary is a 1-cycle CW space K (see, e.g., Fig. 4.1) and in the CW space K' (see, e.g., Fig. 4.2) has finitely-generated cyclic group presentation. □

Example 5. *Each of the collections of nested cycles in Fig. 5 has an Abelian group presentation, e.g.,*

1° *Cycle cycE1 in Fig. 5.1 is represented by $G(\langle 1' \rangle, +)$ with 1 as the generator, i.e., every vertex in the cycle can be reached from vertex 1'.*

2° *Cycle cycE'2 in Fig. 5.2 is represented by $G(\langle 2 \rangle, +)$ with 2 as the generator, i.e., every vertex in the cycle can be reached from vertex 2. ■*

From Theorem 9 in App. C, a Pascal 3-triangle leads to a collection of nested cycles in the CW space in K (Fig. 4.1) and in the CW space in K' (Fig. 4.2), each of which as a free finitely-generated (fg) Abelian group presentation.

Example 6. *Each of the collections of nested cycles in Fig. 4 has a free fg Abelian group presentation, e.g.,*

- 1° The collection of nested cycles in shE in Fig. 4.1 is represented by $G(\langle 1', 2, 3 \rangle, +)$ with vertexes $1', 2, 3$ as the generators, i.e., every vertex in the cycle can be reached from vertexes $1', 2, 3$.
- 2° The collection of nested cycles in shE' in Fig. 4.2 is represented by $G(\langle 1', 2, 3, 4, 5, 3' \rangle, +)$ with vertexes $1', 2, 3, 4, 5, 3'$ as the generators, i.e., every vertex in the cycle can be reached from vertexes $1', 2, 3, 4, 5, 3'$. ■

6. VIEWS AND PROPERTIES OF THE PASCAL 3-TRIANGLE

Let us examine two points on x and y lines in orthogonal coordinate system. One can see clearly, a way between an x -point and a y -point is not inverting way in the first sestet from x to y by horizontal and vertical unit section. For instance, there are four way from the point $x = (3, 0)$ to the point $y = (0, 1)$:

$$\begin{aligned} & \{(3, 0)(2, 0)(1, 0)(0, 0)(0, 1)\} \\ & \{(3, 0)(2, 0)(1, 0)(1, 1)(0, 1)\} \\ & \{(3, 0)(2, 0)(2, 1)(1, 1)(0, 1)\} \\ & \{(3, 0)(3, 1)(2, 1)(1, 1)(0, 1)\} \end{aligned}$$

The number of ways between an x -point and a y -point can be seen on diagonals in Pascal 3-triangle. In table 2, each element is obtained the sum of the elements on previous row from the left to element above.

TABLE 2. Number of paths between x -points and a y -points

X	Y							
	0	1	2	3	4	5	6	7
0	1	1	1	1	1	1	1	1
1	1	2	3	4	5	6	7	8
2	1	3	6	10	15	21	28	36
3	1	4	10	20	35	56	84	120
4	1	5	15	35	70	126	210	330
5	1	6	21	56	126	252	462	792
6	1	7	28	84	210	462	924	1716
7	1	8	36	120	330	792	1716	3432
8	1	9	45	165	495	1287	3003	6435
9	1	10	55	220	715	2002	5005	11440

For example, number 21 in the third row is the conclusion of summing up the elements of the second row: $1 + 2 + 3 + 4 + 5 + 6 = 21$.

One of the properties of this table is each entrance in the diagonal beginning $\{i, j, \dots\}$ reports the number of terms in the expansion of $(a_1 + a_2 + a_3 + \dots + a_{n+3-j})^n$ for $n \geq j - 2$. For instance, diagonal $\{1, 3, 10, 35, 126, \dots\}$ means the number of terms in the expansion of $(a_1 + a_2 + a_3 + \dots + a_{n-1})^n$, for $n \geq 2$, which is $\binom{2n-2}{n-2}$. Obviously, one can be obtained the sequence of Catalan's numbers by dividing each term of this diagonal by 1, 2, 3, 4, ..., respectively.

The i th row of antidiagonals in Table 2 is equivalent to the i th row of the classical Pascal triangle. The term $a_{i,j}$ of Table 2 satisfies $a_{i,j} = a_{i-1,j} + a_{i,j-1}$.

The determinants of matrices generated by the term $a_{1,1}$ are equal to 1. Also, the square matrices obtained from the term $a_{1,1}$ are symmetric in accordance with the diagonal $\{1, 2, 6, 20, 70, \dots\}$.

As well as, if in 2 each row is moved two places to the right according as the prior row, table 3 existed. Take into account each column of Table 3 is equal to

TABLE 3. Rectangular Pascal 3-triangle

1	1	1	1	1	1	1	1	1	1	1	1
		1	2	3	4	5	6	7	8	9	10
				1	3	6	10	15	21	28	36
						1	4	10	20	35	56
								1	5	15	35
										1	6

the corresponding diagonal of the Pascal 3-triangle which is also known as the "deformed" Pascal triangle [21].

6.1. Some Properties of the Pascal 3-triangle Diagonals. It will be called here triple diagonal to each of the different lines of triples of contiguous numbers on the Pascal 3-triangle as shown in Table 1, from right to left and the from first row to bottom. For instance, the second triple diagonal is $\{1 - 3 - 6, 10 - 15 - 21, \dots\}$ these lines are also called 'anti-diagonals'.

It is clear that the n th diagonal terms of the Pascal 3-triangle are definitely the coefficients of the Taylor expansion of function $g_n(x) = \frac{1}{(1-x)^n}$ centered at the origin.

For a small value $x = 10^{-r}$ ($r \in \mathbf{N}$), when the function $g_n(x)$ written as a decimal number has its integer part equal to 1, and its decimal number can be seen as r-uplas showing the first terms of the n th antidiagonal of the Pascal 3-triangle. For instance,

$$g_5(10^{-3}) = \frac{1}{(1 - 10^{-3})^5} = 1.005015035070 \dots \rightarrow \{1, 5, 15, 35, 70, \dots\}$$

which are the fifth anti-diagonal terms of the Pascal 3-triangle. Some characteristic properties of the Pascal 3-triangle are as follow:

1. The second diagonal is positive integer numbers. The third and fourth diagonal equal to the triangular and tetrahedral numbers, respectively. In general, i th diagonal equal to the $(i - 1)$ -dimensional simplex number [7].
2. Each element is obtained the sum of the preceding element at same row and the previous element in the preceding diagonal.
3. The sum of the consecutive terms in the third diagonal is a perfect square.
4. By subtracting the odd number sequence from the triangular number sequence, the triangular number sequence is obtained again.

$$\begin{aligned}
CN_{k,4} &= \sum_{i=1}^4 \frac{i}{8-i} \binom{8-i}{4-i} N_{k,i} = k^3 + 3k^2 + 5k + 6 \\
CN_{k,5} &= \sum_{i=1}^5 \frac{i}{10-i} \binom{10-i}{5-i} N_{k,i} = k^4 + 4k^3 + 9k^2 + 14k + 18 \\
CN_{k,6} &= \sum_{i=1}^6 \frac{i}{12-i} \binom{12-i}{6-i} N_{k,i} = k^5 + 5k^4 + 14k^3 + 31k^2 + 52k + 56
\end{aligned}$$

We can show $\{N_{k,n}\}$ as the $n \times 1$ matrix N_k and the product of the lower triangular matrix C as

$$\begin{aligned}
\begin{pmatrix} CN_{k,1} \\ CN_{k,2} \\ CN_{k,3} \\ CN_{k,4} \\ \dots \end{pmatrix} &= \begin{pmatrix} 1 & & & & \\ 1 & 1 & & & \\ 2 & 2 & 1 & & \\ 5 & 5 & 5 & 3 & \\ \dots & \dots & \dots & \dots & \dots \end{pmatrix} \begin{pmatrix} N_{k,1} \\ N_{k,2} \\ N_{k,3} \\ N_{k,4} \\ \dots \end{pmatrix} = \begin{pmatrix} 1 \\ 1+k \\ k^2+2k+2 \\ k^3+3k^2+5k+6 \\ \dots \end{pmatrix} \\
&= \begin{pmatrix} 1 & & & & \\ 1 & 1 & & & \\ 2 & 2 & 1 & & \\ 5 & 5 & 5 & 3 & \\ \dots & \dots & \dots & \dots & \dots \end{pmatrix} \begin{pmatrix} 1 \\ k \\ k^2 \\ k^3+1 \\ \dots \end{pmatrix}
\end{aligned}$$

where

$$C_{i,j} = \sum_{r=j-1}^{i-1} C_{i-1,r}.$$

The lower triangular matrix $C_{n,n-i}$ is known as the Catalan triangle and its elements verify the formula

$$C_{n,n-i} = \frac{(2n-i)!(i+1)}{(n-i)!(n+1)!} \quad \text{with } 0 \leq i \leq n.$$

9. HANKEL TRANSFORM

Let $A = \{a_0, a_1, a_2, \dots\}$ be a sequence of real numbers [1]. The Hankel transform of A is the sequence of determinants $H_n = \text{Det}[a_{i+j-2}]$, i.e.

$$H_n = \begin{pmatrix} a_0 & a_1 & a_2 & a_3 & \dots \\ a_1 & a_2 & a_3 & a_4 & \dots \\ a_2 & a_3 & a_4 & a_5 & \dots \\ a_3 & a_4 & a_5 & a_6 & \dots \\ \dots & \dots & \dots & \dots & \dots \end{pmatrix}.$$

The upper-left $n \times n$ subdeterminant of H_n is called the Hankel determinant of order n of the sequence A . The Hankel transform of the Catalan sequence is the sequence $\{1, 1, 1, \dots\}$ [20]. In [17], they show that the Hankel transform of the sum of consecutive generalized Catalan numbers is equal to the bisection of the Fibonacci sequence.

Considering the Catalan transform of the k -Narayana sequence, we get:

$$\begin{aligned}
HCN_{k,1} &= \text{Det}[1] = 1 \\
HCN_{k,2} &= \begin{vmatrix} 1 & 1+k \\ 1+k & k^2+2k+2 \end{vmatrix} = 1
\end{aligned}$$

$$HCN_{k,3} = \begin{vmatrix} 1 & 1+k & k^2+2k+2 \\ 1+k & k^2+2k+2 & k^3+3k^2+5k+6 \\ k^2+2k+2 & k^3+3k^2+5k+6 & k^4+4k^3+9k^2+14k+18 \end{vmatrix} = -2k-2$$

10. CONCLUSION

We give some properties and theorems related to the k -Narayana numbers and show Cassini's identity and limit of the k -Narayana sequence. We also calculate the determinant of the k -Narayana numbers and show that their determinant are fixed terms. Also, we give sum of the first terms of the k -Narayana sequence and we deduce many properties of the k -Narayana numbers and relate with the so-called Pascal 3-triangle. As well as we give an expression of k^n as a k -Narayana number. Finally, we give the $CN_{k,n}$ description of the Catalan transformation of k -Narayana $N_{k,n}$ sequences. Furthermore, we write $CN_{k,n}$ transformation as the multiplying of Catalan matrix C which is the lower triangular matrix, and the N_k matrix of $n \times 1$ type. We calculate determinants of matrixes which are formed with $CN_{k,n}$ by using Hankel transform. These identities can be used to develop new identities of sequences.

APPENDIX A. CELL COMPLEXES

A planar Whitehead cell complex K [16] (usually called a CW complex) is a collection of n -dimensional minimal cells $e_\alpha^n, n \in \{0, 1, 2\}$, *i.e.*,

$$K = \{e_\alpha^n \subset \mathbb{R}^2 : n \in \{0, 1, 2\}\}.$$

in the Euclidean plane π .

Definition 2. A cell subcomplex $shE := \{e_\alpha^n\} \in 2^K$ (**shape complex**) is a **closed subcomplex**, provided the subcomplex includes both a nonempty interior (denoted by $int(e_\alpha^n)$) and its boundary (denoted by $bdy(e_\alpha^n)$). In effect, shE is closed, provided

$$shE = int(shE) \cup bdy(shE) \text{ (Closed subcomplex)}.$$

Let 2^π be the collection of all subsets in the Euclidean plane π . In the plane, a Whitehead **Closure-finite Weak (CW)** cell complex $K \in 2^\pi$ has two properties, namely,

- C:** A cell complex K is **closure-finite**, provided each cell $e_\alpha^n \in K$ is contained in a finite subcomplex of K . In addition, each cell $e_\alpha^n \in K$ has a finite number of immediate faces. One cell e_α^n is an **immediate face** of another cell e_α^m , provided $e_\alpha^n \cap e_\alpha^m \neq \emptyset$ [22] (also called a **common face** [13, p. 3]).
- W:** The plane π has a **weak topology** induced by cell complex K , *i.e.*, a subset $S \in 2^\pi$ is closed, if and only if $S \cap e_\alpha^n$ is also closed in e_α^n for each n, α [22, §5.3, p. 65].




A collection $K \in 2^\pi$ is called a **CW complex**, provided it has the closure-finite property and π has the weak topology property induced by K .

Minimal cell planar complexes are given in Table 4.

Remark 4. Closure finite cell complexes with weak topology (briefly, CW complexes) were introduced by J.C.H. Whitehead [24], later formalized in [25]. In this work¹, a **cell complex** K (or complex) [25, §4, p. 221] is a Hausdorff space (union

¹Here, we use $cl(e^n)$ (closure of a cell) and $bdy(e^n)$ (contour of a cell) used in this paper, instead of Whitehead's e^n and $\partial(e^n)$.

TABLE 4. Minimal Planar Cell Complexes

Minimal Complex	Cell $e^n : n \in \{0, 1, 2\}$	Planar Geometry	Interior
	e^0	Vertex	nonempty
	e^1	Edge	line segment w/o end points
	e^2	Filled triangle	nonempty triangle interior w/o edges

of disjoint open cells e, e^n, e_i^n) such that the closure of an n cell $e^n \in K$ (denoted by $cl(e^n)$ is the image of a map $f : \sigma^n \rightarrow cl(e^n)$, where σ^n is a fixed n -simplex and where the boundary $bdy(e^n)$ (otherwise known as the **contour** of a complex) is defined by

$$\text{Complex contour} \rightarrow \text{closure } cl(e^n) \text{ minus } Int(e^n) \text{ interior}$$

$$bdy(e^n) = \overbrace{f(bdy(e^n))} = cl(e^n) - int(e^n).$$

Notice that a subcomplex $X \subset K$ has the weak topology, since X is the union of a finite number intersections $X \cap cl(e)$ for single cells $e \in K$ [25, §5, p. 223]. From a geometric perspective, a cell complex is a triangulation of the CW space K [24, p. 246]. ■

APPENDIX B. SHAPE COMPLEXES

A shape complex has two basic parts, namely, contour and interior, introduced in [15].

Definition 3. A **planar shape contour** of a shape complex shE (denoted by $bdy(shE)$) is a simple, closed curve with no self-loops. ■

Each shape complex shE has a nonempty interior that excludes all points on the shape contour.

Definition 4. **Shape complex interior.**

Set of points bounded by shE contour

$$int(shE) = \overbrace{shE} \setminus bdyE. \quad \blacksquare$$

The fundamental parts of every shape complex are gathered together in the closure of a shape complex, definite using the Hausdorff distance [8] (see, also, [9, §23, p. 128]) between all points in a CW complex K and a shape shE .

Definition 5. Closure of a planar shape shE (denoted by $cl(shE)$) in a CW space K is defined by

$$\text{Hausdorff distance } D(x, shE) = \inf \{\|x - p\| : p \in shE\}.$$

$$\overbrace{cl(shE)} = \inf \{x \in X : D(x, shE) = 0\}.$$

In other words, we have the closure of a planar shape shE is a finite bounded region of the Euclidean plane such that

$$\text{cl}(shE) \text{ includes its contour \& its interior}$$

$$\overbrace{cl(shE)} = \text{bdy}(shE) \cup \text{Int}(shE).$$

APPENDIX C. FINITE 1-CYCLES AND THEIR CYCLIC GROUP PRESENTATION

Finite 1-cycles in a CW space and their cyclic group presentations [16], [4] are briefly considered in this section. In arriving at cyclic group presentation derived from Pascal 3-triangles, we first consider 1-cycles, filled 1-cycles and shape complexes in a CW space.

Definition 6. A **1-cycle** E (denoted by $cycE$) in a CW space is a finite, nonempty collection of path-connected vertexes $\{e^0 \in K : e_i^0, i \geq 1, \text{ is a zero cell}\}$ on 1-cells (edges) with no end vertex.

Definition 7. A **filled 1-cycle** in a CW space is a 1-cycle $cycE$ has a boundary $bdy(cycE)$ (called its **contour**) that is a simple, closed curve with a nonempty interior $int(cycE)$.

Definition 8. In a CW space K , a **shape complex** E (denoted by shE) is a finite region of K bounded by a simple closed curve with nonempty interior.

Theorem 8. Every filled 1-cycle in a CW complex K is a shape complex of K .

Proof. Immediate from Def. 7 and Def. 8. \square

We can select what is known as a generator vertex in each 1-cycle.

Definition 9. A generator vertex v_0 in a 1-cycle $cycE$ is the first vertex at the beginning of every path leading to any other vertex in $cycE$. Let $|cycE|$ be the number of vertexes in $cycE$ and let the set $\langle v_0 \rangle$ be defined [19, p. 21] by

$$\langle v_0 \rangle = \{v \in cycE : v = kv_0, k \in \{1, \dots, |cycE|\}\}.$$

Recall that a **finite group** G is a nonempty set equipped with an associative binary operation \circ (denoted by $G(\circ)$) such that G has the following members:

identity e : $g \circ e = g$ for every $g \in G$.

inverse $-g$: $g \circ -g = e$ for every $g \in G$.

A group G is **Abelian**, provided $g \circ g' = g' \circ g$ for every $g, g' \in G$.

Definition 10. A finite Abelian group G with a generator element $v_0 \in G$ (denoted by $G(\langle v_0 \rangle, \circ)$) is a **cyclic Abelian group**, provided

$$G = \{g \in G : g \in \langle v_0 \rangle\}.$$

A cyclic group with p generators is denoted by \mathbb{Z}_p [6].

Intuitively, every boundary of a cell complex is a cycle [6, §4.8, p. 104]. This intuition leads to the following useful Lemma.

Lemma 2. *Every finite 1-cycle in a CW space has a corresponding Abelian cyclic group.*

Proof. Let $\text{cyc}E$ be a 1-cycle in a CW space K and let v_0 be a vertex in $\text{cyc}E$. From Def. 6, $\text{cyc}E$ is path-connected. Let $+$ be a forward move (traversal) of an edge between adjacent vertexes in $\text{cyc}E$ and let $v \in \langle v_0 \rangle$, i.e., $v = kv_0$. For each pair $v, v' \in \text{cyc}E$, we have

$$v + v' = kv_0 + k'v_0 = (k + k')v_0 = (k' + k)v_0 = v + v'.$$

Consequently, $+$ is Abelian. Similarly, it can be shown that $+$ is associative. For every $v \in \text{cyc}E$, there is a reverse move $-v$ (the additive inverse of v). Let $0v_0 \in \langle v_0 \rangle$ represent 0 moves from v_0 , i.e., $0v_0 = v_0$. The element $0v_0$ serves as an identity element. For each $v = kv_0 \in \langle v_0 \rangle$, we have

$$\begin{aligned} v + 0v_0 &= kv_0 + 0v_0 = (k + 0)v_0 = kv_0 = v, \text{ and} \\ v - v &= kv_0 - kv_0 = (k - k)v_0 = 0v_0 = v_0. \end{aligned}$$

Hence, $\text{cyc}E(\langle v_0 \rangle, +)$ is an Abelian cyclic group. \square

From Lemma 2, we obtain the following result for shape contours defined by the boundary of a collection of nested cycles.

Theorem 9. *The contour of every planar shape complex in a CW space has a cyclic Abelian group presentation.*

Proof. Let $\text{cyc}E$ be a 1-cycle in a CW space K and let 2^K be the collection of all subsets of K . Further, let the mapping $f : 2^K \rightarrow 2^K$ be defined by

$$\begin{aligned} f(\text{cyc}E) &= \{kg : 0 \leq k \leq |\text{cyc}E| \ \& \ g \in \text{cyc}E\} \text{ (from Lemma 2)} \\ &= G(\langle g \rangle, +) \\ &= \mathbb{Z}_1. \end{aligned}$$

Hence, $\text{cyc}E$ has a cyclic Abelian group presentation. \square

From Theorem 8, we obtain the following result for shapes defined by a collection of path-connected vertexes in nested cycles, attached to each other to form a vortex. Recall that a finite free Abelian group has n generators, $n \geq 1$.

Theorem 10. *Every shape complex that is a collection of nested 1-cycles attached to each other pairwise via edges in a CW space, has a free finitely-generated Abelian group presentation.*

Proof. Let $\text{vor}E$ be a vortex, which is a collection n 1-cycles $\{\text{cyc}_iE, 1 \leq i \leq n\}$ attached to each other via edges in a CW space K and let 2^K be the collection of all subsets of K . Each 1-cycle cyc_iE has its distinguished vertex $g_i \in \text{cyc}_iE$ that is a generator of a cyclic group. Further, let the mapping $f : 2^K \rightarrow 2^K$ be defined by

$$\begin{aligned} f(\text{vor}E) &= \{\{kg_i\} : 0 \leq k \leq |\text{cyc}_iE| \ \& \ g_i \in \text{cyc}_iE, 1 \leq i \leq n\} \text{ (from Lemma 2).} \\ &= G(\langle g_1, \dots, g_n \rangle, +) \\ &= \mathbb{Z}_n. \end{aligned}$$

Hence, $\text{vor}E$ has a finite free Abelian group presentation. \square

REFERENCES

1. P. Barry, *A Catalan transform and related transformations on integer sequences*, J. Integer Seq. **8** (2005), no. 4, 1–24, MR2165334.
2. ———, *Generalized catalan numbers associated with a family of pascal-like triangles*, J. Integer Seq. **22** (2019), no. 5, 1–50, Art. 19.5.8, MR4008156.
3. A. Cvetkovi'c, R. Rajkovi'c, and M. Ivkovi'c, *Catalan numbers, and Hankel transform and Fibonacci numbers*, J. Integer Seq. **5** (2002), no. 1, 1–8, Article 02.1.3, MR1919940.
4. A. Cattabriga and M. Mulazzani, *Strongly-cyclic branched coverings of (1,1)-knots and cyclic presentations of groups*, Math. Proc. Cambridge Philos. Soc. **135** (2003), no. 1, 137–146, MR1990837.
5. G. Edgar, *Measure, topology, and fractal geometry*, Springer, New York, 2008, xvi+268 pp. ISBN: 978-0-387-74748-4, MR2356043.
6. P. Giblin, *Graphs, surfaces and homology, 3rd ed.*, Cambridge University Press, Cambridge, GB, 2016, xx+251 pp. ISBN: 978-0-521-15405-5, MR2722281, first edition in 1981, MR0643363.
7. R.L. Graham, D.E. Knuth, and O. Patashnik, *Matematyka konkretna. (polish) [concrete mathematics]*, Wydawnictwo Naukowe PWN, Addison Wesley, Warsaw, Reading, MA, 1998, 719 pp., ISBN: 83-01-12124-6 68-01, MR1678922.
8. F. Hausdorff, *Grundzüge der mengenlehre*, Veit and Company, Leipzig, 1914, viii + 476 pp.
9. ———, *Set theory, trans. by j.r. aumann*, AMS Chelsea Publishing, Providence, RI, 1957, 352 pp.
10. M. Milich, *Kreise für planare matchings (German)*, Bachelor thesis, TU Berlin, 2018.
11. M. Milich, T. Mütze, and M. Pergel, *On flips in planar matchings*, Discrete Appl. Math. **289** (2021), 427–445, MR4184338.
12. S.A. Naimpally and J.F. Peters, *Topology with applications. Topological spaces via near and far*, World Scientific, Singapore, 2013, xv + 277 pp, Amer. Math. Soc. MR3075111.
13. S. Öztun , N. Bildik, and A. Mutlu, *The construction of simplicial groups in digital images*, J. Inequal. Appl. **2013:143** (2013), 351–356, MR3055827.
14. Nārāyana Paṇḍita, *Gaṇita kaumudī (sanskrit)*, 1356 A.D., trans. by P. Singh in 1998, MR1739184.
15. J.F. Peters, *Proximal planar shapes. correspondence between triangulated shapes and nerve complexes*, Bulletin of the Allahabad Mathematical Society **33** (2018), 113–137, MR3793556, Zbl 06937935, Review by D. Leseberg (Berlin).
16. ———, *Computational geometry, topology and physics of digital images with applications. Shape complexes, optical vortex nerves and proximities*, Springer Nature, Cham, Switzerland, 2020, xxv+440 pp., ISBN 978-3-030-22191-1/hbk; 978-3-030-22192-8/ebook, Zbl07098311.
17. P.M. Rajkovi c, M.D. Petkovi c, and P. Barry, *The hankel transform of the sum of consecutive generalized catalan numbers*, Integral Transforms Spec. Funct. **18** (2007), no. 3-4, 70–74, MR2319589.
18. J.L. Ramirez and V.F. Sirvent, *A note on the k-narayana sequence*, Ann. Math. Inform. **45** (2015), 91–105, MR3438816.
19. J.J. Rotman, *The theory of groups. An introduction. fourth edition.*, Springer-Verlag, New York, 1965, 1995, xvi+513 pp. ISBN: 0-387-94285-8, MR1307623.
20. N.J.A. Sloane, *The on-line encyclopedia of integer sequences*, Annales Mathematicae et Informaticae (2020), MR3822822.
21. A. Stakhov, *The generalized golden proportions, a new theory of real numbers, and ternary mirror-symmetrical arithmetic*, Chaos, Solitons and Fractals **33** (2007), no. 2, 315–334, MR2313546.
22. R.M. Switzer, *Algebraic topology – homology and homotopy*, Springer, Berlin, 2002, xii+526 pp., Zbl 1003.55002.
23. M. Tastan and E. Özkan, *Catalan transform of the k-jacobsthal sequence*, Electron. J. Math. Anal. Appl. **8** (2020), no. 2, 720–74, ISSN: 2090-729X, MR4087829.
24. J.H.C. Whitehead, *Simplicial spaces, nuclei and m-groups*, Proceedings of the London Math. Soc. **45** (1939), no. 4, 243–327, MR1576810.
25. ———, *Combinatorial homotopy. I.*, Bull. Amer. Math. Soc. **55** (1949), no. 3, 213–245, MR0030759.

DEPARTMENT OF MATHEMATICS, FACULTY OF ARTS AND SCIENCES, ERZINCAN BINALI YILDIRIM UNIVERSITY, ERZINCAN, TURKEY

Email address: eozkanmath@gmail.com; eozkan@erzincan.edu.tr

GRADUATE SCHOOL OF NATURAL AND APPLIED SCIENCES, ERZINCAN BINALI YILDIRIM UNIVERSITY, ERZINCAN, TURKEY

Email address: bahar_kuloglu@hotmail.com

COMPUTATIONAL INTELLIGENCE LABORATORY, UNIVERSITY OF MANITOBA, WPG, MB, R3T 5V6, CANADA AND DEPARTMENT OF MATHEMATICS, FACULTY OF ARTS AND SCIENCES, ADIYAMAN UNIVERSITY, 02040 ADIYAMAN, TURKEY

Email address: james.peters3@umanitoba.ca