



HAL
open science

Modular Synthesis of 9,10-Dihydroacridines through an ortho-C Alkenylation/Hydroarylation Sequence between Anilines and Aryl Alkynes in Hexafluoroisopropanol

Shengdong Wang, Guillaume Force, Jean-François Carpentier, Yann Sarazin, Christophe Bour, Vincent Gandon, David Leboeuf

► **To cite this version:**

Shengdong Wang, Guillaume Force, Jean-François Carpentier, Yann Sarazin, Christophe Bour, et al.. Modular Synthesis of 9,10-Dihydroacridines through an ortho-C Alkenylation/Hydroarylation Sequence between Anilines and Aryl Alkynes in Hexafluoroisopropanol. *Organic Letters*, 2021, 23 (7), pp.2565-2570. 10.1021/acs.orglett.1c00487. hal-03245060

HAL Id: hal-03245060

<https://hal.science/hal-03245060v1>

Submitted on 15 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Modular Synthesis of 9,10-Dihydroacridines through an *ortho*-C Alkenylation/Hydroarylation Sequence Between Anilines and Aryl Alkynes in Hexafluoroisopropanol.

Shengdong Wang,^{†,‡} Guillaume Force,[†] Jean-François Carpentier,^{||} Yann Sarazin,^{||} Christophe Bour,[†] Vincent Gandon^{*†,⊥} and David Leboeuf^{*§}

[†] Institut de Chimie Moléculaire et des Matériaux d'Orsay (ICMMO), CNRS UMR 8182, Université Paris-Saclay, Bâtiment 420, 91405 Orsay, France.


[‡] The Fifth Affiliated Hospital, Key Laboratory of Molecular Target & Clinical Pharmacology and the State Key Laboratory of Respiratory Disease, School of Pharmaceutical Sciences, Guangzhou Medical University, Guangzhou, Guangdong 511436, China.

^{||} Univ. Rennes, CNRS UMR 6226, Institut des Sciences Chimiques de Rennes (ISCR), 35000 Rennes, France.

[⊥] Laboratoire de Chimie Moléculaire (LCM), CNRS UMR 9168, Ecole Polytechnique, Institut Polytechnique de Paris, route de Saclay, 91128 Palaiseau cedex, France.

[§] Institut de Science et d'Ingénierie Supramoléculaires (ISIS), CNRS UMR 7006, Université de Strasbourg, 8 allée Gaspard Monge, 67000 Strasbourg, France.

Supporting Information Placeholder




ABSTRACT: 9,10-Dihydroacridines are frequently encountered as key scaffolds in OLEDs. Yet, accessing those compounds from feedstock precursors typically requires multiple steps. Herein, a modular one-pot synthesis of 9,10-dihydroacridine frameworks is achieved through a reaction sequence featuring a selective *ortho*-C alkenylation of diarylamines with aryl alkynes followed by an intramolecular hydroarylation of the olefin formed as intermediate. This transformation was accomplished by virtue of the combination of hexafluoroisopropanol and triflimide as a catalyst that triggers the whole process.

In the past decade, 9,10-dihydroacridines have emerged as privileged structural motifs in chemistry owing to their remarkable electronic properties. They are now extensively used in the field of materials science as electron-donor units, notably for organic light-emitting diode (OLED) devices. Hence, they have been employed as an integral part of fluorescent emitters, thermally activated delayed fluorescence (TADF) materials or hole-transport materials.¹ Moreover, they have recently proven particularly effective in other areas such as photocatalysis,² detection of explosives,³ and medicinal chemistry.⁴ Yet, despite the wealth of reports regarding their utilization, the development of efficient and versatile synthetic methods to access those building blocks from readily available precursors remain strikingly sparse,⁵ representing a serious impediment to broadening their range of applications. Currently, their preparation usually requires multi-step synthesis from anilines, which are furthermore not always commercially available (Scheme 1a).^{1j} On this basis, most of the applications reported tend to be limited to simple 9,10-dimethyl and -diphenylacridines. As an alternative, the group of Stephan described, in 2017, an elegant approach to access 9,10-dihydroacridines following a one-pot sequence featuring an

ortho-C-alkenylation of anilines with an aryl alkyne and a subsequent hydroarylation of the styrene intermediate in the presence of a phosphonium dication catalyst (Scheme 1b).^{5c} The selective *ortho*-C-alkenylation of the aniline with the alkyne is the key of the process;^{6,7} however, the applicability of this type of reactivity is still underdeveloped when compared to the *ortho*-C alkylation of anilines with olefins.⁸ It might be explained by the formation of a *sp* hybridized carbenium center, which is a sluggish electrophile compared to its *sp*² carbenium counterpart generated from alkenes.⁹ As a result, the scope exhibited by this reaction sequence proved to be limited. In this context, we questioned whether we could address the limitations of this transformation by employing hexafluoroisopropanol (HFIP) as a solvent.¹⁰ In the past years, HFIP has become a prominent solvent in organic synthesis due to its combination of atypical properties, including strong H-bond donating ability, low nucleophilicity, redox stability, and mild acidity.¹¹ It allowed us to achieve reactions that were unlikely to take place in traditional organic solvents,^{8k,10d,12} emphasizing that, when HFIP was associated with a Lewis acid or a Brønsted acid, the acidity of the corresponding combination could be considerably boosted to even activate unreactive


Scheme 1. Strategies to access 9,10-dihydroacridines. $[P]^+ = [(PhO)P(2(N-Mepy))Ph_2]^{2+}$.



substrates such as highly deactivated styrenes. Recently, we applied this principle to the *ortho*-C alkylation of anilines with alkenes,^{8k} which relies on a concerted-like mechanism that is fostered by the presence of HFIP and prevents common regioselectivity issues associated with this type of process (Scheme 1c). While rather efficient and general, this transformation is still fraught with a few limitations in terms of reactivity. For instance, electron-rich styrenes or triarylamines preferentially led to *para*-adducts due to a facile protonation of the olefin moiety under the reaction conditions. Following this work, we assumed that this strategy could be transposed to aryl alkynes to afford the target 9,10-dihydroacridines. We hypothesized that, since the protonated alkynes (vinyl carbocations) are less reactive than protonated alkenes, a concerted mechanism pathway would be predominant, bypassing the above-mentioned limitations. This strategy would thus enable a straightforward and modular synthesis of 9,10-dihydroacridines so that the properties of the corresponding organic materials could be easily tuned, enabling new applications for this class of compounds. Herein, we disclose our findings regarding this reaction sequence to produce an array of structurally varied 9,10-dihydroacridines in good to high yields and with an excellent functional group compatibility. Moreover, we show that the developed conditions also allow the preparation of xanthenes and thioxanthenes.


We started our investigations by examining the reaction between 1-ethynyl-4-(trifluoromethyl)benzene **1a** (highly

deactivated aryl alkyne) and diphenylamine **2a** in HFIP (See the Supporting Information for more details). We established that using Brønsted acid HNTf₂¹³ as catalyst proved to be optimal to deliver the target 9,10-dihydroacridine **3aa** in 86% yield after 16 h at 80 °C (Equation 1).¹⁴ A survey of the reaction optimization revealed that HFIP is crucial for the reaction outcome as another fluorinated solvent such as trifluoroethanol (TFE) yielded the product **3aa** in a low yield (5%); only traces of **3aa** were observed in common solvents such as 1,2-DCE, toluene and nitromethane. Gratifyingly, the reaction worked smoothly on a larger scale (5 mmol), furnishing **3aa** in 81% yield (1.37 g). An issue regularly mentioned with the utilization of HFIP is its cost, but it is important to stress that, because of its low boiling point, it can be easily recovered by distillation,¹⁵ counterbalancing this issue.




We next examined the scope of this transformation (Scheme 2), first evaluating a variety of terminal aryl alkynes (**1aa-1ua**) in reactions with diphenylamine **2a**. In this respect, the tested aryl alkynes bearing moderate and strong electron-donating and -withdrawing groups at the *para*-position were all tolerated and


Scheme 2. Scope of 9,10-dihydroacridines and other heterocycles.




Scope alkynes




Scope diarylamines




Scope triarylamines




Scope heterocycles



Limitations



Applications



provided the corresponding 9,10-dihydroacridines **3aa-3ka** in high yields, ranging from 73 to 98%. Of note, in the case of strong electron-withdrawing groups such as nitrile and nitro (**1c** and **1d**), the reaction had to be conducted at 120 °C to reach complete conversion of the alkyne starting materials. In the case of strong electron-donating groups such as methoxy and amine (**1j** and **1k**), the reaction proceeded faster than the previous examples (6 vs 16 h), providing the target products **3ja** and **3ka** in 86 and 79% yield, respectively. The use of highly electron-poor alkynes such as 3,5-bis(trifluoromethyl)phenyl derivative required to increase the catalyst loading to 10 mol% to afford 9,10-dihydroacridine **3la** in 83% yield. The presence of an *ortho*-substituent was also well tolerated (**3ma**, 87%). The reaction sequence was not limited to phenyl group but could be also extended to a naphthyl or thienyl group, delivering both **3na** and **3oa** in 80% yield. More importantly, the transformation occurred also for internal alkynes incorporating alkyl substituents (e.g., adducts **3pa** and **3qa** were generated in 73 and 70% yield), albeit at a slower rate than the reactions involving terminal alkynes. Gratifyingly, dialkynes **1s-1t** and trialkyne **1u** were readily accommodated in the reactions to provide multiple-9,10-dihydroacridine-bearing compounds in high yields (71-89%). On the other hand, bis-alkyl alkynes proved to be not competent precursors for the reaction, essentially leading to decomposition of the substrate with only traces of the target product **3ra** detected.

We then explored the scope with respect to the diarylamine component (**2b-2n**), using representative aryl alkynes **1c** (R = CN) and **1g** (R = H) as benchmark precursors. In the case of 2,7-substituted 9,10-dihydroacridines (**3cb-3cj**, **3gd**, **3gf** and **3gi**), the reaction demonstrated a broad functional group compatibility from both symmetrical and dissymmetrical diarylamines to provide the products in medium to high yields (57-95%). Both electron-donating and -withdrawing substituents were amenable to the reaction; notably, bromide substituents were easily introduced to generate products **3ce**, **3ci** and **3gi** (57-68%), which constitute useful platforms for further derivatizations. In addition, 4,5-substituted 9,10-dihydroacridines such as **3ck** could be generated in 73% yield. *N*-alkyl diarylamines such as **1l** also underwent the reaction process in high yields (up to 85%), employing both terminal and internal alkynes (**3al**, **3cl**, **3gl** and **3pl**). By incorporating a naphthyl moiety, we could prepare complex polycyclic structures (**3cm**, **3cn** and **3gn**) in yields ranging from 61 to 87%.

Triarylamines, notably triphenylamine **2o**, displayed a satisfying reactivity with respect to this process to form the products in medium to high yields (41-93%), regardless of the electronic demand of the aryl alkyne and its nature (terminal or internal). Another key feature of this method is that the reaction of diphenyl sulfide **2r** in place of **2a** yielded the corresponding thioxanthene **3cr** in 83% yield. In the same vein, the reaction with diphenyl ether **2s** underwent the diarylation to give xanthene **3gs** in 67% yield. However, it should be stressed that diphenyl ether is less reactive than diarylamines and diphenyl sulfide as it was ineffective with highly deactivated aryl alkyne such as **1c** (10% yield after one week at 120 °C). Regarding the limitations of this approach, we noticed that the use of *meta*-substituted diarylamines (**2s** and **2t**) and triarylamines with non-identical aryl groups led to the formation of a mixture of 2 regioisomers that could not be separated in some cases by flash column chromatography.

To further illustrate the synthetic utility of this method, we carried out few post-functionalizations. As stated above, the reaction sequence with dissymmetrical diarylamines occurred with ease to return the corresponding products in high yields. Those results could be used to our advantage to generate densely functionalized 9,10-dihydroacridines such as **4** incorporating four different aryl units, following a Ullmann C-N cross-coupling.¹⁶ It also represents a simple way to circumvent the reactivity issue mentioned with triarylamines. Additionally, we executed a Pd(II)-catalyzed C-H activation to convert **3ma** into pentacyclic compound **5** in 94% yield.^{17,18}

In summary, we have developed an efficient protocol for the assembly of 9,10-dihydroacridines and related heterocycles from inexpensive precursors through the cooperation of HFIP and a Brønsted acid catalyst. The wide array of substrates tolerated validates the utility of this transformation as a means to provide a rapid access to molecules that could find applications in the field of materials science and photocatalysis. In addition, the viability at scale of this method was also demonstrated.

ASSOCIATED CONTENT

Supporting Information

Experimental procedures, characterization data and NMR spectra of all new compounds (PDF). This material is available free of charge via the Internet at <http://pubs.acs.org/>

FAIR Data is available as Supporting Information for Publication and includes the primary NMR FID files for compounds: **3aa-3'gv**, **4** and **5**.

AUTHOR INFORMATION

Corresponding Authors

* dleboeuf@unistra.fr

* vincent.gandon@universite-paris-saclay.fr

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENT

We gratefully thank the ANR (ANR-17-CE07-0003 funding for SW and ANR-16-CE07-0022 funding for GF), the CNRS, Ecole Polytechnique and Université Paris-Saclay for the support of this work. We used the OCCIGEN high performance cluster of the CINES.

REFERENCES

- (1) For selected applications, see: (a) Park, M. S.; Lee, J. Y. Indolo Acridine-Based Hole-Transport Materials for Phosphorescent OLEDs with Over 20% External Quantum Efficiency in Deep Blue and Green. *Chem. Mater.* **2011**, *23*, 4338-4343. (b) Méhes, G.; Nomura, H.; Zhang, Q.; Nakagawa, T.; Adachi, C. Enhanced Electroluminescence Efficiency in a Spiro-Acridine Derivative through Thermally Activated Delayed Fluorescence. *Angew. Chem. Int. Ed.* **2012**, *51*, 11311-11315. (c) Kim, M.; Lee, J. Y. Improved Power Efficiency in Deep Blue Phosphorescent Organic Light-Emitting Diodes Using an Acridine Core Based Hole Transport Material. *Org. Electron.* **2012**, *13*, 1245-1249. (d) Zhang, Q.; Li, B.; Huang, S.; Nomura, H.; Tanaka, H.; Adachi, C. Efficient Blue Organic Light-Emitting Diodes Employing Thermally Activated Delayed Fluorescence. *Nat. Photonics* **2014**, *8*, 326-332. (e) Zhang, Y.-X.; Zhang, L.; Cui, L.-S.; Gao, C.-H.; Chen, H.; Li, Q.; Jiang, Z.-Q.; Liao, L.-S. Control of Conjugation Degree via Position Engineering to Highly Efficient Phosphorescent Host Materials. *Org. Lett.* **2014**, *16*, 3748-3751. (f) Liu, X.-Y.; Liang, F.;

- Yuan, Y.; Jiang, Z.-Q.; Liao, L.-S. Utilizing 9,10-Dihydroacridine and Pyrazine-Containing Donor-Acceptor Host Materials for Highly Efficient Red Phosphorescent Organic Light-Emitting Diodes. *J. Mater. Res. C* **2016**, *4*, 7869-7874. (g) Zhang, L.; Zhang, Y.-X.; Hu, Y.; Shi, X.-B.; Jiang, Z.-Q.; Wang, Z.-K.; Liao, L. S. Highly Efficient Blue Phosphorescent Organic Light-Emitting Diodes Employing a Host Material with Small Bandgap. *ACS Appl. Mater. Interfaces* **2016**, *8*, 16186-16191. (h) Zeng, W.; Lai, H.-Y.; Lee, W.-K.; Jiao, M.; Shiu, Y.-J.; Zhong, C.; Gong, S.; Zhou, T.; Xie, G.; Sarma, M.; Wong, K.-T.; Wu, C.-C.; Yang, C. Achieving Nearly 30% External Quantum Efficiency for Orange-Red Organic Light Emitting Diodes by Employing Thermally Activated Delayed Fluorescence Emitters Composed of 1,8-Naphthalimide-Acridine Hybrids. *Adv. Mater.* **2017**, *29*, 1704961. (i) Yu, L.; Wu, Z.; Xie, G.; Zeng, W.; Ma, D.; Yang, C. Molecular Design to Regulate the Photophysical Properties of Multifunctional TADF Emitters Towards High-Performance TADF-Based OLEDs with EQES up to 22.4% and Small Efficiency Roll-Offs. *Chem. Sci.* **2018**, *9*, 1385-1391. (j) Liu, X.-Y.; Ma, Y.-Y.; Zhang, W.; Song, B.; Ding, L.; Fung, M.-K.; Fan, J. A Novel Linking Strategy of Using 9,10-Dihydroacridines to Construct Efficient Host Materials for Red Phosphorescent Organic Light-Emitting Diodes. *Chem. Eur. J.* **2018**, *24*, 11755-11762. (k) Wada, Y.; Kubo, S.; Kaji, H. Adamantyl Substitution Strategy for Realizing Solution-Processable Thermally Stable Deep-Blue Thermally Activated Delayed Fluorescence Materials. *Adv. Mater.* **2018**, *30*, 1705641. (l) Yang, Y.; Wang, S.; Zhu, Y.; Wang, Y.; Zhan, H.; Cheng, Y. Thermally Activated Delayed Fluorescence Conjugated Polymers with Backbone-Donor/Pendant-Acceptor Architecture for Nondoped OLEDs with High External Quantum Efficiency and Low Roll-Off. *Adv. Funct. Mater.* **2018**, *28*, 1706916. (m) He, X.; Ren, S.; Liu, H.; Zhao, S.; Liu, F.; Du, C.; Min, J.; Zhang, H.; Lu, P. Efficient Nondoped Pure Blue Organic Light-Emitting Diodes Based on an Anthracene and 9,9-Diphenyl-9,10-dihydroacridine Derivative. *Chem. Asian J.* **2020**, *15*, 163-168.
- (2) (a) Buss, B. L.; Lim, C.-H.; Miyake, G. M. Dimethyl Dihydroacridines as Photocatalysts in Organocatalyzed Atom Transfer Radical Polymerization of Acrylate Monomers. *Angew. Chem. Int. Ed.* **2020**, *59*, 3209-3217. (b) Liu, Y.; Chen, Q.; Tong, Y.; Ma, Y. 9,9-Dimethyl Dihydroacridine-Based Organic Photocatalyst for Atom Transfer Radical Polymerization from Modifying "Unstable" Electron Donor. *Macromolecules* **2020**, *53*, 7053-7062.
- (3) Andrew, T. L.; Swager, T. M. Detection of Explosives via Photocatalytic Cleavage of Nitroesters and Nitramines. *J. Org. Chem.* **2011**, *76*, 2976-2993.
- (4) Bagriantsev, S. N.; Ang, K.-H.; Gallardo-Godoy, A.; Clark, K. A.; Arkin, M. R.; Renslo, A. R.; Minor Jr., D. L. A High-Throughput Functional Screen Identifies Small Molecule Regulators of Temperature- and Mechano-Sensitive K_{2P} Channels. *ACS Chem. Biol.* **2013**, *8*, 1841-1851.
- (5) For selected examples, see: (a) Mahendar, L.; Satyanarayana, G. Copper Catalyzed Coupling of Protecting Group Free and Sterically Hindered 2-Bromobenzyl Tertiary Alcohols with Phenols and Anilines: Facile Synthesis of Xanthenes and Dihydroacridines. *RSC Adv.* **2016**, *6*, 20588-20597. (b) Yoshida, H.; Kuriki, H.; Fujii, S.; Ito, Y.; Osaka, I.; Takaki, K. Aryne-Imine-Aryne Coupling Reaction via [4+2] Cycloaddition between Aza-*o*-Quinone Methides and Arynes. *Asian J. Org. Chem.* **2017**, *6*, 973-976. (c) LaFortune, J. H. W.; Bayne, J. M.; Johnstone, T. C.; Fan, L.; Stephan, D. W. Catalytic Double Hydroarylation of Alkynes to 9,9-Disubstituted 9,10-Dihydroacridine Derivatives by an Electrophilic Phenoxyphosphonium Dication. *Chem. Commun.* **2017**, *53*, 13312-13315. (d) LaFortune, J. H. W.; Szkop, K. M.; Farinha, F. E.; Johnstone, T. C.; Postle, S.; Stephan, D. W. Probing Steric Influences on Electrophilic Phosphonium Cations: A Comparison of $[(3,5-(CF_3)_2C_6H_3)_3PF]^+$ and $[(C_6F_5)_3PF]^+$. *Dalton Trans.* **2018**, *47*, 11411-11419. (e) Smith, A. J.; Dimitrova, D.; Arokianathar, J. N.; Kolodziejczak, K.; Young, A.; Allison, M.; Poole, D. L.; Leach, S. G.; Parkinson, J. A.; Tuttle, T.; Murphy, J. A. New Reductive Rearrangement of *N*-Arylindoles Triggered by the Grubbs-Stoltz Reagent Et_3SiH/KO^tBu . *Chem. Sci.* **2020**, *11*, 3719-3726.
- (6) (a) Arienti, A.; Bigi, F.; Maggi, R.; Marzi, E.; Moggi, P.; Rastelli, M.; Sartori, G.; Tarantola, F. Regioselective Electrophilic Alkylation of Anilines with Phenylacetylene in the Presence of Montmorillonite KSF. *Tetrahedron* **1997**, *53*, 3795-3804. (b) Fedushkin, I. L.; Nikipelov, A. S.; Morozov, A. G.; Skatova, A. A.; Cherkasov, A. V.; Abakumov, G. A. Addition of Alkynes to a Gallium Bis-Amido Complex: Imitation of Transition-Metal-Based Catalytic Systems. *Chem. Eur. J.* **2012**, *18*, 255-266. (c) Fedushkin, I. L.; Moskalev, M. V.; Baranov, E. V.; Abakumov, G. A. Addition of Diphenylacetylene and Methylvinylketone to Aluminum Complex of Redox-Active Diimine Ligand. *J. Organomet. Chem.* **2013**, *747*, 235-240. (d) Mamedov, N.; Peraka, S.; Kodumuri, S.; Chevella, D.; Marri, M. R.; Nama, N. *Ortho*-Alkenylation of Anilines with Aromatic Terminal Alkynes over Nanosized Zeolite Beta. *RSC Adv.* **2015**, *5*, 78374-78378. (e) Chatupheeraphat, A.; Rueping, M.; Magre, M. Chemo- and Regioselective Magnesium-Catalyzed *ortho*-Alkenylation of Anilines. *Org. Lett.* **2019**, *21*, 9153-9157.
- (7) For the synthesis of related compounds starting from olefins, see: Cooper, P.; Crisenza, G. E. M.; Feron, L. F.; Bower, J. F. Iridium-Catalyzed α -Selective Arylation of Styrenes by Dual C-H Activation. *Angew. Chem. Int. Ed.* **2018**, *57*, 14198-14202.
- (8) For selected examples of *ortho*-C alkylation of anilines with olefins, see: (a) Beller, M.; Thiel, O. R.; Trauthwein, H. Catalytic Alkylation of Aromatic Amines with Styrene in the Presence of Cationic Rhodium Complexes and Acid. *Synlett* **1999**, 243-245. (b) Kaspar, L. T.; Fingerhut, B.; Ackermann, L. Titanium-Catalyzed Intermolecular Hydroamination of Vinylarenes. *Angew. Chem. Int. Ed.* **2005**, *44*, 5972-5974. (c) Cherian, A. E.; Domski, G. J.; Rose, J. M.; Lobkovsky, E. B.; Coates, G. W. Acid-Catalyzed *ortho*-Alkylation of Anilines with Styrenes: An Improved Route to Chiral Anilines with Bulky Substituents. *Org. Lett.* **2005**, *7*, 5135-5137. (d) Anderson, L. L.; Arnold, J.; Bergman, R. G. Proton-Catalyzed Hydroamination and Hydroarylation Reactions of Anilines and Alkenes: A Dramatic Effect of Counteranions on Reaction Efficiency. *J. Am. Chem. Soc.* **2005**, *127*, 14542-14543. (e) Crisenza, G. E. M.; Sokolova, O. O.; Bower, J. F. Branch-Selective Alkene Hydroarylation by Cooperative Destabilization: Iridium-Catalyzed *ortho*-Alkylation of Acetanilides. *Angew. Chem. Int. Ed.* **2015**, *54*, 14866-14870. (f) Song, G.; Luo, G.; Oyamada, J.; Luo, Y.; Hou, Z. *ortho*-Selective C-H Addition of *N,N*-Dimethyl Anilines to Alkenes by a Yttrium Catalyst. *Chem. Sci.* **2016**, *7*, 5265-5270. (g) Zhu, W.; Sun, Q.; Wang, Y.; Yuan, D.; Yao, Y. Chemo- and Regioselective Hydroarylation of Alkenes with Aromatic Amines Catalyzed by $[Ph_3C][B(C_6H_5)_4]$. *Org. Lett.* **2018**, *20*, 3101-3104. (h) Schroeter, F.; Lerch, S.; Kaliner, M.; Strassner, T. Cobalt-Catalyzed Hydroarylations and Hydroaminations of Alkenes in Tunable Aryl Alkyl Ionic Liquids. *Org. Lett.* **2018**, *20*, 6215-6219. (i) Grélaud, S.; Cooper, P.; Feron, L. J.; Bower, J. F. Branch-Selective and Enantioselective Iridium-Catalyzed Alkene Hydroarylation via Anilide-Directed C-H Oxidative Addition. *J. Am. Chem. Soc.* **2018**, *140*, 9351-9356. (j) Rank, C. K.; Özkaya, B.; Patureau, F. W. HBF₄- and AgBF₄-Catalyzed *ortho*-Alkylation of Diarylamines and Phenols. *Org. Lett.* **2019**, *21*, 6830-6834. (k) Wang, S.; Force, G.; Guillot, R.; Carpentier, J.-F.; Sarazin, Y.; Bour, C.; Gandon, V.; Lebœuf, D. Lewis Acid/Hexafluoroisopropanol: A Promoter System for Selective *ortho*-C-Alkylation of Anilines with Deactivated Styrene Derivatives and Unactivated Alkenes. *ACS Catal.* **2020**, *10*, 10794-10802.
- (9) Byrne, P. A.; Kobayashi, S.; Würthwein, E.-U.; Ammer, J.; Mayr, H. Why Are Vinyl Cations Sluggish Electrophiles? *J. Am. Chem. Soc.* **2017**, *139*, 1499-1511.
- (10) For examples of hydroarylation processes in HFIP, see: (a) Richmond, E.; Vuković, V. D.; Moran, J. Nucleophilic Ring Opening of Donor-Acceptor Cyclopropanes Catalyzed by a Brønsted Acid in Hexafluoroisopropanol. *Org. Lett.* **2018**, *20*, 574-577. (b) Roy, S.; Motiwala, H. F.; Koshlap, K. M.; Aubé, J. Hexafluoroisopropanol and Acetyl Chloride Promoted Catalytic Hydroarylation with Phenols. *Eur. J. Org. Chem.* **2018**, 306-315. (c) Richmond, E.; Yi, J.; Vuković, V. D.; Sajadi, F.; Rowley, C. N.; Moran, J. Ring-Opening Hydroarylation of Monosubstituted Cyclopropanes Enabled by Hexafluoroisopropanol. *Chem. Sci.* **2018**, *9*, 6411-6416. (d) Qi, C.; Gandon, V.; Lebœuf, D. Calcium(II)-Catalyzed Intermolecular Hydroarylation of Deactivated

Styrenes in Hexafluoroisopropanol. *Angew. Chem. Int. Ed.* **2018**, *57*, 14245-14249. (e) Takahashi, I.; Fujita, T.; Shoji, N.; Ichikawa, J. Brønsted Acid-Catalysed Hydroarylation of Unactivated Alkynes in a Fluoroalcohol-Hydrocarbon Biphasic System: Construction of Phenanthrene Frameworks. *Chem. Commun.* **2019**, *55*, 9267-9270; (f) Nielsen, C. D.-T.; White, A. J. P.; Sale, D.; Bures, J.; Spivey, A. C. Hydroarylation of Alkenes by Protonation/Friedel-Crafts Trapping: HFIP-Mediated Access to Per-aryl Quaternary Stereocenters. *J. Org. Chem.* **2019**, *84*, 14965-14973. (g) Colomer, I. Hydroarylation of Alkenes Using Anilines in Hexafluoroisopropanol. *ACS Catal.* **2020**, *10*, 6023-6029.

(11) For reviews on HFIP, see: (a) Bégue, J.-P.; Bonnet-Delpon, D.; Crousse, B. Fluorinated Alcohols: A New Medium for Selective and Clean Reaction. *Synlett* **2004**, 18-29. (b) Shuklov, I. A.; Dubrovina, N. V.; Börner, A. Fluorinated Alcohols as Solvents, Cosolvents and Additives in Homogeneous Catalysis. *Synthesis* **2007**, 2925-2943. (c) Sugiishi, T.; Matsugi, M.; Hamamoto, H.; Amii, H. Enhancement of Stereoselectivities in Asymmetric Synthesis Using Fluorinated Solvents, Auxiliaries, and Catalysts. *RSC Adv.* **2015**, *5*, 17269-17282. (d) Wencel-Delord, J.; Colobert, F. A Remarkable Solvent Effect of Fluorinated Alcohols on Transition Metal Catalysed C-H Functionalizations. *Org. Chem. Front.* **2016**, *3*, 394-400. (e) Colomer, I.; Chamberlain, A. E. R.; Haughey, M. B.; Donohoe, T. J. Hexafluoroisopropanol as a Highly Versatile Solvent. *Nat. Rev. Chem.* **2017**, *1*, 0088. (f) Sinha, S. K.; Bhattacharya, T.; Maiti, D. Role of Hexafluoroisopropanol in C-H Activation. *React. Chem. Engl.* **2019**, *4*, 244-253. (g) Yu, C.; Sanjosé-Orduna, J.; Patureau, F.; Pérez-Temprano, M. *Chem. Soc. Rev.* **2020**, *49*, 1643-1652. (h) Pozhydaiev, V.; Power, M.; Gandon, V.; Moran, J.; Lebœuf, D. Exploiting Hexafluoroisopropanol (HFIP) in Lewis and Brønsted Acid-Catalyzed Reactions. *Chem. Commun.* **2020**, *56*, 11548-11564.

(12) (a) Lebœuf, D.; Marin, L.; Michelet, B.; Perez-Luna, A.; Guillot, R.; Schulz, E.; Gandon, V. Harnessing the Lewis Acidity of HFIP Through its Cooperation with a Calcium(II) Salt: Application to the

Aza-Piancatelli Reaction. *Chem. Eur. J.* **2016**, *22*, 16165-16171. (b) Qi, C.; Hasenmaile, F.; Gandon, V.; Lebœuf, D. Calcium(II)-Catalyzed Intra- and Intermolecular Hydroamidation of Unactivated Alkenes in Hexafluoroisopropanol. *ACS Catal.* **2018**, *8*, 1734-1739. (c) Qi, C.; Yang, S.; Gandon, V.; Lebœuf, D. Calcium(II)- and Triflimide-Catalyzed Intramolecular Hydroacyloxylation of Unactivated Alkenes in Hexafluoroisopropanol. *Org. Lett.* **2019**, *21*, 7405-7409.

(13) Zhao, W.; Sun, S. Triflimide (HNTf₂) in Organic Synthesis. *Chem. Rev.* **2018**, *118*, 10349-10392.

(14) For preliminary investigations regarding the mechanism of the reaction, notably the key *ortho*-C alkenylation step, see Supporting Information for details.

(15) Vekariya, R. H.; Aubé, J. Hexafluoro-2-propanol-Promoted Intermolecular Friedel-Crafts Acylation Reaction. *Org. Lett.* **2016**, *18*, 3534-3537.

(16) Sambagio, C.; Marsden, S. P.; Blacker, A. J.; McGowan, P. C. Copper Catalysed Ullmann Type Chemistry: From Mechanistic Aspects to Modern Development. *Chem. Soc. Rev.* **2014**, *43*, 3525-3550.

(17) Campeau, L.-C.; Parisien, M.; Jean, A.; Fagnou, K. Catalytic Direct Arylation with Aryl Chlorides, Bromides, and Iodides: Intramolecular Studies Leading to New Intermolecular Reactions. *J. Am. Chem. Soc.* **2005**, *128*, 581-590.

(18) For another approach to synthesize this type of compounds, see: Gu, Z.-Y.; Liu, C.-G.; Wang, S.-Y.; Ji, S.-J. Pd-Catalyzed Intramolecular Heck Reaction, C(sp²)-H Activation, 1,4-Pd Migration, and Aminopalladation: Chemoselective Synthesis of Dihydroindeno[1,2,3-*kl*]acridines and 3-Arylindoles. *Org. Lett.* **2016**, *18*, 2379-2382.