



HAL
open science

Ultrasound combined with microbubbles modulates signal transduction pathways in blood cells

Maja Mujić, Ragnhild Haugse, Spiros Kotopoulos, Andre Sulen, Odd Helge
Gilja, Michiel Postema, Bjørn Tore Gjertsen

► **To cite this version:**

Maja Mujić, Ragnhild Haugse, Spiros Kotopoulos, Andre Sulen, Odd Helge Gilja, et al.. Ultrasound combined with microbubbles modulates signal transduction pathways in blood cells. MedViz Conference 2013, MedViz; Haukeland University Hospital; University of Bergen; Christian Michelsen Research, Jan 2013, Bergen, Norway. pp.119-120. hal-03245682

HAL Id: hal-03245682

<https://hal.science/hal-03245682>

Submitted on 2 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

MedViz Conference 2013

Ultrasound combined with microbubbles modulates signal transduction pathways in blood cells

Maja Mujic¹, Ragnhild Haugse¹, Spiros Kotopoulos^{2,3}, Andre Sulen¹, Odd Helge Gilja³, Michiel Postema^{2,3}, Bjørn Tore Gjertsen^{1,4}

¹ Institute of Medicine, Section for Haematology, University of Bergen, Bergen, Norway

² Department of Physics and Technology, University of Bergen, Bergen, Norway


³ National Centre for Ultrasound in Gastroenterology, Haukeland University Hospital, Bergen, Norway

⁴ Department of Medicine, Hematology Section, Haukeland University Hospital, Norway

Ultrasound is widely used in medical diagnostics and is considered as a fast, cost-effective and safe imaging modality. Gas-filled microbubbles (MB) are introduced into the blood pool in order to enhance the signal-to-noise ratio in clinical contrast-enhanced ultrasonographical (CEUS) imaging [1]. CEUS is also explored for use in gene and drug delivery, by sonoporation [1]. Low-intensity pulsed ultrasound (LIPUS) is involved in activation of signaling pathways involved in cell stress [2], and is also shown to inhibit cancer cell proliferation and induce apoptosis in leukemia cells [3-7]. Limited is known about the signaling pathways simulated by ultrasound.

Intracellular signaling pathways in the monocytoic cell line MOLM-13 and peripheral blood mononuclear cells (PBMCs) were investigated after treatment with ultrasound alone or in combination with a gas-filled microbubble (MB). Three ultrasound intensities (mechanical index 0.16, 0.59, 0.72) for 10 minutes were employed with or without presence of microbubble contrast agent, using the set-up shown in figure 1. Multiparameter flow cytometry with protein modification specific antibodies was performed at predefined time-points ranging from 10 min and up to 24 hours after sonication to determine impact on intracellular signaling pathways.

Figure 1



MOLM-13 and PBMCs responded similarly to ultrasound and MBs, with phosphorylation of p38 and ERK1/2 MAPK with some cell specific differences.

MedViz Conference 2013

Early induction of p38 MAPK signaling was seen in all cell types, but monocytes had a relative increase in phospho-p38 MAPK compared to MOLM-13, T lymphocyte, B lymphocyte and natural killer cells. Induction of phosphorylation of p38 MAPK was dependent on ultrasound intensities and MB presence. In general, increased phosphorylation of p38 MAPK is observed in response to stress stimuli and is involved in various cellular processes including differentiation and induction of cell death, acting upstream of the tumor suppressor p53 [8]. Furthermore, there is an example of downregulated p53 acetylation after increased phosphorylation of p38 MAPK [9], indicating involvement of an established stress pathway in ultrasound exposure.

Changes in phosphorylation of ERK1/2, AKT, NFκB and p53 as well as p53 acetylation were observed, but without alteration in total p53, clonogenicity or viability. It can be concluded that stress related signaling in leukocytes are affected by diagnostic ultrasound in vitro. The concerted upregulation of phosphorylated p38, ERK1/2 and AKT demonstrated in this study is previously described by other stress stimuli [10], suggesting involvement of a common mechanism.

1. Postema, M. and O.H. Gilja, *Contrast-enhanced and targeted ultrasound*. World J Gastroenterol, 2011. **17**(1): p. 28-41.
2. Ikeda, K., et al., *Effects of low-intensity pulsed ultrasound on the differentiation of C2C12 cells*. Life Sci, 2006. **79**(20): p. 1936-43.
3. Lagneaux, L., et al., *Ultrasonic low-energy treatment: a novel approach to induce apoptosis in human leukemic cells*. Exp Hematol, 2002. **30**(11): p. 1293-301.
4. Tabuchi, Y., et al., *Identification of genes responsive to low intensity pulsed ultrasound in a human leukemia cell line Molt-4*. Cancer Lett, 2007. **246**(1-2): p. 149-56.
5. Feril, L.B., Jr., et al., *Apoptosis induced by the sonomechanical effects of low intensity pulsed ultrasound in a human leukemia cell line*. Cancer Lett, 2005. **221**(2): p. 145-52.
6. Ando, H., et al., *An echo-contrast agent, Levovist, lowers the ultrasound intensity required to induce apoptosis of human leukemia cells*. Cancer Lett, 2006. **242**(1): p. 37-45.
7. Zhong, W., et al., *Sonoporation induces apoptosis and cell cycle arrest in human promyelocytic leukemia cells*. Ultrasound Med Biol, 2011. **37**(12): p. 2149-59.
8. Kishi, H., et al., *Osmotic shock induces G1 arrest through p53 phosphorylation at Ser33 by activated p38MAPK without phosphorylation at Ser15 and Ser20*. J Biol Chem, 2001. **276**(42): p. 39115-22.
9. Vahtola, E., et al., *Sirtuin1-p53, forkhead box O3a, p38 and post-infarct cardiac remodeling in the spontaneously diabetic Goto-Kakizaki rat*. Cardiovasc Diabetol, 2010. **9**: p. 5.
10. Chen, Y., et al., *IL-17A stimulates the production of inflammatory mediators via Erk1/2, p38 MAPK, PI3K/Akt, and NF-kappaB pathways in ARPE-19 cells*. Mol Vis, 2011. **17**: p. 3072-7.

MedViz Conference 2013

Notes:

MedViz Conference 2013

Notes: