

Radiomics Model for Mycetoma Grains Classification from Histopathological Microscopic Images Using Partial Least Squares Discriminant Analysis (PLS-DA)

Hyam Omar Ali, Romain Abraham, Guillaume Desoubeaux, Ahmed H Fahal,

Clovis Tauber

► To cite this version:

Hyam Omar Ali, Romain Abraham, Guillaume Desoubeaux, Ahmed H Fahal, Clovis Tauber. Radiomics Model for Mycetoma Grains Classification from Histopathological Microscopic Images Using Partial Least Squares Discriminant Analysis (PLS-DA). 2021. hal-03247688

HAL Id: hal-03247688 https://hal.science/hal-03247688

Preprint submitted on 3 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Radiomics Model for Mycetoma Grains Classification from Histopathological Microscopic Images Using Partial Least Squares Discriminant Analysis (PLS-DA)

Hyam Omar Ali^{1,2,3,4}, Romain Abraham⁴, Guillaume Desoubeaux^{5,6}, Ahmed H.Fahal², Clovis Tauber³,

- 1 Faculty of Mathematical Sciences, University of Khartoum, Sudan
- 2 The Mycetoma Research Center, University of Khartoum, Khartoum, Sudan
- **3** INSERM U1253-iBrain, Tours University, Tours, France
- 4 Institute Denis Poisson (IDP), University of Orleans, Orleans, France
- 5 Parasitology and mycology Department, Bretonneau Hospital, Tours, France
- 6 INSERM U1100-Team 3, Tours University, Tours, France

Abstract

Mycetoma is a chronic granulomatous inflammatory disease that causes severe deformities, disabilities, with many impact on patients and family, particularly in advanced disease stages or when treatment fails. The therapeutic disease strategy heavily relies on the identification of the causative organism and the corresponding classification of the disease as eumycetoma or actinomycetoma. Various diagnostic tools are used for mycetoma differential diagnosis. Histopathology is considered to be an efficient, cost and time-effective tool for mycetoma diagnosis in endemic areas. While histology is currently, the most used diagnostic tool, it requires well-trained pathologists, and that lacks in most rural areas where mycetoma is endemic. In this communication, we present a computational method to effectively differentiate between eumycetoma and actinomycetoma from the grains features in histopathological microscopic images that is based on Radiomics and Partial Least Squares Discrimination Analysis (PLS-DA). In this work, the data were collected from the Mycetoma Research Center of Khartoum, and the proposed approach achieved mycetoma types identification with an accuracy of 91.8% and 0.836 Matthew's Correlation Coefficient (MCC). This computational tool could be of great benefit in rural areas with limited access to specialised clinical centres.

Key words:

Mycetoma, Grains, Radiomics, PLS-DA, Histopathology, Image analysis.

Author summary

Mycetoma is a badly neglected tropical disease that commonly affects poor communities in rural areas. It is classified into actinomycetoma and eumycetoma depending on the causative organisms. Several diagnostic tools are used for the diagnosis of mycetoma causative agents, that include cytology, histopathology, culture, and molecular technique. However, the latter tool requires well-equipped centres that are not available in rural endemic regions. Therefore, cytology, culture, and histopathology are more commonly in use. The histopathological technique is more accurate than cytological one and faster than grains culture. Since the histological technique is operator-dependent, we introduced a novel computational method that has significant discrimination power for mycetoma types using grains features in histopathological microscopic tissue images. We believe that our method will have a valuable impact on mycetoma patients' diagnosis and management.

Introduction

Mycetoma is a WHO recognised neglected tropical disease that was included in the WHO/ 2 NTD list in 2016. It is a chronic granulomatous inflammatory disease. It is reported 3 worldwide but endemic in tropical and sub-tropical areas. Majority of cases occur in 4 the "Mycetoma belt" stretching between the latitudes of 15° South and 30° North. The 5 belt countries are Sudan, Somalia, Senegal, India, Yemen, Mexico, Venezuela, Colombia, 6 Argentina [1,2]. Sudan, India and Mexico reported the greatest number of cases [3,4]. 7 The most susceptible group of mycetoma infection is young adults in remote rural areas. Mycetoma largely affects field labourers, agriculturalists and herdsmen [1, 4, 5]. The lower extremity and hands are the frequently infected sites comparing to the other body 10 sites [2–4]. The infection might spread to involve the deep structures and bone resulting 11 in destruction, deformity, loss of function, and occasionally mortality [5–7]. 12 Mycetoma is an inflammatory, painless, and slowly progressive disease caused by certain 13

types of bacteria (Actinomycetoma) or fungi (Eumycetoma). While the mode of myce-14 toma transmission is still unknown [8], the literature suggests that causative organisms 15 are present in the soil, thorns, or animal dunk and can enter the subcutaneous tissue 16 through minor trauma [2, 8]. Mycetoma is characterised by subcutaneous mass with 17 multiple sinuses discharge grains containing colonies of the causative organism, as shown 18 in Fig1. These grains are considered as a unique characteristic of the disease [4, 6, 7, 9]. 19 Identification of causative organisms plays a significant role in the treatment of mycetoma, 20 which requires prolonged administration of anti-fungal or antibiotics drugs depending 21 on the mycetoma type [1,4,5,10,11]. Incorrect diagnosis of mycetoma can have serious 22 consequences on the patient and the disease prognosis and outcome. Currently, there are 23 many efforts aiming at establishing an early identification of the causative organism 12 24



Fig 1. Massive foot Actinomycetoma showing subcutaneous mass with multiple sinuses and discharge.

- 25 Several diagnostic tools are used to identify causative organisms and to ascertain
- $_{\rm 26}~$ the disease extension alone the tissue planes. These tools include; imaging, cytology,
- ²⁷ histological, culture, and molecular techniques [10, 12]. The imaging techniques such
- ²⁸ as X-ray, CT and MRI, define the extent of disease [13–16], while the other tools are
- $_{\rm 29}$ $\,$ employed to recognise the causative organisms. Grain culture, cytological and histological
- ³⁰ are commonly used in endemic areas [17].
- 31 Grain culture is a core tool for organisms' identification, but it is time-consuming and
- ³² requires expert microbiologists to obtain accurate results. Also, this method is vulnerable

to false-positive results due to contamination. Although molecular techniques provide 33 authenticated results, it is expensive and cannot be afforded by the majority of patients, 34 and require well-equipped infrastructures which are not available in most of the mycetoma 35 endemic area [6,9]. On the other hand, cytological and histopathological techniques 36 are simple, rapid, cheap methods and commonly used in rural areas where most of the 37 affected populations located [2, 6, 9]. However, false-negative results are common in 38 cytology because fine-needle aspiration for cytology (FNAC) is blindly performed, and it 39 is possible to miss grain pockets in the tissues. A recent comprehensive study conducted 40 at the Mycetoma Research Center (MRC) showed that histological technique is more 41 accurate than the cytological one in organism identification [18]. 42 The histopathological method can only differentiate between mycetoma fungal and 43 bacterial types conditionally by the availability of grains in tissue sections [9]; this because 44 the tissue reactions are similar in both types of mycetoma and to other non-specific 45 46 chronic granuloma diseases [6]. This discrimination mainly relies on the knowledge and experience of pathologists on the microscopic appearance of the organism [6, 10, 18-20]. 47 Due to the neglect of mycetoma and its high prevalence in the tropical regions, especially 48 rural areas, it is rare to find well-trained pathologists with adequate experience in 49 mycetoma diagnosis. To tackle this, the present work was conducted to provide an 50 open-source tool for objective identification of the mycetoma causative organisms. This 51 work introduces a new computational mycetoma differential diagnostic method based on 52 histopathological image analysis. 53 The proposed method seeks at identifying causative organisms of mycetoma with an 54 efficiency that is comparable to expert-based diagnosis in specialised clinical centres. 55 We developed a quantitative method for the discrimination of the mycetoma types from 56 grains properties in histopathological microscopic images. In the proposed approach, 57 grains in microscopic images are first characterised through Radiomics features [21–24] 58 and then classified using Partial Least Square Discriminant Analysis (PLS-DA) [25–27]. 59 The radiomics are used to extract a large number of quantitative features, which vary 60 from classical features such as first-order statistics to advanced ones that involve texture 61 and spatial characteristic of the grains. Once the features of the grain are calculated, we 62 estimated a PLS-DA model to identify new variables, which are a linear combination 63 of the original features that can be used to discriminate the grains based on their 64 causative organism. The proposed method can be implemented as a robust, reliable and 65 user-friendly software for the analysis of histopathological images, providing a solution 66 for medical needs for mycetoma diagnosis. 67

⁶⁸ Materials and methods

⁶⁹ Samples and Images Acquisition

Two sets of data were included in this study. The main set of images acquired following 70 a unique protocol that was used to train and validate the model. The secondary set, 71 composed of images acquired in various conditions, to evaluate the robustness of the 72 proposed approach. The main data set included 55 patients with confirmed different 73 mycetoma types. Surgical biopsies were obtained from patients with various mycetoma 74 types, duration and clinical presentations seen at the MRC or from the field surveys in 75 Sudan after written informed consent. Nine patients had biopsies devoid of any grains 76 and were excluded from the study. 77 The patients were randomly selected among patients seen at the MRC during the last five 78 years to ensure homogeneity and accuracy of the diagnosis. There were 31 patients with 79

 $_{\rm 80}$ $\,$ eumycetoma and 24 patients with actinomycetoma. The collected surgical biopsies were

⁸¹ fixed in 10% formal solutions, followed by paraffin-embedded tissue blocks preparation.

- Rotary microtome was used to acquire 2-3 sections with $(3-5)\mu$ thickness. All sections
- were stained with Hematoxylin and Eosin stain (H&E) according to standard routine
- ⁸⁴ laboratory procedures at Bretonneau Hospital (CHRU) through Tissue-Tek Prisma
- ⁸⁵ instrument.
- $_{\tt 86}$ $\,$ The main data set contained a total of 327 tissue microscopic images from the 55 patients
- ⁸⁷ which were used for model training and validation. The number of grains that could
- ⁸⁸ be used for each patient's biopsy varied between one and six. Images were captured in
- ⁸⁹ RGB colour space with Nikon Eclipse 80i microscope by the conditions given in Table 1
- ⁹⁰ and labelled with consideration for the patient ID in order to avoid statistical bias.

ParameterValueKnob 5/10Brightness controlND8 OnND32 OnField diaphragmHighest level

Table 1. Microscopic Acquisition Conditions

Magnification

Dimension and Quality

Colour Field diaphragm knob

Filter

NCB11 Filter

⁹¹ The secondary set of data was composed of 14 actinomycetoma and 14 eumycetoma

10X

 $\frac{800 \times 600}{\text{Enhance and white auto}}$

Highest level

6

Off

- ⁹² photomicrographs. This data set has diverse acquisition parameters and staining method.
- ⁹³ The H&E staining process was performed manually. Olympus microscope was used for
- ₉₄ images capturing with 10X magnification, while lighting and tuning conditions were not
- ⁹⁵ unified. This secondary data set covered several sources of technical variability that can
- ⁹⁶ limit the performance of predictive models [28].

⁹⁷ Preprocessing and Features Extraction

- ⁹⁸ Grains were manually segmented from the tissue photomicrograph using ImageJ software.
- Images were converted into weighted grey images before features extraction in order
- to limit the colour influence from the staining process [29]. Fig.2 shows illustrative eumycetoma and actinomycetoma microscopic images in the first and second column, respectively, along with the segmented grains.
- Radiomics features were composed of 102 variables divided into nine shape descriptors, 18 first-order statistics, and 75 texture features. Shape features were extracted from the grains masks while all other features were based on grey intensity values at the pixel level in microscopic images classes. Features were standardised by auto-scaling and used as an input to build the predictive model. PyRadiomics package version 2.2.0 was used for feature extractions [30].

¹⁰⁹ Modelling and Analysis

A PLS-DA model was adopted for grains classification. It is a supervised classification method that combines the properties of the Partial Least Square (PLS) regression model with a classification technique. For PLS-DA modelling, the features set X of images is analysed. The classes membership is translated into a dummy column vector Y by values of "1" and "-1" that indicates if a sample is eumycetoma (FM) and actinomycetoma (BM). Each row in $X_{m \times n}$ represents the different extracted features (m) of one individual sample, where n indicates number of samples.



Fig 2. Image Preprocessing. In the first row, sections stained with H&E shows mycetoma grains, the second row exhibits the conversion into grey images, and the last one demonstrates the segmentation of grains. (A, C, E): eumycetoma, and (B, D, F): actinomycetoma

The procedure of the proposed PLS-DA model used is given in pseudocode(1) [25-27]. 117 The source of variability was modelled by Latent Variables (LVs) which are linear 118 combinations of the extracted features in X. The maximum variation which secures 119 from X is determined by the weight vector w. The whole set of features X was utilised 120 for grains classification due to the ability of the PLS-DA model in reducing the impact 121 of the irrelevant features. Hence, the loading vectors (p) and (q) are the coefficients 122 assigned to features in their linear combination with various magnitude based on the 123 importance of features, so loading vectors indicate the influence of each feature on each 124 LV. Similarly, X-Score (t) represents the coordinates of samples in the LVs projection. 125 Each LV generates a variation which sums up to the total of variation secured by the 126 other LVs. The residual variation which has not been estimated by the current LV 127 is updated as a new features set. Eventually, the matrix B provides the regression 128 coefficients which describe the relationship between mycetoma grains features X and 129

 $_{130}$ mycetoma classes Y.

Algorithm 1 PLS-DA Model Construction.

Input: Features Set (X), dummy variable (Y), and number of LVs (A)Output: Regression Matrix (B)for *i* in *A* do 1. Weight Vector: w = X'Y. 2. X-Score: $t = \frac{Xw}{\sqrt{\Sigma w^2}}$. 3. X-Loading: $p = \frac{t'X}{\sqrt{\Sigma t^2}}$. 4. Y-Loading: $q = \frac{Y't}{\sqrt{\Sigma t^2}}$. 5. Regression Coefficient: $b_i = w(pw)^{-1}q$. 6. Residual of *X*: $res_x = X - tp$. 7. Residual of *Y*: $res_y = Y - tq$. 8. $X = res_x$. 9. $Y = res_y$. end for Regression Matrix *B*: $B = \{b_1, b_2, \cdots, b_A\}$.

For the prediction purpose, the model utilises the regression coefficients B and features set X of unknown samples to predict whether they are FM or BM as follows:

$Y_{pred} = XB$

Since the PLS-DA model is inherited from the PLS model, the estimated Y_{pred} is never 131 an integer with an exact membership (i.e 1 or -1). Several decision rules can be used to 132 convert the predicted values into their essential classes [25]. In this study, the threshold 133 for classes separation is calculated based on Bayes's theory and used to define perfect 134 class membership. The Bayesian threshold calculation assumes the predicted values of 135 both classes fit into a Gaussian distribution. This gives the probability of any sample 136 belonging to class FM/BM from its predicted value Y_{pred} . The estimated threshold 137 value is selected at the point where the number of false positives and false negatives is 138 minimised [31-34]. 139 The images of the different patients were randomly split into training/validation with 140

¹⁴⁰ The images of the different patients were randomly split into training/valuation with ¹⁴¹ 70/30 proportions. A summary of the data set is presented in Table 2.

Table 2. Data Split.

	\mathbf{FM}	BM	Total
Training	131	98	229
Validation	63	35	98
Total	194	133	327

All the analysis was performed using MATLAB software version R2017b and PLS
 Toolbox software version 88 from Eigenvector Technologies [35].

¹⁴⁴ Quantitative evaluation

The proposed model was assessed in two different ways according to recommended practices [25, 36].

First, we performed Cross-Validation (CV) as an internal validation method to assess 147 the complexity of the model by determining the optimum number of LVs. This allows 148 to evaluate the complexity of the model considering the predictive ability of the model 149 itself [25, 32]. The proposed model was trained with venetian blinds 10 folds-Cross-150 Validation (10 CV), and the minimum CV classification error was considered to select 151 the LVs number [32, 37–39]. In our experiments, the smallest error was associated with 152 6 LVs, Fig.3. However, using the high number of latent variables can often be associated 153 with overfitting. Meaning that the model would have a good performance on the learning 154 data, but would fail on other data sets. To prevent overfitting and to have a stronger 155 generalisation capacity, we opted for a lesser number of variables and considered three 156 latent variables. The difference of CV error between using 6 LVs (0.0556) and 3 LVs 157 (0.07281) was small in the training data, and the model behaved better on testing data. 158



Fig 3. Classification Error: The plot shows the error made by the model according to the involved number of LV.

For further assessment of the model robustness, we employed a permutation test to 159 determine the signification of the relation between grains features set and predicted 160 classes. We conducted a permutation test with 100 cycles whereby the mycetoma classes 161 of the images are randomly shuffled while maintaining the features set unchanged in each 162 cycle and building the model with the same parameters as the original one [25, 40, 41]. 163 Furthermore, it is important to evaluate the effect of samples associated with grains 164 features on the model prediction ability. The residual Q and the Hotelling T^2 values 165 are calculated for this purpose. T^2 value measures the variation in each sample within 166 the PLS-DA model. A larger value indicates a greater influence of the sample on the 167 model. Q demonstrates the goodness of samples fit the model. This allows the detection 168 of outliers, that can be removed from the learning sample set [32, 33]. 169 Variable Importance in Projection (VIP) score was analysed to understand the importance 170

of each feature in the PLS-DA model and how strongly they contribute to the classification.

¹⁷² Usually, it is considered that VIP scores have a threshold value of 1, meaning that the

features which score greater than 1 in the model are significant for the prediction ability [?].

¹⁷⁵ Finally, the performance of the model was evaluated through sensitivity, specificity, area

under the curve (AUC), accuracy, and Matthew's Correlation Coefficient (MCC).

177 **Results**

 $_{178}$ $\,$ The data shown in Fig.4 depict the scores plot and projection of samples with regards to

 $_{179}$ $\,$ the three LVs used for predication model. LVs represent the combinations of the grains

¹⁸⁰ features that best discriminate between eumycetoma and actinomycetoma. The plot

demonstrates a separation for the discriminated classes in the estimated feature space,
where two loose clusters can be identified for the mycetoma classes.



Fig 4. Samples projections: 3D projection of mycetoma images for the three latent variables (LV1, LV2, and LV3)

To detect the outliers, the plot of hotelling T^2 versus Q at 95% confidence levels was used, Fig.5. While a few samples were found slightly higher than the confidence limits for the mycetoma classes, a single FM sample was indicated as a clear outlier with high value for both Q and T^2 . A visual analysis led to consider that the sample exhibited similar properties compared to the other images of the data set, and quantitative analysis showed that the features of the sample were in line with the other samples' features which led us to maintain it within the data set.



Fig 5. Evaluation of Outliers. (A):Residuals vs Hotelling, (B):Outlier FM sample.

Analysing VIPs score of grain features with a threshold value equals to 1 qualified more than 40% of features to be significant for differential diagnosis, Fig.6. Therefore, we

¹⁹² consider 1.2 thresholds to highlight the most important features.

¹⁹³ In our experiments, the role of shape features was trivial, while texture features were ¹⁹⁴ dominant. We observed that most of the significant features were related to variance,

¹⁹⁵ entropy or complexity of grains. The peak of VIPs measures the joint distribution of low

¹⁹⁶ grey value and spatial connectivity between a pixel and its neighbours within the small ¹⁹⁷ surround. This feature is an indicator for the homogeneity of textures and the tendency

¹⁹⁸ for closer blocks to have similar spatial variation. The data shown in Fig.7 illustrates the

¹⁹⁹ great differences in shapes that can be found for similar organisms. This shape features, ²⁰⁰ which obtained a very low VIP, were not important to achieve a good classification. On

the contrary, Fig.8 represents samples of grains that score low and high VIPs for some

 $_{\rm 202}$ $\,$ of the most important features to classify the data. Grey Level Variance measures the

variance in grey-level intensity of the consecutive and adjacent blocks within the grains.
In contrast, difference variance indicates the heterogeneity of texture inside the grains.

In contrast, difference variance indicates the heterogeneity of texture inside the grains. In the three columns, the grains in the top and bottom rows are from different causative

organisms, and the corresponding features values, which were very different, helped the

²⁰⁷ discriminating process.







Fig 7. Diameter of the grains. The first and the second rows represents the grains that have the shortest and longest diameter, respectively. (A, C): actinomycetoma, and (B, D): eumycetoma.



Fig 8. Grains which score greatest and smallest score for selected VIPs. First and second rows show lowest and highest score, respectively. (A, D): Difference variance, (B, E): Small dependence low-gray-level emphasis (peak feature), and (C, F): Grey-level variance.

²⁰⁸ Predicted values for the two classes are given in Fig.9. A threshold of 0.1 is shown

as the horizontal red dashed line and apply for samples classification. This threshold
was calculated using Bayes' theorem. A sample is labelled as BM if it scores value
greater than 0.1 and FM otherwise. With such a threshold, only one sample of BM was
misclassified, while a few FM were misclassified. It indicated that while most samples
were correctly classified, a few of eumycetoma samples presented similar radiomics
features compared to actinomycetoma.



Fig 9. Predictions of the Samples: Estimated class values for samples for discrimination between eumycetoma and actinomycetoma

215 Evaluation of the model

217

²¹⁶ The evaluation results given in Table 3 illustrates that the base on our experiments,

the model is reliable and robust, with similar results obtained between training and

validation. The value of sensitivity and specificity are good estimates of the model, and

²¹⁹ they are compatible with prediction in Fig.9.

	Training set	Validation set
Sensitivity	0.948	0.971
Specificity	0.885	0.889
Accurcy	92%	91.8%
AUC	-	0.9683
Matthew's correlation	0.844	0.836

 Table 3. Estimated metrics for the model.

The estimated threshold for classes discrimination is indicated by a vertical red line in Fig 10B, with the intersection points corresponding to the sensitivity and specificity of the model. ROC curve in Fig 10A displays the sensitivity and the specificity of the model for a similar distributed set of samples with different classification thresholds. A very high AUC of 0.9683 was achieved on our data. The permutation test was used to analyse the reliability of the proposed model and the existence of a real association between grains features and their classes.

²²⁷ The permuted models and original model are compared for the goodness of fit (R^2) and

prediction (Q^2) and tested for over-fitting. The results of the permutation test are given

²²⁹ in Fig.11. The horizontal axis represents the correlation coefficient between the actual



Fig 10. Receiver Operating Characteristic curves (ROC) and threshold plots. (A):Predicted ROC, (B):Modelled Threshold

classes and the permuted classes, while the vertical axis translates the standardised sum 230 of squares that measures the deviation of each model away from the mean value and 231 is expressed in standard deviations. Small values of the sum of squares mean that the 232 model differs considerably from the mean of error, and since the permutation shuffles the 233 samples classes, there should be a great distance from the mean of error. Furthermore, 234 permuting classes might leave a small correlation between original and permuted classes; 235 hence, the right side of the plot represent the original model. The actual model scores 236 significantly higher value compared to permuted models indicating several standard 237 deviations away from the mean of error [40]. The permutation plot complies with the 238 classical validation criteria [41], and it suggests that the proposed model is not over-fitted 239 and that it is unlikely for the relation between classes and features to be random. 240



Fig 11. Permutation Test

With the objective of assessing the model on external data set, the secondary data set was used to evaluate the robustness of the model with images from new patients that have distinct acquisition parameters and slides preparation techniques. Fig.12 shows the classification of test samples. The proposed model achieves 0.892 accuracy, which is quite similar to validation results, confirming the robustness of our approach.



Fig 12. Evaluation of the model. Prediction of images with different acquisition and preparation methods from the training data set.

245

246 Discussion

Proper mycetoma management and treatment require accurate identification of mycetoma 247 causative organisms [8,20]. Currently, histopathological techniques seem to be the optimal 248 methods for identification of mycetoma organisms in terms of performance, cost and 249 time [6,9,18]. However, the ability of pathologists to discriminant between eumycetoma 250 and actinomycetoma is restricted by their knowledge about the microscopic appearance 251 of the organisms. Furthermore, some organisms look very similar to each other [6, 19, 20]. 252 Hence, the judgment is vulnerable to false results. Therefore, in this work, we pioneered 253 a computational method to differentiate between eumycetoma and actinomycetoma from 254 grains quantitative features in histopathological microscopic images. 255

In this study, we use the histopathological microscopic images from surgical biopsies from 55 randomly chosen patients from the MRC. The imaging protocol was set to uniform parameters with a digital optical microscope. As images acquisition and samples preparation method are feasible in many clinical centres that have histopathology departments, we used this technique in the present study.

The performance of our method was evaluated in terms of sensitivity, specificity, accuracy, 261 AUC and MCC. The proposed model achieved an accuracy of 91.8%, sensitivity of 0.971, 262 and specificity of 0.889. The obtained results were in line with the reported results from 263 trained expert pathologist from the MRC [18]. The 92% accuracy of expert pathologists 264 indicates that on our data set and the proposed model can be as efficient as an expert. 265 MCC metric measures the statistical accuracy taking into account the different sizes 266 of classes. It scored 0.836, which indicates a skilful model. To evaluate the robustness 267 of the model, AUC is computed by aggregating the performance of the model across 268 different thresholds. AUC of 0.968 indicated that the proposed model has excellent 269 discrimination ability. AUC and MCC values were homogeneous, which again showed 270 the ability of the model to separate the classes. Furthermore, the permutation test 271 assesses the risk of not predicating mycetoma class for a new sample. The proposed 272

273 model exhibited to be reliable and robust. Therefore, the proposed model found to be

strongly significant since the achieved results are comparable to expert classification analysis performed at the MRC. The proposed method is objective and reproducible and can reduce the need for highly specialised pathologists for diagnosis in endemic areas.

²⁷⁷ In the literature, several studies proposed size, mostly diameter, and the border of the

grains as characteristic features of the mycetoma grains [4,6,9,20]. Eumycetoma has the largest grains, while actinomycetoma grains vary from small to medium. The grains of both mycetoma types can be rounded or oval, while actinomycetoma has irregular borders, vermiform shape, or multi-lobed shape. Based on our results, there are some overlapping features between the causative organisms, and it is inconvenient to describe the grains in terms of size or shape for identification of the causative organism.

The VIPs results concluded that textural features are the most dominant and powerful 284 features for differential diagnosis. The top textural features were difference variance, 285 small dependence low-grey-level emphasis, grey-level variance, and complexity. These 286 287 features illustrate that eumycetoma tends to have a non-uniform and complex pattern within grains, and it is usually composed of connected blocks that are less homogeneous. 288 On the other hand, actinomycetoma grains are compact with a simple or regular pattern. 289 These results are of particular interest due to the reported fact, that eumycetoma grains 290 known to be harder with a coarse texture and tend to be fractured [6,9,10,20]. Hence, 291 we believe that our proposed model provides quantitative features which are quite 292 connected to the qualitative features in use by expert pathologists. In the light of the 293 aforementioned results, we can conclude that the discriminating features of mycetoma 294 types depend on the variation in texture for the adjacent regions within grains. In other 295 words, the homogeneity of textures and the tendency for closer blocks to have similar 296 spatial variation. 297

The results of this work were drawn from a well-defined set of images using uniform 298 image acquisition parameters. However, it was interesting to test the model on different 299 images from new patients with distinct acquisition parameters and slides preparation 300 techniques and to evaluate the robustness of the model. We observed a good classification 301 of this data set. It is noticeable that the misclassified samples belong to actinomycetoma 302 (Nocardia), which is uncommonly seen in Africa [3] and which was not included in 303 the training/validation data set. Certainly, the variability of the data set affects the 304 performance of the model, so this limitation should be investigated carefully in future 305 studies considering the distribution of species. 306

Considering the various subtypes of actinomycetoma and the mycetoma retrospective 307 study [18], it can be inferred that mycetoma can be classified into more than two types. 308 Thus, the proposed approach might be extended to consider more types according to the 309 individual causative organisms. Clearly, bigger sample size is needed to train the model. 310 Despite this, we tested this assumption, and the preliminary result with an accuracy of 311 80.6% is encouraging. The literature reports that smaller grains for two of the tested 312 classes (Actinomadura madurae and Actinomadura pelletierii) are similar [6,18,20]. This 313 strengthens the fact that the classification tasks for these types are challenging and the 314 accuracy we obtained was not odd. Hence, we believe that the limited number of images 315 for these classes strongly affects the performance of the extended model. Accordingly, 316 we suggest increasing the number of images of the different classes for the extended 317 model in upcoming works. 318

Tackling to develop an automated diagnostic model for mycetoma histopathological diagnosis, as another natural perspective, the proposed classification model could be integrated with a segmentation technique. The promising discrimination results fulfilled in this study are encouraging for expanding the model in the future.

In conclusion, the different effective diagnostic tools used for mycetoma diagnosis require

- expert personnel and good set up. In this work, we introduced a novel, simple, and
- $_{325}$ low-cost computational method that could be integrated into routine histopathological

diagnosis procedures. The proposed method uses radiomics in conjunction with PLS DA to effectively discriminate between actinomycetoma and eumycetoma with 91.8%
 accuracy and robustness to samples preparation techniques. This could reduce the need
 for expert pathologists in non-specialised clinical centres to perform the histological
 analysis.

331 Acknowledgments

We are grateful to Dr.Cécila Rousselot-Denis for assistance with the preparation of the histology slides, Mr.Emmanuel Edwar Siddig and Miss Omnia Babekir Abdallah who were in charge of samples collection.

335 References

- WHO: Mycetoma; Accessed January 2020. https://www.who.int/buruli/
 mycetoma/en/.
- Fahal A. Mycetoma: a thorn in the flesh. Transactions of the Royal Society of Tropical Medicine and Hygiene. 2004;98(1):3–11.
- 340 3. van de Sande WW. Global burden of human mycetoma: a systematic review and meta-analysis. PLoS neglected tropical diseases. 2013;7(11):e2550.
- Relhan V, Mahajan K, Agarwal P, Garg VK. Mycetoma: an update. Indian journal of dermatology. 2017;62(4):332.
- 5. Fahal A, Mahgoub ES, Hassan AME, Abdel-Rahman ME. Mycetoma in the
 Sudan: an update from the mycetoma research centre, University of Khartoum,
 Sudan. PLoS neglected tropical diseases. 2015;9(3):e0003679.
- Fahal A. Mycetoma—clinicopathological monograph, Khartoum University Press.
 Khartoum, Sudan. 2006.
- ³⁴⁹ 7. A F. Bailey and Love's Short Practice of Surgery;.
- 8. van de Sande WW, Fahal AH, Goodfellow M, Welsh O, Zijlstra E, et al. The
 mycetoma knowledge gap: identification of research priorities. PLoS neglected
 tropical diseases. 2014;8(3):e2667.
- Ahmed AA, van de Sande W, Fahal AH. Mycetoma laboratory diagnosis. PLoS
 neglected tropical diseases. 2017;11(8):e0005638.
- 10. Emmanuel P, Dumre SP, John S, Karbwang J, Hirayama K. Mycetoma: a
 clinical dilemma in resource limited settings. Annals of clinical microbiology and
 antimicrobials. 2018;17(1):35.
- 11. Efared B, Tahiri L, Boubacar MS, Atsam-Ebang G, Hammas N, Chbani L, et al.
 Mycetoma in a non-endemic area: a diagnostic challenge. BMC clinical pathology.
 2017;17(1):1.
- 12. van de Sande WW, Fahal AH, Goodfellow M, Welsh O, Zijlstra EE, et al. Merits
 and pitfalls of currently used diagnostic tools in mycetoma. PLoS neglected
 tropical diseases. 2014;8(7):e2918.
- 13. Abd El Bagi ME. New radiographic classification of bone involvement in pedal
 mycetoma. American Journal of Roentgenology. 2003;180(3):665–668.

366	14.	El-Bagi MEA, Fahal AH. Mycetoma revisited. Saudi Med J. 2009;30(4):529–533
367 368	15.	Fahal A, Sheik H, Homeida M, Arabi Y, Mahgoub E. Ultrasonographic imaging of mycetoma. British journal of surgery. 1997;84(8):1120–1122.
369 370 371	16.	El Shamy M, Fahal A, Shakir M, Homeida M. New MRI grading system for the diagnosis and management of mycetoma. Transactions of the Royal Society of Tropical Medicine and Hygiene. 2012;106(12):738–742.
372 373 374	17.	Alam K, Maheshwari V, Bhargava S, Jain A, Fatima U, ul Haq E. Histological diagnosis of madura foot (mycetoma): a must for definitive treatment. Journal of global infectious diseases. 2009;1(1):64.
375 376 377 378	18.	Siddig EE, Mhmoud NA, Bakhiet SM, Abdallah OB, Mekki SO, El Dawi NI, et al. The Accuracy of Histopathological and Cytopathological Techniques in the Identification of the Mycetoma Causative Agents. PLoS neglected tropical diseases. 2019;13(8):e0007056.
379 380 381	19.	van de Sande W, Fahal A, Ahmed SA, Serrano JA, Bonifaz A, Zijlstra E, et al Closing the mycetoma knowledge gap. Medical mycology. 2018;56(suppl_1):S153-S164.
382 383	20.	Siddig E, Fahal A. Histopathological Approach in Diagnosis of Mycetoma Causative Agents: A Mini Review. J Cytol Histol. 2017;8:466.
384 385	21.	Kumar V, Gu Y, Basu S, Berglund A, Eschrich SA, Schabath MB, et al. Radiomics: the process and the challenges. Magnetic resonance imaging. 2012;30(9):1234–1248
386 387	22.	Avanzo M, Stancanello J, El Naqa I. Beyond imaging: The promise of radiomics Physica Medica. 2017;38:122–139.
388 389 390	23.	Lambin P, Rios-Velazquez E, Leijenaar R, Carvalho S, Van Stiphout RG, Granton P, et al. Radiomics: extracting more information from medical images using advanced feature analysis. European journal of cancer. 2012;48(4):441–446.
391 392 393	24.	Limkin E, Sun R, Dercle L, Zacharaki E, Robert C, Reuzé S, et al. Promises and challenges for the implementation of computational medical imaging (radiomics) in oncology. Annals of Oncology. 2017;28(6):1191–1206.
394 395 396	25.	Lee LC, Liong CY, Jemain AA. Partial least squares-discriminant analysis (PLS-DA) for classification of high-dimensional (HD) data: a review of contemporary practice strategies and knowledge gaps. Analyst. 2018;143(15):3526–3539.
397 398	26.	Brereton RG, Lloyd GR. Partial least squares discriminant analysis: taking the magic away. Journal of Chemometrics. 2014;28(4):213–225.
399 400 401	27.	Kvalheim OM. Interpretation of partial least squares regression models by means of target projection and selectivity ratio plots. Journal of Chemometrics. 2010;24(7-8):496–504.
402 403 404	28.	McCann MT, Ozolek JA, Castro CA, Parvin B, Kovacevic J. Automated histology analysis: Opportunities for signal processing. IEEE Signal Processing Magazine 2014;32(1):78–87.
405 406	29.	Komura D, Ishikawa S. Machine learning methods for histopathological image analysis. Computational and structural biotechnology journal. 2018;16:34–42.

407 408 409	30.	Van Griethuysen JJ, Fedorov A, Parmar C, Hosny A, Aucoin N, Narayan V, et al. Computational radiomics system to decode the radiographic phenotype. Cancer research. 2017;77(21):e104–e107.
410 411 412 413	31.	dos Santos VHJ, Ramos AS, Pires JP, Engelmann PdM, Lourega RV, Ketzer JM, et al. Discriminant analysis of biodiesel fuel blends based on combined data from Fourier Transform Infrared Spectroscopy and stable carbon isotope analysis. Chemometrics and Intelligent Laboratory Systems. 2017;161:70–78.
414 415	32.	Ballabio D, Consonni V. Classification tools in chemistry. Part 1: linear models. PLS-DA. Analytical Methods. 2013;5(16):3790–3798.
416 417	33.	Ramos PM, Ruisánchez I. Data fusion and dual-domain classification analysis of pigments studied in works of art. Analytica chimica acta. 2006;558(1-2):274–282.
418 419 420	34.	Pérez NF, Ferré J, Boqué R. Calculation of the reliability of classification in discriminant partial least-squares binary classification. Chemometrics and Intelligent Laboratory Systems. 2009;95(2):122–128.
421 422	35.	Wise BM, Gallagher N, Bro R, Shaver J, Windig W, Koch RS. PLS Toolbox 4.0; 2007.
423 424	36.	Esbensen KH, Geladi P. Principles of proper validation: use and abuse of re- sampling for validation. Journal of Chemometrics. 2010;24(3-4):168–187.
425 426 427	37.	Villa JE, Quiñones NR, Fantinatti-Garboggini F, Poppi RJ. Fast discrimination of bacteria using a filter paper–based SERS platform and PLS-DA with uncertainty estimation. Analytical and bioanalytical chemistry. 2019;411(3):705–713.
428 429 430	38.	Pereira LS, Lisboa FL, Neto JC, Valladão FN, Sena MM. Screening method for rapid classification of psychoactive substances in illicit tablets using mid infrared spectroscopy and PLS-DA. Forensic science international. 2018;288:227–235.
431 432 433	39.	Botelho BG, Reis N, Oliveira LS, Sena MM. Development and analytical validation of a screening method for simultaneous detection of five adulterants in raw milk using mid-infrared spectroscopy and PLS-DA. Food chemistry. 2015;181:31–37.
434 435 436	40.	Ahn JK, Kim S, Kim J, Hwang J, Kim KH, Cha HS. A comparative metabolomic evaluation of Behcet's disease with arthritis and seronegative arthritis using synovial fluid. PloS one. $2015;10(8):e0135856$.
437 438	41.	Musharraf SG, Siddiqui AJ, Shamsi T, Choudhary MI, Rahman Au. Serum metabonomics of acute leukemia using nuclear magnetic resonance spectroscopy.

439 Scientific reports. 2016;6:30693.