

Supplementary Tables. Univariate analysis of cellulitis risk factors for patients with secondary and primary lymphedema.

Primary lower limb lymphedema

Characteristic	OR (95% CI)	P value
Male sex	2.8 (1.77-4.46)	<10 ⁻³
Body mass index (kg/m ²)	1.05 (1.02-1.08)	.001
Lymphedema-onset-to-1 st -first consultation interval	1.002 (1.000-1.003)	.004
Age at lymphedema onset	0.986 (0.975-0.996)	.009
Diabetes	1.34 (0.38-4.50)	.64

Secondary upper limb lymphedema

Characteristic	OR (95% CI)	P value
Body mass index (kg/m ²)	1.000 (0.977-1.023)	.98
Diabetes	1.30 (0.71-2.36)	.38
Age at lymphedema onset	0.97 (0.96-0.99)	<10 ⁻⁵
Lymphedema-onset-to-1 st -first consultation interval	1.006 (1.004-1.008)	<10 ⁻⁵
Unilateral	0.29 (0.08-0.88)	.04
Hormonotherapy	0.76 (0.59-0.98)	.03
Radiotherapy	1.94 (1.16-3.35)	.01
Tumorectomy (vs mastectomy)	1.29 (1.01-166)	.04
Chemotherapy	0.88 (0.64-1.22)	.45
Lymph-node excision	1.18 (0.63-2.29)	.62

Secondary lower limb lymphedema

Characteristic	OR (95% CI)	P value
Diabetes	0.65 (0.09-2.73)	.59
Body mass index (kg/m ²)	1.02 (0.98-1.06)	.4
Lymphedema-onset-to-1 st -first consultation interval	1.005 (1.002-1.008)	.003
Chemotherapy	0.91 (0.55-1.49)	.71
Lymph-node excision	1.83 (0.89-4.05)	.11
Unilateral	0.95 (0.56-1.64)	.84
Age at lymphedema onset	0.97 (0.95-0.99)	.002
Male sex	1.25 (0.68-2.27)	.46
Radiotherapy	0.94 (0.57-1.56)	.8