



HAL
open science

Multi-parameters rational solutions to the mKdV equation

Pierre Gaillard

► **To cite this version:**

| Pierre Gaillard. Multi-parameters rational solutions to the mKdV equation. 2021. hal-03251014

HAL Id: hal-03251014

<https://hal.science/hal-03251014v1>

Preprint submitted on 5 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Multi-parameters rational solutions to the mKdV equation

Pierre Gaillard,
Université de Bourgogne,
Institut de mathématiques de Bourgogne,
9 avenue Alain Savary BP 47870
21078 Dijon Cedex, France :
E-mail : Pierre.Gaillard@u-bourgogne.fr

Abstract

N -order solutions to the modified Korteweg-de Vries (mKdV) equation are given in terms of a quotient of two wronskians of order N depending on $2N$ real parameters. When one of these parameters goes to 0, we succeed to get for each positive integer N , rational solutions as a quotient of polynomials in x and t depending on $2N$ real parameters. We construct explicit expressions of these rational solutions for orders $N = 1$ until $N = 6$.

Key Words : mKdV equation, wronskians, rational solutions.

PACS numbers :

33Q55, 37K10, 47.10A-, 47.35.Fg, 47.54.Bd

1 Introduction

We consider the modified Korteweg-de Vries (mKdV) equation

$$u_t - 6u^2u_x + u_{xxx} = 0, \quad (1)$$

with $u_t = \partial_t u$, $u_x = \partial_x u$ and $u_{xxx} = \partial_x^3 u$.

The mKdV equation has many applications in various fields as in the study of waves propagating in plasma [3], the dynamics of traffic flow [4] and fluid mechanics [5]. In particular, it is used in nonlinear optics as for example to model supercontinuum generation in optical fibres [1] or to describe pulses consisting of a few optical cycles [2].

Various methods have been used to construct solutions to the mKdV equation. Hirota [6] constructed the exact soliton for the mKdV equation in 1972. In the same year, Tanaka [7] was the first to solve the mKdV equation by using the inverse scattering technique. Wadati using this same method succeeded to obtain

the exact N -soliton solution for the mKdV [8]. In 1976, rational solutions to the mKdV equation [9] were derived by Ono by using Bäcklund transformation. A limiting procedure based on bilinear results gave other type of solutions [10] in 1978. In 1983, rational solutions to the mKdV equation were determined by recurrence relations in [11] by Tanaka. Solitary wave solutions and cnoidal wave solutions were given in 1994 in [12]. Some classes of periodic solutions of mKdV have been given in [13] in 2004.

More recently, in 2012, solutions to the mKdV equation has been constructed via bilinear Bäcklund transformation in [14] and rational solutions in terms of Wronskians were obtained. In 2016, periodic solutions and rational solution of first and second order were presented in [15] by using a Darboux transformation were constructed.

Here, Darboux transformation is used to construct different type of solutions. Representations of solutions in terms of wronskians of order N depending on $2N$ real parameters are given, using trigonometric or hyperbolic functions. Rational solutions are obtained in performing a passage to the limit when one of these parameters goes to 0. So rational solutions as a quotient of polynomials in x and t , depending on $2N$ parameters are constructed. We give explicit solutions in the simplest cases $N = 1, 2, 3$ and some particular rational solutions for $N = 1$ until 6.

2 N-order solutions to the mKdV equation in terms of wronskians

2.1 N-order solutions in terms of wronskians of hyperbolic sine functions

We consider the mKdV equation

$$u_t - 6u^2u_x + u_{xxx} = 0.$$

We recall that the wronskian of order N of the functions f_1, \dots, f_N is the determinant denoted $W(f_1, \dots, f_N)$, defined by $\det(\partial_x^{i-1} f_j)_{1 \leq i \leq N, 1 \leq j \leq N}$, ∂_x^i being the partial derivative of order i with respect to x and $\partial_x^0 f_j$ being the function f_j .

We consider a_j, b_j arbitrary real numbers $1 \leq j \leq N$. We have the following result :

Theorem 2.1 *Let f_j be the functions defined by*

$$f_j(x, t) = \sinh \left(\frac{1}{2} a_j x - \frac{1}{2} a_j^3 t + b_j \right), \quad \text{for } 1 \leq i \leq N, \quad (2)$$

then the function u defined by

$$u(x, t) = \partial_x \ln \left(\frac{W(\partial_x(f_1), \dots, \partial_x(f_N))}{W(f_1, \dots, f_N)} \right) \quad (3)$$

is a solution to the mKdV equation (1) depending on $2N$ real parameters $a_j, b_j, 1 \leq j \leq N$.

2.2 Some examples of solutions to the mKdV equation with sine hyperbolic generating functions

In the following we only give the solutions of order 1, 2 and 3 in the case of generating hyperbolic sinus functions.

Solution of order 1

Proposition 2.1 *The function u defined by*

$$u(x, t) = \frac{a_1}{\sinh(-a_1 x + a_1^3 t - 2b_1)}$$

is a solution to the mKdV equation (1) with a_1, b_1 arbitrarily real parameters.

Solution of order 2

Proposition 2.2 *The function u defined by*

$$u(x, t) = \frac{n(x, t)}{d(x, t)}, \quad (4)$$

with

$$n(x, t) = -\frac{1}{2}(a_1^2 - a_2^2)[a_2 \sinh(-a_1 x + a_1^3 t - 2b_1) - a_1 \sinh(-a_2 x + a_2^3 t - 2b_2)]$$

and,

$$d(x, t) = 2(-\sinh(-1/2 a_1 x + 1/2 a_1^3 t - b_1) \cosh(-1/2 a_2 x + 1/2 a_2^3 t - b_2) a_2 + \sinh(-1/2 a_2 x + 1/2 a_2^3 t - b_2) \cosh(-1/2 a_1 x + 1/2 a_1^3 t - b_1) a_1) (\cosh(-1/2 a_1 x + 1/2 a_1^3 t - b_1) \sinh(-1/2 a_2 x + 1/2 a_2^3 t - b_2) a_2 - \cosh(-1/2 a_2 x + 1/2 a_2^3 t - b_2) \sinh(-1/2 a_1 x + 1/2 a_1^3 t - b_1) a_1)$$

is a solution to the mKdV equation (1) with a_1, a_2, b_1, b_2 arbitrarily real parameters.

Solution of order 3

In this case of order 3, we only present solution with $a_1 = 1, a_2 = 2, a_3 = 3, b_1 = 0, b_2 = 0, b_3 = 0$ to shorten the paper.

Proposition 2.3 *The function u defined by*

$$u(x, t) = \frac{n(x, t)}{d(x, t)}, \quad (5)$$

with

$$n(x, t) = -150 (\cosh(-1/2 x + 1/2 t))^2 - 384 (\cosh(-x + 4 t - 2))^2 - 54 (\cosh(-3/2 x +$$

$\frac{27}{2}t-3)^2+300 \sinh(-1/2x+1/2t) \cosh(-3/2x+\frac{27}{2}t-3) \cosh(-1/2x+1/2t) \sinh(-3/2x+\frac{27}{2}t-3)-312 \cosh(-3/2x+\frac{27}{2}t-3) \sinh(-x+4t-2) \sinh(-3/2x+\frac{27}{2}t-3) \cosh(-x+4t-2)-180 (\cosh(-1/2x+1/2t))^2(\cosh(-3/2x+\frac{27}{2}t-3))^2+480 (\cosh(-x+4t-2))^2(\cosh(-1/2x+1/2t))^2+288 (\cosh(-x+4t-2))^2(\cosh(-3/2x+\frac{27}{2}t-3))^2-600 \sinh(-1/2x+1/2t) \cosh(-x+4t-2) \cosh(-1/2x+1/2t) \sinh(-x+4t-2)$

and,

$d(x, t) = 2(16 \sinh(-1/2x+1/2t) \cosh(-x+4t-2) \sinh(-3/2x+\frac{27}{2}t-3)-9 \sinh(-1/2x+1/2t) \cosh(-3/2x+\frac{27}{2}t-3) \sinh(-x+4t-2)-5 \cosh(-1/2x+1/2t) \sinh(-x+4t-2) \sinh(-3/2x+\frac{27}{2}t-3))(-16 \cosh(-1/2x+1/2t) \sinh(-x+4t-2) \cosh(-3/2x+\frac{27}{2}t-3)+9 \cosh(-1/2x+1/2t) \sinh(-3/2x+\frac{27}{2}t-3) \cosh(-x+4t-2)+5 \sinh(-1/2x+1/2t) \cosh(-x+4t-2) \cosh(-3/2x+\frac{27}{2}t-3))$

is a solution to the mKdV equation (1).

2.3 Other types of solutions

We obtain similar results with other types of generating functions whose proofs are identical.

2.3.1 Solutions with hyperbolic cosine generating functions

Theorem 2.2 Let h_j, h be the following functions

$$h_j(x, t) = \cosh\left(\frac{1}{2}a_jx - \frac{1}{2}a_j^3t + b_j\right), \quad \text{for } 1 \leq i \leq N, \quad (6)$$

then the function u defined by

$$u(x, t) = \partial_x \ln\left(\frac{W(\partial_x(h_1), \dots, \partial_x(h_N))}{W(h_1, \dots, h_N)}\right) \quad (7)$$

is a solution to the mKdV equation (1) with a_j, b_j $1 \leq j \leq N$ arbitrarily real parameters.

2.3.2 Solutions with trigonometric generating functions

Theorem 2.3 Let g_j be the following functions

$$g_j(x, t) = \cos\left(\frac{1}{2}a_jx + \frac{1}{2}a_j^3t + b_j\right), \quad \text{for } 1 \leq i \leq N, \quad (8)$$

then the function u defined by

$$u(x, t) = \partial_x \ln\left(\frac{W(\partial_x(g_1), \dots, \partial_x(g_N))}{W(g_1, \dots, g_N)}\right) \quad (9)$$

is a solution to the mKdV equation (1) with a_j, b_j $1 \leq j \leq N$ arbitrarily real parameters.

Theorem 2.4 Let k_j be the following functions

$$k_j(x, t) = \sin \left(\frac{1}{2} a_j x + \frac{1}{2} a_j^3 t + b_j \right), \quad \text{for } 1 \leq i \leq N, \quad (10)$$

then the function u defined by

$$u(x, t) = \partial_x \ln \left(\frac{W(\partial_x(k_1), \dots, \partial_x(k_N))}{W(k_1, \dots, k_N)} \right) \quad (11)$$

is a solution to the mKdV equation (1) with a_j, b_j $1 \leq j \leq N$ arbitrarily real parameters.

3 Rational solutions to the mKdV equation

In the following, we replace all parameters a_j and b_j , $1 \leq j \leq N$ by $\hat{a}_j = \sum_{k=1}^N a_k(je)^{2k-1}$ and $\hat{b}_j = \sum_{k=1}^N b_k(je)^{2k-1}$ with e an arbitrary real parameter. We realize this change to obtain rational solutions to the mKdV equation (1); for this, we perform a limit when the parameter e tends to 0.

We get the following result :

Theorem 3.1 Let ψ_j be the functions

$$\psi_j(x, t, e) = \sinh \left(\frac{1}{2} \sum_{k=1}^N a_k(je)^{2k-1} x - \frac{1}{2} \left(\sum_{k=1}^N a_k(je)^{2k-1} \right)^3 t + \sum_{k=1}^N b_k(je)^{2k-1} \right),$$

for $1 \leq j \leq N$,

then the function u defined by

$$u(x, t) = \lim_{e \rightarrow 0} \partial_x \ln \left(\frac{W(\partial_x(\psi_1), \dots, \partial_x(\psi_N))}{W(\psi_1, \dots, \psi_N)} \right) \quad (12)$$

is a rational solution to the mKdV equation (1).

We have similar results with generating sine or trigonometric functions.

We can also give the expression of the rational solutions of the mKdV without the presence of a limit.

We get the following result :

Theorem 3.2 Let ψ, φ_j be the functions

$$\psi(x, t, e) = \sinh \left(\frac{1}{2} \left(\sum_{k=1}^N a_k e^{2k-1} \right) x - \frac{1}{2} \left(\sum_{k=1}^N a_k e^{2k-1} \right)^3 t + \sum_{k=1}^N b_k e^{2k-1} \right),$$

$$\varphi_j(x, t) = \frac{\partial^{2j-1} \psi(x, t, 0)}{\partial_{2j-1} e}, \quad \text{for } 1 \leq j \leq N,$$

then the function v defined by

$$v(x, t) = \partial_x \ln \left(\frac{W(\partial_x(\varphi_1), \dots, \partial_x(\varphi_N))}{W(\varphi_1, \dots, \varphi_N)} \right) \quad (13)$$

is a rational solution to the mKdV equation (1) depending on $2N$ parameters $a_j, b_j, 1 \leq j \leq N$.

We give some examples of rational solutions in the following.

It must be pointed out that these resulting rational solutions are singular.

These results are consequences of the previous result.

3.1 First order rational solutions

We have the following result at order $N = 1$:

Proposition 3.1 *The function v defined by*

$$v(x, t) = \frac{-a_1}{a_1 x + 2b_1}, \quad (14)$$

is a rational solution to the mKdV equation (1) with a_1, b_1 , arbitrarily real parameters.




Figure 1. Solution of order 1 to (1), on the left $a_1 = 1, b_1 = 1$; on the right $a_1 = 1, b_1 = 10^2$.

3.2 Second order rational solutions

Proposition 3.2 *The function v defined by*

$$v(x, t) = \frac{n(x, t)}{d(x, t)}, \quad (15)$$

with

$$n(x, t) = -2a_1^5 x^3 - 12a_1^4 b_1 x^2 - 24a_1^3 b_1^2 x + 2a_1(12a_2 b_1 + 6a_1^4 t - 12b_2 a_1 - 8a_1 b_1^3),$$

and,

$$d(x, t) = a_1^5 x^4 + 8a_1^4 b_1 x^3 + 24a_1^3 b_1^2 x^2 + (24a_1^2 b_1^3 + a_1(24a_2 b_1 + 12a_1^4 t - 24b_2 a_1 + 8a_1 b_1^3))x + 2b_1(24a_2 b_1 + 12a_1^4 t - 24b_2 a_1 + 8a_1 b_1^3)$$

is a rational solution to the mKdV equation (1) dependant on 4 real parameters a_1, a_2, b_1, b_2 .




Figure 2. Solution of order 2 to (1), on the left $a_1 = 1, a_2 = 2, b_1 = 10, b_2 = 2$; in the center $a_1 = 1, a_2 = 2, b_1 = 1, b_2 = 2$; on the right $a_1 = 1, a_2 = 2, b_1 = 10, b_2 = 0$.

3.3 Rational solutions of order three

In this case, we only give the rational solution with some fixed parameters in reason of the length of the general solution. We choose $a_j = b_j = j$ for $1 \leq j \leq 3$. We get the following rational solution given by :

Proposition 3.3 *The function v defined by*

$$v(x, t) = \frac{n(x, t)}{d(x, t)}, \quad (16)$$

with

$$n(x, t) = 3x^8 + 48x^7 + 336x^6 + (1344 + 72t)x^5 + (720t + 3360)x^4 + (2880t + 5376)x^3 + (5760t + 5376 + 4320t^2)x^2 + (5760t + 3072 + 17280t^2)x + 768 + 2304t + 17280t^2$$

and,

$$d(x, t) = -x^9 - 18x^8 - 144x^7 + (-672 - 72t)x^6 + (-2016 - 864t)x^5 + (-4032 - 4320t)x^4 + (-4096 - 9120t + (8 + 12t)(-60t - 160) + 720t^2)x^3 + ((8 + 12t)(-240 - 360t) - 2688 - 11520t + 4320t^2)x^2 + ((8 + 12t)(-192 - 720t) - 768 - 5760t + 8640t^2)x + (8 + 12t)(-64 - 480t + 720t^2)$$

is a rational solution to the mKdV equation (1).




Figure 3. Solution of order 3 to (1), on the left $a_1 = 1, a_2 = 2, a_3 = 3, b_1 = 0, b_2 = 0, b_3 = 0$; in the center $a_1 = 1, a_2 = 2, a_3 = 3, b_1 = 10, b_2 = 2, b_3 = 3$; on the right $a_1 = 1, a_2 = 2, a_3 = 3, b_1 = 10, b_2 = 0, b_3 = 0$.

3.4 Rational solutions of order four

With the same choices of parameters $a_j = b_j = j$ for $1 \leq j \leq 4$, we get the following rational solution given by :

Proposition 3.4 *The function v defined by*

$$v(x, t) = \frac{n(x, t)}{d(x, t)}, \quad (17)$$

with

$$\begin{aligned} n(x, t) = & 4x^{15} + 120x^{14} + 1680x^{13} + (14560 + 600t)x^{12} + (14400t + 87360)x^{11} + \\ & (158400t + 384384)x^{10} + (36000t^2 + 1056000t + 1281280)x^9 + (3294720 + 648000t^2 + \\ & 4752000t)x^8 + (15206400t + 5184000t^2 + 6589440)x^7 + (-2419200t^3 + 10250240 + \\ & 24192000t^2 + 35481600t)x^6 + (12300288 + 72576000t^2 - 29030400t^3 + 60825600t)x^5 + \\ & (-145152000t^3 + 145152000t^2 + 11182080 + 76032000t)x^4 + (67584000t - 36288000t^4 + \\ & 193536000t^2 + 7454720 - 387072000t^3)x^3 + (-217728000t^4 + 40550400t + 3440640 + \\ & 165888000t^2 - 580608000t^3)x^2 + (983040 - 435456000t^4 + 82944000t^2 - 464486400t^3 + \\ & 14745600t)x + 131072 + 2457600t - 217728000t^5 + 18432000t^2 - 154828800t^3 - 290304000t^4 \end{aligned}$$

and,

$$\begin{aligned} d(x, t) = & -x^{16} - 32x^{15} - 480x^{14} + (-4480 - 240t)x^{13} + (-29120 - 6240t)x^{12} + \\ & (-139776 - 74880t)x^{11} + (-358912 + 720t^2 - 462720t + (-160 - 60t)(180t + 960))x^{10} + \\ & ((-160 - 60t)(2520t + 3360) - 1752000t - 696320 + (-360t - 240)(180t + 960) + \\ & 14400t^2)x^9 + (-1013760 + 129600t^2 - 4440960t + (-160 - 60t)(8064 + 15120t) + \\ & (-192 - 720t)(180t + 960) + (-360t - 240)(2520t + 3360))x^8 + (-302400t^3 - 7580160t - \\ & 1064960 + (-192 - 720t)(2520t + 3360) + (-360t - 240)(8064 + 15120t) + (-64 + 720t^2 - \\ & 480t)(180t + 960) + (-160 - 60t)(13440 + 50400t))x^7 + ((-64 + 720t^2 - 480t)(2520t + \\ & 3360) - 4233600t^3 - 8248320t - 753664 + (-160 - 60t)(100800t + 15360) + (-192 - \\ & 720t)(8064 + 15120t) + (-360t - 240)(13440 + 50400t))x^6 + (-25401600t^3 - 5114880t - \end{aligned}$$

$319488 + (-192 - 720t)(13440 + 50400t) + (-360t - 240)(100800t + 15360) + (-64 + 720t^2 - 480t)(8064 + 15120t) + (-160 - 60t)(120960t + 11520))x^5 + ((-64 + 720t^2 - 480t)(13440 + 50400t) + (-192 - 720t)(100800t + 15360) + (-160 - 60t)(302400t^3 + 80640t + 5120) - 61440 - 1382400t - 36288000t^3 + (-360t - 240)(120960t + 11520))x^4 + ((-360t - 240)(302400t^3 + 80640t + 5120) + (-64 + 720t^2 - 480t)(100800t + 15360) + (-192 - 720t)(120960t + 11520) + (-160 - 60t)(1024 + 23040t + 604800t^3))x^3 + ((-64 + 720t^2 - 480t)(120960t + 11520) + (-192 - 720t)(302400t^3 + 80640t + 5120) + (-360t - 240)(1024 + 23040t + 604800t^3))x^2 + ((-64 + 720t^2 - 480t)(302400t^3 + 80640t + 5120) + (-192 - 720t)(1024 + 23040t + 604800t^3))x + (-64 + 720t^2 - 480t)(1024 + 23040t + 604800t^3)$

is a rational solution to the mKdV equation (1).




Figure 4. Solution of order 4 to (1), on the left $a_1 = 1, a_2 = 2, a_3 = 3, a_4 = 4, b_1 = 0, b_2 = 0, b_3 = 0, b_4 = 10$; in the center $a_1 = 1, a_2 = 2, a_3 = 3, a_4 = 4, b_1 = 10, b_2 = 0, b_3 = 0, b_4 = 0$; on the right $a_1 = 1, a_2 = 2, a_3 = 3, a_4 = 4, b_1 = 0, b_2 = 0, b_3 = 0, b_4 = 4$.

3.5 Rational solutions of orders five and six

We choose parameters to get the shorter expression of the solution. For this, we consider $a_j = j$ and $b_j = 0$, for $1 \leq j \leq N$, we get the following rational solution given by :

Order 5

Proposition 3.5 The function v defined by

$$v(x, t) = \frac{n(x, t)}{d(x, t)}, \quad (18)$$

with

$$n(x, t) = 5x^{24} + 2280tx^{21} + 352800t^2x^{18} + 4838400t^3x^{15} + 2794176000t^4x^{12} + 259096320000t^5x^9 + 5120962560000t^6x^6 - 153628876800000t^7x^3 + 460886630400000t^8$$

and,

$d(x, t) = -x^{25} - 600tx^{22} - 100800t^2x^{19} - 6955200t^3x^{16} - 254016000t^4x^{13} + 39626496000t^5x^{10} - 365783040000t^6x^7 + 76814438400000t^7x^4 + 460886630400000t^8x$
is a rational solution to the mKdV equation (1).

Order 6

Proposition 3.6 *The function v defined by*

$$v(x, t) = \frac{n(x, t)}{d(x, t)}, \quad (19)$$

with

$$\begin{aligned}
n(x, t) = & -6x^{35} - 6300tx^{32} - 2419200t^2x^{29} - 381024000t^3x^{26} - 50295168000t^4x^{23} + \\
& 192036096000t^5x^{20} + 1371137725440000t^6x^{17} + 129970029772800000t^7x^{14} - 116143430860800000t^8x^{11} + \\
& 31939443486720000000t^9x^8 + 2146330602307584000000t^{10}x^5 + 128779836138455040000000t^{11}x^2
\end{aligned}$$

and,

$$\begin{aligned}
d(x, t) = & x^{36} + 1260tx^{33} + 544320t^2x^{30} + 110073600t^3x^{27} + 11430720000t^4x^{24} + \\
& 329204736000t^5x^{21} + 119062379520000t^6x^{18} + 7189831434240000t^7x^{15} + 1393721170329600000t^8x^{12} - \\
& 59620294508544000000t^9x^9 - 4292661204615168000000t^{10}x^6 + 128779836138455040000000t^{12}
\end{aligned}$$

is a rational solution to the mKdV equation (1).

For solutions of order 5 and 6 depending respectively on 10 and 12 parameters, the structure of the polynomials of the numerators and denominators is the following.

For order 5, the numerator contains 8370 terms and the denominator 10279; the degree of the numerator in x is 24, in t is 8; the degree of the denominator in x is 25, in t is 8.

For order 6, the numerator contains 100409 terms and the denominator 119620; the degree of the numerator in x is 35, in t is 11; the degree of the denominator in x is 36, in t is 12.

Unlike other equations like the nonlinear Schrödinger equation, we do not see a specific structure of polynomials defining these solutions.

4 Conclusion

Two types of representations of solutions to the mKdV equation have been given. First, we have constructed solutions as a quotient of a wronskian of order N by a wronskian of order N depending on $2N$ real parameters. Then rational solutions to the mKdV equation depending on $2N$ real parameters, performing a passage to the limit when one parameter goes to 0.

We can mention some other recent works about this equation. In [18], first and second-order rational solutions are given as limiting cases of periodic solutions, the second one depending on one real parameter. In [19] the reduced Maxwell-Bloch (RMB) equations are considered and N th-order rational solutions containing several free parameters are presented, in particular explicit

expressions of these solutions from first to second order. Explicit periodic and rational solutions of first and second order are given in [20], and some typical nonlinear wave patterns are shown. The first four exact rational solutions of the set of rational solutions of the mKdV equation are presented in [21]. Multiple periodic solutions of the mKdV equation are given in [22] and in particular first to third-order rational solutions.

All the solutions presented in this article are different from those proposed in previous references and are new. So we get an infinite hierarchy of multiparametric families of rational solutions to the mKdV equation as a quotient of a polynomials in x and t depending on $2N$ real parameters.

It would be relevant to better understand the structure of polynomials defining the rational solutions of this equation.

References

- [1] H. Leblond, P. Grelu, D. Mihalache, Models for supercontinuum generation beyond the slowly-varying-envelope approximation, *Phys. Rev. A*, V. 90, 053816-1-9, 2014
- [2] H. Leblond, D. Mihalache, Few-optical-cycle solitons: Modified Korteweg-de Vries sine-Gordon equation versus other non-slowly-varying-envelope-approximation models, *Phys. Rev. A*, V. 79, 063835-1-7, 2009
- [3] A.H. Khater, O.H. El-Kalaawy, D.K. Callebaut, Bäcklund transformations and exact solutions for Alfvén solitons in a relativistic electronpositron plasma, *Phys. Scripta*, V. 58, 545-548, 1998
- [4] Z.P. Li, Y.C. Liu, Analysis of stability and density waves of traffic flow model in an ITS environment, *Eur. Phys. J. B*, V. 53, 367-374, 2006
- [5] M.A. Helal, Soliton solution of some nonlinear partial differential equations and its applications in fluid mechanics, *Chaos, Solitons and Fractals*, V. 13, 1917-1929, 2002
- [6] R. Hirota, Exact envelope-soliton solutions of a nonlinear wave equation, *Jour. Of Math. Phys.*, V. 14, N. 7, 805-809, 1973
- [7] S. Tanaka, Modified Korteweg. de Vries Equation and Scattering Theory, *Proc. Japan Acad.*, V. 48, 466-469, 1972
- [8] M. Wadati, The Exact Solution of the Modified Korteweg-de Vries Equation, *J. Phys. Soc. Jpn.*, V. 32, 1681-1681, 1972
- [9] Y. Ono, Algebraic Soliton of the Modified Korteweg-de Vries Equation, *Jour. Phys. Soc. Japan*, V. 41, N. 5, 1817-1818, 1976
- [10] M.J. Ablowitz, J. Satsuma, Solitons and rational solutions of nonlinear evolution equations, *Jour. Math. Phys*, V. 19, N. 10, 2180-2186, 1978

- [11] Y. Kametaka, On Rational Similarity Solutions of KdV and m.KdV Equations, Proc. Japan. Acad., V. 59, 407-409, 1983
- [12] S.Y. Lou, L.L. Chen, Solitary wave solutions and cnoidal wave solutions to the combined KdV and mKdV equation, Math Meth. Appl. Sci., V. 17, N.5, 339-347, 1994
- [13] P.G. Kevrekidis, A. Khare, A. Saxena, G. Herring, On some classes of mKdV periodic solutions, J. Phys. A, V. 37, 1095910965, (2004)
- [14] Y.Y. Sun, D.J. Zhang, Rational Solutions with non-zero asymptotics of the modified Korteweg-de Vries Equation, Commun. Theor. Phys., V. 57, N.6, 923-929, 2012
- [15] A. Chowdury, A. Ankiewicz, N. Akhmediev, Periodic and rational solutions of modified Korteweg-de Vries equation, Eur. Phys. J. D, V. 70, N. 104, 1-7, 2016
- [16] V.B. Matveev and M.A. Salle, Darboux transformations and solitons, Series in Nonlinear Dynamics, Springer-Verlag, Berlin, 1991
- [17] M.Yu. Kulikov and G.M. Fraiman, Darboux transformations and solitons, Preprint of the Institute of Applied Physics, Russian Academy of Sciences, N. 404, Nizhni Novgorod, 1996
- [18] A. Chowdury, A. Ankiewicz, N. Akhmediev, Periodic and rational solutions of mKdV equation, Eur. Phys. J. D, V. 70, N. 104, 1-7, 2016
- [19] J. Wei, X. Wang, X.G. Geng, Periodic and rational solutions of the reduced Maxwell Bloch equations, Commun. Nonlinear Sci. Numer. Simulat., V. 59, 1-14, 2018
- [20] X. Wang, J. Zhang, L. Wang, Conservation laws periodic and rational solutions for an extended mKdV equation, Nonlinear Dyn., V. 92, 1507-1516, 2018
- [21] A. Ankiewicz, N. Akhmediev, Rogue wave-type solutions of the mKdV equation and their relation to known NLSE rogue wave solutions, Nonlinear Dyn., V. 91, 1931-1038, 2018
- [22] Zhaquilao, Z.C. Zhou, S.X. Li, Nonsingular complexiton and rational solutions for the mKdV equation and KdV equation, Pacific Jour. Of Appl. Math., V. 6, N. 3, 43-58, 2014
- [23] P. Gaillard, V.B. Matveev, Wronskian addition formula and its applications, Max-Planck-Institut für Mathematik, MPI 02-31, V. 161, 2002
- [24] P. Gaillard, A new family of deformations of Darboux-Pöschl-Teller potentials, Lett. Math. Phys., V. 68, 77-90, 2004

- [25] P. Gaillard, V.B. Matveev, New formulas for the eigenfunctions of the two-particle Calogero-Moser system, *Lett. Math. Phys.*, V. **89**, 1-12, 2009
- [26] P. Gaillard, V.B. Matveev, Wronskian and Casorai determinant representations for Darboux-Pöschl-Teller potentials and their difference extensions, *J. Phys A : Math. Theor.*, V. **42**, 404409-1-16, 2009
- [27] P. Gaillard, Families of quasi-rational solutions of the NLS equation and multi-rogue waves, *J. Phys. A : Meth. Theor.*, V. **44**, 435204-1-15, 2011
- [28] P. Gaillard, Wronskian representation of solutions of the NLS equation and higher Peregrine breathers, *J. Math. Sciences : Adv. Appl.*, V. **13**, N. 2, 71-153, 2012
- [29] P. Gaillard, Degenerate determinant representation of solution of the NLS equation, higher Peregrine breathers and multi-rogue waves, *J. Math. Phys.*, V. **54**, 013504-1-32, 2013
- [30] P. Gaillard, Wronskian representation of solutions of NLS equation and seventh order rogue waves, *J. Mod. Phys.*, V. **4**, N. 4, 246-266, 2013
- [31] P. Gaillard, V.B. Matveev, Wronskian addition formula and Darboux-Pöschl-Teller potentials, *J. Math.*, V. **2013**, ID 645752, 1-10, 2013
- [32] P. Gaillard, Two parameters deformations of ninth Peregrine breather solution of the NLS equation and multi rogue waves, *J. Math.*, V. **2013**, 1-111, 2013
- [33] P. Gaillard, Two-parameters determinant representation of seventh order rogue waves solutions of the NLS equation, *J. Theor. Appl. Phys.*, V. **7**, N. **45**, 1-6, 2013
- [34] P. Gaillard, Six-parameters deformations of fourth order Peregrine breather solutions of the NLS equation, *J. Math. Phys.*, V. **54**, 073519-1-22, 2013
- [35] P. Gaillard, Deformations of third order Peregrine breather solutions of the NLS equation with four parameters, *Phys. Rev. E*, V. **88**, 042903-1-9, 2013
- [36] P. Gaillard, Ten parameters deformations of the sixth order Peregrine breather solutions of the NLS equation, *Phys. Scripta*, V. **89**, 015004-1-7, 2014
- [37] P. Gaillard, The fifth order Peregrine breather and its eight-parameters deformations solutions of the NLS equation, *Commun. Theor. Phys.*, V. **61**, 365-369, 2014
- [38] P. Gaillard, Higher order Peregrine breathers, their deformations and multi-rogue waves, *J. Of Phys. : Conf. Ser.*, V. **482**, 012016-1-7, 2014

- [39] P. Gaillard, M. Gastineau, Eighteen parameter deformations of the Peregrine breather of order ten solutions of the NLS equation, *Int. J. Mod. Phys. C*, V. **26**, N. 2, 1550016-1-14, 2014
- [40] P. Gaillard, Two parameters wronskian representation of solutions of nonlinear Schrödinger equation, eight Peregrine breather and multi-rogue waves, *J. Math. Phys.*, V. **5**, 093506-1-12, 2014
- [41] P. Gaillard, Hierarchy of solutions to the NLS equation and multi-rogue waves, *J. Phys. : Conf. Ser.*, V. **574**, 012031-1-5, 2015
- [42] P. Gaillard, Tenth Peregrine breather solution of the NLS, *Ann. Phys.*, V. **355**, 293-298, 2015
- [43] P. Gaillard, M. Gastineau, The Peregrine breather of order nine and its deformations with sixteen parameters solutions of the NLS equation *Phys. Lett. A*, V. **379**, 1309-1313, 2015
- [44] P. Gaillard, Other $2N-2$ parameters solutions to the NLS equation and $2N+1$ highest amplitude of the modulus of the N -th order AP breather, *J. Phys. A: Math. Theor.*, V. **48**, 145203-1-23, 2015
- [45] P. Gaillard, Multi-parametric deformations of the Peregrine breather of order N solutions to the NLS equation and multi-rogue waves, *Adv. Res.*, V. **4**, N. **5**, 346-364, 2015
- [46] P. Gaillard, Higher order Peregrine breathers solutions to the NLS equation, *Jour. Phys. : Conf. Ser.*, V. **633**, 012106-1-6, 2016
- [47] P. Gaillard, M. Gastineau Patterns of deformations of Peregrine breather of order 3 and 4, solutions to the NLS equation with multi-parameters, *Journal of Theoretical and Applied Physics*, V. **10**,1-7, 2016
- [48] P. Gaillard, M. Gastineau Twenty parameters families of solutions to the NLS equation and the eleventh Peregrine breather, *Commun. Theor. Phys*, V. **65**, N. **2**, 136-144, 2016
- [49] P. Gaillard, Rational solutions to the KPI equation and multi rogue waves, *Annals Of Physics*, V. **367**, 1-5, 2016
- [50] P. Gaillard, M. Gastineau Twenty two parameters deformations of the twelfth Peregrine breather solutions to the NLS equation, *Adv. Res.*, V. **10**, 83-89, 2016
- [51] P. Gaillard, Towards a classification of the quasi rational solutions to the NLS equation, *Theor. And Math. Phys.*, V. **189**, N. **1**, 1440-1449, 2016
- [52] P. Gaillard, Fredholm and Wronskian representations of solutions to the KPI equation and multi-rogue waves, *Jour. of Math. Phys.*, V. **57**, 063505-1-13, doi: 10.1063/1.4953383, 2016

- [53] P. Gaillard, M. Gastineau Families of deformations of the thirteenth Peregrine breather solutions to the NLS equation depending on twenty four parameters, *Jour. Of Bas. And Appl. Res. Int.*, V. **21**, N. 3, 130-139, 2017
- [54] P. Gaillard, From Fredholm and Wronskian representations to rational solutions to the KPI equation depending on $2N^2$ parameters, *Int. Jour. of Appl. Sci. And Math.*, V. **4**, N. 3, 60-70, 2017
- [55] P. Gaillard, Families of Rational Solutions of Order 5 to the KPI Equation depending on 8 Parameters, *New Hor. in Math. Phys.*, V. **1**, N. 1, 26-31, 2017
- [56] P. Gaillard, 6-th order rational solutions to the KPI Equation depending on 10 parameters, *Jour. Of Bas. And Appl. Res. Int.*, V. **21**, N. 2, 92-98, 2017
- [57] P. Gaillard, N -Order rational solutions to the Johnson equation depending on $2N - 2$ parameters, *Int. Jour. of Adv. Res. in Phys. Sci.*, V. **4**, N. 9, 19-37, 2017
- [58] P. Gaillard, Families of rational solutions to the KPI equation of order 7 depending on 12 parameters, *Int. Jour. of Adv. Res. in Phys. Sci.*, V. **4**, N. 11, 24-30, 2017
- [59] P. Gaillard, Rational solutions to the Johnson equation and rogue waves, *Int. Jour. of Inn. In Sci. and Math.*, V. **6**, N. 1, 14-19, 2018
- [60] P. Gaillard, Multiparametric families of solutions of the KPI equation, the structure of their rational representations and multi-rogue waves, *Theo. And Mat. Phys.*, V. **196**, N. 2, 1174-1199, 2018
- [61] P. Gaillard, The Johnson Equation, Fredholm and Wronskian representations of solutions and the case of order three, *Adv. In Math. Phys.*, V. **2018**, 1-18, 2018
- [62] P. Gaillard, Families of Solutions of Order 5 to the Johnson Equation Depending on 8 Parameters, *NHIMP*, V. **2**, N. 4, 53-61, 2018
- [63] P. Gaillard, Multiparametric families of solutions to the Johnson equation, *J. Phys. : Conf. Series*, V. **1141**, 012102-1-10, 2018
- [64] P. Gaillard, Rational solutions to the Boussinesq equation, *Fund. Jour. Of Math. And Appl.*, V. , 109-112, 2019
- [65] P. Gaillard, Differential relations for the solutions to the NLS equation and their different representations, *Comm. In Adv. Math. Sci.*, V. **2**, N. 4, 1-4, 2019,
- [66] P. Gaillard, Multi-Parametric Families of Solutions of Order N to the Boussinesq and KP Equations and the Degenerate Rational Case, *UJMA*, V. **3**, N. 2, 44-52, 2020,