



HAL
open science

Note on Artin's Conjecture on Primitive Roots

Sankar Sitaraman

► To cite this version:

| Sankar Sitaraman. Note on Artin's Conjecture on Primitive Roots. 2021. hal-03251183v1

HAL Id: hal-03251183

<https://hal.science/hal-03251183v1>

Preprint submitted on 6 Jun 2021 (v1), last revised 20 Dec 2021 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Note on Artin's Conjecture on Primitive Roots

Sankar Sitaraman

Howard University, Washington, DC

To Gerald, Ralph, Sundar and Susan

Abstract

E. Artin conjectured that any integer $a > 1$ which is not a perfect square is a primitive root modulo p for infinitely many primes p . Let $f_a(p)$ be the multiplicative order of the non-square integer a modulo the prime p . M. R. Murty and S. Srinivasan [10] showed that if $\sum_{p < x} \frac{1}{f_a(p)} = O(x^{1/4})$ then Artin's conjecture is true for a . We relate the Murty-Srinivasan condition to sums involving the cyclotomic periods from the subfields of $\mathbb{Q}(e^{2\pi i/p})$ corresponding to the subgroups $\langle a \rangle \subseteq \mathbb{F}_p^*$.

1. Introduction

Let $a > 1$ be an integer which is not a perfect square and let $f_a(p)$ be the multiplicative order of a modulo a given prime p . For such a the Murty-Srinivasan [10] condition $\sum_{p < x} \frac{1}{f_a(p)} = O(x^{1/4})$ applies, and one could try to evaluate this sum by looking at the average value of $\omega(\Phi_j(a))$ where $\Phi_j(x)$ is the j -th cyclotomic polynomial and $\omega(n)$ is the number of distinct primes dividing n . We were led to this idea by our thinking of $f_a(p)$ as the lengths of the orbits of the points on the unit circle S^1 under the "Frobenius" map $\sigma_a : x \rightarrow x^a$. This meant looking at the torsion of the group of roots of unity. Note that for a given prime p , $f_a(p)$ is the order of this map acting on the multiplicative subgroup generated by $\zeta_p = e^{\frac{2\pi i}{p}}$. It is well known that if $f_a(p) = j$ then $p \mid \Phi_j(a)$ where $\Phi_j(x)$ is the j -th cyclotomic polynomial. See, for instance, [15], Chapter 2.

Since the $\Phi_j(a)$ divide $a^j - 1$ and we are only concerned with $p < x$ we actually use $\sum \omega_x(a^j - 1)$ where $\omega_x(m)$ for integral m is the number of primes upto x that divide it. The argument to approach the Murty-Srinivasan condition using the average value of $\omega(\Phi_j(a))$ is also in Murty-Wong [11], Section 5, albeit implicitly. There an inequality for a sum that contains $S(x) = \sum_{p < x} \frac{1}{f_a(p)}$ is used to give heuristics for the largest prime factor of $a^j - 1$, $j \leq x$.

We show how the sum $\sum_{j \leq x} \omega_x(a^j - 1)$ can be related to cyclotomic periods after expressing it in terms of finite Fourier series (exponential sums) and give some bounds for these sums, from which one could easily get bounds for $\sum_{p < x} \frac{1}{f_a(p)}$.

More precisely, our main result is the following:

Theorem 1 For a and $f_a(p)$ as above, with $\eta_{i,p}$ the cyclotomic periods from the subfield of $\mathbb{Q}(e^{2\pi i/p})$ corresponding to the subgroup $\langle a \rangle \subseteq \mathbb{F}_p^*$, we get the following bounds for any $u \leq x$ where x is a fixed positive integer:

$$A : \sum_{j \leq u} \omega_x(a^j - 1) \leq O(u \log \log x) + u \sum_{p \leq x} \frac{\sum_{i=1}^{\frac{p-1}{f_a(p)}} |\eta_{i,p}|}{p}.$$

$$B : \sum_{j \leq u} \omega_x(a^j - 1) \leq O(u \log \log x) + 2\pi u \sum_{p \leq x} \frac{\left(\sum_{k=1}^{p-2} \left| \sum_{m=1}^k \eta_{h_m,p} \right| \right)}{p^2}$$

where $\eta_{h_m,p} = \sum_{j=1}^{f_a(p)} \zeta_p^{h_m a^j}$ with $h_m \equiv m f_a(p) \pmod{p}$, $m \in [1, p-1]$.

Much progress has been made on Artin's conjecture by Hooley [8], Gupta-Murty [5], Heath-Brown [7] and others. Heath-Brown proved that Artin's conjecture is true for one of 2, 3 and 5.

Our initial motivation was to come up with a Dirichlet series that would help in obtaining information about the asymptotic behavior of invariants of cyclotomic fields such as class numbers. We hope to provide such a framework that would enable us to calculate the asymptotics of $f_a(p)$ for a given a more precisely, as indicated in Sitaraman [12]. For instance, Artin conjectured that $f_a(p) = p - 1$ with a certain non-zero density C (equal to 0.374 approximately) known now as the Artin constant.

The author is grateful to Niranjana Ramachandran and Larry Washington for their hospitality and conversations during his visit to the University of Maryland, College Park, in 2014-15. He would also like to thank Francois Ramarosan and Dani Szpruch for helpful conversations.

2. From the Murty-Srinivasan criterion to Cyclotomic Periods.

2a. Reduction to estimate of $\sum \omega_x(a^j - 1)$

Let $S(x) = \sum_{p < x} \frac{1}{f_a(p)}$. Let $N(j)$ to be the number of primes $p < x$ with a of order j modulo p . (i.e, $f_a(p) = j$). Then we have $S(x) \leq \sum_{j < x} \frac{N(j)}{j}$. Since as mentioned before $f_a(p) = j \implies p | \Phi_j(a)$ we get

$$S(x) = \sum_{j < x} \frac{N(j)}{j} \leq \sum_{j < x} \frac{\omega_x(\Phi_j(a))}{j}.$$

Here $\omega_x(m)$ is a “truncated” version of the usual $\omega(m)$ that denotes the number of distinct prime factors of m in that it only counts primes less than x .

Since $\Phi_j(a)$ is a factor of $a^j - 1$ we have $\sum_{j < x} \frac{\omega_x(\Phi_j(a))}{j} \leq \sum_{j < x} \frac{\omega_x(a^j - 1)}{j}$. Since it is easier to deal with $a^j - 1$ we consider the sum $\sum_{j < x} \frac{\omega_x(a^j - 1)}{j} = W(x)$.

Initially we tried to use sieve theoretic techniques to deal with $W(x)$. H. Halberstam [6] obtained an Erdős-Kac type result for $\omega(f(m))$ where $f(x) \in \mathbb{Z}[x]$ is an irreducible polynomial. Namely he showed that $\sum_{m \leq n} \frac{\omega(f(m)) - A(n)}{n\sqrt{A(n)}}$ is asymptotically normally distributed,

where $A(n) = \sum_{p \leq n} \frac{r(p)}{p}$ with $r(p)$ being the number of solutions to $f(k) \equiv 0 \pmod p$ with $0 \leq k < p$. Halberstam needs $r(p)$ to be uniformly bounded, and that is possible because the polynomial is fixed. Here we need to average over several polynomials of varying degree. Even using a later more general version of Halberstam’s result proved by Granville-Soundararajan [4] we found it difficult to get a decent estimate for $W(x)$.

On the other hand A. T. Felix [3] has shown that on average $\sum 1/f_a(p)$ is about $\log x$. So actual bound could be much smaller than $O(x^{1/4})$. This is also the conclusion drawn in Murty-Wong.

The best bound we know for $\sum_{p < x} \frac{1}{f_a(p)}$ is $\sqrt{x}/(\log x)^{1+\delta}$ where δ is some small positive quantity, given by P. Erdős and M. R. Murty [2]. The bounds A and B, numerically at least, seem to give better estimates.

2b. Some Notation

In what follows $f_a(p)$ will be denoted simply as f_p .

$$e_p = \frac{p-1}{f_p}.$$

g is a primitive generator of \mathbb{F}_p^* and so $a \equiv g^{e_p} \pmod p$.

g_i is the least element of $\{1, 2, \dots, p-1\}$ such that $g_i \equiv g^i \pmod p$.

$\eta_{i,p} = \sum_{j=1}^{f_p} \zeta_p^{g^j g^{e_p j}} = \sum_{j=1}^{f_p} \zeta_p^{g^i a^j} = \sum_{j=1}^{f_p} \zeta_p^{g_i a^j}$ are the cyclotomic periods for $i = 1, 2, \dots, e_p$. They are contained in the subfield of $\mathbb{Q}(\zeta_p)$ corresponding to the subgroup $\langle a \rangle$.

H_p is the subgroup $\langle a \rangle$ generated by a in \mathbb{F}_p^* .

2c. Proof of bounds A and B

Using summation by parts, one could reduce the estimate of $W(x)$ to that of $D(u)$ for $u \leq x$ for positive integers u , where $D(u) = \sum_{j \leq u} \omega_x(a^j - 1)$.

Now it is easy to see that

$$D(u) = \sum_{j \leq u} \omega_x(a^j - 1) = \sum_{p \leq x} \frac{1}{p} \sum_{j \leq u} \sum_{k=1}^p \zeta_p^{k(a^j-1)}. \quad (1)$$

Bound A:

Note that, for a fixed prime p we have, on the one hand, $\sum_{j \leq u} \omega_p(a^j - 1) = \left\lfloor \frac{u}{f_p} \right\rfloor$. Here we abused the notation a little bit to make $\omega_p(m)$ equal 1 if p divides m and 0 otherwise. This will be the case regardless of whether f_p divides u or not. Moreover, if $f_p > u$, then clearly p will not divide any of the $a^j - 1, j \leq u$, and $\left\lfloor \frac{u}{f_p} \right\rfloor = 0$.

For the part of the sum $D(u)$ with $k = p$ we have $\sum_{p < x} \frac{1}{p} \sum_{j \leq u} \sum_{k=1}^p \zeta_p^{k(a^j-1)} = \sum_{p < x} \sum_{j \leq u} \frac{1}{p}$ and this is $O(u \log \log x)$.

In what follows we focus on the $k < p$ part. Also let $u \equiv u_f \pmod{f_p}$ with $0 \leq u_f < f_p$. There will be no confusion owing to the dependence on p .

$$\begin{aligned} D(u) &= O(u \log \log x) + \sum_{p \leq x} \frac{1}{p} \sum_{j=1}^u \sum_{k=1}^{p-1} \zeta_p^{(a^j-1)k} \\ &= O(u \log \log x) + \sum_{p \leq x} \frac{1}{p} \sum_{j=1}^{u_f} \sum_{k=1}^{p-1} \zeta_p^{(a^j-1)k} \\ &\quad + \sum_{p \leq x} \frac{1}{p} \left\lfloor \frac{u}{f_p} \right\rfloor \sum_{H_p g_i \in \mathbb{F}_p^*/H_p} \left(\sum_{k \in H_p g_i} \zeta_p^{-k} \left(\sum_{j=1}^{f_p} \zeta_p^{a^j k} \right) \right). \end{aligned} \quad (2)$$

In the last term of the sum in (2) above we used the facts that the sum over j is periodic with period f_p and also that the inner sum $\sum_{j=1}^{f_p} \zeta_p^{a^j k}$ is identical for k in the same coset.

In the second term of the sum above, for a fixed j we have $\sum_{k=1}^{p-1} \zeta_p^{(a^j-1)k} = -1$ if $f_p \nmid j$ which would be the case for $j \leq u_f < f_p$. So we get

$$D(u) = O(u \log \log x) + \sum_{p \leq x} \left(-\frac{u_f}{p} \right) + \sum_{p \leq x} \frac{1}{p} \left\lfloor \frac{u}{f_p} \right\rfloor \sum_{H_p g_i \in \mathbb{F}_p^*/H_p} \left(\sum_{k \in H_p g_i} \zeta_p^{-k} \eta_{i,p} \right)$$

The second term of the sum is $O(u \log \log x)$ and for the last term we bound trivially and then since each coset (i.e, each $\eta_{i,p}$) appears f_p times, we get the bound A:

$$D(u) = O(u \log \log x) + \sum_{p \leq x} \frac{1}{p} \left\lfloor \frac{u}{f_p} \right\rfloor \sum_{i=1}^{e_p} f_p |\eta_{i,p}|$$

$$\implies D(u) \leq O(u \log \log x) + u \sum_{p \leq x} \frac{\sum_{i=1}^{e_p} |\eta_{i,p}|}{p}.$$

Note: The inner sum of equation (2), namely $\sum_{H_p g_i \in \mathbb{F}_p^*/H_p} \left(\sum_{k \in H_p g_i} \zeta_p^{-k} \left(\sum_{j=1}^{f_p} \zeta_p^{a^j k} \right) \right)$, can be seen to equal $\sum_i \bar{\eta}_{i,p} \eta_{i,p} = \sum_i |\eta_{i,p}|^2$ and this is known to be equal to $p - f_p$. See, for instance, Thaine [14], p. 36. So, left as it is, this sum does not give anything new. Although this equation actually gives a formula to compute f_p in terms of the cyclotomic periods, one expects it to be hard to work with, given that f_p is hard to compute even on average.

Even though it might be bigger than original sum, the bound A (as well as bound B below) at least eliminates f_p from the estimate.

Bound B:

To get bound B we use summation by parts. Since, as seen above, the sum for $j = 1$ to u_f contributes only $O(u \log \log x)$ as does the sum for $k = p$ for each p , we simply focus on the remainder of the expression for $D(u)$. We call this part $D(u)'$.

$$D(u)' = \sum_{p \leq x} \frac{1}{p} \left\lfloor \frac{u}{f_p} \right\rfloor \sum_{H_p g_i \in \mathbb{F}_p^*/H_p} \left(\sum_{k \in H_p g_i} \zeta_p^{-k} \eta_{i,p} \right)$$

Before summing by parts, first we rearrange the sum so that it goes from $g_i = 1$ to $g_i = p - 1$.

$$D(u)' = \sum_{p \leq x} \frac{1}{p} \left\lfloor \frac{u}{f_p} \right\rfloor \left(\sum_{g_i=1}^{p-1} \zeta_p^{-g_i} \eta_{i,p} \right)$$

Now we rearrange the indices again and try to replace the g_i with $h_m \in [1, p-1]$ as m goes from 1 to $p - 1$ such that $h_{m+1} - h_m \equiv f_p \pmod{p}$. This can be done by letting $h_m \equiv m f_p \pmod{p}$. Since $p \nmid f_p$ the map $m \rightarrow h_m$ will result in a permutation of $\{1, 2, \dots, p - 1\}$.

$$\text{In particular, } \sum_{g_i} \eta_{i,p} = \sum_{i=1}^{p-1} \eta_{i,p} = \sum_{m=1}^{p-1} \eta_{h_m,p}.$$

Summing by parts,

$$D(u)' = \sum_{p \leq x} \frac{1}{p} \left\lfloor \frac{u}{f_p} \right\rfloor \left[\left(\sum_{g_i} \eta_{i,p} \right) \zeta_p^{-h_{p-1}} + \left(\sum_{k=1}^{p-2} \left(\sum_{m=1}^k \eta_{h_m,p} \right) (\zeta_p^{-h_k} - \zeta_p^{-h_{k+1}}) \right) \right]. \quad (3)$$

Now note that $|\zeta_p^{-h_k} - \zeta_p^{-h_{k+1}}| = \left| 2 \sin \left(\frac{\pi(h_{k+1} - h_k)}{p} \right) \right| \leq \frac{2\pi f_p}{p}$ because $h_{k+1} - h_k = f_p$,

for $k = 1, 2, \dots, p - 2$. Also note that, since $\sum_{i=1}^{e_p} \eta_{i,p} = -1$, and there are f_p full sets of $\eta_{i,p}$, we have $\sum_{g_i} \eta_{i,p} = -f_p$. Using these two computations, we get

$$|D(u)'| \leq \sum_{p \leq x} \frac{1}{p} \left\lfloor \frac{u}{f_p} \right\rfloor \left[f_p + \left(\sum_{k=1}^{p-2} \left| \sum_{m=1}^k \eta_{h_m, p} \right| \frac{2\pi f_p}{p} \right) \right] \leq u \sum_{p \leq x} \frac{1}{p} + 2\pi u \sum_{p \leq x} \frac{\left(\sum_{k=1}^{p-2} \left| \sum_{m=1}^k \eta_{h_m, p} \right| \right)}{p^2}$$

Adding the $O(u(\log \log x))$ terms from the sum $D(u)$ that were left out, we get bound B:

$$D(u) \leq O(u \log \log x) + 2\pi u \sum_{p \leq x} \frac{\left(\sum_{k=1}^{p-2} \left| \sum_{m=1}^k \eta_{h_m, p} \right| \right)}{p^2}$$

Due to the increased possibility of cancellations, we think bound B would be better than bound A. Numerically that seems to be the case.

2d. Estimates for Bounds A and B

Clearly, the best estimates for the bounds A and B would be obtained using estimates for the average values of $|\eta_{i,p}|$ and $\left| \sum_{m=1}^k \eta_{h_m, p} \right|$ over the cyclotomic periods as well as over the primes. Here we restrict ourselves to seeing what estimates can be obtained for a fixed prime. As such, in this section we will drop the subscript p from $\eta_{i,p}$ and $\eta_{h_m, p}$.

Bound A:

As mentioned in the Note in the previous section, we have $\sum_i \bar{\eta}_i \eta_i = \sum_i |\eta_i|^2 = p - f_p$. From this we see that all the cyclotomic periods have absolute values bounded by \sqrt{p} , and possibly the absolute values are much less than \sqrt{p} , at least on average. Indeed, we have (looking at the general term from the sum in bound A):

$$\begin{aligned} \left(\sum_i |\eta_i| \right)^2 &\leq e_p \sum_i |\eta_i|^2 \implies \frac{1}{p} \sum_i |\eta_i| \leq \frac{\sqrt{e_p}}{p} \sqrt{\sum_i |\eta_i|^2} \\ &\implies \frac{1}{p} \sum_i |\eta_i| \leq \sqrt{\frac{p-1}{f_p} \frac{\sqrt{p-f_p}}{p}} = \sqrt{\frac{1}{f_p} - \frac{1}{p}} \times \sqrt{1 - \frac{f_p}{p}}. \end{aligned} \quad (4)$$

(The first inequality above follows from the elementary fact that the average of a set of positive numbers is smaller than the square root of the average of their squares).

As expected the estimate from equation (4) does not yield a better estimate for $D(u)$ (and hence for $\sum 1/f_p$) than the existing ones.

On the other hand there are several estimates that show that $\max_i |\eta_i|$ is smaller than f_p when f_p itself is small. For instance we have

$$\max_i |\eta_i| < p^{-3\alpha/8} f_p \text{ when } f_p > p^{\frac{1}{3} + \alpha}.$$

See Bourgain [1] for a survey of such results. H. Montgomery, R. C. Vaughan and T. D. Wooley [9] conjecture that $\max_i |\eta_i| < \min\left(\sqrt{p}, C\sqrt{f_p \times \log p}\right)$, with the constant C independent of p .

Suppose $\max_i |\eta_i| < M_p$. When one uses such an estimate in bound A one gets

$$\begin{aligned} D(u) &= \sum_{j \leq u} \omega_x(a^j - 1) \leq O(u \log \log x) + u \sum_{p \leq x} \frac{\sum_{i=1}^{e_p} |\eta_{i,p}|}{p} \\ &\leq O(u \log \log x) + u \sum_{p \leq x} \frac{1}{p} \left(\frac{p-1}{f_p} (M_p) \right) \leq O(u \log \log x) + u \sum_{p \leq x} \frac{M_p}{f_p}. \end{aligned}$$

Since one key problem is that we don't know the amount of primes with f_p in a given interval $p^a \leq f_p \leq p^b$ it is difficult to average the bound A over the primes p . Even otherwise, it appears that using existing estimates for the maximum of the periods it would be difficult to get a useful estimate for $D(u)$ and hence $\sum_p 1/f_p$. One does need a good estimate for the averages of the $|\eta_i|$ as opposed to their maximum.

Bound B:

For bound B we have the following very basic estimate in the form of a trigonometric sum:

We first estimate $\left| \sum_{m=1}^k \eta_{h_m, p} \right|$. As before we omit p in the subscript and denote $h_{m,p}$ by h_m . We get:

$$\begin{aligned} \left| \sum_{m=1}^k \eta_{h_m} \right| &= \left| \sum_{m=1}^k \sum_{j=1}^{f_p} \zeta_p^{h_m a^j} \right| = \left| \sum_{m=1}^k \sum_{j=1}^{f_p} \zeta_p^{h_m a^j} \right| \\ \implies \left| \sum_{m=1}^k \eta_{h_m} \right| &= \left| \sum_{j=1}^{f_p} \sum_{m=1}^k \zeta_p^{m f_p a^j} \right| = \left| \sum_{j=1}^{f_p} \zeta_p^{f_p a^j} \sum_{m=0}^{k-1} \zeta_p^{m f_p a^j} \right| \leq \sum_{j=1}^{f_p} \left| \frac{\sin(\pi a^j f_p k / p)}{\sin(\pi a^j f_p / p)} \right|. \end{aligned}$$

Putting everything together, we get for Bound B the following estimate in terms of a trigonometric sum:

$$D(u) \leq O(u \log \log x) + 2\pi u \sum_{p \leq x} \frac{1}{p^2} \left(\sum_{k=1}^{p-2} \sum_{j=1}^{f_p} \left| \frac{\sin(\pi a^j f_p k / p)}{\sin(\pi a^j f_p / p)} \right| \right).$$

2e. Implications for Artin's Conjecture

If $D(u)$ is of order $ux^{1/4}/(\log x)$ or less then after summation by parts we get that $W(x)$ is of order $x^{1/4}$ and that would verify the Murty-Srinivasan condition and hence Artin's conjecture.

Now, when $f_p > p^{3/4}$ the $\sum 1/f_p$ is at most $O(x^{1/4}/(\log x))$. So main problem is when $f_p < p^{3/4}$.

Erdős and Murty [2] show that for most primes f_p is of order bigger than \sqrt{p} . In the bounds given here most of the contribution would come from the primes p with small f_p and for such p the approximation of $\sin(\pi f_p/p)$ by $\pi f_p/p$ would be better.

Finally, the full sums of bounds A and B seem to be at least as big as $ux^{1/4}$, based on numerical evidence.

Nevertheless, one could try to use the expression for $D(u)$ in terms of cyclotomic periods (equation (2) and (3)) without taking absolute values as we did here (to get the bounds A and B). If it is possible to sum over the primes first before summing over the k 's a la Voronoi in his estimate for the Dirichlet divisor problem (cf. for instance, Tenenbaum [13] I.6.4) one could make progress on the conjecture.

In this direction would the deep estimates of exponential sums over finite fields developed by Bourgain, Konyagin, et al be useful ? See for instance Bourgain's survey article [1]. Perhaps the same methods would yield good estimates for average sums for the cyclotomic periods.

References

- [1] J. Bourgain, Sum-Product theorems and applications, *Festschrift in honor of the sixtieth birthday of Melvyn B. Nathanson, Springer*,(2010), XI.
- [2] P. Erdős and M. R. Murty, On the order of a (mod p), Number theory (Ottawa, ON, 1996),*CRM Proc. Lecture Notes 19, Amer. Math. Soc., Providence, RI* (1999), 87 - 97.
- [3] A. T. Felix, Generalizing The Titchmarsh Divisor Problem, *International Journal of Number Theory*, **08**, (2012) No. 03, pp. 613-629.
- [4] A. Granville and K. Soundararajan, Sieving and the Erdős-Kac Theorem, *Equidistribution in Number Theory, An Introduction, Granville and Rudnick, editors*, Proceedings of the NATO Advanced Study Institute on Equidistribution in Number Theory, Montreal, Canada, 11–22 July 2005, pp. 15-27.
- [5] R. Gupta and M. Ram Murty, A remark on Artin's conjecture, *Inventiones Math.*, **78** (1984) 127-130.
- [6] H. Halberstam, On the distribution of additive number theoretic functions (II and III) *J. Lond. Math. Soc.*, **31** (1956), Part 1, No. 121, 1-27.
- [7] D.R. Heath-Brown, Artin's conjecture for primitive roots, *Quart. J. Math. Oxford*, (**2**) **37** (1986) 27-38.
- [8] C. Hooley, On Artin's Conjecture, *J. reine angew. Math.*, **226** (1967) 209-220.

- [9] H. Montgomery, R. C. Vaughan, T. D. Wooley, Some remarks on Gauss sums associated with k th powers, *Math. Proc. Camb. Phil. Soc.* , **118**, (1995), 21-33.
- [10] R. Murty and S. Srinivasan, Some remarks on Artin's Conjecture, *Canad. Math. Bull.*, **30** (1), (1987) 80-85.
- [11] R. Murty and S. Wong, The ABC conjecture and prime divisors of the Lucas and Lehmer sequences, *Number Theory for the Millenium, M. Bennett et al, editors*, **3**, 2002, pp. 43-54.
- [12] S. Sitaraman, Mean values of certain multiplicative functions and Artin's Primitive Roots Conjecture, *Hardy-Ramanujan Journal*, **37** (2015) 1-7.
- [13] G. Tenenbaum, Introduction to analytic and probabilistic number theory, *Cambridge Studies in Advanced Mathematics*, **46** 1995.
- [14] F. Thaine, Properties that characterize Gaussian periods and cyclotomic numbers, *Proc. Amer. Math. Soc.* , **124**(1) (1996) 35-45.
- [15] L. C. Washington, Introduction to Cyclotomic Fields, 2nd ed., Springer-Verlag (1997).