



HAL
open science

Fully algebraic domain decomposition preconditioners with adjustable spectral bounds

Loïc Gouarin, Nicole Spillane

► **To cite this version:**

Loïc Gouarin, Nicole Spillane. Fully algebraic domain decomposition preconditioners with adjustable spectral bounds. 2021. hal-03258644v1

HAL Id: hal-03258644

<https://hal.science/hal-03258644v1>

Preprint submitted on 11 Jun 2021 (v1), last revised 18 Jun 2021 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Fully algebraic domain decomposition preconditioners with adjustable spectral bounds ^{*}

Loïc Gouarin[†] Nicole Spillane[‡]

June 11, 2021

Abstract

In this article a new family of preconditioners is introduced for symmetric positive definite linear systems. The new preconditioners, called the AWG preconditioners (for Algebraic-Woodbury-GenEO) are constructed algebraically. By this, we mean that only the knowledge of the matrix \mathbf{A} for which the linear system is being solved is required. Thanks to the GenEO spectral coarse space technique, the condition number of the preconditioned operator is bounded theoretically from above. This upper bound can be made smaller by enriching the coarse space with more spectral modes.

The novelty is that, unlike in previous work on the GenEO coarse spaces, no knowledge of a partially non-assembled form of \mathbf{A} is required. Indeed, the spectral coarse space technique is not applied directly to \mathbf{A} but to a low-rank modification of \mathbf{A} of which a suitable non-assembled form is known by construction. The extra cost is a second (and to this day rather expensive) coarse solve in the preconditioner. One of the AWG preconditioners has already been presented in the short preprint [37]. This article is the first full presentation of the larger family of AWG preconditioners. It includes proofs of the spectral bounds as well as numerical illustrations.

1 Introduction

Throughout this article we consider the problem of finding $\mathbf{x}_* \in \mathbb{R}^n$ that is the solution of the following linear system

$$\mathbf{A}\mathbf{x}_* = \mathbf{b}, \text{ where } \mathbf{A} \in \mathbb{R}^{n \times n} \text{ is symmetric positive and definite (spd),} \quad (1)$$

for a given right hand side $\mathbf{b} \in \mathbb{R}^n$.

The applications we have in mind are ones for which \mathbf{A} is typically sparse and the number n of unknowns is very large. Hence, we study parallel solvers and more specifically preconditioners for the preconditioned conjugate gradient (PCG) method [34][Section 9.2]. Our objective is to propose a new preconditioner \mathbf{H} for solving (1) in such a way that the condition number of $\mathbf{H}\mathbf{A}$ is bounded from above by a *small enough* constant chosen by the user. This guarantees that the PCG method will converge in *sufficiently few* iterations [41][Lemma C.10]. Two-level domain decomposition preconditioners already exist that satisfy such a nice property (specifically the spectral coarse space methods described below). These methods, however, rely on the knowledge

^{*}Received... Accepted... Published online on... Recommended by...

[†]CNRS, CMAP, École Polytechnique, Institut Polytechnique de Paris, 91128 Palaiseau Cedex, France (loic.gouarin@cmmap.polytechnique.fr)

[‡]CNRS, CMAP, École Polytechnique, Institut Polytechnique de Paris, 91128 Palaiseau Cedex, France (nicole.spillane@cmmap.polytechnique.fr)

of some partially unassembled form of matrix \mathbf{A} . The additional challenge that we tackle in this work is that the new preconditioner must rely only on the knowledge of the matrix \mathbf{A} , and this is the meaning of the word *algebraic* as it is in Algebraic Multigrid (AMG) [8, 33, 44].

Very generally speaking, domain decomposition methods partition the domain Ω in which the solution is sought into smaller spaces, indexed by $s \in \llbracket 1, N \rrbracket$, and characterized by prolongation matrices $\mathbf{R}^{s\top}$ that satisfy $\sum_{s=1}^N \text{range}(\mathbf{R}^{s\top}) = \Omega$. One-level domain decomposition preconditioners then approximate \mathbf{A}^{-1} by a sum (interpolated by the $\mathbf{R}^{s\top}$) of inverses of well-chosen problems $\tilde{\mathbf{A}}^s$. Two-level domain decomposition methods have an additional space called the coarse space (generated by the columns of a matrix $\mathbf{R}^{0\top}$). A coarse problem $\tilde{\mathbf{A}}^0$ is solved in the coarse space. As an example, an application of the two-level additive preconditioner to a vector $\mathbf{x} \in \mathbb{R}^n$ takes the form: $\sum_{s=1}^N \mathbf{y}^s$ where

$$\mathbf{y}^s = \mathbf{R}^{s\top} (\tilde{\mathbf{A}}^s)^{-1} \mathbf{R}^s \mathbf{x} \quad \text{and} \quad \mathbf{y}^0 = \mathbf{R}^{0\top} (\tilde{\mathbf{A}}^0)^{-1} \mathbf{R}^0 \mathbf{x}.$$

prolongation (by 0) to Ω
local solve
restriction to Ω^s
coarse interpolation
coarse solve
interpolation back into Ω

The notation in the previous line should seem completely natural to readers who have already studied domain decomposition. The coarse contribution \mathbf{y}^0 is written separately to insist on the fact that it plays a different role to the other \mathbf{y}^s . Usually, the coarse space is computed for a given choice of the \mathbf{R}^s and $\tilde{\mathbf{A}}^s$.

The choice of the coarse space is in fact a very crucial topic in domain decomposition. Over the last decade, the range of symmetric positive definite problems for which two-level domain decomposition preconditioners can be made scalable and robust by a good choice of the coarse space has become very large by the development of so-called spectral coarse spaces. The following list gives an overview of some of these contributions: [28, 29, 13, 15, 16, 14, 38, 39, 17, 1, 21] for Additive Schwarz, [45] for Additive Average Schwarz, [40, 1] for BDD and FETI, [26, 24, 10, 23, 11, 31, 25, 46] for BDDC and/or FETI-DP, [20] for Optimized Schwarz and [27] in the context of boundary element methods.

In this article it is particularly referred to the GenEO coarse spaces [38, 39, 40] to which one of the authors contributed. The abstract theory of course spaces of the GenEO family in [36] is applied within the definition and analysis of the new Algebraic-Woodbury-GenEO (AWG) preconditioners.

The spectral coarse spaces mentioned above, and in particular the GenEO coarse spaces, are computed by solving one or two generalized eigenvalue problems in each subdomain, then selecting either the lowest or highest-frequency eigenvectors and prolongating them to the global domain. To the best of the authors' knowledge, all of the spectral coarse spaces for which there are no assumptions on the shapes of subdomains and the distribution of material parameters require the knowledge of a set of symmetric positive semi-definite matrices \mathbf{N}^s that satisfy

$$\exists C > 0 \text{ such that } \sum_{s=1}^N \langle \mathbf{x}, \mathbf{R}^{s\top} \mathbf{N}^s \mathbf{R}^s \mathbf{x} \rangle \leq C \langle \mathbf{x}, \mathbf{A} \mathbf{x} \rangle \quad \forall \mathbf{x} \in \mathbb{R}^n. \quad (2)$$

The matrices \mathbf{N}^s enter directly into the coarse space construction *via* the choice of matrix pencil for the generalized eigenvalue problems. In other words they play an essential role.

For matrices arising from discretized PDEs the above assumption is far from unnatural and the matrices \mathbf{N}^s are not expensive to compute as long as it is known at assembly that they are

required. As an example, if \mathbf{A} arises from the finite element discretization (with bases functions $\{\phi_k\}_{k=1,\dots,n}$) of the Laplace equation over some domain $\omega \subset \mathbb{R}^2$ or 3 , then the coefficients in \mathbf{A} are $\int_{\Omega} \nabla \phi_i \cdot \nabla \phi_j$. Assuming that the degrees of freedom selected by the restriction matrix \mathbf{R}^s are those in some $\omega^s \subset \omega$, \mathbf{N}^s can be taken to be the matrix whose coefficients are $\int_{\omega^s} \nabla \phi_i \cdot \nabla \phi_j$ (for the basis functions ϕ_i and ϕ_j whose support intersects ω^s). This is how condition (2) is usually satisfied. These \mathbf{N}^s are sometimes called the local Neumann matrices. Partial assembly over subdomains is neither hard conceptually nor expensive computationally and the purpose of this article is not to rule it out when it is possible. There are however many cases where only \mathbf{A} is known or available without writing many more lines of code. Then, the unassembled information is simply lost and the GenEO coarse spaces can't be computed. This is quite a common scenario: the problem may have been assembled by another user or with another piece of software. In this case only black box algorithms can be used.

Direct solvers, like MUMPS [3, 2], belong to the category of black box solvers and they are the most efficient up to a certain problem size. More closely related to domain decomposition are the multigrid algorithms, a very well-established set of solvers that solves the problem by iterating over the original (fine) problem as well as coarser and coarser approximations of it. The original multigrid algorithm [7, 9] is often referred to as geometric multigrid as it requires more information on the problem than just the matrix \mathbf{A} . Our objective with this algorithm is not unlike the objective of Algebraic Multigrid (AMG) which is to make multigrid applicable in cases where less information about the problem is accessible or some assumptions are not satisfied (see [8, 33] for the original contributions or [44] for a unified presentation and theory of most multigrid algorithms, many of which have not been mentioned). Our ambition here is to propose an algorithm as easy to apply as are the Algebraic Multigrid algorithms.

In this article, a new preconditioner (with several variants) is proposed for the cases where \mathbf{A} is already assembled. It is a domain decomposition preconditioner with two coarse space. The methodology is the following:

- The problem matrix \mathbf{A} is split into symmetric but possibly indefinite matrices \mathbf{B}^s as $\mathbf{A} = \sum_{s=1}^N \mathbf{R}^{s\top} \mathbf{B}^s \mathbf{R}^s$.
- The positive parts \mathbf{A}_+^s of the matrices \mathbf{B}^s are computed and assembled to form a global matrix $\mathbf{A}_+ = \sum_{s=1}^N \mathbf{R}^{s\top} \mathbf{A}_+^s \mathbf{R}^s$. By construction, a splitting of \mathbf{A}_+ into spsd matrices (*i.e.*, a suitable partially unassembled form) of \mathbf{A}_+ is known. Consequently, two-level preconditioners with GenEO coarse spaces can be computed for \mathbf{A}_+ by applying the abstract theory in [36].
- Finally, the Woodbury matrix identity relates the inverses of \mathbf{A} and of \mathbf{A}_+ and makes apparent that a good preconditioner for \mathbf{A} can be obtained by adding a second coarse space to a GenEO preconditioner for \mathbf{A}_+ .

Full theory for the condition number of the new preconditioned operators is given as well as numerical illustrations. The outline of the remainder of this article is as follows. In Section 2, some elements of the Abstract Schwarz theory [41] are recalled in their fully algebraic form. For readers less familiar with domain decomposition, the general form of a one-level and a two-level domain decomposition preconditioner is recalled. In Section 3, the new operator \mathbf{A}_+ is introduced and several preconditioners with their GenEO coarse spaces are considered. For each one, the spectral bounds are given by applying a result from [36]. Then, in Section 4, it is noticed that \mathbf{A}_+ is a low rank modification of \mathbf{A} and that applying the Woodbury matrix identity makes apparent how to add a second coarse space to the preconditioners for \mathbf{A}_+ in order to get a preconditioner for \mathbf{A} that satisfies the same convergence bounds. Each of these new preconditioners is indexed by one or two parameters that can be adjusted to decrease the condition number of the preconditioned operator by enriching the GenEO coarse space with more spectral modes. Some comments are also

made about the implementation of the new preconditioners. Finally, Section 5 presents some preliminary numerical results with the objective of confirming the theoretical results and illustrating the practical behaviour of the new algebraic preconditioners.

2 Abstract Schwarz Framework in the Algebraic Setting

An algebraic version of the abstract Schwarz framework is introduced in this section. This means that all the domain decomposition-type operators are written only in terms of vectors in \mathbb{R}^n .

2.1 Subdomains

Let $\Omega = \llbracket 1, n \rrbracket$ be the set of all indices in \mathbb{R}^n .

Definition 1 (Partition of Ω). *A set $(\Omega^s)_{s=1, \dots, N}$ of $N \in \mathbb{N}$ subsets of $\Omega = \llbracket 1, n \rrbracket$ is called a partition of Ω if*

$$\Omega = \bigcup_{s=1}^N \Omega^s.$$

The partition is said to have at least minimal overlap if Assumption 1 is satisfied.

Assumption 1 (Minimal overlap). *For any pair of indices $(i, j) \in \llbracket 1, n \rrbracket^2$, denoting by A_{ij} the coefficient of \mathbf{A} at the i -th line and j -th column,*

$$A_{ij} \neq 0 \Rightarrow (\exists s \in \llbracket 1, N \rrbracket \text{ such that } \{i, j\} \subset \Omega^s).$$

The usual global-to-local restriction matrices are defined next.

Definition 2. *For each $s = 1, \dots, N$, let n^s be the cardinality of Ω^s . Then, let $\mathbf{R}^s \in \mathbb{R}^{n^s \times n}$ be the restriction matrix defined by: \mathbf{R}^s is zero everywhere except for the block formed by the columns in Ω^s which is the $n^s \times n^s$ identity matrix.*

By simply performing the multiplications it can be proved that

$$\mathbf{R}^{s\top} \mathbf{R}^s \text{ is diagonal and that } \mathbf{R}^s \mathbf{R}^{s\top} = \mathbf{I} \text{ (the identity matrix in } \mathbb{R}^{n^s} \text{)}.$$

2.2 Partition of unity, coloring constant

In the construction and analysis of the preconditioners, two more elements from the abstract Schwarz theory are needed: the partition of unity matrices and the coloring constants.

Assumption 2 (Partition of unity matrices). *Let $\{\mathbf{D}^s \in \mathbb{R}^{n^s \times n^s}; s = 1, \dots, N\}$ be a family of matrices that satisfies*

$$\mathbf{I} = \sum_{s=1}^N \mathbf{R}^{s\top} \mathbf{D}^s \mathbf{R}^s, \text{ with } \mathbf{I} \text{ the } n \times n \text{ identity matrix.} \quad (3)$$

One way to fulfill Assumption 2 is to choose the following partition of unity matrices.

Definition 3 (Possible choice of partition of unity \mathbf{D}^s). *First, let $\mathbf{D} \in \mathbb{R}^{n \times n}$ be the non-singular diagonal matrix defined by*

$$\mathbf{D} := \left(\sum_{t=1}^N \mathbf{R}^{t\top} \mathbf{R}^t \right)^{-1}.$$

Then, for each $s \in \llbracket 1, N \rrbracket$, let $\mathbf{D}^s \in \mathbb{R}^{n^s \times n^s}$ be defined by

$$\mathbf{D}^s := \mathbf{R}^s \mathbf{D} \mathbf{R}^{s\top}.$$

The matrices in the above definition satisfy Assumption 2 (see [36][Lemma 4]). Their coefficients are the inverses of the multiplicity of each degree of freedom. Next, the coloring constant is defined in agreement with [41][Section 2.5.1]. The dependency of the coloring constant on the matrix with respect to which orthogonality is taken is written explicitly.

Definition 4 (Coloring constant). *Let $\mathbf{M} \in \mathbb{R}^{n \times n}$ be a symmetric matrix. Let $\mathcal{N}(\mathbf{M}) \in \mathbb{N}$ be such that there exists a set $\{\mathcal{C}_j; 1 \leq j \leq \mathcal{N}(\mathbf{M})\}$ of pairwise disjoint subsets of $\llbracket 1, N \rrbracket$ satisfying*

$$\llbracket 1, N \rrbracket = \bigcup_{1 \leq j \leq \mathcal{N}(\mathbf{M})} \mathcal{C}_j \quad \text{and} \quad \forall j \in \llbracket 1, \mathcal{N}(\mathbf{M}) \rrbracket : \{s, t\} \subset \mathcal{C}_j \Rightarrow (\mathbf{R}^s \mathbf{M} \mathbf{R}^{t\top} = \mathbf{0} \text{ or } s = t).$$

2.3 Abstract Schwarz preconditioners

One-level abstract Schwarz preconditioners are of the form:

$$\mathbf{H} := \sum_{s=1}^N \mathbf{R}^{s\top} \tilde{\mathbf{A}}^s \dagger \mathbf{R}^s. \quad (4)$$

where for each $s = 1, \dots, N$, it is assumed that

$$\tilde{\mathbf{A}}^s \in \mathbb{R}^{n^s \times n^s} \text{ is an spsd matrix, and that } \tilde{\mathbf{A}}^{s\dagger} \text{ is the pseudo-inverse of } \tilde{\mathbf{A}}^s.$$

Two-level domain decomposition preconditioner have two extra ingredients compared to the one-level method that they are based on: a coarse space and a coarse solver. Let's assume that the coarse space is denoted by V^0 and that the interpolation operator \mathbf{R}^0 satisfies Assumption 3.

Assumption 3. *A basis for the coarse space V^0 is stored in the lines of a matrix denoted \mathbf{R}^0 :*

$$V^0 = \text{range}(\mathbf{R}^{0\top}); \quad \mathbf{R}^0 \in \mathbb{R}^{n^0 \times n}; \quad n^0 = \dim(V^0); \quad n^0 < n.$$

The most common choice for the coarse solver, and the one that we wish to introduce, is the so called *exact* solver, where the word *exact* is with respect to the problem being solved. If the matrix in the linear system is an spd matrix $\tilde{\mathbf{A}}$ then the matrix that is inverted during the coarse solve is $\mathbf{R}^0 \tilde{\mathbf{A}} \mathbf{R}^{0\top}$.

With the same $\tilde{\mathbf{A}}^s$ and \mathbf{R}^0 , there are still at least two two-level preconditioners: the two-level additive preconditioner (denoted by \mathbf{H}_{ad}), and the hybrid preconditioner (denoted by \mathbf{H}_{hyb} and also called the deflated preconditioner). They are defined as follows:

$$\mathbf{H}_{ad} := \mathbf{H} + \mathbf{R}^{0\top} (\mathbf{R}^0 \tilde{\mathbf{A}} \mathbf{R}^{0\top})^{-1} \mathbf{R}^0, \quad (5)$$

and

$$\mathbf{H}_{hyb} := \mathbf{\Pi} \mathbf{H} \mathbf{\Pi}^\top + \mathbf{R}^{0\top} (\mathbf{R}^0 \tilde{\mathbf{A}} \mathbf{R}^{0\top})^{-1} \mathbf{R}^0;$$

where

$$\mathbf{\Pi} := \mathbf{I} - \mathbf{R}^{0\top} (\mathbf{R}^0 \tilde{\mathbf{A}} \mathbf{R}^{0\top})^{-1} \mathbf{R}^0 \tilde{\mathbf{A}}; \quad \mathbf{I} \text{ is the } n \times n \text{ identity matrix.} \quad (6)$$

The generic notation $\tilde{\mathbf{A}}$ has very deliberately been used in the previous equations instead of \mathbf{A} . Indeed, the two level preconditioner in this article is a preconditioner for a new matrix that will be denoted \mathbf{A}_+ . The next section gives the definition of \mathbf{A}_+ and the choices of $\tilde{\mathbf{A}}^s$ and V^0 that make the characterization of an abstract two-level preconditioner for \mathbf{A}_+ complete.

3 A new matrix \mathbf{A}_+ and its two level GenEO preconditioners

This section introduces a lot of the new operators and notation. Since it is not known how to algebraically find local spsd matrices that satisfy (2), it is chosen to relax the assumption by allowing the matrices to be indefinite. Precisely, in subsection 3.1 it is assumed that symmetric matrices \mathbf{B}^s are known such that $\mathbf{A} = \sum_{s=1}^N \mathbf{R}^s \mathbf{B}^s \mathbf{R}^s$ and one possible (algebraic) choice of \mathbf{B}^s is given. In Section 3.2, each \mathbf{B}^s is in turn split into an spsd part and a symmetric negative semi-definite part as $\mathbf{B}^s = \mathbf{A}_+^s - \mathbf{A}_-^s$. Finally, the spsd parts are assembled to form $\mathbf{A}_+ = \sum_{s=1}^N \mathbf{R}^s \mathbf{A}_+^s \mathbf{R}^s$. The resulting matrix \mathbf{A}_+ is shown to be symmetric positive definite and, by construction, it satisfies (2) with $\mathbf{N}^s = \mathbf{A}_+^s$ and $C = 1$. In other words, a very nice characteristic of \mathbf{A}_+ is that the abstract GenEO theory applies to defining and analyzing two level preconditioners for \mathbf{A}_+ . Four of these are considered in Section 3.4.

3.1 A splitting of \mathbf{A} into symmetric matrices

The matrices \mathbf{B}^s in the assumption below are the starting point for the new preconditioners. To make the construction complete, an example of such matrices, is given below. It is (of course) constructed algebraically and is the one used in numerical computations. This choice is however far from unique.

Assumption 4. *Let's assume that there exists a family of symmetric matrices $\mathbf{B}^s \in \mathbb{R}^{n^s \times n^s}$ for $s = 1, \dots, N$ such that*

$$\mathbf{A} = \sum_{s=1}^N \mathbf{R}^s \mathbf{B}^s \mathbf{R}^s.$$

Such a family of matrices can always be chosen under Assumption 1 (minimal overlap). Indeed, one possible choice is given in the new definition.

Definition 5 (Possible choice of matrix \mathbf{B}^s). *Let $\mathbf{S}(\mathbf{A})$ be the $n \times n$ boolean matrix that shares the same sparsity pattern as \mathbf{A} :*

$$(\mathbf{S}(\mathbf{A}))_{ij} := \begin{cases} 1 & \text{if } A_{ij} \neq 0 \\ 0 & \text{otherwise} \end{cases} \quad \text{for any } i, j \in \llbracket 1, n \rrbracket.$$

Then let \mathbf{M}_μ be the matrix that counts the number of subdomains that each pair of indices in $\{\{i, j\}; A_{ij} \neq 0\}$ belongs to:

$$\mathbf{M}_\mu := \sum_{s=1}^N \mathbf{R}^s \mathbf{S}(\mathbf{A}) \mathbf{R}^s \mathbf{R}^s \mathbf{R}^s,$$

and let \mathbf{B} be the Hadamard division of \mathbf{A} by \mathbf{M}_μ

$$B_{ij} := \begin{cases} A_{ij}/M_{\mu ij} & \text{if } A_{ij} \neq 0 \\ 0 & \text{otherwise} \end{cases} \quad \text{for any } i, j \in \llbracket 1, n \rrbracket.$$

Finally, set \mathbf{B}^s to be the block of \mathbf{B} corresponding to degrees of freedom in Ω^s :

$$\mathbf{B}^s := \mathbf{R}^s \mathbf{B} \mathbf{R}^s \mathbf{R}^s \mathbf{R}^s.$$

Note that from the previous definition only the notation \mathbf{B}^s will be reused further on in the article. We next check that these matrices \mathbf{B}^s are indeed suitable.

Theorem 1. *Let \mathbf{A} be an order n spd matrix, let $(\Omega^s)_{s=1, \dots, N}$ represent the partition into subdomains and let $(\mathbf{R}^s)_{s=1, \dots, N}$ be the set of restriction matrices from Definition 2. Under Assumption 1, the matrices \mathbf{B}^s from Definition 5 satisfy Assumption 4.*

Proof. First, we justify the fact that \mathbf{M}_μ counts the multiplicity of the pairs of degrees of freedom $\{i, j\}$ for which $A_{ij} \neq 0$:

$$(M_\mu)_{ij} = \left(\sum_{s=1}^N \mathbf{R}^{s\top} \mathbf{R}^s \mathbf{S}(\mathbf{A}) \mathbf{R}^{s\top} \mathbf{R}^s \right)_{ij} = \sum_{\{s; \{i, j\} \subset \Omega^s\}} \left(\mathbf{R}^{s\top} \mathbf{R}^s \mathbf{S}(\mathbf{A}) \mathbf{R}^{s\top} \mathbf{R}^s \right)_{ij} = \sum_{\{s; \{i, j\} \subset \Omega^s\}} (S(\mathbf{A}))_{ij}$$

so, for any $i, j \in \llbracket 1, n \rrbracket$,

$$(M_\mu)_{ij} = \begin{cases} \#\{s; \{i, j\} \subset \Omega^s\} & \text{if } A_{ij} \neq 0, \\ 0 & \text{otherwise.} \end{cases}$$

With a similar calculation, it can then be checked that the \mathbf{B}^s form a splitting of \mathbf{A} :

$$\left(\sum_{s=1}^N \mathbf{R}^{s\top} \mathbf{B}^s \mathbf{R}^s \right)_{ij} = \left(\sum_{s=1}^N \mathbf{R}^{s\top} \mathbf{R}^s \mathbf{B}^s \mathbf{R}^{s\top} \mathbf{R}^s \right)_{ij} = \sum_{\{s; \{i, j\} \subset \Omega^s\}} B_{ij}$$

so, for any $i, j \in \llbracket 1, n \rrbracket$,

$$\left(\sum_{s=1}^N \mathbf{R}^{s\top} \mathbf{B}^s \mathbf{R}^s \right)_{ij} = \begin{cases} \sum_{\{s; \{i, j\} \subset \Omega^s\}} A_{ij} / M_{\mu_{ij}} = A_{ij} & \text{if } A_{ij} \neq 0, \\ 0 = A_{ij} & \text{otherwise.} \end{cases}$$

It can be concluded that $\mathbf{A} = \sum_{s=1}^N \mathbf{R}^{s\top} \mathbf{B}^s \mathbf{R}^s$. \square

Remark 1. Notice that, if the minimal overlap condition is not satisfied, then there exists a pair of indices $\{i, j\}$ for which $A_{ij} \neq 0$ but $\{s; \{i, j\} \subset \Omega^s\} = \emptyset$. This leads to $\left(\sum_{s=1}^N \mathbf{R}^{s\top} \mathbf{B}^s \mathbf{R}^s \right)_{ij} = 0$ and shows that it is impossible that Assumption 4 be satisfied without the minimal overlap condition (no matter how the matrices \mathbf{B}^s are chosen).

3.2 Definition of \mathbf{A}_+

The first step in defining the very important matrix \mathbf{A}_+ is to split \mathbf{B}^s into its positive part and its negative semi-definite part.

Definition 6 (Local splitting of the splitting). Let \mathbf{B}^s , for $s \in \llbracket 1, N \rrbracket$, be a family of matrices that satisfy Assumption 4. For each s , let a diagonalization of \mathbf{B}^s be written as

$$\mathbf{B}^s = \mathbf{V}^s \mathbf{\Lambda}^s \mathbf{V}^{s\top}; \text{ with } \mathbf{V}^s \text{ orthogonal and } \mathbf{\Lambda}^s \text{ diagonal.}$$

Assume, without loss of generality, that the diagonal values $(l_{ii}^s)_{i=1, \dots, n^s}$ of $\mathbf{\Lambda}^s$ (which are the eigenvalues of \mathbf{B}^s) are sorted in non-decreasing order. Let n_+^s be the number of positive eigenvalues and $n_-^s = n^s - n_+^s$ be the number of non-positive eigenvalues. Let $\mathbf{V}_-^s, \mathbf{V}_+^s, \mathbf{\Lambda}_-^s, \mathbf{\Lambda}_+^s$ be the blocks of \mathbf{V}^s and $\mathbf{\Lambda}^s$ that satisfy

$$\mathbf{\Lambda}^s = \begin{pmatrix} \mathbf{\Lambda}_-^s & \mathbf{0} \\ \mathbf{0} & \mathbf{\Lambda}_+^s \end{pmatrix}, \quad \mathbf{V}^s = [\mathbf{V}_-^s | \mathbf{V}_+^s], \quad \mathbf{\Lambda}_+^s \text{ is spd,} \quad -\mathbf{\Lambda}_-^s \text{ is spsd.}$$

Finally, define the two following matrices in $\mathbb{R}^{n^s \times n^s}$

$$\mathbf{A}_+^s := \mathbf{V}_+^s \mathbf{\Lambda}_+^s \mathbf{V}_+^{s\top} \text{ and } \mathbf{A}_-^s := -\mathbf{V}_-^s \mathbf{\Lambda}_-^s \mathbf{V}_-^{s\top}.$$

It is clear that, for each $s \in \llbracket 1, N \rrbracket$, both matrices \mathbf{A}_+^s and \mathbf{A}_-^s are spsd matrices and that $\mathbf{B}^s = \mathbf{A}_+^s - \mathbf{A}_-^s$. Next, global matrices are computed by assembling the local components with the usual restriction and prolongation operators \mathbf{R}^s and $\mathbf{R}^{s\top}$.

Definition 7 (Global counterparts of the splitting matrices). *Let the global matrices $\mathbf{A}_+ \in \mathbb{R}^{n \times n}$ and $\mathbf{A}_- \in \mathbb{R}^{n \times n}$ be defined by*

$$\mathbf{A}_+ := \sum_{s=1}^N \mathbf{R}^s \mathbf{A}_+^s \mathbf{R}^s, \text{ and } \mathbf{A}_- := \sum_{s=1}^N \mathbf{R}^s \mathbf{A}_-^s \mathbf{R}^s.$$

Theorem 2. *The matrices \mathbf{A}_+ and \mathbf{A}_- from Definition 7 satisfy the following three properties*

- (i) $\mathbf{A} = \mathbf{A}_+ - \mathbf{A}_-$,
- (ii) *the matrix \mathbf{A}_- is symmetric positive semi-definite,*
- (iii) *the matrix \mathbf{A}_+ is symmetric positive definite.*

Proof. By definition, $\mathbf{A}_+ = \sum_{s=1}^N \mathbf{R}^s \mathbf{V}_+^s \mathbf{\Lambda}_+^s \mathbf{V}_+^{s \top} \mathbf{R}^s$ and $\mathbf{A}_- = -\sum_{s=1}^N \mathbf{R}^s \mathbf{V}_-^s \mathbf{\Lambda}_-^s \mathbf{V}_-^{s \top} \mathbf{R}^s$ where the matrices $\mathbf{\Lambda}_+^s$ ($s \in \llbracket 1, N \rrbracket$) are diagonal matrices with positive entries and the matrices $\mathbf{\Lambda}_-^s$ ($s \in \llbracket 1, N \rrbracket$) are diagonal matrices with non-positive entries. Consequently, \mathbf{A}_+ and \mathbf{A}_- are both spsd and item (ii) is proved. Moreover, by Definition 6 and Assumption 4, item (i) holds:

$$\begin{aligned} \mathbf{A}_+ - \mathbf{A}_- &= \sum_{s=1}^N \mathbf{R}^s \mathbf{V}_+^s \mathbf{\Lambda}_+^s \mathbf{V}_+^{s \top} \mathbf{R}^s + \sum_{s=1}^N \mathbf{R}^s \mathbf{V}_-^s \mathbf{\Lambda}_-^s \mathbf{V}_-^{s \top} \mathbf{R}^s \\ &= \sum_{s=1}^N \mathbf{R}^s \mathbf{B}^s \mathbf{R}^s \\ &= \mathbf{A}. \end{aligned}$$

Finally, it has already been argued that \mathbf{A}_+ is spsd so, to prove item (iii), it remains only to confirm that the kernel of \mathbf{A}_+ is restricted to the zero vector. Let $\mathbf{x} \in \mathbb{R}^n$ such that $\mathbf{A}_+ \mathbf{x} = \mathbf{0}$,

$$0 = \langle \mathbf{x}, \mathbf{A}_+ \mathbf{x} \rangle = \langle \mathbf{x}, \mathbf{A} \mathbf{x} \rangle + \langle \mathbf{x}, \mathbf{A}_- \mathbf{x} \rangle \geq \langle \mathbf{x}, \mathbf{A} \mathbf{x} \rangle,$$

and this last term equals $\mathbf{0}$ only if $\mathbf{x} = \mathbf{0}$ which ends the proof. \square

Remark 2. *The zero eigenvalues of \mathbf{B}^s are in $\mathbf{\Lambda}_-^s$. Another possibility would be to put them into $\mathbf{\Lambda}_+^s$.*

The previously defined matrix \mathbf{A}_+ is an spd matrix for which we have knowledge of spsd local matrices $\mathbf{N}^s = \mathbf{A}_+^s$ that satisfy (2) with $C = 1$. This means that it fits right into the abstract GenEO theory [36] and, hence, a variety of two level preconditioners with guaranteed convergence rates can be defined. First, the one level preconditioners to which to apply GenEO are chosen and then the GenEO coarse spaces are given. In terms of the two ingredients still missing in the abstract two-level preconditioners from Section 2.3: the local solvers \mathbf{A}^s are defined in Section 3.3 and the coarse interpolation operators \mathbf{R}^0 are defined in Section 3.4.

3.3 One-level preconditioners for \mathbf{A}_+

In order to define a one-level preconditioner in our framework it remains only to choose the matrices $\tilde{\mathbf{A}}^s$ (*i.e.*, the local solvers) in the abstract form (4). Three types of local solvers are introduced as all three are natural choices for \mathbf{A}_+ : the exact local solver ($\tilde{\mathbf{A}}^s = \mathbf{R}^s \mathbf{A}_+ \mathbf{R}^{s \top}$), the matrix in the spsd splitting with adequate weights ($\tilde{\mathbf{A}}^s = \mathbf{D}^{s-1} \mathbf{A}_+^s \mathbf{D}^{s-1}$), and what would be the exact solver if we were solving a problem with \mathbf{A} since this is, after all, our endgame ($\tilde{\mathbf{A}}^s = \mathbf{R}^s \mathbf{A} \mathbf{R}^{s \top}$).

Definition 8. Let three one-level preconditioners be defined by:

$$\mathbf{H}_+^{AS} := \sum_{s=1}^N \mathbf{R}^s \mathbf{A}_+ \mathbf{R}^{s\top} (\mathbf{R}^s \mathbf{A}_+ \mathbf{R}^{s\top})^{-1} \mathbf{R}^s,$$

$$\mathbf{H}^{AS} := \sum_{s=1}^N \mathbf{R}^s \mathbf{A} \mathbf{R}^{s\top} (\mathbf{R}^s \mathbf{A} \mathbf{R}^{s\top})^{-1} \mathbf{R}^s,$$

and

$$\mathbf{H}^{NN} := \sum_{s=1}^N \mathbf{R}^s \mathbf{D}^s (\mathbf{A}_+^s)^\dagger \mathbf{D}^s \mathbf{R}^s = \sum_{s=1}^N \mathbf{R}^s \mathbf{D}^s \mathbf{V}_+^s (\mathbf{A}_+^s)^{-1} \mathbf{V}_+^{s\top} \mathbf{D}^s \mathbf{R}^s,$$

where \mathbf{D}^s are the partition of unity matrices from Definition 3.

It is recalled that, for $s \in \llbracket 1, N \rrbracket$, \mathbf{A}_+^s , \mathbf{V}_+^s , and \mathbf{A}_+^s were introduced in Definition 6, \mathbf{D}^s in Definition 3, and \mathbf{A}_+ in Definition 7.

Lemma 1. The one-level preconditioners \mathbf{H}_+^{AS} , \mathbf{H}^{AS} and \mathbf{H}^{NN} from Definition 8 are spd.

Proof. The preconditioners \mathbf{H}_+^{AS} and \mathbf{H}^{AS} are usual Additive Schwarz preconditioners for spd matrices so they are spd. For the third preconditioner, it is obvious that \mathbf{H}^{NN} is spsd. Moreover, let $\mathbf{x} \in \text{Ker}(\mathbf{H}^{NN})$ then

$$0 = \langle \mathbf{x}, \mathbf{H}^{NN} \mathbf{x} \rangle = \sum_{s=1}^N \langle \mathbf{D}^s \mathbf{R}^s \mathbf{x}, \mathbf{A}_+^s \dagger \mathbf{D}^s \mathbf{R}^s \mathbf{x} \rangle.$$

For this to hold, each term in the sum of non-negative terms must also be zero so:

$$\text{for any } s \in \llbracket 1, N \rrbracket : \mathbf{D}^s \mathbf{R}^s \mathbf{x} \in \text{Ker}(\mathbf{A}_+^s \dagger) = \text{Ker}(\mathbf{A}_+^s).$$

Let's prolongate $(\mathbf{A}_+^s) \mathbf{D}^s \mathbf{R}^s \mathbf{x} = \mathbf{0}$ to the global domain with $\mathbf{R}^{s\top}$, sum over s and inject the definition of \mathbf{D}^s (Definition 3 in which \mathbf{D} is diagonal) to obtain

$$\mathbf{0} = \sum_{s=1}^N \mathbf{R}^s \mathbf{A}_+^s \mathbf{D}^s \mathbf{R}^s \mathbf{x} = \sum_{s=1}^N \mathbf{R}^s \mathbf{A}_+^s \mathbf{R}^s \mathbf{D} \mathbf{R}^{s\top} \mathbf{R}^s \mathbf{x} = \sum_{s=1}^N \mathbf{R}^s \mathbf{A}_+^s \mathbf{R}^s \mathbf{D} \mathbf{x} = \mathbf{A}_+ \mathbf{D} \mathbf{x}.$$

Finally, the non singularity of \mathbf{A}_+ and of \mathbf{D} allow to conclude that $\mathbf{x} = \mathbf{0}$ which ends the proof. \square

Remark 3. For the proof of the non-singularity of \mathbf{H}^{NN} , the definition of the partition of unity matrices \mathbf{D}^s was used (Definition 3). A general proof does not go through for all partitions of unity (i.e., if the \mathbf{D}^s in Definition 8 are replaced by another family of matrices that satisfy Assumption 2). However, it is not likely that the \mathbf{D}^s could and would be chosen in a way that makes \mathbf{H}^{NN} singular. In other words, this is a technical restriction and other choices of partition of unity matrices should definitely be explored. All parts of the article that are not related to \mathbf{H}^{NN} are not concerned by this technical restriction.

3.4 Two level preconditioners for \mathbf{A}_+ with GenEO

Next, the GenEO coarse spaces that correspond to solving a linear system for \mathbf{A}_+ with each of the one level preconditioners are introduced. The corresponding spectral bounds for the preconditioned operators are given. The proofs consist in giving the adequate references to [36]. The information is organized with one theorem per choice of one-level preconditioner. First, some very useful notation is chosen to designate a normalized basis of the high (or low) frequency eigenvectors with respect to a certain matrix pencil and a certain threshold.

Definition 9. Let $m \in \mathbb{N}^*$, let $\mathbf{M}_A \in \mathbb{R}^{m \times m}$ be an spsd matrix, let $\mathbf{M}_B \in \mathbb{R}^{m \times m}$ be an spd matrix. Let $(\lambda_k, \mathbf{y}_k)_{k=1, \dots, m}$ be the (ordered and \mathbf{M}_B -normalized) eigenpairs of the generalized eigenproblem associated with matrix pencil $(\mathbf{M}_A, \mathbf{M}_B)$, i.e.,

$$\lambda_k \in \mathbb{R}, \quad \mathbf{y}_k \in \mathbb{R}^m, \quad \langle \mathbf{y}_k, \mathbf{M}_B \mathbf{y}_k \rangle = 1, \quad \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_m, \quad \text{and} \quad \mathbf{M}_A \mathbf{y}_k = \lambda_k \mathbf{M}_B \mathbf{y}_k.$$

For any scalar $\tau > 0$, set $m_L := \min \{k \in \llbracket 0, m-1 \rrbracket; \lambda_{k+1} \geq \tau\}$ if $\lambda_m \geq \tau$, and $m_L := m$ otherwise. Then define the two following matrices by concatenating eigenvectors

$$\mathbf{Y}_L(\tau, \mathbf{M}_A, \mathbf{M}_B) := [\mathbf{y}_1 | \dots | \mathbf{y}_{m_L}], \quad \text{and} \quad \mathbf{Y}_H(\tau, \mathbf{M}_A, \mathbf{M}_B) := [\mathbf{y}_{m_L+1} | \dots | \mathbf{y}_m].$$

The subscripts $*_L$ and $*_H$ refer to the words *low* and *high* depending on which end of the spectrum is selected. The definition is in agreement with the definition in [36].

Theorem 3 (Two-level preconditioners with GenEO for \mathbf{H}_+^{AS}). For any $\tau_b > 1$, let $V_{AS,+}^0(\tau_b)$ be defined by

$$V_{AS,+}^0(\tau_b) := \sum_{s=1}^N \text{range}(\mathbf{R}^{s\top} \mathbf{Y}_L(\tau_b^{-1}, \mathbf{D}^{s-1} \mathbf{A}_+^s \mathbf{D}^{s-1}, \mathbf{R}^s \mathbf{A}_+ \mathbf{R}^{s\top})),$$

and assume that a corresponding interpolation matrix $\mathbf{R}_{AS,+}^0(\tau_b)$ is defined to satisfy Assumption 3. Then, the coarse projector as well as the hybrid and additive preconditioners are defined naturally as

$$\begin{aligned} \mathbf{\Pi}_+^{AS}(\tau_b) &:= \mathbf{I} - \mathbf{R}_{AS,+}^0(\tau_b)^\top (\mathbf{R}_{AS,+}^0(\tau_b) \mathbf{A}_+ \mathbf{R}_{AS,+}^0(\tau_b)^\top)^{-1} \mathbf{R}_{AS,+}^0(\tau_b) \mathbf{A}_+, \\ \mathbf{H}_{+,hyb}^{AS}(\tau_b) &:= \mathbf{\Pi}_+^{AS}(\tau_b) \mathbf{H}_+^{AS} \mathbf{\Pi}_+^{AS}(\tau_b)^\top + \mathbf{R}_{AS,+}^0(\tau_b)^\top (\mathbf{R}_{AS,+}^0(\tau_b) \mathbf{A}_+ \mathbf{R}_{AS,+}^0(\tau_b)^\top)^{-1} \mathbf{R}_{AS,+}^0(\tau_b), \\ \mathbf{H}_{+,ad}^{AS}(\tau_b) &:= \mathbf{H}_+^{AS} + \mathbf{R}_{AS,+}^0(\tau_b)^\top (\mathbf{R}_{AS,+}^0(\tau_b) \mathbf{A}_+ \mathbf{R}_{AS,+}^0(\tau_b)^\top)^{-1} \mathbf{R}_{AS,+}^0(\tau_b). \end{aligned}$$

The eigenvalues of the preconditioned operators are bounded as follows

$$\begin{aligned} 1/\tau_b &\leq \lambda(\mathbf{H}_{+,hyb}^{AS}(\tau_b) \mathbf{A}_+) \leq \mathcal{N}(\mathbf{A}_+) \\ 1/((1 + 2\mathcal{N}(\mathbf{A}_+))\tau_b) &\leq \lambda(\mathbf{H}_{+,ad}^{AS}(\tau_b) \mathbf{A}_+) \leq \mathcal{N}(\mathbf{A}_+) + 1, \end{aligned}$$

where $\mathcal{N}(\mathbf{A}_+)$ is the coloring constant with respect to matrix \mathbf{A}_+ (see Definition 4).

Proof. This results from an application of [36][Corollary 3] (for the hybrid preconditioner) and [36][Corollary 4] (for the additive preconditioner) under [36][Assumption 7] and with $\mathcal{N}' = 1$, $\mathbf{M}^s = \mathbf{D}^{s-1} \mathbf{A}_+^s \mathbf{D}^{s-1}$, $\tilde{\mathbf{A}}^s = \mathbf{R}^s \mathbf{A}_+ \mathbf{R}^{s\top}$ and the alternate formulation for the coarse space given in [36][Definition 5]. \square

Note that, if $0 < \tau_b \leq 1$, there is also a spectral result which is slightly longer to state (it involves min and max).

Theorem 4 (Two-level preconditioner with GenEO for \mathbf{H}^{NN}). For any $0 < \tau_\sharp < 1$, let $V_{NN}^0(\tau_\sharp)$ be defined by

$$V_{NN}^0(\tau_\sharp) := \sum_{s=1}^N \text{range}(\mathbf{R}^{s\top} \mathbf{Y}_L(\tau_\sharp, \mathbf{D}^{s-1} \mathbf{A}_+^s \mathbf{D}^{s-1}, \mathbf{R}^s \mathbf{A}_+ \mathbf{R}^{s\top})),$$

and assume that a corresponding interpolation matrix $\mathbf{R}_{NN}^0(\tau_\sharp)$ is defined to satisfy Assumption 3. Then, the coarse projector as well as the hybrid preconditioner are defined naturally as

$$\begin{aligned} \mathbf{\Pi}^{NN}(\tau_\sharp) &:= \mathbf{I} - \mathbf{R}_{NN}^0(\tau_\sharp)^\top (\mathbf{R}_{NN}^0(\tau_\sharp) \mathbf{A}_+ \mathbf{R}_{NN}^0(\tau_\sharp)^\top)^{-1} \mathbf{R}_{NN}^0(\tau_\sharp) \mathbf{A}_+, \\ \mathbf{H}_{hyb}^{NN}(\tau_\sharp) &:= \mathbf{\Pi}^{NN}(\tau_\sharp) \mathbf{H}^{NN} \mathbf{\Pi}^{NN}(\tau_\sharp)^\top + \mathbf{R}_{NN}^0(\tau_\sharp)^\top (\mathbf{R}_{NN}^0(\tau_\sharp) \mathbf{A}_+ \mathbf{R}_{NN}^0(\tau_\sharp)^\top)^{-1} \mathbf{R}_{NN}^0(\tau_\sharp). \end{aligned}$$

The eigenvalues of the preconditioned operator are bounded as follows

$$1 \leq \lambda(\mathbf{H}_{hyb}^{NN}(\tau_{\sharp})\mathbf{A}_+) \leq \mathcal{N}(\mathbf{A}_+)/\tau_{\sharp}.$$

where $\mathcal{N}(\mathbf{A}_+)$ is the coloring constant with respect to matrix \mathbf{A}_+ (see Definition 4).

Proof. This results from an application of [36][Corollary 2] with $\mathcal{N}' = 1$, $\tau_b \rightarrow 1$ (as in [36][Section 5.2.3]), $\tilde{\mathbf{A}}^s = \mathbf{D}^{s-1}\mathbf{A}_+\mathbf{D}^{s-1}$ and $\mathbf{R}^s\mathbf{A}_+\mathbf{R}^{s\top}$ as the exact local solver (because we are solving for \mathbf{A}_+). \square

Note that, if $\tau_{\sharp} \geq 1$, there is also a spectral result which is slightly longer to state (it involves min and max). There is no spectral result for the two-level additive preconditioner so it is not considered in the theorem.

Theorem 5 (Two-level preconditioner with GenEO for \mathbf{H}^{AS}). *For any $\tau_b > 1$ and $0 < \tau_{\sharp} < 1$, let $V_{AS}^0(\tau_{\sharp}, \tau_b)$ be defined by*

$$V_{AS}^0(\tau_{\sharp}, \tau_b) := \sum_{s=1}^N \text{range}(\mathbf{R}^{s\top}\mathbf{Y}_L(\tau_b^{-1}, \mathbf{D}^{s-1}\mathbf{A}_+\mathbf{D}^{s-1}, \mathbf{R}^s\mathbf{A}\mathbf{R}^{s\top})) \\ + \sum_{s=1}^N \text{range}(\mathbf{R}^{s\top}\mathbf{Y}_L(\tau_{\sharp}, \mathbf{R}^s\mathbf{A}\mathbf{R}^{s\top}, \mathbf{R}^s\mathbf{A}_+\mathbf{R}^{s\top})),$$

and assume that a corresponding interpolation matrix $\mathbf{R}_{AS}^0(\tau_{\sharp}, \tau_b)$ is defined to satisfy Assumption 3. Then, the coarse projector as well as the hybrid preconditioner are defined naturally as

$$\mathbf{\Pi}^{AS}(\tau_{\sharp}, \tau_b) := \mathbf{I} - \mathbf{R}_{AS}^0(\tau_{\sharp}, \tau_b)^{\top}(\mathbf{R}_{AS}^0(\tau_{\sharp}, \tau_b)\mathbf{A}_+\mathbf{R}_{AS}^0(\tau_{\sharp}, \tau_b)^{\top})^{-1}\mathbf{R}_{AS}^0(\tau_{\sharp}, \tau_b)\mathbf{A}_+, \\ \mathbf{H}_{hyb}^{AS}(\tau_{\sharp}, \tau_b) := \mathbf{\Pi}^{AS}(\tau_{\sharp}, \tau_b)\mathbf{H}^{AS}\mathbf{\Pi}^{AS}(\tau_{\sharp}, \tau_b)^{\top} + \mathbf{R}_{AS}^0(\tau_{\sharp}, \tau_b)^{\top}(\mathbf{R}_{AS}^0(\tau_{\sharp}, \tau_b)\mathbf{A}_+\mathbf{R}_{AS}^0(\tau_{\sharp}, \tau_b)^{\top})^{-1}\mathbf{R}_{AS}^0(\tau_{\sharp}, \tau_b).$$

The eigenvalues of the preconditioned operator are bounded as follows

$$1/\tau_b \leq \lambda(\mathbf{H}_{hyb}^{AS}(\tau_{\sharp}, \tau_b)\mathbf{A}_+) \leq \mathcal{N}(\mathbf{A}_+)/\tau_{\sharp}.$$

where $\mathcal{N}(\mathbf{A}_+)$ is the coloring constant with respect to matrix \mathbf{A}_+ (see Definition 4).

Proof. This results from [36][Corollary 2] with $\mathcal{N}' = 1$, $\mathbf{M}^s = \mathbf{D}^{s-1}\mathbf{A}_+\mathbf{D}^{s-1}$, $\tilde{\mathbf{A}}^s = \mathbf{R}^s\mathbf{A}\mathbf{R}^{s\top}$, $\mathbf{R}^s\mathbf{A}_+\mathbf{R}^{s\top}$ as the exact local solver (because we are solving for \mathbf{A}_+), and the alternate formulation for the coarse space given in [36][Definition 5]. \square

Note that, if $0 < \tau_b \leq 1$, there is also a spectral result which is slightly longer to state (it involves min and max). The choice $\tau_{\sharp} \geq 1$ would lead to having all the vectors in the coarse space so this is excluded. There is no spectral result for the two-level additive preconditioner so it is not considered in the theorem.

Remark 4. *For all four considered choices of two-level preconditioner, it is easy to see that $\sum_{s=1}^N \mathbf{R}^{s\top}\mathbf{D}^s\mathbf{V}_-^s$ is included in the coarse space as $\text{Ker}(\mathbf{D}^{s-1}\mathbf{A}_+\mathbf{D}^{s-1}) = \mathbf{D}^s\mathbf{V}_-^s$.*

4 New fully algebraic preconditioners for \mathbf{A}

In the previous section, four two-level preconditioners for \mathbf{A}_+ (indexed by the choice of one or two thresholds) have been introduced with their spectral bounds: $\mathbf{H}_{+,hyb}^{AS}(\tau_b)$, $\mathbf{H}_{+,ad}^{AS}(\tau_b)$, $\mathbf{H}_{hyb}^{NN}(\tau_{\sharp})$, and $\mathbf{H}_{hyb}^{AS}(\tau_{\sharp}, \tau_b)$. In this section, let \mathbf{H}_2 denote any one of them and let $[\lambda_{\min}(\mathbf{H}_2\mathbf{A}_+), \lambda_{\max}(\mathbf{H}_2\mathbf{A}_+)]$ be an interval that contains all eigenvalues of the preconditioned operator $\mathbf{H}_2\mathbf{A}_+$. The subscript $*_2$ was chosen to refer to two-level preconditioners. We may now set aside the choice of a preconditioner for \mathbf{A}_+ and come back to our original problem of finding a preconditioner for \mathbf{A} .

4.1 Woodbury matrix identity for $\mathbf{A} = \mathbf{A}_+ - \mathbf{A}_-$

The new preconditioner for \mathbf{A} arises from the realization that \mathbf{A} can be viewed as a low rank modification of \mathbf{A}_+ and adding a term to \mathbf{H}_2 accordingly.

Theorem 6. *The rank of \mathbf{A}_- , which we denote by n_- , satisfies $n_- \leq \sum_{s=1}^N n^s - n$.*

Proof. By definition of \mathbf{A}_+ and \mathbf{A}_- , it holds that

$$\text{rank}(\mathbf{A}_+) + \text{rank}(\mathbf{A}_-) \leq \sum_{s=1}^N n^s,$$

and $\text{rank}(\mathbf{A}_+) = n$ as \mathbf{A}_+ is non-singular. \square

With words, the rank of \mathbf{A}_- is at most the difference between the number of degrees of freedom and the number of degrees of freedom multiplied by their multiplicity. If there is little overlap (while still satisfying the minimal overlap condition) then the rank of \mathbf{A}_- is small compared to the rank n of \mathbf{A} ($n_- \ll n$). In practice it is possible, and desirable that n_- be much smaller even than $\sum_{s=1}^N n^s - n$. Following this observation it is natural to write \mathbf{A} as a low rank modification of \mathbf{A}_+ .

To this end, let's introduce the factors in the diagonalization of \mathbf{A}_- .

Definition 10. *Let $\mathbf{\Lambda}_- \in \mathbb{R}^{n_- \times n_-}$ and $\mathbf{V}_- \in \mathbb{R}^{n \times n_-}$ be the diagonal matrix and the orthogonal matrix that are obtained by removing the null part of \mathbf{A}_- from its diagonalization in such a way that*

$$\mathbf{A}_- = \mathbf{V}_- \mathbf{\Lambda}_- \mathbf{V}_-^\top.$$

Remark 5. *The diagonalization of \mathbf{A}_- is not required in the numerical implementation (see Section 4.4).*

It now holds that $\mathbf{A} = \mathbf{A}_+ - \mathbf{V}_- \mathbf{\Lambda}_- \mathbf{V}_-^\top$ with \mathbf{A} , \mathbf{A}_+ , $\mathbf{\Lambda}_-$ spd matrices and \mathbf{V}_- a full rank matrix. The Woodbury matrix identity [43] applied to computing the inverse of \mathbf{A} gives

$$\mathbf{A}^{-1} = \mathbf{A}_+^{-1} + \mathbf{A}_+^{-1} \mathbf{V}_- (\mathbf{\Lambda}_-^{-1} - \mathbf{V}_-^\top \mathbf{A}_+^{-1} \mathbf{V}_-)^{-1} \mathbf{V}_-^\top \mathbf{A}_+^{-1}. \quad (7)$$

Remark 6. *The Woodbury matrix identity is also called the Sherman-Morrison-Woodbury Formula (e.g., in [18][Section 2.1.4]). The formula is correct since \mathbf{A}_+ is non-singular and $\mathbf{\Lambda}_-^{-1} - \mathbf{V}_-^\top \mathbf{A}_+^{-1} \mathbf{V}_-$ is also non-singular.*

4.2 First Algebraic-Woodbury-GenEO (AWG) preconditioner for \mathbf{A}

The Woodbury matrix identity leads, rather straightforwardly, to a new preconditioner for the original matrix \mathbf{A} that is defined in the following theorem.

Theorem 7 (AWG preconditioner for \mathbf{A} with inexact coarse space). *Given a preconditioner \mathbf{H}_2 for \mathbf{A}_+ such that the eigenvalues of $\mathbf{H}_2 \mathbf{A}_+$ are in the interval $[\lambda_{\min}(\mathbf{H}_2 \mathbf{A}_+), \lambda_{\max}(\mathbf{H}_2 \mathbf{A}_+)]$. Let the new preconditioner for \mathbf{A} be defined as*

$$\mathbf{H}_{3, \text{inex}} := \mathbf{H}_2 + \mathbf{A}_+^{-1} \mathbf{V}_- (\mathbf{\Lambda}_-^{-1} - \mathbf{V}_-^\top \mathbf{A}_+^{-1} \mathbf{V}_-)^{-1} \mathbf{V}_-^\top \mathbf{A}_+^{-1}.$$

The eigenvalues of the new preconditioned operator satisfy

$$\lambda(\mathbf{H}_{3, \text{inex}} \mathbf{A}) \in [\min(1, \lambda_{\min}(\mathbf{H}_2 \mathbf{A}_+)), \max(1, \lambda_{\max}(\mathbf{H}_2 \mathbf{A}_+))]. \quad (8)$$

Proof. The estimate for the eigenvalues of $\mathbf{H}_2\mathbf{A}_+$ is equivalent to

$$\lambda_{\min}(\mathbf{H}_2\mathbf{A}_+)\langle \mathbf{x}, \mathbf{A}_+^{-1}\mathbf{x} \rangle \leq \langle \mathbf{x}, \mathbf{H}_2\mathbf{x} \rangle \leq \lambda_{\max}(\mathbf{H}_2\mathbf{A}_+)\langle \mathbf{x}, \mathbf{A}_+^{-1}\mathbf{x} \rangle, \forall \mathbf{x} \in \mathbb{R}^n.$$

Adding, $\langle \mathbf{x}, \mathbf{A}_+^{-1}\mathbf{V}_- (\mathbf{D}_-^{-1} - \mathbf{V}_-^\top \mathbf{A}_+^{-1}\mathbf{V}_-)^{-1} \mathbf{V}_-^\top \mathbf{A}_+^{-1}\mathbf{x} \rangle$ to each term, it holds that

$$\min(1, \lambda_{\min}(\mathbf{H}_2\mathbf{A}_+))\langle \mathbf{x}, \mathbf{A}_+^{-1}\mathbf{x} \rangle \leq \langle \mathbf{x}, \mathbf{H}_{3, \text{inex}}\mathbf{x} \rangle \leq \max(1, \lambda_{\max}(\mathbf{H}_2\mathbf{A}_+))\langle \mathbf{x}, \mathbf{A}_+^{-1}\mathbf{x} \rangle, \forall \mathbf{x} \in \mathbb{R}^n,$$

where the Woodbury matrix identity (7) was applied. This is equivalent to (8). \square

We have just introduced new preconditioners called the AWG preconditioners with an inexact coarse space. The plural in the previous sentence comes from the fact that there are many possible choices for \mathbf{H}_2 (including the four from Section 3) and that for each one there are parameters that can be adjusted. These new preconditioners are purely algebraic and they have guaranteed spectral bounds when applied to solving linear system $\mathbf{A}\mathbf{x} = \mathbf{b}$. The condition number of $\mathbf{H}_{3, \text{inex}}\mathbf{A}$ can be made smaller by enriching the coarse space in \mathbf{H}_2 . Next, two other AWG preconditioners are defined which have an exact coarse spaces in the sense of the Abstract Schwarz theory: their coarse operator is of the form $\mathbf{R}^0\mathbf{A}\mathbf{R}^{0\top}$.

4.3 Two more AWG preconditioners for \mathbf{A}

Solving a problem with the preconditioner introduced in the previous section requires it to be computationally feasible to compute multiply by $(\mathbf{A}_+^{-1}\mathbf{V}_-)$ and its transpose. Looking at the Woodbury identity, we realize (and prove it in the lemma below) that $\text{range}(\mathbf{A}_+^{-1}\mathbf{V}_-) = \text{range}(\mathbf{A}^{-1}\mathbf{V}_-)$. This opens up new possibilities: if it is possible to compute $\text{range}(\mathbf{A}^{-1}\mathbf{V}_-)$, it is also possible to project \mathbf{A} -orthogonally onto the space that is ℓ_2 -orthogonal to $\text{range}(\mathbf{V}_-)$ which is exactly $\text{Ker}(\mathbf{A}_-)$ (see Π_3 in Definition 11 below). On that space $\mathbf{A}_{|\text{Ker}(\mathbf{A}_-)} = (\mathbf{A}_+)_{|\text{Ker}(\mathbf{A}_-)}$ and we can fall back onto a known and efficient preconditioner \mathbf{H}_2 for \mathbf{A}_+ .

Lemma 2. *With \mathbf{A}_+ from Definition 7 and \mathbf{V}_- from Definition 10, the following property holds*

$$\text{range}(\mathbf{A}_+^{-1}\mathbf{V}_-) = \text{range}(\mathbf{A}^{-1}\mathbf{V}_-).$$

Proof. It follows from the Woodbury identity that

$$\text{range}(\mathbf{A}^{-1}\mathbf{V}_-) = \text{range}\left(\mathbf{A}_+^{-1}\mathbf{V}_-(\mathbf{I} + (\mathbf{A}_-^{-1} - \mathbf{V}_-^\top \mathbf{A}_+^{-1}\mathbf{V}_-)^{-1} \mathbf{V}_-^\top \mathbf{A}_+^{-1}\mathbf{V}_-)\right),$$

where \mathbf{I} is the $n_- \times n_-$ identity matrix. Proving the result in the lemma comes down to proving that the range of $(\mathbf{I} + (\mathbf{A}_-^{-1} - \mathbf{V}_-^\top \mathbf{A}_+^{-1}\mathbf{V}_-)^{-1} \mathbf{V}_-^\top \mathbf{A}_+^{-1}\mathbf{V}_-)$ is the whole of \mathbb{R}^{n_-} or, equivalently, that its kernel is restricted to $\mathbf{0}$ (the zero vector in \mathbb{R}^{n_-}). This last step is achieved as follows. Let $\mathbf{y} \in \mathbb{R}^{n_-}$,

$$\begin{aligned} & \left(\mathbf{I} + (\mathbf{A}_-^{-1} - \mathbf{V}_-^\top \mathbf{A}_+^{-1}\mathbf{V}_-)^{-1} \mathbf{V}_-^\top \mathbf{A}_+^{-1}\mathbf{V}_-\right) \mathbf{y} = \mathbf{0} \\ \Leftrightarrow & (\mathbf{A}_-^{-1} - \mathbf{V}_-^\top \mathbf{A}_+^{-1}\mathbf{V}_-) \mathbf{y} = -\mathbf{V}_-^\top \mathbf{A}_+^{-1}\mathbf{V}_- \mathbf{y} \\ \Leftrightarrow & \mathbf{A}_-^{-1} \mathbf{y} = \mathbf{0} \\ \Leftrightarrow & \mathbf{y} = \mathbf{0}. \end{aligned}$$

\square

Definition 11. Let $\mathbf{W} \in \mathbb{R}^{n \times n}$ be such that $\text{range}(\mathbf{W}) = \text{range}(\mathbf{A}_+^{-1}\mathbf{V}_-)$ and let

$$\mathbf{\Pi}_3 := \mathbf{I} - \mathbf{W}(\mathbf{W}^\top \mathbf{A}\mathbf{W})^{-1}\mathbf{W}^\top \mathbf{A}.$$

Assume that \mathbf{H}_2 is a given preconditioner for \mathbf{A}_+ such that the eigenvalues of $\mathbf{H}_2\mathbf{A}_+$ are in the interval $[\lambda_{\min}(\mathbf{H}_2\mathbf{A}_+), \lambda_{\max}(\mathbf{H}_2\mathbf{A}_+)]$. Let two new preconditioner for \mathbf{A} be defined as

$$\mathbf{H}_{3,ad} := \mathbf{H}_2 + \mathbf{W}(\mathbf{W}^\top \mathbf{A}\mathbf{W})^{-1}\mathbf{W}^\top,$$

and

$$\mathbf{H}_{3,hyb} := \mathbf{\Pi}_3\mathbf{H}_2\mathbf{\Pi}_3^\top + \mathbf{W}(\mathbf{W}^\top \mathbf{A}\mathbf{W})^{-1}\mathbf{W}^\top.$$

Theorem 8. Let $\mathbf{\Pi}_3$, $\mathbf{H}_{3,ad}$ and $\mathbf{H}_{3,hyb}$ be as in Definition 11. The operator $\mathbf{\Pi}_3$ is an \mathbf{A} -orthogonal projection operator that satisfies

$$\text{Ker}(\mathbf{\Pi}_3) = \text{range}(\mathbf{A}^{-1}\mathbf{V}_-) \text{ and } \text{range}(\mathbf{\Pi}_3) = \text{Ker}(\mathbf{A}_-). \quad (9)$$

Moreover, the new preconditioned operators satisfy the spectral bounds :

$$\lambda_{\min}(\mathbf{H}_2\mathbf{A}_+) \leq \lambda(\mathbf{H}_2\mathbf{A}\mathbf{\Pi}_3) \leq \lambda_{\max}(\mathbf{H}_2\mathbf{A}_+) \text{ if } \lambda(\mathbf{H}_2\mathbf{A}\mathbf{\Pi}_3) \neq 0 \quad (10)$$

$$\min(1, \lambda_{\min}(\mathbf{H}_2\mathbf{A}_+)) \leq \lambda(\mathbf{H}_{3,hyb}\mathbf{A}) \leq \max(1, \lambda_{\max}(\mathbf{H}_2\mathbf{A}_+)) \quad (11)$$

$$\min(1, \lambda_{\min}(\mathbf{H}_2\mathbf{A}_+)) \leq \lambda(\mathbf{H}_{3,ad}\mathbf{A}) \leq (\lambda_{\max}(\mathbf{H}_2\mathbf{A}_+) + 1). \quad (12)$$

where we recall that \mathbf{H}_2 can be chosen as one of the two level preconditioners from Section ?? in such a way that $\lambda_{\min}(\mathbf{H}_2\mathbf{A}_+)$ and $\lambda_{\max}(\mathbf{H}_2\mathbf{A}_+)$ are known and controlled by the choice of the coarse space.

Note that a bound (10) for the projected and preconditioned operator has been included in the theorem (equation (??)).

Proof. We begin by proving (9). Let $\mathbf{x} \in \mathbb{R}^n$, \mathbf{x} is in the kernel of $\mathbf{\Pi}_3$ if:

$$\mathbf{\Pi}_3\mathbf{x} = 0 \Leftrightarrow \mathbf{x} = \mathbf{W}(\mathbf{W}^\top \mathbf{A}\mathbf{W})^{-1}\mathbf{W}^\top \mathbf{A}\mathbf{x} \Leftrightarrow \mathbf{x} \in \text{range}(\mathbf{W}) = \text{range}(\mathbf{A}_+^{-1}\mathbf{V}_-) = \text{range}(\mathbf{A}^{-1}\mathbf{V}_-).$$

The last equality comes from Lemma 2. Moreover, $\mathbf{\Pi}_3$ is an \mathbf{A} -orthogonal projection so

$$\text{range}(\mathbf{\Pi}_3) = (\text{Ker}(\mathbf{\Pi}_3))^{\perp \mathbf{A}} = (\text{range}(\mathbf{A}^{-1}\mathbf{V}_-))^{\perp \mathbf{A}} = (\text{range}(\mathbf{V}_-))^{\perp \ell_2} = \text{Ker}(\mathbf{V}_-^\top) = \text{Ker}(\mathbf{A}_-),$$

by definition of \mathbf{V}_- in Definition 10. A direct consequence of this result, that is frequently used in the remainder of the proof, is the identity $\mathbf{A}_+\mathbf{\Pi}_3 = \mathbf{A}\mathbf{\Pi}_3$.

We now move onto proving the spectral bounds starting with (10) for the projected and preconditioned operator $\mathbf{H}_2\mathbf{A}\mathbf{\Pi}_3$. Let $(\lambda, \mathbf{y}) \in \mathbb{R} \times \mathbb{R}^n$ be an eigenpair of the matrix $\mathbf{H}_2\mathbf{A}\mathbf{\Pi}_3 (= \mathbf{H}_2\mathbf{A}_+\mathbf{\Pi}_3)$ meaning that:

$$\mathbf{y} \neq \mathbf{0} \text{ and } \mathbf{H}_2\mathbf{A}_+\mathbf{\Pi}_3\mathbf{y} = \lambda\mathbf{y}.$$

Taking the inner product by $\mathbf{A}_+\mathbf{\Pi}_3\mathbf{y}$ gives

$$\langle \mathbf{A}_+\mathbf{\Pi}_3\mathbf{y}, \mathbf{H}_2\mathbf{A}_+\mathbf{\Pi}_3\mathbf{y} \rangle = \lambda \langle \mathbf{A}_+\mathbf{\Pi}_3\mathbf{y}, \mathbf{y} \rangle = \lambda \langle \mathbf{A}\mathbf{\Pi}_3\mathbf{y}, \mathbf{y} \rangle = \lambda \langle \mathbf{A}\mathbf{\Pi}_3\mathbf{y}, \mathbf{\Pi}_3\mathbf{y} \rangle = \lambda \langle \mathbf{A}_+\mathbf{\Pi}_3\mathbf{y}, \mathbf{\Pi}_3\mathbf{y} \rangle.$$

Moreover, since \mathbf{A}_+ and \mathbf{H}_2 are spd, the spectral bound for $\mathbf{H}_2\mathbf{A}_+$ is equivalent to

$$\lambda_{\min}(\mathbf{H}_2\mathbf{A}_+) \langle \mathbf{x}, \mathbf{A}_+\mathbf{x} \rangle \leq \langle \mathbf{x}, \mathbf{A}_+\mathbf{H}_2\mathbf{A}_+\mathbf{x} \rangle \leq \lambda_{\max}(\mathbf{H}_2\mathbf{A}_+) \langle \mathbf{x}, \mathbf{A}_+\mathbf{x} \rangle, \quad \forall \mathbf{x} \in \mathbb{R}^n. \quad (13)$$

In particular, for $\mathbf{x} = \mathbf{\Pi}_3\mathbf{y}$ we get

$$\lambda_{\min}(\mathbf{H}_2\mathbf{A}_+) \langle \mathbf{\Pi}_3\mathbf{y}, \mathbf{A}_+\mathbf{\Pi}_3\mathbf{y} \rangle \leq \underbrace{\langle \mathbf{\Pi}_3\mathbf{y}, \mathbf{A}_+\mathbf{H}_2\mathbf{A}_+\mathbf{\Pi}_3\mathbf{y} \rangle}_{=\lambda \langle \mathbf{A}_+\mathbf{\Pi}_3\mathbf{y}, \mathbf{\Pi}_3\mathbf{y} \rangle} \leq \lambda_{\max}(\mathbf{H}_2\mathbf{A}_+) \langle \mathbf{\Pi}_3\mathbf{y}, \mathbf{A}_+\mathbf{\Pi}_3\mathbf{y} \rangle.$$

Finally, there are two possibilities, either $\langle \mathbf{A}_+ \mathbf{\Pi}_3 \mathbf{y}, \mathbf{\Pi}_3 \mathbf{y} \rangle = 0$, so $\mathbf{y} \in \text{Ker}(\mathbf{\Pi}_3)$ and $\lambda = 0$, or $\langle \mathbf{A}_+ \mathbf{\Pi}_3 \mathbf{y}, \mathbf{\Pi}_3 \mathbf{y} \rangle \neq 0$ and $\lambda \in [\lambda_{\min}(\mathbf{H}_2 \mathbf{A}_+), \lambda_{\max}(\mathbf{H}_2 \mathbf{A}_+)]$. In other words (10) holds.

Next, we prove the spectral bound with the hybrid preconditioner $\mathbf{H}_{3,hyb}$ from (11). Let $\mathbf{x} \in \mathbb{R}^n$, we add the term

$$\langle \mathbf{x}, \mathbf{A} \mathbf{W} (\mathbf{W}^\top \mathbf{A} \mathbf{W})^{-1} \mathbf{W}^\top \mathbf{A} \mathbf{x} \rangle = \langle \mathbf{x}, \mathbf{A} (\mathbf{I} - \mathbf{\Pi}_3) \mathbf{x} \rangle = \langle (\mathbf{I} - \mathbf{\Pi}_3) \mathbf{x}, \mathbf{A} (\mathbf{I} - \mathbf{\Pi}_3) \mathbf{x} \rangle$$

to estimate (13) evaluated at $\mathbf{\Pi}_3 \mathbf{x}$ and obtain

$$\begin{aligned} \lambda_{\min}(\mathbf{H}_2 \mathbf{A}_+) \langle \mathbf{\Pi}_3 \mathbf{x}, \mathbf{A} \mathbf{\Pi}_3 \mathbf{x} \rangle + \langle (\mathbf{I} - \mathbf{\Pi}_3) \mathbf{x}, \mathbf{A} (\mathbf{I} - \mathbf{\Pi}_3) \mathbf{x} \rangle &\leq \\ \langle \mathbf{\Pi}_3 \mathbf{x}, \mathbf{A} \mathbf{H}_2 \mathbf{A} \mathbf{\Pi}_3 \mathbf{x} \rangle + \langle \mathbf{x}, \mathbf{A} \mathbf{W} (\mathbf{W}^\top \mathbf{A} \mathbf{W})^{-1} \mathbf{W}^\top \mathbf{A} \mathbf{x} \rangle &\leq \\ \lambda_{\max}(\mathbf{H}_2 \mathbf{A}_+) \langle \mathbf{\Pi}_3 \mathbf{x}, \mathbf{A} \mathbf{\Pi}_3 \mathbf{x} \rangle + \langle (\mathbf{I} - \mathbf{\Pi}_3) \mathbf{x}, \mathbf{A} (\mathbf{I} - \mathbf{\Pi}_3) \mathbf{x} \rangle & \end{aligned}$$

where $\mathbf{A}_+ \mathbf{\Pi}_3 = \mathbf{A} \mathbf{\Pi}_3$ as also been applied. This then implies that

$$\min(1, \lambda_{\min}(\mathbf{H}_2 \mathbf{A}_+)) \langle \mathbf{x}, \mathbf{A} \mathbf{x} \rangle \leq \langle \mathbf{x}, \mathbf{A} \mathbf{H}_{3,hyb} \mathbf{A} \mathbf{x} \rangle \leq \max(1, \lambda_{\max}(\mathbf{H}_2 \mathbf{A}_+)) \langle \mathbf{x}, \mathbf{A} \mathbf{x} \rangle.$$

and the eigenvalue estimate in the theorem holds because \mathbf{A} and $\mathbf{H}_{3,hyb}$ are spd.

Finally, we prove the spectral bound with the additive preconditioner $\mathbf{H}_{3,ad}$ from (11). Matrices \mathbf{A}_+ and \mathbf{H}_2 are both spd so the fact that all eigenvalues are not greater than $\lambda_{\max}(\mathbf{H}_2 \mathbf{A}_+)$ is equivalent to

$$\langle \mathbf{x}, \mathbf{H}_2 \mathbf{x} \rangle \leq \lambda_{\max}(\mathbf{H}_2 \mathbf{A}_+) \langle \mathbf{x}, \mathbf{A}_+^{-1} \mathbf{x} \rangle, \text{ for any } \mathbf{x} \in \mathbb{R}^n.$$

Moreover, $\langle \mathbf{x}, \mathbf{A}_+^{-1} \mathbf{x} \rangle \leq \langle \mathbf{x}, \mathbf{A}^{-1} \mathbf{x} \rangle$ for any $\mathbf{x} \in \mathbb{R}^n$ so

$$\langle \mathbf{x}, \mathbf{H}_2 \mathbf{x} \rangle \leq \lambda_{\max}(\mathbf{H}_2 \mathbf{A}_+) \langle \mathbf{x}, \mathbf{A}^{-1} \mathbf{x} \rangle, \text{ for any } \mathbf{x} \in \mathbb{R}^n.$$

It also holds that

$$\langle \mathbf{x}, \mathbf{A} \mathbf{W} (\mathbf{W}^\top \mathbf{A} \mathbf{W})^{-1} \mathbf{W}^\top \mathbf{A} \mathbf{x} \rangle = \langle (\mathbf{I} - \mathbf{\Pi}_3) \mathbf{x}, \mathbf{A} (\mathbf{I} - \mathbf{\Pi}_3) \mathbf{x} \rangle \leq \langle \mathbf{x}, \mathbf{A} \mathbf{x} \rangle, \text{ for any } \mathbf{x} \in \mathbb{R}^n.$$

Adding the last two results together gives us

$$\langle \mathbf{x}, \mathbf{H}_{3,ad} \mathbf{x} \rangle \leq (\lambda_{\max}(\mathbf{H}_2 \mathbf{A}_+) + 1) \langle \mathbf{x}, \mathbf{A}^{-1} \mathbf{x} \rangle, \text{ for any } \mathbf{x} \in \mathbb{R}^n,$$

or in other words, all eigenvalues of $\mathbf{H}_{3,ad} \mathbf{A}$ are not greater than $(\lambda_{\max}(\mathbf{H}_2 \mathbf{A}_+) + 1)$. For the smallest eigenvalue of $\mathbf{H}_{3,ad} \mathbf{A}$, we can look at $\mathbf{H}_{3,ad}$ in the abstract Schwarz framework. Indeed

$$\mathbf{H}_{3,ad} = \mathbf{I} \mathbf{H}_2 \mathbf{I} + \mathbf{W} (\mathbf{W}^\top \mathbf{A} \mathbf{W})^{-1} \mathbf{W}^\top,$$

is an abstract Schwarz solver for 2 subspaces \mathbb{R}^n and \mathbb{R}^{n-} with prolongation operators \mathbf{I} (identity matrix in \mathbb{R}^n) and \mathbf{W} , and with local solvers \mathbf{H}_2 and $(\mathbf{W}^\top \mathbf{A} \mathbf{W})^{-1}$. We know that \mathbf{H}_2 is spd so the classical stable splitting result from Toselli Widlund [41] applies (or from [36]): all eigenvalues of $\mathbf{H}_{3,ad} \mathbf{A}$ are larger than C_0^{-2} if, for any $\mathbf{x} \in \mathbb{R}^n$, there exist $\mathbf{z}_+ \in \mathbb{R}^n$ and $\mathbf{z}_- \in \mathbb{R}^{n-}$ that satisfy

$$\mathbf{z}_+ + \mathbf{W} \mathbf{z}_- = \mathbf{x} \text{ and } \langle \mathbf{z}_+, \mathbf{H}_2^{-1} \mathbf{z}_+ \rangle + \langle \mathbf{z}_-, \mathbf{W}^\top \mathbf{A} \mathbf{W} \mathbf{z}_- \rangle \leq C_0^2 \langle \mathbf{x}, \mathbf{A} \mathbf{x} \rangle.$$

The following splitting is proposed: $\mathbf{z}_+ = \mathbf{\Pi}_3 \mathbf{x}$ and $\mathbf{z}_- = (\mathbf{W}^\top \mathbf{A} \mathbf{W})^{-1} \mathbf{W}^\top \mathbf{A} \mathbf{x}$. We first check that they do split \mathbf{x} :

$$\mathbf{z}_+ + \mathbf{W} \mathbf{z}_- = (\mathbf{I} - \mathbf{\Pi}_3) \mathbf{x} + \mathbf{W} (\mathbf{W}^\top \mathbf{A} \mathbf{W})^{-1} \mathbf{W}^\top \mathbf{A} \mathbf{x} = \mathbf{x}.$$

We then check the stability of the splitting:

$$\begin{aligned} \langle \mathbf{z}_+, \mathbf{H}_2^{-1} \mathbf{z}_+ \rangle + \langle \mathbf{z}_-, \mathbf{W}^\top \mathbf{A} \mathbf{W} \mathbf{z}_- \rangle &= \langle \mathbf{\Pi}_3 \mathbf{x}, \mathbf{H}_2^{-1} \mathbf{\Pi}_3 \mathbf{x} \rangle + \langle (\mathbf{W}^\top \mathbf{A} \mathbf{W})^{-1} \mathbf{W}^\top \mathbf{A} \mathbf{x}, \mathbf{W}^\top \mathbf{A} \mathbf{W} (\mathbf{W}^\top \mathbf{A} \mathbf{W})^{-1} \mathbf{W}^\top \mathbf{A} \mathbf{x} \rangle \\ &\leq (\lambda_{\min}(\mathbf{H}_2 \mathbf{A}_+))^{-1} \langle \mathbf{\Pi}_3 \mathbf{x}, \mathbf{A}_+ \mathbf{\Pi}_3 \mathbf{x} \rangle + \langle (\mathbf{I} - \mathbf{\Pi}_3) \mathbf{x}, \mathbf{A} (\mathbf{I} - \mathbf{\Pi}_3) \mathbf{x} \rangle \\ &\leq (\lambda_{\min}(\mathbf{H}_2 \mathbf{A}_+))^{-1} \langle \mathbf{\Pi}_3 \mathbf{x}, \mathbf{A} \mathbf{\Pi}_3 \mathbf{x} \rangle + \langle (\mathbf{I} - \mathbf{\Pi}_3) \mathbf{x}, \mathbf{A} (\mathbf{I} - \mathbf{\Pi}_3) \mathbf{x} \rangle \\ &\leq \max(1, \lambda_{\min}(\mathbf{H}_2 \mathbf{A}_+)^{-1}) \langle \mathbf{x}, \mathbf{A} \mathbf{x} \rangle. \end{aligned}$$

Finally, we have proved that all eigenvalues of $\mathbf{H}_{3,ad}$ are greater than or equal to $\min(1, \lambda_{\min}(\mathbf{H}_2 \mathbf{A}_+))$. \square

Again, the preconditioners $\mathbf{H}_{3,ad}$ and $\mathbf{H}_{3,hyb}$ are families of preconditioners that are computed algebraically and lead to guaranteed spectral bounds when applied to solving $\mathbf{A}\mathbf{x} = \mathbf{b}$. The condition numbers can be made smaller by enriching the coarse space in \mathbf{H}_2 .

4.4 Remarks on the implementation of the new preconditioners

Below, some important remarks are made about the implementation of $\mathbf{H}_{3,ad}$ and $\mathbf{H}_{3,hyb}$.

1. It is not necessary to factorize \mathbf{A}_- as suggested by the definition of \mathbf{V}_- in Definition 10. Indeed, Definition 11 requires only a basis \mathbf{W} of $\text{range}(\mathbf{A}_+^{-1} \mathbf{V}_-)$ to generate the second coarse space. A natural choice is to recall that \mathbf{V}_- is generated by the eigenvectors of \mathbf{B}^s that correspond to negative eigenvalues (once prolonged to Ω by $\mathbf{R}^{s\top}$). If the matrices \mathbf{B}^s are non-singular then this is the same as computing

$$\mathbf{W} = \mathbf{A}_+^{-1} [\mathbf{R}^{1\top} \mathbf{V}_-^1 \mid \dots \mid \mathbf{R}^{N\top} \mathbf{V}_-^N],$$

with the \mathbf{V}_-^s from Definition 6. If the \mathbf{B}^s are singular, it is necessary to first remove the columns in \mathbf{V}_-^s that correspond to zero eigenvalues. It may also be necessary (although we haven't observed it in practice) to remove some linear dependencies between the columns. This is rather standard and can be done either when computing \mathbf{W} or when factorizing the coarse problem $\mathbf{W}^\top \mathbf{A} \mathbf{W}$.

2. The computation of \mathbf{W} is one of the bottlenecks of the algorithm: many systems must be solved for the global matrix \mathbf{A}_+ . In our current implementation these linear systems are solved one after the other with PCG preconditioned by \mathbf{H}_2 . Since \mathbf{H}_2 is a *good* preconditioner for \mathbf{A}_+ this takes *few* iterations. It must be explored whether computational efficiency could be improved with block CG methods [30] or adaptive multipreconditioning [35].
3. All four choices of preconditioners \mathbf{H}_2 for \mathbf{A}_+ that are considered in the article require that the action of $(\mathbf{A}_+^s)^\dagger$ be implemented in order to compute the corresponding GenEO coarse space. Instead of computing the full diagonalization of \mathbf{B}^s , it is sufficient to compute its negative eigenvalues and corresponding eigenvectors (*i.e.*, \mathbf{A}_-^s and \mathbf{V}_-^s) and to recall that

$$\mathbf{A}_+^s = (\mathbf{I} - \mathbf{V}_-^s \mathbf{V}_-^{s\top}) \mathbf{B}^s (\mathbf{I} - \mathbf{V}_-^s \mathbf{V}_-^{s\top})$$

which also implies that

$$\mathbf{A}_+^{s\dagger} = (\mathbf{I} - \mathbf{V}_-^s \mathbf{V}_-^{s\top}) \mathbf{B}^{s\dagger} (\mathbf{I} - \mathbf{V}_-^s \mathbf{V}_-^{s\top}).$$

Since \mathbf{B}^s is symmetric, it can be factorized using MUMPS [3, 2].

4. All four choices of preconditioners \mathbf{H}_2 for \mathbf{A}_+ that are considered in the article require that the action of $\mathbf{R}^s \mathbf{A}_+ \mathbf{R}^{s\top}$ be implemented in order to compute the corresponding GenEO coarse space. As $\mathbf{R}^s \mathbf{A}_+ \mathbf{R}^{s\top}$ is a dense matrix, it is never assembled. Instead, the action of $\mathbf{R}^s \mathbf{A}_+ \mathbf{R}^{s\top}$ is computed as

$$\mathbf{R}^s \mathbf{A}_+ \mathbf{R}^{s\top} = \mathbf{R}^s \mathbf{A} \mathbf{R}^{s\top} + \mathbf{R}^s \mathbf{A}_- \mathbf{R}^{s\top} = \mathbf{R}^s \mathbf{A} \mathbf{R}^{s\top} - \sum_{t=1}^N \mathbf{R}^s \mathbf{R}^{t\top} \mathbf{V}_-^t \mathbf{A}_-^t \mathbf{V}_-^{t\top} \mathbf{R}^t \mathbf{R}^{s\top}.$$

In the sum, all terms for which $\mathbf{R}^t \mathbf{R}^{s\top}$ is zero are zero.

5. If $\mathbf{H}_+^{\text{AS}} := \sum_{s=1}^N \mathbf{R}^{s\top} (\mathbf{R}^s \mathbf{A}_+ \mathbf{R}^{s\top})^{-1} \mathbf{R}^s$ is chosen as a one-level preconditioner for \mathbf{A}_+ then it is necessary to compute the action of $(\mathbf{R}^s \mathbf{A}_+ \mathbf{R}^{s\top})^{-1}$. This is done by applying the Woodbury matrix identity to the formula just above.

5 Numerical Results

In this section, numerical results are presented for the new AWG preconditioners with exact coarse spaces: $\mathbf{H}_{3,ad}$ and $\mathbf{H}_{3,hyb}$. The theoretical convergence bounds are checked, and the behaviour of the new preconditioners is illustrated for the first time. Some comparisons to non-algebraic domain decomposition preconditioners with standard or GenEO coarse spaces are performed. The linear systems that are considered result from discretizing a two-dimensional linear elasticity problem with Q_1 finite elements. All details are given below.

All the results presented were obtained with petsc4py [12], a Python port to the PETSc libraries [4, 5, 6]. The eigensolves are performed by SLEPc [22] and the matrix factorizations (for the local and coarse problems) are performed by MUMPS [3, 2]. Our code is available on github [19].

Let $\omega = [0, 3] \times [0, 3] \subset \mathbb{R}^2$ be the computational domain. Let $\partial\omega_D$ be the left hand side boundary of ω and let $\mathcal{V} = \{\mathbf{v} \in H^1(\omega)^2; \mathbf{v} = \mathbf{0} \text{ on } \partial\omega_D\}$. A solution $\mathbf{u} \in \mathcal{V}$ is sought such that

$$\int_{\omega} 2\mu\varepsilon(\mathbf{u}) : \varepsilon(\mathbf{v}) dx + \int_{\omega} L \operatorname{div}(\mathbf{u}) \operatorname{div}(\mathbf{v}) dx = \int_{\omega} \mathbf{g} \cdot \mathbf{v} dx, \text{ for all } \mathbf{v} \in \mathcal{V}, \quad (14)$$

where, for $i, j = 1, 2$, $\varepsilon_{ij}(\mathbf{u}) = \frac{1}{2} \left(\frac{\partial u_i}{\partial x_j} + \frac{\partial u_j}{\partial x_i} \right)$, δ_{ij} is the Kronecker symbol, $\mathbf{g} = (0, -9.81)^\top$ and the Lamé coefficients are functions of Young's modulus E and Poisson's ratio ν : $\mu = \frac{E}{2(1+\nu)}$, $L = \frac{E\nu}{(1+\nu)(1-2\nu)}$. It is well known (see, *e.g.*, [32]) that the solution of (14) in a heterogeneous medium is challenging due to ill-conditioning. Unless otherwise specified, the coefficient distribution that is considered here is the following: for any $(x, y) \in \omega$,

$$\nu(x, y) = 0.3 \quad \text{and} \quad E(x, y) = \begin{cases} 10^{11} & \text{if } (\text{floor}(y) - y) \in [1/7, 2/7] \cup [3/7, 4/7] \cup [5/7, 6/7] \\ 10^7 & \text{otherwise} \end{cases}$$

This is represented in Figure 1.

The computational domain is discretized by a uniform mesh with element size $h = 1/21$ so there are

$$n = 8192 \text{ degrees of freedom.}$$

The boundary value problem is solved numerically with Q_1 finite elements. Let \mathcal{V}_h be the space of Q_1 finite elements that satisfy the Dirichlet boundary condition. Let $\{\phi_k\}_{k=1}^n$ be a basis of \mathcal{V}_h . The linear system that is to be solved is

$$\text{Find } \mathbf{x}_* \in \mathbb{R}^n \text{ such that } \mathbf{A}\mathbf{x}_* = \mathbf{b},$$

with $A_{ij} = \int_{\omega} [2\mu\varepsilon(\phi_i) : \varepsilon(\phi_j) + L \operatorname{div}(\phi_i) \operatorname{div}(\phi_j)] dx$ and $\mathbf{b}_i = \int_{\omega} \mathbf{g} \cdot \phi_i dx$.

Unless specified otherwise, for each computation, the domain ω is split into

$$N = 9$$

unit square subdomains that overlap only at the interface and this in turn gives the partition of the degrees of freedom into $\Omega^1, \dots, \Omega^9$. With Q_1 finite elements, for the overlap to be restricted to the shared subdomain boundaries is enough to ensure the minimal overlap condition. There are 504 degrees of freedom that are shared by more than one subdomain. All linear systems, are solved with PCG up to a relative residual tolerance of 10^{-10} . The preconditioner is specified for each test case.

The matrix $\mathbf{A}_+^{-1}\mathbf{V}_-$ is computed by solving n_- linear systems for \mathbf{A}_+^{-1} . This is also done with PCG preconditioned by \mathbf{H}_2 up to a relative residual tolerance of 10^{-10} (unless specified otherwise).

Unless it is otherwise specified (*e.g.*, in the next paragraph), the AWG preconditioner under study is

$$\mathbf{H}_{3,ad} \text{ with } \mathbf{H}_2 = \mathbf{H}_{hyb}^{\text{NN}}(\tau_{\#}).$$




Figure 1: Solution: ω has been displaced by 10^5 multiplied by the solution to the PDE. The colors show the distribution of coefficients. Dark: $E = E_1 = 10^{11}$ - Light : $E = E_2 = 10^7$

Comparison of $\mathbf{H}_{3,ad}$ and $\mathbf{H}_{3,hyb}$ for all variants of \mathbf{H}_2 The test case is solved with the eight AWG preconditioners (for fixed values of threshold). Specifically, there are two ways of incorporating the second coarse space leading to $\mathbf{H}_{3,ad}$ (additive) and $\mathbf{H}_{3,hyb}$ (hybrid) as well as, for each one, four choices for \mathbf{H}_2 : $\mathbf{H}_{hyb}^{NN}(\tau_{\sharp})$, $\mathbf{H}_{hyb}^{AS}(\tau_{\sharp}, \tau_{\flat})$, $\mathbf{H}_{+,ad}^{AS}(\tau_{\flat})$ and $\mathbf{H}_{+,hyb}^{AS}(\tau_{\flat})$. The values for selecting eigenvalues in the GenEO coarse spaces are set to $\tau_{\flat} = 10$ and $\tau_{\sharp} = 0.1$. The results are shown in Table 1. Results for classical (non algebraic) domain decomposition preconditioners with GenEO coarse spaces are incorporated too as a matter of comparison. The quantity n_- (size of the second coarse space) is the same for all AWG preconditioners as it depends only on \mathbf{A} . The size of the GenEO coarse space is also the same for all choices of \mathbf{H}_2 . For $\mathbf{H}_{hyb}^{NN}(\tau_{\sharp})$, $\mathbf{H}_{hyb}^{AS}(\tau_{\sharp}, \tau_{\flat})$, $\mathbf{H}_{+,ad}^{AS}(\tau_{\flat})$ it is in fact the same eigenvalue problem being solved so as long as $\tau_{\sharp} = \tau_{\flat}^{-1}$ this was expected. For the last choice $\mathbf{H}_2 = \mathbf{H}_{hyb}^{AS}(\tau_{\sharp}, \tau_{\flat})$, it is not entirely surprising that the size of the coarse space is not too different as there are connections between the GenEO eigenproblems.

For all AWG preconditioners, the extreme eigenvalues of the preconditioned operators behave as predicted. All AWG preconditioners reduce the condition number of the preconditioned operator to a very small value below 20. The GenEO coarse spaces constructed by AWG are of almost the same size as the classical (non GenEO) coarse spaces which is very satisfying. Of course the AWG preconditioners bear the cost of the extra coarse space.

For $\mathbf{H}_2 = \mathbf{H}_{hyb}^{NN}(\tau_{\sharp})$ the first 20 eigenvalues that are computed in each subdomain for the GenEO eigenproblem in \mathbf{H}_2 are plotted in Figure 2. It appears that choosing larger values of τ_{\sharp} could increase significantly the size of the coarse space without improving the condition number much as the eigenvalues are quite clustered. The eigenvectors that are selected for the coarse space are plotted in Figure 3. Since the unknowns are displacements, they have been represented by applying the deformation to the subdomain. The colors show the values of E . The influence of the hard layers can be seen but it is not easy to make any conclusions about these.

In all that follows we focus on the choice of preconditioner:

$$\mathbf{H}_{3,ad} \text{ with } \mathbf{H}_2 = \mathbf{H}_{hyb}^{NN}(\tau_{\sharp}).$$

	κ	It	λ_{\min}	λ_{\max}	$\#V_0$	n_-
New AWG preconditioners:						
$\mathbf{H}_{3,ad}$ with $\mathbf{H}_2 = \mathbf{H}_{hyb}^{NN}(0.1)$	9.09	26	1.0	9.1	57	48
$\mathbf{H}_{3,ad}$ with $\mathbf{H}_2 = \mathbf{H}_{hyb}^{AS}(0.1, 10)$	12.2	26	0.33	4.0	57	48
$\mathbf{H}_{3,ad}$ with $\mathbf{H}_2 = \mathbf{H}_{+,hyb}^{AS}(10)$	12.3	25	0.33	4.0	57	48
$\mathbf{H}_{3,ad}$ with $\mathbf{H}_2 = \mathbf{H}_{+,ad}^{AS}(10)$	16.8	31	0.24	4.0	57	48
$\mathbf{H}_{3,hyb}$ with $\mathbf{H}_2 = \mathbf{H}_{hyb}^{NN}(0.1)$	9.09	27	1.0	9.1	57	48
$\mathbf{H}_{3,hyb}$ with $\mathbf{H}_2 = \mathbf{H}_{hyb}^{AS}(0.1, 10)$	12.1	25	0.33	4.0	57	48
$\mathbf{H}_{3,hyb}$ with $\mathbf{H}_2 = \mathbf{H}_{+,hyb}^{AS}(10)$	12.2	25	0.33	4.0	57	48
$\mathbf{H}_{3,hyb}$ with $\mathbf{H}_2 = \mathbf{H}_{+,ad}^{AS}(10)$	16.7	29	0.24	4.0	57	48
Non-algebraic methods:						
Hybrid AS + GenEO ($\tau = 10$)	26.5	43	0.15	4.0	55	0
Additive AS + GenEO ($\tau = 10$)	50.0	58	0.080	4.0	55	0
BNN with GenEO ($\tau = 0.1$)	11.1	29	1.0	11.1	55	0
One-level AS	34772	151	0.000115	4.0	0	0

Table 1: Comparison between all AWG preconditioners for fixed τ_{\sharp} and τ_{\flat} . Data is also included for classical GenEO and the one-level method. κ : condition number of preconditioned operator, It : number of iterations, λ_{\min} : smallest eigenvalue of preconditioned operator, λ_{\max} : largest eigenvalue of preconditioned operator, $\#V_0$: dimension of GenEO coarse space, $n_- = \text{rank}(\mathbf{A}_-)$: dimension of second coarse space.




Figure 2: For $\mathbf{H}_2 = \mathbf{H}_{hyb}^{NN}(\tau_{\sharp})$: 20 smallest eigenvalues of the GenEO eigenproblem in each subdomain




Figure 3: For $\mathbf{H}_2 = \mathbf{H}_{hyb}^{\text{NN}}(\tau_{\sharp})$: plot of the 9 eigenvalues that are selected in the central subdomain represented as deformations. The colors correspond to the two values of E .




Figure 4: Condition number with respect to coarse space size for $\mathbf{H}_{3,Ad}$ with $\mathbf{H}_2 = \mathbf{H}_{hyb}^{NN}(\tau_{\sharp})$ and the comparable classical GenEO coarse space. $\tau_{\sharp} \in [0, 0.001, 0.01, 0.05, 0.1, 0.2, 0.5]$. Left: $\nu = 0.3$. Right: $\nu = 0.4$.

With this choice of \mathbf{H}_2 , the theory predicts that the smallest eigenvalue of the preconditioned operator is larger than 1 and this bound is observed to be sharp. For this reason we no longer report on the extreme only eigenvalues. Instead we only give values of the condition number κ .

Influence of the threshold τ_{\sharp} For this test we study the influence of τ_{\sharp} . When τ_{\sharp} increases, more vectors are selected for the coarse space and the condition number bound decreases. AWG is compared to classical Neumann Neumann GenEO (which is not algebraic). Figure 4 is a plot of the condition number of the preconditioned operator versus the size of the GenEO coarse space. Recall that GenEO has the disadvantage of not being algebraic but the AWG method has the disadvantage of having a second coarse space (of size 48). For AWG, the size of the coarse space cannot go below n_- . The study was performed by running the simulations for $\tau_{\sharp} \in [0, 0.001, 0.01, 0.05, 0.1, 0.2, 0.5]$. Two values of ν are considered $\nu = 0.3$ and $\nu = 0.4$. For $\nu = 0.3$ the AWG coarse space to achieve a condition number of 10 is almost the same as with classical GenEO. When $\nu = 0.4$ more vectors are required for AWG.

Influence of ν The value of ν varies between 0.2 and 0.49. Young's modulus is constant in the domain and $E = 10^{11}$. The threshold is $\tau_{\sharp} = 0.05$ (so slightly smaller than previously). As shown in Table 2, increasing ν has quite a dramatic effect on n_- (even without going near the incompressible limit). This is rather disappointing. The classical GenEO does not suffer from this problem (away from the incompressible limit $\nu \rightarrow 0.5$).

Influence of E The threshold is set back to $\tau_{\sharp} = 0.1$ in all that follows. This time the values E_1 and E_2 of Young's modulus in the dark and light parts of ω represented in Figure 1 are varied. The results are in Table 3

Influence of the accuracy of \mathbf{W} The second coarse space for the AWG preconditioners is computed by solving n_- linear systems: $\mathbf{A}_+^{-1} \mathbf{R}^{s\top} \mathbf{v}^s$ for all vectors \mathbf{v}^s that correspond to a negative eigenvalue of \mathbf{B}^s and all subdomains. Until now, we have solved with very high accuracy: the relative residual tolerance $rtol$ was set to 10^{-10} . In this experiment we vary $rtol$. Table 4

AWG					Classical GenEO				
ν	κ	It	$\#V_0$	n_-	ν	κ	It	$\#V_0$	n_-
0.20	19.7	33	21	12	0.20	17.2	33	21	0
0.30	20.3	32	29	19	0.30	17.6	36	21	0
0.35	18.6	32	47	25	0.35	19.1	37	21	0
0.40	25.8	39	98	70	0.40	20.1	39	24	0
0.45	27.1	29	115	110	0.45	33.7	46	28	0
0.49	16.8	25	362	357	0.49	34.9	51	94	0

Table 2: The influence of Poisson’s ratio ν is studied when E is constant and equal to 10^{11} . The threshold is $\tau_{\sharp} = 0.05$. ν : Poisson’s ratio, κ : condition number of preconditioned operator, It : number of iterations, $\#V_0$: dimension of GenEO coarse space, $n_- = \text{rank}(\mathbf{A}_-)$: dimension of second coarse space. Recall that classical GenEO is not algebraic.

AWG					Classical GenEO				
(E_1, E_2)	κ	It	$\#V_0$	n_-	(E_1, E_2)	κ	It	$\#V_0$	n_-
$(10^5, 10^{11})$	10.8	22	95	75	$(10^5, 10^{11})$	8.6	23	90	0
$(10^7, 10^{11})$	10.8	23	95	75	$(10^7, 10^{11})$	8.6	26	87	0
$(10^9, 10^{11})$	10.4	24	94	73	$(10^9, 10^{11})$	8.5	25	85	0
$(10^{11}, 10^{11})$	12.2	29	35	19	$(10^{11}, 10^9)$	11.2	30	52	0
$(10^{11}, 10^9)$	8.0	26	59	48	$(10^{11}, 10^{11})$	13.7	32	28	0
$(10^{11}, 10^7)$	9.0	26	57	48	$(10^{11}, 10^7)$	11.1	29	55	0
$(10^{11}, 10^5)$	8.4	29	57	48	$(10^{11}, 10^5)$	12.7	30	55	0

Table 3: The influence of E and of the jump between E_1 and E_2 is studied. (E_1, E_2) : values of Young’s modulus in the layers of coefficients, κ : condition number of preconditioned operator, It : number of iterations, $\#V_0$: dimension of GenEO coarse space, $n_- = \text{rank}(\mathbf{A}_-)$: dimension of second coarse space. Recall that classical GenEO is not algebraic.

$\nu = 0.3$					$\nu = 0.4$				
$rtol$	κ	It	$\#V_0$	n_-	$rtol$	κ	It	$\#V_0$	n_-
10^{-10}	9.0	26	57	48	10^{-10}	9.4	29	100	74
10^{-2}	9.0	27	57	48	10^{-2}	9.4	30	100	74
0.05	11.1	31	57	48	0.05	12.0	33	100	74
0.1	12.2	32	57	48	0.1	17.4	36	100	74
0.5	400.8	40	57	48	0.5	1563.3	88	100	74
0.9	706.8	64	57	48	0.9	2142.1	100	100	74

Table 4: Influence of the accuracy of \mathbf{W} ($rtol$) up to which the linear systems with \mathbf{A}_+ preconditioned by \mathbf{H}_2 are solved during the setup of the second coarse space. $rtol$: tolerance, κ : condition number of preconditioned operator, It : number of iterations, $\#V_0$: dimension of GenEO coarse space, $n_- = \text{rank}(\mathbf{A}_-)$: dimension of second coarse space. Recall that classical GenEO is not algebraic.

AWG					Classical GenEO				
E	κ	It	$\#V_0$	n_-	E	κ	It	$\#V_0$	n_-
$E = 10^{11}$	12.2	29	35	19	$E = 10^{11}$	13.7	32	28	0
9 layers	9.0	26	57	48	9 layers	11.1	29	55	0
3 layers	9.8	29	43	25	3 layer	9.9	31	35	0
$E = 10^7$	12.2	29	35	19	$E = 10^7$	13.7	32	28	0

Table 5: The number of layers of the harder coefficient varies. Two cases with homogeneous E are also considered. κ : condition number of preconditioned operator, It : number of iterations, $\#V_0$: dimension of GenEO coarse space, $n_- = \text{rank}(\mathbf{A}_-)$: dimension of second coarse space. Recall that classical GenEO is not algebraic.

shows how increasing $rtol$ affects the condition number of \mathbf{A} preconditioned by AWG. Two cases have been studied with different Poisson’s ratios: $\nu = 0.3$ and $\nu = 0.4$. Up to $rtol = 10^{-2}$ there is no change compared to $rtol = 10^{10}$. In fact, up to $rtol = 0.5$, the condition number is hardly degraded. It must be kept in mind that this is a case with few subdomains and small changes to κ on such a case could become more problematic with more subdomains. Still, the conclusion is optimistic: the linear solves with \mathbf{A}_+ do not need to be too precise. Of course a theoretical study of how precise they should be is necessary.

Varying number of channels This time it is the number of layers of the harder coefficient that varie. The two extreme cases of homogenous hard and soft material are also considered. The results are shown in table 5. Heterogeneous distributions require larger coarse spaces.

Long domain with layers of coefficients This testcase (represented in Figure 5) is often studied in domain decomposition articles and presentations. Its drawback is that it doesn’t have any crosspoints (degrees of freedom that belong to more than two subdomains) which is why it was not considered up until now. In this paragraph, weak scalability is studied. The results are shown in Table 6.

- $\omega = [N, 1]$,
- $h = 1/14$ (mesh size),
- $\nu = 0.3$ (Poisson’s ratio),




Figure 5: Solution: ω has been displaced by 10^5 multiplied by the solution to the PDE. The colors show the distribution of coefficients. Dark: $E = E_1 = 10^{11}$ - Light : $E = E_2 = 10^7$

- $E = \begin{cases} E_1 = 10^{11} & \text{if } y \in [1/7; 2/7] \cup [3/7; 4/7] \cup [5/7; 6/7], \\ E_2 = 10^7 & \text{otherwise,} \end{cases}$ (Young's modulus),
- $\mathbf{H}_2 = \mathbf{H}_{hyb}^{NN}(\tau_{\sharp})$ with $\tau_{\sharp} = 0.1$.

With the AWG preconditioner, the condition number hardly increases with the number of subdomains. The sizes of the coarse spaces increase linearly as expected. This points toward weak scalability.

AWG					Classical GenEO				
N	κ	It	$\#V_0$	n_-	N	κ	It	$\#V_0$	n_-
2	12.6	15	8	8	2	9.5	15	7	0
4	9.8	16	26	20	4	11.9	19	19	0
8	9.0	15	62	44	8	12.6	23	43	0
16	14.7	23	218	170	16	14.1	31	194	0
32	14.8	19	442	346	32	14.1	31	394	0

Table 6: The number of subdomains increases, the problem size is proportional to the number of subdomains. Weak scalable behaviour would be for the number of iterations to remain constant and this is what is observed. N : number of subdomains, κ : condition number of preconditioned operator, It : number of iterations, $\#V_0$: dimension of GenEO coarse space, $n_- = \text{rank}(\mathbf{A}_-)$: dimension of second coarse space. Recall that classical GenEO is not algebraic.

6 Conclusion

In this article new preconditioners called AWG for Algebraic Woodbury-GenEO have been introduced. Combined with PCG, they are algebraic domain decomposition methods with two coarse spaces. Fast convergence can be guaranteed by adjusting some user chosen thresholds and enriching one of the coarse spaces accordingly. Numerical results have been presented as a proof of concept and to illustrate the behaviour of the AWG preconditioners on some simple test cases. Further numerical simulations must be performed to assess the overall efficiency of the AWG preconditioners. Some possible improvements of the AWG preconditioners are still under investigation. This includes decreasing the size n_- of the second coarse space by other choices of \mathbf{B}^s , finding sparse approximations for the second coarse space and coarse solve, and perhaps, injecting some information into the preconditioner like the near kernel of \mathbf{A} in a way inspired by smoothed aggregation [42].

References

- [1] E. Agullo, L. Giraud, and L. Poirel. Robust preconditioners via generalized eigenproblems for hybrid sparse linear solvers. *SIAM Journal on Matrix Analysis and Applications*, 40(2):417–439, 2019.
- [2] P. Amestoy, A. Buttari, J.-Y. L’Excellent, and T. Mary. Performance and Scalability of the Block Low-Rank Multifrontal Factorization on Multicore Architectures. *ACM Transactions on Mathematical Software*, 45:2:1–2:26, 2019.
- [3] P. Amestoy, I. S. Duff, J. Koster, and J.-Y. L’Excellent. A fully asynchronous multifrontal solver using distributed dynamic scheduling. *SIAM Journal on Matrix Analysis and Applications*, 23(1):15–41, 2001.
- [4] S. Balay, S. Abhyankar, M. F. Adams, J. Brown, P. Brune, K. Buschelman, L. Dalcin, A. Dener, V. Eijkhout, W. D. Gropp, D. Karpeyev, D. Kaushik, M. G. Knepley, D. A. May, L. C. McInnes, R. T. Mills, T. Munson, K. Rupp, P. Sanan, B. F. Smith, S. Zampini, H. Zhang, and H. Zhang. PETSc Web page. <https://www.mcs.anl.gov/petsc>, 2021.
- [5] S. Balay, S. Abhyankar, M. F. Adams, J. Brown, P. Brune, K. Buschelman, L. Dalcin, A. Dener, V. Eijkhout, W. D. Gropp, D. Karpeyev, D. Kaushik, M. G. Knepley, D. A. May, L. C. McInnes, R. T. Mills, T. Munson, K. Rupp, P. Sanan, B. F. Smith, S. Zampini, H. Zhang, and H. Zhang. PETSc users manual. Technical Report ANL-95/11 - Revision 3.15, Argonne National Laboratory, 2021.
- [6] S. Balay, W. D. Gropp, L. C. McInnes, and B. F. Smith. Efficient management of parallelism in object oriented numerical software libraries. In E. Arge, A. M. Bruaset, and H. P. Langtangen, editors, *Modern Software Tools in Scientific Computing*, pages 163–202. Birkhäuser Press, 1997.
- [7] A. Brandt. Multi-level adaptive solutions to boundary-value problems. *Mathematics of computation*, 31(138):333–390, 1977.
- [8] A. Brandt. Algebraic multigrid (amg) for sparse matrix eqations. *Sparsity and its Applications*, pages 257–284, 1984.
- [9] W. L. Briggs, V. E. Henson, and S. F. McCormick. *A multigrid tutorial*. SIAM, 2000.
- [10] J. G. Calvo and O. B. Widlund. An adaptive choice of primal constraints for BDDC domain decomposition algorithms. *Electron. Trans. Numer. Anal.*, 45:524–544, 2016.

- [11] L. B. Da Veiga, L. F. Pavarino, S. Scacchi, O. B. Widlund, and S. Zampini. Adaptive selection of primal constraints for isogeometric BDDC deluxe preconditioners. *SIAM Journal on Scientific Computing*, 39(1):A281–A302, 2017.
- [12] L. D. Dalcin, R. R. Paz, P. A. Kler, and A. Cosimo. Parallel distributed computing using python. *Advances in Water Resources*, 34(9):1124–1139, 2011.
- [13] V. Dolean, F. Nataf, R. Scheichl, and N. Spillane. Analysis of a two-level Schwarz method with coarse spaces based on local Dirichlet-to-Neumann maps. *Comput. Methods Appl. Math.*, 12(4):391–414, 2012.
- [14] Y. Efendiev, J. Galvis, R. Lazarov, and J. Willems. Robust domain decomposition preconditioners for abstract symmetric positive definite bilinear forms. *ESAIM Math. Model. Numer. Anal.*, 46(5):1175–1199, 2012.
- [15] J. Galvis and Y. Efendiev. Domain decomposition preconditioners for multiscale flows in high-contrast media. *Multiscale Model. Simul.*, 8(4):1461–1483, 2010.
- [16] J. Galvis and Y. Efendiev. Domain decomposition preconditioners for multiscale flows in high contrast media: reduced dimension coarse spaces. *Multiscale Model. Simul.*, 8(5):1621–1644, 2010.
- [17] M. J. Gander and A. Loneland. Shem: An optimal coarse space for ras and its multiscale approximation. In *Domain decomposition methods in science and engineering XXIII*, pages 313–321. Springer, 2017.
- [18] G. H. Golub and C. F. van Loan. *Matrix Computations*. JHU Press, fourth edition, 2013.
- [19] L. Gouarin and N. Spillane. Petsc4py geneo code, 2021.
- [20] R. Haferssas, P. Jolivet, and F. Nataf. An additive Schwarz method type theory for Lions’s algorithm and a symmetrized optimized restricted additive Schwarz method. *SIAM Journal on Scientific Computing*, 39(4):A1345–A1365, 2017.
- [21] A. Heinlein, A. Klawonn, J. Knepper, and O. Rheinbach. Adaptive gds coarse spaces for overlapping schwarz methods in three dimensions. *SIAM Journal on Scientific Computing*, 41(5):A3045–A3072, 2019.
- [22] V. Hernandez, J. E. Roman, and V. Vidal. SLEPc: A scalable and flexible toolkit for the solution of eigenvalue problems. *ACM Trans. Math. Software*, 31(3):351–362, 2005.
- [23] H. H. Kim, E. Chung, and J. Wang. BDDC and FETI-DP preconditioners with adaptive coarse spaces for three-dimensional elliptic problems with oscillatory and high contrast coefficients. *Journal of Computational Physics*, 349:191–214, 2017.
- [24] A. Klawonn, M. Kuhn, and O. Rheinbach. Adaptive coarse spaces for FETI-DP in three dimensions. *SIAM Journal on Scientific Computing*, 38(5):A2880–A2911, 2016.
- [25] A. Klawonn, M. Lanser, and O. Rheinbach. Toward extremely scalable nonlinear domain decomposition methods for elliptic partial differential equations. *SIAM Journal on Scientific Computing*, 37(6):C667–C696, 2015.
- [26] A. Klawonn, P. Radtke, and O. Rheinbach. A comparison of adaptive coarse spaces for iterative substructuring in two dimensions. *Electron. Trans. Numer. Anal.*, 45:75–106, 2016.

- [27] P. Marchand, X. Claeys, P. Jolivet, F. Nataf, and P.-H. Tournier. Two-level preconditioning for h h-version boundary element approximation of hypersingular operator with GenEO. *Numerische Mathematik*, 146(3):597–628, 2020.
- [28] F. Nataf, H. Xiang, and V. Dolean. A two level domain decomposition preconditioner based on local Dirichlet-to-Neumann maps. *C. R. Math. Acad. Sci. Paris*, 348(21-22):1163–1167, 2010.
- [29] F. Nataf, H. Xiang, V. Dolean, and N. Spillane. A coarse space construction based on local Dirichlet-to-Neumann maps. *SIAM J. Sci. Comput.*, 33(4):1623–1642, 2011.
- [30] D. P. O’Leary. The block conjugate gradient algorithm and related methods. *Linear algebra and its applications*, 29:293–322, 1980.
- [31] C. Pechstein and C. R. Dohrmann. A unified framework for adaptive BDDC. *Electron. Trans. Numer. Anal*, 46(273-336):3, 2017.
- [32] C. Pechstein and R. Scheichl. Analysis of FETI methods for multiscale PDEs. Part II: interface variation. *Numer. Math.*, 118(3):485–529, 2011.
- [33] J. W. Ruge and K. Stüben. Algebraic multigrid. In *Multigrid methods*, pages 73–130. SIAM, 1987.
- [34] Y. Saad. *Iterative methods for sparse linear systems*. SIAM, 2003.
- [35] N. Spillane. An adaptive multipreconditioned conjugate gradient algorithm. *SIAM journal on Scientific Computing*, 38(3):A1896–A1918, 2016.
- [36] N. Spillane. An abstract theory of domain decomposition methods with coarse spaces of the GenEO family. working paper or preprint, Mar. 2021.
- [37] N. Spillane. Toward a new fully algebraic preconditioner for symmetric positive definite problems. working paper or preprint, Mar. 2021.
- [38] N. Spillane, V. Dolean, P. Hauret, F. Nataf, C. Pechstein, and R. Scheichl. A robust two-level domain decomposition preconditioner for systems of PDEs. *C. R. Math. Acad. Sci. Paris*, 349(23-24):1255–1259, 2011.
- [39] N. Spillane, V. Dolean, P. Hauret, F. Nataf, C. Pechstein, and R. Scheichl. Abstract robust coarse spaces for systems of PDEs via generalized eigenproblems in the overlaps. *Numer. Math.*, 126(4):741–770, 2014.
- [40] N. Spillane and D. J. Rixen. Automatic spectral coarse spaces for robust FETI and BDD algorithms. *Int. J. Numer. Meth. Engng.*, 95(11):953–990, 2013.
- [41] A. Toselli and O. Widlund. *Domain decomposition methods—algorithms and theory*, volume 34 of *Springer Series in Computational Mathematics*. Springer-Verlag, Berlin, 2005.
- [42] P. Van, M. Brezina, J. Mandel, et al. Convergence of algebraic multigrid based on smoothed aggregation. *Numerische Mathematik*, 88(3):559–579, 2001.
- [43] M. A. Woodbury. *Inverting modified matrices*. Statistical Research Group, 1950.
- [44] J. Xu and L. Zikatanov. Algebraic multigrid methods. *Acta Numerica*, 26:591–721, 2017.
- [45] Y. Yu, M. Dryja, and M. Sarkis. From Additive Average Schwarz Methods to Non-overlapping Spectral Additive Schwarz Methods. *arXiv preprint arXiv:2012.13610*, 2020.

- [46] S. Zampini. PCBDDC: a class of robust dual-primal methods in PETSc. *SIAM Journal on Scientific Computing*, 38(5):S282–S306, 2016.