



HAL
open science

A composite Pliocene record of sea surface temperature in the central Mediterranean (Capo Rossello composite section – South Sicily)

Catherine Beltran, Marie-Alexandrine Sicre, Christian Ohneiser, Matilde Sainz

► To cite this version:

Catherine Beltran, Marie-Alexandrine Sicre, Christian Ohneiser, Matilde Sainz. A composite Pliocene record of sea surface temperature in the central Mediterranean (Capo Rossello composite section – South Sicily). *Sedimentary Geology*, 2021, 420, pp.105921. 10.1016/j.sedgeo.2021.105921. hal-03261629

HAL Id: hal-03261629

<https://hal.science/hal-03261629>

Submitted on 15 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Running head: Evolution of the Equatorial Pacific mean state over the last 10 Myr

2
3 Title: Evolution of the zonal gradients across the Equatorial Pacific during the Miocene-
4 Pleistocene

5 CATHERINE BELTRAN^{1,3}, GABRIELLE ROUSSELLE¹, MARC DE RAFÉLIS², MARIE-
6 ALEXANDRINE SICRE⁴, NATHALIE LABOURDETTE¹, STEFAN SCHOUTEN^{5,6}

7
8 ¹Sorbonne Universités (UPMC, Université Paris 06), UMR 7193, IStEP, Case Postale 116 - 4, place
9 Jussieu 75005 Paris, France.

10 ²Université Paul Sabatier, UMR 5563, Géosciences Environnements Toulouse - 14, avenue Edouard
11 Belin 31400 Toulouse, France.

12 ³Department of Marine Science, University of Otago, PO Box 56, Dunedin 9054, New Zealand.
13 Present address. **Corresponding author: catherine.beltran@otago.ac.nz**

14 ⁴Sorbonne Universités (UPMC, Université Paris 06)-CNRS-IRD-MNHN, LOCEAN Laboratory, 4
15 place Jussieu, F-75005 Paris, France.

16 ⁵NIOZ Royal Netherlands Institute for Sea Research, Department of Marine Microbiology and
17 Biogeochemistry, and University of Utrecht, P.O. Box 59, 1790 AB Den Burg (Texel), The
18 Netherlands

19 ⁶Departement of Earth Sciences, Utrecht University, Heidelberglaan 8, 3584 CS Utrecht, The
20 Netherlands

21
22 **ABSTRACT**

23 Combining $U_{37}^{k'}$ - and TEX_{86} -derived temperatures, and oxygen isotopic values of mixed-
24 layer and thermocline species from the IODP site U1338 (East Equatorial Pacific) and ODP
25 Site 806 (West Equatorial Pacific) we assess the evolution of the zonal sea-surface
26 temperature gradients and thermocline depth across the equatorial Pacific from the Late

27 Miocene through the Pleistocene. Data suggest a long-term shoaling of the thermocline along
28 the equator throughout the Miocene-Pliocene that accelerated around 5.3 Ma. We identify a
29 critical transition at about 3.8 Ma from a El-Niño-like-dominated mean state during the Late
30 Miocene and Early Pliocene to a La-Niña-like-dominated state during the Late Pliocene-
31 Pleistocene. This transition coincides with the restriction of the Indonesian seaway and the
32 onset of ice growth in the northern hemisphere and in Antarctica that led to the long-term
33 strengthening of the Walker circulation and affected low-latitude zonal gradient.

34

35 **KEY WORDS:** Equatorial Pacific, Miocene Pliocene climate, biomarker proxies, planktonic
36 foraminifera, coccoliths.

37

38

INTRODUCTION

39 The equatorial Pacific Ocean plays an important role in the global climate by influencing the
40 heat transport from low to mid- and high latitudes. The modern structure of the equatorial
41 Pacific is characterized by a pronounced east-west asymmetry of the thermocline depth and of
42 sea-surface temperature (SSTs) with the warm pool (~ 29°C) in the west and the cold tongue
43 (~ 23°C) in the east (Fig. 1). The thermocline in the West Equatorial Pacific is deep and
44 controls the flux of atmospheric latent heat (Tian et al., 2001) whereas in the east it is shallow
45 and heat is absorbed (Boccaletti et al., 2004). The SSTs across the equatorial Pacific are
46 affected by the El Niño–Southern Oscillation (Cane, 1998), one of the most important
47 component of the global climate system, which alternates at inter-annual time scale between
48 warm “El Niño” and cold “La Niña” phases.

49 The El Niño–Southern Oscillation is sensitive to the SST distribution across the equatorial
50 Pacific through the link between temperatures and the strength of the trade winds (Neelin et

51 al., 1998). Changes in the meridional SST gradient in the equatorial Pacific have important
52 consequences on the local and global climate.

53 The modern setting of the equatorial Pacific was established during the mid- to late Miocene
54 (LaRivière et al., 2012) through the progressive closure of the Central American Seaway
55 (CAS) and the Indonesian Seaway (Keigwin, 1982; Haug and Tiedemann, 1998; Srinivasan
56 and Sinha, 1998; Schneider and Schmittner, 2006) together with Northern Hemisphere
57 Glaciation (Mudelsee and Raymo, 2005) (Fig. 2). During this period, the onset of the modern
58 thermohaline circulation and equatorial upwelling led to the progressive cooling of deep -
59 water (Woodruff and Savin, 1989; Ravelo et al., 2004), upper-ocean stratification, and
60 thermocline rise. Nathan and Leckie (2009) dated the proto warm pool in the West Equatorial
61 Pacific (WEP) between 11.6 and 10 Ma. The thermocline depth asymmetry set up around 4.8
62 Ma and induced a well-documented cold-water tongue in the Eastern Equatorial Pacific (EEP)
63 between 4.4 and 3.6 Ma (Chaisson and Ravelo, 2000; Wara et al., 2005; Steph et al., 2010;
64 Rousselle et al., 2013).

65 Although numerous studies (e.g., Raymo et al., 1996; Haywood and Valdes, 2004; Ravelo et
66 al., 2004, 2006; Rickaby and Halloran, 2005; Wara et al. 2005; Barreiro et al., 2005; Fedorov
67 et al. 2006, 2013; Dekens et al. 2008; Brierley et al., 2009) established that the equatorial
68 Pacific zonal gradients were weaker during the early Pliocene warm period (El Niño-like, El
69 Padre) and stronger during the mid-Holocene (La Niña-like; Koutavas et al. 2006), there is no
70 consensus on the long-term evolution of the equatorial Pacific mean state since the Miocene
71 (Fig. 2). This is a major limitation to predict El Niño–Southern Oscillation, in that its
72 behavior is closely related to long-term changes in tropical Pacific oceanography that are still
73 not fully understood. Some authors suggest that the equatorial thermal state remained
74 unchanged over the last 12 Myr (Zhang et al., 2014) whereas others identified alternating El
75 Niño-like and La Niña-like conditions (Wara et al., 2005; Ravelo et al., 2006; Nathan and

76 Leckie, 2009; Kamikuri et al., 2009; Drury et al., 2018) or a progressive shoaling of the
77 thermocline over the last 13 Myr (LaRiviere et al., 2012).

78 Because the past variability of the equatorial Pacific mean state is also considered to have
79 played a role in major climatic transitions (Yin and Battisti, 2001), reconstructing the long-
80 term evolution of the thermal structure in this region is crucial.

81 While most published studies focus on very specific time intervals (e.g., Medina-Elizalde and
82 Lea, 2010; Ford et al., 2015), the purpose of this study is to examine the long-term evolution
83 of equatorial Pacific surface conditions over the last 10 Myr. To do so, surface and subsurface
84 temperature records from IODP Site U1338 (EEP) (Fig. 1) are generated from both alkenones
85 ($U_{37}^{k'}$) produced by coccolithophores (marine unicellular haptophyte algae, Brassell et al.,
86 1986) and glycerol- dialkyl glycerol- tetraethers (GDGTs, TEX_{86}^H) produced by
87 *Thaumarchaeota* (Archaea, Schouten et al., 2002; Kim et al., 2010). Additional hydrological
88 information is provided by the oxygen-isotope composition of calcareous nannofossils
89 *Noelaerhabdaceae* spp. calcifying in the photic zone and the thermocline-dweller
90 foraminifera *Globorotalia menardii* (Martin, 1999; Spero et al., 2003) recovered from the
91 ODP Site 806 (WEP) and IODP Site U1338 (Fig. 1).

92 **OCEANOGRAPHIC SETTING**

93 The equatorial Pacific surface ocean circulation, controlled by the Walker cell and Trade
94 winds, results in two westward surface currents, the North and South Equatorial Currents,
95 respectively (SEC and NEC; Fig. 1). The easterly winds pile warm waters in the WEP and
96 create a subsurface eastward circulation known as the Equatorial Under Current (EUC),
97 which brings cold and nutrient-rich waters to the surface in the EEP. The subsequent thinning
98 of the mixed-layer and formation of a cold tongue in the EEP produce a strong W-E
99 asymmetry of the equatorial thermocline depth, deeper in the West Pacific than in the East
100 Pacific (Fig. 1). The modern equatorial SST gradient across the Pacific varies in response to

101 the inter-annual El Niño-Southern Oscillation. During El Niño years, the trade winds weaken,
102 the thermocline deepens in the EEP, and the equatorial upwelling attenuates. As a
103 consequence, the equatorial SST gradient is reduced, mean temperatures are warmer, and the
104 extratropical heat distribution is impacted (Molnar and Cane, 2002; McPhaden et al., 2006).
105 Since the Miocene, the Pacific zonal temperature gradients and the thermocline tilt have
106 varied. Extended periods of reduced equatorial SST gradient and deep thermocline that mimic
107 the modern El Niño events are called permanent El Niño-like states. Similarly, periods of
108 long-term strong equatorial Pacific zonal gradients are called La Niña-like states.

109 **MATERIALS AND METHODS**

110 *IODP Site U1338 and ODP Site 806*

111 The IODP Site U1338 (2° 30.469' N, 117° 58.178' W, 4200 meters water depth, leg 321) was
112 drilled in the EEP, in the modern cold tongue (Pälike et al., 2010) (Fig. 1). Despite the
113 northward drift of the Pacific plate, this site remained in the equatorial band for the past 10
114 Myr (Pälike et al., 2010). Sediments at Site U1338 were deposited above the calcite
115 compensation depth (Pälike et al., 2012) and are composed of well-preserved biogenic
116 material, predominantly nannofossil oozes, with variable abundances of diatoms, radiolarians,
117 and foraminifera (Pälike et al., 2010). The age model of the core was constructed from
118 biostratigraphic (nannofossils, foraminifers, diatoms, and radiolarians) and paleomagnetic
119 data (Pälike et al., 2010; Backman et al., 2016). Average sedimentation rates are relatively
120 high for the studied time interval (~ 27 m/Myr).

121 ODP Site 806 (0° 19,1' N, 159° 21,7' E, 2520 meters water depth, leg 130) is located on the
122 northeast margin of the Ontong Java plateau in the area of the WEP warm pool (Fig. 1).
123 Sediments are composed mostly of well-preserved calcareous nannofossils and planktic
124 foraminifera. The age model is based exclusively on biostratigraphic data of calcareous
125 nannofossils, foraminifera, diatoms, and radiolarians established by Kroenke et al. (1991),

126 later refined by Takayama (1993) and Chaisson and Leckie (1993). Ages are corrected based
127 on biostratigraphic data of Backman et al. (2016) (see Supplement). The average
128 sedimentation rate at this site is higher than at U1338, ranging from ~ 45 m/Myr between 9
129 and 5 Ma to ~ 29 m/Myr during the Pliocene.

130 Proxy records at both sites are generated for the period between 10 and 0.18 Ma using
131 samples from sections U1338B-22H7W through U1338B-1H2W and sections 806B-37X3W
132 through 806B-1H1W. The average sampling resolution of these records varies between 100
133 kyr and 400 kyr.

134 *Oxygen Stable-Isotope Measurements*

135 To determine changes in the thermocline depth in the EEP and WEP through time, we
136 measured the oxygen-isotope ratios ($\delta^{18}\text{O}$) in the calcite of calcareous-nannofossil-enriched
137 fractions and of the upper-thermocline-dweller planktonic foraminifera *Globorotalia*
138 *menardii*. Calcareous nannofossils calcify mainly in the photic zone and thus are used to
139 derive surface-water properties (Roth, 1986) whereas *G. menardii* is selected to assess
140 subsurface water changes.

141 $\delta^{18}\text{O}$ values are measured on calcareous nannofossils *Noelaerhabdaceae* spp. isolated from
142 the bulk sediment following the procedure of Minoletti et al. (2009). $\delta^{18}\text{O}$ measurements are
143 performed on the 2-5 μm fractions at both sites. For site U1338, we use the $\delta^{18}\text{O}$ data previously
144 published by Rousselle et al. (2013) based on the same protocol as in the present study.
145 $\delta^{18}\text{O}$ values are measured using a Finnigan Delta E mass spectrometer at 50°C, at a temporal
146 resolution of 400 kyr. The quality of the granulometric separation based on 200 particles
147 counting in smear slides is estimated to be at least 75% except between 8 and 9 Ma at Site
148 806, where *Sphenolitus abies* dominated. These measurements are thus not considered here.

149 In addition, five to seven specimens of *G. menardii* were hand-picked from the > 160 μm
150 fractions for isotopic measurements. Analyses are performed on a VG ISOPRIME at 90°C

151 with a precision of 0.1‰ and expressed with reference to the VPDB international standard. 45
152 samples at Site U1338 and 99 samples at Site 806 were analyzed. The average temporal
153 resolution over the last 10 Myr is comprised between 120 and 150 ka.

154 *Alkenone Analyses*

155 Alkenone-derived SSTs of Site U1338 previously reported by Rousselle et al., (2013) are
156 supplemented in the present study with 23 additional samples selected from the Pliocene
157 interval (U1338B-8H5W to U1338B-1H2W) to reach a temporal resolution of 100 kyr.
158 Sedimentary alkenones are analyzed following the procedure described by Ternois et al.
159 (2000). About 5 g of freeze-dried sediments are extracted in a mixture of CH₂Cl₂ / CH₃OH.
160 Alkenones are isolated from the total lipid extract by silica-gel chromatography using
161 solvents of increasing polarity. The fraction containing alkenones is concentrated, transferred
162 into clean glass vials, and evaporated under a nitrogen stream. Gas-chromatography analyses
163 are performed using a Varian 3400CX series equipped with a septum programmable injector
164 (SPI) and a flame ionization detector (FID). We use a fused silica capillary column
165 (Chrompack CP Sil5CB, 50 m long, 0.32 mm internal diameter, 0.25 μm film thickness) and
166 helium as a carrier gas. The alkenone unsaturation ratio $U^{K'}_{37}$ ($C_{37:2}/(C_{37:2} + C_{37:3})$) is converted
167 into SSTs using the Conte et al. (2006) calibration: $T = -0.957 + 54.293 (U^{K'}_{37}) - 52.894$
168 $(U^{K'}_{37})^2 + 28.321 (U^{K'}_{37})^3$, which provides more accurate estimates in warm-temperature
169 range than Prahl et al. (1988). Average external precision of the SSTs using this calibration
170 has been estimated at ~ 1.2°C (Conte et al., 2006). Replicate analyses indicate internal
171 precision of ± 0.5°C. Because of the limited amount of sediment material at Site 806 (where
172 alkenone concentrations are extremely low), previously published data from Pagani et al.
173 (2010) provide data covering the past 5 Myr. Note that we recalculated SSTs from the $U^{K'}_{37}$
174 values of Pagani et al. (2010) using the Conte et al. (2006) calibration to avoid calibration
175 bias between sites (see Supplement).

176

GDGT Analyses

177 Forty samples from Site U1338 were analyzed for GDGTs at the Netherlands Institute for Sea
178 Research (NIOZ, Texel, The Netherlands). About 5 g of sediment are freeze dried and
179 extracted in a mixture of CH₂Cl₂ and CH₃OH. The total lipid extracts are separated following
180 the protocol described in Schouten et al. (2002), using Al₂O₃ column chromatography to
181 separate non-polar compounds from the polar GDGTs. Polar fractions are then filtered
182 through a 0.4 μm pore-size filter, evaporated under nitrogen, and diluted to a concentration of
183 2 mg/ml. GDGTs are analyzed using an Agilent HP1100 (HPLC) equipped with a Prevail
184 Cyano column (3 μm, 150 mm x 2.1 mm) following Schouten et al. (2007). The TEX₈₆ index
185 is calculated using peak areas of the following GDGTs as defined by Schouten et al. (2002):
186 $TEX_{86} = [GDGT2] + [GDGT3] + [GDGT5'] / [GDGT1] + [GDGT2] + [GDGT3] + [GDGT5']$. The
187 TEX_{86}^H , defined as the log(TEX₈₆), is used to calculate temperatures over the past 9 Myr.
188 Temperatures are calculated using the calibration established for temperatures above 15°C by
189 Kim et al. (2010) ($T = 68.4 * TEX_{86}^H + 38.6$). The error on temperature estimates is ± 2.5°C.
190 Other calibrations have been proposed for TEX₈₆ temperatures, in particular based on
191 Bayesian statistics (Tierney and Tingley, 2015) but these yield results similar to those shown
192 here, whereas the Kim et al. (2010) calibration is consistent with other studies from this area
193 (e.g., Seki et al., 2010; Zhang et al., 2014). TEX₈₆^H - derived temperatures are obtained at a
194 mean temporal resolution of 230 kyr. At Site 806, because of the very low GDGT
195 concentrations, we use previously published data from Zhang et al. (2014) covering the past
196 12 Myr for comparison with the data from Site U1338.

197

RESULTS

198

Stable Isotopes of the Calcareous Nannofossil Fine Fractions and G. menardii

199

δ¹⁸O Record at Site U1338.---

200 The oxygen stable-isotope signatures of the *Noelaerhabdaceae*-enriched fine fractions
201 ($\delta^{18}\text{O}_{\text{Noelaerhabdaceae}}$) vary between -1.5‰ and 0.5‰ (Fig. 3A). $\delta^{18}\text{O}_{\text{Noelaerhabdaceae}}$ progressively
202 increase between 8 Ma and 5 Ma, before declining to -1.3‰ until 2.7 Ma and rising again to
203 0.4‰. The $\delta^{18}\text{O}_{\text{Noelaerhabdaceae}}$ record is similar to the higher resolution $\delta^{18}\text{O}$ record of the
204 surface dweller *G. sacculifer* from the nearby Site 847 covering only the last 5 Myr (Wara et
205 al., 2005) (Figs. 1, 3A).

206 $\delta^{18}\text{O}$ values measured on the thermocline-dwelling planktonic foraminifera *G. menardii*
207 ($\delta^{18}\text{O}_{\text{G.menardii}}$) over the last 8 Myr (Figure 3A) demonstrate that values vary between -1.4‰
208 and 1.25‰ and display trends similar to those of $\delta^{18}\text{O}_{\text{Noelaerhabdaceae}}$ until 5 Ma. Then,
209 $\delta^{18}\text{O}_{\text{G.menardii}}$ become more positive than $\delta^{18}\text{O}_{\text{Noelaerhabdaceae}}$ and still increase until 0.4 Ma, as
210 also recorded in the deep thermocline dweller *G. tumida* $\delta^{18}\text{O}$ values ($\delta^{18}\text{O}_{\text{G.tumida}}$) at Site 847
211 (Figs. 1, 3A; Wara et al., 2005).

212 **$\delta^{18}\text{O}$ Record at Site 806.---**

213 $\delta^{18}\text{O}_{\text{Noelaerhabdaceae}}$ at site 806 vary within a range similar (between -2.8 and 0.9‰) to site
214 U1338 (Fig. 3B). $\delta^{18}\text{O}_{\text{Noelaerhabdaceae}}$ values are relatively stable (around 0‰) between 7 Ma
215 and 4.4 Ma and rapidly decline to more negative values between 4.5 Ma and 4 Ma, after
216 which values remain stable until 0.38 Ma. The $\delta^{18}\text{O}_{\text{G.menardii}}$ record at Site 806 shows a long-
217 term decrease between 9.7 Ma and 7 Ma and becomes stable until 1.9 Ma. At 3.7 Ma oxygen-
218 isotope values decrease sharply by 1‰, and again between 2.5 Ma and 1.8 Ma. The
219 $\delta^{18}\text{O}_{\text{G.menardii}}$ values then increase to higher values (-0.4 ‰) until 0.5 Ma and further to 0.5‰
220 at 0.2 Ma. Overall, $\delta^{18}\text{O}_{\text{Noelaerhabdaceae}}$ and $\delta^{18}\text{O}_{\text{G.menardii}}$ reveal more positive values in the EEP
221 than in the WEP and include larger-amplitude variations.

222 *EEP Temperature Reconstructions.*

223 Alkenone-derived SSTs ($\text{SST}_{\text{U}^{\text{K}^2}_{37}}$) at Site U1338 are relatively stable and warm ($\sim 27^\circ\text{C}$)
224 between 10 Ma and 6.8 Ma (Fig. 4A). They cool by 1°C between 6.8 Ma and 5.6 Ma, and

225 show a broad warming until 3.7 Ma. From 3.7 Ma onwards, SSTs depict another cooling of
226 about 4°C and superimposed cold episodes of 1 to 2°C amplitude around 2.9 Ma, 1.7 Ma, and
227 0.5 Ma.

228 From the Late Miocene to the Pleistocene, TEX₈₆-derived temperatures (T_{TEX}) indicate a
229 long-term cooling. Between 8 and 5 Ma, T_{TEX} and the SST_{U^{K'}₃₇} vary in the same range of
230 values (between 24°C and 27°C; Fig. 4A), although they are slightly cooler. From approx. 4.5
231 Ma, the two temperature proxies show marked temperature difference. Note that T_{TEX}
232 increase around 7.5 Ma and 4.8 Ma, reaching values close to SST_{U^{K'}₃₇}.
233 SST_{U^{K'}₃₇} at Site U1338 are warmer than T_{TEX} by 2 to 3°C on average. Both progressively cool
234 down between 9 Ma and 0.18 Ma but T_{TEX} decreases by 5°C while SST_{U^{K'}₃₇} cools by only
235 3°C (Fig. 4A).

236 DISCUSSION

237 *Surface and Subsurface Water Temperatures in the Equatorial Pacific*

238 Warmer SST_{U^{K'}₃₇} than T_{TEX}, as observed at Site U1338, is reported in several studies (e.g.,
239 Huguet et al., 2007; Lee et al., 2008; Leider et al., 2010; Lopes dos Santos et al., 2010;
240 Rommerskirchen et al., 2011; Seki et al., 2012; Zhang et al., 2014) and explained by different
241 depth habitat or season of production. However, because seasonality is weak at low latitudes,
242 it is unlikely to explain a temperature differences of up to 4°C, at Site U1338. Unlike
243 haptophytes thriving in the photic zone, *Thaumarchaeota*, the sources of the GDGTs, occur
244 throughout the water column (e.g., Karner et al., 2001; Huguet et al., 2007), and the export of
245 GDGTs can be integrated for a larger part of the upper part of the ocean, likely including a
246 part of the thermocline. Wuchter et al. (2005) reported maximum concentrations of GDGTs
247 between 100 and 200 m depth in the Pacific Ocean, coinciding with high ammonium
248 concentrations below the photic zone (Murray et al., 1999; Karner et al., 2001; Wuchter et al.,
249 2005; Huguet et al., 2007; Pearson et al., 2007). GDGT producers can thus preferentially

250 develop in subsurface or deeper waters (Huguet et al., 2007; Kim et al., 2008; Lee et al.,
251 2008; Huguet et al., 2011; Kim et al., 2012). The difference between T_{TEX} and $\text{SST}_{\text{U}^{\text{K}'37}}$ in the
252 EEP may then account for different production or export depth (Lopes dos Santos et al., 2010;
253 Rommerskirchen et al., 2011; Schouten et al., 2013).

254 The interpretation of T_{TEX} temperatures across the equatorial Pacific is complicated by both the
255 upwelling in the EEP and the different trophic conditions between the EEP and the WEP that
256 influence the export depth of GDGTs (Seki et al., 2012). Numerous studies indicate that
257 upwelling impacts on the depth of production and export of GDGTs to the sediment (Wuchter
258 et al., 2005; Huguet et al., 2007; Lee et al., 2008; Lopes dos Santos et al., 2010; Seki et al.,
259 2012). In the absence of upwelling, T_{TEX} reflects the surface temperatures (Sinninghe Damsté
260 et al., 2002; Wakeham et al., 2004) whereas under upwelling conditions, T_{TEX} represent
261 thermocline temperatures (Wakeham et al., 2002; Wuchter et al., 2005; Huguet et al., 2007).

262 Furthermore, the nitrate maximum across the modern equatorial Pacific exhibits a clear
263 asymmetry in the depth. Könneke et al. (2005) found that the production of *Thaumarchaeota*
264 occurs in the subsurface nitrate maximum, which thus is deeper in oligotrophic regions such
265 as the WEP (Brzezinski, 1988). As a consequence, within the same basin T_{TEX} may reflect
266 surface and subsurface conditions in response to different environmental factors (Hertzberg et
267 al., 2016).

268 In the EEP, the $T_{\text{TEX}}-\text{SST}_{\text{U}^{\text{K}'37}}$ temperature gradient (ΔT) is proposed to monitor changes of
269 the thermocline depth (Seki et al., 2012) assuming that ΔT is driven by the depth migration of
270 GDGT producers in response to upwelling activity. Therefore, at Site U1338, warming T_{TEX}
271 and reduced ΔT reflect a deep thermocline and surface export of GDGTs, whereas a more
272 pronounced cooling of T_{TEX} compared to $\text{SST}_{\text{U}^{\text{K}'37}$, (increased ΔT) reflects a shallow
273 thermocline, upwelling activity, and subsurface export of GDGTs.

274 In the WEP, the interpretation of $T_{\text{TEX}}\text{-SST}_{\text{U}^{\text{K}'37}}$ is not straightforward either. At Site 806,
275 $\text{SST}_{\text{U}^{\text{K}'37}$ calculated from Pagani et al., (2010) are warm and stable, averaging 28 to 28.5°C
276 over the last 5 Myr (Fig. 4C) except for a 2°C sharp cooling at ~ 1.5 Ma. High-resolution
277 Mg/Ca-derived SSTs ($\text{SST}_{\text{Mg/Ca}}$) corrected from Mg/Ca_{seawater} (Medina-Elizalde et al., 2008)
278 from the same site show markedly warmer conditions in the surface of the WEP especially
279 between 5 and 3.5 Ma, reaching up to 32°C. This difference with the alkenone-derived
280 temperatures can be attributed to (1) the alkenone calibration being less accurate in the
281 warmer temperature range (Conte et al., 2006), thus leading to the underestimation of
282 $\text{SST}_{\text{U}^{\text{K}'37}$ (Tierney and Tingley 2018) or (2) the uncertainties on the seawater Mg/Ca
283 correction. However, from 3 Ma onwards, $\text{SST}_{\text{U}^{\text{K}'37}$ and $\text{SST}_{\text{Mg/Ca}}$ temperatures display
284 similar trends and values, except between 1.8 Ma and 1.2 Ma, when they diverge. Since this
285 discrepancy coincides with isotopic changes (Fig. 3) we can reasonably conclude that the
286 difference between the two records during the late Pliocene reflects oceanographic changes.
287 During the early Pliocene, cooler $\text{SST}_{\text{U}^{\text{K}'37}$ appear to be linked to a calibration limitation of
288 the proxy at warm temperatures.

289 Comparison between the T_{TEX} record (Zhang et al., 2014) and subsurface temperatures
290 derived from Mg/Ca_{*G.tumida*} (Ford et al., 2015) at ODP Site 806, highlights that from 5 to 3.5
291 Ma both reconstructions agree well within their uncertainties (Fig. 4C). From 3.5 to 2 Ma,
292 T_{TEX} match with $\text{SST}_{\text{U}^{\text{K}'37}$ and Mg/Ca-derived SSTs from the surface dweller *G. ruber*
293 (Mg/Ca_{*G.ruber*} Medina-Elizalde et al., 2008), whereas Mg/Ca_{*G.tumida*} temperatures are
294 systematically cooler than T_{TEX} (Fig. 4C). After 1.3 Ma, T_{TEX} values are intermediate between
295 Mg/Ca_{*G.ruber*} and Mg/Ca_{*G.tumida*} temperatures, while $\text{SST}_{\text{U}^{\text{K}'37}$ tend to agree with Mg/Ca_{*G.ruber*}
296 SSTs. It is likely that in the WEP, the T_{TEX} reflect production in different water depth in
297 response to trophic changes and environmental conditions. We speculate that during the warm
298 early Pliocene and after 1.3 Ma, T_{TEX} reflect subsurface conditions and that during the Late

299 Pliocene, the export of GDGTs derived mainly from surface waters. This interpretation
300 nuances the interpretations of Zhang et al. (2014), who considered that T_{TEX} reflect surface
301 temperatures and could be used as an alternative to $U^{\text{K}'}_{37}$ under warm conditions.

302 *Changes in the Equatorial Pacific Mean State over the Last 10 Myr*

303 To investigate the evolution of the surface stratification across the Equatorial Pacific, we
304 calculate the oxygen-isotope gradients between *Noelaerhabdaceae* and *G. menardii* ($\Delta\delta^{18}\text{O}$)
305 (Fig. 4B) (Nathan and Leckie, 2009; Beltran et al., 2014). Note that in doing so, the sea-water
306 $\delta^{18}\text{O}$ component is factored out. Higher (lower) $\Delta\delta^{18}\text{O}$ values reflect a shallower (deeper)
307 thermocline (Ravelo and Fairbanks, 1992; Ravelo and Shackleton, 1995; Farrell et al., 1995;
308 Nathan and Leckie, 2009, Beltran et al., 2014).

309 Variations in the gradient between T_{TEX} and $\text{SST}_{U^{\text{K}'}_{37}}$ also can be used to evaluate changes in
310 the thermocline depth. T_{TEX} and $\text{SST}_{U^{\text{K}'}_{37}}$ values similar to those observed between 7.3 Ma
311 and 6.8 Ma, and around 4.8 Ma at IODP Site U1338, indicate a deep thermocline (Fig. 4A).
312 Some intervals are characterized by an increase in ΔT (such as, for example, between 6.8 Ma
313 and 5.3 Ma at IODP Site U1338) associated with a cooling trend of both proxies. These
314 features indicate that GDGTs are produced deeper and that surface waters are concomitantly
315 cooling, which is consistent with upwelling activity.

316 Based on temperatures and isotopic gradients between surface and subsurface waters from the
317 EEP and WEP, the data reveal a pronounced oceanographic change at *ca.* 5 Ma with the
318 thermocline progressively rising (Fig. 4A, 4B).

319 **Late Miocene [9 – 5.3 Ma] Interval.---**

320 During the late Miocene period, east and west $\Delta\delta^{18}\text{O}$ are reduced and stable, indicating a deep
321 thermocline across the equatorial Pacific (Fig. 4B). From 10 Ma to 6.3 Ma, data indicate
322 warm SSTs ($\sim 27\text{ }^\circ\text{C}$) in the EEP (Fig. 4A). This result coincides with the low abundances of
323 upwelling radiolarian species reported between 9 and 7 Ma by Kamikuri et al. (2009). Similar

324 SST_{U^K37} and T_{TEX} values at U1338 between 7.3 and 6.8 Ma also is consistent with a thick
325 mixed layer in the EEP (Fig. 4A; Seki et al., 2012). In the WEP, Nathan and Leckie (2009)
326 reported a similar decrease in the abundance of surface dwellers in favor of thermocline
327 species, which suggests a deep thermocline. This reduced zonal gradient in SSTs and
328 thermocline tilt across the equatorial Pacific corresponds to El Niño-like conditions, in
329 agreement with earlier findings of Herbert et al. (2016), who indicate an expanded and weak
330 Hadley circulation before 8 Ma.

331 The long-term Late Miocene El Niño-like state was interrupted by a brief interval (6.5 - 5.3
332 Ma) during which the thermocline shoaled in the EEP, as shown by the $\delta^{18}\text{O}$ gradient (Fig.
333 4B). The SST cooling and decrease of surface salinity recorded at Site U1338 (Rousselle et
334 al., 2013), supported by higher abundances of upwelling siliceous species during this interval
335 (Kamikuri et al., 2009), suggest the first establishment of the cold tongue and of upwelling in
336 the EEP. Increasing abundances of surface-dweller species at Site 806 (Nathan and Leckie,
337 2009) (Fig. 3 and 4), this interval further supports the idea of a weak and transient La Niña
338 like period. Drury et al. (2018) suggested a cooling of the thermocline waters related to the
339 global late Miocene cooling (Federov et al., 2006; Drury et al., 2017). Although the data do
340 not rule out this hypothesis, the T_{TEX} cooling is supportive of a change in depth of GDGT
341 production or export in response to the onset of the EEP upwelling.

342 **Plio-Pleistocene [from 5.3 Ma] Interval.---**

343 The oxygen-isotope and temperature data indicate a long-term gradual thermocline shoaling
344 and sea-surface cooling in the EEP (Fig. 4A, 4B) throughout this interval. In the WEP, the
345 thermocline shoaled between 5.3 and 4 Ma, returning to a deep position between 4 and 1.8
346 Ma and shoaling again after 1.8 Ma. This interval is characterized by the initiation of a
347 progressive asymmetry in the thermocline depth across the equatorial Pacific.

348 After 5.3 Ma, SSTs at IODP Site U1338 were warm (~ 27 °C), in agreement with a deep
349 thermocline until 4.5 Ma (Fig. 4A). This result is supported by a rapid increase of T_{TEX} at
350 U1338. Moreover, abundances of upwelling radiolarian species decreased again in the EEP
351 (Kamikuri et al., 2009). In the WEP, the proto-warm pool period ended (Nathan and Leckie,
352 2009). This interval coincides with the Messinian salinity crisis (Hilgen et al., 2007), during
353 which the isolation of the Mediterranean Sea would have triggered a reduction of the North
354 Atlantic Deep Water formation (NADW) at ~ 6 Ma because of changing water density of the
355 North Atlantic waters (Pérez-Asensio et al., 2012). According to model simulations, this
356 process would result in a deepening of the thermocline in the equator and El Niño-like
357 conditions (Rühlemann et al., 2004; Timmermann et al., 2005; Barreiro et al., 2008), thus
358 providing a possible explanation to the results herein.

359 Numerous studies have focused on the magnitude of Pliocene equatorial Pacific warmth (4.5
360 – 3.5 Ma) and the factors responsible for it (e.g., Raymo et al., 1996; Haywood and Valdes,
361 2004; Ravelo et al., 2004; Barreiro et al., 2008; Rickaby and Halloran, 2005; Brierley et al.,
362 2009; Scroxton et al., 2011; Ford et al., 2015). It is accepted that during the mid-Piacenzian
363 Warm Period, persistent El-Niño like conditions prevailed. This interval, also known as El
364 Padre mean state (Ford et al., 2015), is characterized by a deep and warm thermocline and
365 warm surface conditions across the equatorial Pacific.

366 The low-resolution record presented here suggests a brief interval centered around 4 Ma when
367 the thermocline deepened in the east and shoals in the west. SSTs in the EEP reach 30°C and
368 around 32°C in the WEP (Fig. 4A, 4C, $\text{Mg}/\text{Ca}_{G.ruber}$ SSTs; Medina-Elizalde et al., 2008).
369 From 3.5 Ma, a pronounced zonal gradient of the thermocline depth and of the SSTs
370 developed across the Pacific. The SST cooling trend and shoaling of the thermocline
371 identified in the EEP coincides with the permanent establishment of the cold tongue
372 (Groeneveld et al., 2006; Dekens et al., 2007; Steph et al., 2010; Rousselle et al., 2013). This

373 result is in agreement with Chaisson and Ravelo (2000) and Kamikuri et al., (2009), who
374 report higher abundances of subsurface-dweller foraminifera and radiolarian upwelling
375 species in the EEP at that time. Simultaneous with the changes in the EEP, the warm pool
376 became established and the thermocline deepened in the WEP, as suggested by warmer
377 surface waters and reduced ΔT (Fig. 4C, Wara et al., 2005; Medina-Elizalde et al., 2008) and
378 planktonic foraminifera abundances (Cannariato and Ravelo, 1997; Chaisson and Ravelo,
379 2000; Li et al., 2006). This La Niña-like pattern is consistent with earlier studies (Rickaby and
380 Halloran 2005; Kamikuri et al., 2009; Steph et al., 2010; Ford et al., 2012 and Rousselle et al.,
381 2013) in favor of an active equatorial upwelling and a strong Walker cell.
382 Between 1.8 and 1.3 Ma, data indicate that the thermocline tilt and the SST gradient across
383 the Pacific were reinforced (Fig. 4). This result is in agreement with the setting of the modern
384 cold tongue identified by Martinez-Garcia et al. (2011).

385 *Evolution of the Pacific Mean State in a Regional and Global Context*

386 Prior research has attributed changes in the thermocline tilt to local tectonic events such as the
387 opening and closing of the Panama and Indonesian oceanic gateways (e.g., LaRiviere et al.,
388 2012), and changes in sea-surface-temperature zonal gradients to global climate perturbations
389 or changes in regional wind patterns and strength (Hovan, 1995; Ford et al., 2012).
390 However, there is still a strong debate about the timing of the closure of these oceanic
391 gateways and their role in controlling the equatorial Pacific mean climate state. Montes et al.,
392 (2012, 2015), Osborne et al. (2014), and Sepulchre et al. (2014) suggested that the Central
393 American Seaway was restricted by 10-11 Ma, which contradicts the commonly accepted late
394 Pliocene timing of its closure (Keigwin, 1982; Duque-Caro, 1990; Haug and Tiedemann,
395 1998; Haug et al., 2001). On the other hand, Molnar and Cronin (2015) showed a marked
396 restriction of the Indonesian Seaway from 5 Ma. This interpretation implies that the changes

397 of the equatorial Pacific mean state during the Miocene would have been controlled mainly by
398 global climate changes.

399 The brief La Niña-like interval between 6.5 and 5.3 Ma may be attributable to the
400 combination of the restricted flow through the Central American Seaway (Keigwin, 1982;
401 Keller et al., 1989) and the 25 m sea-level drop at ~ 6.5 Ma (Haq et al., 1987; Hardenbol et
402 al., 1998; Haq and Schutter, 2008) due to the expansion of ice sheets in Greenland (Larsen et
403 al., 1994), Alaska (Bradshaw et al., 2012) and Antarctica (Billups, 2002; Williams et al. 2010)
404 and the 2 Myr-long Northern Hemisphere glaciation (Larsen et al., 1994). This global cooling
405 is thought to have contributed to the reinforcement of the equatorial upwelling and EUC.

406 Data from sediments younger than ~ 5 Ma indicate an important change in the equatorial
407 Pacific mean state with the dominance of La Niña-like intervals. The onset of this long-term
408 state coincides with a drastic restriction of the Indonesian Seaway (Srinivasan and Sinha,
409 1998; Molnar and Cronin, 2015), and changes in the global climate. The Indonesian Seaway
410 restriction is thought to have contributed to the cooling of the eastern tropical Pacific over the
411 last 5 Myr through the strengthening of the Walker circulation, leading to the increase in the
412 Pacific zonal temperature gradient (Fig. 3, 4).

413 The La Niña-like event between 1.8 and 1.3 Ma coincides with pronounced ice growth in
414 Antarctica (McKay et al., 2012) and the development of the northern-hemisphere ice sheets.
415 The gradual increase of the pole-to-equator temperature gradient is thought to have reinforced
416 the Walker cell, and affected the low-latitude zonal gradient (Molnar and Cane, 2002; McKay
417 et al., 2012). Furthermore, modelling experiments show a positive feedback of the Pliocene
418 low-latitude cooling on the global cooling (Fedorov et al., 2006). The shoaling of the
419 thermocline increased the climate sensitivity of this area by strengthening the temperature
420 gradient through the tropics and subsequently the Walker and Hadley cells (Brierley et al.,
421 2009; Brierley and Fedorov, 2010). These processes led to the final setting of upwelling areas

422 (Philander and Fedorov, 2003; Dekens et al., 2007; Brierley et al., 2009; Brierley and
423 Fedorov, 2010).

424

CONCLUSION

425 This study demonstrates the complexity of reconstructing the evolution of the equatorial
426 Pacific mean climate state and oceanographic features and the need of a multi-proxy approach
427 to get a comprehensive understanding of the mechanism at play over the past 10 Myr.
428 Combined isotopic zonal gradients between the surface (*Noelaerhabdaceae*) and subsurface
429 (*G. menardii*) waters from the east and west equatorial Pacific reveals important steps towards
430 the establishment of what is known today as El Niño–Southern Oscillation. The record
431 suggests a general shoaling of the thermocline along the equator since the Late Miocene. The
432 data show the appearance of the cold tongue at ~ 6.8 Ma before its proper establishment
433 around 4.4 to 3.6 Ma (Chaisson 1995; Cannariato and Ravelo, 1997; Chaisson and Ravelo,
434 2000; Molnar and Cane, 2002; Ravelo et al., 2004, 2006; Sato et al., 2008; Steph et al., 2010;
435 Rousselle et al., 2013) and its modern configuration ~ 1.8 Ma. This result is in agreement with
436 the findings of LaRivière et al. (2012) and Zhang et al., (2014), who suggested progressive
437 shoaling of the thermocline over the last 13 Myr.

438 The data indicate two critical transitions at about 5.3 Ma, when the thermocline globally
439 started shoaling across the equatorial Pacific and around 3.8 Ma from an El-Niño-like-
440 dominated mean state during the Late Miocene to a La-Niña-like-dominated state during the
441 Plio-Pleistocene. These transitions coincide with marked regional and global changes.
442 Locally, the restriction of the Indonesian Seaway and subsequent intensification of the Walker
443 circulation increased the Pacific zonal temperature gradient over the last 5 Myr. From a global
444 point of view, the growth of ice sheets in the northern hemisphere and in Antarctica most
445 likely affected the Walker cell and the low-latitude zonal gradient.

446

447

ACKNOWLEDGMENTS

448 We thank the European Consortium for Ocean Research Drilling (ECORD) for the ECORD
449 Grant. We are also grateful to Anhelique Mets and Jort Ossebaar for support with the TEX₈₆
450 measurements. We finally want to thank Fabrice Minoletti for providing us the separation
451 technique development necessary to then perform the isotopic measurements on
452 *Noelaerhabdaceae* calcite. We also thank the Ocean and International Ocean drilling Program
453 (ODP and IODP) for providing samples from Sites 806 and U1338. SS and AS were
454 supported by the Netherlands Earth System Science Center funded by the Dutch Ministry
455 Science and Education (OCW). MAS was supported by CNRS (Centre National de la
456 Recherche Scientifique) for salary.

457

REFERENCES

- 458 Backman, J., Baldauf, J.G., Ciummelli, M., and Raffi, I., 2016, Data report: a revised
459 biomagnetostratigraphic age model for Site U1338, IODP Expedition 320/321:
460 Proceedings of the Integrated Ocean Drilling Program, 320/321,
461 <http://doi.org/10.2204/iodp.proc.320321.219.2016>.
- 462 Barreiro, M., Philander, G., Pacanowski, R., and Fedorov, A., 2005, Simulations of warm
463 tropical conditions with application to middle Pliocene atmospheres: *Climate
464 Dynamics*, v. 26, p. 349–365.
- 465 Barreiro, M., Fedorov, A., Pacanowski, R., and Philander, S.G., 2008, Abrupt climate
466 changes: how freshening of the Northern Atlantic affects the thermohaline and wind-
467 driven oceanic circulations: *Annual Review of Earth and Planetary Sciences*, v. 36, p.
468 33-58.
- 469 Beltran, C., Roussele, G., Backman, J., Wade, B.S., and Sicre, M.A., 2014,
470 Paleoenvironmental trigger for the development of bloom forming calcareous
471 nanofossils: an example from the Late Miocene Eastern Equatorial Pacific:
472 *Palaeoceanography*, v. 29 (3), p. 210-222.
- 473 Billups, K., 2002, Late Miocene through early Pliocene deep water circulation and climate
474 change viewed from the sub-Antarctic South Atlantic: *Palaeogeography,
475 Palaeoclimatology, Palaeoecology*, v. 185(3-4), p. 287-307.
- 476 Boccaletti, G., Pacanowski, R.C., Philander, S.G.H., and Fedorov, A.V., 2004, The thermal
477 structure of the upper ocean: *Journal of Physical Oceanography*, v. 34, p. 888–902
- 478 Bradshaw, C.D., Lunt, D.J., Flecker, R., Salzmann, U., Pound, M.J., Haywood, A.M., and
479 Eronen, J.T., 2012, The relative roles of CO₂ and palaeogeography in determining
480 late Miocene climate: Results from a terrestrial model-data comparison: *Climate of
481 the Past*, v. 8, p. 1257–1285.

482 Brassell, S.C., Eglinton, G., Marlowe, I.T., Pflaumann, U., and Sarnthein, M., 1986,
483 Molecular stratigraphy: a new tool for climatic assessment: *Nature*, v. 320, p. 129-133.

484 Brierley, C.M., and Fedorov, A.V., 2010, Relative importance of meridional and zonal sea
485 surface temperature gradients for the onset of the ice ages and Pliocene-Pleistocene
486 climate evolution: *Paleoceanography*, v. 25, PA2214, doi: 10.1029/2009PA001809.

487 Brierley, C., Fedorov, A.V., Liu, Z., Herbert, T.D., Lawrence, K.T., and LaRivière, J.P.,
488 2009, Greatly expanded tropical warm pool and weakened Hadley circulation in the
489 early Pliocene: *Science*, v. 323, p. 1714-1718.

490 Brzezinski, M.A., 1988, Vertical distribution of ammonium in stratified oligotrophic waters:
491 *Limnology and Oceanography*, v. 33, p. 1176-1182.

492 Cane, M.A., 1998, A role for the tropical Pacific: *Science*, v. 282(5386), p. 59-61, doi:
493 10.1126/science.282.5386.59.

494 Cannariato, K.G., and Ravelo, A.C., 1997, Pliocene-Pleistocene evolution of eastern tropical
495 Pacific surface water circulation and thermocline depth: *Paleoceanography*, v. 12(6),
496 p. 805-820.

497 Chaisson W.P., 1995, Planctonic foraminiferal assemblages and paleoceanographic change in
498 the trans-tropical Pacific Ocean: a comparison of West (leg 130) and East (leg 138),
499 latest Miocene to Pleistocene: *Proceedings of the Ocean Drilling Program, Scientific*
500 *Results*, v. 138, p. 555-581.

501 Chaisson, W.P., and Leckie, R.M., 1993, High-resolution Neogene planktonic foraminifer
502 biostratigraphy of Site 806, Ontong Java Plateau (western equatorial Pacific), *in*
503 Berger, W.H., Kroenke, L.W., Mayer, L.A., et al. eds., *Proceedings of Ocean Drilling*
504 *Program, Scientific Results*, 130, p. 137-178.

505 Chaisson, W.P., and Ravelo, A.C., 2000, Pliocene development of the east-west hydrographic
506 gradient in the equatorial Pacific: *Paleoceanography*, v. 15(5), p. 497-505.

507 Conte, M.H., Sicre, M-A., Rühlemann, C., Weber, J.C., Schulte, S., Schulz-Bull, D., and
508 Blanz, T., 2006, Global calibration of the alkenone unsaturation index ($U^{K'}_{37}$) with
509 surface water production temperature and a comparison of the core top integrated
510 production temperatures recorded by $U^{K'}_{37}$ with overlying sea surface temperatures:
511 *Geochemistry Geophysics Geosystems*, v. 7(2), p. 1-22.

512 Dekens, P.S., Ravelo, A.C., and McCarthy, M.D., 2007, Warm upwelling regions in the
513 Pliocene warm period: *Paleoceanography*, v. 22, p. 1-12.

514 Dekens, P.S., Ravelo, A.C., McCarthy, M.D., and Edwards, C.A., 2008, A 5 million year
515 comparison of Mg/Ca and alkenone paleothermometers: *Geochemistry, Geophysics,*
516 *Geosystems*, v. 9, doi:10.1029/2007GC001931.

517 Drury, A.J., Westerhold, T., Frederichs, T., Tian, J., Wilkens, R., Channell, J.E.T., Evans, H.,
518 John, C.M., Lyle, M., and Röhl, U., 2017, Late Miocene climate and time scale
519 reconciliation: Accurate orbital calibration from a deep-sea perspective: *Earth and*
520 *Planetary Science Letters*, v. 475, p. 254–266, doi:10.1016/j.epsl.2017.07.038.

521 Drury, A.J., Geoff, L., Gray, W.R., Lyle, M., Westerhold, T., Shevenell, A., and John, C.,
522 2018, Deciphering the State of the Late Miocene to Early Pliocene Equatorial
523 Pacific: *Paleoceanography and Paleoclimatology*, v. 33, doi:10.1002/2017PA003245.

524 Duque-Caro, H., 1990, Neogene stratigraphy, paleoceanography and paleobiogeography in
525 northwest South America and the evolution of the Panama Seaway: *Palaeogeography,*
526 *Palaeoclimatology, Palaeoecology*, v. 77, p. 203-234.

527 Farrell, J.W., Raffi, I., Janecek, T.R., Murray, D.W., Levitan, M., Dadey, K.A., Emeis, K.,
528 Lyle, M., Flores, J.-A., and Hovan, S., 1995, Late Neogene sedimentation patterns in
529 the eastern equatorial Pacific Ocean: *Proceedings of the Ocean Drilling Program,*
530 *Scientific Results*, v. 138, p. 717-756.

531 Fedorov, A.V., Dekens, P.S., McCarthy, M., Ravelo, A.C., DeMenocal, P.B., Barreiro, M., Pa
532 canowki, R.C., and Philander, S.G., 2006, The Pliocene paradox (mechanisms for a
533 permanent El Niño): *Science*, v. 312, p. 1485–1489.

534 Fedorov, A.V., Brierley, C.M., Lawrence, K.T., Liu, Z., Dekens, P.S., and Ravelo, A.,
535 2013, Patterns and mechanisms of early Pliocene warmth: *Nature*, v. 496, p. 43–49,
536 doi:10.1038/nature12003.

537 Ford, H.L., Ravelo, A.C., and Hovan, S., 2012, A deep Eastern Equatorial Pacific thermocline
538 during the early Pliocene warm period: *Earth and Planetary Science Letters*, v. 355-
539 356, p. 152-161.

540 Ford, H.L., Ravelo, A.C., Dekens, P.S., LaRiviere, J.P., and Wara, M.W., 2015, The
541 evolution of the equatorial thermocline and the early Pliocene El Padre mean state:
542 *Geophysical Research Letters*, v. 42, p. 4878–4887, doi:10.1002/2015GL064215.

543 Groeneveld, J., Steph, S., Tiedemann, R., Garbe, D., Nürnberg, D., and Sturn, A., 2006,
544 Pliocene mixed-layer oceanography for site 1241, using combined Mg/Ca and $\delta^{18}\text{O}$
545 analyses of *Globigerinoides sacculifer*: *Proceedings of the Ocean Drilling Program*,
546 *Scientific Results*, v. 202, p. 1-27.

547 Haq, B.U., and Schutter, S.R., 2008, A chronology of Paleozoic sea-level changes: *Science*, v.
548 322(5898), p. 64-68.

549 Haq, B.U., Hardenbol, J., and Vail, P.R., 1987, Chronology of fluctuating sea levels since the
550 Triassic (250 million years ago to present): *Science*, v. 235, p. 1156-1167.

551 Hardenbol, J., Thierry, J., Farley, M.B., Jacquin, T., De Graciansky, P.C., and Vail, P., 1998,
552 Mesozoic and Cenozoic sequence chronostratigraphic framework of European basins,
553 in De Graciansky, P.-C., Hardenbol, J., Jacquin, T., Vail, P.R., and Farley, M.B., eds.,
554 Mesozoic and Cenozoic Sequence Stratigraphy of European Basins: SEPM, Special
555 Publication 60, p. 3-13.

556 Haug, G.H., and Tiedemann, R., 1998, Effect of the formation of the Isthmus of Panama on
557 Atlantic Ocean thermohaline circulation: *Nature*, v. 393, p. 673-676.

558 Haug, G.H., Tiedemann, R., Zahn, R., and Ravelo, A.C., 2001, Role of Panama uplift on
559 oceanic freshwater balance: *Geology*, v. 29, p. 207-210.

560 Haywood, A.M, and Valdes, P.J., 2004, Modelling middle Pliocene warmth: contribution of
561 atmosphere, oceans and cryosphere: *Earth and Planetary Science Letters*, v. 218, p.
562 363–377.

563 Herbert, T.D., Lawrence, K.T., Tzanova, A., Peterson, L.C., Caballero-Gill, R., and Kelly,
564 C.S., 2016, Late Miocene global cooling and the rise of modern ecosystems: *Nature*
565 *Geoscience*, v. 9, p. 843–847, doi:10.1038/NGEO2813.

566 Hertzberg J.E., Schmidt M.W., Bianchi, T.S., Smith, R.W., Shields, M.R., and Marcantonio,
567 F., 2016, Comparison of eastern tropical Pacific TEX₈₆ and *Globigerinoides ruber*
568 Mg/Ca derived sea surface temperatures: Insights from the Holocene and Last Glacial
569 Maximum: *Earth and Planetary Science Letters*, v. 434, p. 320-332.

570 Hilgen, F., Kuiper, K., Krijgsman, W., Snel, E., and Van Der Laan, E., 2007, Astronomical
571 tuning as the basis for high resolution chronostratigraphy: the intricate history of the
572 Messinian Salinity Crisis: *Stratigraphy*, v. 4(2-3), p. 231-238.

573 Hovan, S.A., 1995, Late Cenozoic atmospheric circulation intensity and climatic history
574 recorded by eolian deposition in the eastern equatorial Pacific Ocean, Leg
575 138: *Proceedings of the Ocean Drilling Program, Scientific Results*, v. 138, p. 615–
576 625.

577 Huguet, C., Schimmelmann, A., Thunell, R., Lourens, L.J., Sinninghe Damsté, J., and
578 Schouten, S., 2007, A study of the TEX₈₆ paleothermometer in the water column and
579 sediments of the Santa Barbara Basin, California: *Paleoceanography*, v. 22, p. 9.

580 Huguet, C., Martrat, B., Grimalt, J.O., Sinninghe Damsté, J.S., and Schouten, S., 2011,

581 Coherent millennial scale patterns in U^{K}_{37} and TEX^{H}_{86} temperature records during the
582 penultimate interglacial-to-glacial cycle in the western Mediterranean:
583 *Paleoceanography*, v. 26, p. 10.

584 Kamikuri, S., Motoyama, I., Nishi, H., and Iwai, M., 2009, Evolution of Eastern Warm Pool
585 and upwelling processes since the middle Miocene based on analysis of radiolarian
586 assemblages: response to Indonesian and Central American Seaways: *Palaogeography*,
587 *Palaeoclimatology, Palaeoecology*, v. 280, p. 469-479.

588 Karner, M.B., DeLong, E.F., and Karl, D.M., 2001, Archaeal dominance in the mesopelagic
589 zone of the Pacific Ocean: *Nature*, v. 409, p. 507–510.

590 Keigwin, L.D., 1982, Isotope paleoceanography of the Caribbean and east Pacific: Role of
591 Panama uplift in late Neogene time: *Science*, v. 217, p. 350-353.

592 Keller, G., Zenker, C.E., and Stone, S.M., 1989, Late Neogene history of the Pacific -
593 Caribbean gateway: *Journal of South American Earth Sciences*, v. 2(1), p. 73-108.

594 Kim, J.H., Schouten, S., Hopmans, E.C., Donner, B., and Sinninghe Damsté, J.S., 2008,
595 Global sediment core-top calibration of the TEX_{86} paleothermometer in the ocean:
596 *Geochimica et Cosmochimica Acta*, v. 72(4), p. 1154-1173.

597 Kim, J.H., Van Der Meer, J., Schouten, S., Helmke, P., Willmott, V., Sangiorgi, F., Koç, N.,
598 Hopmans, E.C., and Sinninghe Damsté, J.S., 2010, New indices and calibrations
599 derived from the distribution of crenarchaeal isoprenoid tetraether lipids: implications
600 for past sea surface temperature reconstructions: *Geochimica et Cosmochimica Acta*,
601 v. 74, p. 4639-4654.

602 Kim, J.H., Romero, O.E., Lohmann, G., Donner, B., Laepple, T., Haam, E., and Sinninghe
603 Damsté, J.S., 2012, Pronounced subsurface cooling of North Atlantic waters off
604 Northwest Africa during Dansgaard - Oeschger interstadials: *Earth and Planetary
605 Science Letters*, v. 339-340, p. 95-102.

606 Könnenke, M., Bernhard, A.E., De La Torre, J.R., Walker, C.B., Waterbury, J.B., and Stahl,
607 D.A., 2005, Isolation of an autotrophic ammonia-oxidizing marine
608 Archaeon: *Nature*, v. 437, p. 543–546.

609 Koutavas, A., Demenocal, G., Olive, C., and Lynch-Stieglitz, J., 2006, Mid-Holocene El Niño
610 - Southern Oscillation (ENSO) attenuation revealed by individual foraminifera in
611 eastern tropical Pacific sediments: *Geology*, v. 34, p. 993–996.

612 Kroenke, L.W., Berger, W.H., and Janecek, T.R., 1991, Site 806: Proceedings of the Ocean
613 Drilling Program, Initial reports, v. 130, p. 291-367.

614 LaRivière, J.P., Ravelo, A.C., Crimmins, A., Dekens, P.S., Ford, H.L., Lyle, M., and Wara,
615 M.W., 2012, Late Miocene decoupling of oceanic warmth and atmospheric carbon
616 dioxide forcing: *Nature*, v. 486, p. 97-100.

617 Larsen, H.C., Saunders, A.D., Clift, P.D., Beget, J., Wei, W., and Spezzaferri, S., 1994, Seven
618 million years of glaciation in Greenland: *Science*, v. 264, p. 952–955.

619 Lee, K.E., Kim, J.H., Wilke, I., Helmke, P., and Schouten, S., 2008, A study of the alkenone,
620 TEX₈₆, and planktonic foraminifera in the Benguela Upwelling System: Implications
621 for past sea surface temperatures estimates: *Geochemistry, Geophysics, Geosystems*,
622 v. 9(10), p. 19.

623 Leider, A., Hinrichs, K.U., Mollenhauer, G., and Versteegh, G.J.M., 2010, Core-top
624 calibration of the lipid-based U^K₃₇ and TEX₈₆ temperature proxies on the southern
625 Italian shelf (SW Adriatic Sea, Gulf of Taranto): *Earth and Planetary Science Letters*,
626 v. 300(1-2), p. 112-124.

627 Li, Q., Li, B., Zhong, G., McGowran, B., Zhou, Z., Wang, J., and Wang, P., 2006, Late
628 Miocene development of the western Pacific warm pool: planktonic foraminifer and
629 oxygen isotopic evidence: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v.
630 237, p. 465-482.

631 Lopes Dos Santos, R.A., Prange, M., Castañeda, I.S., Schefuß, E., Mulitza, S., Schulz, M.,
632 Niedermeyer, E.M., Sinninghe Damsté, J.S., and Schouten, S., 2010, Glacial -
633 interglacial variability in Atlantic meridional overturning circulation and thermocline
634 adjustments in the tropical North Atlantic: *Earth and Planetary Science Letters*, v. 300,
635 p. 407-414.

636 Martin, R.E., 1999, Taphonomy and temporal resolution of foraminiferal assemblages, *in* Sen
637 Gupta, B.K., ed., *Modern Foraminifera*: Kluwer Academic Publishers, 371 p.

638 Martinez-Garcia, A., Rosell-Mele, A., Jaccard, S.L., Geibert, W., Sigman, D.M., and Haug,
639 G.H., 2011, Southern Ocean dust–climate coupling over the past four million years:
640 *Nature*, v. 476, p. 312-315, doi:10.1038/nature10310

641 McKay, R., Naish, T., Carter, L., Riesselman, C., Dunbar, R., Sjunneskog, C., Winter, D.,
642 Sangiorgi, F., Warren, C., Pagani, M., Schouten, S., Willmott, V., Levy, R., DeConto,
643 R., and Powell, R.D., 2012, Antarctic and Southern Ocean influences on Late Pliocene
644 global cooling: *National Academy of Sciences Proceedings (United States of*
645 *America)*, v. 109 (17), p. 6423-6428.

646 McPhaden, M.J., Zebiak, S.E., and Glantz, M.H., 2006, ENSO as an integrating concept in
647 *Earth science: Science*, v. 314, p. 1740–1745.

648 Medina-Elizalde, M., and Lea, D.W., 2010, Late Pliocene equatorial Pacific:
649 *Paleoceanography*, v. 25, PA2208, doi:10.1029/2009PA001780.

650 Medina-Elizalde, M., Lea, D.W., and Fantle, M.S., 2008, Implications of seawater Mg/Ca
651 variability for Plio-Pleistocene tropical climate reconstruction: *Earth and Planetary*
652 *Science Letters*, v. 269, p. 584–594.

653 Minoletti, F., Hermoso, M., and Gressier, V., 2009, Separation of sedimentary micron-sized
654 particles for palaeoceanography and calcareous nannoplankton biogeochemistry:
655 *Nature Protocols*, v. 4(1), p. 14 - 24.

656 Molnar, P., and Cane, M.A., 2002, El Niño's tropical climate and teleconnections as a
657 blueprint for pre-Ice Age climate: *Paleoceanography*, v. 172, p. 1-13.

658 Molnar, P., and Cronin, T.W., 2015, Growth of the maritime continent and its possible
659 contribution to recurring Ice Ages: *Paleoceanography*, v. 30, p. 196–225,
660 doi:10.1002/2014PA002752.

661 Montes, C., Cardona, A., McFadden, R., Morón, S., Silva, J.C., Restrepo-Moreno, S.,
662 Ramirez, D., Hoyos, N., Wilson, J. Farris, D., Bayona, G., Jaramillo, C., Valencia, V.,
663 Bryan, J., and Flores, J.A., 2012, Evidence for middle Eocene and younger land
664 emergence in Central Panama: Implications for Isthmus closure: *Geological Society of*
665 *America, Bulletin*, v. 124, p. 780-799.

666 Montes, C., Cardona, A., Jaramillo, C., Pardo, A., Silva, J.C., Valencia, V., Ayala, C., Pérez-
667 Angel, L.C., Rodriguez-Parra, L.A., Ramirez, V., and Niño, H., 2015, Middle
668 Miocene closure of the Central American Seaway: *Science*, v. 348, p. 226-229, doi:
669 10.1126/science.aaa2815.

670 Mudelsee, M., and Raymo, M.E., 2005, Slow dynamics of the Northern Hemisphere
671 glaciation: *Paleoceanography*, v. 20, PA4022, doi:10.1029/2005PA001153.

672 Murray, A.E., Blakis, A., Massana, R., Strawzewski, S., Passow, U., Alldredge, A., and
673 Delong, E.F., 1999, A time series assessment of planktonic archaeal variability in the
674 Santa Barbara Channel: *Aquatic Microbial Ecology*, v. 20, p. 129–145.

675 Nathan, S.A., and Leckie, R.M., 2009, Early history of the Western Pacific Warm Pool during
676 the middle to late Miocene (~13,2-5,8 Ma): Role of sea-level change and implications
677 for equatorial circulation: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 274,
678 p. 140-159.

679 Neelin, J.D., Battisti, D.S., Hirst, A.C., Jin, F.-F., Wakata, Y., Yamagata, T., and Zebiak,
680 S.E., 1998, ENSO Theory: *Journal of Geophysics Research*, v. 103, p. 14,261–14,290.

681 Osborne, A.H., Newkirk, D.R., Groeneveld, J., Martin, E.E., Tiedemann, R., and Frank, M.,
682 2014, The seawater neodymium and lead isotope record of the final stages of Central
683 American Seaway closure: *Paleoceanography*, v. 29(7), p. 715–729,
684 doi:10.1002/2014PA002676.

685 Pagani, M., Liu, Z., LaRiviere, J., and Ravelo, A.C., 2010, High Earth-system climate
686 sensitivity determined from Pliocene carbon dioxide concentrations: *Nature*
687 *Geoscience*, v. 3, p. 27-30.

688 Pälike, H., Lyle, M., Nishi, H., Raffi, I., Gamage, K., Klaus, A., and the Expedition 320/321
689 Scientists, 2010, Site 1338: Proceedings of the International Ocean Drilling Program
690 320/321, Expedition Reports, v. 321, 141 p.

691 Pälike, H., Lyle, M.W., Nishi, H., Raffi, I., Ridgwell, A., Gamage, K., Klaus, A., Acton, G.,
692 Anderson, L., Backman, J., Baldauf, J., Beltran, C., Bohaty, S.M., Bown, P., Busch,
693 W., Channell, J.E., Chun, C.O., Delaney, M., Dewangan, P., Dunkley Jones, T.,
694 Edgar, K.M., Evans, H., Fitch, P., Foster, G.L., Gussone, N., Hasegawa, H., Hathorne,
695 E.C., Hayashi, H., Herrle, J.O., Holbourn, A., Hovan, S., Hyeong, K., Iijima, K., Ito,
696 T., Kamikuri, S., Kimoto, K., Kuroda, J., Leon-Rodriguez, L., Malinverno, A., Moore,
697 T.C.J.R., Murphy, B.H., Murphy, D.P., Nakamura, H., Ogane, K., Ohneiser, C.,
698 Richter, C., Robinson, R., Rohling, E.J., Romero, O., Sawada, K., Scher, H.,
699 Schneider, L., Sluijs, A., Takata, H., Tian, J., Tsujimoto, A., Wade, B.S., Westerhold,
700 T., Wilkens, R., Williams, T., Wilson, P.A., Yamamoto, Y., Yamamoto, S., Yamazaki,
701 T., and Zeebe, R.E., 2012, A Cenozoic record of the equatorial Pacific carbonate
702 compensation depth: *Nature*, v. 488(7413), p. 609–614, doi:10.1038/nature11360.

703 Pearson, P.N., Van Dongen, B.E., Nicholas, C.J., Pancost, R.D., Schouten, S., Singano, J.M.,
704 and Wade, B.S., 2007, Stable warm tropical climate through the Eocene epoch:
705 *Geology*, v. 35(3), p. 211-214.

706 Pérez-Asensio, J.N., Schmiedl, G., Aguirre, J., and Civis, J., 2012, Impact of restriction of the
707 Atlantic-Mediterranean gateway on the Mediterranean Outflow Water and eastern
708 Atlantic circulation during the Messinian: *Paleoceanography*, v. 27 (3), PA3222,
709 doi:10.1029/2012PA002309.

710 Philander, S., and Fedorov, A., 2003, Role of tropics in changing the response to Milankovich
711 forcing some three million years ago: *Paleoceanography*, v. 18,
712 doi:10.1029/2002PA000837.

713 Pisias, N.G., Mayer, L.A., Janecek, T.R., Palmer-Julson, A., and Van Andel, T.H., 1995,
714 *Proceedings of the Ocean Drilling Program, Scientific results*, v. 138.

715 Prah, F.G., Muehlhausen, L.A., and Zahnle, D.L., 1988, Further evaluation of long-chain
716 alkenones as indicators of paleoceanographic conditions: *Geochimica et*
717 *Cosmochimica Acta*, v. 52, p. 2303–2310.

718 Ravelo, A.C., and Fairbanks, R.G., 1992, Oxygen isotopic composition of multiple species of
719 planktonic foraminifera: recorders of the modern photic zone temperature gradient:
720 *Paleoceanography*, v. 7, p. 815-831.

721 Ravelo, A.C., and Shackleton, N.J., 1995, Evidence for surface-water circulation changes at
722 Site 851 in the eastern tropical Pacific Ocean: *Proceedings of the Ocean Drilling*
723 *Program, Scientific Results*, v. 138, p. 503-514.

724 Ravelo, A.C., Andreasen, D., Lyle, M., Olivarez, L., and Wara, M.W., 2004, Regional climate
725 shifts caused by gradual global cooling in the Pliocene Epoch: *Nature*, v. 429, p. 263-
726 267.

727 Ravelo, A.C., Dekens, P.S., and McCarthy, M., 2006, Evidence for El Niño-like conditions
728 during the Pliocene: *Geological Society of America Today*, v. 16(3), p. 4-11.

729 Raymo, M.E., Grant, B., Horowitz, M., and Rau, G.H., 1996, Mid-Pliocene warmth: stronger
730 greenhouse and stronger conveyor: *Marine Micropaleontology*, v. 27, p. 313-
731 326, doi:10.1016/0377-8398(95)00048-8.

732 Rickaby, R.E.M., and Halloran, P., 2005, Cool La Niña the warmth of the Pliocene: *Science*,
733 v. 307, p. 1948-1952.

734 Rommerskirchen, F., Condon, T., Mollenhauer, G., Dupont, L., and Schefuss, E., 2011,
735 Miocene to Pliocene development of surface and subsurface temperatures in the
736 Benguela Current system: *Paleoceanography*, v. 26, 15 p.

737 Roth, P.H., 1986, Mesozoic paleoceanography of the North Atlantic and Tethys Oceans,
738 North Atlantic: *Paleoceanography*, v. 21, p. 299-320.

739 Rousselle, G., Beltran, C., Sicre, M.-A., Raffi, I., and De Raféllis, M., 2013, Sea-surface
740 condition changes in the Equatorial Pacific during the Mio-Pliocene as inferred from
741 coccoliths geochemistry: *Earth and Planetary Science Letters*, v. 361, p. 412-421.

742 Rühlemann, C., Mulitza, S., Lohmann, G., Paul, A., Prange, M., and Wefer, G., 2004,
743 Intermediate depth warming in the tropical Atlantic related to weakened thermohaline
744 circulation: Combining paleoclimate data and modelling results for the last
745 deglaciation: *Paleoceanography*, v. 19, 10 p.

746 Sato, K., Oda, M., Chiyonobu, S., Kimoto, K., Domitsu, H., and Ingle, J.C., 2008,
747 Establishment of the western Pacific warm pool during the Pliocene: Evidence from
748 planktic foraminifera, oxygen isotopes, and Mg/Ca ratios: *Palaeogeography,*
749 *Palaeoclimatology, Palaeoecology*, v. 265, p. 140-147.

750 Schneider, B., and Schmittner, A., 2006, Simulating the impact of the Panamanian seaway
751 closure on ocean circulation, marine productivity and nutrient cycling: *Earth and*
752 *Planetary Science Letters*, v. 246, p. 367-380.

753 Schouten, S., Hopmans, E.C., Schefuss, E., and Sinninghe Damsté, J.S., 2002, Distributional
754 variations in marine crenarchaeotal membrane lipids: a new tool for reconstructing
755 ancient sea water temperatures?: *Earth and Planetary Science Letters*, v. 204, p. 265-
756 274.

757 Schouten, S., Huguet, C., Hopmans, E.C., Kienhuis, M.V.M., and Sinninghe Damsté, J.S.,
758 2007, Improved analytical methodology and constraints on analysis of the TEX₈₆
759 paleothermometer by high performance liquid chromatography/atmospheric pressure
760 chemical ionization-mass spectrometry: *Analytical Chemistry*, v. 79, p. 2940-2944.

761 Schouten, S., Hopmans, E.C., and Sinninghe Damsté, J.S., 2013, The organic geochemistry of
762 glycerol dialkyl glycerol tetraether lipids: a review: *Organic geochemistry*, v. 54, p.
763 19-61.

764 Scroxton, N., Bonham, S.G., Rickaby, R.E.M., Lawrence, S.H.F., Hermoso, M., and
765 Haywood, A.M., 2011, Persistent El Niño–Southern Oscillation variation during the
766 Pliocene Epoch: *Paleoceanography*, v. 26, PA2215, doi:10.1029/2010PA002097.

767 Seki, O., Foster, G.L., Schmidt D.N., Mackensen, A., Kawamura, K., and Pancost, R.D.,
768 2010, Alkenone and boron-based Pliocene pCO₂ records: *Earth and Planetary Science*
769 *Letters*, v. 292, p. 201-211, doi:10.1016/j.epsl.2010.01.037.

770 Seki, O., Schmidt, D.N., Schouten, S., Hopmans, E.C, Sinninghe Damsté, J.S., and Pancost,
771 R.D., 2012, Paleoceanographic changes in the Eastern Equatorial Pacific over the last
772 10 Myr: *Paleoceanography*, v. 27, PA3224, doi:10.1029/2011PA002158.

773 Sepulchre, P., Arsouze, T., Donnadieu, Y., Dutay, J.C., Jaramillo, C., Le Bras, J., Martin, E.,
774 Montes, C., and Waite, A., 2014, Consequences of shoaling of the Central American
775 Seaway determined from modeling Nd isotopes: *Paleoceanography*, v. 29, p. 176-189,
776 doi:10.1002/2013PA002501.

777 Sinninghe Damsté, J.S., Hopmans, E.C., Schouten, S., Van Duin, A.C.T., and Geenevasen,
778 J.A.J., 2002, Crenarchaeol: the characteristic core glycerol dibiphytanyl glycerol
779 tetraether membrane lipid of cosmopolitan pelagic crenarchaeota: *Journal of Lipid*
780 *Research*, v. 43, p. 1641–1651.

781 Spero, H.J., Mielke, K.M., Kalve, E.M., Lea, D.W., and Pak, D.K., 2003, Multispecies
782 approach to reconstructing eastern equatorial Pacific thermocline hydrography during
783 the past 360 kyr: *Paleoceanography*, v. 18(1), p. 1-16.

784 Srinivasan, M.S., and Sinha, D.K., 1998, Early Pliocene closing of the Indonesian Seaway:
785 evidence from north-east Indian Ocean and Tropical Pacific deep sea cores: *Journal of*
786 *Asian Earth Sciences*, v. 16(1), p. 29-44.

787 Steph, S., Tiedemann, R., Prange, M., Groeneveld, J., Schulz, M., Timmermann, A.,
788 Nürnberg, D., Rühlemann, C., Saukel, C., and Haug, G.H., 2010, Early Pliocene
789 increase in thermohaline overturning: A precondition for the development of the
790 modern equatorial Pacific cold tongue: *Paleoceanography*, v. 252, 17 p.

791 Takayama, T., 1993, Notes on Neogene calcareous nannofossil biostratigraphy of the Ontong
792 Java Plateau and size variations of *Reticulofenestra* coccoliths, *in* Berger, W.H.,
793 Kroenke, L.W., Mayer, L.A., et al. eds., *Proceedings of the Ocean Drilling Program:*
794 *Scientific Results*, v. 130, p. 179–229.

795 Ternois, Y., Sicre, M.-A., and Paterne, M., 2000, Climatic changes along the northwestern
796 African continental margin over the last 30 kyrs: *Geophysical Research Letters*, v.
797 27(1), p. 133-136.

798 Tian, B., Zhang, G.J., and Ramanathan, V., 2001, Heat balance in the Pacific warm pool
799 atmosphere during TOGA COARE and CEPEX: *Journal of Climate*, v. 14(8), p.
800 1881–1893.

801 Tierney, J.E., and Tingley, M.P., 2015, A TEX₈₆ surface sediment database and extended
802 Bayesian calibration: Scientific Data, 2150029, doi: 10.1038/sdata.2015.29.

803 Tierney, J.E., and Tingley, M.P., 2018, BAYSPLINE: A new calibration for the alkenone
804 paleothermometer: *Paleoceanography and Paleoclimatology*, v. 33, p. 281–301,
805 doi:10.1002/2017PA003201.

806 Timmermann, A., An, S.I., Krebs, U., and Goosse, H., 2005, ENSO Suppression due to
807 weakening of the North Atlantic thermohaline circulation: *Journal of Climate*, v. 18, p.
808 3122-3139.

809 Wakeham, S.G., Peterson, M.L., Hedges, J.I., and Lee, C., 2002, Lipid biomarker fluxes in
810 the Arabian Sea, with a comparison to the equatorial Pacific Ocean: *Deep-Sea*
811 *Research II*, v. 49, p. 2265–2301.

812 Wakeham, S.G., Hopmans, E.C., Schouten, S., and Sinninghe Damsté J.S., 2004, Archaeal
813 lipids and anaerobic oxidation of methane in euxinic water columns: A comparative
814 study of the Black Sea and Cariaco Basin: *Chemical Geology*, v. 205, p. 427-442.

815 Wara, M.W., Ravelo, A.C., and Delaney, M.L., 2005, Permanent El Niño-Like Conditions
816 During the Pliocene Warm Period: *Science*, v. 309, p. 758-761.

817 Williams, T., Van De Flierdt, T., Hemming, S.R., Chung, E., Roy, M., and Goldstein, S.L.,
818 2010, Evidence for iceberg armadas from East Antarctica in the Southern Ocean
819 during the late Miocene and early Pliocene: *Earth and Planetary Science Letters*, v.
820 290, p. 351–361.

821 Woodruff, F., and Savin, S.M., 1989, Miocene deep water oceanography: *Paleoceanography*,
822 v. 4(1), p. 87-140.

823 Wuchter, C., Schouten, S., Wakeham, S.G., and Sinninghe Damsté, J.S., 2005, Temporal and
824 spatial variation in tetraether membrane lipids of marine Crenarchaeota in particulate

825 organic matter: Implications for TEX₈₆ paleothermometry: *Paleoceanography*, v. 20,
826 PA3013, doi:10.1029/2004PA001110.

827 Yin, J.H., and Battisti, D. S., 2001, The importance of tropical sea surface temperature
828 patterns in simulations of Last Glacial Maximum climate: *Journal of Climate*, v. 14, p.
829 565–581.

830 Zhang, Y.G., Pagani, M., and Liu, Z., 2014, A 12-million-year temperature history of the
831 tropical Pacific Ocean: *Science*, v. 344, p. 84-86.

832

833

FIGURE CAPTIONS

834 **Figure 1. A)** Modern equatorial Pacific sea-surface temperatures (SSTs) from Ocean Data
835 View. The white dots show the location of the study sites, ODP 806 and IODP U1338. The
836 gray dots indicate the sites of previously published oxygen-isotope records (ODP 847 and
837 1241). WPWP, Western Pacific Warm Pool; EPWP, Eastern Pacific Warm Pool; NEC, North
838 Equatorial Current; SEC, South Equatorial Current; EUC, Equatorial Under Current; NECC,
839 North Equatorial Counter Current, PC, Peru Current (modified after Pisias et al., 1995). **B)**
840 Cross sections of temperature (colors) and currents (vectors) averaged over 2.5°S-2.5°N
841 (modified after US CLIVAR).

842

843 **Figure 2.** Comparison of El Niño-like (in red) and La Niña-like (in blue) time periods
844 inferred from previous studies using different proxies from the EEP and WEP over the past 12
845 Myrs.

846

847 **Figure 3. A)** $\delta^{18}\text{O}$ records in the EEP. $\delta^{18}\text{O}_{\text{Noelaerhabdaceae}}$ at IODP Site U1338 from Rousselle
848 et al. (2013), $\delta^{18}\text{O}_{G. \text{sacculifer}}$ and $\delta^{18}\text{O}_{G. \text{tumida}}$ at site 847 from Wara et al. (2005), $\delta^{18}\text{O}_{G. \text{menardii}}$
849 at Site U1338. **B)** $\delta^{18}\text{O}$ records in the WEP. $\delta^{18}\text{O}_{\text{Noelaerhabdaceae}}$ and $\delta^{18}\text{O}_{G. \text{menardii}}$ at ODP Site
850 806.

851

852 **Figure 4. A)** Temperature records at IODP Site U1338 (EEP): $\text{U}^{\text{K}'}_{37}$ -derived SST and TEX_{86} -
853 derived temperatures. The shaded areas represent the error bars of the temperature estimates
854 (light blue: TEX_{86} -derived temperatures and dark blue: $\text{U}^{\text{K}'}_{37}$ -derived SST). The arrows (black
855 and blue) highlight the temperature trends. **B)** Isotopic gradient ($\Delta\delta^{18}\text{O}$) between surface and
856 thermocline waters at IODP site U1338 (EEP) and ODP Site 806 (WEP). The evolution of the
857 thermocline depth in the EEP and the WEP over the last 10 Ma according to the $\Delta\delta^{18}\text{O}$ is also

858 shown. C) Temperature records at ODP Site 806: $U^{K'}_{37}$ -derived SST (Pagani et al., 2010) in
859 green, TEX_{86} -derived temperatures (Zhang et al., 2014) in red, $Mg/Ca_{G.ruber}$ -derived SST
860 (Medina-Elizalde et al., 2008) in brown and $Mg/Ca_{G.tumida}$ -derived temperature (Ford et al.,
861 2015) in orange.