



Landfast ice: a major driver of reproductive success in a polar seabird

Sara Labrousse, Alexander Fraser, Michael Sumner, Frédéric Le Manach,
Christophe Sauser, Isabella Horstmann, Eileen Devane, Karine Delord, Stéphanie
Jenouvrier, Christophe Barbraud

► To cite this version:

Sara Labrousse, Alexander Fraser, Michael Sumner, Frédéric Le Manach, Christophe Sauser, et al.. Landfast ice: a major driver of reproductive success in a polar seabird. *Biology Letters*, 2021, 17 (6), pp.20210097. <10.1098/rsbl.2021.0097>. <hal-03263429>

HAL Id: hal-03263429

<https://hal.science/hal-03263429v1>

Submitted on 17 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Landfast ice: a major driver of reproductive success in a polar seabird

Biology Letters

Sara LABROUSSE^{1,2*}, Alexander D. FRASER^{3,4}, Michael SUMNER⁵, Frédéric LE MANACH⁶, Christophe SAUSER⁷,
Isabella HORSTMANN², Eileen DEVANE², Karine DELORD⁷, Stéphanie JENOUVRIER^{2#}, Christophe BARBRAUD^{7#}

1 Sorbonne Universités, UPMC Univ., Paris 06, UMR 7159, LOCEAN-IPSL F-75005, Paris, France

2 Department of Biology, Woods Hole Oceanographic Institution, Woods Hole, MA, USA

3 Australian Antarctic Program Partnership, Institute for Marine and Antarctic Studies, University of Tasmania,
Hobart, Tasmania 7001, Australia

4 Institute for Marine and Antarctic Studies, University of Tasmania, Hobart 7001, Tasmania, Australia

5 Australian Antarctic Division, Channel Highway, Kingston, Tasmania, 7050, Australia

6 BLOOM, 62bis avenue Parmentier, 75011 Paris, France

7 Centre d'Études Biologiques de Chizé (CEBC), CNRS UMR 7372, 79360 Villiers en Bois, France

These authors contributed equally: S. Jenouvrier, C. Barbraud.

***Corresponding author**

Email: sara.labrousse@gmail.com

Keywords

Emperor penguin; sea ice; breeding success; climate window analysis; non-linear effect

24 **Abstract**

25

26 In a fast-changing world, polar ecosystems are threatened by climate variability. Understanding the roles of
27 fine-scale processes, linear and non-linear effects of climate factors on the demography of polar species is
28 crucial for anticipating the future state of these fragile ecosystems. While the effects of sea ice on polar marine
29 top predators are increasingly being studied, little is known about the impacts of landfast ice on this species
30 community. Based on a unique 39-year time-series of satellite imagery, *in situ* meteorological conditions and
31 on the world's longest dataset of emperor penguin *Aptenodytes forsteri* breeding parameters, we studied the
32 effects of fine-scale variability of landfast ice and weather conditions on this species reproductive success. We
33 found that longer distances to the landfast ice edge (i.e. foraging areas) negatively affected the overall
34 breeding success but also the fledging success. Climate window analyses suggested that chick mortality was
35 particularly sensitive to landfast ice variability between August and November. Snowfall in May also affected
36 hatching success. Given the sensitivity of landfast ice to storms and changes in wind direction, important future
37 repercussions on the breeding habitat of emperor penguins are to be expected in the context of climate
38 change.

39

1. Introduction

Polar ecosystems are subject to local and regionally contrasted sea ice trends due to climate change [1,2]. Given the complexity of these trends, which are tightly linked to the atmosphere and the ocean dynamics, there is an urgent need to measure and forecast how polar marine populations will respond to sea ice habitat changes [3,4]. Among the studies that have investigated the impacts of climate change and variability on population dynamics in the Southern Ocean [5,6], a thorough understanding of the fine-scale processes by which climate affects the population dynamics of polar organisms is still lacking, thereby preventing the scientific community from improving model projections to correctly assess the future states of polar populations and ecosystems. Given that population dynamics are driven by several demographic components whose sensitivities to climatic factors vary [7,8], it is important to investigate the links between climate and each demographic component. Determining the spatial and temporal scales at which climate variability affects biological parameters is also of prime importance [9]. Also crucial for improving projections, long-term multi-decadal biological series are required to detect non-linear effects of climate on populations [10–13]. The obtention of such long time-series is however often limited by logistical challenges associated with conducting long-term studies in these remote and extreme areas.

Many Antarctic marine top predators, such as seals and seabirds, are intricately linked to landfast ice (LFI), i.e. the narrow band of coastal, compact sea ice held in place by ice shelves and grounded icebergs [14], throughout their breeding period [15–17]. Therefore, LFI variability, such as extreme extent or early break up, can profoundly impact their breeding areas and breeding success [19,20]. However, functional relationships between LFI variability and demographic parameters of polar marine predators remain poorly known due to the scarcity of biological datasets and the difficulty to characterize LFI variability over long time periods.

To improve our understanding of how polar species will respond to future climate changes, we explored the role LFI variability and *in situ* meteorological conditions have on the overall breeding success, but also the fledging and hatching success of a unique sea ice sentinel species [20], the emperor penguin (*Aptenodytes forsteri*). We used the longest historical time-series of Antarctic LFI collected by the Advanced Very High

66 Resolution Radiometer (AVHRR) and the Moderate-Resolution Imaging Spectroradiometer (MODIS), covering
67 the years 1979 to 2017, i.e. since the inception of modern satellite monitoring. We also used the world's
68 longest time-series of emperor penguin breeding parameters, collected at Pointe Géologie, Adélie Land, since
69 1952. The novelty of this research, while relying on previous studies (e.g. [21–26]), lies in i) assessing the
70 climate effect on different components of the reproduction, ii) using the longest time-series available for LFI
71 and emperor penguin reproduction, iii) taking into account the relative contribution of fine-scale processes
72 (local LFI and *in situ* meteorological conditions), iv) exploring different time windows of these effects, and v)
73 testing non-linear effects.

74

75 **2. Material and methods**

76

77 (a) Landfast ice data

78 Three sources of satellite imagery were used to cover the 1979–2017 period and aggregate LFI data (electronic
79 supplementary material; see figure S1 for examples):

80 1) 1979–1991: visible (when available) or thermal infrared images from AVHRR's Global Area Coverage

81 (GAC) mode (spatial resolution of 4 kilometres per pixel; km/px).

82 2) 1992–1999: visible (when available) or thermal infrared images from the AVHRR Coastal Atlas of East

83 Antarctica [27] (resolution of 1.1 km/px).

84 3) 2000–2017: LFI maps from Moderate-Resolution Imaging Spectroradiometer (MODIS) images

85 (resolution of 1 km), classified by Ref. [28].

86 Distances between the penguin colony location and the nearest landfast ice edge (LFIE) (i.e., proxy for access

87 to the ocean) and landfast ice areas (LFIA) were extracted from the images.

88

89 (b) Meteorological data

90 Meteorological data were obtained from the French weather station of Dumont D'Urville. Three

91 parameters were used in this study: the number of days per month with i) temperatures under -10°C, ii) winds

above 28 m/s, and iii) snowfall. We hypothesised that egg loss during incubation and chick mortality could be enhanced during cold and windy conditions caused by katabatic winds and winter storms [15]. Heat loss due to cold temperatures and strong winds, which could be enhanced by snowfall, may increase chick mortality.

(c) Reproductive data

Data are similar to those used by Refs. [21,29] with updated estimates (electronic supplementary material). From count data we estimated 'breeding success' as the number of fledged chicks divided by the number of breeding pairs; 'hatching success' as the number of breeding pairs minus the number of dead eggs divided by the number of breeding pairs; and 'fledging success' as the number of fledged chicks divided by the number of breeding pairs minus the number of dead eggs. Breeding success was estimated over the period 1979–2017; hatching and fledging success over the period 1983–2017.

(d) Climate window analysis

We performed a 'climate window analysis' using the R package *climwin*, following the steps described in Ref. [30]. Climate window analyses determine, without any *a priori* hypothesis, the best climate window(s) (i.e. candidate models) that identify potential climate signals between biological and climate data. Two datasets were analysed: one that contained our monthly climate data, i.e. landfast ice or meteorological data covering the 1979–2017 period, and one that contained information on the response variable, i.e. breeding, hatching, and fledging success. For each climate window, a model was computed. Akaike Information Criteria (AICs) were used for ranking and comparing different candidate climate windows, and then for assessing the best models, their uncertainty, explanatory power, and applicability. Details on the analysis and outputs of the analysis are provided in the electronic supplementary material. The full dataset and codes can be found on Dryad [31].

3. Results

(a) Reproduction time-series

Hatching success was the most stable reproductive parameter (mean \pm SD = 0.82 ± 0.07 , CV = 8.3 %), while fledging success (0.65 ± 0.30) and breeding success (0.53 ± 0.25) were more variable (CV = 46.4 % and 46.5 %, respectively; figure 1a). Hatching success increased during the study period (slope = 0.051 ± 0.007 (SE), $p < 0.001$), while fledging success (slope = -0.003 ± 0.052 , $p = 0.96$) and breeding success (0.026 ± 0.040 , $p = 0.52$) remained stable (figure 1a). Variations in both fledging and breeding successes seemed to co-vary with the LFIA, but even more so with the distance to the nearest LFIE (figure 1b-c).

(b) Climate window analysis

Breeding success was higher for shorter distances to the LFIE between August and November ($p_{\text{randomization}} = 0.006$; adjusted $R^2 = 0.4$), while the LFIA did not have a significant influence (i.e. based on the randomization test; table 1, figure 2a). The number of days per month with temperatures under -10°C , with winds above 28 m/s, and with snowfall did not influence the breeding success (table 1). Neither the LFIA nor the number of days per month with winds above 28 m/s or temperature below -10°C had an influence on the hatching success (table 1). However, the hatching success appeared to be influenced by the number of days with snowfall in May ($p_{\text{randomization}} = 0.0003$, adjusted $R^2 = 0.3$; table 1, figure 2c). This relationship was non-linear, with hatching success increasing with the proportion of days with snowfall per month up to 37% and remaining stable or decreasing slightly for higher proportions. Finally, fledging success was higher for shorter distances to the LFIE in November ($p_{\text{randomization}} = 0.035$, adjusted $R^2 = 0.5$; table 1, figure 2b), while the LFIA, the number of days per month with temperature below -10°C , with winds above 28 m/s, and with snowfall did not have a significant influence (table 1). Fledging success declined non-linearly with the nearest distance to the LFIE, with an accelerated decline for distances greater than ca. 50 km.

4. Discussion

142 We showed that, over 39 years, different components of the reproduction of an Antarctic seabird were
143 affected by fine-scale LFI and *in situ* meteorological conditions at different times of its breeding season, and,
144 importantly, these relationships were non-linear.

145 Adult emperor penguins during the breeding season forage and hunt by diving at the edge of the LFI
146 in cracks, flaw leads, and polynyas [32]. Longer distances between the colony and foraging grounds accessed
147 by the LFIE imply lower chick-feeding frequency, and thus lower chick growth with negative consequences on
148 fledging and breeding success. Using historical AVHRR and recent MODIS images, our study brings important
149 and novel results. First, we identified that distance to nearest LFIE particularly affected fledging success in
150 November (and the second-best model identified a window between August and November), indicating that
151 chick mortality was the main cause of declining breeding success with increasing distance to LFIE. Second, this
152 relationship was nonlinear, with over 50% chick mortality when the distance to LFIE exceeded ca. 65 km. Non
153 linearity could be detected by extending the time series from 8 years in a previous work [26] to nearly 40 years
154 in our study. Third, we identified that the best climate window explaining the relationships between distance
155 to LFIE (i.e. foraging grounds) and breeding success was between August and November, suggesting chicks
156 were particularly sensitive to environmental variability during this period of high energetic demands for body
157 growth [33,34].

158 Reproduction has been monitored at extremely few other emperor penguin colonies. Surprisingly, no
159 relationship was found between LFI and breeding success of emperor penguins at Taylor Glacier colony [35].
160 Although this may depict the complex interactions between environment and penguin foraging behaviour and
161 their consequences for breeding performances, ref. [35] used distance to LFIE in April and September, and our
162 time windows analysis indicated that these months did not represent the full critical period for fledging and
163 breeding success. Nevertheless, this highlights the need to monitor multiple sites in order to understand how
164 sea ice variability, and especially LFI, is affecting the global emperor penguin population.

165 Our study supports previous findings that it is crucial to consider both fine-scale climate processes and
166 fine-scale temporal windows when investigating the relationships between climate variability and
167 demographic traits [9,36]. Despite the diversity of studies that have investigated the effect of climate change

on polar species, there is a strong need to account for the factors that control population dynamics at local/regional scales in order to understand how they may modulate the effects of large-scale environmental variations on long-term population trend [13]. For example, ref. [37] compared the influence of environmental factors on the breeding success of snow petrels (*Pagodroma nivea*) at Casey station with the colony of Adélie Land, and showed that despite similarities in the biological processes controlling snow petrel breeding success, the correlation of large-scale environmental factors with breeding success differed substantially between the two colonies, likely due to the effects of the environmental factors at the local/regional scale.

Landfast ice variability may have important indirect effects that we did not consider in this study. For example, LFI break-ups could contribute to the phytoplankton seeding process (e.g. [38–40]) and may drive a phytoplankton bloom associated with trophic cascades. This could in turn benefit emperor penguins through bottom-up processes with a temporal lag depending on the timing within the breeding period. In the Arctic, longer temporal lags between sea ice melting and phytoplankton bloom resulted in rapidly decreasing breeding performance for little auks (*Alle alle*) and Brünnich's guillemots (*Uria lomvia*) [41]. Thus, considering local to regional-scale phenology in the development of potential phytoplankton blooms in responses to LFI variability may help understand climate-driven environmental impacts on seabirds.

During breeding, individual emperor penguins do not use a fixed nest site as do other penguins. Therefore, the colony is mobile during the breeding season and can move several hundred meters or even a few kilometers. Therefore, the selection of nest site (and experience to nest site) is not relevant for this species. However, there might be selection for sites where colonies are situated, as these sites are generally occupied for long time periods (several decades at least), as our results suggest a strong selection pressure from environmental factors such as LFIE. Nevertheless, the environmental factors affecting colony site selection have not been investigated and quantified to date.


Finally, none of the meteorological variables, except snow falls for the hatching success, had an influence on reproductive parameters. The positive relationship between the number of days with snowfall in May and the hatching success may be associated with the hydration of males during their long fasting period of ca. four months. We speculate that important snowfalls in May allow males to supplement their water

194 intake by eating snow, thus decreasing dehydration potentially leading to the abandonment of the egg before
195 it hatches. Indeed, field observations during winter indicate that male emperor penguins eat snow all along
196 the incubation period ([42]; CB, pers. obs.).

197 Our results bring new insights on the proximate mechanisms through which a poorly known polar
198 habitat feature, LFI, affects demographic parameters of polar top predators. We note that, although we might
199 be able to better predict the future state of polar populations once such fine-scale processes are fully
200 understood, population projections based on sea ice models (e.g. [43]) remain hampered by the fact that these
201 models project sea ice extent but do not provide information on LFI dynamics yet. Important future
202 repercussions on the breeding habitat of emperor penguins and ultimately their persistence are to be
203 expected in the context of climate change [2] given the sensitivity of LFI to storms and changes in wind
204 direction [44], as well as the recently observed strong and opposed LFI trends in adjacent regions [45]. Given
205 the demographic sensitivity of emperor penguins associated with postglacial warming leading to a major
206 southward expansion [46], major shifts such as decline or extinction of emperor penguin populations are
207 expected under anthropogenic climate change.

208

209



213 **Fig. 1** Times-series of the emperor penguin reproductive parameters (panel a; 95% confidence intervals in
214 grey) and LFI conditions (LFIA, panel b; nearest distance to LFIE, panel c) at Pointe Géologie, 1979-2017. Pink
215 rectangles highlight years for which breeding success was below 25%.




Fig. 2 Relationships between emperor penguin reproductive parameters (breeding, fledging, and hatching successes, panels a, b, and c respectively) and climate variables from 1979 to 2017 at Pointe Géologie obtained from the climate window analysis. The best climate window was August to November for (a), November for (b), and May for (c).

222 **Table 1.** Summary of the climate window analysis.

Climate variables	Biological variable	Period considered	Years	Best climate window	<i>p</i> -value best model ¹	<i>p</i> -value after randomisation	Fit selected [alternative fit]	Sign of the relation	R ² after randomisation (k = 10)
Nearest open water (LFIE)	Breeding success	May-Nov.	1979-2017	Aug.- Nov.	1.48e-05	0.006	linear, AIC = -125.8826 [quadratic, AIC = -125.082]	-	0.386
	Hatching success	May-Aug.	1983-2017						
	Fledging success	May-Nov.	1983-2017	Nov.	x = 0.500 x ² = 0.025	0.035	quadratic, AIC = -107.642 [linear, AIC = -105.3069]	-	0.530
Landfast ice area (136°-146° E)	Breeding success	May-Nov.	1979-2017	NS	0.003	0.499		NS	
	Hatching success	May-Aug.	1983-2017	NS	0.007	0.715		NS	
	Fledging success	May-Nov.	1983-2017	NS	0.0001	0.266		NS	
Nb. of days/month with temperatures under -10° C	Breeding success	May-Nov.	1979-2017	NS	NS	/		NS	
	Hatching success	May-Aug.	1983-2017	NS	NS	/		NS	
	Fledging success	May-Nov.	1983-2017	NS	NS	/		NS	
Nb. of days/month with winds above 28 m/s	Breeding success	May-Nov.	1979-2017	NS	NS	/		NS	
	Hatching success	May-Aug.	1983-2017	NS	NS	/		NS	
	Fledging success	May-Nov.	1983-2017	NS	NS	/		NS	
Nb. of days/month with snowfall	Breeding success	May-Nov.	1979-2017	NS	0.044	0.926		NS	
	Hatching success	May-Aug.	1983-2017	May	x = 0.003 x ² = 0.017	0.0003	quadratic, AIC = -198.8435 [linear, AIC = -194.5369]	+ (bell shape)	0.321
	Fledging success	May-Nov.	1983-2017	NS	NS	/		NS	

¹ For quadratic relationships, *p*-values for the linear and quadratic terms are given as x and x² respectively.

223

224

225 **Funding**

226 This work was supported by Exp  and IPEV program 109, WHOI's postdoctoral scholar award, NSF awards
227 1744794 and 1643901, the AAD, the Australian Government's Antarctic Science Collaboration Initiative
228 program, and the Australian Research Council's Special Research Initiative for Antarctic Gateway Partnership
229 (Project ID SR140300001). This study is a contribution to the Project SENSEI funded by Foundation BNP Paribas.

230

231 **Acknowledgments**

232 We warmly thank all the wintering fieldworkers for conducting counts and monitoring at Pointe G ologie since
233 1979, R. Massom for all the initial discussions and D. Iles for methodological advice.

234

- 236 1. Parkinson CL, Cavalieri DJ. 2012 Antarctic sea ice variability and trends, 1979-2010. *The Cryosphere* **6**,
237 871–880. (doi:10.5194/tc-6-871-2012)
- 238 2. Meredith M *et al.* 2019 Chapter 3: Polar Regions. IPCC Special Report on the Ocean and Cryosphere
239 in a Changing Climate.
- 240 3. Post E *et al.* 2009 Ecological Dynamics Across the Arctic Associated with Recent Climate Change.
241 *Science* **325**, 1355–1358. (doi:10.1126/science.1173113)
- 242 4. Massom RA, Stammerjohn SE. 2010 Antarctic sea ice change and variability – Physical and ecological
243 implications. *Polar Sci.* **4**, 149–186. (doi:10.1016/j.polar.2010.05.001)
- 244 5. Constable AJ *et al.* 2014 Climate change and Southern Ocean ecosystems I: how changes in physical
245 habitats directly affect marine biota. *Glob. Change Biol.* **20**, 3004–3025. (doi:10.1111/gcb.12623)
- 246 6. Bestley S *et al.* 2020 Marine Ecosystem Assessment for the Southern Ocean: Birds and Marine
247 Mammals in a Changing Climate. *Front. Ecol. Evol.* **8**, 338. (doi:10.3389/fevo.2020.566936)
- 248 7. Pardo D, Jenouvrier S, Weimerskirch H, Barbraud C. 2017 Effect of extreme sea surface temperature
249 events on the demography of an age-structured albatross population. *Phil Trans R Soc B* **372**, 20160143.
- 250 8. Pacoureau N, Delord K, Jenouvrier S, Barbraud C. 2019 Demographic and population responses of an
251 apex predator to climate and its prey: a long-term study of South Polar Skuas. *Ecol. Monogr.* **89**.
252 (doi:10.1002/ecm.1388)
- 253 9. van de Pol M, Cockburn A. 2011 Identifying the Critical Climatic Time Window That Affects Trait
254 Expression. *Am. Nat.* **177**, 698–707. (doi:10.1086/659101)
- 255 10. Bjørnstad ON, Grenfell BT. 2001 Noisy Clockwork: Time Series Analysis of Population Fluctuations in
256 Animals. **293**, 7.
- 257 11. Mysterud A, Stenseth NC, Yoccoz NG, Langvatn R, Steinheim G. 2001 Nonlinear effects of large-scale
258 climatic variability on wild and domestic herbivores. **410**, 4.
- 259 12. Clark TJ. 2020 Nonlinear population dynamics are ubiquitous in animals. **4**, 17.
- 260 13. Iles DT, Lynch H, Ji R, Barbraud C, Delord K, Jenouvrier S. 2020 Sea ice predicts long-term trends in
261 Adélie penguin population growth, but not annual fluctuations: Results from a range-wide multiscale
262 analysis. *Glob. Change Biol.* **26**, 3788–3798. (doi:10.1111/gcb.15085)
- 263 14. Fraser AD, Massom RA, Michael KJ, Galton-Fenzi BK, Lieser JL. 2012 East Antarctic Landfast Sea Ice
264 Distribution and Variability, 2000–08. *J. Clim.* **25**, 1137–1156. (doi:10.1175/JCLI-D-10-05032.1)
- 265 15. Jouventin P. 1975 Mortality parameters in emperor penguins *Aptenodytes forsteri*. *Biol. Penguins*
266 *Lond. Macmillan*, 435–446.
- 267 16. Ainley D. 2002 *The Adélie Penguin: Bellwether of Climate Change*. Columbia University Press.
- 268 17. LaRue M *et al.* 2019 Physical and ecological factors explain the distribution of Ross Sea Weddell seals
269 during the breeding season. *Mar. Ecol. Prog. Ser.* **612**, 193–208. (doi:10.3354/meps12877)
- 270 18. Barbraud C, Delord K, Weimerskirch H. 2015 Extreme ecological response of a seabird community to
271 unprecedented sea ice cover. *R. Soc. Open Sci.* **2**, 140456. (doi:10.1098/rsos.140456)
- 272 19. Fretwell PT, Trathan PN. 2019 Emperors on thin ice: three years of breeding failure at Halley Bay.
273 *Antarct. Sci.* **31**, 133–138. (doi:10.1017/S0954102019000099)
- 274 20. Boersma PD. 2008 Penguins as Marine Sentinels. *BioScience* **58**, 597–607. (doi:10.1641/B580707)
- 275 21. Barbraud C, Weimerskirch H. 2001 Emperor penguins and climate change. *Nature* **411**, 183–186.
276 (doi:10.1038/35075554)
- 277 22. Jenouvrier S, Barbraud C, Weimerskirch H. 2005 Long-term contrasted responses to climate of two
278 Antarctic seabirds species. *Ecology* **86**, 2889–2903. (doi:10.1890/05-0514)
- 279 23. Jenouvrier S, Weimerskirch H, Barbraud C, Park Y-H, Cazelles B. 2005 Evidence of a shift in the
280 cyclicity of Antarctic seabird dynamics linked to climate. *Proc. R. Soc. B Biol. Sci.* **272**, 887–895.
281 (doi:10.1098/rspb.2004.2978)
- 282 24. Jenouvrier S, Caswell H, Barbraud C, Holland M, Strøve J, Weimerskirch H. 2009 Demographic
283 models and IPCC climate projections predict the decline of an emperor penguin population. *Proc. Natl. Acad.*
284 *Sci.* **106**, 1844–1847.

- 285 25. Jenouvrier S, Holland M, Stroeve J, Barbraud C, Weimerskirch H, Serreze M, Caswell H. 2012 Effects
286 of climate change on an emperor penguin population: analysis of coupled demographic and climate models.
287 *Glob. Change Biol.* **18**, 2756–2770. (doi:10.1111/j.1365-2486.2012.02744.x)
- 288 26. Massom R, Hill K, Barbraud C, Adams N, Ancel A, Emmerson L, Pook M. 2009 Fast ice distribution in
289 Adélie Land, East Antarctica: interannual variability and implications for emperor penguins *Aptenodytes*
290 *forsteri*. *Mar. Ecol. Prog. Ser.* **374**, 243–257. (doi:10.3354/meps07734)
- 291 27. Michael KJ, Hill K, Australia. Antarctic Division, Antarctic CRC, Commonwealth Bureau of
292 Meteorology (Australia). 2003 *Sea ice atlas : East Antarctica : AVHRR imagery 1992-1999, August 2003*.
293 Antarctic CRC.
- 294 28. Fraser AD, Massom RA, Ohshima KI, Willmes S, Kappes PJ, Cartwright J, Porter-Smith R. 2020 High-
295 resolution mapping of circum-Antarctic landfast sea ice distribution, 2000–2018. *Earth System Science Data*
296 **12**, 2987–2999.
- 297 29. Barbraud C, Gavrilov M, Mizin Y, Weimerskirch H. 2011 Comparison of emperor penguin declines
298 between Pointe Géologie and Haswell Island over the past 50 years. *Antarct. Sci.* **23**, 461–468.
299 (doi:10.1017/S0954102011000356)
- 300 30. Bailey LD, van de Pol M. 2016 climwin: An R Toolbox for Climate Window Analysis. *PLOS ONE* **11**,
301 e0167980. (doi:10.1371/journal.pone.0167980)
- 302 31. Labrousse S, Fraser A, Sumner M, Le Manach F, Sauser C, Horstmann I, Devane E, Delord K,
303 Jenouvrier S, Barbraud C. 2021 Data from: Landfast ice: a major driver of reproductive success in a polar
304 seabird. Dryad Digital Repository. (<https://doi.org/10.5061/dryad.cnp5hqc47>)
- 305 32. Labrousse S *et al.* 2019 Dynamic Fine-Scale Sea Icescape Shapes Adult Emperor Penguin Foraging
306 Habitat in East Antarctica. *Geophys. Res. Lett.* **46**, 11206–11218. (doi:10.1029/2019GL084347)
- 307 33. Prevost J. 1961 Expéditions polaires françaises. In (ed P J), pp. 1–204. Hermann Press, Paris, France.
- 308 34. Mougin J-L. 1966 *Observations écologiques à la colonie de manchots empereurs de Pointe Géologie*
309 *(Terre Adélie) en 1964...*
- 310 35. Robertson G, Wienecke B, Emmerson L, Fraser AD. 2014 Long-Term Trends in the Population Size
311 and Breeding Success of Emperor Penguins at the Taylor Glacier Colony, Antarctica. *Polar Biology* **37**, 251–
312 259.
- 313 36. Barreau E, Ropert-Coudert Y, Delord K, Barbraud C, Kato-Ropert A. 2019 Scale matters: sea ice and
314 breeding success of Adélie penguins. *Polar Biol.* **42**, 1405–1410. (doi:10.1007/s00300-019-02531-2)
- 315 37. Olivier F, Franeker JA van, Creuwels JCS, Woehler EJ. 2005 Variations of snow petrel breeding success
316 in relation to sea-ice extent: detecting local response to large-scale processes? *Polar Biol.* **28**, 687–699.
317 (doi:10.1007/s00300-005-0734-5)
- 318 38. McMinn A. 1996 Preliminary investigation of the contribution of fast-ice algae to the spring
319 phytoplankton bloom in Ellis Fjord, eastern Antarctica. *Polar Biol.* **16**, 301–307.
320 (doi:10.1007/s003000050057)
- 321 39. Riaux-Gobin C, Poulin M, Dieckmann G, Labruno C, Vétion G. 2011 Spring phytoplankton onset after
322 the ice break-up and sea-ice signature (Adélie Land, East Antarctica). *Polar Res.* **30**, 5910.
323 (doi:10.3402/polar.v30i0.5910)
- 324 40. Mangoni O, Saggiomo M, Modigh M, Catalano G, Zingone A, Saggiomo V. 2009 The role of platelet
325 ice microalgae in seeding phytoplankton blooms in Terra Nova Bay (Ross Sea, Antarctica): a mesocosm
326 experiment. *Polar Biol.* **32**, 311–323. (doi:10.1007/s00300-008-0507-z)
- 327 41. Ramírez F, Tarroux A, Hovinen J, Navarro J, Afán I, Forero MG, Descamps S. 2017 Sea ice phenology
328 and primary productivity pulses shape breeding success in Arctic seabirds. *Sci. Rep.* **7**, 4500.
329 (doi:10.1038/s41598-017-04775-6)
- 330 42. Le Maho Y. 1977 The emperor penguin: A strategy to live and breed in the cold: Morphology,
331 physiology, ecology, and behavior distinguish the polar emperor penguin from other penguin species,
332 particularly from its close relative, the king penguin. *Am. Sci.* **65**, 680–693.
- 333 43. Jenouvrier S *et al.* 2019 The Paris Agreement objectives will likely halt future declines of emperor
334 penguins. *Glob. Change Biol.* **26**, 1170–1184. (doi:10.1111/gcb.14864)
- 335 44. Zhai MX, Zhao TC, Hui FM. 2019 Anomalous extensive landfast sea ice in the vicinity of Inexpressible

336 Island, Antarctica. *Adv Polar Sci* **30**, 406–411.
337 45. Fraser AD *et al.* 2021 18 year record of circum-Antarctic landfast sea ice distribution allows detailed
338 baseline characterisation, reveals trends and variability. *Cryosphere Discuss.* **2021**, 1–23. (doi:10.5194/tc-
339 2021-121)
340 46. Cole TL *et al.* 2019 Receding ice drove parallel expansions in Southern Ocean penguins. *Proc. Natl.*
341 *Acad. Sci.* **116**, 26690. (doi:10.1073/pnas.1904048116)
342

343