

Effects of elevated temperature and $p\text{CO}_2$ on the respiration, biomineralization and photophysiology of the giant clam *Tridacna maxima*

Chloé Brahmi^{1,*}, Leila Chapron², Gilles Le Moullac³, Claude Soyez³, Benoît Beliaeff³, Claire E. Lazareth⁴, Nabila Gaertner-Mazouni¹ and Jeremie Vidal-Dupiol^{3,5}

¹Univ. Polynésie française, IFREMER, ILM, IRD, EIO UMR 241, F-98702 Faa'a, Tahiti, Polynésie française

²School of Earth Sciences, The Ohio State University, Columbus, OH 43210, USA

³IFREMER, IRD, Institut Louis-Malardé, Univ. Polynésie française, EIO, F-98719 Taravao, Tahiti, Polynésie française, France

⁴Laboratoire de Biologie des Organismes et Ecosystèmes Aquatiques (BOREA) MNHN, CNRS, IRD, SU, UCN, UA, Muséum National d'Histoire Naturelle, 61 Rue Buffon, CP53, 75231, Paris Cedex 05, France

⁵IHPE, Univ. Montpellier, CNRS, Ifremer, Univ. Perpignan Via Domitia, Montpellier France

*Corresponding author: Univ. Polynésie française, EIO UMR 241, BP 6570, F-98702 Faa'a, Tahiti, French Polynesia. Tel: +689 40 866 479. Email: chloe.brahmi@upf.pf

Many reef organisms, such as the giant clams, are confronted with global change effects. Abnormally high seawater temperatures can lead to mass bleaching events and subsequent mortality, while ocean acidification may impact biomineralization processes. Despite its strong ecological and socio-economic importance, its responses to these threats still need to be explored. We investigated physiological responses of 4-year-old *Tridacna maxima* to realistic levels of temperature (+1.5°C) and partial pressure of carbon dioxide ($p\text{CO}_2$) (+800 μatm of CO_2) predicted for 2100 in French Polynesian lagoons during the warmer season. During a 65-day crossed-factorial experiment, individuals were exposed to two temperatures (29.2°C, 30.7°C) and two $p\text{CO}_2$ (430 μatm , 1212 μatm) conditions. The impact of each environmental parameter and their potential synergetic effect were evaluated based on respiration, biomineralization and photophysiology. Kinetics of thermal and/or acidification stress were evaluated by performing measurements at different times of exposure (29, 41, 53, 65 days). At 30.7°C, the holobiont O_2 production, symbiont photosynthetic yield and density were negatively impacted. High $p\text{CO}_2$ had a significant negative effect on shell growth rate, symbiont photosynthetic yield and density. No significant differences of the shell microstructure were observed between control and experimental conditions in the first 29 days; however, modifications (i.e. less-cohesive lamellae) appeared from 41 days in all temperature and $p\text{CO}_2$ conditions. No significant synergetic effect was found. Present thermal conditions (29.2°C) appeared to be sufficiently stressful to induce a host acclimatization response. All these observations indicate that temperature and $p\text{CO}_2$ are both forcing variables affecting *T. maxima*'s physiology and jeopardize its survival under environmental conditions predicted for the end of this century.

Key words: Giant clams, ocean acidification, photosynthetic yield, respiration, symbionts, thermal stress

Editor: Steven Cooke

Received 7 July 2020; Revised 18 January 2021; Editorial Decision 17 May 2021; Accepted 22 May 2021

Cite as: Brahmi C, Chapron L, Le Moullac G, Soyez C, Beliaeff B, Lazareth CE, Gaertner-Mazouni N, Vidal-Dupiol J (2021) Effects of elevated temperature and $p\text{CO}_2$ on the respiration, biomineralization and photophysiology of the giant clam *Tridacna maxima*. *Conserv Physiol* 9(1): coab041; doi:10.1093/conphys/coab041.

Introduction

Anthropocene era is characterized by human activities releasing gigatons of CO₂ in the atmosphere and contributing to induce global climate change (Broeker *et al.*, 1979; Caldeira and Wickett, 2003; Sabine *et al.*, 2004; Zeebe *et al.*, 2008; IPCC, 2014). The transfer of CO₂ from the atmosphere to the ocean results consequently to the increase of dissolved CO₂ in the seawater inducing significant pH decreases. On the other hand, due to the ‘greenhouse gas’ property of CO₂, terrestrial and sea surface temperature rise constantly (i.e. global warming). According to the last Intergovernmental Panel on Climate Change report (IPCC, 2019), warming had already reached +1°C by comparison to the pre-industrial period and may reach +1.5°C around 2040 if the current rate of emission is maintained. Enrichment of dissolved CO₂ in the ocean modifies the carbonate chemistry by decreasing carbonate ion concentration ([CO₃²⁻]) and releasing protons that respectively decrease calcium carbonate saturation state (Ω) and seawater pH, i.e. ocean acidification (Kleypas *et al.*, 1999; Caldeira and Wickett, 2003). Indeed, pH had already decreased by 0.1 pH unit and this drop may reach -0.3 by 2100 (according to the RCP (Representative Concentration Pathway) 8.5 scenario; IPCC, 2019).

CO₃²⁻ is a carbonate ion form involved in biologically controlled calcification process, i.e. biomineralization (Orr *et al.*, 2005) of many marine organisms such as scleractinian corals and molluscs constructing calcium carbonate structures (e.g. exoskeleton, shell, test, spicule). This characteristic of their physiology make them highly sensitive to ocean acidification (Hoegh-Guldberg *et al.*, 2007; Ries *et al.*, 2009; Kroeker *et al.*, 2013; Gazeau *et al.*, 2013; Parker *et al.*, 2013), especially when this skeleton is made of aragonite, which is more sensitive to dissolution than calcite (Morse *et al.*, 2007). Among marine calcifying molluscs (e.g. oysters, mussels, giant clams, abalones, limpets) negative impacts of temperature and *p*CO₂ have been demonstrated on the survival, growth, biomineralization processes and other key physiological functions on different stages of their life cycle (Bougrier *et al.*, 1995; Kurihara, 2008; McClintock *et al.*, 2009; Gazeau *et al.*, 2010; Welladsen *et al.*, 2010; Melzner *et al.*, 2011; Rodolfo-Metalpa *et al.*, 2011; Schwartzmann *et al.*, 2011; Talmage and Gobler, 2011; Liu *et al.*, 2012; Watson *et al.*, 2012; Kurihara *et al.*, 2013; Fitzer *et al.*, 2014; Le Moullac *et al.*, 2016a, b; Meng *et al.*, 2018; Wessel *et al.*, 2018; Avignon *et al.*, 2020). However, effects of each stressor and their potential synergetic effect on giant clam’s physiology are still poorly understood.

Giant clams produce an aragonitic shell composed of a prismatic inner layer and a crossed-lamellar outer layer (Pätzold *et al.*, 1991; Faylona *et al.*, 2011; Agbaje *et al.*, 2017; Gannon *et al.*, 2017). These organisms form an extracellular phototrophic mutualistic symbiosis with dinoflagellates of the Symbiodiniaceae family (Holt *et al.*, 2014; LaJeunesse *et al.*, 2018). The symbionts are hosted in the mantle part exposed to the light, in a tubular system (Z-tubules) directly connected to the stomach (Norton *et al.*, 1992; Holt *et al.*,

2014). The resulting mixotrophic organisms therefore acquire nutrients from heterotrophic (seawater filtration) and photoautotrophic pathways (Klumpp *et al.*, 1992; Hawkins and Klumpp, 1995). Nutrients provided by symbionts (such as glucose; Ishikura *et al.*, 1999) may account for a major part of the giant clam’s energy needs (Klumpp *et al.*, 1992; Klumpp and Griffiths, 1994; Klumpp and Lucas, 1994; Hawkins and Klumpp, 1995; Elfwing *et al.*, 2002; Yau and Fan, 2012; Holt *et al.*, 2014; Soo and Todd, 2014).

Despite their ecological and socio-economic importance, effects of thermal stress and acidification on giant clams still need to be investigated. Thermal stress has shown to decrease fertilization success in *T. maxima* (Armstrong *et al.*, 2020), decrease oxygen production in *Tridacna derasa* and *Tridacna gigas* as well as respiration rates in *T. derasa* species (Blidberg *et al.*, 2000). In contrast, an increase of respiration with a high photosynthetic rate in the holobiont was reported for *Tridacna squamosa* (Elfwing *et al.*, 2001). Thermal stress also reduces the abundance of symbionts in *T. gigas* (Leggat *et al.*, 2003) and *Tridacna crocea* (Zhou *et al.*, 2019) and a decrease of photosynthate export from symbionts to the host (Leggat *et al.*, 2003). In *T. maxima*, heat stress was shown to induce changes in the fatty acid composition, lipid pathways and the overexpression of genes encoding reactive oxygen species (ROS) scavengers (Dubousquet *et al.*, 2016). In symbionts hosted in *T. gigas*, high-light levels and heat stress caused a decrease of symbiont density, cell size and chlorophyll content (Buck *et al.*, 2002). Toonen *et al.* (2012) demonstrated that low pH combined with high nutrient concentration have different impacts on shell growth rate depending on the *Tridacna* species. Moreover, Watson (2015) showed that high-light irradiance condition (i.e. high photosynthetically active radiation level) may limit the impact of *p*CO₂ on shell growth and total animal mass gain. Regarding the potential synergetic effect of temperature and *p*CO₂ parameters, the only study carried out showed that the survival of *T. squamosa* juveniles may decrease by ocean warming and acidification (Watson *et al.*, 2012). To our knowledge, no study had investigated the impacts of both parameters, and their potential synergetic effect, on several key physiological parameters of both *T. maxima* host and its symbionts. To fill this gap, our experimental approach was designed to better understand the physiological mechanisms underlying the response of giant clams to global climate changes.

In French Polynesia, *T. maxima* (Röding, 1798) is one of the most emblematic and patrimonial species. It represents an important food resource for inhabitants of remote atolls and giant clam fishery and aquaculture activities generate substantial incomes for local fishermen and farmers (Van Wynsberge *et al.*, 2016; Andréfouët *et al.*, 2017). However, wild and cultivated giant clam stocks are largely threatened by environmental disturbances such as abnormally high sea surface temperature inducing mass mortality events as reported in Tuamotu atolls by Addessi (2001) and Andréfouët *et al.* (2013). Like the symbiotic coral species, bleaching events (e.g. the symbiosis dissociation) affecting giant clam have been

Table 1: Measured and calculated parameters of seawater for all treatments

Treatments	Temperature (°C)	$p\text{CO}_2$ (μatm)	pH_{NBS}	Salinity (‰)	A_T ($\mu\text{mol/kg SW}$)	DIC ($\mu\text{mol/kg SW}$)	$\Omega_{\text{aragonite}}$
Control	29.2 (± 0.1)	428 (± 21)	8.19 (± 0.01)	35	2422 (± 170)	2071 (± 85)	4.17
Thermal stress	30.7 (± 0.1)	431 (± 25)	8.19 (± 0.01)	35	2409 (± 157)	2051 (± 89)	4.14
Acidification stress	29.2 (± 0.1)	1210 (± 41)	7.81 (± 0.03)	35	2466 (± 90)	2306 (± 34)	1.99
Acidification and thermal stress	30.7 (± 0.1)	1213 (± 29)	7.81 (± 0.03)	35	2423 (± 110)	2252 (± 58)	2.02

Total alkalinity (A_T) is given in mean (\pm SD) based on weekly measurements for each experimental tank and for each condition. $p\text{CO}_2$, $\Omega_{\text{aragonite}}$ and dissolved inorganic carbon (DIC) were calculated using the CO_2SYS software.

recorded several times in the Indo-Pacific Region (Addessi, 2001; Buck *et al.*, 2002; Leggat *et al.*, 2003; Andréfouët *et al.*, 2013, 2015; Junchompoo *et al.*, 2013) in association to seawater temperature increase by a few degrees above the seasonal maximum (Addessi, 2001; Andréfouët *et al.*, 2013). More recently, two mass bleaching events of giant clams occurring in Reao and Tatakoto (Tuamotu islands) were linked to a prolonged exposure to high temperature ($\geq 30^\circ\text{C}$ over several weeks) (Andréfouët *et al.*, 2017; see Supplementary Fig. S1).

Based on a 65-day crossed-factorial experiment, we investigated physiological responses of 4-year-old *T. maxima* to temperature and $p\text{CO}_2$ conditions in the French Polynesian lagoons during the present warmer season and those predicted for 2100 by the IPCC 2014: $+1.5^\circ\text{C}$ (RCP 4.5 scenario) and $+800 \mu\text{atm}$ of CO_2 (RCP 8.5 scenario). The effect of each parameter and their potential synergetic effect were evaluated based on respiration, biomineralization and photophysiology by analyzing the holobiont O_2 production and respiration, growth rate, ultrastructure of the shell, symbiont density and photosynthetic yield. In addition, the kinetics of thermal and acidification stress were assessed by performing analyses at different time of exposure (i.e. 29, 41, 53, 65 days).

Materials and methods

Biological material

Two hundred juvenile *T. maxima* (4-year-old, $\sim 5\text{--}6$ cm height), with brownish/dark-green colour, were collected from the cultivated stock in Reao lagoon (Tuamotu islands). Then, they were exported to the Centre Ifremer du Pacifique in Tahiti where they were acclimatized in outdoor tank continuously renewed with natural, unfiltered, lagoonal seawater providing a constant input of microflora and fauna at natural concentration. Each individual was placed onto a petri dish on which byssal gland further developed for attachment. All individuals directly opened up right after their transfer into the outdoor tank. No visual sign of stress was observed during the acclimation period except for 4 individuals that died

within the first 3 days (corresponding to a 2% mortality rate). Specimens used in this study were collected and held under a special permit (MEI #284) delivered by the French Polynesian government.

Experimental design and rearing system

To study the impact of temperature, $p\text{CO}_2$, and their synergetic effect on the physiology of the giant clams and their symbionts, 4 experimental conditions were set up by applying 2 temperatures (29.2°C and 30.7°C) and 2 levels of $p\text{CO}_2$ ($430 \pm 22 \mu\text{atm}$ and $1212 \pm 35 \mu\text{atm}$). The tested conditions were as follows: (i) control: 29.2°C , $430 \mu\text{atm}$; (ii) acidification stress: 29.2°C , $1212 \mu\text{atm}$; (iii) thermal stress: 30.7°C , $430 \mu\text{atm}$; and (iv) acidification and thermal stress: 30.7°C , $1212 \mu\text{atm}$ (Table 1). Temperature was set and maintained with an electronic controller (Hobby Biotherm Professional) connected to an aquarium heater (Shego). The pH was manipulated by bubbling CO_2 in water tanks. This was controlled by a pH-stat system (Dennerle) that continuously monitored pH (calibrated to NIST scale) and temperature to control the quantity of CO_2 to maintain the desired pH. The light was set to obtain a photosynthetically active radiation of $200 \pm 20 \mu\text{mol}$ of photons $\text{m}^{-2} \cdot \text{s}^{-1}$ on a 12:12 h light/dark photoperiod.

After a 3-week acclimation period in an outdoor tank, 96 clams were randomly distributed in the experimental tanks 1 week before starting the experiment. For each condition, we used a 500-l tank containing 4 tanks of 30 l (ecological replicates) renewed with natural unfiltered lagoonal seawater at a flow rate of 50 l/h providing a constant input of microflora and fauna at natural concentration. Each 30-l tank contained 6 clams (biological replicates). To avoid physiological shock, targeted temperature and $p\text{CO}_2$ were linearly achieved over 7 days. To evaluate the kinetics of the thermal and/or acidification stress, analyses were performed at 4 different times of exposure, i.e. 29, 41, 53 and 65 days. In total, 64 clams were used for data acquisition corresponding to 4 individuals per condition (1 individual per 30-l tank) and per time of exposure.

Monitoring of temperature, pH and water quality

To insure the stability of experimental conditions, temperature and pH parameters were measured twice a day for each tank at 8:00 am and 4:00 pm using a mercury thermometer certified ISO 9001 ($\pm 0.1^\circ\text{C}$ accuracy) and a pH-meter Consort P603 (± 0.01 accuracy). Total alkalinity (TA) was weekly titrated using a 0.01-N HCl solution and a titrator (Schott Titroline Easy). Levels of $p\text{CO}_2$ and aragonite saturation state were calculated from temperature, pH (NBS scale), salinity and mean TA using the CO₂SYS software (van Heuven *et al.*, 2009). All parameters including seawater carbonate chemistry are reported in Table 1.

Holobiont O₂ consumption and production measurements

Giant clams were placed in an ecophysiological measurement system (EMS) to monitor O₂ consumption and production. The EMS consisted of five open-flow chambers. Four giant clams were individually placed into four chambers, while an empty shell was placed into a fifth chamber used as a control. EMS chambers contained water at the same temperature and $p\text{CO}_2$ conditions as in the experimental tanks. The light energy and photoperiod conditions were the same as for the acclimation tanks. Flow rate in all chambers was constantly maintained at 12 l.h⁻¹. Each chamber was equipped with a two-way electromagnetic valve activated by an automaton (FieldPoint National Instruments). When the electro-valve was opened, the water released from the chamber was analyzed for 3 min using an oxygen sensor (OXI 538, Cellox 325, WTW, Weilheim, Germany) to quantify dissolved oxygen. Oxygen measurements were performed over 48 h. The first 8 h of measurement were discarded due to the animal acclimatization to the chamber. In each chamber, the cycle was completed within 3 min: the first 2 min served to stabilize the measurement and an average of oxygen data was performed on the last minute of acquisition. This cycle was followed by another time frame of 3 min in the control chamber following the sequence specimen #1, control, specimen #2, control, specimen #3, control, specimen #4, control.

Respiration rate (RR) and production rate (PR) were calculated from the data obtained during night time and day time, respectively, using differences in oxygen concentrations between the control and experimental chambers. RR and PR = $V(\text{O}_1 - \text{O}_2)$, where O₁ is the oxygen concentration in the control chamber, O₂ is the oxygen concentration in the experimental chamber and V is the water flow rate. RR and PR data were normalized to tissue dry weight. Once normalized, the terminology becomes O₂ consumption for RR and O₂ production for PR; both expressed in mg O₂.h⁻¹.g⁻¹ dry weight.

After O₂ production and O₂ consumption analyses were completed, a piece of the mantle was dissected for further

symbiont fluorescence and density analyses (see Section Fluorescence and density measurements of symbionts). The remaining soft tissues were frozen and lyophilized for RR and PR data normalization.

Fluorescence and density measurements of symbionts

Potential effect of temperature and $p\text{CO}_2$ conditions on the photophysiology of the symbionts was studied by comparing fluorescence yield of photosystem II (PSII) between all experimental conditions. After clams were sacrificed, a 1 × 2-cm mantle fragment was dissected. The tissue fragment was gently swiped using tissue paper to remove excess mucus and symbionts were collected by doing 5 smears using a sterilized razor blade. Collected symbionts were diluted into 5-ml of 0.2- μm filtered seawater and placed at the obscurity for 10 min to inactivate the PSII before light excitation. Samples were homogenized and 3 ml of the homogenate were collected, placed into a quartz-glass cuvette and analyzed with AquaPen fluorometer (APC-100, Photon System Instruments®, Czech Republic) at a 450-nm wavelength. The minimal fluorescence (F_0) and the maximal fluorescence (F_m) were measured. The quantum yield of photosynthesis was calculated as F_v/F_m where F_v is the variable fluorescence ($F_v = F_m - F_0$).

In addition, symbiont densities were evaluated from mantle fragments. For each individual, a circular (5 mm in diameter) piece of mantle was collected using a punch. The piece was weighed, grounded in 0.2- μm filtered seawater and homogenized. Then, 20 μl of the tissue extract were immediately collected and placed into Malassez cells for symbiont counting under optical microscope. For each sample, counting was performed on 4 replicates (3 columns per replicate). Data are expressed in number of symbionts/mg of mantle tissue.

Calcein labelling for evaluating daily shell extension rate

To study the impact of temperature and $p\text{CO}_2$ on shell growth rate, the mineralization front of giant clams was marked using calcein fluorochrome, which is irreversibly precipitated at the CaCO₃ mineralization site. Before the experiment starts, giant clams were immersed in a 100-mg.l⁻¹ calcein solution (Sigma Aldrich) (calcein diluted in 1- μm filtered-seawater) for 8 h in the dark. During the labelling procedure, the bath of calcein solution was aerated using bubblers and water current was created via pumps. Calcein-labelled specimens were then placed into the experimental tanks. At the end of the experiment, for each individual, a 5-mm thick section was cut along the maximal shell growth axis through the right valve using a Swap Top Inland® diamond saw. All sections obtained were polished and observed under epifluorescence with a Leitz Dialux® 22 microscope. The distance between the fluorescent calcein mark and the edge of the shell formed during the experiment was measured following the maximal

growth direction. Daily shell extension rate (expressed in $\mu\text{m}\cdot\text{day}^{-1}$) was obtained by dividing the measured distance by the number of days of incubation in experimental conditions.

Shell scanning electron microscopy study

To characterize temperature and $p\text{CO}_2$ effect on the shell ultrastructure, a scanning electron microscopy (SEM) study was carried out. For each individual, a 10-mm thick shell section was cut facing the section used for calcein observations. The section was then fractured along the width, 2 cm below the growing part of the shell (marginal part), using a small chisel and a hammer. Then, the apical fragment was longitudinally fractured and one piece was sonicated in tap water for 10 s, air dried and an additional drying was done overnight at 35°C . Sample was placed on a stub covered with carbon tape, gold-coated and observed at 15 kV using a Hitachi TM3030 SEM at the Université de la Polynésie française. For each condition, and for each time of exposure, two samples were selected based on their daily shell extension rate, i.e. samples showing the lowest and the highest rate. To evaluate the impact of temperature and/or $p\text{CO}_2$ on the shell ultrastructure, SEM observations were performed for each specimen in two different zones of the crossed-lamellar outer layer. Zone 1 corresponds to the shell formed *in situ* (i.e. in the shell region located before the calcein mark), while zone 2 corresponds to the shell formed during the experiment. Observations were made at the ultrastructural level and focused on the aspect and integrity of the lamellae of the crossed-lamellar outer layer. In addition, to check the potential effect of temperature and $p\text{CO}_2$ on the external shell surface, the periostracum (i.e. the organic layer covering the shell external surface) of two individuals per experimental condition (65 days of exposure) was observed.

Statistical analyses and data processing

Normality of data distribution and homogeneity of variance were tested with the Shapiro–Wilk test and the Bartlett test, respectively. Production and consumption of O_2 data followed the conditions of application of parametric tests, but photosynthetic yields and symbiont densities were transformed using Box Cox transformation while shell extension rates were square root transformed to meet these conditions. Comparisons were done using a three-way ANOVA with interactions (fixed factors: time of exposure, temperature and $p\text{CO}_2$). Tukey post hoc comparisons were done at $\alpha = 0.05$ for all analyses. Correlations between physiological parameters were tested using Pearson method with a threshold of $r = 0.25$ ($\alpha = 0.05$).

For all physiological parameters, i.e. O_2 respiration, O_2 consumption, symbiont photosynthetic yield and density, means ($\pm\text{SD}$) were calculated based on the four biological replicates for each condition and each time of exposure. For the daily shell extension rate, means ($\pm\text{SD}$) were calculated for each condition and each time of exposure based on the




Figure 1: Variations of O_2 production and O_2 consumption acquired over a 48-h period after 29 days of exposure to different temperature/ $p\text{CO}_2$ conditions: (A) 29.2°C , $430 \mu\text{atm}$ of CO_2 ; (B) 30.7°C , $430 \mu\text{atm}$ of CO_2 ; (C) 29.2°C , $1212 \mu\text{atm}$ of CO_2 ; and (D) 30.7°C , $1212 \mu\text{atm}$ of CO_2 . Data are expressed in $\text{mg O}_2\cdot\text{h}^{-1}\cdot\text{g}^{-1}$ tissue dry weight (dw) and correspond to day-time and night-time acquisitions for four replicates per condition (each color line represents one replicate).

number of biological replicates displaying a detectable calcein mark.

Results

Effect of temperature and $p\text{CO}_2$ on the holobiont oxygen balance

For all sampling time (i.e. 29, 41, 53, 65 days), a cyclic pattern of O_2 production and consumption was observed following the circadian cycle (Fig. 1). This pattern corresponds to oxygen photosynthetically produced and heterotrophically consumed by the holobiont during the day and the night, respectively. No mortality was observed in all tested temperature/ $p\text{CO}_2$ conditions during the whole experiment.

Mean values of normalized O_2 production and consumption are shown in Fig. 2. Tukey post hoc test results are




Figure 2: Graphs reporting data of (A) oxygen production, (B) oxygen consumption, (C) symbiont photosynthetic yield, (D) symbiont density and (E) daily shell extension rate obtained for each temperature/pCO₂ experimental condition and time of exposure (black and grey columns correspond to 430 μatm and 1212 μatm, respectively). Data are given in mean (+SD) calculated from four replicates for all physiological parameters except for daily shell extension rate for which numbers of replicates are specified.

reported in Table 3. The ANOVA has indicated that oxygen production of the holobiont during day time was significantly altered at 30.7°C ($P = 0.009$) but not at 1212 μatm of CO₂ ($P = 0.362$) (Table 2). In addition, O₂ production was higher

at 29.2°C than at 30.7°C for all times of exposure (Fig. 2A) and decreased over time ($P = 0.030$, Table 2). The night-time O₂ consumption, however, was not significantly influenced by temperature ($P = 0.590$) neither by pCO₂ ($P = 0.361$) nor

Table 2: Results from the three-way ANOVA performed on holobiont O₂ production and consumption data, symbiont photosynthetic yield, symbiont density and giant clam shell extension rate

		O ₂ production (n = 64)	O ₂ consumption (n = 64)	Photosynthetic yield (n = 64)	Symbiont density (n = 64)	Shell extension rate (n = 55)
Time	F	3.264	0.740	3.566	6.035	4.695
	P	0.030*	0.533	0.021*	0.001***	0.007**
Temperature	F	7.551	0.294	65.200	79.158	0.466
	P	0.009**	0.590	<0.0001***	<0.0001***	0.499
pCO ₂	F	0.846	0.851	6.125	8.665	7.409
	P	0.362	0.361	0.017*	0.005**	0.010**
Time × temperature	F	0.413	0.965	3.393	1.528	0.319
	P	0.744	0.417	0.025*	0.219	0.812
Time × pCO ₂	F	1.765	0.476	0.024	1.033	0.251
	P	0.167	0.700	0.995	0.386	0.860
Temperature × pCO ₂	F	1.334	1.029	0.100	1.989	0.026
	P	0.254	0.315	0.753	0.165	0.872
Time × temperature × pCO ₂	F	1.216	0.148	2.153	0.474	0.336
	P	0.314	0.930	0.106	0.702	0.799

The three fixed factors are pCO₂, temperature and time of exposure. Asterisks denote significant differences (*P ≤ 0.05, **P ≤ 0.01, ***P ≤ 0.001).

by time of exposure ($P = 0.533$) and remained stable throughout the whole experiment (Fig. 2B). No interaction effect between the three tested factors was found to affect the holobiont oxygen balance (Table 2).

Symbiont density and photosynthesis under different temperature/pCO₂ conditions

After 2 months of exposure, the symbiont photosynthetic yield and density were significantly impacted at 30.7°C ($P < 0.0001$), high pCO₂ (1212 μatm) ($P = 0.017$ and $P = 0.005$, respectively) and by the time of exposure ($P = 0.021$ and $P = 0.001$, respectively). During the whole experiment and for both pCO₂ conditions, the means of photosynthetic yield (Fig. 2C) and symbiont density (Fig. 2D) tended to be higher at 29.2°C than at 30.7°C. However, an inter-individual variability was observed.

Concerning the interaction parameters, only interaction between temperature and time of exposure parameters had a significant effect on the symbiont photosynthetic yield ($P = 0.025$) (Table 2). No parameter interaction was found to affect the symbiont density (Table 2).

Effect of pCO₂ on daily shell extension rate

Calcein mark was detectable in 55 over 64 shell sections. Statistical analyses on shell extension rate showed that pCO₂ and the time of exposure had a significant effect on the shell extension rate ($P = 0.010$ and $P = 0.007$, respectively, Table 2).

As shown in Fig. 2E, the mean values of shell extension rates tended to be lower at high pCO₂.

Effect of long-term exposure to temperature and pCO₂ on the periostracum and the shell microstructure

As a first approach, SEM observations of the periostracum (external shell surface) were made in order to evaluate potential effect of the treatments. These observations were performed at two different places of the shell: (i) the ventral extremity part and (ii) a lower part of the shell. Two individuals per experimental condition (65 days of exposure) were studied. For all the samples, the periostracum was of the same colour (based on visual observations). Under SEM, at the ventral extremity in control condition (29.2°C, 430 μatm), the periostracum did not appear altered (Supplementary Fig. S2A, C; Table 4), in the lower part it displayed a ‘mushy’ aspect with numerous micro-borer galleries (Supplementary Fig. S2B, D). These features were observed in all the other experimental conditions (i.e. 29.2°C, 1212 μatm; 30.7°C, 430 μatm; 30.7°C, 1212 μatm; Fig. S2) and for all the samples studied, which may signify that the treatments did not significantly affect the periostracum.

To strengthen the results above ultrastructural observations of shell fractures were done in two zones: zone 1 corresponding to the shell formed *in situ* (i.e. before the experiment) and zone 2 corresponding to the shell formed during the experimental conditions. In the control condition




Figure 3: Effects of experimental conditions on the ultrastructure of the *T. maxima* shell outer layer investigated via SEM. (A, B, D, E, G, H, J, K, M, N: scale bar, 10 μm ; C, F, I, L, O: scale bar, 2 μm). Zone 1 and zone 2 correspond to the shell formed *in situ* (i.e. before the experiment) and in the experimental conditions, respectively. At 29 days, no significant differences between zones 1 and 2 are observed (A–C: 29.2°C, 430 μatm ; D–F: 29.2°C, 1212 μatm). In both zones, lamellae are well cohesive, displaying elongated shape with a smooth surface and slightly rounded outlines. This latter microstructural feature is observed in all shells formed *in situ* (zone 1), for all temperature/ $p\text{CO}_2$ conditions and all time of exposure (A, D, G, J, M). At 41 days, lamellae in zone 2 are less cohesive with or without pronounced granular aspect (H, I: 29.2°C, 430 μatm ; K, L: 30.7°C, 430 μatm ; N, O: 30.7°C, 1212 μatm).

(i.e. 29.2°C, 430 μatm), after 29 days of exposure, the lamellae formed before the experiment (zone 1) and during the experiment (zone 2) were well cohesive and displayed an elongated shape, a smooth surface and sharp or slightly rounded outlines (Fig. 3A–C). At 29 days, no difference was observed between both zones for all temperature/ $p\text{CO}_2$ conditions (example given for 29.2°C, 1212 μatm in Fig. 3D–F; Table 4).

From 41 days and after, and for all temperature/ $p\text{CO}_2$ experimental conditions, the shell formed before the experiment (zone 1) displayed the same microstructural features like the ones described above (example given for 29.2°C, 430 μatm ; 30.7°C, 430 μatm ; and 30.7°C, 1212 μatm at 41 days in Fig. 3G, J, M; Table 4). At 41 days, the lamellae observed in the majority of shells formed during the experiment (zone 2),

Table 3: Results from Tukey post hoc tests following the three-way ANOVA performed on analyzed physiological parameters

		O ₂ production	Photosynthetic yield	Symbiont density	Shell extension rate
Time	29 days	a	a	a	a
	41 days	ab	ab	ab	ab
	53 days	ab	ab	b	b
	65 days	b	b	b	b
Temperature	29.2 °C	a	a	a	-
	30.7 °C	b	b	b	-
pCO ₂	430 μatm	-	a	a	a
	1212 μatm	-	b	b	b
Time × temperature	29 days × 29.2°C	-	ab	-	-
	41 days × 29.2°C	-	a	-	-
	53 days × 29.2°C	-	a	-	-
	65 days × 29.2°C	-	a	-	-
	29 days × 30.7°C	-	cd	-	-
	41 days × 30.7°C	-	abc	-	-
	53 days × 30.7°C	-	bcd	-	-
	65 days × 30.7°C	-	d	-	-

The effects of significant parameters were tested as time of exposure alone and combined to the temperature. The letter annotations correspond to the significances between conditions ($P < 0.05$).

in all temperature/pCO₂ conditions, appeared less cohesive (Fig. 3N, O showing shell formed at 30.7°C, 1212 μatm; Table 4) and some of them additionally displayed a pronounced granular aspect of the surface even in the control condition (Fig. 3H, I and Fig. 3K, L showing shell formed at 29.2°C, 430 μatm and 30.7°C, 430 μatm, respectively; Table 4). These modifications were observed for the majority of shells formed in all temperature/pCO₂ conditions, at 53 and 65 days (Table 4). Two states classifying ultrastructural differences between the shell formed before and during the experiment were defined as follows: (i) ND, no difference observed between zones 1 and 2; and (ii) D, difference observed between zones 1 and 2 meaning that the lamellae were less cohesive in zone 2 compared to zone 1 (see Table 4). In total, 19 samples over 32 displayed differences in lamellae aspect between both zones.

Effect of time of exposure on oxygen balance, biomineralization and symbiont photophysiology

Time of exposure to thermal and acidification stress was shown to have a significant impact on O₂ production ($P=0.030$), photosynthetic yield ($P=0.021$), symbiont density ($P=0.001$) and shell growth rate ($P=0.007$) (Table 2). Post hoc tests showed that the kinetics of thermal or acidification stress varies depending on the physiological

parameter measured: O₂ production and photosynthetic yield were significantly different between 29 and 65 days while symbiont density and shell growth rate were both impacted from 53 days of exposure (Table 3).

Relationship between physiological parameters

To establish if relationships exist between the various physiological parameters measured, a correlation matrix was generated (Table 5). The photosynthetic yield of symbionts was strongly correlated to their density ($r=0.667$). O₂ production was also strongly correlated to symbiont density ($r=0.492$) and photosynthetic yield ($r=0.331$). Concerning the circadian functioning of the holobiont, its nocturnal oxygen need was strongly correlated to the diurnal O₂ production ($r=0.629$). No significant correlation was found between shell extension rate and other physiological parameters.

Discussion

In the present study, we investigated the physiological responses of *T. maxima* (i.e. 4-year-old specimens) to temperature and pCO₂ conditions in the French Polynesian lagoons under current warmer season conditions and those predicted for 2100 by the IPCC 2014 (+1.5°C and +800 μatm of CO₂). Present thermal conditions were stressful enough to induce an acclimatization process characterized

Table 4: Compilation of ultrastructural observations of lamellae of the shell outer layer and the periostracum using SEM

Time of exposure	Temperature (°C)	$p\text{CO}_2$ (μatm)	Sample	Lamellae of outer shell layer	Periostracum
29 days	29.2	430	A	ND	
			B	ND	
	30.7	430	A	ND	
			B	ND	
	29.2	1212	A	ND	
			B	ND	
	30.7	1212	A	ND	
			B	ND	
41 days	29.2	430	A	D	
			B	D	
	30.7	430	A	D	
			B	D	
	29.2	1212	A	D	
			B	D	
	30.7	1212	A	D	
			B	ND	
53 days	29.2	430	A	D	
			B	D	
	30.7	430	A	D	
			B	D	
	29.2	1212	A	ND	
			B	D	
	30.7	1212	A	D	
			B	ND	
65 days	29.2	430	A	D	No alteration
			B	D	No alteration
	30.7	430	A	D	No alteration
			B	ND	No alteration
	29.2	1212	A	D	No alteration
			B	D	No alteration
	30.7	1212	A	ND	No alteration
			B	D	No alteration

'A' and 'B' represent the two samples observed for each treatment and time of exposure; 'ND' means that no significant difference is observed between the shell formed before and during the experiment for the lamellae of the outer shell layer; 'D' means that a notable difference is observed between both parts of the shell (i.e. the lamellae are less cohesive). For the periostracum, 'no alteration' means that the periostracum covering the ventral extremity part of the shell is well preserved.

by a slight regulation of the symbiont density without the collapse of the photosynthesis. However, the predicted thermal and $p\text{CO}_2$ conditions were much more stressful and induced strong disturbances of the processes linked to the phototrophic symbiosis and the biomineralization.

Indeed, the 30.7°C treatments have significantly impacted the holobiont by inducing a strong bleaching response illustrated by the reduction of its O_2 production, symbiont density and photosynthetic yield. High $p\text{CO}_2$ (+800 μatm of CO_2) was shown to alter the symbiont photosynthetic yield and

Table 5: Correlation matrix integrating the different physiological parameters monitored, i.e. O₂ production and consumption, shell extension rate and symbiont photosynthetic yield and density

	Photosynthetic yield (n = 64)	Symbiont density (n = 64)	Shell extension rate (n = 55)	O ₂ production (n = 64)
Symbiont density	0.667*			
Shell extension rate	-0.032	0.168		
O ₂ production	0.331*	0.492*	0.221	
O ₂ consumption	0.163	0.226	0.129	0.629*

Correlation was tested using the Pearson method (threshold of $r = 0.250$, $\alpha = 0.05$) and significant relationships between parameters are indicated by an asterisk (*).

density and affect its biomineralization process by decreasing the shell growth rate.

No synergetic effect between temperature and pCO₂ on giant clam and symbiont physiological parameters

Increase in atmospheric CO₂ impacts both the seawater temperature and pH inducing ocean warming and acidification (Sabine *et al.*, 2004; IPCC, 2014). Therefore, the analysis of the synergetic effect of both stressors on marine bivalve's physiology is crucial. Watson *et al.* (2012) reported in *T. squamosa* juveniles a decrease of giant clam's survival rate induced by exposure to high-temperature or high-pCO₂ conditions. The lowest survival rate (i.e. <20%) was observed under the highest pCO₂ condition (+600 μ atm of CO₂) at +1.5°C and +3°C, after 60 days of exposure.

In our study, no synergetic effect was detected for all measured physiological parameters. Moreover, no mortality was observed in all tested temperature/pCO₂ conditions during the whole experiment which leads us to suggest that a temperature of 30.7°C and a pCO₂ of 1212 μ atm seem to be non-lethal, at least over 65 days of exposure. However, we cannot exclude that some specimens, especially the bleached individuals, may have died after longer exposure to the experimental conditions.

Effect of temperature on holobiont oxygen balance, symbiont photophysiology and shell ultrastructure

Temperature is one of the most important parameters driving bivalve physiology (Aldridge *et al.*, 1995; Bougrier *et al.*, 1995; Watson *et al.*, 2012; Gazeau *et al.*, 2013; Le Moulac *et al.*, 2016a; Latchère *et al.*, 2017). However, since giant clams live in symbiosis with photosynthetic symbionts (Klumpp *et al.*, 1992; Klumpp and Griffiths, 1994; Soo and Todd, 2014), the comparison at the metabolic level with non-symbiotic bivalves can be misleading. Discussion on elevated temperature effect on the holobiont oxygen balance should therefore integrate the effect on both giant clam and its symbionts (Jones and Hoegh-Guldberg, 2001). Even though physiological and molecular responses to thermal stress may

differ, the comparison with scleractinian corals, which are also symbiotic and calcifying organisms, makes sense to better understand the impact of temperature on the holobiont physiology.

We observed that high temperature (+1.5°C) significantly reduced the holobiont O₂ production, the density and the photosynthetic yield of symbionts from 29 days of exposure. Additionally, partial or total bleaching was observed for the majority of individuals exposed to 30.7°C (at both ambient and high pCO₂, see Supplementary Fig. S3). This suggests that thermal stress has a significant impact on the holobiont photophysiology. The reduction of the symbiont photosynthetic yield could reflect photoinhibition. Photoinhibition was previously linked to the degradation of the D1 protein of the reaction center of the PSII altering the photosynthetic apparatus functioning in symbiotic corals and sea anemone subjected to thermal stress (Warner *et al.*, 1999; Smith *et al.*, 2005; Richier *et al.*, 2006; Ferrier-Pagès *et al.*, 2007). Decrease of photosynthetic activity reduced the symbiont O₂ production and consequently, impacted the holobiont O₂ production. Comparable results were obtained from a 24-h experiment showing that an increase of 3°C affected the oxygen production of *T. gigas* and *T. derasa* (Blidberg *et al.*, 2000). Photoinhibition led to the production of ROS by symbionts, which are known to pass through cellular membranes, cause oxidative damages (Lesser *et al.*, 1990; Lesser, 1996; Downs *et al.*, 2002) and impact the PSII (Richter *et al.*, 1990). In corals, Downs *et al.* (2002) showed that high levels of oxidative damage associated with coral bleaching. Indeed, ROS such as hydrogen peroxide (H₂O₂) may play a role in signalling molecule activating the symbiosis dissociation (Fitt *et al.*, 2001; Smith *et al.*, 2005; Perez and Weis, 2006). Expulsion of symbionts by the host may be a strategy to limit oxidative stress and damage to ultimately survive environmental stress (Downs *et al.*, 2002; Perez and Weis, 2006). Two recent works support this hypothesis. Firstly, Zhou *et al.* (2019) related a decrease in symbiont density in response to an excess of oxidative stress in thermally stressed *T. crocea*. Secondly, Dubousquet *et al.* (2016) showed that *T. maxima* overexpressed genes encoding ROS scavengers in response to thermal stress. Interestingly, the cellular and molecular mechanisms enhanced before the bleaching response (e.g. the loose of the symbionts) seem similar between giant clams and corals and conduct to

the same phenomenon of symbiosis dissociation. However, since giant clams and corals display an extra- and an intracellular symbiosis, respectively, the mechanisms of symbiosis dissociation may differ. This constitutes an interesting example of evolutionary convergence of a stress response in two very distant organisms. In this context, it would be interesting to test if the bleaching response in giant clam can be adaptive as it was proposed for corals (Fautin and Buddemeier, 2004) and if variations in giant clam thermotolerance are correlated to the composition of the symbiotic population (Baker, 2003).

Concerning the effect of temperature on the ultrastructure of the crossed-lamellar structure, the lamellae formed in all experimental conditions in the first 29 days were well cohesive with elongated shape and a slight granular aspect, which is consistent with the description made by Faylona *et al.* (2011) in *T. maxima*, Belda *et al.* (1993) in *T. gigas* and Agbaje *et al.* (2017) in *T. derasa*. However, after 29 days, the lamellae observed in the majority of the shells formed in both temperature conditions (whatever the $p\text{CO}_2$ condition) appeared to be less cohesive with pronounced granular aspect. We suggest that these features are due to a lack of organic matrix between the lamellae (i.e. inter-lamellar organic matrix) and embedding the nano-grains forming the lamellae in the crossed-lamellar structure. Indeed, the biomineralization of calcium carbonate structures involves the transport of ions to the mineralization site and the synthesis of macromolecules referred to as 'organic matrix' (Allemand *et al.*, 2004). The organic matrix, consisting of 0.9 wt% and 1.83 wt% in *T. derasa* and *T. gigas* shell, respectively (Agbaje *et al.*, 2017, 2019), is mainly composed of macromolecules commonly found in molluscan shell organic matrix such as lipids, polysaccharides and proteins displaying various levels of glycosylation (Marin *et al.* 2012; Agbaje *et al.*, 2017). The formation of these macromolecules may be energetically costly for molluscs such as marine gastropods (Palmer, 1992). We suggest that in our tested experimental temperature, while oxygen balance is significantly altered, less energy may be available and allocated to the synthesis of macromolecules involved in biomineralization, which may explain the lack of organic matrix embedded into the giant clam shell.

Effect of $p\text{CO}_2$ on giant clam biomineralization and symbiont photophysiology

Results of the present study have indicated that in *T. maxima* high- $p\text{CO}_2$ condition (+800 μatm) induced a significant decrease of the shell extension rate. This observation is in accordance with those obtained for giant clams in three different experiments using various CO_2 enrichment levels and exposure durations. Watson (2015) has exposed juveniles *T. squamosa* to +250 and +550 μatm of CO_2 for 8 weeks under various light levels. Under mid-light level, the author has reported a reduced survival and growth while under high-light levels the growth only was affected. Kurihara and

Shikota (2018) also reported a negative effect of $p\text{CO}_2$ on shell growth (i.e. shell height) in juveniles *T. crocea* when exposed to +600 μatm and +1600 μatm for 4 weeks. Toonen *et al.* (2012) demonstrated that young specimens of *T. maxima* and *T. squamosa* had lower shell growth rates, compared to those reported in the literature under natural $p\text{CO}_2$ /pH conditions, when kept 1 year in +350 to +1000 μatm $p\text{CO}_2$ conditions.

As mentioned above, high- $p\text{CO}_2$ condition (+800 μatm) affected giant clam's shell growth rate and also photosynthetic yield and density of symbionts. Negative effect of $p\text{CO}_2$ on the shell growth rate may be linked to two non-exclusive hypotheses: (i) the physiological adjustment needed to thrive with a seawater at 1212 μatm of CO_2 instead of 430 μatm and/or (ii) an alteration of symbiont photophysiology leading to a potential reduction of the 'light-enhanced calcification' (LEC) phenomenon. Concerning the former, the increase of CO_2 dissolution in the water column leads to the modification of the carbonate chemistry equilibrium and to the increase of H^+ concentration (Cyronak *et al.*, 2016). These strong environmental changes affect the whole organism's homeostasis, but more particularly physiological functions, such as biomineralization, where the control of carbonate concentration and pH are the most essential. To form their mineralized structures, calcifying organisms modify the carbonate composition at the mineralization site to promote CaCO_3 precipitation. These modifications are performed by several enzymatic reactions, including the removal of H^+ to locally increase the pH at the mineralization site for maintaining chemical conditions to enhance CaCO_3 precipitation (Allemand *et al.*, 2011; Taylor *et al.*, 2012). Corals are known to regulate the pH at their mineralization site (Venn *et al.*, 2013; Holcomb *et al.*, 2014) but a decrease of coral calcification was also linked to a decline in pH in the calcifying fluid (Ries, 2011; McCulloch *et al.*, 2012). The maintenance of biomineralization under these non-optimal environmental conditions are therefore energetically costly, which may result in a general decrease of growth (Vidal-Dupiol *et al.*, 2013). In our case, we suggest that giant clam *T. maxima* exposed to high $p\text{CO}_2$ may allocate more energy to maintain a proper pH of the extrapallial fluid for nucleation and deposition of aragonite. Regarding the LEC (Vandermeulen *et al.*, 1972), this phenomenon observed in Symbiodiniaceae-host symbiosis has been extensively described in corals and represents the capacity of symbionts to stimulate the host calcification (Allemand *et al.*, 2004). The symbionts stimulate the host metabolism and calcification by providing energy resources and/or O_2 (Chalker and Taylor, 1975). Moreover, symbionts may promote the aragonite precipitation by providing inorganic carbon, nitrogen and phosphorus and by synthesizing molecules used as precursor for the synthesis of skeletal organic matrix (Pearse and Muscatine, 1971; Cuif *et al.*, 1999; Furla *et al.*, 2000; Muscatine *et al.*, 2005). They also facilitate CaCO_3 precipitation by influencing the dissolved inorganic carbon (DIC) equilibrium by removing the CO_2 via photosynthesis (Goreau, 1959). Such phenomenon

has been also described in giant clams (Ip *et al.*, 2006; Ip *et al.*, 2017). In *T. squamosa*, LEC increased the pH and reduced the ammonia concentration at the interface between the inner mantle and the shell in the extrapallial fluid, where the biomineralization occurs (Ip *et al.*, 2006). Recently, Chew *et al.* (2019) reported a light-enhanced expression of carbonic anhydrase, i.e. CA4-like, in the inner mantle of *T. squamosa* and suggested that this enzyme is involved in giant clam biomineralization by catalyzing the conversion of HCO_3^- to CO_2 . In this context, an altered photophysiology of the symbionts can rationally alter LEC and consequently results in a decrease of the shell growth rate. Finally, one can suggest that under acidification stress, giant clam may reduce some physiological functions such as biomineralization and allocate more energy to essential functions for its survival. In our study, temperature also altered photophysiology and holobiont O_2 production, but did not significantly affect shell growth rate. Therefore, the most plausible hypothesis explaining the negative effect of $p\text{CO}_2$ on shell growth rate may be related to the low aragonite saturation state at high $p\text{CO}_2$.

Concerning the effect of $p\text{CO}_2$ on the shell microstructural integrity, in the temperate bivalve *Mytilus edulis*, an exposure to +150, +350 and +600 μatm of CO_2 for 6 months induced disorientation in the shell of the newly formed calcite crystals of the prismatic layer (Fitzer *et al.*, 2014). In the giant clam *T. maxima*, high $p\text{CO}_2$ had no significant effect on the integrity of the aragonitic lamellae of the crossed-lamellar layer during the first 29 days of exposure. From 41 days of exposure, its potential impact remains unresolved as differences were noticed in the shells formed under future high-temperature, high- $p\text{CO}_2$ and even under today's temperature/ $p\text{CO}_2$ conditions. The fact that differences were reported for shells formed in all experimental conditions from 41 days suggests that they may be due to a long-term exposure to 29.2°C and 30.7°C.

Conclusion

This study enables the evaluation of *T. maxima*'s physiological responses to realistic temperature and $p\text{CO}_2$ predicted at the end of this century. We demonstrated that high temperature mimicking temperature encountered during the warmer months of the year has a significant negative impact on symbiont densities and photosynthetic capacities. This negative impact on the symbiont physiology induces a decrease in the net holobiont O_2 production. Therefore, by influencing symbiont physiology, the temperature may affect the energetic needs of the giant clam host. The high $p\text{CO}_2$ has a negative impact on shell growth rate, symbiont densities and photosynthetic capacities. Shell microstructure is not affected by temperature nor by the $p\text{CO}_2$ in the first 29 days of exposure. However, for all temperature/ $p\text{CO}_2$ conditions, a longer exposure (≥ 41 days) modified the shell ultrastructure. These observations support our hypothesis that 29.2°C is a temperature that already affects giant clam metabolism,

at least over a long-term exposure. However, no synergetic effect was found between temperature and $p\text{CO}_2$ parameters. All these observations suggest that temperature and $p\text{CO}_2$ influence different physiological functions and that giant clam populations may dramatically suffer from the temperature and $p\text{CO}_2$ conditions predicted for the next decades. This is especially true for the temperature since the populations of giant clams (wild and farmed) of the Tuamotu Archipelago are already confronted with temperatures of $\sim 29.2^\circ\text{C}$ every year. Additionally, the threshold of 30.7°C applied in our study and that will be encountered annually in the future corresponds to what is measured currently during abnormally warm years (Fig S1). During the heatwave event in 2016, 80% of the farmed clam populations have bleached resulting in a high level of mortality (personal communication). To complement these results on the effects of temperature on the giant clams, it is now essential to conduct integrative analyses that will take into account the acclimatization and adaptive potential of the whole holobiont. The tools of transcriptomic, genomic and epigenomic in association with ecologically relevant experiments both at the individual and population levels will be particularly relevant to address these questions (Torda *et al.*, 2017). These will allow a better understanding of the fundamental physiological processes of the holobiont and its response to future changes. Results from these studies may help in adapting local policies and management to maintain sustainability of giant clam populations and their exploitation, especially in the Eastern Tuamotu islands where bleaching events have been observed at an increasing and alarming rate.

Funding

This work was supported by the Ifremer Institution (Politique de site program, GECO project) and the Université de la Polynésie française (MAPIKO and CLAMS projects).

Conflicts of interest

The authors declare no financial and personal conflict of interest.

Acknowledgments

We would like to acknowledge the two anonymous reviewers for their valuable comments. We also would like to thank Celine Lafabrie for fruitful discussions and Mickael Mege and Alexia Pihier for their technical assistance. Georges Remoissenet (Direction of Marine Resources of French Polynesia) is thanked for sharing information about bleaching events in Eastern Tuamotu islands. The Direction of Marine Resources and the Ministry of Economic Recovery, Blue Economy and Digital Policy of French Polynesia are also

thanked for the special permit to collect and hold small specimens of *T. maxima* species. This study is set within the framework of the ‘Laboratoires d’Excellence (LabEX)’ TULIP (ANR-10-LABX-41).

References

- Addressi L (2001) Giant clam bleaching in the lagoon of Takapoto atoll (French Polynesia). *Coral Reefs* 19: 220.
- Agbaje OBA, Wirth R, Morales LFG, Shirai K, Kosnik M, Watanabe T, Jacob DE (2017) Architecture of crossed-lamellar bivalve shells: the southern giant clam (*Tridacna derasa*, Röding, 1798). *R Soc Open Sci* 4: 170622. doi:10.1098/rsos.170622.
- Agbaje OBA, Thomas DE, Dominguez JG, McInerney BV, Kosnik MA, Jacob DE (2019) Biomacromolecules in bivalve shells with crossed lamellar architecture. *J Mater Sci* 54: 4952–4969.
- Aldridge DW, Payne BS, Miller AC (1995) Oxygen-consumption, nitrogenous excretion, and filtration-rates of *Dreissena polymorpha* at acclimation temperatures between 20 and 32 degrees. *Can J Fish Aquat Sci* 52: 1761–1767.
- Allemand D, Ferrier-Pagès C, Furla P, Houlbrèque F, Puverel S, Reynaud S, Tambutté E, Tambutté S, Zoccola D (2004) Biomineralisation in reef-building corals: from molecular mechanisms to environmental control. *C R Palevol* 3: 453–467.
- Allemand D, Tambutté E, Zoccola D, Tambutté S (2011) Coral calcification, cells to reefs. In Z Dubinsky, N Stambler, eds, *Coral Reefs: An Ecosystem in Transition*. Springer, Dordrecht, pp. 119–150
- Andréfouët S, Van Wynsberge S, Gaertner-Mazouni N, Menkes C, Gilbert A, Remoissenet G (2013) Climate variability and massive mortalities challenge giant clam conservation and management efforts in French Polynesia atolls. *Biol Conserv* 160: 190–199.
- Andréfouët S, Duthel C, Menkes CE, Bador M, Lengaigne M (2015) Mass mortality events in atoll lagoons: environmental control and increased future vulnerability. *Glob Change Biol* 21: 95–205.
- Andréfouët S, Van Wynsberge S, Kabbadj L, Wabnitz CCC, Menkes C, Tamata T, Pahuatini M, Tetairekie I, Teaka I, Ah Scha T et al. (2017) Adaptive management for the sustainable exploitation of lagoon resources in remote islands: lessons from a massive El Nino-induced giant clam bleaching event in the Tuamotu atolls (French Polynesia). *Environ Conserv* 45: 30–40.
- Armstrong EJ, Dubousquet V, Mills SC, Stillman JH (2020) Elevated temperature, but not acidification, reduces fertilization success in the small giant clam. *Mar Biol* 167: 8. doi:10.1007/s00227-019-3615-0.
- Avignon S, Auzoux-Bordenave S, Martin S, Dubois P, Badou A, Coheleach M, Richard N, Giglio S, Malet L, Servili A et al. (2020) An integrated investigation of the effects of ocean acidification on adult abalone (*Haliotis tuberculata*). *ICES J Mar Sci* 77: 757–772.
- Baker AC (2003) Flexibility and specificity in coral-algal symbiosis: diversity, ecology. *Annu Rev Ecol Evol Syst* 34: 661–689.
- Belda CA, Cuff C, Yellowlees D (1993) Modification of shell formation in the giant clam *Tridacna gigas* at elevated nutrient levels in sea water. *Mar Biol* 117: 251–257.
- Blidberg E, Elfving T, Planhnan P, Tedengren M (2000) Water temperature influences on physiological behaviour in three species of giant clams (Tridacnidae). In *Proceeding 9th International Coral Reef Symposium*, Bali, Indonesia, 23–27 October 2000
- Bougrier S, Geairon P, Deslous-Paoli JM, Bacher C, Jonquières G (1995) Allometric relationships and effects of temperature on clearance and oxygen consumption rates of *Crassostrea gigas* (Thunberg). *Aquaculture* 134: 143–154.
- Broecker WS, Takahashi T, Simpson HJ, Peng T-H (1979) Fate of fossil fuel carbon dioxide and the global carbon budget. *Science* 206: 409–418.
- Buck BH, Rosenthal H, Saint-Paul U (2002) Effect of increased irradiance and thermal stress on the symbiosis of *Symbiodinium microadriaticum* and *Tridacna gigas*. *Aquat Living Resour* 15: 107–117.
- Caldeira K, Wickett ME (2003) Anthropogenic carbon and ocean pH. *Nature* 425: 365.
- Chalker BE, Taylor DL (1975) Light-enhanced calcification, and the role of oxidative phosphorylation in calcification of the coral *Acropora cervicornis*. *Proc R Soc Lond B* 190: 323–331.
- Chew SF, Koh CZY, Hiong KC, Choob CYL, Wong WP, Neo ML, Ip Y (2019) Light-enhanced expression of Carbonic Anhydrase 4-like supports shell formation in the fluted giant clam *Tridacna squamosa*. *Gene* 683: 101–112.
- Cuif J-P, Dauphin Y, Freiwald A, Gautret P, Zibrowius H (1999) Biochemical markers of zooxanthellae symbiosis in soluble matrices of skeleton of 24 Scleractinia species. *Comp Biochem Physiol A Mol Integr Physiol* 123: 269–278.
- Cyronak T, Schulz KG, Jokiel PL (2016) The Omega myth: what really drives lower calcification rates in an acidifying ocean. *ICES J Mar Sci* 73: 558–562.
- Downs CA, Fauth JE, Halas JC, Dustan P, Bemiss J, Woodley CM (2002) Oxidative stress and seasonal coral bleaching. *Free Radic Biol Med* 33: 533–543.
- Dubousquet V, Gros E, Berteaux-Lecellier V, Viguier B, Raharivelomanana P, Bertrand C, Lecellier G (2016) Changes in fatty acid composition in the giant clam *Tridacna maxima* in response to thermal stress. *Biol Open* 5: 1400–1407.
- Elfving T, Plantman P, Tedengren WE (2001) Responses to temperature, heavy metal and sediment stress by the giant clam *Tridacna squamosa*. *Mar Freshw Behav Physiol* 34: 239–248.
- Elfving T, Blidberg E, Tedengren M (2002) Physiological responses to copper in giant clams: a comparison of two methods in revealing effects on photosynthesis in zooxanthellae. *Mar Environ Res* 54: 147–155.
- Fautin D, Buddemeier R (2004) Adaptive bleaching: a general phenomenon. *Hydrobiologia* 530: 459–467.

- Faylona MGP, Lazareth CE, Sémah A-M, Caquineau S, Boucher H, Ronquillo WP (2011) Preliminary study on the preservation of giant clam (*Tridacnidae*) shells from the Balobok Rockshelter archaeological site. *Geoarchaeology* 26: 888–901.
- Ferrier-Pagès C, Richard C, Forcioli D, Allemand D, Pichon M, Shick JM (2007) Effects of temperature and UV radiation increases on the photosynthetic efficiency in four scleractinian coral species. *Biol Bull* 213: 76–87.
- Fitt WK, Brown BE, Warner ME, Dunne RP (2001) Coral bleaching: interpretation of thermal tolerance limits and thermal thresholds in tropical corals. *Coral Reefs* 20: 51–65.
- Fitzer SC, Phoenix VR, Cusack M, Kamenos NA (2014) Ocean acidification impacts mussel control on biomineralisation. *Sci Rep* 4: 6218. doi:10.1038/srep06218.
- Furla P, Galgani I, Durand I, Allemand D (2000) Sources and mechanisms of inorganic carbon transport for coral calcification and photosynthesis. *J Exp Biol* 203: 3445–3457.
- Gannon ME, Pérez-Huerta A, Aharon P, Street SC (2017) A biomineralization study of the Indo-Pacific giant clam *Tridacna gigas*. *Coral Reefs* 36: 503–517.
- Gazeau F, Gattuso J-P, Dawber C, Pronker A, Peene F, Peene J, Heip C, Middelburg J (2010) Effect of ocean acidification on the early life stages of the blue mussel *Mytilus edulis*. *Biogeosciences* 7: 2051–2060.
- Gazeau F, Parker LM, Comeau S, Gattuso J-P, O'Connor W, Martin S, Portner H-O, Ross PM (2013) Impacts of ocean acidification on marine shelled molluscs. *Mar Biol* 160: 2207–2245.
- Goreau TF (1959) The physiology of skeleton formation in corals. I. A method for measuring the rate of calcium deposition by corals under different conditions. *Biol Bull* 116: 59–75.
- Hawkins AJS, Klumpp DW (1995) Nutrition of the giant clam *Tridacna gigas* (L.). II. Relative contributions of filter-feeding and the ammonium-nitrogen acquired and recycled by symbiotic algae towards total nitrogen requirements for tissue growth and metabolism. *J Exp Mar Biol Ecol* 190: 263–290.
- Hoegh-Guldberg O, Mumby PJ, Hooten AJ, Steneck RS, Greenfield P, Gomez E, Harvell CD, Sale PF, Edwards AJ, Caldeira K *et al.* (2007) Coral reefs under rapid climate change and ocean acidification. *Science* 318: 1737–1742.
- Holcomb M, Venn A, Tambutté E, Tambutté S, Allemand D, Trotter J, McCulloch M (2014) Coral calcifying fluid pH dictates response to ocean acidification. *Sci Rep* 4: 5207. doi: 10.1038/srep05207.
- Holt AL, Vahidinia S, Gagnon YL, Morse DE, Sweeney AM (2014) Photosymbiotic giant clams are transformers of solar flux. *J R Soc Interface* 11: 20140678. doi:0.1098/rsif.2014.0678.
- Ip YK, Loong AM, Hiong KC, Wong WP, Chew SF, Reddy K, Sivaloganathan B, Ballantyne JS (2006) Light induces an increase in the pH of and a decrease in the ammonia concentration in the extrapallial fluid of the giant clam *Tridacna squamosa*. *Physiol Biochem Zool* 79: 656–664.
- Ip YK, Hiong KC, Goh EJK, Boo MV, Choo YL, Ching B, Wong WP, Chew SF (2017) The whitish inner mantle of the giant clam, *Tridacna squamosa*, expresses an apical Plasma Membrane Ca^{2+} -ATPase (PMCA) which displays light-dependent gene and protein expressions. *Front Physiol* 8: 781. doi:10.3389/fphys.2017.00781.
- IPCC (2014) Summary for policymakers. In CB Field, VR Barros, DJ Dokken, KJ Mach, MD Mastrandrea, TE Bilir, M Chatterjee, KL Ebi, YO Estrada, RC Genova *et al.*, eds, *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, UK and New York, NY, USA, pp. 1–32
- IPCC (2019) Summary for Policymakers. In H-O Pörtner, DC Roberts, V Masson-Delmotte, P Zhai, M Tignor, E Poloczanska, K Mintenbeck, M Nicolai, A Okem, J Petzold *et al.*, eds, *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate*. In press.
- Ishikura M, Adachi K, Maruyama T (1999) Zooxanthellae release glucose in the tissue of a giant clam, *Tridacna crocea*. *Mar Biol* 133: 665–673.
- Jones RJ, Hoegh-Guldberg O (2001) Diurnal changes in the photochemical efficiency of the symbiotic dinoflagellates (*Dinophyceae*) of corals: photoprotection, photoinactivation and the relationship to coral bleaching. *Plant Cell Environ* 24: 89–99.
- Junchompoo C, Sinrapasan N, Penpain C, Patsorn P (2013) Changing sea-water temperature effects on giant clams bleaching, Mannai Island, Rayong province, Thailand. In *Proceedings of the Design Symposium on Conservation of Ecosystem (The 12th SEASTAR2000 workshop)*, pp. 71–76
- Kleypas JA, Buddemeier RW, Archer D, Gattuso J-P, Langdon C, Opdyke BN (1999) Geochemical consequences of increased atmospheric carbon dioxide on coral reefs. *Science* 284: 118–120.
- Klumpp DW, Griffiths CL (1994) Contributions of phototrophic and heterotrophic nutrition to the metabolic and growth requirements of four species of giant clam (*Tridacnidae*). *Mar Ecol Prog Ser* 115: 103–115.
- Klumpp DW, Lucas JS (1994) Nutritional ecology of the giant clams *Tridacna tevoroa* and *T. derasa* from Tonga – Influence of light on filter-feeding and photosynthesis. *Mar Ecol Prog Ser* 107: 147–156.
- Klumpp DW, Bayne BL, Hawkins AJS (1992) Nutrition of the giant clam *Tridacna gigas*. Contribution of filter feeding and photosynthates to respiration and growth. *J Exp Mar Biol Ecol* 155: 105–122.
- Kroecker KJ, Kordas RL, Crim R, Hendriks IE, Ramajo L, Singh GS, Duarte CM, Gattuso J-P (2013) Impacts of ocean acidification on marine organisms: quantifying sensitivities and interaction with warming. *Glob Change Biol* 19: 1884–1896.
- Kurihara H (2008) Effects of CO₂-driven ocean acidification on the early developmental stages of invertebrates. *Mar Ecol Prog Ser* 373: 275–284.
- Kurihara H, Shikota T (2018) Impact of increased seawater pCO₂ on the host and symbiotic algae of juvenile giant clam *Tridacna crocea*. *Galaxea J Coral Reef Stud* 20: 19–28.

- Kurihara T, Yamada H, Inoue K, Iwai K, Hatta M (2013) Impediment to symbiosis establishment between giant clams and *Symbiodinium* algae due to sterilization of seawater. *PLoS One* 8: 1–8. doi:10.1371/journal.pone.0061156.
- LaJeunesse TC, Parkinson JE, Gabrielson PW, Jeong HJ, Reimer JD, Voolstra CR, Santos SR (2018) Systematic revision of Symbiodiniaceae highlights the antiquity and diversity of coral endosymbionts. *Curr Biol* 28: 2570–2580.e6.
- Latchère O, Le Moullac G, Gaetner-Mazouni N, Fievet J, Magré K, Saulnier D (2017) Influence of preoperative food and temperature conditions on pearl biogenesis in *Pinctada margaritifera*. *Aquaculture* 479: 176–187.
- Leggat W, Buck BH, Grice A, Yellowlees D (2003) The impact of bleaching on the metabolic contribution of dinoflagellate symbionts to their giant clam host. *Plant Cell Environ* 26: 1951–1961.
- Le Moullac G, Soyez C, Latchère O, Vidal-Dupiol J, Fremery J, Saulnier D, Lo Yat A, Belliard C, Mazouni-Gaertner N, Gueguen Y (2016a) *Pinctada margaritifera* responses to temperature and pH: acclimation capabilities and physiological limits. *Estuar Coast Shelf Sci* 182: 261–269.
- Le Moullac G, Soyez C, Vidal-Dupiol J, Belliard C, Fievet J, Sham-Koua M, Lo Yat A, Saulnier D, Gaertner-Mazouni N, Gueguen Y (2016b) Impact of pCO₂ on the energy, reproduction and growth of the shell of the pearl oyster *Pinctada margaritifera*. *Estuar Coast Shelf Sci* 182: 274–282.
- Lesser MP (1996) Elevated temperatures and ultraviolet radiation cause oxidative stress and inhibit photosynthesis in symbiotic dinoflagellates. *Limnol Oceanogr* 41: 271–283.
- Lesser MP, Stochaj WR, Tapley DW, Schick JM (1990) Bleaching in coral reef anthozoans: effects of irradiance, ultraviolet radiation and temperature on the activities of protective enzymes against active oxygen. *Coral Reefs* 8: 225–232.
- Liu W, Huang X, Lin J, He M (2012) Seawater acidification and elevated temperature affect gene expression patterns of the pearl oyster *Pinctada fucata*. *PLoS One* 7: 1–7. doi:10.1371/journal.pone.0033679.
- Marin F, Le Roy N, Marie B (2012) Formation and mineralization of mollusk shell. *Front Biosci* 4: 1099–1125.
- McClintock JB, Angus RA, McDonald MR, Amsler CD, Catledge SA, Vohra YK (2009) Rapid dissolution of shells of weakly calcified Antarctic benthic macroorganisms indicates high vulnerability to ocean acidification. *Antarct Sci* 21: 449–456.
- McCulloch M, Falter J, Trotter J, Montagna P (2012) Coral resilience to ocean acidification and global warming through pH up-regulation. *Nat Clim Change* 2: 623–627.
- Melzner F, Stange P, Trübenbach K, Thomsen J, Casties I, Panknin U, Gorb S, Gutowska M (2011) Food supply and seawater pCO₂ impact calcification and internal shell dissolution in the blue mussel *Mytilus edulis*. *PLoS One* 6: 1–9. doi:10.1371/journal.pone.0024223.
- Meng Y, Guo Z, Fitzer SC, Upadhyay A, Chan VBS, Li C, Cusack M, Yao H, Yeung KWK, Thiyagarajan V (2018) Ocean acidification reduces hardness and stiffness of the Portuguese oyster shell with impaired microstructure: a hierarchical analysis. *Biogeosciences* 15: 6833–6846.
- Morse JW, Arvidson RD, Lüttge A (2007) Calcium carbonate formation and dissolution. *Chem Rev* 107: 342–381.
- Muscatine L, Goiran C, Land L, Jaubert J, Cuif J-P, Allemand D (2005) Stable isotopes (delta C-13 and delta N-15) of organic matrix from coral skeleton. *Proc Natl Acad Sci U S A* 102: 1525–1530.
- Norton JH, Shepherd MA, Long HM, Fitt WK (1992) The zooxanthellal tubular system in the giant clam. *Biol Bull* 183: 503–506.
- Orr JC, Fabry VJ, Aumont O, Bopp L, Doney SC, Feely RA, Gnanadesikan A, Gruber N, Ishida A, Joos F *et al.* (2005) Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. *Nature* 437: 681–686.
- Palmer AR (1992) Calcification in marine molluscs: how costly is it? *Proc Natl Acad Sci U S A* 89: 1379–1382.
- Parker LM, Ross PM, O'Connor WA, Pörtner HO, Scanes E, Wright JM (2013) Predicting the response of molluscs to the impact of ocean acidification. *Biology* 2: 651–692.
- Pätzold J, Heinrichs JP, Wolschendorf K, Wefer G (1991) Correlation of stable oxygen isotope temperature record with light attenuation profiles in reef-dwelling *Tridacna* shells. *Coral Reefs* 10: 65–69.
- Pearse VB, Muscatine L (1971) Role of symbiotic algae (zooxanthellae) in coral calcification. *Biol Bull* 141: 350–363.
- Perez S, Weis V (2006) Nitric oxide and cnidarian bleaching: an eviction notice mediates breakdown of a symbiosis. *J Exp Biol* 209: 2804–2810.
- Richier S, Sabourault C, Courtiade J, Zucchini N, Allemand D, Furla P (2006) Oxidative stress and apoptotic events during thermal stress in the symbiotic sea anemone. *FEBS J* 273: 4186–4198.
- Richter C, Rühle W, Wild A (1990) Studies on the mechanisms of photosystem II photoinhibition II. The involvement of toxic oxygen species. *Photosynth Res* 24: 237–243.
- Ries JB (2011) A physiochemical framework for interpreting the biological calcification response to CO₂-induced ocean acidification. *Geochim Cosmochim Acta* 75: 4053–4064.
- Ries JB, Cohen AL, McCorkle DC (2009) Marine calcifiers exhibit mixed responses to CO₂-induced ocean acidification. *Geology* 37: 1131–1134.
- Rodolfo-Metalpa R, Houlbreque F, Tambutte E, Boisson F, Baggini C, Patti FP, Jeffree R, Fine M, Foggo A, Gattuso J-P *et al.* (2011) Coral and mollusc resistance to ocean acidification adversely affected by warming. *Nat Clim Change* 1: 308–312.
- Sabine CL, Feely RA, Gruber N, Key RM, Lee K, Bullister JL, Wanninkhof R, Wong CS, Wallace DWR, Tilbrook B *et al.* (2004) The oceanic sink for anthropogenic CO₂. *Science* 305: 367–371.
- Schwartzmann C, Durrieu M, Sow M, Ciret P, Lazareth CE, Massabuau J-C (2011) In situ giant clam growth rate behavior in relation to temperature: a one-year couple study of high-frequency noninvasive valvometry and sclerochronology. *Limnol Oceanogr* 56: 1940–1951.

- Smith DJ, Suggett DJ, Baker NR (2005) Is photoinhibition of zooxanthellae photosynthesis the primary cause of thermal bleaching in corals? *Glob Change Biol* 11: 1–11.
- Soo P, Todd PA (2014) The behaviour of giant clams. *Mar Biol* 161: 2699–2717.
- Talmage SC, Gobler CJ (2011) Effects of elevated temperature and carbon dioxide on the growth and survival of larvae and juveniles of three species of northwest Atlantic bivalves. *PLoS One* 6: 1–12. doi:10.1371/journal.pone.0026941.
- Taylor AR, Brownlee C, Wheeler GL (2012) Proton channels in algae: reasons to be excited. *Trends Plant Sci* 17: 675–684.
- Toonen RJ, Nakayama T, Ogawa T, Rossiter A, Delbeek JC (2012) Growth of cultured giant clams (*Tridacna spp.*) in low pH, high nutrient seawater: species-specific effects of substrate and supplemental feeding under acidification. *J Mar Biol Assoc UK* 92: 731–740.
- Torda G, Donelson JM, Aranda M, Barshis DJ, Bay L, Berumen ML, Bourne DG, Cantin N, Foret S, Matz M *et al.* (2017) Rapid adaptive responses to climate change in corals. *Nat Clim Chang* 7: 627.
- Vandermeulen JH, Davis ND, Muscatine L (1972) The effect of inhibitors of photosynthesis on zooxanthellae in corals and other marine invertebrates. *Mar Biol* 16: 185–191.
- van Heuven S, Pierrot D, Lewis E, Wallace DWR (2009) MATLAB Program developed for CO₂ system calculations, ORNL/CDIAC-105b, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, US Department of Energy, Oak Ridge, Tennessee.
- Van Wynsberge S, Andréfouët S, Gaertner-Mazouni N, Wabnitz CCC, Gilbert A, Remoissenet G, Payri C, Fauvelot C (2016) Drivers of density for the exploited giant clam *Tridacna maxima*: a meta-analysis. *Fish Fish* 17: 567–584.
- Venn A, Tambutté E, Holcomb M, Laurent J, Allemand D, Tambutté S (2013) Impact of seawater acidification on pH at the tissue-skeleton interface and calcification in reef corals. *Proc Natl Acad Sci U S A* 110: 1634–1639.
- Vidal-Dupiol J, Zoccola D, Tambutté E, Grunau C, Cosseau C, Smith KM, Freitag M, Dheilily NM, Allemand D, Tambutté S (2013) Genes related to ion-transport and energy production are upregulated in response to CO₂-driven pH decrease in corals: new insights from transcriptome analysis. *PLoS One* 8: 1–11. doi: 10.1371/journal.pone.0058652.
- Warner ME, Fitt WK, Schmidt GW (1999) Damage to photosystem II in symbiotic dinoflagellates: a determinant of coral bleaching. *Proc Natl Acad Sci U S A* 96: 8007–8012.
- Watson S-A (2015) Giant clams and rising CO₂: Light may ameliorate effects of ocean acidification on a solar-powered animal. *PLoS One* 10: 1–18. doi: 10.1371/journal.pone.0128405.
- Watson S-A, Southgate PC, Miller GM, Moorhead JA, Knauer J (2012) Ocean acidification and warming reduce juvenile survival of the fluted giant clam *Tridacna squamosa*. *Molluscan Res* 32: 177–180.
- Welladsen HM, Southgate PC, Heimann K (2010) The effects of exposure to near-future levels of ocean acidification on shell characteristics of *Pinctada fucata* (Bivalvia: Pteriidae). *Molluscan Res* 30: 125–130.
- Wessel N, Martin S, Badou A, Dubois P, Huchette S, Julia V, Nunes F, Harney E, Paillard C, Auzoux-Bordenave S (2018) Effect of CO₂-induced ocean acidification on the early development and shell mineralization of the European abalone (*Haliotis tuberculata*). *J Exp Mar Biol Ecol* 508: 52–63.
- Yau AJY, Fan TY (2012) Size-dependent photosynthetic performance in the giant clam *Tridacna maxima*, a mixotrophic marine bivalve. *Mar Biol* 159: 65–75.
- Zeebe RE, Zachos JC, Caldeira K, Tyrrell T (2008) Oceans-Carbon emissions and acidification. *Science* 321: 51–52.
- Zhou Z, Liu Z, Wang L, Luo J, Li H (2019) Oxidative stress, apoptosis activation and symbiosis disruption in giant clam *Tridacna crocea* under high temperature. *Fish Shellfish Immunol* 84: 451–457.