



HAL
open science

The development and maintenance of resident macrophages

Elisa Gomez Perdiguero, Frederic Geissmann

► **To cite this version:**

Elisa Gomez Perdiguero, Frederic Geissmann. The development and maintenance of resident macrophages. *Nature Immunology*, 2016, 17 (1), pp.2-8. 10.1038/ni.3341 . hal-03265984

HAL Id: hal-03265984

<https://hal.science/hal-03265984v1>

Submitted on 12 Jul 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Published in final edited form as:

Nat Immunol. 2016 January ; 17(1): 2–8. doi:10.1038/ni.3341.

Development and maintenance of resident macrophages

Elisa Gomez Perdiguero¹ and Frederic Geissmann²

¹ Macrophages and endothelial cells, Department of Developmental and Stem Cell Biology, CNRS URA 2578, Institut Pasteur, 25 rue du Docteur Roux, Paris 75015, France

² Immunology program, Memorial Sloan Kettering Cancer Center, 417 E 68th St, New York NY 10065, USA

Abstract

The molecular and cellular mechanisms that underlie the many roles of macrophages in health and disease states *in vivo* remain poorly understood. The purpose of this Review is to present and discuss current knowledge on the developmental biology of macrophages, as it underlies the concept of a layered myeloid system composed of ‘resident’ macrophages that mostly originate from yolk sac progenitors and of ‘passenger’ or ‘transitory’ myeloid cells that originate and renew from bone marrow hematopoietic stem cells, and to provide a framework to investigate the functions of macrophages *in vivo*.

The current ‘mononuclear phagocyte system’ model¹ has guided many investigators studying the functions of macrophages over the past 30 years. It postulates that tissue macrophages are maintained from a constant supply of bone marrow derived circulating blood monocytes that extravasate into tissues². This model was initially based on the observation that blood leucocytes recruited in the inflamed peritoneum differentiate into macrophages. Subsequent transplantation experiments in irradiated recipients demonstrated that bone marrow hematopoietic stem cells (HSC) and HSC-derived myeloid progenitors can indeed give rise to tissue macrophages or dendritic cells in mice and humans^{3, 4, 5, 6, 7}. Therefore, investigations designed to study the molecular and cellular basis for macrophage phenotypic diversity and functions have largely focused on the activation, or polarization, of monocyte-derived macrophage by external cues, such as cytokines and microbe-associated signals^{8, 9, 10}. Nevertheless, it remains currently unclear whether polarization stages of monocytes are stable or represent transient states of activation, whether they account for the phenotypic and functional heterogeneity of macrophages *in vivo* in health or disease states, and how many such polarized states exist^{10, 11}.

A new model of macrophage development is emerging based on a number of independent observations that two distinct populations of macrophages, which can be distinguished by their progenitors, developmental history, turnover and mechanisms of maintenance, coexist within the tissues of an adult mouse. The macrophage system of a mouse can now be described as a ‘layered’ system where ‘resident’ macrophages that develop in embryos independently of HSC¹² persist in adult tissues independently of adult

HSC^{12, 13, 14, 15, 16, 17, 18}, and coexist with ‘passenger’ leucocytes such as monocytes and classical dendritic cells that originate and renew from bone marrow HSCs and myeloid progenitors^{4, 6, 19}. Resident macrophages include spleen red pulp macrophages, lung alveolar macrophages, epidermal Langerhans cells, brain microglia, liver Kupffer cells, large peritoneal macrophages, and F4/80^{bright} pancreatic, kidney and cardiac macrophages. Many ‘resident’ macrophages are long-lived in mice and can proliferate within their tissue of residence, a mechanism involved in their maintenance in adults^{20, 21, 22, 23, 24, 25, 26, 27}. Nevertheless, bone marrow-derived progenitors also contribute to resident subsets in the lamina propria, spleen, brain, skin, heart, liver and kidney in a proportion that varies with the tissue considered, the age of the mice, and pathological processes^{12, 13, 14, 15, 16, 17, 18, 28, 29}.

Therefore, the purpose of this Review is to present and discuss current knowledge on the developmental biology of resident macrophages, as it underlies the concept of a layered myeloid system composed of resident macrophages distinct from passenger macrophages and myeloid cells, and provides a new framework and experimental tools to characterise the functions of macrophages *in vivo*, in health and disease settings.

The original concept of ‘primitive macrophages’

During ontogeny, macrophages were found in the mouse yolk sac and the embryo before the initiation of HSC-derived hematopoiesis and before monocytes could be detected in the fetal blood. These ‘primitive macrophages’ were described in a series of seminal works by M. Naito and K. Takahashi in the 1980’s^{20, 30, 31} that reported the presence of large cells with macrophage morphology in blood islands of the yolk sac at E9 and in the fetal liver at E10. They were described as immature because of the lack of detectable phagocytic activity and a weak immunoreactivity for F4/80, as detected by electron microscopy^{20, 30}. In addition to the ‘immature macrophages’, these early studies also reported the existence of ‘fetal macrophages’ from E10 to E17, as slightly smaller cells with pseudopodia, phagocytic activity and frequent mitotic figures³¹. At E17, the ‘fetal macrophages’ were reported as more differentiated, as they acquired ultrastructural features similar to those observed in adult tissue macrophages^{30, 32}. These early studies noted that the remaining difference between fetal and adult tissue macrophages was the presence of numerous mitotic figures in fetal macrophages throughout development. It was also noted that poorly differentiated myeloid cells are detected in the blood vessels of the yolk sac at E10²⁰, where immature stages of the monocyte and neutrophil differentiation series can be detected from E12 onwards, and bona-fide neutrophils and monocytes, based on ultrastructural features and PO-cytochemistry, are present in the fetal liver at E15 and E16 respectively³⁰. They therefore proposed that fetal macrophages develop without a monocytic intermediate, and in parallel with the monocyte and granulocyte lineage.

In the recent literature, this early work was sometimes reported as evidence for the early colonization of the fetus by ‘yolk-sac macrophages’ before their replacement by monocyte-derived macrophages. However, this body of work described the identification of proliferating non-phagocytic macrophage precursors (described as ‘immature macrophages’) that originate in the yolk sac, colonize the embryo and finish their differentiation *in situ* within the fetal tissues^{20, 30, 31}. Their description did not correspond to a particular wave of

hematopoietic progenitors ('primitive' or 'definitive'), as discovered subsequently. However, the description of erythro-myeloid progenitors (EMP)^{33, 34, 35, 36} (see below) and recent fate mapping data are in accordance with the authors original interpretation of their morphological data. Indeed, genetic pulse labeling of *Csfr1*⁺ Kit⁺ CD45^{lo} AA4.1⁺ EMP present in the yolk sac at E8.5 is followed by the detection of labeled CD11b⁺ cells with a negative or faint immunoreactivity for F4/80 at E9.5 in both the yolk sac and embryo and then by the detection of F4/80^{bright} cells starting from E10.5¹⁸.

More recent works have identified three successive but overlapping waves of hematopoietic progenitors during development, all of which have the potential to give rise to fetal macrophage³⁴. They can be distinguished by the anatomical site (extra-embryonic yolk sac versus intra-embryonic) where they are generated, expand and differentiate, their differentiation potential and their transcription factor dependency (**Table 1** and **Fig. 1**). Here we will focus on the myeloid lineages. Erythroid lineages have been reviewed in³⁷.

Primitive hematopoiesis is a source of macrophages in embryos

Primitive hematopoiesis is the source of the first hematopoietic cells detected in the developing embryo. It is characterized by the emergence of progenitors with limited potential for erythrocytes and macrophages and which can be found in the extra-embryonic yolk-sac blood islands in mice³⁶. These progenitors are thought to arise directly from the posterior plate mesoderm, and clonal analysis of the blood islands reveals that the endothelial cells lining the blood islands and the hematopoietic cells they contain do not share a common origin³⁸. This distinguishes them from the subsequent 'definitive' hematopoietic waves, which are thought to arise from a hemogenic endothelium. Accordingly, primitive progenitors and their daughter cells are not affected when transcription factors required for the endothelial-hematopoietic transition, such as RUNX1, are absent (Table 1).

In zebrafish embryos, the 'primitive' wave can give rise to erythrocytes and macrophages, as well as neutrophils, in the absence of *Runx1*^{39, 40, 41}. In the mouse embryo, the *in vivo* characterization of the macrophage progeny of the primitive wave is still hampered by experimental constraints. However, (rare) progenitors with restricted macrophage potential *in vitro* can be found in the yolk sac at the neural plate stage between E7.5 and E8^{36, 34}.

Erythro-Myeloid Progenitors constitute the first wave of definitive hematopoiesis and give rise to most resident macrophages

The first 'definitive' progenitors emerge in the yolk sac of mouse embryos at E8.25^{34, 36}. Termed erythro-myeloid progenitors (EMP), these progenitors are phenotypically defined as Kit⁺, AA4.1⁺ (CD93), CD41⁺, VE-cadherin⁺ (VE-Cad), CD16/32⁺ (FCγII/III receptors) and CD45^{low}^{33, 35} (Table 1). Examination of their erythroid progeny led to their classification as 'definitive' progenitors⁴². However, they can be distinguished from HSCs by the lack of lymphoid potential, both *in vitro* and *in vivo*, the lack of long-term repopulating potential³⁵ and the lack of cell-surface expression of Sca-1³⁵. The EMP-derived hematopoiesis is sufficient to support survival of embryos lacking HSCs until birth⁴³. EMP emerge from the

precursors can be distinguished genetically, and overall the layering of the hematopoietic system and in particular of macrophage development appears to stand on a robust genetic basis. Primitive hematopoiesis occurs in the absence of *Runx1* and *Myb*^{63, 64}, while EMP are *Runx1*-dependent for their emergence⁶⁴ but *Myb*-independent for their myeloid differentiation^{12, 18, 49} and Notch1-independent⁶¹. In contrast, fetal and adult HSC require *Runx1*, *Myb* and *Notch1* (**Table 1**)^{39, 40, 58, 59, 61}. Based on these data, several reports have investigated the lineage of origin of fetal (primitive) and adult macrophages and the mechanisms that may account for their persistence in adults.

Resident macrophages originate in majority from yolk sac EMP

The development of fetal F4/80^{bright} macrophages is unaltered in *Myb*-deficient mice and adult F4/80^{bright} macrophages are largely maintained in adult mice in the absence of *Myb*, and independently of bone marrow progenitors¹². In addition, adult tissue resident F4/80^{bright} macrophages were traced back to *Myb*-independent *Csf1r*-expressing (*Csf1r*⁺) precursors present in the early mouse embryo at E8.5 using Cre-mediated hydroxy-tamoxifen (OH-TAM)-induced pulse labeling^{65, 66} in *Csf1r-MeriCreMer* mice¹² (see below).

These data suggested that F4/80^{bright} macrophages do not originate from HSC, but from yolk sac progenitors such as either primitive progenitors or EMP⁴⁸. In accordance, brain microglia develop from yolk sac progenitors¹⁴, later functionally identified as EMP⁴⁸. Following pulse-labeling of *Csf1r*-expressing cells with OH-TAM at E8.5, the labeling efficiency of F4/80^{bright} macrophages was 30% at day 10.5, and remained high (~20%) in microglia, but progressively decreased from E12.5 onwards in other organs to reach a plateau at birth, maintained even in one year-old animals (5% for the liver Kupffer cells, 1% for epidermal Langerhans cells (^{12, 18, 67} and unpublished data from E. Gomez Perdiguero and F. Geissmann). This may be explained by the observation that pulse labeling with OH-TAM at E8.5 only labels the very first EMP to emerge in the yolk sac, while EMP continue to emerge in the yolk sac until at least E10.5¹⁸, and therefore the labeling percentage of the macrophages in embryonic tissues may reflect the time at which their colonization by fetal macrophages is completed. However, it is also possible that this “dilution” reflects a contribution of another progenitor, such as fetal HSC or HSC-derived monocytes to several fetal macrophage populations^{17, 67, 68, 69}. A contribution of fetal monocytes is unlikely, as they also derive from EMP until E16.5^{18, 35}. However, a contribution of fetal HSC is still debated^{17, 68, 69, 70}.

The main argument against a significant contribution of fetal HSCs come from experiments using a *Tie2-MeriCreMer* fate mapping model, where the expression of Cre recombinase is inducible in *Tie2*-expressing cells^{18, 71}. *Tie2* (also known as *Tek*) is expressed in the hemogenic endothelium, endothelial cells and hematopoietic progenitors in the yolk sac⁷², aorta-gonado-mesonephros region, fetal liver HSC and adult HSC⁵². Pulse labeling with tamoxifen at E7.5, E8.5, E9.5 and E10.5 resulted in labeling of adult HSC, lymphocytes, monocytes and neutrophils with the same efficiency, as expected. In contrast, fetal macrophages and adult resident F4/80^{bright} macrophages in all organs were labeled with a decreasing efficiency (high at E7.5 to very low at E10.5) indicating that fetal and adult

F4/80^{bright} macrophages originate in majority from a progenitor that does not express *Tie2* after E10.5, with a minor contribution of progenitors (such as HSC) expressing *Tie2* after E10.5.

Using fate mapping models where the expression of Cre recombinase is either inducible or constitutive in *Csf1r*-expressing cells (*Csf1r-MerCremer* and *Csf1r-iCre* mice), the macrophage progenitor itself was identified as a CD45^{lo}Kit⁺AA4.1⁺ EMP that appears in the yolk sac from E8.5 ¹⁸. These EMPs migrate in the fetal liver, where they expand and persist as Kit⁺ progenitors until at least E16.5 ¹⁸. These EMPs give rise to erythrocytes, monocytes, granulocytes and mast cells until late fetal development ^{18,35} in accordance with the earlier observation that HSC-independent hematopoiesis is not only necessary, but also sufficient to support survival until the birth of embryos lacking HSCs ⁴³. They also give rise to F4/80^{bright} fetal macrophages in all embryonic tissues from E10.5 onwards ¹⁸. An EMP origin of the majority of resident macrophages was also favoured in an independent report ⁶⁷.

Altogether these data support a major contribution from yolk-sac EMPs to fetal and adult resident macrophages (Fig.1 Fig.2). EMP-derived erythrocytes, monocytes, granulocytes and mast cells are ultimately replaced by HSC derived myeloid cells in the late embryo ¹⁸, while EMP-derived macrophages persist all through fetal and post-natal development as tissue-resident macrophages ^{12, 18} (**Fig. 2**). It was also speculated that a first wave of *Myb*-independent EMP give rise to microglia and a second wave of *Myb*-dependent EMPs give rise to other macrophages respectively ⁶⁷. However, there is no available data to support the existence of two genetically distinct populations of EMP, and this hypothesis is not in accordance with previous reports that *Myb* is expressed in early yolk sac-EMP ^{36, 48} and with the persistence of fetal resident macrophages in all *Myb*-deficient embryonic tissues ¹².

The contribution of “primitive” hematopoiesis to microglia

Primitive hematopoiesis could also contribute to the “primitive” and fetal macrophages. Early (E7.5) pulse-labelling of progenitors using several different reporters such as *Runx1*, *Csf1r*, *Tie2* and *Kit* ^{12, 14, 18, 70}, all result in a more efficient labelling of adult microglia in comparison with other adult tissue resident macrophages. As mentioned above, this may relate either to an earlier termination of brain colonization by macrophages in comparison to other organs, or the existence of several subsets of EMPs. However, it might also reflect the contribution of primitive hematopoiesis to microglia. It will be interesting to determine whether primitive progenitors are labelled in these models, and to investigate whether macrophages produced by primitive hematopoiesis can persist in adult mice. *Myb*-independency does not allow to genetically distinguish primitive from EMP-derived hematopoiesis and it is not yet known whether *Runx1* is required for the differentiation of macrophages in mice. Altogether, the lack of genetic dissection and the low temporal resolution of classical fate mapping approaches in the mouse has not yet allowed independent tracking of the respective progeny of the two progenitor populations. A better understanding of the molecular control of primitive progenitors and EMP commitment and differentiation is required to develop better tools in order to characterize the possible

contribution of “primitive” hematopoiesis to the development of fetal macrophages, and possibly adult macrophages in mice.

Contribution of HSCs to populations of resident macrophages in adult mice varies depending on organs and time

The analysis of HSC contribution to adult resident macrophages in the absence of bone marrow irradiation was performed by several groups using *Flt3-Cre* mice^{12, 17, 18, 73} based on the observation that *Flt3* (encoded by *Flik2*) is expressed by HSC-derived progenitors^{71, 74}, but not by fetal macrophages¹². HSC contribution was also studied in several independent models of bone marrow transplantation without irradiation such as *Mx-1-Cre Myb^{fl/fl}* and *Kit^{W/W^v}* *Rag2^{-/-}* *γc^{-/-}*^{12, 18} and in parabiotic mice^{15, 75}. Overall, these analyses revealed that HSC contribution to resident macrophages was different between organs and frequently increased with age. HSC contribution to adult resident macrophages was either minor (<5%) in brain, liver, epidermis^{12, 14, 48}, small but increasing with age in lung, heart and spleen^{18, 75}, and sometimes predominant after weaning in the case of the gut lamina propria^{28, 76} (**Fig.1** and **Fig. 2**). It is not yet known whether the precursor(s) of the HSC-derived resident macrophages are monocytes or earlier myeloid progenitors, except in the case of the gut, where Ly6C⁺ monocytes are proposed to be the immediate precursor of gut lamina propria macrophages^{28, 76}.

Of note, a recent report investigated the origin of resident macrophages using a fate mapping model where the expression of the Cre recombinase was inducible in *Kit*-expressing cells (*Kit-MeriCreMer* mice)⁷⁰. Importantly, this study represents an independent confirmation that most adult resident macrophages originate from embryonic and fetal progenitors and persist in adults independently of adult HSCs, as *Kit*⁺ progenitors do not seem to contribute to resident tissue macrophages after E16.5⁷⁰. The authors also conclude that most resident macrophages, except microglia, are descendants of fetal HSCs and not of EMP, based on the observation that labeling of *Kit*⁺*Sca1*⁺ EMP at day E7.5 results in the labeling of microglia only, while labeling of *Kit*⁺ progenitors at E8.5 or E9.5 results in contrast in the labeling of all adult resident macrophages and HSC-derived cells. However, EMP do not express *Sca1* (**Table 1** and³⁵), while EMP and fetal HSC both express *Kit* and co-exist in the fetal liver until E16.5^{18, 35}. Thus, *Kit* expression and *Kit*-based fate mapping may not allow to discriminate between EMP and fetal HSC.

Recent work in chicken suggests that in contrast to mice, yolk sac-derived macrophages are not retained after hatching and, surprisingly, that tissue macrophage in adult birds originate from a macrophage-restricted, self-renewing progenitor cell in bone marrow that does not give rise to any other lineage⁷³. These findings, if confirmed, would indicate that myelopoiesis in birds is markedly different from its counterparts in mice, zebrafish and human, because a macrophage-restricted, self-renewing progenitor cell in the bone marrow or its equivalent was not identified in these species.

Maintenance of macrophages by local proliferation or recruitment

As indicated above, a proportion of alveolar macrophages, kidney macrophages and heart macrophages appear to be replaced during the normal ageing process^{12, 18, 75}. A partial replacement of resident macrophages is also observed following gamma irradiation, bone marrow transplantation or adoptive transfer experiments^{3, 4, 77} and in macrophage depletion studies, such as intravenous injections of clodronate liposomes. Liver Kupffer cells are also reported to be replaced by bone marrow derived progenitors following, for example, the death of Kupffer cells in severe experimental *Listeria* infection²⁹. However, resident macrophages such as Langerhans cells, Kupffer cells, microglia, alveolar macrophages, peritoneal macrophages and splenic macrophages all have the potential to proliferate^{15, 22, 23, 27, 78, 79, 80, 81, 82}. In some cases, resident macrophages can immediately replenish themselves following a severe depletion^{15, 23, 82}. A growing body of evidence now clearly shows that fully differentiated resident macrophages can re-enter cell cycle and maintain their numbers by local proliferation^{23, 24, 25, 26, 27}. Macrophage proliferation increases dramatically during inflammatory conditions, such as skin inflammation, where up to 30% of Langerhans cells express the proliferation marker Ki67²². During an acute inflammatory response, tissue-resident peritoneal macrophages that survive undergo a transient and intense proliferative burst *in situ* to repopulate the tissue²³. In response to helminthes infection, resident macrophages in the pleura and peritoneal cavities expand by local proliferation⁸¹. In glucan-induced granuloma formation in monocytopenic mice, Kupffer cells proliferate actively and contribute to granuloma formation⁷⁸. After conditional ablation of microglia in adult mice, the microglial compartment is reconstituted within one week of depletion by CNS-resident cells, independent of bone-marrow-derived precursors⁸². However, it is yet unclear whether the tissue-resident macrophage populations contain a local 'progenitor compartment', or whether all macrophages have the capacity to undergo mitosis. The mechanisms that may underlie this ability of resident macrophages to self-maintain and to proliferate are still unclear. An important role of the Maf family of transcription factors is likely, as combined deficiency for the transcription factors MafB and c-Maf enables the KLF4- and c-Myc-dependent proliferation of differentiated macrophages in culture in response to Csf1⁸³. In the case of epidermal Langerhans cell, proliferation *in vivo* was shown to be driven by an (unknown) keratinocyte-derived signal²². In the case of nematode infection, macrophage proliferation was impaired in the absence of IL-4^{81, 84}. The cellular and molecular mechanisms that control macrophage maintenance and expansion in adult mice, and the consequences of ageing on these processes, are an area of active investigation, as their elucidation will shed light on the pathophysiology of inflammation.

Conclusions and perspectives on macrophage functions

A revised model for macrophage development and functions is therefore emerging, which distinguishes the resident macrophages that develop during embryogenesis from the passenger myeloid cells that originate and renew from bone marrow HSC. This distinction between passenger and resident macrophages is in line with earlier kinetic studies of the splenic-myeloid compartment in adult mice, which proposed a dual origin for myeloid cells, with half of myeloid cells renewing from blood and the other half being produced within the

spleen⁸⁵. It also explains the observations that Langerhans cells do not exchange between parabiotic mice¹³ and that fetal microglia persists in adult mice¹⁴.

This distinction is relevant to disease pathogenesis, as it is quite clear that the functions of HSC-derived monocytes can be very different from those of tissue-resident macrophages within the same inflammatory environment. The two cell types have distinct expansion mechanisms and distinct functions in the brain during experimental auto-immune encephalitis (EAE)⁸⁶. Infiltrating monocytes are recruited via extravasation from blood vessels and produce inflammatory mediators important for disease progression, but do not persist after the resolution of inflammation, while, in contrast, activated resident microglia proliferate locally, persist and return to quiescence following remission⁸⁶. Following helminth parasite infection, pleural resident macrophages adopt anti-inflammatory phenotypes, while passenger macrophages exhibit pro-inflammatory responses⁸¹. However, in a model of chronic neurodegeneration, proliferating resident microglial cells contribute to neuronal damage during the development of the disease⁸⁷. Thus, it might be inaccurate to describe resident macrophages as always being 'anti-inflammatory' as opposed to passenger macrophages being 'pro-inflammatory'. The respective roles of resident and passenger macrophages remain to be studied in detail, and most experimental models available, at present, do not easily elucidate the roles of resident and recruited macrophages in adult mouse tissues. Future work should help characterize the roles of (EMP-derived) resident macrophages, separate them from the function of HSC-derived passenger macrophages and characterize the underlying molecular mechanisms involved.

It is also important to recognize that the distinct subsets of resident macrophages associated with different tissues, for example microglia, Langerhans cells, Kupffer cells, peritoneal macrophages, red pulp macrophages or alveolar macrophages have different phenotypes, transcriptional profiles and chromatin landscape^{88, 89}, require specific growth factors and rely on distinct transcription factors for their differentiation or maintenance. For example, colony stimulating factor 1 (CSF1/M-CSF) is essential for Kupffer cells⁹⁰, TGF β 1 and IL-34 for microglia and Langerhans cells^{88, 91, 92, 93}, colony stimulating factor 2 (CSF2/GM-CSF) for the differentiation or maintenance of alveolar macrophages in mice and humans^{94, 95, 96, 97, 98, 99}, and retinoic acid and GATA6 for large peritoneal macrophages^{88, 100, 101}. SPIC is important for the development of red pulp macrophages¹⁰², while LXR-deficient mice are defective in the generation of marginal zone and metallophilic macrophages¹⁰³.

Tissue-specific environment are important for the maintenance of macrophages identities^{88, 89}. Several studies also suggest that individual resident macrophage subsets follow a differentiation program during embryogenesis and early post-natal life that result in the establishment of stable phenotypic and functional subsets^{22, 48}. Future works investigating the the respective contribution and interplay of lineage-specific 'hard-wired' differentiation programs and of the local tissue micro-environment for the differentiation, maintenance and activation of tissue-specific resident macrophages will contribute to the understanding of their functions.

REFERENCES

1. van Furth R, Cohn ZA, Hirsch JG, Humphrey JH, Spector WG, Langevoort HL. The mononuclear phagocyte system: a new classification of macrophages, monocytes, and their precursor cells. *Bull World Health Organ.* 1972; 46(6):845–852. [PubMed: 4538544]
2. van Furth R, Cohn ZA. The origin and kinetics of mononuclear phagocytes. *J Exp Med.* 1968; 128(3):415–435. [PubMed: 5666958]
3. Katz SI, Tamaki K, Sachs DH. Epidermal Langerhans cells are derived from cells originating in bone marrow. *Nature.* 1979; 282(5736):324–326. [PubMed: 503208]
4. Fogg DK, Sibon C, Miled C, Jung S, Aucouturier P, Littman DR, et al. A clonogenic bone marrow progenitor specific for macrophages and dendritic cells. *Science.* 2006; 311(5757):83–87. [PubMed: 16322423]
5. Liu K, Waskow C, Liu X, Yao K, Hoh J, Nussenzweig M. Origin of dendritic cells in peripheral lymphoid organs of mice. *Nature immunology.* 2007; 8(6):578–583. [PubMed: 17450143]
6. Liu K, Victora GD, Schwickert TA, Guermonprez P, Meredith MM, Yao K, et al. In vivo analysis of dendritic cell development and homeostasis. *Science.* 2009; 324(5925):392–397. [PubMed: 19286519]
7. Auffray C, Fogg DK, Narni-Mancinelli E, Senechal B, Trouillet C, Saederup N, et al. CX3CR1+ CD115+ CD135+ common macrophage/DC precursors and the role of CX3CR1 in their response to inflammation. *The Journal of experimental medicine.* 2009; 206(3):595–606. [PubMed: 19273628]
8. Gordon S, Martinez FO. Alternative activation of macrophages: mechanism and functions. *Immunity.* 2010; 32(5):593–604. [PubMed: 20510870]
9. Gordon S. Alternative activation of macrophages. *Nature reviews Immunology.* 2003; 3(1):23–35.
10. Murray PJ, Allen JE, Biswas SK, Fisher EA, Gilroy DW, Goerdt S, et al. Macrophage activation and polarization: nomenclature and experimental guidelines. *Immunity.* 2014; 41(1):14–20. [PubMed: 25035950]
11. Schultze JL, Freeman T, Hume DA, Latz E. A transcriptional perspective on human macrophage biology. *Seminars in immunology.* 2015; 27(1):44–50. [PubMed: 25843246]
12. Schulz C, Gomez Perdiguero E, Chorro L, Szabo-Rogers H, Cagnard N, Kierdorf K, et al. A lineage of myeloid cells independent of Myb and hematopoietic stem cells. *Science.* 2012; 336(6077):86–90. [PubMed: 22442384]
13. Merad M, Manz MG, Karsunky H, Wagers A, Peters W, Charo I, et al. Langerhans cells renew in the skin throughout life under steady-state conditions. *Nature immunology.* 2002; 3(12):1135–1141. [PubMed: 12415265]
14. Ginhoux F, Greter M, Leboeuf M, Nandi S, See P, Gokhan S, et al. Fate mapping analysis reveals that adult microglia derive from primitive macrophages. *Science.* 2010; 330(6005):841–845. [PubMed: 20966214]
15. Hashimoto D, Chow A, Noizat C, Teo P, Beasley MB, Leboeuf M, et al. Tissue-resident macrophages self-maintain locally throughout adult life with minimal contribution from circulating monocytes. *Immunity.* 2013; 38(4):792–804. [PubMed: 23601688]
16. Yona S, Kim KW, Wolf Y, Mildner A, Varol D, Breker M, et al. Fate mapping reveals origins and dynamics of monocytes and tissue macrophages under homeostasis. *Immunity.* 2013; 38(1):79–91. [PubMed: 23273845]
17. Epelman S, Lavine KJ, Beaudin AE, Sojka DK, Carrero JA, Calderon B, et al. Embryonic and adult-derived resident cardiac macrophages are maintained through distinct mechanisms at steady state and during inflammation. *Immunity.* 2014; 40(1):91–104. [PubMed: 24439267]
18. Gomez Perdiguero E, Klapproth K, Schulz C, Busch K, Azzoni E, Crozet L, et al. Tissue-resident macrophages originate from yolk-sac-derived erythro-myeloid progenitors. *Nature.* 2015; 518(7540):547–551. [PubMed: 25470051]
19. Geissmann F, Manz MG, Jung S, Sieweke MH, Merad M, Ley K. Development of monocytes, macrophages, and dendritic cells. *Science.* 2010; 327(5966):656–661. [PubMed: 20133564]
20. Takahashi K, Yamamura F, Naito M. Differentiation, maturation, and proliferation of macrophages in the mouse yolk sac: a light-microscopic, enzyme-cytochemical, immunohistochemical, and ultrastructural study. *Journal of leukocyte biology.* 1989; 45(2):87–96. [PubMed: 2536795]

21. Alliot F, Godin I, Pessac B. Microglia derive from progenitors, originating from the yolk sac, and which proliferate in the brain. *Brain research Developmental brain research*. 1999; 117(2):145–152. [PubMed: 10567732]
22. Chorro L, Sarde A, Li M, Woollard KJ, Chambon P, Malissen B, et al. Langerhans cell (LC) proliferation mediates neonatal development, homeostasis, and inflammation-associated expansion of the epidermal LC network. *The Journal of experimental medicine*. 2009; 206(13):3089–3100. [PubMed: 19995948]
23. Davies LC, Rosas M, Smith PJ, Fraser DJ, Jones SA, Taylor PR. A quantifiable proliferative burst of tissue macrophages restores homeostatic macrophage populations after acute inflammation. *European journal of immunology*. 2011; 41(8):2155–2164. [PubMed: 21710478]
24. Lawson LJ, Perry VH, Gordon S. Turnover of resident microglia in the normal adult mouse brain. *Neuroscience*. 1992; 48(2):405–415. [PubMed: 1603325]
25. Ghigo C, Mondor I, Jorquera A, Nowak J, Wienert S, Zahner SP, et al. Multicolor fate mapping of Langerhans cell homeostasis. *The Journal of experimental medicine*. 2013; 210(9):1657–1664. [PubMed: 23940255]
26. Kanitakis J, Morelon E, Petruzzo P, Badet L, Dubernard JM. Self-renewal capacity of human epidermal Langerhans cells: observations made on a composite tissue allograft. *Experimental dermatology*. 2011; 20(2):145–146. [PubMed: 20707812]
27. Bouwens L, Baekeland M, De Zanger R, Wisse E. Quantitation, tissue distribution and proliferation kinetics of Kupffer cells in normal rat liver. *Hepatology*. 1986; 6(4):718–722. [PubMed: 3733004]
28. Bain CC, Bravo-Blas A, Scott CL, Gomez Perdiguero E, Geissmann F, Henri S, et al. Constant replenishment from circulating monocytes maintains the macrophage pool in the intestine of adult mice. *Nature immunology*. 2014; 15(10):929–937. [PubMed: 25151491]
29. Bleriot C, Dupuis T, Jouvion G, Eberl G, Disson O, Lecuit M. Liver-resident macrophage necroptosis orchestrates type 1 microbicidal inflammation and type-2-mediated tissue repair during bacterial infection. *Immunity*. 2015; 42(1):145–158. [PubMed: 25577440]
30. Naito M, Takahashi K, Nishikawa S. Development, differentiation, and maturation of macrophages in the fetal mouse liver. *Journal of leukocyte biology*. 1990; 48(1):27–37. [PubMed: 2358750]
31. Takahashi K, Takahashi H, Naito M, Sato T, Kojima M. Ultrastructural and functional development of macrophages in the dermal tissue of rat fetuses. *Cell and tissue research*. 1983; 232(3):539–552. [PubMed: 6883456]
32. Deimann W, Fahimi HD. Peroxidase cytochemistry and ultrastructure of resident macrophages in fetal rat liver. A developmental study. *Developmental biology*. 1978; 66(1):43–56. [PubMed: 751843]
33. Bertrand JY, Giroux S, Golub R, Klaine M, Jalil A, Boucontet L, et al. Characterization of purified intraembryonic hematopoietic stem cells as a tool to define their site of origin. *Proc Natl Acad Sci U S A*. 2005; 102(1):134–139. [PubMed: 15623562]
34. Bertrand JY, Jalil A, Klaine M, Jung S, Cumano A, Godin I. Three pathways to mature macrophages in the early mouse yolk sac. *Blood*. 2005; 106(9):3004–3011. [PubMed: 16020514]
35. McGrath KE, Frame JM, Fegan KH, Bowen JR, Conway SJ, Catherman SC, et al. Distinct Sources of Hematopoietic Progenitors Emerge before HSCs and Provide Functional Blood Cells in the Mammalian Embryo. *Cell reports*. 2015; 11(12):1892–1904. [PubMed: 26095363]
36. Palis J, Robertson S, Kennedy M, Wall C, Keller G. Development of erythroid and myeloid progenitors in the yolk sac and embryo proper of the mouse. *Development*. 1999; 126(22):5073–5084. [PubMed: 10529424]
37. Palis J. Primitive and definitive erythropoiesis in mammals. *Frontiers in physiology*. 2014; 53
38. Ueno H, Weissman IL. Clonal analysis of mouse development reveals a polyclonal origin for yolk sac blood islands. *Developmental cell*. 2006; 11(4):519–533. [PubMed: 17011491]
39. Kissa K, Herbomel P. Blood stem cells emerge from aortic endothelium by a novel type of cell transition. *Nature*. 2010; 464(7285):112–115. [PubMed: 20154732]
40. Okuda T, van Deursen J, Hiebert SW, Grosveld G, Downing JR. AML1, the target of multiple chromosomal translocations in human leukemia, is essential for normal fetal liver hematopoiesis. *Cell*. 1996; 84(2):321–330. [PubMed: 8565077]

41. Le Guyader D, Redd MJ, Colucci-Guyon E, Murayama E, Kissa K, Briolat V, et al. Origins and unconventional behavior of neutrophils in developing zebrafish. *Blood*. 2008; 111(1):132–141. [PubMed: 17875807]
42. England SJ, McGrath KE, Frame JM, Palis J. Immature erythroblasts with extensive ex vivo self-renewal capacity emerge from the early mammalian fetus. *Blood*. 2011; 117(9):2708–2717. [PubMed: 21127173]
43. Chen MJ, Li Y, De Obaldia ME, Yang Q, Yzaguirre AD, Yamada-Inagawa T, et al. Erythroid/myeloid progenitors and hematopoietic stem cells originate from distinct populations of endothelial cells. *Cell stem cell*. 2011; 9(6):541–552. [PubMed: 22136929]
44. Chen MJ, Yokomizo T, Zeigler BM, Dzierzak E, Speck NA. Runx1 is required for the endothelial to haematopoietic cell transition but not thereafter. *Nature*. 2009; 457(7231):887–891. [PubMed: 19129762]
45. Lux CT, Yoshimoto M, McGrath K, Conway SJ, Palis J, Yoder MC. All primitive and definitive hematopoietic progenitor cells emerging before E10 in the mouse embryo are products of the yolk sac. *Blood*. 2008; 111(7):3435–3438. [PubMed: 17932251]
46. Rampon C, Huber P. Multilineage hematopoietic progenitor activity generated autonomously in the mouse yolk sac: analysis using angiogenesis-defective embryos. *The International journal of developmental biology*. 2003; 47(4):273–280. [PubMed: 12755332]
47. Kieusseian A, Brunet de la Grange P, Burlen-Defranoux O, Godin I, Cumano A. Immature hematopoietic stem cells undergo maturation in the fetal liver. *Development*. 2012; 139(19):3521–3530. [PubMed: 22899849]
48. Kierdorf K, Erny D, Goldmann T, Sander V, Schulz C, Perdiguero EG, et al. Microglia emerge from erythromyeloid precursors via Pu.1- and Irf8-dependent pathways. *Nature neuroscience*. 2013; 16(3):273–280. [PubMed: 23334579]
49. Sumner R, Crawford A, Mucenski M, Frampton J. Initiation of adult myelopoiesis can occur in the absence of c-Myb whereas subsequent development is strictly dependent on the transcription factor. *Oncogene*. 2000; 19(30):3335–3342. [PubMed: 10918590]
50. Bartunek P, Kralova J, Blendinger G, Dvorak M, Zenke M. GATA-1 and c-myc crosstalk during red blood cell differentiation through GATA-1 binding sites in the cmyb promoter. *Oncogene*. 2003; 22(13):1927–1935. [PubMed: 12673198]
51. Davidson CJ, Tirouvanziam R, Herzenberg LA, Lipsick JS. Functional evolution of the vertebrate Myb gene family: B-Myb, but neither A-Myb nor C-Myb, complements drosophila Myb in hemocytes. *Genetics*. 2005; 169(1):215–229. [PubMed: 15489525]
52. Cumano A, Godin I. Ontogeny of the hematopoietic system. *Annual review of immunology*. 2007; 25:745–785.
53. Orkin SH, Zon LI. Hematopoiesis: an evolving paradigm for stem cell biology. *Cell*. 2008; 132(4):631–644. [PubMed: 18295580]
54. Bertrand JY, Chi NC, Santos B, Teng S, Stainier DY, Traver D. Haematopoietic stem cells derive directly from aortic endothelium during development. *Nature*. 2010; 464(7285):108–111. [PubMed: 20154733]
55. Boisset JC, van Cappellen W, Andrieu-Soler C, Galjart N, Dzierzak E, Robin C. In vivo imaging of haematopoietic cells emerging from the mouse aortic endothelium. *Nature*. 2010; 464(7285):116–120. [PubMed: 20154729]
56. Zovein AC, Turlo KA, Ponc RM, Lynch MR, Chen KC, Hofmann JJ, et al. Vascular remodeling of the vitelline artery initiates extravascular emergence of hematopoietic clusters. *Blood*. 2010; 116(18):3435–3444. [PubMed: 20699440]
57. Delassus S, Cumano A. Circulation of hematopoietic progenitors in the mouse embryo. *Immunity*. 1996; 4(1):97–106. [PubMed: 8574856]
58. Mucenski ML, McLain K, Kier AB, Swerdlow SH, Schreiner CM, Miller TA, et al. A functional c-myc gene is required for normal murine fetal hepatic hematopoiesis. *Cell*. 1991; 65(4):677–689. [PubMed: 1709592]
59. Mukoyama Y, Chiba N, Mucenski ML, Satake M, Miyajima A, Hara T, et al. Hematopoietic cells in cultures of the murine embryonic aorta-gonad-mesonephros region are induced by c-Myb. *Current biology : CB*. 1999; 9(15):833–836. [PubMed: 10469571]

60. Soza-Ried C, Hess I, Netuschil N, Schorpp M, Boehm T. Essential role of c-myb in definitive hematopoiesis is evolutionarily conserved. *Proceedings of the National Academy of Sciences of the United States of America*. 2010; 107(40):17304–17308. [PubMed: 20823231]
61. Kumano K, Chiba S, Kunisato A, Sata M, Saito T, Nakagami-Yamaguchi E, et al. Notch1 but not Notch2 is essential for generating hematopoietic stem cells from endothelial cells. *Immunity*. 2003; 18(5):699–711. [PubMed: 12753746]
62. Muller AM, Medvinsky A, Strouboulis J, Grosveld F, Dzierzak E. Development of hematopoietic stem cell activity in the mouse embryo. *Immunity*. 1994; 1(4):291–301. [PubMed: 7889417]
63. Tober J, McGrath KE, Palis J. Primitive erythropoiesis and megakaryopoiesis in the yolk sac are independent of c-myb. *Blood*. 2008; 111(5):2636–2639. [PubMed: 18174377]
64. Yokomizo T, Ogawa M, Osato M, Kanno T, Yoshida H, Fujimoto T, et al. Requirement of Runx1/AML1/PEBP2alphaB for the generation of haematopoietic cells from endothelial cells. *Genes to cells : devoted to molecular & cellular mechanisms*. 2001; 6(1):13–23. [PubMed: 11168593]
65. Qian BZ, Li J, Zhang H, Kitamura T, Zhang J, Campion LR, et al. CCL2 recruits inflammatory monocytes to facilitate breast-tumour metastasis. *Nature*. 2011; 475(7355):222–225. [PubMed: 21654748]
66. Metzger D, Clifford J, Chiba H, Chambon P. Conditional site-specific recombination in mammalian cells using a ligand-dependent chimeric Cre recombinase. *Proceedings of the National Academy of Sciences of the United States of America*. 1995; 92(15):6991–6995. [PubMed: 7624356]
67. Hoeffel G, Chen J, Lavin Y, Low D, Almeida FF, See P, et al. C-Myb(+) erythromyeloid progenitor-derived fetal monocytes give rise to adult tissue-resident macrophages. *Immunity*. 2015; 42(4):665–678. [PubMed: 25902481]
68. Hoeffel G, Wang Y, Greter M, See P, Teo P, Malleret B, et al. Adult Langerhans cells derive predominantly from embryonic fetal liver monocytes with a minor contribution of yolk sac-derived macrophages. *The Journal of experimental medicine*. 2012
69. Williams M, De Kleer I, Henri S, Post S, Vanhoutte L, De Prijck S, et al. Alveolar macrophages develop from fetal monocytes that differentiate into long-lived cells in the first week of life via GM-CSF. *The Journal of experimental medicine*. 2013; 210(10):1977–1992. [PubMed: 24043763]
70. Sheng J, Ruedl C, Karjalainen K. Most Tissue-Resident Macrophages Except Microglia Are Derived from Fetal Hematopoietic Stem Cells. *Immunity*. 2015; 43(2):382–393. [PubMed: 26287683]
71. Busch K, Klapproth K, Barile M, Flossdorf M, Holland-Letz T, Schlenner SM, et al. Fundamental properties of unperturbed haematopoiesis from stem cells in vivo. *Nature*. 2015; 518(7540):542–546. [PubMed: 25686605]
72. Ema M, Yokomizo T, Wakamatsu A, Terunuma T, Yamamoto M, Takahashi S. Primitive erythropoiesis from mesodermal precursors expressing VE-cadherin, PECAM-1, Tie2, endoglin, and CD34 in the mouse embryo. *Blood*. 2006; 108(13):4018–4024. [PubMed: 16926294]
73. Garceau V, Balic A, Garcia-Morales C, Sauter KA, McGrew MJ, Smith J, et al. The development and maintenance of the mononuclear phagocyte system of the chick is controlled by signals from the macrophage colony-stimulating factor receptor. *BMC biology*. 2015; 13:12. [PubMed: 25857347]
74. Buza-Vidas N, Woll P, Hultquist A, Duarte S, Lutteropp M, Bouriez-Jones T, et al. FLT3 expression initiates in fully multipotent mouse hematopoietic progenitor cells. *Blood*. 2011; 118(6):1544–1548. [PubMed: 21628405]
75. Molawi K, Wolf Y, Kandalla PK, Favret J, Hagemeyer N, Frenzel K, et al. Progressive replacement of embryo-derived cardiac macrophages with age. *The Journal of experimental medicine*. 2014; 211(11):2151–2158. [PubMed: 25245760]
76. Varol C, Vallon-Eberhard A, Elinav E, Aychek T, Shapira Y, Luche H, et al. Intestinal lamina propria dendritic cell subsets have different origin and functions. *Immunity*. 2009; 31(3):502–512. [PubMed: 19733097]
77. Hickey WF, Vass K, Lassmann H. Bone marrow-derived elements in the central nervous system: an immunohistochemical and ultrastructural survey of rat chimeras. *Journal of neuropathology and experimental neurology*. 1992; 51(3):246–256. [PubMed: 1583531]

78. Yamada M, Naito M, Takahashi K. Kupffer cell proliferation and glucan-induced granuloma formation in mice depleted of blood monocytes by strontium-89. *Journal of leukocyte biology*. 1990; 47(3):195–205. [PubMed: 2307905]
79. Golde DW, Byers LA, Finley TN. Proliferative capacity of human alveolar macrophage. *Nature*. 1974; 247(440):373–375. [PubMed: 4817856]
80. Sawyer RT, Strausbauch PH, Volkman A. Resident macrophage proliferation in mice depleted of blood monocytes by strontium-89. *Laboratory investigation; a journal of technical methods and pathology*. 1982; 46(2):165–170.
81. Jenkins SJ, Ruckerl D, Cook PC, Jones LH, Finkelman FD, van Rooijen N, et al. Local macrophage proliferation, rather than recruitment from the blood, is a signature of TH2 inflammation. *Science*. 2011; 332(6035):1284–1288. [PubMed: 21566158]
82. Bruttger J, Karram K, Wortge S, Regen T, Marini F, Hoppmann N, et al. Genetic Cell Ablation Reveals Clusters of Local Self-Renewing Microglia in the Mammalian Central Nervous System. *Immunity*. 2015; 43(1):92–106. [PubMed: 26163371]
83. Aziz A, Soucie E, Sarrazin S, Sieweke MH. MafB/c-Maf deficiency enables self-renewal of differentiated functional macrophages. *Science*. 2009; 326(5954):867–871. [PubMed: 19892988]
84. Jenkins SJ, Ruckerl D, Thomas GD, Hewitson JP, Duncan S, Brombacher F, et al. IL-4 directly signals tissue-resident macrophages to proliferate beyond homeostatic levels controlled by CSF-1. *The Journal of experimental medicine*. 2013; 210(11):2477–2491. [PubMed: 24101381]
85. van Furth R, Diesselhoff-den Dulk MM. Dual origin of mouse spleen macrophages. *The Journal of experimental medicine*. 1984; 160(5):1273–1283. [PubMed: 6491600]
86. Ajami B, Bennett JL, Krieger C, McNagny KM, Rossi FM. Infiltrating monocytes trigger EAE progression, but do not contribute to the resident microglia pool. *Nature neuroscience*. 2011; 14(9):1142–1149. [PubMed: 21804537]
87. Gomez-Nicola D, Franssen NL, Suzzi S, Perry VH. Regulation of microglial proliferation during chronic neurodegeneration. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 2013; 33(6):2481–2493. [PubMed: 23392676]
88. Gosselin D, Link VM, Romanoski CE, Fonseca GJ, Eichenfield DZ, Spann NJ, et al. Environment drives selection and function of enhancers controlling tissue-specific macrophage identities. *Cell*. 2014; 159(6):1327–1340. [PubMed: 25480297]
89. Lavin Y, Winter D, Blecher-Gonen R, David E, Keren-Shaul H, Merad M, et al. Tissue-resident macrophage enhancer landscapes are shaped by the local microenvironment. *Cell*. 2014; 159(6):1312–1326. [PubMed: 25480296]
90. Yamamoto T, Kaizu C, Kawasaki T, Hasegawa G, Umezumi H, Ohashi R, et al. Macrophage colony-stimulating factor is indispensable for repopulation and differentiation of Kupffer cells but not for splenic red pulp macrophages in osteopetrotic (*op/op*) mice after macrophage depletion. *Cell and tissue research*. 2008; 332(2):245–256. [PubMed: 18335245]
91. Butovsky O, Jedrychowski MP, Moore CS, Cialic R, Lanser AJ, Gabriely G, et al. Identification of a unique TGF-beta-dependent molecular and functional signature in microglia. *Nature neuroscience*. 2014; 17(1):131–143. [PubMed: 24316888]
92. Wang Y, Szretter KJ, Vermi W, Gilfillan S, Rossini C, Cella M, et al. IL-34 is a tissue-restricted ligand of CSF1R required for the development of Langerhans cells and microglia. *Nature immunology*. 2012; 13(8):753–760. [PubMed: 22729249]
93. Borkowski TA, Letterio JJ, Farr AG, Udey MC. A role for endogenous transforming growth factor beta 1 in Langerhans cell biology: the skin of transforming growth factor beta 1 null mice is devoid of epidermal Langerhans cells. *The Journal of experimental medicine*. 1996; 184(6):2417–2422. [PubMed: 8976197]
94. Lieschke GJ, Stanley E, Grail D, Hodgson G, Sinickas V, Gall JA, et al. Mice lacking both macrophage- and granulocyte-macrophage colony-stimulating factor have macrophages and coexistent osteopetrosis and severe lung disease. *Blood*. 1994; 84(1):27–35. [PubMed: 8018921]
95. Shibata Y, Berclaz PY, Chronos ZC, Yoshida M, Whitsett JA, Trapnell BC. GM-CSF regulates alveolar macrophage differentiation and innate immunity in the lung through PU.1. *Immunity*. 2001; 15(4):557–567. [PubMed: 11672538]

96. Happle C, Lachmann N, Skuljec J, Wetzke M, Ackermann M, Brenning S, et al. Pulmonary transplantation of macrophage progenitors as effective and long-lasting therapy for hereditary pulmonary alveolar proteinosis. *Science translational medicine*. 2014; 6(250):250ra113.
97. Willinger T, Rongvaux A, Takizawa H, Yancopoulos GD, Valenzuela DM, Murphy AJ, et al. Human IL-3/GM-CSF knock-in mice support human alveolar macrophage development and human immune responses in the lung. *Proceedings of the National Academy of Sciences of the United States of America*. 2011; 108(6):2390–2395. [PubMed: 21262803]
98. Akagawa KS, Kamoshita K, Tokunaga T. Effects of granulocyte-macrophage colony-stimulating factor and colony-stimulating factor-1 on the proliferation and differentiation of murine alveolar macrophages. *J Immunol*. 1988; 141(10):3383–3390. [PubMed: 3053898]
99. Chen BD, Mueller M, Chou TH. Role of granulocyte/macrophage colony-stimulating factor in the regulation of murine alveolar macrophage proliferation and differentiation. *J Immunol*. 1988; 141(1):139–144. [PubMed: 3288696]
100. Okabe Y, Medzhitov R. Tissue-specific signals control reversible program of localization and functional polarization of macrophages. *Cell*. 2014; 157(4):832–844. [PubMed: 24792964]
101. Rosas M, Davies LC, Giles PJ, Liao CT, Kharfan B, Stone TC, et al. The transcription factor Gata6 links tissue macrophage phenotype and proliferative renewal. *Science*. 2014; 344(6184):645–648. [PubMed: 24762537]
102. Kohyama M, Ise W, Edelson BT, Wilker PR, Hildner K, Mejia C, et al. Role for Spi-C in the development of red pulp macrophages and splenic iron homeostasis. *Nature*. 2009; 457(7227):318–321. [PubMed: 19037245]
103. N AG, Guillen JA, Gallardo G, Diaz M, de la Rosa JV, Hernandez IH, et al. The nuclear receptor LXRalpha controls the functional specialization of splenic macrophages. *Nature immunology*. 2013; 14(8):831–839. [PubMed: 23770640]




Figure 1.

Time scale of hematopoietic progenitors development in the mouse embryo. There are three successive but overlapping waves of hematopoietic progenitors during development, all of which have the potential to give rise to fetal macrophages. They can be distinguished by the hematopoietic niches exist where the progenitors expand and differentiate during development (upper panel), the yolk sac (primitive progenitors in green and EMP in blue), the fetal liver (EMP and HSC), and finally the bone marrow (HSC, orange). While primitive progenitors are only described in a small time window (middle panel), definitive progenitors (including EMPs and HSCs) co-exist during most fetal development in the fetal liver. Only HSC-derived hematopoiesis then shifts to the bone marrow niche. The three waves of progenitor can also be distinguished by their differentiation potential *in vivo* (lower panel). Primitive progenitors are restricted to the erythroid or the myeloid lineages, while EMP have both erythroid and myeloid potential. EMP-derived hematopoiesis gives rise to erythrocytes, macrophages, monocytes, granulocytes and mast cells and is sufficient to support survival of embryos lacking HSCs until birth. EMP-derived macrophages persist after birth in most tissues as resident macrophages, with the exception of the intestine lamina propria. HSCs are distinguished from other progenitors by their capacity to perform long-term repopulation of all leukocytes lineages when transplanted into a conventional irradiated recipient mouse.




Figure 2. Schematic of myeloid development and maintenance in the mouse embryo and adult (see text). Circular red arrows indicate self renewal potential. Primitive progenitors do not share a common origin with endothelial cells lining the blood islands in the yolk sac (YS) and they emerge the absence of *Runx1* and *Myb*. Within definitive hematopoiesis, HSC and EMP arise from distinct hemogenic endothelial cells through a *Runx1*-dependent endothelial to hematopoietic transition. Both EMP and HSC express the transcription factor *Myb*, and while no fetal or adult HSC-derived hematopoiesis can occur in the absence of *Myb*, EMPs are

Myb-independent for their myeloid differentiation. EMP-derived hematopoiesis gives rise to erythrocytes and short-lived myeloid cells (monocytes, granulocytes, mast cells) that are replaced by HSC-derived cells late during gestation. EMP-derived macrophages colonise all tissues during fetal development where they specialize to their tissue of residency after birth and can persist throughout adult life by local proliferation (red arrow). Depending on the age and environmental challenges, HSC-derived cells can contribute to adult tissue resident populations.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1

phenotype, differentiation potential, and molecular features of hematopoietic progenitors of

Progenitor name	Primitive		Definitive	
	Mp	EMp	HSC	
Origin	Posterior plate mesoderm	hemogenic endothelium	Arterial hemogenic endothelium	
Progenitor surface phenotype	Kit ^{lo} CD41 ^{lo}	Kit ⁺ CD41 ⁺ VE-Cad ⁺ CD16/32 ⁺ AA4.1 ⁺ Tie2 ⁺ CD45 ^{lo} Sca1 ⁻	Lin ⁻ Kit ⁺ Sca1 ⁺ CD41 ⁺ VE- Cad ⁺ Tie2 ⁺ CD45 ⁺	
time window of generation	E7.0-?	E8.25-E10.5	E9.5-E12.5	
Anatomical location of	Generation	YS (blood islands)	YS	AGM (and large arteries)
	Expansion	YS	FL	FL
	Differentiation/commitment	YS	YS and FL	FL, bone marrow
Differentiation potential			Erythroid	Erythroid
		Myeloid	Myeloid	Myeloid
				Lymphoid
Macrophage lineage		Pu.1-dependent	Pu.1-dependent	Pu.1-dependent
		Runx1-independent	Runx1 -dependent	Runx1 -dependent
		Myb-independent	Myb-independent	Myb-dependent
			Notch1 independent	Notch1-dependent
Macrophage self-renewal	?	yes		
progenitor long-term self-renewal	no (transient)	no (transient)		yes
TF expression	Pu.1 (sp1)	n/a	+	+
	Myb	n/a	+	+
	Gata2	n/a	+	+
	Gata3	n/a	-	+
	Lmo2	n/a	-	+
TF requirement	Pu.1 (sp1)	yes	yes	no (**)
	Myb	no	no (*)	yes
	Runx1	no	yes	yes
	CBFb	no	yes	yes
	Scl/tal-1	yes	yes	yes
	Notch1	no	no	yes

(*) red blood cell differentiation is compromised

(**) macrophage differentiation is compromised